1 NK Cell Exhaustion in Wilson's Disease Revealed by Single-cell RNA Sequencing Predicts the Prognosis

2   of Cholecystitis

Yong Jin[1, #], Jiayu Xing[1, #], Chenyu Dai[2, #], Lei Jin[3, 4], Wanying Zhang[1], Qianqian Tao[3, 4], Mei Hou[1], Ziyi Li[3, 4], Wen Yang[1, 5, *], Qiyu Feng[1, *], Hongyang Wang[1, 5, *] & Qingsheng Yu[2, 3, *]

[1] Cancer Research Center, The First Affiliated Hospital of USTC, Division of Life Sciences and Medicine,

University of Science and Technology of China, Hefei, Anhui, 230036, P. R. China.

[2] Department of Cadre Cardiology, The First Affiliated Hospital of Anhui University of Chinese Medicine,

Hefei, Anhui, 230031, P. R. China.

[3] Department of General Surgery, The First Affiliated Hospital of Anhui University of Chinese Medicine,

Hefei, Anhui, 230031, P. R. China.

[4] Institute of Chinese Medicine Surgery, Anhui Academy of Chinese Medicine, Hefei, Anhui, 230031, P. R.

China.

[5] International Co-operation Laboratory on Signal Transduction, Eastern Hepatobiliary Surgery Hospital,

Second Military Medical University, Shanghai 200438, China; National Center for Liver Cancer, Second

Military Medical University, Shanghai 201805, China.

[#] Contributed equally.

[*] Corresponding author.

Correspondence to Qingsheng Yu (qsy6312@163.com), Hongyang Wang (hywangk@vip.sina.com), or

Qiyu Feng (qiyufeng@ustc.edu.cn).

Abstract

Metabolic abnormalities associated with liver disease have a significant impact on the risk and prognosis of

cholecystitis.   However, the underlying mechanism is complicated and remains to be elucidated.   In

particular, the effect of metabolic abnormalities on the progression of cholecystitis through the regulation of

immune cell function is poorly understood.   In this study, we investigated this issue using Wilson's disease

(WD) as a model.   Wilson's disease is a genetic disorder characterized by impaired mitochondrial function

and abnormal copper metabolism.   Our retrospective clinical study of over 600 patients with WD found

that they have a significantly higher incidence of cholecystitis and a poorer prognosis.   The immune cell

landscape in the hepatic mesenchymal stromal microenvironment using single-cell RNA sequencing

showed that the tissue immune microenvironment is altered in patients with WD, mainly a major change in

the constitution and function of the innate immune system, including enhanced antigen presentation process,

activation of the immune response, and activation of lymphocytes.   Exhaustion of natural killer (NK) cells

is the fundamental factor, supported by an increase in the expression of the inhibitory receptors NKG2A

and TIGIT and a decrease in the expression of cytotoxic molecules.   Clinical tissue and blood samples

verified increased $NKG2A^+$ and $TIGIT^+$ NK cells and decreased $IFN\gamma^+$ NK cells in WD.   Further

bioinformatic analysis has confirmed a positive correlation between NK cell exhaustion and poor prognosis

in cholecystitis and other inflammatory diseases.   The study demonstrated abnormal function of liver

mesenchymal immune cells triggered by specific metabolic dysfunction in WD, with a focus on the

correlation between NK cell exhaustion and poor healing of cholecystitis.   Our findings highlight the

immune cell dysfunction due to metabolic changes in hepatocytes and provide new insights into the

improvement of inflammatory diseases by assessing immune cell function.

Keywords: NK cell exhaustion, immune cell dysfunction, metabolic abnormalities, Wilson's disease,

cholecystitis

Introduction

Cholecystitis refers to a group of inflammatory diseases with different etiologies and clinical courses[1]. While the corresponding pathological mechanisms of cholecystitis are complicated, systemic inflammation is widely recognized as a primary catalyst of cholecystitis pathology[2-6]. Cholecystectomy is typically the preferred treatment option for patients with severe cholecystitis and gallstones, whereas medication is prescribed for those with mildly symptomatic cholecystitis or calculous cholecystitis. However, the effectiveness of alternative or conservative treatments can vary greatly. Furthermore, a substantial percentage of patients face the possibility of complications or recurrence[7,8]. Therefore, it is important to assess the prognosis of pharmacological treatment in patients with cholecystitis through prospective evaluation. Inflammatory diseases, ranging from common inflammatory patterns to rare chronic illnesses, are characterized by a defective immune system and an exaggerated inflammatory cascade response. Compelling evidence supports that the proportion, gene expression profile, and functional status of immune cells hold significant prognostic value[9-14]. Therefore, it is critical to comprehensively comprehend the immune cell dysfunction linked with cholecystitis, its corresponding altered characteristics, and their prognostic implications.

Natural killer (NK) cells are the major innate lymphocyte subset and are found in most organs[15,16]. They produce lytic granules that eliminate infected or cancerous cells, coordinate the inflammation process, form immunological memory, and modulate antigen-presenting cell function in inflammatory diseases[17-19]. In the context of tumors and chronic infections, NK cells exhibit an exhausted phenotype, termed NK cell exhaustion, which is characterized by downregulated expression of activating receptors (e.g., NKG2D, CD16), upregulated expression of inhibitory receptors (e.g., NKG2A, PD-1, TIGIT, Tim-3), decreased production of effector cytokines, and impaired cytolytic activity[20,21]. Recently, blockade of NKG2A and TIGIT has been shown to reverse NK cell exhaustion and enhance the therapeutic effects of anti-tumor and anti-infective immunotherapies[21-23]. In addition, NK cell exhaustion has been reported to predict poor prognosis in hepatocellular carcinoma and acute myeloid leukemia; however, the correlation of NK cell exhaustion with prognosis in cholecystitis and other inflammatory diseases remains largely unexplored[24-26]. The liver plays a critical role in immune defense. It contains a substantial population of diverse immune

cells, particularly NK cells[27-29]. The liver modulates these cells to mount immune responses upon

encountering specific inflammatory triggers[28,30]. Furthermore, liver pathologies affect mesenchymal

immune cell function by means of regulating intercellular signaling, thereby influencing systemic

inflammatory responses. Liver disease commonly impacts the development and outcome of

inflammation-associated diseases such as cholecystitis[31-33]. Wilson's disease (WD) is a rare genetic

disorder caused by mutations in the gene encoding the transmembrane copper-transporting P-type ATPase

protein[34]. It leads to deficits in mitochondrial structure and function, as well as specific alterations in liver

metabolism, resulting in a variety of liver diseases. Additionally, WD is commonly associated with

inflammatory conditions in the clinic[35-38]. Thus, WD, with its unique etiology, serves as an excellent

model for exploring the regulation of immune cell function by hepatic metabolic abnormalities.

In this retrospective clinical study of over six hundred WD patients, it was found that they had a

significantly higher cholecystitis incidence than the general population. This, in turn, is associated with a

poor prognosis for cholecystitis in these patients. A thorough examination of the immune cell landscape

in the hepatic mesenchymal stromal microenvironment using single-cell RNA sequencing (scRNA-seq)

revealed that the tissue immune microenvironment is altered in patients with WD. Specifically, there were

significant changes in the composition and function of the innate immune system, including an increase of

mononuclear phagocytes (MPs) and NK cells, as well as enhanced pro-inflammatory characteristics.

Additionally, there was an enhancement in the biological processes of antigen presentation, activation of

immune response, and activation of lymphocytes. One key event among these changes was the

exhaustion of NK cells, as demonstrated by an increase in the expression of inhibitory receptors NKG2A

and TIGIT and a decrease in the expression of cytotoxic molecules. Multiplex immunohistochemistry and

flow cytometry techniques confirmed the exhaustion of NK cells in both clinical liver tissue and peripheral

blood samples. Finally, our bioinformatics analysis has confirmed a positive correlation between NK cell

exhaustion and poor prognosis in cholecystitis and other inflammatory diseases. Our study has

demonstrated abnormal liver mesenchymal immune cell function triggered by specific metabolic

dysfunction in WD, focusing on NK cell exhaustion and its correlation with poor healing of cholecystitis.

These findings provide foundation for further studies of effect of hepatocytes metabolic changes on

immune cell function and insights into improvement of inflammatory diseases by assessing immune cell

function.

Results

WD causes poor prognosis of cholecystitis

WD is a genetic disorder resulting from mutations in the *ATP7B* gene, exhibiting significant genetic

heterogeneity with over 600 mutations or polymorphisms[39]. Through sequencing DNA samples from 20

patients with WD, we found that c. 2333G>T (p. R778L) accounted for the majority (65%) of the three

most common mutations in Chinese, while c. 2804C>T (p. T935M) and c. 2975C>T (p. P992L) were not

detected (Figure 1a)[40]. In hepatocytes, ATP7B deficiency in the trans-Golgi network (TGN) leads to

abnormalities in mitochondrial function and copper metabolism, and ultimately to a wide range of liver

diseases (Figure 1b)[41-44]. Typical clinical manifestations are liver fibrosis and Kayser-Fleischer Ring in

the cornea that caused by Cu accumulation in the cornea (Figure 1c, d)[41]. Additionally, metabolic

symptoms of WD patients, such as a higher plasma total cholesterol (T-CHOL) concentration and a lower

risk of hyperglycemia (Cohort 1 and Cohort 2), are observed (Figure 1e, Figure S1a). Mitochondrial

stress and glycolysis assays revealed that knockout of the *ATP7B* (ATP7B-KO) in human hepatocyte cell

lines WRL 68 significantly impaired the basal oxygen consumption rate (OCR) and capacity of ATP

production, while the extracellular acidification rate (ECAR) of glycolysis was slightly increased and

glycolysis capacity remained unchanged (Figure S1b, c). These results demonstrate that WD is a disease

characterized by abnormal hepatocyte metabolism.

Interestingly, chronic cholecystitis emerged as the most common complication, with a notably higher

incidence in WD patients (49.51%) compared to HBV patients (9.46%) in Cohort 2 (Figure 1e). Despite

the comparable incidence of gallbladder in the two groups and the significantly reduced plasma

concentration of total bilirubin in WD patients (Cohort 1 and Cohort 2), this finding holds (Figure 1e).

Moreover, in cholecystitis patients, the proportion of cases with abnormal index was significantly higher

when combined with WD, which included white blood cell (WBC), neutrophil (NEU), and lymphocyte

(LYM) counts (Cohort 3, Figure 1e). Overall, these findings suggest that cholecystitis outcomes are

negatively affected by WD and that there are immune abnormalities that affect the prognosis of

cholecystitis in WD patients.

Single-cell profiling of non-parenchymal cells in the liver of WD patients

The hepatic mesenchymal cells from three WD patients and three liver hemangioma patients were analyzed

using the 10X Genomics platform for scRNA-seq (Figure 2a) to better understand the complicated immune

abnormalities in WD patients. 57,384 high-quality single transcriptomes were generated, with 30,068

cells in the case group and 27,316 in the control group.   We validated twenty-six principal components *via*

PCA and substantiated the grouping through UMAP dimensionality reduction cluster graph (Figure 2b, c,

Figure S2a).   We then utilized individual cell expression profiles and reference to cell marker gene

expression to implement an automated annotation method, which was further complemented by manual

review and fine-tuning.   This initial annotation process resulted in the classification and display of 12

significant cell clusters.   The UMAP algorithm was used to segregate and display the clusters.   The study

identified various cell types, including hepatocytes[45,46], cholangiocytes[45], endothelial cells (ECs)[47-49],

hepatic stellate cells (HepSCs)[46,50], proliferating cells[48,51,52], B cells[53,54], plasma cells[52,53], T and NK cells[53,55],

neutrophils[56], mast cells[48,55,56], mononuclear phagocytes (MPs)[57], and plasmacytoid dendritic cells

(pDCs)[48,58] (Figure 2d, e and Figure S2b, c).   Figure S2d exhibited the top 10 differently expressed genes

(DEGs) for each cell type.

Based on the initial annotation, there were notable variations in the quantities and ratios of different cell

types between the case group and control group.   Our study observed that T and NK lineages were the

primary immune cell population among all the examined samples, accounting for 56.54% in CASE and

73.55% in CON, followed by MPs which constituted 17.33% in CASE and 9.97% in CON (Figure 2f,

Figure S2e, and Supplementary Table S1).   The significant increase in the proportion of MPs and decrease

in T and NK cells suggest that MPs likely dominate the immune microenvironment remodeling in patients

with WD, while lymphocytes play a minor role.   B cell proportions (4.53% in CASE and 2.27% in CON),

plasma cell proportions (2.36% in CASE and 0.42% in CON), and pDC proportions (0.42% in CASE and

0.16% in CON) were elevated, while neutrophil proportions (3.53% in CASE and 6.45% in CON) declined.

The proportion of remaining cells is shown in Supplementary Table S1.

The immune microenvironment was altered in WD patients

To demonstrate dissimilarities in the composition of the immune microenvironment between the two

groups of individuals in a more detailed manner, a secondary subtype clustering and annotation process was

applied to MPs and T and NK cells (Figure 3a, b, and Figure S3a, b).   MPs are myeloid immune cells that

serve phagocytic and antigen-presenting functions. These cells, which include monocytes, macrophages,

and dendritic cells, play an essential role in determining the nature of the immune response[59].   A total of

6,962 MPs were identified (4,439 in CASE and 2,523 in CON), and they were subdivided into 7 clusters

through annotation (Figure 3a, Figure S3a, and Supplementary Table S1).   Cell marker genes for

annotating cell subtypes were depicted in Figure S3c and d, and Figure S3e demonstrated the top 10 DEGs

between the two groups[46,48,49,51,52,55-58,60-65].   The DEGs were related to various cell types, including

proliferating cells, macrophages, monocytes, mature dendritic cells (MatureDCs), conventional type 1

dendritic cells (cDC1), conventional type 2 dendritic cells (cDC2), and Kupffer cells (KCs).   In the case

group, there was a significant increase in both macrophages (25.03% in CASE versus 14.33% in CON) and

KCs, the resident macrophages in the liver (26.03% in CASE versus 22.96% in CON), suggesting their

involvement in promoting hepatic inflammation in metabolic liver disease (Figure 3a)[66].   By contrast, the

percentages of monocytes (35.45% in CASE and 41.78% in CON), cDC1 (2.14% in CASE and 5.99% in

CON), and cDC2 (10.39% in CASE and 14.13% in CON) were decreased in the case group (Figure 3a).

Seventeen clusters were obtained through reclustering for the largest population of T and NK cells, totaling

34,339 cells (15,403 cells in CASE and 18,936 in CON)[53,56,67-82]. The clusters included ILC3_IL1R1,

NKT_GNLY, NKT_IFNG, NKT_KLRG1, NK_FCER1G, NK_KLRC1, NKT_XCL2, NK_FCGR3A,

CD4NaiveT_CCR7, CD4Tmem_CD40LG, CD4Treg_CTLA4, CD8MAIT_KLRB1, CD8MAIT_SLC4A10,

CD8Teff_GZMH, CD8Teff_GZMK, CD8Teff_ISG15, and CD8Teff_MT1X (Figure 3b, Figure S3b,

Supplementary Table S1).   The proportion of NK cells (40.97% and 17.07% in CASE and CON,

respectively) and CD4Treg (0.92% and 0.31% in CASE and CON, respectively) was significantly increased

in the case group.   On the other hand, a decrease of NKT cells (28.72% in CASE and 35.26% in CON),

CD8MAIT (7.76% in CASE and 22.14% in CON) and CD8Teff (14.95% in CASE and 17.88% in CON)

was observed in the case group. CD4 regulatory T cells are thought to inhibit T cell activation in

autoimmune and virus-related liver diseases, indicating a correlation between elevated CD4 regulatory T

cells and reduced CD8 T cells in WD patients[83].   The distribution of other types of cells between the two

groups is shown in Figure 3b and Supplementary Table S1.

To systematically evaluate the key functional alteration, we analyzed the top 20 differently expressed genes

in all immune cell populations.   We found that the expression of genes encoding complement C1q (*C1QA*,

C1QB, and *C1QC*) and *HLA-DRA* was significantly upregulated in the case group. Conversely, the

expression of genes encoding heat shock protein (*HSPA1A*, *HSPA1B*, *HSPM1*, *HSPA6*, *HSP90AB1*, *HSPE1*,

and *HSPD1*) was markedly downregulated in the case group (Figure 3c).   The Gene Ontology (GO)

analysis results demonstrate that WD enhances the abilities of antigen-presenting cells to activate immune

response and related lymphocytes, through upregulating antigen processing and presentation, activation of

immune response, and positive regulation of lymphocyte activation.   However, heat shock proteins

delivering antigens may be responsible for the downregulation of lymphocytes differentiation and T cell

activation[84].   Next, we analyzed the cellular communication network using Cellchat.   Our findings reveal

that CD8Teff, KCs, NKT, CD8MAIT, and monocytes were the primary partners in intercellular

communication (Figure 3e).   Furthermore, CD8Teff and NK cells showed stronger signals in the case

group, while neutrophils exhibited weaker signals, in both outgoing and incoming (Figure 3e).

Collectively, these results demonstrate that the immune microenvironment in the liver of WD patients is

remodeled, with an obvious enhancement in both quantity and function of innate immunocytes.

The dysfunction of main immune cells in WD patients

To obtain a detail analysis of function comparison in immune cells between two groups, we scored and

evaluated the core functional signature of major cell types. We further clustered macrophages, which

accounted for 14.33% of the MPs in the control group and increased to 25.03% in the case group, resulting

in the identification of four distinct subtypes (Figure S4a).   Interestingly, these four subtypes in the

patients with WD exhibited distinct gene expression profiles.   Macrophages_3 in WD displayed elevated

expression levels of S100A family genes (*S100A8*, *S100A9*, *S100A12*, *S100A6*, *VCAN*, *FCN1*, and *S100A4*),

indicating a potential contribution to MDSC-like macrophages. In contrast, Macrophages_4 in WD were

characterized by upregulation of lipid metabolism genes (*FABP5*, *APOC1*, *CTSD*, and *PLA2G7*) (Figure

S4b)[85,86]. By scoring various gene signatures, we compared the M1 polarization, M2 polarization,

pro-inflammatory signature, and interferon response signature of four subtypes within two groups (Figure

4a). These four subtypes displayed consistently overlapped M1/M2 signatures. Significantly, the

Macrophages_3 subtype exhibited a more pronounced pro-inflammatory signal in the case group, while the

other subtypes showed a slight decrease. In addition, the interferon response signature showed slight

elevation in four subtypes within the case group. Kupffer cells, which are liver-resident macrophages with

unique functions in maintaining homeostasis, represent the most abundant population of tissue-resident

macrophages in the human body[66]. KCs were also further clustered and three subtypes were obtained

(Figure S4c). Significant differential expression of genes related to the inflammatory response, including

IL1B, *CXCL2*, *CCL3C1*, *CXCL8*, and *CCL2*, were identified between the two groups (Figure S4d).

Similarly, KCs subtypes in two groups also showed overlapping M1/M2 signatures, but all subtypes in the

case group showed varying degrees of increase in pro-inflammatory and interferon-responsive signatures

(Figure 4b).

Neutrophils, the most abundant leukocytes in human blood, respond to inflammatory stimuli by migrating

to the site of inflammation. Its immunological functions, such as phagocytosis, reactive oxygen species

generation, degranulation, and production of cytokines and chemokines, play a crucial role in the resolution

of inflammation and tissue repair[87,88]. In the case group, a lower percentage of neutrophils (3.53% in

CASE and 6.45% in CON) was observed, and the functional characteristics were subsequently evaluated.

The results showed that the maturation, chemotaxis, phagocytosis, and chemokine activity scores were

decreased in the CASE group, while the activation of the type I interferon signaling pathway was enhanced

(Figure 4c).

Since the biological process of antigen processing and presentation was upregulated, we calculated the

relative expression intensity of genes encoding MHC molecules and complement molecules in cDC1, cDC2,

and mature DCs (Figure 4d). Notably, genes encoding MHC class I molecules (*HLA-A*, *-B*, *-C*, *-E*, and *-F*)

were significantly upregulated in the mature DCs in the case group, while genes encoding MHC class II

molecules (*HLA-DQA1*, *-DQA2*, *-DQB1*, *-DQB2*, *-DPA1*, *-DPB1*, *-DRA*, *-DRB1*, and *-DRB5*) were all

downregulated in the mature DCs.   Monocytes accounted for the highest proportion (35.45% in CASE and

41.78% in CON), and most of them were CD14$^+$ classical monocytes (Figure S4e).   Compared to the

control group, *S100A8*, *S100A9*, and *S100A12* were the top 3 upregulated genes in classical monocytes of

WD patients (Figure S4f).   *S100A8* and *S100A9* are confirmed to play vital roles in modulating the

inflammatory responses by recruiting leukocytes and inducing cytokine secretion during inflammation, and

as an alarmin, their elevation indicates inflammation in the local environment[89,90].   GO analysis of DEGs

in monocytes showed a decrease in biological processes involving cytokine production, response to

lipopolysaccharide and bacteria, as well as lymphocyte activation (Figure 4e).

For T cells, we performed GO analysis on DEGs in CD8 T cells between two groups.   Notably, the

biological processes linked to energy and lipid metabolism were upregulated in several subtypes, including

CD4Tmem_CD40LG, CD4Treg_CTLA4, CD8MAIT_KLRB1, CD8MAIT_SLC4A10, CD8Teff_GZMH,

and CD8Teff_GZMK.   In addition, biological processes associated with response to copper ion were

upregulated in the CD8Teff_MT1X subtype (Figure S4g).   Meanwhile, T cell subtypes showed

comparable cytotoxicity scores, and there were slight changes between two groups (Figure 4f).   However,

CD8 T cells in the case group exhibited lower levels of exhaustion, except for CD8Teff_GZMH and

CD8Teff_ISG15 (Figure 4f).   It was also discovered that while the proportion of CD4Treg_CTAL4 was

increased in the case group, the Treg signature was slightly lower (Figure 4g).   Therefore, these results

suggest that WD patients exhibit variations in immune cell function and certain biological processes.

NK cell exhaustion in WD patients

The KEGG analysis of DEGs in NK cells between the case and control groups revealed down-regulation of

the natural killer cell-mediated cytotoxicity, along with the NF-kappa B signaling pathway and chemokine

signaling pathway, in the case group (Figure S5a).   Interestingly, GO analysis showed upregulated

biological processes of oxidative phosphorylation and ATP metabolic process, implying the altered

metabolic program in NK cells and immune microenvironment (Figure S5b).   Since the primary effect of

NK cells is cytotoxicity on target cells, we compared the gene expression levels of activating receptors,

inhibitory receptors, and effector molecules between two groups. The results showed increased

expression in *KLRC1* and *TIGIT*, and decreased expression in *FCGR3A*, *TNF*, *IFNG*, *GZMB*, *GNLY*, and

CCL4 (Figure 5a). KLRC1 and TIGIT are inhibitory receptors expressed on NK cell surface, and their

activation inhibits NK cell functions[21,91]. It was collectively indicated that WD patients have NK cells

with exhausted status, which results in poor effector function.

In addition, trajectory analysis showed that NK cells in two groups exhibited a distinct differentiation

trajectory. Most of the NK cells in the control group differentiated towards cell fate 1, while those in the

case group displayed a higher tendency for differentiation towards cell fate 2, with cell grouping occurring

at the final stage (Figure 5b). To ascertain the gene expression profile and the respective cell statuses of

the two outcomes, we conducted pseudotime analysis of gene expression in two differentiating directions.

The results showed that the gene expression of *KLRC1* (included in gene cluster 1) was gradually increased

along the way to cell fate 2, while the gene expression of *FCGR3A*, *TNF*, and *GZMB* (included in gene

cluster 3) was weakened in the same direction, suggesting that cell fate 2 is a more exhausted state of NK

cells (Figure S5c). Consequently, these results demonstrate that NK cells differentiate towards an

exhausted status in WD patients. The development, maturation, and function of NK cells are strictly

regulated by transcription factors (TFs)[92]. Next, to assess the transcription factors (TFs) underlying

differences in NK cells between WD patients and control groups, we applied PYSCENIC analysis in NK

cells. We found that the gene expression of T-bet (encoded by *TBX21*) was downregulated and Eomes

(encoded by *EOMES*) was upregulated in the case group (Figure S5d). These two constitute a pair of TFs

that play significantly complementary roles in directing the transcriptional program of NK cells.

Consistent with previous studies, T-bet expression was downregulated in exhausted NK cells, whereas

Eomes expression was increased[24,93,94]. Thus, it was suggested that the downregulation of *TBX21* and

upregulation of *EOMES* contributed to the exhaustion of NK cells in WD patients.

To validate NK cell exhaustion in WD patients, we performed multiplex immunohistochemistry (mIHC)

staining on liver tissue sections. The results indicate that NKG2A$^+$ cells were more frequently observed in

CD56$^+$ NK cells in WD patients (Figure 5c). Moreover, flow cytometry also confirmed a significant

increase in the percentage of NKG2A$^+$ and TIGIT$^+$ NK cells and a significant decrease in the percentage of

IFNγ$^+$ NK cells in peripheral blood lymphocytes from WD patients compared to control groups (Figure 5d,

e).   Meanwhile, the proportion of NK cells in lymphocytes from peripheral blood and the proportion of

CD56^{Bright} NK cells in total NK cells were also calculated.   The significantly higher proportion of NK cells

in WD patients was consistent with the scRNA sequencing data, and more CD56^{Bright} NK cells suggested a

relatively weaker effector function in WD patients (Figure S5e).   Therefore, these results indicated that,

despite the increased proportion and the upregulated oxidative phosphorylation, NK cell exhibited

exhausted status in WD patients.

NK cell exhaustion predicts poor prognosis of cholecystitis

Cholecystitis patients with WD exhibit poorer clinical outcomes.   There is evidence demonstrating that

NK cell exhaustion is correlated with poor liver cancer prognosis[24,26,95].   Thus, we hypothesize that NK

cell exhaustion also contributes to the worse prognosis of cholecystitis patients with WD.

Prognostic models for inflammatory related diseases in GEO were developed using machine learning

methods, including random survival forest, ROC curve, and Kaplan-Meier survival curves.   A logistic

regression algorithm was used to construct a multigene prediction model based on GSE166915

(cholecystitis) and other inflammatory diseases, namely GSE91035 (pancreatitis), GSE75037 (lung

adenocarcinoma), and GSE41804 (hepatocellular carcinoma).   Using stepwise regression analysis, 12

genes, *KLRC1*, *TIGIT*, *HAVCR2*, *PDCD1*, *CD96*, *LAG3*, *FCGR3A*, *NCR3*, *NCR2*, *NCR1*, *KLRK1*, and

PRF1, were selected to obtain the best model.   The results showed that the predictive model constructed

from these genes had good prognostic performance, with AUC of 0.77, 0.80, 0.97, 0.90 (Figure S6a-d).

The AUCs of the models in GSE75037 and GSE41084 were relatively high, 0.97 and 0.90, respectively.

Univariate Cox regression analysis revealed that the *TIGIT* (in GSE166915) and *PDCD1* (in GSE75037)

were risk genes with hazard ratio (HR) > 1, while *NCR2* (in GSE91035), *HAVCR2*, *KLRC1*, *NCR3* (in

GSE75037), *NCR3*, *LAG3* (in GSE41804) were protective gene with HR < 1 among 12 differently

expressed markers of NK cell exhaustion (Figure S6a-d).   Consistently, the Kaplan-Meier survival curves

suggested a shorter survival time in the group with high markers of NK cell exhaustion than in the group

with low markers in the GSE166915 (*P*= 6.1e-6), GSE91035 (*P*= 1.9e-5), GSE75037 (*P*= 3.8e-16), and

GSE41804 (*P*= 3.0e-3) datasets (Figure 6a-d). Patients with cholecystitis, pancreatitis, lung

adenocarcinoma, and hepatocellular carcinoma who show high markers of NK cell exhaustion are

associated with poor survival compared to those with low markers of NK cell exhaustion.

Discussion

Immune cell dysfunction impacts the development, progression, and prognosis of diseases and serves as the

basis for immunotherapy. The current understanding of immune cell dysfunction in inflammatory

diseases is limited, which hinders the development of effective therapeutic strategies. Our retrospective

clinical analysis indicates that WD contributes to the unfavorable prognosis of cholecystitis. We

developed an atlas of immune cell dysfunction linked to specific metabolic disruption in WD patients,

including 19 major cell types. The immune microenvironment change primarily encompasses increased

proportion and strengthened function in innate immunocytes, including upregulated antigen processing and

presentation and activation of immune responses. Furthermore, we conducted a comprehensive

assessment and comparison of immune cell function in WD and control group. This includes an

evaluation of pro-inflammatory and immune regulatory function in macrophages and KCs, phagocytosis

and chemokine activity in neutrophils, and cytotoxicity in T cells, etc. We identified NK cell exhaustion

in the WD group, with upregulation of *NKG2A* and *TIGIT* and downregulation of *CD16*, *TNF*, *IFNG*,

GZMB, *GNLY*, and *CCL4*. The validation of our findings in patient liver tissues and blood samples was

conducted using immunohistochemistry assays and flow cytometry. Our findings indicated that the NK

cell exhaustion signature has the ability to predict and prognosticate inflammatory diseases such as

cholecystitis, pancreatitis, lung adenocarcinoma, and hepatocellular carcinoma. These results unveil novel

perspectives on the immune cell dysfunction role in inflammatory diseases.

WD represents a model of metabolic abnormality in hepatocytes caused by *ATP7B* mutation. Therefore,

immune cell dysfunction was induced by the extracellular signals within the reshaped tissue

microenvironment. Emerging research indicates that metabolic crosstalk in the tumor microenvironment

disrupts metabolic and signaling pathways in immune cells, leading to dysfunction[96-100]. Targeting

metabolic pathways to enhance the efficacy of immunotherapy is a promising approach. Our study found

multiple metabolic abnormalities in immune cells, including the widely upregulated ATP metabolic process

in T and NK cells. Interestingly, glycolysis and oxidative phosphorylation contribute to maintain the

cytotoxic ability and promote expansion of NK cells[101-103]. However, high rate of oxidative

phosphorylation, increased proportion, and exhausted status were simultaneously observed in NK cells of

WD patients. Further studies are needed to elucidate the metabolic landscape, mechanisms underlying

metabolic alterations, and their role in immune cell dysfunction. Defective ATP7B leads to excessive

copper accumulation in the system's circulation. A recent study showed that excess copper induces cell

death in a manner dependent on mitochondrial respiration[104]. Considering the impact on mitochondrial

respiration processes in immune cells, it is worth investigating the role of copper in the physiology of

immune cells. Further research is expected to establish a foundation for improving immunotherapies

through metabolic or ionophore interventions.

Methods

Patients and clinical samples

All the clinical data were obtained from The First Affiliated Hospital of Anhui University of Chinese

Medicine. The whole blood samples and liver tissue samples were collected from patients at The First

Affiliated Hospital of Anhui University of Chinese Medicine. The study was approved by the Ethics

Committee of the First Affiliated Hospital of Anhui University of Traditional Chinese Medicine and the Ethics

Committee of the First Affiliated Hospital of USTC.

Sanger sequencing

The sanger sequencing on 20 WD patients were conducted by Sangon Biotech (Shanghai) Co., Ltd.

The primers to detect mutations are:

R778L-F, 5'-AGCCTTCACTGTCCTTGTCTTTC-3';

R778L-R, 5'-ATTAGCTGGGATTTCAGAAGTAGTG-3';

T935M-F, 5'-ATGCTTGTGGTGTTTTATTTCTTC-3';

T935M-R, 5'-AGAAGCAAGCAAATAAAATGTAATG-3';

P992L-F, 5'-TCTCAACCTGCCTCTGACTCTG-3';

P992L-R, 5'-TGGTGGCTACTCTGTTGCTACTG-3'.

Cell culture, gene knockout, and metabolic assays

WRL 68 cells were cultured using DMEM high-glucose media supplemented with 10% FBS at 37 °C with

5% CO_2.   The lentivirus-mediated CRISPR-Cas9 KO vector targeting human *ATP7B* was obtained from

OBiO Technology (Shanghai) Co., Ltd.   Cells at 30% to 40% confluency were incubated in a medium

containing optimal dilutions of lentivirus, then subjected to puromycin selection to obtain the stable

transfected cells.   OCRs and ECARs were measured using Seahorse XF Cell Mito Stress Test Kit

(103015-100, Agilent) and Seahorse XF Glycolysis Stress Test Kit (103020-100, Agilent) according to

manuals in Agilent XFe96 Analyzer, respectively.

Single-cell RNA sequencing

In the cell preparation section: For the quality check and counting of single cell suspension, the cell

survival rate is generally above 80%.   The cells that have passed the test are washed and resuspended to

prepare a suitable cell concentration of 700~1,200 cells/μL for 10x Genomics ChromiumTM.   The system

is operated on the machine.

Steps in GEM Creation & Thermal cycling: GEMs (Gel Bead in Emulsion) were constructed for single cell

separation according to the number of cells to be harvested.   After GEMs were normally formed, GEMs

were collected for reverse transcription in a PCR machine for labeling.

Post Cycling Cleanup & cDNA Amplication: The GEMs were oil-treated, and the amplified cDNA was

purified by magnetic beads, and then subjected to cDNA amplification and quality inspection.

Library Preparation & Quantification: The 3'Gene Expression Library was constructed with the

quality-qualified cDNA.   After fragmentation, adaptor ligation, sample index PCR, etc., the library is

finally quantitatively examined.

Sequencing: The final library pool was sequenced on the Illumina Hiseq instrument using 150-base-pair

paired-end reads.

Cell type annotation and differentially expressed genes (DEGs) analysis

The cell type identity of each cluster was determined with the expression of canonical markers found in the

DEGs using SynEcoSys database. The markers and references are shown in Supplementary Table S4.

To identify DEGs, we used the Seurat FindMarkers function based on Wilcox likelihood-ratio test with

default parameters, and selected the genes expressed in more than 10% of the cells in a cluster and with an

average log (Fold Change) value greater than 0.25 as DEGs. For the cell type annotation of each cluster,

we combined the expression of canonical markers found in the DEGs with knowledge from literatures, and

displayed the expression of markers of each cell type with heatmaps that were generated with Seurat

Heatmap function. Doublet cells were identified as expressing markers for different cell types, and

removed manually.

Pathway enrichment analysis

To investigate the potential functions of DEGs, the Gene Ontology (GO) and Kyoto Encyclopedia of Genes

and Genomes (KEGG) analysis were used with the "clusterProfiler" R package 3.16.1[105]. Pathways with

p_adj value less than 0.05 were considered as significantly enriched. Gene Ontology gene sets including

molecular function (MF), biological process (BP), and cellular component (CC) categories were used as

reference.

Cell-cell Interaction Analysis

CellChat (version 0.0.2) was used to analyze the intercellular communication networks from scRNA-seq

data[106]. A CellChat object was created using the R package process. Cell information was added into the

meta slot of the object. The ligand-receptor interaction database was set, and the matching receptor

inference calculation was performed.

Trajectory analysis

Cell differentiation trajectory was reconstructed with Monocle2[107]. Highly-variable genes (HVGs) were

used to sort cells in order of spatial-temporal differentiation. We used DDRTree to perform

FindVairableFeatures and dimension-reduction. Finally, the trajectory was visualized by

plot_cell_trajectory function. Next, CytoTRACE (a computational method that predicts the differentiation

state of cells from single-cell RNA-sequencing data using gene Counts and Expression) was used to predict

the differentiation potential[108].

Transcription factor regulatory network analysis

Transcription factor network was constructed by pyscenic v0.11.0 using scRNA expression matrix and

transcription factors in AnimalTFDB[109]. First, GRNBoost2 predicted a regulatory network based on the

co-expression of regulators and targets. CisTarget was then applied to exclude indirect targets and to

search transcription factor binding motifs. After that, AUCell was used for regulon activity quantification

for every cell.

UCell gene set scoring

Gene set scoring was performed using the R package UCell v 1.1.0[110]. UCell scores are based on the

Mann-Whitney U statistic by ranking query genes in order of their expression levels in individual cells.

Because UCell is a rank-based scoring method, it is suitable to be used in large datasets containing multiple

samples and batches. The gene sets for signature scoring are listed in Supplementary Table S3.

Multiplex immunohistochemistry

The procedure of deparaffinization, dehydration, antigen retrieval, endogenous peroxidases quench, block,

antibody incubation was conducted as standard immunohistochemistry. Solution of tyramide signal

amplification (TSA) (WAS1101010, BioMed World) was applied on the slides and antibodies were

removed by microwave heating. Repeat block, antibody incubation, TSA, and microwave heating to

perform next cycle of staining. DAPI (WAS1301050, BioMed World) was used to stain nuclei and

antifade mountant (WAS1302050, BioMed World) was applied. The primary antibody to CD56 (ab75813,

Abcam) and NKG2A (ab260035, Abcam) and second antibody (WAS1201100, BioMed World) were used.

Image of immunofluorescence-stained slides were acquired with Nikon Eclipse Ci-L and 3DHISTECH

Pannoramic MIDI II.

Flow cytometry

Lymphocytes were isolated from the whole blood using density gradient centrifugal method (P8610,

Solarbio). Isolated cells were blocked (564219, BD Pharmingen) and stained with antibodies against CD3

(560835, BD Pharmingen), CD56 (555518, BD Pharmingen), NKG2A (375114, Biolegend), and TIGIT

(568672, BD Pharmingen). For intracellular cytokine staining, cells were stimulated for 4 h with

Leukocyte Activation Cocktail and BD GolgiPlug (550583, BD Pharmingen) according to manuals. After

stimulation, cells were blocked, stained for surface markers, fixed and permeabilized with

Fixation/Permeablization Kit (554714, BD Pharmingen). Fixed cells were stained with antibodies against

IFNγ (552887, BD Pharmingen).

Construction and verification of prognostic model

On the basis of the markers of NK cell exhaustion, univariate Cox analysis was conducted in cholecystitis,

pancreatitis, lung adenocarcinoma, and hepatocellular carcinoma patients to screen survival-related markers

of NK cell exhaustion, and markers with P values < 0.05 were retained. We conducted multivariate

analysis to identify the optimal prognostic markers for the prognostic model. The risk scores of the

patients were calculated based on the normalized gene expression levels and the Cox regression coefficients

of selected markers. Time-dependent receiver operating characteristic (ROC) curves were utilized to

verify the prognostic performance of the model for overall survival (OS). For Kaplan-Meier survival, we

used the R software package maxstat (Maximally selected rank statistics with multiple p-value

approximations version: 0.7-25) to calculate the optimal cutoff value for RiskScore. We set the minimum

group sample size to be greater than 25% and the maximum group sample size to be less than 75%,

ultimately obtaining the optimal cutoff value. Based on this, patients were divided into high and low

groups, and further analyzed the prognostic differences between the two groups using the R software

package's survival function.   The logrank test method was used to evaluate the significance of prognostic

differences between different groups of samples, and significant prognostic differences were ultimately

observed.

Statistical analysis

Statistical analyses were performed using appropriate tests as indicated in legends (unpaired two-tailed

t-test, chi-square test, and Wilcox test).

Data availability

sc-RNA sequening data that support the findings of this study have been deposited in Gene Expression

Omnibus (GEO) with the accession numbers GSE254082.

Author contributions

Y.W., Q.F., H.W., and Q.Y. conceived the concept for this study.   Y.J., J.X., L.J., and M.H. analyzed data

and implemented experiments.   C.D. conducted the bioinformatic analysis.   L.J., Q.T., and Z.L. collected

the clinical data and clinical samples.   Y.J., J.X., and C.D. wrote the manuscript.   W.Z. and Q. F.

proofread the manuscript.

Fundings

This work was supported by grants from National Natural Science foundation of China (Project:

2021KY340; NSFC: 81773112 and 82073186 for Q.F.).   This work was supported by Construction project

of the State Administration of Traditional Chinese Medicine in 2021 (Teaching Letter [2021] No. 270 for

Q.Y.). and Anhui Province Clinical Key Specialty Project in 2022 (Anhui Weichuan [2022] No. 297 for

Q.Y.).

Competing interests

The authors declare no competing interests.

References

References

1   Elwood, D. R. Cholecystitis. *Surg Clin North Am* **88**, 1241-1252, viii (2008).

https://doi.org:10.1016/j.suc.2008.07.008

2   Huffman, J. L. & Schenker, S. Acute acalculous cholecystitis: a review. *Clin Gastroenterol Hepatol* **8**,

15-22 (2010). https://doi.org:10.1016/j.cgh.2009.08.034

3   Markaki, I., Konsoula, A., Markaki, L., Spernovasilis, N. & Papadakis, M. Acute acalculous

cholecystitis due to infectious causes. *World J Clin Cases* **9**, 6674-6685 (2021).

https://doi.org:10.12998/wjcc.v9.i23.6674

4   Laurila, J. J. *et al.* Histopathology of acute acalculous cholecystitis in critically ill patients.

Histopathology **47**, 485-492 (2005).

https://doi.org:https://doi.org/10.1111/j.1365-2559.2005.02238.x

5   Fu, Y. T., Pang, L. W., Dai, W. L., Wu, S. D. & Kong, J. Advances in the Study of Acute Acalculous

Cholecystitis: A Comprehensive Review. *Digest Dis* **40**, 468-478 (2022).

https://doi.org:10.1159/000520025

6   Abu-Sbeih, H. *et al.* Case series of cancer patients who developed cholecystitis related to immune

checkpoint inhibitor treatment. *J Immunother Cancer* **7** (2019). https://doi.org:ARTN 118

10.1186/s40425-019-0604-2

7   Loozen, C. S., Oor, J. E., van Ramshorst, B., van Santvoort, H. C. & Boerma, D. Conservative

treatment of acute cholecystitis: a systematic review and pooled analysis. *Surg Endosc* **31**, 504-515

(2017). https://doi.org:10.1007/s00464-016-5011-x

8   van Dijk, A. H. *et al.* Systematic review of antibiotic treatment for acute calculous cholecystitis. *Brit J*

Surg **103**, 797-811 (2016). https://doi.org:10.1002/bjs.10146

9   Buonacera, A., Stancanelli, B., Colaci, M. & Malatino, L. Neutrophil to Lymphocyte Ratio: An

Emerging Marker of the Relationships between the Immune System and Diseases. *Int J Mol Sci* **23**

(2022). https://doi.org:10.3390/ijms23073636

10   McKinney, E. F. *et al.* A CD8+ T cell transcription signature predicts prognosis in autoimmune

disease. *Nat Med* **16**, 586-591, 581p following 591 (2010). https://doi.org:10.1038/nm.2130

11   McKinney, E. F., Lee, J. C., Jayne, D. R., Lyons, P. A. & Smith, K. G. T-cell exhaustion,

co-stimulation and clinical outcome in autoimmunity and infection. *Nature* **523**, 612-616 (2015).

https://doi.org:10.1038/nature14468

12   Sun, Y. *et al.* Single-cell landscape of the ecosystem in early-relapse hepatocellular carcinoma. *Cell*

184, 404-421 e416 (2021). https://doi.org:10.1016/j.cell.2020.11.041

13   Fridman, W. H., Zitvogel, L., Sautes-Fridman, C. & Kroemer, G. The immune contexture in cancer

prognosis   and   treatment.   *Nat   Rev   Clin   Oncol*   **14**,   717-734   (2017).

https://doi.org:10.1038/nrclinonc.2017.101

14   Wouters, M. C. A. & Nelson, B. H. Prognostic Significance of Tumor-Infiltrating B Cells and Plasma

Cells   in   Human   Cancer.   *Clin   Cancer   Res*   **24**,   6125-6135   (2018).

https://doi.org:10.1158/1078-0432.CCR-18-1481

15   Abel, A. M., Yang, C., Thakar, M. S. & Malarkannan, S. Natural Killer Cells: Development,

Maturation,   and   Clinical   Utilization.   *Front   Immunol*   **9**,   1869   (2018).

https://doi.org:10.3389/fimmu.2018.01869

16   Crinier, A., Narni-Mancinelli, E., Ugolini, S. & Vivier, E. SnapShot: Natural Killer Cells. *Cell* **180**,

1280-1280 e1281 (2020). https://doi.org:10.1016/j.cell.2020.02.029

17   Highton, A. J., Schuster, I. S., Degli-Esposti, M. A. & Altfeld, M. The role of natural killer cells in liver

inflammation.   *Semin*   *Immunopathol*   **43**,   519-533   (2021).

https://doi.org:10.1007/s00281-021-00877-6

18   Parisi, L. *et al.* Natural Killer Cells in the Orchestration of Chronic Inflammatory Diseases. *J Immunol*

Res **2017**, 4218254 (2017). https://doi.org:10.1155/2017/4218254

19   Earls, R. H. & Lee, J. K. The role of natural killer cells in Parkinson's disease. *Exp Mol Med* **52**,

1517-1525 (2020). https://doi.org:10.1038/s12276-020-00505-7

20   Bi,   J.   &   Tian,   Z.   NK   Cell   Exhaustion.   *Front*   *Immunol*   **8**,   760   (2017).

https://doi.org:10.3389/fimmu.2017.00760

21   Zhang, Q. *et al.* Blockade of the checkpoint receptor TIGIT prevents NK cell exhaustion and elicits

potent   anti-tumor   immunity.   *Nat*   *Immunol*   **19**,   723-732   (2018).

https://doi.org:10.1038/s41590-018-0132-0

22   Liu, X. *et al.* Immune checkpoint HLA-E:CD94-NKG2A mediates evasion of circulating tumor cells

from   NK   cell   surveillance.   *Cancer*   *Cell*   **41**,   272-287   e279   (2023).

https://doi.org:10.1016/j.ccell.2023.01.001

23   Zhang, C. *et al.* NKG2A is a NK cell exhaustion checkpoint for HCV persistence. *Nat Commun* **10**,

1507 (2019). https://doi.org:10.1038/s41467-019-09212-y

24   Sun, H. *et al.* Human CD96 Correlates to Natural Killer Cell Exhaustion and Predicts the Prognosis

568   of   Human   Hepatocellular   Carcinoma.   *Hepatology*   **70**,   168-183   (2019).

569   https://doi.org:10.1002/hep.30347

570   25   Liu, G. *et al.* Increased TIGIT expressing NK cells with dysfunctional phenotype in AML patients

571   correlated   with   poor   prognosis.   *Cancer   Immunol   Immunother*   **71**,   277-287   (2022).

572   https://doi.org:10.1007/s00262-021-02978-5

573   26   Sun, C. *et al.* High NKG2A expression contributes to NK cell exhaustion and predicts a poor

574   prognosis   of   patients   with   liver   cancer.   *Oncoimmunology*   **6**,   e1264562   (2017).

575   https://doi.org:10.1080/2162402X.2016.1264562

576   27   Crispe, I. N. The liver as a lymphoid organ. *Annu Rev Immunol* **27**, 147-163 (2009).

577   https://doi.org:10.1146/annurev.immunol.021908.132629

578   28   Kubes, P. & Jenne, C. Immune Responses in the Liver. *Annu Rev Immunol* **36**, 247-277 (2018).

579   https://doi.org:10.1146/annurev-immunol-051116-052415

580   29   Racanelli, V. & Rehermann, B. The liver as an immunological organ. *Hepatology* **43**, S54-62 (2006).

581   https://doi.org:10.1002/hep.21060

582   30   Hardy, T., Oakley, F., Anstee, Q. M. & Day, C. P. Nonalcoholic Fatty Liver Disease: Pathogenesis

583   and   Disease   Spectrum.   *Annu   Rev   Pathol*   **11**,   451-496   (2016).

584   https://doi.org:10.1146/annurev-pathol-012615-044224

585   31   Armstrong, M. J., Adams, L. A., Canbay, A. & Syn, W. K. Extrahepatic complications of nonalcoholic

586   fatty liver disease. *Hepatology* **59**, 1174-1197 (2014). https://doi.org:10.1002/hep.26717

32   Matyas, C., Haskó, G., Liaudet, L., Trojnar, E. & Pacher, P. Interplay of cardiovascular mediators,

oxidative stress and inflammation in liver disease and its complications. *Nature Reviews Cardiology*

18, 117-135 (2021). https://doi.org:10.1038/s41569-020-0433-5

33   Knab, L. M., Boller, A. M. & Mahvi, D. M. Cholecystitis. *Surg Clin North Am* **94**, 455-470 (2014).

https://doi.org:10.1016/j.suc.2014.01.005

34   Bandmann, O., Weiss, K. H. & Kaler, S. G. Wilson's disease and other neurological copper

disorders. *Lancet Neurol* **14**, 103-113 (2015). https://doi.org:10.1016/S1474-4422(14)70190-5

35   Huster, D. *et al.* Consequences of copper accumulation in the livers of the Atp7b-/- (Wilson disease

gene) knockout mice. *Am J Pathol* **168**, 423-434 (2006). https://doi.org:10.2353/ajpath.2006.050312

36   Medici, V. *et al.* Wilson's disease: changes in methionine metabolism and inflammation affect global

DNA methylation in early liver disease. *Hepatology* **57**, 555-565 (2013).

https://doi.org:10.1002/hep.26047

37   Huster, D. Structural and metabolic changes in Atp7b-/- mouse liver and potential for new

interventions in Wilson's disease. *Ann N Y Acad Sci* **1315**, 37-44 (2014).

https://doi.org:10.1111/nyas.12337

38   Aggarwal, A. & Bhatt, M. Wilson disease. *Curr Opin Neurol* **33**, 534-542 (2020).

https://doi.org:10.1097/WCO.0000000000000837

39   Cheng, N. *et al.* Spectrum of ATP7B mutations and genotype-phenotype correlation in large-scale

Chinese patients with Wilson Disease. *Clin Genet* **92**, 69-79 (2017).

https://doi.org:10.1111/cge.12951

40   Yang, G. M. *et al.* Prevalent Pathogenic Variants of ATP7B in Chinese Patients with Wilson's

Disease: Geographical Distribution and Founder Effect. *Genes (Basel)* **12** (2021).

https://doi.org:10.3390/genes12030336

41   Członkowska, A. *et al.* Wilson disease. *Nat Rev Dis Primers* **4**, 21 (2018).

https://doi.org:10.1038/s41572-018-0018-3

42   Roberts, E. A., Robinson, B. H. & Yang, S. Mitochondrial structure and function in the untreated

Jackson toxic milk (tx-j) mouse, a model for Wilson disease. *Mol Genet Metab* **93**, 54-65 (2008).

https://doi.org:10.1016/j.ymgme.2007.08.127

43   Zischka, H. & Lichtmannegger, J. Pathological mitochondrial copper overload in livers of Wilson's

disease patients and related animal models. *Ann N Y Acad Sci* **1315** (2014).

https://doi.org:10.1111/nyas.12347

44   Muchenditsi, A. *et al.* Systemic deletion of Atp7b modifies the hepatocytes' response to copper

overload in the mouse models of Wilson disease. *Sci Rep* **11**, 5659 (2021).

https://doi.org:10.1038/s41598-021-84894-3

45   He, S. *et al.* Single-cell transcriptome profiling of an adult human cell atlas of 15 major organs.

Genome Biol **21**, 294 (2020). https://doi.org:10.1186/s13059-020-02210-0

46   Sun, Y. *et al.* Single-cell landscape of the ecosystem in early-relapse hepatocellular carcinoma. *Cell*

184 (2021). https://doi.org:10.1016/j.cell.2020.11.041

47   Li, Q. *et al.* Single-cell transcriptome profiling reveals vascular endothelial cell heterogeneity in

human skin. *Theranostics* **11**, 6461-6476 (2021). https://doi.org:10.7150/thno.54917

48   Sathe, A. *et al.* Single-Cell Genomic Characterization Reveals the Cellular Reprogramming of the

Gastric Tumor Microenvironment. *Clinical Cancer Research : an Official Journal of the American*

Association   For   Cancer   Research **26**,   2640-2653   (2020).

https://doi.org:10.1158/1078-0432.CCR-19-3231

49   Wang, L. *et al.* Single-cell reconstruction of the adult human heart during heart failure and recovery

reveals the cellular landscape underlying cardiac function. *Nat Cell Biol* **22**, 108-119 (2020).

https://doi.org:10.1038/s41556-019-0446-7

50   Bhattacharjee, S. *et al.* Tumor restriction by type I collagen opposes tumor-promoting effects of

cancer-associated fibroblasts. *J Clin Invest* **131** (2021). https://doi.org:10.1172/JCI146987

51   Dumas, S. J. *et al.* Single-Cell RNA Sequencing Reveals Renal Endothelium Heterogeneity and

Metabolic   Adaptation   to   Water   Deprivation. *J   Am   Soc   Nephrol* **31**,   118-138   (2020).

https://doi.org:10.1681/ASN.2019080832

52   Durante, M. A. *et al.* Single-cell analysis reveals new evolutionary complexity in uveal melanoma.

Nature Communications **11**, 496 (2020). https://doi.org:10.1038/s41467-019-14256-1

53   Gong, L. *et al.* Comprehensive single-cell sequencing reveals the stromal dynamics and

tumor-specific characteristics in the microenvironment of nasopharyngeal carcinoma. *Nature*

Communications **12**, 1540 (2021). https://doi.org:10.1038/s41467-021-21795-z

54   Wang, X. *et al.* Comparative analysis of cell lineage differentiation during hepatogenesis in humans

and   mice   at   the   single-cell   transcriptome   level. *Cell   Res* **30**,   1109-1126   (2020).

https://doi.org:10.1038/s41422-020-0378-6

55   Zeng, Y. *et al.* Single-Cell RNA Sequencing Resolves Spatiotemporal Development of Pre-thymic

Lymphoid Progenitors and Thymus Organogenesis in Human Embryos. *Immunity* **51** (2019).

https://doi.org:10.1016/j.immuni.2019.09.008

56   Wu, F. *et al.* Single-cell profiling of tumor heterogeneity and the microenvironment in advanced

non-small   cell   lung   cancer.   *Nature   Communications*   **12**,   2540   (2021).

https://doi.org:10.1038/s41467-021-22801-0

57   Ramachandran, P. *et al.* Resolving the fibrotic niche of human liver cirrhosis at single-cell level.

Nature **575**, 512-518 (2019). https://doi.org:10.1038/s41586-019-1631-3

58   Pombo Antunes, A. R. *et al.* Single-cell profiling of myeloid cells in glioblastoma across species and

disease stage reveals macrophage competition and specialization. *Nat Neurosci* **24**, 595-610

(2021). https://doi.org:10.1038/s41593-020-00789-y

59   Strauss, O., Dunbar, P. R., Bartlett, A. & Phillips, A. The immunophenotype of antigen presenting

cells of the mononuclear phagocyte system in normal human liver--a systematic review. *J Hepatol*

62, 458-468 (2015). https://doi.org:10.1016/j.jhep.2014.10.006

60   Singh, M. *et al.* High-throughput targeted long-read single cell sequencing reveals the clonal and

transcriptional   landscape   of   lymphocytes.   *Nature   Communications*   **10**,   3120   (2019).

https://doi.org:10.1038/s41467-019-11049-4

61   Halbritter, F. *et al.* Epigenomics and Single-Cell Sequencing Define a Developmental Hierarchy in

Langerhans   Cell   Histiocytosis.   *Cancer   Discov*   **9**,   1406-1421   (2019).

https://doi.org:10.1158/2159-8290.CD-19-0138

62 Bassez, A. *et al.* A single-cell map of intratumoral changes during anti-PD1 treatment of patients

with breast cancer. *Nature Medicine* **27**, 820-832 (2021).

https://doi.org:10.1038/s41591-021-01323-8

63 Popescu, D.-M. *et al.* Decoding human fetal liver haematopoiesis. *Nature* **574**, 365-371 (2019).

https://doi.org:10.1038/s41586-019-1652-y

64 Chen, Y.-P. *et al.* Single-cell transcriptomics reveals regulators underlying immune cell diversity and

immune subtypes associated with prognosis in nasopharyngeal carcinoma. *Cell Res* **30**, 1024-1042

(2020). https://doi.org:10.1038/s41422-020-0374-x

65 Wang, J. *et al.* Liver Immune Profiling Reveals Pathogenesis and Therapeutics for Biliary Atresia.

Cell **183** (2020). https://doi.org:10.1016/j.cell.2020.10.048

66 Dixon, L. J., Barnes, M., Tang, H., Pritchard, M. T. & Nagy, L. E. Kupffer cells in the liver. *Compr*

Physiol **3**, 785-797 (2013). https://doi.org:10.1002/cphy.c120026

67 Cella, M. *et al.* Subsets of ILC3-ILC1-like cells generate a diversity spectrum of innate lymphoid

cells in human mucosal tissues. *Nature Immunology* **20**, 980-991 (2019).

https://doi.org:10.1038/s41590-019-0425-y

68 Elmentaite, R. *et al.* Cells of the human intestinal tract mapped across space and time. *Nature* **597**,

250-255 (2021). https://doi.org:10.1038/s41586-021-03852-1

69 Liu, Y. *et al.* Single-cell analysis reveals immune landscape in kidneys of patients with chronic

transplant rejection. *Theranostics* **10**, 8851-8862 (2020). https://doi.org:10.7150/thno.48201

70 Wang, L. *et al.* Nanoparticle enhanced combination therapy for stem-like progenitors defined by

single-cell transcriptomics in chemotherapy-resistant osteosarcoma. *Signal Transduction and*

Targeted Therapy **5**, 196 (2020). https://doi.org:10.1038/s41392-020-00248-x

71   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by single-cell

RNA-seq. *Science (New York, N.Y.)* **352**, 189-196 (2016). https://doi.org:10.1126/science.aad0501

72   Jaeger, N. *et al.* Single-cell analyses of Crohn's disease tissues reveal intestinal intraepithelial T

cells heterogeneity and altered subset distributions. *Nature Communications* **12**, 1921 (2021).

https://doi.org:10.1038/s41467-021-22164-6

73   Depuydt, M. A. C. *et al.* Microanatomy of the Human Atherosclerotic Plaque by Single-Cell

Transcriptomics. *Circ Res* **127**, 1437-1455 (2020).

https://doi.org:10.1161/CIRCRESAHA.120.316770

74   Andor, N. *et al.* Single-cell RNA-Seq of follicular lymphoma reveals malignant B-cell types and

coexpression of T-cell immune checkpoints. *Blood* **133**, 1119-1129 (2019).

https://doi.org:10.1182/blood-2018-08-862292

75   Heming, M. *et al.* Neurological Manifestations of COVID-19 Feature T Cell Exhaustion and

Dedifferentiated Monocytes in Cerebrospinal Fluid. *Immunity* **54** (2021).

https://doi.org:10.1016/j.immuni.2020.12.011

76   Xin, Z. *et al.* The immune landscape of human thymic epithelial tumors. *Nature Communications* **13**,

5463 (2022). https://doi.org:10.1038/s41467-022-33170-7

77   Zhao, J. *et al.* Single-cell RNA sequencing reveals the heterogeneity of liver-resident immune cells

in human. *Cell Discovery* **6**, 22 (2020). https://doi.org:10.1038/s41421-020-0157-z

78   Williams, D. W. *et al.* Human oral mucosa cell atlas reveals a stromal-neutrophil axis regulating

tissue immunity. *Cell* **184** (2021). https://doi.org:10.1016/j.cell.2021.05.013

79   Madissoon, E. *et al.* scRNA-seq assessment of the human lung, spleen, and esophagus tissue

stability after cold preservation. *Genome Biol* **21**, 1 (2019).

https://doi.org:10.1186/s13059-019-1906-x

80   Zhang, F. *et al.* Adaptive immune responses to SARS-CoV-2 infection in severe versus mild

individuals. *Signal Transduction and Targeted Therapy* **5**, 156 (2020).

https://doi.org:10.1038/s41392-020-00263-y

81   Fernandez, D. M. *et al.* Single-cell immune landscape of human atherosclerotic plaques. *Nature*

Medicine **25**, 1576-1588 (2019). https://doi.org:10.1038/s41591-019-0590-4

82   Luoma, A. M. *et al.* Molecular Pathways of Colon Inflammation Induced by Cancer Immunotherapy.

Cell **182** (2020). https://doi.org:10.1016/j.cell.2020.06.001

83   Zhang, X. *et al.* Persistence of cirrhosis is maintained by intrahepatic regulatory T cells that inhibit

fibrosis resolution by regulating the balance of tissue inhibitors of metalloproteinases and matrix

metalloproteinases. *Transl Res* **169**, 67-79 e61-62 (2016). https://doi.org:10.1016/j.trsl.2015.10.008

84   Shevtsov, M. & Multhoff, G. Heat Shock Protein-Peptide and HSP-Based Immunotherapies for the

Treatment of Cancer. *Front Immunol* **7**, 171 (2016). https://doi.org:10.3389/fimmu.2016.00171

85   Zhang, Q. *et al.* Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma.

Cell **179**, 829-845 e820 (2019). https://doi.org:10.1016/j.cell.2019.10.003

86   Wang, X. *et al.* Single-cell dissection of remodeled inflammatory ecosystem in primary and

metastatic   gallbladder   carcinoma.   *Cell   Discov*   **8**,   101   (2022).

https://doi.org:10.1038/s41421-022-00445-8

87   Tang, J., Yan, Z., Feng, Q., Yu, L. & Wang, H. The Roles of Neutrophils in the Pathogenesis of Liver

Diseases. *Front Immunol* **12**, 625472 (2021). https://doi.org:10.3389/fimmu.2021.625472

88   Cho, Y. & Szabo, G. Two Faces of Neutrophils in Liver Disease Development and Progression.

Hepatology **74**, 503-512 (2021). https://doi.org:10.1002/hep.31680

89   Wang,   S.   *et   al.*   S100A8/A9   in   Inflammation.   *Front   Immunol*   **9**,   1298   (2018).

https://doi.org:10.3389/fimmu.2018.01298

90   Chen, Y., Ouyang, Y., Li, Z., Wang, X. & Ma, J. S100A8 and S100A9 in Cancer. *Biochim Biophys*

Acta Rev Cancer **1878**, 188891 (2023). https://doi.org:10.1016/j.bbcan.2023.188891

91   André, P. *et al.* Anti-NKG2A mAb Is a Checkpoint Inhibitor that Promotes Anti-tumor Immunity by

Unleashing Both T and NK Cells. *Cell* **175**, 1731-+ (2018). https://doi.org:10.1016/j.cell.2018.10.014

92   Brillantes, M. & Beaulieu, A. M. Transcriptional control of natural killer cell differentiation.

Immunology **156**, 111-119 (2019). https://doi.org:10.1111/imm.13017

93   Zhang, J., Rousseaux, N. & Walzer, T. Eomes and T-bet, a dynamic duo regulating NK cell

differentiation. *Bioessays* **44**, e2100281 (2022). https://doi.org:10.1002/bies.202100281

94   da Silva, I. P. *et al.* Reversal of NK-cell exhaustion in advanced melanoma by Tim-3 blockade.

Cancer Immunol Res **2**, 410-422 (2014). https://doi.org:10.1158/2326-6066.CIR-13-0171

95   Sun, C., Sun, H., Zhang, C. & Tian, Z. NK cell receptor imbalance and NK cell dysfunction in HBV

infection   and   hepatocellular   carcinoma.   *Cell   Mol   Immunol*   **12**,   292-302   (2015).

https://doi.org/10.1038/cmi.2014.91

96   Hosseini, R. *et al.* Cancer exosomes and natural killer cells dysfunction: biological roles, clinical

significance and implications for immunotherapy. *Molecular Cancer* **21** (2022). https://doi.org:ARTN

15

10.1186/s12943-021-01492-7

97   Park, J., Hsueh, P. C., Li, Z. Y. & Ho, P. C. Microenvironment-driven metabolic adaptations guiding

CD8+T   cell   anti-tumor   immunity.   *Immunity*   **56**,   32-42   (2023).

https://doi.org/10.1016/j.immuni.2022.12.008

98   Yan, C. S. *et al.* Exhaustion-associated cholesterol deficiency dampens the cytotoxic arm of

antitumor immunity. *Cancer Cell* **41**, 1276-+ (2023). https://doi.org/10.1016/j.ccell.2023.04.016

99   Bader, J. E., Voss, K. & Rathmell, J. C. Targeting Metabolism to Improve the Tumor

Microenvironment   for   Cancer   Immunotherapy.   *Mol   Cell*   **78**,   1019-1033   (2020).

https://doi.org/10.1016/j.molcel.2020.05.034

100   Riera-Domingo, C. *et al.* Immunity, Hypoxia, and Metabolism-the Menage a Trois of Cancer:

Implications   for   Immunotherapy.   *Physiol   Rev*   **100**,   1-102   (2020).

https://doi.org/10.1152/physrev.00018.2019

101   Wu, S. Y., Fu, T., Jiang, Y. Z. & Shao, Z. M. Natural killer cells in cancer biology and therapy.

Molecular Cancer **19** (2020). https://doi.org:ARTN 120

10.1186/s12943-020-01238-x

102   Poznanski, S. M. & Ashkar, A. A. What Defines NK Cell Functional Fate: Phenotype or Metabolism?

767    *Frontiers in Immunology* **10** (2019). https://doi.org:ARTN 1414

768 10.3389/fimmu.2019.01414

769 103 O'Brien, K. L. & Finlay, D. K. Immunometabolism and natural killer cell responses. *Nat Rev Immunol*

770    **19**, 282-290 (2019). https://doi.org:10.1038/s41577-019-0139-2

771 104 Tsvetkov, P. *et al.* Copper induces cell death by targeting lipoylated TCA cycle proteins. *Science*

772    **375**, 1254-+ (2022). https://doi.org:10.1126/science.abf0529

773 105 Yu, G. C., Wang, L. G., Han, Y. Y. & He, Q. Y. clusterProfiler: an R Package for Comparing

774    Biological Themes Among Gene Clusters. *Omics* **16**, 284-287 (2012).

775    https://doi.org:10.1089/omi.2011.0118

776 106 Jin, S. Q. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature*

777    *Communications* **12** (2021). https://doi.org:ARTN 1088

778 10.1038/s41467-021-21246-9

779 107 Qiu, X. J. *et al.* Single-cell mRNA quantification and differential analysis with Census. *Nat Methods*

780    **14**, 309-+ (2017). https://doi.org:10.1038/Nmeth.4150

781 108 Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of developmental potential.

782    *Science* **367**, 405-+ (2020). https://doi.org:10.1126/science.aax0249

783 109 van de Sande, B. *et al.* A scalable SCENIC workflow for single-cell gene regulatory network analysis.

784    *Nat Protoc* **15**, 2247-2276 (2020). https://doi.org:10.1038/s41596-020-0336-2

785 110 Andreatta, M. & Carmona, S. J. UCell: Robust and scalable single-cell gene signature scoring.

786    *Comput Struct Biotec* **19**, 3796-3798 (2021). https://doi.org:10.1016/j.csbj.2021.06.043

Figure 1

a.

Genotypes of ATP7B in 20 WD cases

Mutant alleles*	Amino acid change	No. (%)
c.2333G>T	p.R778L	13 (65%)
c.2804C>T	p.T935M	0 (0%)
c.2975C>T	p.P992L	0 (0%)
Unknow	Unknow	7 (35%)

* Reference. Seq: NM_000053.4

b.

c.

Control   WD

Masson-trichrome

d.

Kayser-Fleischer ring

e.

Clinical statistics of the patients.

Cohort 1			
	WD (n=92)	HBV (n=95)	P value
TBIL - µmol/L (≤26)	14.90±6.59	20.46±13.94	0.0007
TG - µmol/L (≤1.7)	0.98±0.58	0.99±0.71	0.7699
T-CHOL - µmol/L (3-5.7)	4.09±0.75	3.84±1.13	0.0743

Cohort 2			
	WD (n=618)	HBV (n=370)	P value
Hyperglycemia – no. (%)	33 (5.34%)	42 (11.35%)	0.0006
Gallstone – no. (%)	65 (10.52%)	39 (10.54%)	0.9910
Cholecystitis – no. (%)	306 (49.51%)	35 (9.46%)	0.0001

Cohort 3			
	Cholecystitis & WD (n=34)	Cholecystitis (n=44)	P value
WBC – no. (%)	12 (35.29%)	1 (2.27%)	0.0004
NEU – no. (%)	10 (29.41%)	5 (11.36%)	0.0449
LYM – no. (%)	11 (32.35%)	6 (13.64%)	0.0471
TBIL – no. (%)	5 (14.71%)	6 (13.64%)	0.8929
ALB – no. (%)	10 (29.41%)	10 (22.73%)	0.5026
ALT – no. (%)	18 (52.94%)	28 (63.64%)	0.3410
AST – no. (%)	9 (26.47%)	11 (25.00%)	0.8827

WD, Wilson disease; HBV, Hepatitis B virus; ; TBIL, Total bilirubin; TG, Triglyceride; T-CHOL, Total cholesterol; WBC, White blood cell; NEU, Neutrophil; LYM, Lymphocyte; ALB, Albumin; ALT, Alanine aminotransferase; AST, Aspartate aminotransferase.

Figure 1. Clinical manifestation of WD patients and WD causes poor prognosis of cholecystitis. **a.**
DNA sequencing result of 20 WD patients was summarized in the graph. **b.** The schematic diagram
illustrates copper transport disturbances caused by ATP7B mutations, along with structural and functional
impairments in mitochondria. Created with BioRender.com. **c.** Masson-trichrome staining of liver
tissues of patients with WD and control. **d.** The picture of Kayser-Fleischer (KF) ring in the cornea of a
patient with WD, captured during clinical examination. **e.** The table of clinical statistics of patients in 3
cohorts. Data are mean± S.D. Unpaired two-tailed t-test and chi-square test.

Figure 2

798   **Figure 2. Single-cell profiling of non-parenchymal cells in the liver of WD patients.**   **a.** The workflow

799   of scRNA-seq.   Created with BioRender.com.   **b.** UMAP visualization of 26 principal components of all

800   the single cells.   **c.** UMAP visualization of 26 principal components by samples.   **d.** UMAP visualization

801   of 12 main cell types.   **e.** The dot plot showing the expression of top 3 marker genes in 12 main cell types.

Figure 3

Figure 3. The altered immune microenvironment in WD patients.   **a.** UMAP visualization of MPs

colored and labeled by cell subtypes (left).   The bar chart showing the average percentage of different

subtypes in case group (n=3) and control group (n=3) (right).   Data are mean ± S.D.   **b.** UMAP

visualization of T and NK cells colored and labeled by cell subtypes (left).   The bar chart showing the

average percentage of different subtypes in case group (n=3) and control group (n=3) (right).   Data are

mean± S.D.   **c.** The heatmap showing the top 20 differently expressed genes (DEGs) for all immune cells.

d. The dot plot showing GO enrichment on DEGs of all immune cells between the case and control group.

BP, biological processes; CC, cellular components; MF, molecular function.   **e.** The scatter showing the

strength of all cell subtypes in signal incoming and outgoing. Upper, the case group; down, the control group.

Figure 4

Figure 4. The dysfunction of main immune cells in WD patients. **a.** The violin plots displaying M1

polarization, M2 polarization, pro-inflammatory signature, and interferon responsed signature of the 4

subtypes within macrophages between two groups by gene scoring. **b.** The violin plots displaying M1

polarization, M2 polarization, pro-inflammatory signature, and interferon responsed signature of the 3

subtypes within Kupffer cells between two groups by gene scoring. **c.** The violin plots displaying

maturation, phagocytosis, chemotaxis, chemokine activity and Type I interferon signaling pathway scores within

neutrophils between two groups. **d.** The heatmap showing the relative expression level of genes in each

cell subtype between the case and control group. **e.** The dot plot showing biological process of GO

enrichment on differentially expressed genes (DEGs) in monocytes between the case and control group.

Upper, upregulation; down, downregulation. **f.** The violin plots displaying cytotoxicity and exhaustion

scores of the 6 cell subtypes within CD8 T cells between two groups by gene scoring. **g.** The violin plots

displaying Treg scores of the 3 cell subtypes within CD4 T cells between two groups by gene scoring.

Figure 5

Figure 5. Identification of NK cell exhaustion in WD patients.   **a.** The violin plots showing the relative

expression level of genes in NK cells between the case and control group.   ***, *P*<0.001; Wilcox test.   **b.**

Trajectories of NK cells in the case and control group.   **c.** Representative immunofluorescence staining for

NK cells (CD56) and expression of NKG2A in liver tissues of WD and control.   Scale bars, 100 μm for

40X and 50 μm for 100X.   NKG2A^{+} NK cells are highlighted by arrows.   Quantifications were

performed by assessing three random fields per slide.   Data are mean± S.D.   Unpaired two-tailed t-test.

d. Gating strategy applied in flow cytometry.   **e.** The bar charts showing the percentage of NKG2A^{+} (WD,

n=5; Control, n=5), TIGIT^{+} (WD, n=3; Control, n=3), and IFNγ^{+} (WD, n=4; Control, n=4) NK cells.   Data

are mean± S.D.   Unpaired two-tailed t-test.

Figure 6

836   **Figure 6. NK cell exhaustion predicts poor prognosis of inflammatory diseases.**   **a.-d.** Kaplan-Meier

837   survival curves for cholecystitis, pancreatitis, lung adenocarcinoma, and hepatocellular carcinoma

838   according to the markers of NK cell exhaustion in the GSE166915, GSE91035, GSE75037, and GSE41804

839   datasets.

Figure 1

a.

Genotypes of ATP7B in 20 WD cases

Mutant alleles*	Amino acid change	No. (%)
c.2333G>T	p.R778L	13 (65%)
c.2804C>T	p.T935M	0 (0%)
c.2975C>T	p.P992L	0 (0%)
Unknow	Unknow	7 (35%)

* Reference. Seq: NM_000053.4

b.

c.

Masson-trichrome | Control | WD

d.

Kayser-Fleischer ring

e.

Clinical statistics of the patients.

Cohort 1

	WD (n=92)	HBV (n=95)	*P* value
TBIL - µmol/L (≤26)	14.90±6.59	20.46±13.94	0.0007
TG - µmol/L (≤1.7)	0.98±0.58	0.99±0.71	0.7699
T-CHOL - µmol/L (3-5.7)	4.09±0.75	3.84±1.13	0.0743

Cohort 2

	WD (n=618)	HBV (n=370)	*P* value
Hyperqlycemia – no. (%)	33 (5.34%)	42 (11.35%)	0.0006
Gallstone – no. (%)	65 (10.52%)	39 (10.54%)	0.9910
Cholecystitis – no. (%)	306 (49.51%)	35 (9.46%)	0.0001

Cohort 3

	Cholecystitis & WD (n=34)	Cholecystitis (n=44)	*P* value
WBC – no. (%)	12 (35.29%)	1 (2.27%)	0.0004
NEU – no. (%)	10 (29.41%)	5 (11.36%)	0.0449
LYM – no. (%)	11 (32.35%)	6 (13.64%)	0.0471
TBIL – no. (%)	5 (14.71%)	6 (13.64%)	0.8929
ALB – no. (%)	10 (29.41%)	10 (22.73%)	0.5026
ALT – no. (%)	18 (52.94%)	28 (63.64%)	0.3410
AST – no. (%)	9 (26.47%)	11 (25.00%)	0.8827

WD, Wilson disease; HBV, Hepatitis B virus; ; TBIL, Total bilirubin; TG, Triglyceride; T-CHOL, Total cholesterol; WBC, White blood cell; NEU, Neutrophil; LYM, Lymphocyte; ALB, Albumin; ALT, Alanine aminotransferase; AST, Aspartate aminotransferase.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6