Observing one-divalent-metal-ion dependent and histidine-promoted His-Me family I-PpoI nuclease catalysis in crystallo

Caleb Chang[1], Grace Zhou[1], and Yang Gao[1,*]

[1]Department of Biosciences, Rice University, Houston, Texas, 77005, USA

[*]Correspondence: yg60@rice.edu , +1 (713) 348-2619

Abstract

Metal-ion-dependent nucleases play crucial roles in cellular defense and biotechnological applications. Time-resolved crystallography has resolved catalytic details of metal-ion-dependent DNA hydrolysis and synthesis, uncovering the essential roles of multiple metal ions during catalysis. The histidine-metal (His-Me) superfamily nucleases are renowned for binding one divalent metal ion and requiring a conserved histidine to promote catalysis. Many His-Me family nucleases, including homing endonucleases and Cas9 nuclease, have been adapted for biotechnological and biomedical applications. However, it remains unclear how the single metal ion in His-Me nucleases, together with the histidine, promotes water deprotonation, nucleophilic attack, and phosphodiester bond breakage. By observing DNA hydrolysis *in crystallo* with His-Me I-PpoI nuclease as a model system, we proved that only one divalent metal ion is required during its catalysis. Moreover, we uncovered several possible deprotonation pathways for the nucleophilic water. Interestingly, binding of the single metal ion and water deprotonation are concerted during catalysis. Our results reveal catalytic details of His-Me nucleases, which is distinct from multi-metal-ion-dependent DNA polymerases and nucleases.

Introduction

Mg^{2+}-dependent nucleases play fundamental roles in DNA replication and repair [1-4], RNA processing [5-8], as well as immune response and defense [9-11]. Moreover, they are widely employed for genome editing in biotechnological and biomedical applications [12,13]. These nucleases are proposed to cleave DNA through a SN$_2$ type reaction, in which a water molecule, or sometimes, a tyrosine side chain [14], initiates the nucleophilic attack on the scissile phosphate with the help of metal ions [15]. Metal ions can orient and stabilize the binding of the negatively charged nucleic acid backbone [16], promoting proton transfer, nucleophilic attack, and stabilization of the transition state. As a highly varied family of enzymes, Mg^{2+}-dependent nucleases can be broadly categorized by the number of metal ions captured in their active site. So far, Mg^{2+}-dependent nucleases with one and two metal ions have been observed. In the two-Mg^{2+}-ion dependent nuclease, a metal ion binds on the leaving group side of the scissile phosphate (Me$^{2+}_B$) while the other binds on the nucleophile side (Me$^{2+}_A$). In one-metal ion dependent nucleases, only the metal ion corresponding to the B site in two-metal ion dependent nucleases is present [17]. Moreover, transiently bound metal ions have been identified in the previously thought two-metal-ion RNaseH [18] and EndoV nucleases [19] *via* time-resolved X-ray crystallography, proposed to play key roles in various stages of their catalysis. Similarly, catalysis in the one-metal ion dependent APE1 nuclease has been observed *in crystallo*, but mechanistic details regarding its metal-ion dependence have not been thoroughly explored [20-22]. There also exist three-Zn^{2+} dependent nucleases, with two Zn^{2+} binding in the A-equivalent and B-equivalent positions, while the third Zn^{2+} coordinating the sp oxygen on the nucleophile side of the scissile phosphate [23]. However, it remains unclear how the single metal ion in one-metal-ion dependent nucleases is capable of aligning the substrate, promoting deprotonation and nucleophilic attack, and stabilizing the pentacovalent transition state.

A large subfamily of one-metal-ion dependent nucleases consist of histidine-metal (His-Me) nucleases that perform critical tasks in biological pathways such as apoptosis [24,25], extracellular defense [26,27], intracellular immunity (CRISPR-Cas9) [28,29], and intron homing (homing endonuclease) [30,31]. Despite sharing poor sequence homology, the structural cores and active sites of His-Me nucleases are highly conserved and thus are proposed to catalyze DNA hydrolysis through a similar mechanism [15,31,32]. Surrounded by two beta sheets and an alpha helix, the His-Me nucleases active sites all contain a single metal ion, a histidine, and an asparagine (Fig. 1A, S1A, and S2). The asparagine residue helps to stabilize metal ion binding whereas the strictly conserved histidine is suggested to deprotonate a nearby water for nucleophilic attack towards the scissile phosphate [33-36]. To this day, the dynamic reaction process of DNA hydrolysis by His-Me nucleases has never been visualized, and the mechanism of single metal ion-dependent and histidine-promoted catalysis remains unclear. Emerging genome editing and disease treatment involving CRISPR-Cas9 emphasize the importance of understanding the catalytic mechanism of DNA hydrolysis by His-Me nucleases [37-39]. Recent crystal and cryo-electron microscope structures of Cas9 have captured the His-Me family Cas9 histidine-asparagine-histidine (HNH) active site engaged with DNA before and after cleavage [40-43]. However, due to the relatively low resolution and the static nature of the structures, key catalytic details, such as metal ion dependence, transition-state stabilization, and alternative deprotonation pathways, remain elusive. Moreover, the large size and multiple conformational checkpoints during Cas9 catalysis [40-43] hinder *in crystallo* observation of Cas9 catalysis.

Fig. 1. Observing His-Me family I-PpoI catalyze DNA hydrolysis *in crystallo*.
(**A**) Model of one-metal-ion dependent and histidine promoted His-Me enzyme catalysis and transition state stabilization. (**B**) Metal ion soaking setup for *in crystallo* observation of DNA hydrolysis with I-PpoI. (**C-E**) Structural intermediates of I-PpoI *in crystallo* DNA cleavage showcasing the pre-reaction state in (**C**) and product states in (**D**) and (**E**). The $2F_o$-F_c map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ (σ values represent r.m.s. density values). (**E**) The F_o-F_c omit map for the up conformation of the product (violet) was contoured at 3.0 σ.

I-PpoI is a well-characterized intron-encoded homing endonuclease member of the *physarum polycephalum* slimemold. By 1999, Stoddard and colleagues were able to capture intermediate structures of I-PpoI complexed with DNA before and after product formation [33,44]. We herein employed I-PpoI as a model system and applied time-resolved crystallography to observe the catalytic process of His-Me nuclease. By determining over 40 atomic resolution structures of I-PpoI during its reaction process, we show that one and only one divalent metal ion is involved in DNA hydrolysis. Moreover, we uncover several possible deprotonation pathways for the nucleophilic water. Notably, metal ion binding and water deprotonation are highly concerted during catalysis. Our findings provide mechanistic insights into one-metal-ion dependent nucleases, enhancing future design and engineering of these enzymes for emerging biomedical applications.

Results
Preparation of the I-PpoI system for in crystallo DNA hydrolysis

We sought to implement I-PpoI for *in crystallo* metal ion soaking (Fig. 1B), which has been successful in elucidating the catalytic mechanisms of DNA polymerases [45-50], nucleases [18-21], and glycosylase [51]. First, a complex of I-PpoI and a palindromic DNA was crystalized at pH 6 with 0.2 M sodium malonate. Similar to previous studies, a dimer of I-PpoI was found in the asymmetric unit, with both active sites engaged for catalysis and DNA in the middle bent by 55° (Fig. S1A). The two I-PpoI molecules were almost identical and thus served as internal controls for evaluating the reaction process (Fig. S1B). Within each I-PpoI active site, a water molecule (nucleophilic water, WatN) existed 3.6 Å from the scissile phosphate, near the imidazole side chain of His98. On the leaving group side of the scissile phosphate, the metal exhibited an octahedral geometry, being coordinated by three water molecules, two oxygen atoms from the scissile phosphate, and the conserved asparagine (Fig. 1C).

Next, we removed malonate in the crystallization buffer, which may chelate metal ions and hinder metal ion diffusion [52], by equilibrating the crystals in 200 mM NaCl buffer at pH 6, 7, or 8 for 30 min. The diffraction quality of the crystals was not affected during the soaking. The structures of I-PpoI equilibrated at pH 6, 7, or 8 in the presence of 200 mM NaCl showed no signs of product formation and appeared similar to the malonate and previous reported I-PpoI structures [33,44] (Fig. S3A). To confirm if a monovalent metal ion binds in the pre-reaction state (PRS), we soaked the I-PpoI crystals in buffer containing Tl^+ and detected anomalous electron density at the metal ion binding site without product formation (Fig. 3SB). Our results support that the monovalent metal ion can bind within the active site without initiating reaction, in corroboration with our biochemical assays (Fig. S3C).

Witnessing DNA hydrolysis in crystallo by I-PpoI

To initiate the chemical reaction *in crystallo*, we transferred the I-PpoI crystals equilibrated in NaCl buffer to a reaction buffer with 500 µM Mg^{2+} (Fig. 1B). After 600 s soaking, we saw a significant negative F_o-F_c peak on the leaving group side of the scissile phosphate atom as well as a positive F_o-F_c peak on the other side (Fig. S3D), indicating that DNA hydrolysis was occurring *in crystallo*. After DNA cleavage, the newly generated phosphate group shifted 1 Å towards the WatN (Fig. 1D). Furthermore, an additional product state was observed after soaking the crystals in 500 µM Mg^{2+} for 600 s and 1800 s, at which the newly formed phosphate shifted 3 Å away from the metal ion to form a hydrogen bond with Arg61 (Fig. 1E). Our results confirm that the implementation of I-PpoI with *in crystallo* Me^{2+} soaking is feasible for observing the I-PpoI catalytic process and dissecting the mechanism of His-Me nucleases.

With an established *in crystallo* reaction system, we next monitored the reaction process by soaking I-PpoI crystals in buffer containing 500 µM Mg^{2+} pH 7, for 10-1200 s. Density between the reactant phosphate and nucleophilic water increased along with longer soaking time, indicating the generation of phosphate products (Fig. 2A). During the reaction, the coordination distances of the metal ion ligands decreased 0.1-0.2 Å when Mg^{2+} exchanged with Na^+, which is consistent with their preferred geometry [53] (Fig. 2B, C). At reaction time 160 s, 45% of product had been generated (Fig. 2D), which later plateaued to 65% at 300 s. During the reaction process, the sugar ring of the reactant and product DNA that resides around 3 Å away from the scissile phosphate, remained in a C3´-endo conformation. As Mg^{2+} soaking time increased from 10-160 s, we observed the WatN approaching the scissile phosphate (Fig. 2E). At the same time, the conserved His98 sidechain proposed to deprotonate the WatN was slightly rotated.

142
Fig. 2. *In crystallo* **DNA hydrolysis by I-PpoI.**
(**A**) Structures of I-PpoI during *in crystallo* catalysis after 500 µM Mg^{2+} soaking for 0 s, 20 s,
40 s, 80 s, 160 s, 320 s. The F_o-F_c omit map for the product phosphate (green) was contoured at
3.0 σ. (**B-D**) I-PpoI complexes featuring metal ion coordination when bound with Na^+ in (**B**)
Mg^{2+} in the earlier time point of the reaction process in (**C**), and Mg^{2+} in the later time point of
the reaction process in (**D**). The $2F_o$-F_c map for Me^{2+}, DNA, waters (red spheres), and catalytic
residues (blue) was contoured at 2.0 σ. (**E**) Alignment of the WatN and rotation in His98 during I-
PpoI reaction.

151
152

A single divalent metal ion was captured during DNA hydrolysis
Throughout the reaction process with Mg^{2+}, we found that the electron density for the Mg^{2+} metal
ion strongly correlated (R^2=0.97) with product formation (Fig. S4A, C), which may suggest that
saturation of this metal ion site is required and sufficient for catalysis. However, Mg^{2+}'s similar
size to Na^+ makes it suboptimal for quantifying metal ion binding. To thoroughly investigate
metal-ion dependence, we repeated the *in crystallo* soaking experiment with Mn^{2+}, which is more
electron rich and can be unambiguously assigned based on its electron density and anomalous
signal. With 500 µM Mn^{2+} in the reaction buffer, we found that the reaction process and the
product conformation were similar to that for Mg^{2+}. After 160 s Mn^{2+} soaking, clear anomalous
signal was present at the metal ion binding site, confirming the binding of Mn^{2+} (Fig. S4F). For
crystal structures of I-PpoI with partial product formation, Mn^{2+} signal at the metal ion site
correlated with product formation with a R^2 of 0.98 (Fig. 3A and S4B). To further search if
additional and transiently-bound divalent ions participate in the reaction, we soaked the I-PpoI
crystals in high concentration of Mn^{2+} (200 mM) for 600 s, at which 80% product formed within
the active site. However, apart from the single metal ion binding site, we do not detect anomalous
signal for additional Mn^{2+}, despite such high concentration of Mn^{2+} (Fig. 3B and S5). The strong
correlation between product phosphate formation with Mn^{2+} binding and the absence of additional
anomalous density peaks in heavy Mn^{2+} soaked crystals suggest that one and only one divalent
metal ion is involved in I-PpoI DNA hydrolysis.

Fig. 3. Detection of Me^{2+} binding during DNA hydrolysis *in crystallo*.
(**A**) Correlation (R^2) between the newly formed phosphate and Mn^{2+} binding at pH 6. The points represent the mean of duplicate measurements for the electron density of the reaction product phosphate within two I-PpoI molecules in the asymmetric unit while the errors bars represent the standard deviation. (**B**) Additional Mn^{2+} binding sites were not detected after 0.9765 Å X-ray diffraction in the I-PpoI active site after 10 min soaking in 200 mM Mn^{2+}. The 2F$_o$-F$_c$ map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ. The anomalous map for Mn^{2+} was contoured at 3.0 σ.

To examine if additional monovalent metal ions participate during DNA hydrolysis, we titrated Tl$^+$, which can replace monovalent Na$^+$ or K$^+$ and yields anomalous signal [54,55], at up to 100 mM concentration in the biochemical assay. Apart from precipitation that occurred at 50 and 100 mM Tl$^+$, we found that product conversion remained unaffected with increasing Tl$^+$ in solution (Fig. S4G). Similarly, increasing Na$^+$ concentration in solution did not increase product conversion (Fig. S4H). In addition, only one Tl$^+$ was detected by its anomalous signal in the pre-reaction state (Fig. S3B). The biochemical and structural results indicate that additional monovalent metal ion may not be necessary for DNA hydrolysis. However, crystal deterioration limited us from soaking the I-PpoI crystals in high concentration of Tl$^+$ for *in crystallo* reaction.

pH-dependence of I-PpoI DNA cleavage
During DNA hydrolysis, deprotonation of the WatN is required for the nucleophilic attack and phosphodiester bond breakage. This proton transfer has been proposed to be mediated by the highly conserved His98 [33,44] that lies within 3 Å from the WatN. We speculated that the ability of His98 to activate the nearby WatN and mediate proton transfer would be affected by pH. The DNA cleavage assay revealed that I-PpoI cleavage activity increased with pH with a pKa of 8.3 (Fig. 4A), which is much higher than the pKa of histidine, but the histidine pKa and the proton transfer process may be affected by the active site environment. To explore how pH affects the active site configuration and catalysis, we conducted *in crystallo* soaking experiments with Mg^{2+} at pH 6 and pH 8 in addition to the pH 7 data series in Figure 2. Consistent with in solution experiments, higher pH resulted in faster product formation *in crystallo* (Fig. 4B), indicating that metal ion binding may also be affected by pH. To test this, we performed Mg^{2+} titration at different pH in solution. Our results showed that over 100-times higher concentration of Mg^{2+} was needed to yield 50% product in pH 6 versus pH 8 (Fig. 4C), confirming that pH affects metal-ion dependent I-PpoI catalysis. Likewise, the reactions *in* crystallo at higher pH reached 50% product formation at shorter soaking times (320 s at pH 6, 160 s at pH 7, and 80 s at pH 8, respectively). Interestingly, the crystal structures that contained 30-35% product at pH 6 and pH 8 were nearly identical (Fig. 4D, E). The positions of the His98 residue and the WatN were practically superimposable (Fig. 4F). At all pH, Mg^{2+} binding strongly correlated with product

formation (R^2 greater than 0.95), suggesting that low pH reduces the overall reaction rate without altering the reaction pathway (Fig. S4D, E).

Fig. 4. Effect of pH on I-PpoI DNA hydrolysis.
(**A**) DNA hydrolysis by WT I-PpoI with increasing pH in solution. (**B**) DNA hydrolysis by WT I-PpoI *in crystallo* at pH 6, 7, and 8. The points represent the mean of duplicate measurements for the electron density of the reaction product phosphate after a period of Mg^{2+} soaking within two I-PpoI molecules in the asymmetric unit. The errors bars represent the standard deviation. (**C**) The effect of pH on metal ion dependence in solution. The points represent the mean of triplicate measurements for the percentage of cleaved DNA product while the errors bars represent the standard deviation. (**D**) Structure of I-PpoI *in crystallo* DNA hydrolysis at pH 6 after 320 s of 500 µM Mg^{2+} soaking. (**E**) Structure of I-PpoI *in crystallo* DNA hydrolysis at pH 8 after 20 s of 500 µM Mg^{2+} soaking. (**D, E**) The $2F_o$-F_c map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ. (**F**) Structural comparison of the active site after 500 µM Mg^{2+} soaking for 320 s at pH 6 (purple), 160 s at pH 7 (yellow), and 20 s at pH 8 (green).

Nucleophilic water deprotonation pathway during I-PpoI DNA cleavage
We next investigated the deprotonation pathway with mutagenesis. Because His98 has been proposed to primarily activate the nucleophilic water, we first mutated His98 to alanine (Fig. 5A). As expected, cleavage activity of H98A I-PpoI drastically dropped. Our assays showed that H98A I-PpoI displayed residual activity but required a reaction time of 1 hr to be comparable to WT I-PpoI (Fig. 5B), similar to the H98Q mutant in previous studies [56]. Furthermore, varying the pH resulted in a sigmoidal activity curve of I-PpoI pH dependence, corresponding to a pKa of 7.9 (Fig. 5C). The significantly reduced reaction rate and the shift of pH dependence suggest that His98 plays a key role in pH sensing and water deprotonation. On the other hand, the low but existing activity of H98A I-PpoI suggests the presence of alternative general bases for proton transfer. Another histidine (His78) resides on the nucleophile side 3.8 Å from the WatN, with a cluster of water molecules in between (Fig. 5A). We hypothesized that His78 may substitute His98 as the proton acceptor. The DNA cleavage assays revealed that the single mutant, H78A I-PpoI, had an activity similar to WT, whereas the double mutant (H78A/H98A I-PpoI) exhibited much lower activity than that of H98A (Fig. 5B). Furthermore, varying the pH for H78A/H98A I-

PpoI resulted in a sigmoidal activity curve that corresponded to a pKa of 8.7 (Fig. 5C). In consistent with previous speculations, our results confirm the possibility of His78 as an alternative general base [56]. The residual activity and pH dependence of H78A/H98A I-PpoI indicated that something else was still activating the nucleophilic water in the absence of any nearby histidine. Furthermore, we found that titrating imidazole in H98A and H78A/H98A I-PpoI partially rescued cleavage activity (Fig. S6A, B), similar as observed in Cas9 and EndA nuclease [57,58]. Collectively, our results indicate that His98 is the primary proton acceptor like previous simulation results [59] but at the same time, I-PpoI can use alternative pathways to activate the nucleophilic WatN.

Fig. 5. Nucleophilic water deprotonation during I-PpoI DNA hydrolysis.
(**A**) Active site environment surrounding WatN. His78 exists near the WatN besides His98. (**B**) In solution DNA hydrolysis activity of various I-PpoI histidine mutants. The points represent the mean of triplicate measurements for the percentage of cleaved reaction product while the errors bars (too small to see) represent the standard deviation. (**C**) DNA hydrolysis by H98A I-PpoI (blue) and H78A/H98A I-PpoI (green) at various pH in solution. (**D**) Structure of H98A I-PpoI active site after 0.9765 Å X-ray diffraction after 1 mM Mn^{2+} soaking for 1800 s. The anomalous map for Mn^{2+} was contoured at 2.0 σ. The $2F_o$-F_c map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ. (**B, C**) The points represent the mean of triplicate measurements for the percentage of generated reaction product while the errors bars represent the standard deviation.

We next sought to understand the mechanism of His98-promoted hydrolysis from a structural standpoint. In our *in crystallo* soaking experiments, we equilibrated H98A I-PpoI crystals in 500 µM Mg^{2+}. The structure looked nearly identical to the Na^+ structure and previous H98A I-PpoI structure with Mg^{2+}. As shown earlier, the coordination environment of Na^+ and Mg^{2+} was quite similar. To confirm divalent metal ion binding, the H98A I-PpoI crystals were soaked for 1800 s in 500 µM Mn^{2+}, the same concentration used to initiate DNA hydrolysis by WT I-PpoI. However, to our surprise, the metal ion binding site was devoid of any anomalous signal. Increasing the Mn^{2+} concentration to 1 mM still did not produce anomalous density at the Me^{2+} binding site (Fig. 5D and S6C). The results indicate that metal ion binding can be altered by perturbing His98 and possibly water deprotonation, even though the metal ion binding site and His98 exist 7 Å apart without direct interaction. Although we tried soaking the H98A I-PpoI crystals in 1 mM Mg/Mn^{2+} and 100 mM imidazole for 15 h, metal ion binding, imidazole binding

279 or product formation were not detected (Fig. S6D), possibly due to the difficulty of imidazole
280 diffusion within the lattice. Our results indicate that perturbing the deprotonation pathway not
281 only affects nucleophilic attack but also metal binding, suggesting that I-PpoI catalyzes DNA
282 catalysis *via* a concerted mechanism.

Discussion

Time-resolved crystallography can visualize time-dependent structural changes and elucidate
mechanisms of enzyme catalysis with unparalleled detail *in crystallo*, especially for light-
dependent enzymes, in which the reactions can be synchronously initiated by light pulses [60-63]. In
complementary, recently advanced metal ion diffusion-based time-resolved crystallographic
techniques have uncovered rich dynamics at the active site and transient metal ion binding during
the catalytic processes of metal-ion-dependent DNA polymerases [45-50], nucleases [18,19] and
glycosylase [51]. In addition to metal ions captured in static structures, these transient metal ions
have been shown to play critical roles in catalysis, such as water deprotonation for nucleophilic
attack in RNaseH [18], bond breakage and product stabilization in polymerases [45-48,50,64,65] and
alignment of the substrate and nucleophilic water in MutT [66]. Interestingly, one and only one
metal ion was captured within the I-PpoI active site during catalysis, even when high
concentration (200 mM Mn^{2+}) of metal ion was tested (Fig. 3). The observed location of the Me^{2+}
on the leaving group side of the scissile phosphate corresponds to the Me^{2+}_B in two metal-ion
dependent nucleases [17] (Fig. S2). However, the metal ion is unique in its environment and role.
First, it is coordinated by a water cluster and asparagine side chain (sometimes with an additional
aspartate residue) rather than the acidic aspartate and glutamic acid clusters that outline the active
sites of RNaseH and APE1 nuclease [67] as well as DNA polymerases (Fig. S7). Even in the
absence of divalent Mg^{2+} or Mn^{2+}, the active site including the scissile phosphate was already
well-aligned in I-PpoI, which is again different from RNaseH. Instead, Leu116 and the beta sheet
consisting of Arg61, Gln63, Lys65, Thr67 [44,68] helped to position the DNA optimally towards the
active site (Fig. S1A, C). Second, single metal ion binding is strictly correlated with product
formation in all conditions, at different pH and with different mutants (Fig. 3A and S4A, B, C-E)
[58]. Thus, similar to the third metal ion in DNA polymerases and RNaseH, the metal ion in I-PpoI
is not required for substrate alignment but is essential for catalysis. We suspect that the single
metal ion helps stabilize the transition state and reduce the electronegative buildup of DNA,
thereby promoting DNA hydrolysis.

Proton transfer by a general base is essential for a SN_2-type nucleophilic attack. Such
deprotonation of the nucleophilic water has been attributed to His98, which is highly conserved in
His-Me nucleases. Existing close to the nucleophilic water at 2.6 Å, His98 is perfectly positioned
to mediate the proton transfer. Moreover, due to the bulky presence of His98 and beta sheet
protein residues, there is no space for an additional metal ion at the nucleophilic side (Fig. S5).
The H98A mutation significantly reduced catalytic activity and altered pH-dependence. But since
H98A I-PpoI showed residual activity, His78, imidazole, or its surrounding waters may still serve
as alternative general bases for accepting the proton, similar to Pol η, in which primer 3'-OH
deprotonation can occur through multiple pathways [50]. However, the order of events regarding
metal ion binding, water deprotonation, and nucleophilic attack remains unanswered. Based on
our *in crystallo* observations, water deprotonation and metal ion binding appeared to be highly
correlated (Fig 6A). Lowering the pH not only reduced reaction rate but also slowed metal ion
binding (Fig. 4B, C). Moreover, the metal ion was not observed *in crystallo* when His98 was
removed (Fig. 5D). As there is no direct interaction between His98 and the Me^{2+} binding site, the
divalent metal ion may be sensitive to the charge potential of the substrate scissile phosphate,
which may be indirectly affected by the deprotonated state of the nucleophilic water. Conversely,
binding of the divalent metal ion may alter the local electrostatic environment and affect His98

deprotonation. Consistently, previous molecular dynamics simulation of Cas9 has suggested that
the histidine pKa is highly sensitive to active site changes [69]. Without a proper proton acceptor,
the metal ion may be prone for dissociation without the reaction proceeding, and thus stable Mg^{2+}
binding was not observed *in crystallo* without His98 (Fig. 6B). On the other hand, optimal
alignment of the metal ion and WatN within the active site, labeled as metal-binding state, leads
to irreversible bond breakage (Fig. 6A). In summary, our experimental observations suggest a
concerted mechanism for one-metal-ion promoted DNA hydrolysis, offering guidance for future
computational analysis of enzyme catalysis [69] and the rational design and engineering of
nucleases [70] for biotechnological and biomedical applications.
338

340   **Fig. 6. Catalytic model of His-Me nuclease DNA hydrolysis.**
341   Proposed model of His-Me nuclease DNA hydrolysis in which Me^{2+} binding, proton transfer, and
342   nucleophilic attack are concerted (solid arrow) in the presence of the primary proton acceptor in
343   (**A**) versus unfavored (dashed arrows) in the absence of the primary proton acceptor in (**B**).
344

Materials and Methods
Protein expression and purification
WT, His78Ala, His98Ala, H78A/H98A *physarum polycephalum* I-PpoI (residues 1-162) were
cloned into a modified pET28p vector with a N-terminal 6-histidine tag and a PreScission
Protease cleavage site. For protein expression, this I-PpoI plasmid was transformed into BL21
DE3 *E. coli* cells, which were grown in a buffer that contained (10 g/L glucose, 40 g/L α-lactose,
10% glycerol) for 24 hours (20˚C). The cell paste was collected *via* centrifugation and re-
suspended in a buffer that contained 50 mM Tris (pH 7.5), 1 M NaCl, 1mM MgCl$_2$, 10 mM
imidazole, 2 mM ß-mercaptoethanol (BME), and 5% glycerol. After sonication, I-PpoI was
loaded onto a HisTrap HP column (GE Healthcare), which was pre-equilibrated with a buffer that
contained 50 mM Tris (pH 7.5), 1 M NaCl, 1mM MgCl$_2$, 10 mM imidazole, 2 mM ß-
mercaptoethanol (BME), and 5% glycerol. The column was washed with 300 mL of buffer to
remove non-specific bound proteins and was eluted with buffer that contained 50 mM Tris (pH
7.5), 1 M NaCl, 1mM MgCl$_2$, 300 mM imidazole, and 2 mM ß-mercaptoethanol (BME). The
eluted I-PpoI was incubated with PreScission Protease to cleave the N-terminal 6-histidine-tag.
Afterwards, I-PpoI was desalted to
50 mM Tris (pH 7.5), 167 mM NaCl, 1mM MgCl$_2$, 2 mM ß-mercaptoethanol (BME), and 5%
glycerol and was loaded onto a Heparin column (GE Healthcare) equilibrated with 50 mM Tris
(pH 7.5) and 167 mM NaCl. The protein was eluted with an increasing salt (NaCl) gradient,
concentrated, and stored at 40% glycerol at -80˚C.
365

DNA hydrolysis assays

DNA hydrolysis activity of varying time was assayed by the following: The reaction mixture contained 100 nM WT I-PpoI, 50 mM NaCl, 100 mM Tris (pH7.5), 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 µM EDTA, and 4% glycerol. The hydrolysis assays were executed using a palindromic 5′-fluorescein- labelled DNA duplex (5′-TTG ACT CTC TTA AGA GAG TCA-3′). Reactions were initiated by adding 10 mM $MgCl_2$ to the reaction mixture for 0-1 hr at 37 °C and were stopped by mixing with equal volume of a quench buffer, which contain 80% formamide, 100 mM EDTA (pH 8.0), 0.2 mg/ml xylene cyanol, and 0.2 mg/ml bromophenol.

The DNA hydrolysis activity at different pH was assayed by the following: The reaction mixture contained 100-3000 nM WT, H98A, and H78A/H98A I-PpoI, 50 mM NaCl, 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 µM EDTA, and 4% glycerol. Reactions were initiated by adding 50 mM MES (pH5.5-6.5), 50 mM HEPES (pH6.5-7.5), or 50 mM Tris (pH7.5-9.5) together with 10 mM $MgCl_2$ to the reaction mixture for 30 min at 37 °C and were stopped by adding equal volume of quench buffer.

The DNA hydrolysis activity with different mutants was assayed by the following: The reaction mixture contained 50 mM NaCl, 100 mM Tris (pH7.5), 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 mM $MgCl_2$, 10 µM EDTA, and 4% glycerol. Reactions were initiated by adding 0-3000 nM of WT, H78A, H98A, H78A/H98A I-PpoI to the reaction mixture for 15 s or 1 h at 37 °C and stopped by adding equal volume of quench buffer.

The DNA hydrolysis activity with different metal ions was assayed by the following: The reaction mixture contained 100 nM WT I-PpoI, 50 mM NaCl, 100 mM Tris (pH7.5), 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 µM EDTA, and 4% glycerol. Reactions were initiated by adding 10 mM of $MgCl_2$, $MnCl_2$, $CaCl_2$, $NiCl_2$, and $ZnCl_2$ to the reaction mixture for 15 s at 37 °C and were stopped by adding equal volume of quench buffer.

The DNA hydrolysis activity with Tl^+ or additional Na^+ was assayed by the following: The reaction mixture contained 100-3000 nM WT I-PpoI, 100 mM Tris (pH7.5), 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 µM EDTA, and 4% glycerol. Reactions were conducted at 37 °C for 30 min by adding 0-350 mM TlCl or NaCl together with 10 mM $MgCl_2$ to the reaction mixture and were stopped by adding equal volume of quench buffer.

The DNA hydrolysis activity with imidazole was assayed by the following: The reaction mixture contained 100-3000 nM H98A and H78A/H98A I-PpoI, 50 mM NaCl, 100 mM Tris (pH7.5), 1.5 mM DTT, 0.05 mg/mL BSA, 50 nM DNA, 10 µM EDTA, and 4% glycerol. Reactions were conducted at 37 °C for 30 min by adding 0-100 mM imidazole together with 10 mM $MgCl_2$ to the reaction mixture and were stopped by adding equal volume of quench buffer.

For all reactions, after heating the quenched reaction mix to 97 °C for 5 min and immediately placing on ice, reaction products were resolved on 22.5% polyacrylamide urea gels. The gels were visualized by a Sapphire Biomolecular Imager and quantified using the built-in software. Quantification of percentage cleaved and graphic representation were executed by Graph Prism.

Crystallization

WT or H98A I-PpoI in a buffer containing 20 mM Tris 7.5, 300 mM NaCl, 3 mM DTT, and 0.1 mM EDTA was added with (5′-TTG ACT CTC TTA AGA GAG TCA-3′) DNA at a molar molar of 1:1.5 for I-PpoI and DNA and added with 3-folds volume of buffer that contained 20 mM Tris 7.5, 3 mM DTT, and 0.1 mM EDTA. This I-PpoI-DNA complex was then cleaned with a

Superdex 200 10/300 GL column (GE Healthcare) with a buffer that contained 20 mM Tris 7.5, 150 mM NaCl, 3 mM DTT, and 0.1 mM EDTA. The I-PpoI-DNA complex was concentrated to 2.8 mg/mL I-PpoI (confirmed by Bradford assay). All crystals were obtained using the hanging-drop vapour-diffusion method against a reservoir solution containing 0.1 M MES (pH 6.0), 0.2 M sodium malonate, and 20% (w/v) PEG3350 at room temperature within 4 days.

To identify the monovalent Me^+ species that binds during the pre-reaction state, WT I-PpoI crystals were transferred and incubated in a buffer containing 0.1M MES (pH 6.0), 70 mM thallium acetate and 20% (w/v) PEG3350 for 30 min. Afterwards, the crystals were quickly dipped in a cryo-solution supplemented with 20% (w/v) glycerol and flash- cooled in liquid nitrogen.

Chemical reaction *in crystallo*
The WT I-PpoI crystals were first transferred and incubated in a pre-reaction buffer containing 0.1M MES (pH 6.0 or 7.0) or 0.1M Tris (pH 8.0), 0.2 M NaCl, and 20% (w/v) PEG3350 for 30 min. The chemical reaction was initiated by transferring the crystals into a reaction buffer containing 0.1M MES (pH 6.0 or 7.0) or 0.1M Tris (pH 8.0), 0.2 M NaCl, and 20% (w/v) PEG3350, and 500 µM $MgCl_2$ or $MnCl_2$. After incubation for a desired time period, the crystals were quickly dipped in a cryo-solution supplemented with 20% (w/v) glycerol and flash- cooled in liquid nitrogen.

To observe any additional Me^{2+} binding sites during DNA hydrolysis, WT I-PpoI crystals were first transferred and incubated in a pre-reaction buffer containing 0.1M MES (pH 6.0), 0.2 M NaCl, and 20% (w/v) PEG3350 for 30 min. The chemical reaction was initiated by transferring the crystals into a reaction buffer containing 0.1M MES (pH 6.0), 0.2 M NaCl, and 20% (w/v) PEG3350, and 200 mM $MnCl_2$. After incubation for 600 s, the crystals were quickly dipped in a cryo-solution supplemented with 20% (w/v) glycerol and flash- cooled in liquid nitrogen.

The metal ion soaking experiments with His98Ala I-PpoI were performed following the similar protocol as that of WT I-PpoI. His98Ala I-PpoI crystals were first incubated in a pre-reaction buffer containing 0.1M MES 7.0, 0.2 M NaCl, and 20% (w/v) PEG3350 for 30 min, followed by 1800s incubation in a reaction buffer containing 0.1M MES 7.0, 0.2 M NaCl, and 20% (w/v) PEG3350, and 1 mM $MnCl_2$. To observe whether soaking in imidazole can initiate the reaction in the absence of His98, His98Ala I-PpoI crystals were first transferred and incubated in a buffer containing 0.1M MES (pH 6.0), 0.2 M NaCl, 1 mM $MnCl_2$, and 20% (w/v) PEG3350 for 30 min. The crystals were then transferred into a reaction buffer containing 0.1M MES (pH 6.0), 0.2 M NaCl, 1 mM $MgCl_2$ or $MnCl_2$, and 100 mM imidazole, and 20% (w/v) PEG3350. After incubation for a desired time period, the crystals were quickly dipped in a cryo-solution supplemented with 20% (w/v) glycerol and flash- cooled in liquid nitrogen.

Data collection and Refinement
Diffraction data were collected at 100 K on LS-CAT beam lines 21-D-D, 21-ID-F, and 21-ID-G at 1.1 Å or 0.97 Å at the Advanced Photon Source (Argonne National Laboratory) or beamlines 5.0.3. at 0.97 Å at ALS. Data were indexed in space group P3$_1$21, scaled with XSCALE and reduced using XDS [71]. Isomorphous I-PpoI structures with Na$^+$ PDB ID 1CZ0 was used as initial models for refinement using PHENIX [72] and COOT [73].

Occupancies were assigned for the reaction product until there were no significant F_o-F_c peaks. Occupancies were assigned for the metal ions, following the previous protocol [46] until 1) there were no significant F_o-F_c peaks, 2) the B value had roughly similar values to its ligand 3) it

matched the occupancy of the reaction product. For the structures in which some F_o-F_c peaks were present around the Me^{2+} binding sites or reaction product, no change in the assigned occupancy was executed when a 10% change in occupancy (e.g. 100% to 90%) failed to significantly change the intensity of the F_o-F_c peaks. Source data of the electron densities in r.m.s. density are provided as a Source Data file. Each structure was refined to the highest resolution data collected, which ranged between 1.42-2.2 Å. Software applications used in this project were compiled and configured by SBGrid [74]. Source data of data collection and refinement statistics are summarized in **Table S1A-E**. All structural figures were drawn using PyMOL (http:/ /www.pymol.org).

Calculation of electron density

Electron density (r.m.s.d) from the F_o-F_c map of the product phosphate and Me^{2+} were calculated by running a round of B-factor refinement in PHENIX after omitting the reaction product phosphate and Me^{2+} atoms from phase calculation in COOT. All structures from the same experiment (Mg^{2+}-pH 6.0, Mg^{2+}-pH 7.0, Mg^{2+}-pH 8.0, Mn^{2+}-pH 6.0) were refined to the lowest resolution in the same experiment group (1.80 Å for Mg^{2+}-pH 6.0, 1.59 Å for Mg^{2+}-pH 7.0, 1.70 Å for Mg^{2+}-pH 8.0, and 1.80 Å for Mn^{2+}-pH 6.0).

References

Kao, H.-I. & Bambara, R. A. The Protein Components and Mechanism of Eukaryotic Okazaki Fragment Maturation. *Critical Reviews in Biochemistry and Molecular Biology* **38**, 433-452, doi:10.1080/10409230390259382 (2003).

Shen, B. *et al.* Multiple but dissectible functions of FEN-1 nucleases in nucleic acid processing, genome stability and diseases. *Bioessays* **27**, 717-729, doi:10.1002/bies.20255 (2005).

Marti, T. M. & Fleck, O. DNA repair nucleases. *Cellular and Molecular Life Sciences CMLS* **61**, 336-354, doi:10.1007/s00018-003-3223-4 (2004).

Mimitou, E. P. & Symington, L. S. DNA end resection: many nucleases make light work. *DNA Repair (Amst)* **8**, 983-995, doi:10.1016/j.dnarep.2009.04.017 (2009).

Patel, A. A. & Steitz, J. A. Splicing double: insights from the second spliceosome. *Nature Reviews Molecular Cell Biology* **4**, 960-970, doi:10.1038/nrm1259 (2003).

Abelson, J., Trotta, C. R. & Li, H. tRNA splicing. *J Biol Chem* **273**, 12685-12688, doi:10.1074/jbc.273.21.12685 (1998).

Chu, C. Y. & Rana, T. M. Small RNAs: regulators and guardians of the genome. *J Cell Physiol* **213**, 412-419, doi:10.1002/jcp.21230 (2007).

Moore, M. J. & Proudfoot, N. J. Pre-mRNA processing reaches back to transcription and ahead to translation. *Cell* **136**, 688-700, doi:10.1016/j.cell.2009.02.001 (2009).

James, R., Kleanthous, C. & Moore, G. R. The biology of E colicins: paradigms and paradoxes. *Microbiology (Reading)* **142 (Pt 7)**, 1569-1580, doi:10.1099/13500872-142-7-1569 (1996).

Sorek, R., Kunin, V. & Hugenholtz, P. CRISPR — a widespread system that provides acquired resistance against phages in bacteria and archaea. *Nature Reviews Microbiology* **6**, 181-186, doi:10.1038/nrmicro1793 (2008).

Tock, M. R. & Dryden, D. T. The biology of restriction and anti-restriction. *Curr Opin Microbiol* **8**, 466-472, doi:10.1016/j.mib.2005.06.003 (2005).

Adli, M. The CRISPR tool kit for genome editing and beyond. *Nat Commun* **9**, 1911, doi:10.1038/s41467-018-04252-2 (2018).

Carroll, D. Genome Engineering with Targetable Nucleases. *Annu. Rev. Biochem.* **83**, 409-439, doi:10.1146/annurev-biochem-060713-035418 (2014).

14 Grindley, N. D. F., Whiteson, K. L. & Rice, P. A. Mechanisms of Site-Specific
Recombination. *Annu. Rev. Biochem.* **75**, 567-605,
doi:10.1146/annurev.biochem.73.011303.073908 (2006).
15 Yang, W. Nucleases: diversity of structure, function and mechanism. *Q Rev Biophys* **44**,
1-93, doi:10.1017/S0033583510000181 (2011).
16 Chen, Y., Gao, T., Wang, Y. & Yang, G. Investigating the Influence of Magnesium Ions
on p53-DNA Binding Using Atomic Force Microscopy. *Int J Mol Sci* **18**,
doi:10.3390/ijms18071585 (2017).
17 Yang, W. An equivalent metal ion in one- and two-metal-ion catalysis. *Nat Struct Mol*
Biol **15**, 1228-1231, doi:10.1038/nsmb.1502 (2008).
18 Samara, N. L. & Yang, W. Cation trafficking propels RNA hydrolysis. *Nat Struct Mol*
Biol **25**, 715-721, doi:10.1038/s41594-018-0099-4 (2018).
19 Wu, J., Samara, N. L., Kuraoka, I. & Yang, W. Evolution of Inosine-Specific
Endonuclease V from Bacterial DNase to Eukaryotic RNase. *Mol Cell* **76**, 44-56.e43,
doi:10.1016/j.molcel.2019.06.046 (2019).
20 Freudenthal, B. D., Beard, W. A., Cuneo, M. J., Dyrkheeva, N. S. & Wilson, S. H.
Capturing snapshots of APE1 processing DNA damage. *Nat Struct Mol Biol* **22**, 924-931,
doi:10.1038/nsmb.3105 (2015).
21 Whitaker, A. M., Flynn, T. S. & Freudenthal, B. D. Molecular snapshots of APE1
proofreading mismatches and removing DNA damage. *Nat Commun* **9**, 399,
doi:10.1038/s41467-017-02175-y (2018).
22 Dupureur, C. M. One is enough: insights into the two-metal ion nuclease mechanism from
global analysis and computational studies. *Metallomics* **2**, 609-620,
doi:10.1039/C0MT00013B (2010).
23 Garcin, E. D. *et al.* DNA apurinic-apyrimidinic site binding and excision by endonuclease
IV. *Nat Struct Mol Biol* **15**, 515-522, doi:10.1038/nsmb.1414 (2008).
24 Arnoult, D. *et al.* Mitochondrial release of AIF and EndoG requires caspase activation
downstream of Bax/Bak☐mediated permeabilization. *The EMBO Journal* **22**, 4385-4399,
doi:https://doi.org/10.1093/emboj/cdg423 (2003).
25 Lin, J. L. J. *et al.* Oxidative Stress Impairs Cell Death by Repressing the Nuclease Activity
of Mitochondrial Endonuclease G. *Cell Reports* **16**, 279-287,
doi:https://doi.org/10.1016/j.celrep.2016.05.090 (2016).
26 Cheng, Y. S., Hsia, K. C., Doudeva, L. G., Chak, K. F. & Yuan, H. S. The crystal
structure of the nuclease domain of colicin E7 suggests a mechanism for binding to
double-stranded DNA by the H-N-H endonucleases. *J Mol Biol* **324**, 227-236,
doi:10.1016/s0022-2836(02)01092-6 (2002).
27 Hsia, K. C. *et al.* DNA binding and degradation by the HNH protein ColE7. *Structure* **12**,
205-214, doi:10.1016/j.str.2004.01.004 (2004).
28 Doudna, J. A. & Charpentier, E. The new frontier of genome engineering with CRISPR-
Cas9. *Science* **346**, 1258096, doi:doi:10.1126/science.1258096 (2014).
29 Ran, F. A. *et al.* Genome engineering using the CRISPR-Cas9 system. *Nature Protocols* **8**,
2281-2308, doi:10.1038/nprot.2013.143 (2013).
30 Chevalier, B. S. & Stoddard, B. L. Homing endonucleases: structural and functional
insight into the catalysts of intron/intein mobility. *Nucleic Acids Res* **29**, 3757-3774,
doi:10.1093/nar/29.18.3757 (2001).
31 Stoddard, B. L. Homing endonuclease structure and function. *Q Rev Biophys* **38**, 49-95,
doi:10.1017/S0033583505004063 (2005).
32 Wu, C. C., Lin, J. L. J. & Yuan, H. S. Structures, Mechanisms, and Functions of His-Me
Finger Nucleases. *Trends Biochem Sci* **45**, 935-946, doi:10.1016/j.tibs.2020.07.002
(2020).

Galburt, E. A. *et al.* A novel endonuclease mechanism directly visualized for I-PpoI. *Nat Struct Biol* **6**, 1096-1099, doi:10.1038/70027 (1999).

Pommer, A. J. *et al.* Mechanism and cleavage specificity of the H-N-H endonuclease colicin E911Edited by J. Karn. *Journal of Molecular Biology* **314**, 735-749, doi:https://doi.org/10.1006/jmbi.2001.5189 (2001).

Li, C. L. *et al.* DNA binding and cleavage by the periplasmic nuclease Vvn: a novel structure with a known active site. *The EMBO Journal* **22**, 4014-4025, doi:https://doi.org/10.1093/emboj/cdg377 (2003).

Shen, B. W., Landthaler, M., Shub, D. A. & Stoddard, B. L. DNA Binding and Cleavage by the HNH Homing Endonuclease I-HmuI. *Journal of Molecular Biology* **342**, 43-56, doi:https://doi.org/10.1016/j.jmb.2004.07.032 (2004).

Schwank, G. *et al.* Functional Repair of CFTR by CRISPR/Cas9 in Intestinal Stem Cell Organoids of Cystic Fibrosis Patients. *Cell Stem Cell* **13**, 653-658, doi:10.1016/j.stem.2013.11.002 (2013).

Sharma, G., Sharma, A. R., Bhattacharya, M., Lee, S.-S. & Chakraborty, C. CRISPR-Cas9: A Preclinical and Clinical Perspective for the Treatment of Human Diseases. *Molecular Therapy* **29**, 571-586, doi:10.1016/j.ymthe.2020.09.028 (2021).

Khan, F. A. *et al.* CRISPR/Cas9 therapeutics: a cure for cancer and other genetic diseases. *Oncotarget* **7**, 52541-52552, doi:10.18632/oncotarget.9646 (2016).

Sun, W. *et al.* Structures of Neisseria meningitidis Cas9 Complexes in Catalytically Poised and Anti-CRISPR-Inhibited States. *Mol Cell* **76**, 938-952.e935, doi:10.1016/j.molcel.2019.09.025 (2019).

Zhang, Y. *et al.* Catalytic-state structure and engineering of Streptococcus thermophilus Cas9. *Nature Catalysis* **3**, 813-823, doi:10.1038/s41929-020-00506-9 (2020).

Zhu, X. *et al.* Cryo-EM structures reveal coordinated domain motions that govern DNA cleavage by Cas9. *Nat Struct Mol Biol* **26**, 679-685, doi:10.1038/s41594-019-0258-2 (2019).

Das, A. *et al.* Coupled catalytic states and the role of metal coordination in Cas9. *Nature Catalysis* **6**, 969-977, doi:10.1038/s41929-023-01031-1 (2023).

Flick, K. E., Jurica, M. S., Monnat, R. J. & Stoddard, B. L. DNA binding and cleavage by the nuclear intron-encoded homing endonuclease I-PpoI. *Nature* **394**, 96-101, doi:10.1038/27952 (1998).

Freudenthal, Bret D., Beard, William A., Shock, David D. & Wilson, Samuel H. Observing a DNA Polymerase Choose Right from Wrong. *Cell* **154**, 157-168, doi:10.1016/j.cell.2013.05.048
10.1016/j.cell.2013.05.048</p> (2013).

Gao, Y. & Yang, W. Capture of a third Mg2+ is essential for catalyzing DNA synthesis. *Science* **352**, 1334-1337, doi:10.1126/science.aad9633 (2016).

Nakamura, T., Zhao, Y., Yamagata, Y., Hua, Y.-j. & Yang, W. Watching DNA polymerase η make a phosphodiester bond. *Nature* **487**, 196-201, doi:10.1038/nature11181 (2012).

Vyas, R., Reed, A. J., Tokarsky, E. J. & Suo, Z. Viewing Human DNA Polymerase β Faithfully and Unfaithfully Bypass an Oxidative Lesion by Time-Dependent Crystallography. *J. Am. Chem. Soc.* **137**, 5225-5230, doi:10.1021/jacs.5b02109 (2015).

Chim, N., Meza, R. A., Trinh, A. M., Yang, K. & Chaput, J. C. Following replicative DNA synthesis by time-resolved X-ray crystallography. *Nat Commun* **12**, 2641, doi:10.1038/s41467-021-22937-z (2021).

Gregory, M. T., Gao, Y., Cui, Q. & Yang, W. Multiple deprotonation paths of the nucleophile 3′-OH in the DNA synthesis reaction. *Proceedings of the National Academy of Sciences* **118**, e2103990118, doi:10.1073/pnas.2103990118 (2021).

Demir, M. *et al.* Structural snapshots of base excision by the cancer-associated variant MutY N146S reveal a retaining mechanism. *Nucleic Acids Research* **51**, 1034-1049, doi:10.1093/nar/gkac1246 (2023).

Deerfield, D. W., Fox, D. J., Head-Gordon, M., Hiskey, R. G. & Pedersen, L. G. Interaction of calcium and magnesium ions with malonate and the role of the waters of hydration: a quantum mechanical study. *J. Am. Chem. Soc.* **113**, 1892-1899, doi:10.1021/ja00006a004 (1991).

Cowan, J. A. Structural and catalytic chemistry of magnesium-dependent enzymes. *Biometals* **15**, 225-235, doi:10.1023/A:1016022730880 (2002).

Auffinger, P., D'Ascenzo, L. & Ennifar, E. in *The Alkali Metal Ions: Their Role for Life* (eds Astrid Sigel, Helmut Sigel, & Roland K. O. Sigel) 167-201 (Springer International Publishing, 2016).

Kiser, P. D., Lorimer, G. H. & Palczewski, K. Use of thallium to identify monovalent cation binding sites in GroEL. *Acta Crystallogr Sect F Struct Biol Cryst Commun* **65**, 967-971, doi:10.1107/s1744309109032928 (2009).

Eastberg, J. H., Eklund, J., Monnat, R. & Stoddard, B. L. Mutability of an HNH Nuclease Imidazole General Base and Exchange of a Deprotonation Mechanism. *Biochemistry* **46**, 7215-7225, doi:10.1021/bi700418d (2007).

Furuhata, Y. & Kato, Y. Asymmetric Roles of Two Histidine Residues in Streptococcus pyogenes Cas9 Catalytic Domains upon Chemical Rescue. *Biochemistry* **60**, 194-200, doi:10.1021/acs.biochem.0c00766 (2021).

Moon, A. F. *et al.* Structural insights into catalytic and substrate binding mechanisms of the strategic EndA nuclease from Streptococcus pneumoniae. *Nucleic Acids Research* **39**, 2943-2953, doi:10.1093/nar/gkq1152 (2010).

Maghsoud, Y., Jayasinghe-Arachchige, V. M., Kumari, P., Cisneros, G. A. & Liu, J. Leveraging QM/MM and Molecular Dynamics Simulations to Decipher the Reaction Mechanism of the Cas9 HNH Domain to Investigate Off-Target Effects. *Journal of Chemical Information and Modeling* **63**, 6834-6850, doi:10.1021/acs.jcim.3c01284 (2023).

Wilson, M. A. Mapping Enzyme Landscapes by Time-Resolved Crystallography with Synchrotron and X-Ray Free Electron Laser Light. *Annu Rev Biophys* **51**, 79-98, doi:10.1146/annurev-biophys-100421-110959 (2022).

Branden, G. & Neutze, R. Advances and challenges in time-resolved macromolecular crystallography. *Science* **373**, doi:10.1126/science.aba0954 (2021).

Maestre-Reyna, M. *et al.* Visualizing the DNA repair process by a photolyase at atomic resolution. *Science* **382**, eadd7795, doi:doi:10.1126/science.add7795 (2023).

Christou, N.-E. *et al.* Time-resolved crystallography captures light-driven DNA repair. *Science* **382**, 1015-1020, doi:doi:10.1126/science.adj4270 (2023).

Reed, A. J. & Suo, Z. Time-Dependent Extension from an 8-Oxoguanine Lesion by Human DNA Polymerase Beta. *J. Am. Chem. Soc.* **139**, 9684-9690, doi:10.1021/jacs.7b05048 (2017).

Chang, C., Lee Luo, C. & Gao, Y. In crystallo observation of three metal ion promoted DNA polymerase misincorporation. *Nat Commun* **13**, 2346, doi:10.1038/s41467-022-30005-3 (2022).

Nakamura, T. & Yamagata, Y. Visualization of mutagenic nucleotide processing by *Escherichia coli* MutT, a Nudix hydrolase. *Proceedings of the National Academy of Sciences* **119**, e2203118119, doi:doi:10.1073/pnas.2203118119 (2022).

Tsutakawa, S. E. *et al.* Conserved Structural Chemistry for Incision Activity in Structurally Non-homologous Apurinic/Apyrimidinic Endonuclease APE1 and

Endonuclease IV DNA Repair Enzymes*. *Journal of Biological Chemistry* **288**, 8445-8455, doi:https://doi.org/10.1074/jbc.M112.422774 (2013).

Galburt, E. A. *et al.* Conformational Changes and Cleavage by the Homing Endonuclease I-PpoI: A Critical Role for a Leucine Residue in the Active Site. *Journal of Molecular Biology* **300**, 877-887, doi:https://doi.org/10.1006/jmbi.2000.3874 (2000).

Nierzwicki, Ł. *et al.* Principles of target DNA cleavage and the role of Mg2+ in the catalysis of CRISPR-Cas9. *Nat Catal* **5**, 912-922, doi:10.1038/s41929-022-00848-6 (2022).

Ashworth, J. *et al.* Computational redesign of endonuclease DNA binding and cleavage specificity. *Nature* **441**, 656-659, doi:10.1038/nature04818 (2006).

Kabsch, W. XDS. *Acta crystallographica. Section D, Biological crystallography* **66**, 125-132, doi:10.1107/S0907444909047337 (2010).

Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221, doi:10.1107/s0907444909052925 (2010).

Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta Crystallogr D Biol Crystallogr* **66**, 486-501, doi:10.1107/s0907444910007493 (2010).

Morin, A. *et al.* Collaboration gets the most out of software. *eLife* **2**, e01456, doi:10.7554/eLife.01456 (2013).

Acknowledgments

Our sincere appreciation to the members of the Gao lab, and Drs. Phillips, Nikonowicz, and Lu who serve on CC's thesis committee. We thank the APS LS-CAT beam technicians and research scientists Drs. Anderson, Wawrzak, Brunzelle, and Focia. This research used resources of the Advanced Photon Source, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-06CH11357. Use of the LS-CAT Sector 21 was supported by the Michigan Economic Development Corporation and the Michigan Technology Tri-Corridor (Grant 085P1000817). The Berkeley Center for Structural Biology is supported in part by the Howard Hughes Medical Institute. The Advanced Light Source is a Department of Energy Office of Science User Facility under Contract No. DE-AC02-05CH11231. The Pilatus detector on 5.0.1. was funded under NIH grant S10OD021832. The ALS-ENABLE beamlines are supported in part by the National Institutes of Health, National Institute of General Medical Sciences, grant P30 GM124169.

Funding: This work is supported by CPRIT (RR190046) and the Welch Foundation (C-2033-20200401) to YG, a predoctoral fellowship from the Houston Area Molecular Biophysics Program (NIH Grant No. T32 GM008280, Program Director Dr. Theodore Wensel) to CC.

Author contributions: YG conceived the project. CC carried out all time-resolved crystallography experiments, data collection and processing. CC carried out protein purification, and CC and GZ executed protein crystallization. CC and GZ performed the biochemical assays. CC and YG wrote the manuscript.

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: The coordinates, density maps, and structure factors for all the
structures have been deposited in Protein Data Bank (PDB) under accession codes:
8VMO, 8VMP, 8VMQ, 8VMR, 8VMS, 8VMT, 8VMU, 8VMV, 8VMW, 8VMX, 8VMY,
8VMZ, 8VN0, 8VN1, 8VN2, 8VN3, 8VN4, 8VN5, 8VN6, 8VN7, 8VN8, 8VN9, 8VNA,
8VNB, 8VNC, 8VND, 8VNE, 8VNF, 8VNG, 8VNH, 8VNJ, 8VNK, 8VNL, 8VNM,
8VNN, 8VNO, 8VNP, 8VNQ, 8VNR, 8VNS, 8VNT, 8VNU. All data are available in the
main text or the supplementary materials.
720

 Supplementary Materials

Fig. S1. Overall structure and catalytic core of homing endonuclease I-PpoI.
(**A**) Homing endonuclease I-PpoI (PDB ID 1CZO) binding as a dimer to bend DNA at 55˚. The overall endonuclease is colored in pink while the DNA is colored in orange. The catalytic core comprised of an alpha helix and two beta sheets are highlighted in red. The metal ion binding site is depicted as a purple sphere while the Zn^{2+} binding sites are depicted as turquoise spheres. The beta sheets involved in DNA binding are depicted in light green. (**B**) Monomer of homing endonuclease I-PpoI (PDB ID 1CZO) (green) superimposed on top of the other I-PpoI monomer (purple) within the same unit cell, resulting in a RMSD of 0.215. (**C**), Structural comparison of the active site of WT I-PpoI (PDB ID 1CZ0) versus Leu116Ala I-PpoI (PDB ID 1EVW). The DNA fails to dock tightly towards the Mg^{2+} in the active site of Leu116Ala I-PpoI.

Fig. S2. Active sites of His-Me superfamily nucleases.
Carbon atoms of residues within the active site are colored in pink. The metal ions are depicted by green spheres while waters are depicted by red spheres.

Fig. S3. Establishing I-PpoI for *in crystallo* studies.
(**A**) Structural comparison of the active site after soaking in NaCl in purple, sodium malonate in
yellow and PDB ID 1CZ0 in green. (**B**) Tl[+] anomalous signal was detected after 0.9765 Å X-ray
diffraction at the I-PpoI metal ion binding site after 1800 s soaking in 70 mM Tl[+]. The $2F_o$-F_c map
for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ. The
anomalous map for Tl[+] was contoured at 3.0 σ. (**C**) *In crystallo* metal ion assay of 10 mM
additional metal ions on I-PpoI DNA hydrolysis. (**D**) Negative F_o-F_c peaks (red) were detected on
the leaving group side of the scissile phosphate while positive F_o-F_c peaks (green) were detected
on the nucleophile side after 600 s Mg^{2+} soaking. The $2F_o$-F_c map for Me^{2+}, DNA, waters (red
spheres), and catalytic residues (blue) was contoured at 2.0 σ. The negative F_o-F_c map for the
reactant phosphate (red) and the positive F_o-F_c map for the product phosphate (green) were
contoured at 3.0 σ.
752

Fig. S4. Additional metal ions are not required for DNA hydrolysis by I-PpoI.
(**A**) Structures of I-PpoI during *in crystallo* catalysis after 500 µM Mg^{2+} soaking for 0 s, 20 s,
40 s, 80 s, 160 s, 320 s. The F$_o$-F$_c$ omit maps for the product phosphate (green mesh) and Mg^{2+}
(purple mesh) were contoured at 3.0 σ. (**B**) Structures of I-PpoI during *in crystallo* catalysis after
500 µM Mn^{2+} soaking for 0 s, 20 s, 40 s, 80 s, 160 s, 320 s. The F$_o$-F$_c$ omit maps for the
product phosphate (green mesh) and Mn^{2+} (purple mesh) were contoured at 3.0 σ. Correlation
(R^2) between the newly formed phosphate and Mg^{2+} binding *in crystallo* at pH 7 in (**C**), pH 6 in
(**D**), and pH 8 in (**E**). (**C-E**) The points represent the mean of duplicate measurements for the
electron density of the reaction product phosphate within two I-PpoI molecules in the asymmetric
unit while the errors bars represent the standard deviation. (**F**) Mn^{2+} binding during DNA

hydrolysis as revealed by anomalous signal of Mn^{2+} after 0.9786 Å X-ray diffraction. The $2F_o$-F_c
map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ.
The anomalous map for Mn^{2+} was contoured at 3.0 σ. **(G)** Tl^+ concentration on in solution DNA
cleavage. Precipitation was detected when Tl^+ was at or greater than 150 mM. **(H)** Additional Na^+
concentration on in solution DNA cleavage. (**G, H**) The errors bars represent the standard
deviation while the points represent the mean of triplicate measurements for cleaved DNA
product.

772
Fig. S5. Speculative ligand environment of the transient Me^{2+} in I-PpoI (A) in comparison to EndoV (B) and RNaseH (C).
(**A**) The 2F$_o$-F$_c$ map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at 2.0 σ. The DNA phosphate conformation within the active site of I-PpoI is looser in comparison to that in EndoV and RNaseH to bind an additional Me^{2+}. Carbon atoms of residues within the active site are colored in pink. The Me^{2+} are depicted by green spheres while waters are depicted by red spheres. (**A, B**) the DNA reactant state is colored in yellow while the DNA product state is colored in blue.

Fig. S6. Histidine I-PpoI mutants and partial rescued cleavage activity by imidazole.

(**A**) In solution titration of imidazole on DNA cleavage activity by H98A I-PpoI. (**B**) In solution

titration of imidazole on DNA cleavage activity by H78A/H98A I-PpoI. (**A, B**) The errors bars

represent the standard deviation while the points represent the mean of duplicate measurements

for cleaved DNA product. (**C**) Structure of H98A I-PpoI active site after 1 mM Mn^{2+} soaking for

1800 s. (**D**) Structure of H98A I-PpoI active site after 100 mM imidazole for 15 hrs following 1
mM Mn^{2+} soaking for 1800 s. The anomalous map for Mn^{2+} was contoured at 2.0 σ. (**A, D**) The
$2F_o$-F_c map for Me^{2+}, DNA, waters (red spheres), and catalytic residues (blue) was contoured at
2.0 σ.
792

793
794 **Fig. S7. Active sites of Pol η in (A), RNaseH in (B), and APE1 in (C).**
795 Carbon atoms of residues within the active site are colored in pink. The Me^{2+} are depicted by
796 green spheres while waters are depicted by red spheres.
797

Table S1. Crystal Diffraction and refinement data.
(**A**) I-PpoI-DNA complex reaction with 500 μM Mg^{2+} at pH 7.0.

	PRS (pH 7.0)	10 s	20 s	40 s
PDB Code	8VMO	8VMP	8VMQ	8VMR
Data collection				
Wavelength (Å)	0.9786	0.9787	0.9787	0.9787
Space group	$P3_121$	$P3_121$	$P3_121$	$P3_121$
Cell dimensions				
a, b, c (Å)	114.083	113.58	113.58	113.58
	114.083	113.58	113.58	113.58
	88.123	87.99	87.99	87.99
$α, β, γ$ (°)	90, 90, 120	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	49.4 - 1.68	37.18 - 1.45	37.18 - 1.48	37.18 - 1.5
	(1.74 - 1.68)	(1.502 - 1.45)	(1.533 - 1.48)	(1.554 - 1.5)
R_{sym} or R_{merge}[1]	0.09259 (0.9459)	0.08503 (0.7265)	0.07876 (0.9328)	0.08952 (1.033)
$I/σI$[1]	15.60 (2.31)	13.95 (2.24)	16.47 (2.20)	14.32 (1.95)
$CC^{1/2}$ [1]	0.999 (0.838)	0.998 (0.836)	0.999 (0.803)	0.999 (0.814)
Completeness (%)	99.99 (100.00)	99.50 (98.81)	99.90 (99.21)	99.97 (99.86)
Redundancy[1]	11.0 (10.9)	9.0 (8.3)	10.9 (10.1)	10.9 (10.2)
No. unique reflections[1]	75539 (7463)	115178 (11274)	108855 (10719)	104681 (10370)
Refinement				
Me_1	1.0 Na^+	1.0 Na^+	0.1 Mg^{2+}	0.2 Mg^{2+}
			0.9 Na^+	0.8 Na^+
Me_2	1.0 Na^+	1.0 Na^+	1.0 Na^+	1.0 Na^+
$Product_1$	-	-	0.1	0.2
$Product_2$	-	-	-	-
B-factors				
Me_1/Lig_1 [2]	20.7/22.0	15.3/16.6	15.9/17.5	15.4/17.2
Me_2/Lig_2 [2]	16.7/21.0	15.3/16.3	15.2/17.0	15.5/17.5
Protein	24.69	19.91	20.18	20.6
DNA	27.55	22.71	22.55	23.27
Ligand	19.88	15.38	15.64	15.79
Water	31.07	25.52	26.83	26.55
Resolution (Å)	1.68	1.45	1.48	1.5
No. reflections	75534 (7463)	115174 (11274)	108850 (10719)	104674 (10370)
R_{work}/R_{free}	0.17/0.19	0.18/0.19	0.17/0.19	0.18/0.19
Wilson B	22.59	17.58	17.71	17.79
Ramachandran				
Favored (%)	99.06	99.69	99.38	100
Outlier (%)	0	0	0	0
R.m.s. deviations				
Bond lengths (Å)	0.009	0.008	0.01	0.008
Bond angles (°)	1.08	1.07	1.19	1.09

[1]Data in the highest resolution shell is shown in the parenthesis.
[2]B-factor of metal ions and their protein nucleotide ligands.

	80 s	160 s	320 s	600 s
PDB Code	8VMS	8VMT	8VMU	8VMV
Data collection				
Wavelength (Å)	0.9787	0.9787	0.9787	0.9787
Space group	$P3_121$	$P3_121$	$P3_121$	$P3_121$
Cell dimensions				
a, b, c (Å)	113.58	113.58	113.58	113.58
	113.58	113.58	113.58	113.58
	87.99	87.99	87.99	87.99
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	37.18 - 1.42	42.93 - 1.481	37.18 - 1.52	37.18 - 1.59
	(1.471 - 1.42)	(1.534 - 1.481)	(1.574 - 1.52)	(1.574 - 1.52)
R_{sym} or R_{merge}[1]	0.09766 (0.8326)	0.07917 (0.9958)	0.08757 (0.8598)	0.1071 (0.8005)
$I/\sigma I$[1]	12.89 (1.95)	17.59 (2.49)	14.07 (2.11)	12.43 (2.24)
$CC^{1/2}$ [1]	0.998 (0.82)	0.999 (0.811)	0.999 (0.851)	0.998 (0.869)
Completeness (%)	99.89 (99.08)	99.98 (99.99)	99.91 (99.37)	99.97 (100.00)
Redundancy[1]	10.8 (9.9)	11.0 (10.6)	10.9 (10.7)	11.0 (11.1)
No. unique reflections[1]	123050 (12056)	108688 (10781)	100572 (9925)	88017 (8714)
Refinement				
Me_1	0.25 Mg^{2+}	0.5 Mg^{2+}	0.7 Mg^{2+}	1.0 Mg^{2+}
	0.75 Na^+	0.5 Na^+	0.3 Na^+	
Me_2	0.2 Mg^{2+}	0.4 Mg^{2+}	0.6 Mg^{2+}	0.9 Mg^{2+}
	0.8 Na^+	0.6 Na^+	0.4 Na^+	0.1 Na^+
$Product_1$	0.25	0.5	0.7	1.0
$Product_2$	0.2	0.4	0.6	0.9
B-factors				
Me_1/Lig_1 [2]	15.4/17.8	15.1/18.0	15.0/17.6	13.8/15.9
Me_2/Lig_2 [2]	15.8/17.5	15.8/17.5	16.3/17.4	14.4/15.7
Protein	20.05	20.07	20.42	19.41
DNA	22.25	22.28	23.31	22.02
Ligand	15.54	15.38	15.93	14.57
Water	26.41	26.15	26.35	24.89
Resolution (Å)	1.42	1.48	1.52	1.52
No. reflections	123047 (12056)	108678 (10781)	100559 (9925)	88005 (8714)
R_{work}/R_{free}	0.18/0.19	0.18/0.20	0.18/0.19	0.18/0.20
Wilson B	17.52	17.17	17.21	16.85
Ramachandran				
Favored (%)	99.69	99.69	99.38	99.69
Outlier (%)	0	0	0	0
R.m.s. deviations				
Bond lengths (Å)	0.008	0.009	0.009	0.022
Bond angles (°)	1.06	1.1	1.12	1.55

303   [1]Data in the highest resolution shell is shown in the parenthesis.

304   [2]B-factor of metal ions and their protein nucleotide ligands.

(**B**) I-PpoI-DNA complex reaction with 500 μM Mg^{2+} at pH 6.0.

ph6	PRS (pH 6.0)	10 s	20 s
PDB Code	8VMW	8VMX	8VMY
Data collection			
Wavelength (Å)	0.9765	0.9765	0.9765
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.88	113.88	113.88
	113.88	113.88	113.88
	88.22	88.22	88.22
$α, β, γ$ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	47.84 - 1.6	47.84 - 1.45	43.04 - 1.53
	(1.657 - 1.6)	(1.502 - 1.45)	(1.585 - 1.53)
R_{sym} or R_{merge}[1]	0.06892 (0.8392)	0.06899 (0.8012)	0.08801 (0.7433)
$I/σI$[1]	21.67 (2.41)	20.39 (2.42)	14.38 (1.81)
$CC^{1/2}$ [1]	0.999 (0.829)	0.999 (0.826)	0.999 (0.845)
Completeness (%)	99.88 (99.99)	99.95 (99.66)	99.74 (97.54)
Redundancy[1]	9.9 (9.6)	9.9 (9.3)	9.7 (9.2)
No. unique reflections[1]	86963 (8596)	116644 (11543)	99254 (9632)
Refinement			
Me_1	1.0 Na^+	1.0 Na^+	0.05 Mg^{2+}
			0.95 Na^+
Me_2	1.0 Na^+	1.0 Na^+	1.0 Na^+
$Product_1$	-	-	0.05
$Product_2$	-	-	-
B-factors			
Me_1/Lig_1 [2]	14.5/16.6	13.1/14.7	13.0/13.9
Me_2/Lig_2 [2]	115.0/15.7	13.1/14.2	13.2/14.8
Protein	18.25	17.45	17.75
DNA	21.07	20.15	20.36
Ligand	14.37	13.05	13.2
Water	24.16	23.17	23.99
Resolution (Å)	1.6	1.45	1.53
No. reflections	86958 (8596)	116640 (11543)	99251 (9632)
R_{work}/R_{free}	0.18/0.20	0.18/0.18	0.18/0.19
Wilson B	15.76	15.32	15.54
Ramachandran			
Favored (%)	99.06	98.75	99.06
Outlier (%)	0	0	0
R.m.s. deviations			
Bond lengths (Å)	0.008	0.008	0.01
Bond angles (°)	1.11	1.1	1.16

 [1]Data in the highest resolution shell is shown in the parenthesis.

 [2]B-factor of metal ions and their protein nucleotide ligands.

	40 s	80 s	160 s
PDB Code	8VMZ	8VN0	8VN1
Data collection			
Wavelength (Å)	0.9765	0.9765	0.9765
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.88	113.88	113.88
	113.88	113.88	113.88
	88.22	88.22	88.22
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	43.04 - 1.57	43.04 - 1.6	43.04 - 1.79
	(1.626 - 1.57)	(1.657 - 1.6)	(1.854 - 1.79)
R_{sym} or R_{merge}[1]	0.06753 (0.8942)	0.1128 (0.8972)	0.1035 (0.8058)
$I/\sigma I$[1]	22.77 (2.47)	12.65 (2.74)	16.88 (2.64)
$CC^{1/2}$[1]	1 (0.824)	0.997 (0.816)	0.999 (0.849)
Completeness (%)	99.64 (96.64)	99.96 (99.98)	99.88 (99.48)
Redundancy[1]	10.0 (9.6)	9.9 (9.7)	9.9 (8.9)
No. unique reflections[1]	91821 (8875)	87042 (8595)	62370 (6119)
Refinement			
Me_1	0.1 Mg^{2+}	0.2 Mg^{2+}	0.3 Mg^{2+}
	0.9 Na^+	0.8 Na^+	0.7 Na^+
Me_2	0.1 Mg^{2+}	0.2 Mg^{2+}	0.3 Mg^{2+}
	0.9 Na^+	0.8 Na^+	0.7 Na^+
$Product_1$	0.1	0.2	0.3
$Product_2$	0.1	0.2	0.3
B-factors			
Me_1/Lig_1 [2]	12.9/14.8	13.0/14.9	14.7/16.6
Me_2/Lig_2 [2]	13.0/14.5	12.9/14.0	14.2/15.1
Protein	17.67	17.32	17.42
DNA	19.82	19.28	20.43
Ligand	13.12	12.76	14
Water	23.98	24.15	23.84
Resolution (Å)	1.57	1.6	1.79
No. reflections	91816 (8875)	87034 (8595)	62365 (6118)
R_{work}/R_{free}	0.18/0.20	0.18/0.20	0.18/0.20
Wilson B	14.96	14.21	16.34
Ramachandran			
Favored (%)	99.38	99.38	99.38
Outlier (%)	0	0	0
R.m.s. deviations			
Bond lengths (Å)	0.008	0.009	0.008
Bond angles (°)	1.14	1.13	1.09

[1] Data in the highest resolution shell is shown in the parenthesis.

[2] B-factor of metal ions and their protein nucleotide ligands.

	320 s	600 s	1200 s
PDB Code	8VN2	8VN3	8VN4
Data collection			
Wavelength (Å)	0.9765	0.9765	0.9765
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.88	113.88	113.88
	113.88	113.88	113.88
	88.22	88.22	88.22
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	43.04 - 1.63	43.04 - 1.63	43.04 - 1.75
	(1.688 - 1.63)	(1.688 - 1.63)	(1.813 - 1.75)
R_{sym} or R_{merge}[1]	0.09884 (0.8855)	0.08108 (0.8274)	0.1042 (0.8633)
$I/\sigma I$[1]	13.94 (1.90)	19.75 (2.62)	13.09 (1.81)
$CC^{1/2}$ [1]	0.999 (0.815)	0.999 (0.844)	0.998 (0.843)
Completeness (%)	99.87 (99.42)	99.98 (99.99)	99.93 (99.73)
Redundancy[1]	9.8 (9.7)	10.0 (10.2)	9.7 (9.8)
No. unique reflections[1]	82333 (8100)	82420 (8146)	66724 (6607)
Refinement			
Me_1	0.6 Mg^{2+}	0.7 Mg^{2+}	0.95 Mg^{2+}
	0.4 Na^+	0.3 Na^+	0.05 Na^+
Me_2	0.4 Mg^{2+}	0.65 Mg^{2+}	0.95 Mg^{2+}
	0.6 Na^+	0.35 Na^+	0.05 Na^+
$Product_1$	0.6	0.65/0.05	0.80/0.15
$Product_2$	0.4	0.6/0.05	0.70/0.25
B-factors			
Me_1/Lig_1 [2]	14.3/16.0	13.9/16.2	13.5/15.2
Me_2/Lig_2 [2]	14.7/15.0	13.9/15.8	12.6/15.4
Protein	17.93	17.52	18.08
DNA	20.32	20.1	20.75
Ligand	13.85	13.49	13.19
Water	25.15	24.55	24.66
Resolution (Å)	1.63	1.63	1.75
No. reflections	82327 (8099)	82412 (8145)	66716 (6607)
R_{work}/R_{free}	0.18/0.19	0.18/0.19	0.18/0.20
Wilson B	14.97	14.9	15.67
Ramachandran			
Favored (%)	99.38	99.38	99.38
Outlier (%)	0	0	0
R.m.s. deviations			
Bond lengths (Å)	0.008	0.009	0.009
Bond angles (°)	1.17	1.16	1.13

[1]Data in the highest resolution shell is shown in the parenthesis.
[2]B-factor of metal ions and their protein nucleotide ligands.

(C) I-PpoI-DNA complex reaction with 500 μM Mg^{2+} at pH 8.0.

	PRS (pH 8.0)	10 s	20 s
PDB Code	8VN5	8VN6	8VN7
Data collection			
Wavelength (Å)	0.97872	0.9787	0.9787
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.65	113.65	113.65
	113.65	113.65	113.65
	88.31	88.31	88.31
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	47.79 - 1.653	42.99 - 1.541	42.99 - 1.67
	(1.712 - 1.653)	(1.596 - 1.541)	(1.73 - 1.67)
R_{sym} or R_{merge}[1]	0.0808 (1.057)	0.07081 (0.9637)	0.08404 (1.07)
$I/\sigma I$[1]	17.76 (2.20)	20.06 (2.26)	16.93 (2.14)
$CC^{1/2}$ [1]	0.999 (0.822)	0.999 (0.806)	0.999 (0.811)
Completeness (%)	99.69 (97.12)	99.70 (97.14)	99.58 (95.98)
Redundancy[1]	11.1 (10.8)	11.0 (10.8)	11.1 (10.6)
No. unique reflections[1]	78663 (7600)	96789 (9311)	76126 (7246)
Refinement			
Me_1	1.0 Na^+	0.15 Mg^{2+}	0.35 Mg^{2+}
		0.85 Na^+	0.65 Na^+
Me_2	1.0 Na^+	0.15 Mg^{2+}	0.25 Mg^{2+}
		0.85 Na^+	0.75 Na^+
$Product_1$	-	0.15	0.35
$Product_2$	-	0.15	0.25
B-factors			
Me_1/Lig_1 [2]	17.3/19.3	16.7/18.6	16.3/18.4
Me_2/Lig_2 [2]	17.1/18.4	16.9/18.3	16.6/18.4
Protein	21.73	21.68	21.36
DNA	24.16	23.65	23.57
Ligand	17.09	16.87	16.49
Water	28.9	28.62	28.61
Resolution (Å)	1.65	1.54	1.67
No. reflections	78651 (7597)	96776 (9311)	76108 (7241)
R_{work}/R_{free}	0.17/0.19	0.18/0.20	0.18/0.20
Wilson B	18.36	19.47	18.36
Ramachandran			
Favored (%)	99.38	99.06	99.69
Outlier (%)	0	0.31	0
R.m.s. deviations			
Bond lengths (Å)	0.008	0.008	0.009
Bond angles (°)	1.12	1.1	1.11

 [1]Data in the highest resolution shell is shown in the parenthesis.
 [2]B-factor of metal ions and their protein nucleotide ligands.

	40 s	80 s	160 s
PDB Code	8VN8	8VN9	8VNA
Data collection			
Wavelength (Å)	0.9787	0.9787	0.9787
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.65	113.65	113.65
	113.65	113.65	113.65
	88.31	88.31	88.31
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	47.79 - 1.6	42.99 - 1.69	34.87 - 1.543
	(1.657 - 1.6)	(1.75 - 1.69)	(1.598 - 1.543)
R_{sym} or R_{merge}[1]	0.08194 (1.012)	0.08947 (0.9124)	0.08164 (0.9551)
$I/\sigma I$[1]	17.51 (2.23)	15.59 (2.16)	16.71 (2.22)
$CC^{1/2}$ [1]	0.999 (0.79)	0.999 (0.845)	0.999 (0.818)
Completeness (%)	99.74 (97.60)	99.72 (97.37)	99.98 (99.98)
Redundancy[1]	11.0 (10.8)	11.0 (10.5)	11.0 (11.0)
No. unique reflections[1]	86597 (8362)	73645 (7158)	96719 (9584)
Refinement			
Me_1	0.45 Mg^{2+}	0.65 Mg^{2+}	0.95 Mg^{2+}
	0.55 Na^+	0.35 Na^+	0.05 Na^+
Me_2	0.3 Mg^{2+}	0.5 Mg^{2+}	0.7 Mg^{2+}
	0.7 Na^+	0.5 Na^+	0.3 Na^+
$Product_1$	0.45	0.65	0.95
$Product_2$	0.3	0.5	0.7
B-factors			
Me_1/Lig_1 [2]	16.8/19.4	16.7/19.0	16.3/18.3
Me_2/Lig_2 [2]	17.3/18.5	17.7/18.9	16.7/18.6
Protein	22.2	22.85	22.03
DNA	24.43	25.14	24.45
Ligand	17.18	17.61	16.84
Water	29.42	29.72	29.13
Resolution (Å)	1.6	1.69	1.54
No. reflections	86585 (8362)	73632 (7154)	96712 (9584)
R_{work}/R_{free}	0.18/0.19	0.18/0.19	0.18/0.20
Wilson B	19.09	19.87	19.48
Ramachandran			
Favored (%)	99.69	100	99.69
Outlier (%)	0	0	0
R.m.s. deviations			
Bond lengths (Å)	0.009	0.009	0.009
Bond angles (°)	1.15	1.13	1.17

320 [1]Data in the highest resolution shell is shown in the parenthesis.

321 [2]B-factor of metal ions and their protein nucleotide ligands.

	240 s	320 s	600 s
PDB Code	8VNB	8VNC	8VND
Data collection			
Wavelength (Å)	0.9787	0.9787	0.9787
Space group	$P3_121$	$P3_121$	$P3_121$
Cell dimensions			
a, b, c (Å)	113.65	113.65	113.65
	113.65	113.65	113.65
	88.31	88.31	88.31
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	42.99 - 1.72	47.79 - 1.623	47.79 - 1.602
	(1.781 - 1.72)	(1.681 - 1.623)	(1.659 - 1.602)
R_{sym} or R_{merge}[1]	0.1043 (0.9116)	0.08664 (0.9454)	0.09777 (0.9793)
$I/\sigma I$[1]	14.41 (2.29)	16.52 (2.41)	14.55 (2.57)
$CC^{1/2}$ [1]	0.999 (0.844)	0.999 (0.839)	0.998 (0.827)
Completeness (%)	99.95 (99.68)	99.98 (100.00)	99.98 (100.00)
Redundancy[1]	11.0 (10.9)	11.0 (11.1)	11.2 (11.1)
No. unique reflections[1]	70052 (6923)	83221 (8204)	86506 (8558)
Refinement			
Me_1	$1.0\ Mg^{2+}$	$1.0\ Mg^{2+}$	$1.0\ Mg^{2+}$
Me_2	$1.0\ Mg^{2+}$	$1.0\ Mg^{2+}$	$1.0\ Mg^{2+}$
$Product_1$	1.0	1.0	1.0
$Product_2$	1.0	1.0	1.0
B-factors			
Me_1/Lig_1 [2]	15.3/16.0	14.3/15.4	14.3/14.9
Me_2/Lig_2 [2]	15.2/16.2	14.1/15.6	13.4/15.2
Protein	21.5	21.23	20.45
DNA	24.62	24.06	23.37
Ligand	16.41	15.79	15.13
Water	27.43	27.6	26.89
Resolution (Å)	1.72	1.62	1.6
No. reflections	70046 (6922)	83212 (8204)	86496 (8558)
R_{work}/R_{free}	0.18/0.20	0.18/0.19	0.18/0.19
Wilson B	19.61	18.78	17.95
Ramachandran			
Favored (%)	100	99.69	99.69
Outlier (%)	0	0	0
R.m.s. deviations			
Bond lengths (Å)	0.009	0.009	0.008
Bond angles (°)	1.16	1.16	1.1

323   [1]Data in the highest resolution shell is shown in the parenthesis.

324   [2]B-factor of metal ions and their protein nucleotide ligands.

 (D) I-PpoI-DNA complex reaction with 500 μM Mn^{2+} at pH 6.0.

	10 s	20 s	40 s	80 s
PDB Code	8VNE	8VNF	8VNG	8VNH
Data collection				
Wavelength (Å)	0.9786	0.9786	0.9786	0.9786
Space group	$P3_121$	$P3_121$	$P3_121$	$P3_121$
Cell dimensions				
a, b, c (Å)	114.06	114.06	114.06	114.06
	114.06	114.06	114.06	114.06
	88.02	88.02	88.02	88.02
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	32.93 - 1.57	37.33 - 1.5	34.37 - 1.6	49.39 - 1.76
	(1.626 - 1.57)	(1.554 - 1.5)	(1.657 - 1.6)	(1.823 - 1.76)
R_{sym} or R_{merge}[1]	0.07871 (0.9045)	0.08083 (1.097)	0.07564 (0.9743)	0.1045 (1.001)
$I/\sigma I$[1]	18.11 (2.53)	16.89 (2.09)	19.55 (2.40)	13.57 (2.13)
$CC^{1/2}$[1]	0.999 (0.841)	0.999 (0.821)	0.999 (0.809)	0.998 (0.832)
Completeness (%)	99.98 (100.00)	99.99 (100.00)	99.97 (100.00)	99.70 (97.24)
Redundancy[1]	11.1 (11.1)	11.0 (11.0)	11.1 (11.0)	11.0 (10.9)
No. unique reflections[1]	92218 (9154)	105592 (10465)	87153 (8627)	65495 (6311)
Refinement				
Me_1	0.05 Mn^{2+}	0.1 Mn^{2+}	0.15 Mn^{2+}	0.2 Mn^{2+}
	0.95 Na^+	0.9 Na^+	0.85 Na^+	0.8 Na^+
Me_2	0.05 Mn^{2+}	0.1 Mn^{2+}	0.15 Mn^{2+}	0.2 Mn^{2+}
	0.95 Na^+	0.9 Na^+	0.85 Na^+	0.8 Na^+
$Product_1$	0.05	0.1	0.15	0.2
$Product_2$	0.05	0.1	0.15	0.2
B-factors				
Me_1/Lig_1 [2]	15.6/18.4	16.4/18.5	16.2/18.7	14.3/17.8
Me_2/Lig_2 [2]	15.9/17.2	16.8/17.7	16.3/17.4	15.0/16.6
Protein	20.95	21.18	21.5	21.18
DNA	23.45	23.56	23.85	23.64
Ligand	16.09	16.72	16.52	15.39
Water	27.18	27.29	28.05	27.3
Resolution (Å)	1.57	1.5	1.6	1.76
No. reflections	92213 (9154)	105589 (10465)	87145 (8627)	65484 (6311)
R_{work}/R_{free}	0.18/0.20	0.18/0.20	0.18/0.19	0.18/0.21
Wilson B	18.8	18.7	18.63	18.68
Ramachandran				
Favored (%)	99.38	99.06	99.06	99.38
Outlier (%)	0	0	0	0
R.m.s. deviations				
Bond lengths (Å)	0.01	0.01	0.009	0.009
Bond angles (°)	1.17	1.23	1.14	1.13

 [1]Data in the highest resolution shell is shown in the parenthesis.
 [2]B-factor of metal ions and their protein nucleotide ligands.

	120 s	160 s	240 s	320 s
PDB Code	8VNJ	8VNK	8VNL	8VNM
Data collection				
Wavelength (Å)	0.9786	0.9786	0.9786	0.9786
Space group	$P3_121$	$P3_121$	$P3_121$	$P3_121$
Cell dimensions				
a, b, c (Å)	114.06	114.06	114.06	114.06
	114.06	114.06	114.06	114.06
	88.02	88.02	88.02	88.02
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	37.33 - 1.61	34.37 - 1.61	34.37 - 1.64	34.37 - 1.59
	(1.668 - 1.61)	(1.668 - 1.61)	(1.699 - 1.64)	(1.647 - 1.59)
R_{sym} or R_{merge}[1]	0.09422 (0.8953)	0.08262 (1.086)	0.07715 (0.814)	0.06716 (0.9081)
$I/\sigma I$[1]	15.78 (2.26)	21.04 (2.30)	17.91 (2.66)	22.99 (2.63)
$CC^{1/2}$[1]	0.999 (0.812)	0.999 (0.8)	0.999 (0.825)	1 (0.829)
Completeness (%)	99.98 (100.00)	99.97 (100.00)	99.30 (99.98)	99.97 (99.99)
Redundancy[1]	11.0 (11.1)	11.1 (11.1)	9.1 (9.0)	11.2 (11.1)
No. unique reflections[1]	85576 (8465)	85577 (8465)	80455 (8029)	88820 (8795)
Refinement				
Me_1	0.25 Mn^{2+}	0.4 Mn^{2+}	0.6 Mn^{2+}	0.8 Mn^{2+}
	0.75 Na^+	0.6 Na^+	0.4 Na^+	0.2 Na^+
Me_2	0.25 Mn^{2+}	0.45 Mn^{2+}	0.65 Mn^{2+}	0.8 Mn^{2+}
	0.75 Na^+	0.55 Na^+	0.35 Na^+	0.2 Na^+
$Product_1$	0.25	0.4	0.6	0.8
$Product_2$	0.25	0.45	0.65	0.8
B-factors				
Me_1/Lig_1[2]	16.8/19.1	15.8/18.9	15.2/17.0	15.6/17.0
Me_2/Lig_2[2]	16.6/17.7	15.5/17.9	14.4/17.1	14.3/16.4
Protein	21.3	21.4	20.27	21.02
DNA	24.16	24.16	23.05	23.84
Ligand	16.75	16.34	15.37	15.48
Water	27.58	27.91	26.79	27.97
Resolution (Å)	1.61	1.61	1.64	1.59
No. reflections	85571 (8465)	85566 (8465)	80444 (8028)	88812 (8794)
R_{work}/R_{free}	0.18/0.20	0.18/0.19	0.18/0.19	0.17/0.19
Wilson B	19.46	18.85	17.28	18.12
Ramachandran				
Favored (%)	99.38	99.38	100	100
Outlier (%)	0	0	0	0
R.m.s. deviations				
Bond lengths (Å)	0.009	0.009	0.008	0.008
Bond angles (°)	1.17	1.13	1.04	1.07

[1]Data in the highest resolution shell is shown in the parenthesis.

[2]B-factor of metal ions and their protein nucleotide ligands.

	480 s	600 s
PDB Code	8VNN	8VNO
Data collection		
Wavelength (Å)	0.9786	0.9786
Space group	*P3₁21*	*P3₁21*
Cell dimensions		
a, b, c (Å)	114.06	114.06
	114.06	114.06
	88.02	88.02
α, β, γ (°)	90, 90, 120	90, 90, 120
Resolution (Å)[1]	37.33 - 1.792	43.07 - 1.7
	(1.856 - 1.792)	(1.761 - 1.7)
R_{sym} or R_{merge}[1]	0.1312 (1.102)	0.08224 (0.9698)
$I/\sigma I$[1]	13.34 (2.57)	19.05 (2.56)
$CC^{1/2}$ [1]	0.998 (0.831)	0.999 (0.849)
Completeness (%)	99.97 (99.98)	99.97 (99.97)
Redundancy[1]	11.1 (11.1)	11.1 (11.2)
No. unique reflections[1]	62269 (6153)	72812 (7213)
Refinement		
Me_1	0.8 Mn^{2+}	0.9 Mn^{2+}
	0.2 Na^+	0.1 Na^+
Me_2	0.8 Mn^{2+}	0.9 Mn^{2+}
	0.2 Na^+	0.1 Na^+
$Product_1$	0.8	0.9
$Product_2$	0.8	0.9
B-factors		
Me_1/Lig_1 [2]	16.3/17.1	16.2/17.2
Me_2/Lig_2 [2]	14.7/16.7	15.5/17.2
Protein	22.06	21.53
DNA	25.03	24.79
Ligand	16.66	16.45
Water	28.18	27.81
Resolution (Å)	1.79	1.7
No. reflections	62260 (6152)	72799 (7211)
R_{work}/R_{free}	0.18/0.19	0.18/0.19
Wilson B	19.72	19.27
Ramachandran		
Favored (%)	100	99.69
Outlier (%)	0	0
R.m.s. deviations		
Bond lengths (Å)	0.01	0.009
Bond angles (°)	1.22	1.21

333 [1]Data in the highest resolution shell is shown in the parenthesis.
334 [2]B-factor of metal ions and their protein nucleotide ligands.
335

(**E**) Other I-PpoI-DNA complexes.

	I-PpoI 0.2 M sodium malonate	His98Ala I-PpoI 1 mM Mn^{2+} 1800 s	His98Ala I-PpoI 1 mM Mn^{2+} Imidazole, 15 h	I-PpoI 200 mM Mn^{2+} 600 s
PDB Code	8VNP	8VNQ	8VNR	8VNS
Data collection				
Wavelength (Å)	1.1272	0.9765	0.9765	0.9765
Space group	$P3_121$	$P3_121$	$P3_121$	$P3_121$
Cell dimensions				
a, b, c (Å)	113.695	114.646	114.09	117.905
	113.695	114.646	114.09	117.905
	87.994	89.057	88.412	84.743
α, β, γ (°)	90, 90, 120	90, 90, 120	90, 90, 120	90, 90, 120
Resolution (Å)[1]	34.28 - 1.79	43.36 - 1.93	43.13 - 1.98	43.73 - 2.111
	(1.854 - 1.79)	(1.999 - 1.93)	(2.051 - 1.98)	(2.186 - 2.111)
R_{sym} or R_{merge}[1]	0.05373 (0.1686)	0.21 (0.782)	0.1138 (0.8958)	0.1301 (0.6187)
$I/\sigma I$[1]	46.56 (15.80)	6.69 (1.37)	11.19 (1.62)	10.97 (2.08)
$CC^{1/2}$ [1]	1 (0.995)	0.989 (0.848)	0.998 (0.848)	0.996 (0.905)
Completeness (%)	99.91 (99.95)	99.50 (95.26)	99.74 (98.19)	99.61 (96.97)
Redundancy[1]	20.5 (20.0)	9.8 (8.9)	9.8 (9.4)	9.7 (9.5)
No. unique reflections[1]	62043 (6149)	50811 (4824)	46448 (4491)	39247 (3780)
Refinement				
Me_1	1.0 Na	1.0 Na	1.0 Na	0.7 Mn^{2+}
				0.3 Na^+
Me_2	1.0 Na	1.0 Na	1.0 Na	0.8 Mn^{2+}
				0.2 Na^+
$Product_1$	-	-	-	0.7
$Product_2$	-	-	-	0.8
B-factors				
Me_1/Lig_1 [2]	12.8/14.6	25.1/24.6	35.9/33.1	27.3/30.5
Me_2/Lig_2 [2]	12.7/13.2	22.3/22.3	34.6/32.5	29.0/29.7
Protein	16.91	29.29	36.05	36.11
DNA	20.32	32.76	39.19	42.08
Ligand	13.7	23.81	33.27	29.82
Water	22.25	32.52	39.31	33.85
Resolution (Å)	1.79	1.93	1.98	2.11
No. reflections	62036 (6149)	50810 (4824)	46440 (4491)	39239 (3780)
R_{work}/R_{free}	0.17/0.19	0.19/0.21	0.19/0.22	0.22/0.25
Wilson B	16.51	30.06	35.07	35.57
Ramachandran				
Favored (%)	99.06	99.38	99.38	99.38
Outlier (%)	0	0	0	0
R.m.s. deviations				
Bond lengths (Å)	0.007	0.01	0.009	0.01
Bond angles (°)	0.96	1.13	1.08	1.14

337 [1]Data in the highest resolution shell is shown in the parenthesis.

338 [2]B-factor of metal ions and their protein nucleotide ligands.

	I-PpoI 500 μM Mg^{2+} 1800 s	I-PpoI 70 mM Tl^+ 1800 s
PDB Code	8VNT	8VNU
Data collection		
Wavelength (Å)	1.1271	0.9765
Space group	*$P3_121$*	*$P3_121$*
Cell dimensions		
a, b, c (Å)	114.129	113.574
	114.129	113.574
	88.74	87.695
$α, β, γ$ (°)	90, 90, 120	90, 90, 120
Resolution (Å)[1]	33.02 - 1.62	47.67 - 2.2
	(1.678 - 1.62)	(2.279 - 2.2)
R_{sym} or R_{merge}[1]	0.1009 (1.138)	0.1265 (0.9511)
$I/σI$[1]	23.30 (2.78)	12.59 (2.17)
$CC^{1/2}$[1]	0.999 (0.867)	0.998 (0.837)
Completeness (%)	99.96 (99.96)	99.91 (100.00)
Redundancy[1]	20.2 (20.6)	9.9 (10.0)
No. unique reflections[1]	84795 (8426)	33498 (3304)
Refinement		
Me_1	1.0 Mg^{2+}	0.2 Tl^+ 0.9 Na^+
Me_2	1.0 Mg^{2+}	0.2 Tl^+ 0.8 Na^+
$Product_1$	0.75/0.25	0
$Product_2$	0.60/0.40	0
B-factors		
Me_1/Lig_1[2]	19.2/21.8	80/37.4
Me_2/Lig_2[2]	20.5/22.8	73.7/36.0
Protein	23.32	35.3
DNA	26.33	37.63
Ligand	19.65	59.35
Water	30.34	39.55
Resolution (Å)	1.62	2.2
No. reflections	84783 (8426)	33486 (3306)
R_{work}/R_{free}	0.18/0.20	0.18/0.22
Wilson B	21.49	36.08
Ramachandran		
Favored (%)	98.75	98.75
Outlier (%)	0	0.31
R.m.s. deviations		
Bond lengths (Å)	0.008	0.009
Bond angles (°)	1.06	1.04

340 [1]Data in the highest resolution shell is shown in the parenthesis.

341 [2]B-factor of metal ions and their protein nucleotide ligands.