Dorsal prefrontal cortex drives perseverative behavior in mice

Lebedeva A.,[1] Wang Y.,[1] Funnell L.,[2] Terry B.,[2] Oh Y. J.,[2] Miller K.,[1,3,4,*] Harris K.D.[2,*]

[1] UCL Sainsbury Wellcome Centre. [2] UCL Institute of Neurology. [3] UCL Institute of Ophthalmology
[4] Google DeepMind. * Co-senior authors

Perseveration – repeating one choice when others would generate larger rewards – is a common behavior, but neither its purpose nor neuronal mechanisms are understood. Here we demonstrate a neural correlate and causal role of dorsal prefrontal cortex (area MOs) in perseveration in mice performing a dynamic reward learning task. An auditory go cue signaled mice to turn a wheel either left or right, with the reward probability of each action switching in blocks. Mice perseverated, gaining suboptimal reward, but were faster when making repeated choices. Neuropixels recordings found neurons whose activity correlated with perseveration and predicted rapid reaction times, almost exclusively in MOs. Optogenetically inhibiting MOs reduced perseveration and slowed reaction times. In contrast, inactivating medial prefrontal cortex at choice time did not have any effect but impaired learning if delivered after reward delivery. In this task, therefore, MOs encodes a perseverative decision variable, and is necessary for mediating the effect of this decision variable on choice and reaction time.

The psychologist Edward Thorndike proposed two laws of learning (Thorndike, 1911). The "law of effect" holds that when reward follows an action, that action becomes more likely in the future. The less well-known "law of exercise" holds that just performing an action, even without reward, increases the probability this action will be performed again.

The law of effect has inspired substantial research in psychology and neuroscience. Much of this work has applied computational models inspired by reinforcement learning (RL; Sutton & Barto, 1998; Botvinick et al., 2020) to predict how reward history shapes choices, and interpret neural activity recorded during learning tasks (Averbeck & O'Doherty, 2022; Costa & Averbeck, 2020; Daw, 2011; Ito & Doya, 2011; Lee et al., 2012). This work has identified correlates of RL-related cognitive variables in frontal cortex (Barraclough et al., 2004; Daw et al., 2006,Behrens et al., 2007; Seo et al., 2007; Sul et al., 2010; Sul et al., 2011; Bari et al., 2019; Costa & Averbeck, 2020), suggesting a key role of frontal cortex in reward-guided learning and decision-making.

The law of exercise has received comparatively less attention. This law predicts "perseverative" behavior: animals will repeat even actions not followed by reward. Perseveration occurs in many species and tasks (e.g. Aarts et al., 1998; Gold et al., 2008; Lau & Glimcher, 2005; Padoa-Schioppa, 2013; Riefer et al., 2017), and can be quantitatively described by computational models (Ashby et al., 2007; Ashby et al., 2010; Miller et al., 2019; Bogacz, 2020). The benefits of perseveration to an animal are unclear. One possibility is that repeatedly performing an action makes it cognitively simpler or physically easier (Ashby et al., 2007; Gershman, 2020; Wood & Rünger, 2016), consistent with the observations that repeated actions are performed more reliably and quickly (Bertelson, 1965; Gore et al., 2002; Hikosaka et al., 1995), and that responses under time pressure are more likely to be perseverative (Hardwick et al., 2019; Lai & Gershman, 2024).

Relatively little is known about the neural mechanisms of perseveration. A first key question is to identify brain regions that correlate with, and have a causal role in, perseverative behaviors. A second key question is whether perseverative and reward-guided decision-making operate in parallel, with separate neural circuits or individual neurons devoted to implementing each, or whether they rely on partially or fully overlapping sets of circuits.

We address both questions using a dynamic reward learning task (the "probabilistic reversal learning" task; Samejima et al., 2005; Ito & Doya, 2009; Kim et al., 2009) in head-fixed mice. Choices were largely predicted by past choices, with a much smaller influence of past rewards, and could be well approximated by perseverative behavior models. Neural activity in secondary motor cortex (MOs) correlated with perseveration, including during the inter-trial period when no movements were allowed, but such correlations were largely absent from the other frontal regions. Optogenetically inactivating MOs, but not medial prefrontal cortex (mPFC), during choice attenuated both reward-guided behavior and perseveration, and also weakened the perseverative effect of the current choice on future trials. Inactivating MOs following the reward had no discernible effect, but inactivating mPFC following the reward slowed learning, if delivered for multiple trials in succession.

Together, these results indicate that, uniquely in frontal cortex, the secondary motor cortex carries a perseveration variable that is used to guide choice in our task. The perseverative learning rule may depend on neural activity in MOs, but the reward-guided learning rule seems to depend instead on activity elsewhere in the brain.

Results

Choices in a dynamic reward-learning task are driven largely by perseveration

We trained mice in a head-fixed probabilistic reversal learning task (Figure 1). The mouse was head-fixed, with forepaws on a rubber wheel, and a tube near its mouth for delivering liquid rewards (Figure 1A). To initiate each trial, the mouse held the wheel still for a 200ms Fixation Period. An auditory Go Cue signaled the mouse should turn the wheel either clockwise or counterclockwise (from the mouse's perspective; we call these left and right choices). After turning the wheel a sufficient distance, the mouse received either a liquid reward or auditory white noise indicating that no reward would be delivered. On each trial, one turn direction (Correct) led to 80% reward probability, while the other (Incorrect) led to 20% reward probability (Figure 1B). Which action (left or right) was correct remained stable for a block of trials, then reversed. The first block of each session lasted between 1 and 175 trials, with subsequent blocks lasting between 125 and 175 trials, each sampled independently from a uniform distribution (Figure 1C).

Mice performed suboptimally in the task. To show this, we compared their choices to those of an ideal observer model, which optimally infers based on history which side is most likely to be Correct, and deterministically chooses this side. The ideal observer earned reward on 77% of trials, substantially outperforming the mice at 62±0.6% (n=10 mice; mean ± S.E.). Following a block switch, mice persisted in choosing the previously-Correct option for much longer than the ideal observer, taking 11.3 ± 0.2 (n=10 mice; mean ± S.E.) vs. 2.2 trials to reach 50% accuracy, and reaching an asymptotic accuracy of 75% ± 0.01% vs 97% correct choices by 100 trials into the block (Figure 1D,E). We characterized the interplay of reward seeking and perseveration using trial-history regression (Corrado et al., 2005; Lau & Glimcher, 2005) to predict upcoming choices using the history of previous rewards, choices, and reward-choice interactions (i.e. 1 for a rewarded right choice or unrewarded left choice, -1 for an unrewarded right or rewarded left choice), which indicate the likely correct side. Plotting the weights of the past choices and the interaction between the past choices and rewards revealed that the perseverative weight of a past choice was higher than the reward-seeking weight of that choice, which was not the case for the ideal observer model (Figure 1F). Overall, these diagnostics reveal that mouse behavior is strongly perseverative in this task.

Figure 1 | Probabilistic reversal learning task. A, The mouse is head-fixed with paws on a wheel and a spout for delivering reward. **B**, To start a trial, mice must hold the wheel still for 200ms. An auditory Go Cue (0.1 s, 5 kHz) indicates trial start but provides no information about which choice will be rewarded. Feedback (reward or white noise) is given after mice turn the wheel 18mm in either direction, with no time limit. Reward probability depends on the choice, switching in blocks. **C**, Example behavioral session. **D,** learning curves aligned to block switches for mice (black), PR model (red) and ideal observer model (blue). **E**, Lengths of unrewarded trial sequences. **F**, Trial-history regression weights, predicting choice on trial N from choice, rewards, and their interactions on trials N-5 to N-1. Left: choice weights; right: interaction weights. **G**, Quality of fit of 13 probabilistic behavioral models to observed choices sequences, assessed by geometric mean likelihood on held-out trials. Boxes show distribution across sessions. Central mark: median; box edges: 25th and 75th percentiles, green +: outliers, whiskers: range excluding outliers. **H**, Internal state of the PR model fitted to the example session of Fig. 1C.

Mouse choices are well fit by a perseverative behavior model

We quantitatively summarized the interplay of reward-seeking and perseveration by fitting probabilistic cognitive models. Because it was *a priori* unclear which of the many models in the literature would provide the best fit, we evaluated the performance of 13 using cross-validated likelihood (Figure 1G). The difference in fit of the 13 models was highly significant ($p=8\times10^{-40}$, ANOVA), but there was no significant difference in the fits of the five top models, all of which captured perseverative behavior: PR model (Miller et al., 2021); differential forgetting Q-learning (Ito & Doya, 2009); ideal observer with perseveration (Beron et al., 2022); Ideal-observer Bayesian filters with perseveration for either a generic probabilistic reversal-learning task or a generic drifting reward probability task (see Methods). To further evaluate these models, we analyzed synthetic behavioral datasets generated by each model, using parameters fit to each mouse. The five winning models produced similar behavioral sequences to the mice, whether assayed by blockwise learning curves (PR Model: Figure 1D, red curves; remaining models, Supplementary Figure 1), distribution of unrewarded sequence lengths (PR model, Figure 1E; remaining models, Supplementary Figure 1), or trial-history regression weights (PR Model: Figure 1F). In contrast, the eight non-winning models showed behavior that deviates in some way from that of the mice (Supplementary Figure 1), falsifying these models (Palminteri et al., 2017). Although the five winning models were not mathematically equivalent, the choice probabilities they predicted were highly correlated (Supplementary figure 2), suggesting that they captured similar aspects of behavior.

Given that the five perseverative models were equally successful, we chose the simplest among them, which we call the PR model (perseveration/reward learning; Miller et al., 2021). Although this model's performance was not significantly higher than the other winning models', it is the simplest model which encodes reward-learning and perseverative components in separate decision variables, which will facilitate later analyses of the neural correlates of these. The PR model has two decision variables: P which follows the law of exercise and mediates perseverative behavior, and R which follows the law of effect and mediates reward-seeking behavior. On each trial, the PR model agent chooses right over left with log odds given by the sum of the two decision variables:

$$\log\left(\frac{P(C_t = +1)}{P(C_t = -1)}\right) = P_t + R_t + b \tag{3}$$

where b is a parameter governing fixed left/right bias, C_t is the choice on trial t (-1 or +1 for left or right).

The perseverative variable P is initialized to zero on the first trial, and updated according to:

$$P_{t+1} = P_t + \alpha_p\left(C_t\lambda_p - P_t\right) \tag{1}$$

where α_p is a learning rate parameter governing the extent to which an individual choice affects P, and λ_p is a magnitude parameter governing the largest possible magnitude of P. This rule therefore updates P towards $-\lambda_p$ following left choices and $+\lambda_p$ following right choices, so P can be thought of as a running average of previous choices (Figure 1H, gold line).

The reward-seeking decision variable R is likewise initialized to zero, and updated according to:

$$R_{t+1} = R_t + \alpha_r(\lambda_r C_t F_t - R_t) \tag{2}$$

where F_t is the feedback on trial t (-1 or +1 for unrewarded or rewarded), α_r and λ_r are learning rate and magnitude parameters, which can differ from the perseveration parameters α_p and λ_p. The rule updates R towards $-\lambda_r$ following rewarded left choices or unrewarded right choices, and towards $+\lambda_r$ following rewarded right choices or unrewarded left choices. Thus, R can be thought of as a running average of the choice-reward interaction, which provides an estimate of the correct choice (Figure 1H, orange line). The

Figure 2 | Responses are faster for repeated actions, where the perseverative variable has large magnitude. **A**, Same example session as Fig. 1C, with response times shown in gray. **B**, wheel position on two example trials (dashed: a right turn following LLRLR; solid: a right turn following RRRRR) with different values of the perseverative variable (P). **C**, Cumulative probability density of response times for repeat trials (red) and switch trials (blue), truncated at 4s. Thin lines, individual mice; thick line, average. **D**, Mean response time for trials with different values of perseveration variable *P* in the PR model (mean ± s.e., n=10 mice). **E**, Mean response time for trials with different values of reward seeking variable *R*.

overall magnitudes of *P* and *R* – determined by the parameters λ_p and λ_r – therefore govern the relative influence of perseveration and reward seeking on choice.

We fit the model parameters to the observed behavior of each mouse by maximum likelihood, and found that the perseverative process had more than three times the influence of the reward-seeking process (ratio of λ_p to λ_r : 3.67 ± 0.44, mean ± S.E.M., n=10 mice; λ_p greater than λ_r in 10/10 mice), and the learning rate of the perseverative process was somewhat lower than that of the reward-seeking process, meaning that it was influenced by a larger number of past trials (ratio of α_r to α_p : 1.33 ± 0.13 (mean ± S.E.M., n=10 mice, Supplementary figure 3).

Figure 3 | Example MOs neurons showing cue-period correlates of upcoming choices. A, Histological electrode track reconstruction for one example mouse. **B**, Reconstructed probe trajectories for all recordings. **C**, Top: raster plots for four example units from the probe in A, trials ordered chronologically and colored by trial type. Bottom: peristimulus time histograms for the same units. **D,** Difference in z-scored activity prior to right and left choices in an example session, for all neurons as a function of depth. Pink arrows (left): units from C. **E,** Neural preference for future choice side vs. depth. X-axis: side preference index $\frac{F_R-F_L}{F_R+F_L+0.1}$, where F_R and F_L are mean spikes count in fixation periods prior to right and left choices. Cyan curve: running standard deviation (25 units) of side preference as a function of depth.

Perseverative choices are associated with faster responses

Previous work has found that repeated choices are made faster (Bertelson, 1965; Cho et al., 2002; Hikosaka et al., 1995). To test this in our task, we quantified each trial's response time as the time between go cue

onset and when the wheel turn magnitude passed the threshold required to register a choice (Figure 2A, B). We found that response times were longer on "switch" trials, where the mouse selected a different action than on the previous trial, than on "repeat" trials, where it selected the same action (Figure 2C; switch trials 2.5±0.3 mean±s.e., repeat trials 4.5±0.4 mean±s.e., p = 0.002, signed rank test, n=10 mice). This was not limited to the switch trials themselves: response times remained slow for three trials following the switch and were also slow for trials immediately prior to the switch (Supplementary figure 4A). We confirmed that when the perseveration variable P had large absolute value, indicating a sequence of choices to the same side, response times were shorter (Pearson correlation -0.15±0.02, mean ± s.e. over n=10 mice; p = 3x10^{-4}, t-test; Figure 2D). The reward seeking variable R did not significantly correlate with response time (-0.001±0.02, mean ± s.e. over n=10 mice, p = 0.96 t-test; Figure 2E). Thus, response times in our task are shorter when mice perseverate.

Recordings across the forebrain reveal correlates of future choice specifically in MOs

To search for neural correlates of perseveration and reinforcement learning, we used Neuropixels probes (Jun et al., 2017; Steinmetz et al., 2021) to record across the frontal cortex, as well as several related brain regions, while mice performed the task (Figure 3). After automatic spike sorting (Pachitariu et al., 2016; Pachitariu et al., 2023) and manual curation, our dataset comprised 19,893 units from 42 sessions in 10 mice. We obtained a sufficient number of units and sessions for detailed analysis in six major brain regions: secondary motor area (MOs, 2,580 units in 15 sessions), medial prefrontal cortex (mPFC, 4,255 units in 14 sessions; combining prelimbic, infralimbic, and anterior cingulate), dorsomedial striatum (DMS, 1,287 units in 9 sessions), orbital areas (OFC, 2,400 units in 11 sessions; combining medial and lateral orbital), olfactory areas (OLF, 2,509 units in 15 sessions; combining olfactory bulb, olfactory nucleus, taenia tecta, dorsal peduncular area, piriform area, nucleus of the lateral olfactory tract, cortical amygdalar area, piriform-amygdalar area and postpiriform transition area), and dorsal hippocampus (DH, 2,176 units in 15 sessions; combining CA1, CA3, and dentate gyrus) (Figure 3B). Other regions recorded include lateral septal nucleus (646 units in 6 sessions), retrosplenial area (185 units in 5 sessions) and nucleus accumbens (429 units in 6 sessions) (Supplementary figure 5), but the smaller number of sessions and neurons for these latter areas precluded rigorous statistical analysis.

Consistent with previous work (Lee et al., 2012; Steinmetz et al., 2019), we found units in all brain regions whose activity correlated with choice and feedback after these events had begun. However, neurons whose activity during the fixation period predicted the upcoming choice were rarer and appeared far more common in MOs than in any other region (Figure 3C-E). A similar distinction has been observed previously in rats performing a related freely-moving task (Sul et al., 2011; Lee et al., 2012) and interpreted as indicating a specific role for MOs in value-guided action selection.

Verifying the statistical significance of a correlation between neural firing and choices requires care. Both behavioral and neural timeseries show temporal autocorrelations, including due to artifacts such as electrode drift. Commonly used statistical tests assuming that observations are statistically independent of one another can produce spuriously significant "nonsense correlations" in this situation, even in the absence of genuine correlations (Yule, 1926; Harris, 2021b; Elber-Dorozko & Loewenstein, 2018). For example, consider a unit whose rate was artifactually higher at the start of the experiment due to electrode drift, in a session that started with left choice block. A t-test would show significantly higher rates in left than right choice trials, even if the unit carried no genuine information about the choice. Furthermore, because simultaneously recorded units are correlated, tests which assume independence of units may also yield false positives. We solved this problem using a "session permutation" test (Harris, 2021b), which treats an entire recorded session as the unit of independent variability. We predict the sequence of choices

Figure 4 | Upcoming movements are encoded specifically in MOs. A, Decoding of upcoming choice and model variables from population activity. Each column shows decoding of choice (logistic regression), perseveration variable *P* (linear regression); and reward-seeking variable *R* (linear regression). X-axis: time epoch; fixation period (0-200 ms before Go Cue), and subsequent 200 ms epochs from movement onset. Black dots: mean prediction of actual value across sessions; gray violins, null distribution from session permutation. Red stars: $p < 0.01$, session permutation test. **B,** Decoding upcoming choice from fixation-period activity of equally-sized random subsamples of neurons from each region. X-axis: number of units in subsample; Y-axis, mean log-likelihood relative to permuted sessions. Red stars: MOs significantly different to other regions ($p < .01$; t-test).

in each session from the sequence of neural population activity vectors using logistic regression and compute a single test statistic summing the log likelihood of all sessions. We compare this statistic to a null distribution obtained by rerunning after permuting the session labels of the behavioral data. This therefore tests the null hypothesis that neural activity in a particular session predicts that session's choices no better than the choices of a different session, without assuming independence between trials or units within a session.

This analysis confirmed that fixation-period population activity in MOs predicted the upcoming choice. We applied this analysis to neural activity in each of our brain regions, both during the fixation period before movement had begun, and in 200ms timebins following movement initiation. Once the mouse had begun to turn the wheel, the direction of choice could robustly be decoded from neural activity in any of our brain regions, but during the fixation period, the direction of upcoming choice could be robustly decoded only from MOs; the significance of coding from other regions was at most marginal (p>0.01) (Figure 4A). This did not result from different numbers of units recorded in the different regions: repeating the analysis using an equal number of units from each area, still yielded better prediction from MOs than all other areas (Figure 4B). Thus, neural activity in MOs bears a relationship to upcoming choice that is unique among the brain regions recorded, suggesting that it may play a distinct role in the choice process.

MOs population activity during the fixation period correlated significantly with the PR model's perseveration variable P, but not the reward learning variable R. To show this we predicted each session's sequence of P or R values from the sequence of neural population activity vectors using linear regression, and summarized this prediction by averaging the mean squared error across sessions. We compared this statistic to a null distribution obtained by permuting the session labels of the behavioral data. For time epochs following movement onset, P could be decoded from population activity of most brain regions (Figure 4A, second row). However, MOs was the only region whose activity during the fixation period could robustly predict P; the significance of coding from other regions was at most marginal (p>0.01). The reward-learning variable R could not be predicted from any brain region during the fixation period and was not robustly predicted during the movement except from a few areas including MOs and orbital cortex (Figure 4A, third row). We additionally used the fact that rewards were randomly delivered to test for a causal effect of rewards on subsequent population activity. This found that reward had a significant causal effect on activity in all recorded regions, that persisted even through the subsequent trial (Supplementary Figure 6). However in MOs, the interaction of choice and feedback – which updates the reward-seeking decision variable R – had a much more transient causal effect that was no longer significant by the time of the subsequent fixation period (Supplementary Figure 7). We conclude that MOs activity in the fixation period robustly encodes the perseverative decision variable P, but has little or no coding of the reward-seeking decision variable R.

Secondary motor cortex activity reflects a perseverative decision variable

The fact that MOs population activity correlates with both upcoming choice and the P variable, which are themselves correlated, has multiple possible interpretations. First, MOs might simply encode the upcoming choice in a binary way, reflecting a decision made prior to the fixation period; since P is correlated with the choice, MOs activity would correlate with P as well. Alternatively, MOs might encode a continuous perseveration variable beyond the prediction of choice coding alone.

Examining the activity of individual MOs units revealed cells whose activity varied with P beyond their dependence on upcoming choice (Figure 5). An example MOs unit was more active on average prior to left than to right choices but showed further dependence on choice history (Figure 5A). This unit fired essentially no fixation-period spikes after a long sequence of right choices, but it was active following a sequence of left choices, even if the next choice was to the right. Similarly, the mean activity of neurons predicting either left or right choices showed a correlation with P even among trials with the same upcoming choice (Figure 5B).

Figure 5 | MOs encodes perseveration variable better than upcoming choice. A, Top, example unit whose fixation-period activity correlates with *P*. Bottom, response times in the same session. **B,** Mean activity of left- or right-preferring units during fixation period, as a function of *P* (x-axis; logistic scale) and choice direction (color). Each point corresponds to a trial, curves to a lowess smoothing of the points. **C,** Mean square error (MSE) of predicting *P, R,* or *P+R* after projecting out choice. Black line: actual value; gray histogram, null distribution from session permutation. Stars: p<.01, session permutation test for partial correlation. **D,** Response time as a function of neural choice direction prediction, obtained by logistic regression of choice direction from fixation-period population activity. Brown and green points: single trials with right or left choices; pink curve: running median.

To establish the statistical significance of these results, we used a variant of the session permutation test, designed to identify partial correlations in timeseries with correlation between trials and neurons (Harris & Yuan, 2024). This showed that there was a significant correlation between fixation-period MOs population activity and the perseverative decision variable P, even after the common effect of upcoming choice was accounted for (Figure 5C). Applying this same analysis to the reward-seeking decision variable R did not reveal a significant relationship, consistent with our earlier result that R could not be predicted during the fixation period (Figure 4A). We also asked whether MOs activity could also predict the summed decision variable $P+R$ when controlling for upcoming choice, and found that it could (Figure 5C, right). These results are therefore consistent with either the hypothesis that MOs delay-period activity reflects the perseverative decision variable specifically or that it reflects the aggregate summed decision variable. Since this aggregate variable is dominated by P (Figure 1H, Supplementary figure 3), both hypotheses indicate that MOs activity strongly encodes a perseverative decision variable, and encodes the reward-seeking variable weakly, if at all.

We hypothesized that when MOs activity during the fixation period strongly predicts a given action, such as after a long sequence of choices in the same direction, this should indicate a degree of confidence or preparation of the action, leading to shorter reaction times. This was supported by an example neuron that appeared to fire strongly during the fixation period prior to trials with short reaction times (Figure 5A). To test the hypothesis statistically, we predicted the mouse's upcoming choice from fixation-period MOs population activity during the fixation period using logistic regression. We reasoned that the weighted sum of activity found by this classifier (prior to passing through the logistic function) would form an estimate of confidence on each trial: when many MOs units are "voting" for a right or left choice, this predictor would have a large absolute value (positive or negative). Even though the predictor was trained without reference to reaction times, reactions were faster on trials when the predictor had large magnitude. (Figure 5D; p=0.054, sign rank test, n=10 mice). Thus, when pre-cue MOs activity is of the form that strongly indicates either a left or right turn, the motion is executed rapidly, as would be the case if this MOs activity subserved motor preparation for an action to be released as soon as the go cue sounds.

Inhibition of MOs activity reduces perseveration and slows responses

To investigate the causal role of MOs in the task, we used a transgenic mouse line (Ai32xPV-Cre) which expressed a Channelrhodopsin 2 in Parvalbumin-positive interneurons (Guo et al., 2014; Olsen et al., 2012). We implanted six mice bilaterally with cannulas allowing light delivery to the surface of MOs; a further six mice had cannulas in the same AP and DV location, but 1mm deeper, to study the effects of medial prefrontal cortex (mPFC) inactivation, and also to act as a control for inactivation spreading further than anticipated (Li et al., 2019). After surgery, the mice were trained to perform the task. We defined two behavioral windows for optogenetic inhibition: the "choice period", from go cue onset until a choice was registered, and the "outcome period", for 1.6 seconds from the time of feedback (Figure 6B). We applied 473 nm light to suppress neural activity during the choice period on a randomly-selected 20% of trials, and during the outcome period on an independently-selected 20% of trials. Our dataset comprised 44,243 trials from 82 sessions in MOs, and 48,614 trials from 92 sessions in mPFC.

Inactivating MOs during the choice period reduced both accuracy of the mice's choices, and the probability of repeating the previous choice (accuracy in MOs paired t-test, n=6 mice, p = 0.01; repeat probability in MOs paired t-test, n=6 mice, p = 0.01; Figure 6 D-E). In contrast, no significant effect was seen when mPFC was suppressed during the choice period (accuracy in mPFC paired t-test, n=6 mice, p =0.2; repeat probability in mPFC paired t-test, n=6 mice, p =0.12; Figure 6 D-E), and the effect of MOs suppression was

Figure 6 | MOs activity is required for perseveration and rapid responses. A, Location of light-delivery cannulas in two groups of mice. **B**, Timing of optogenetic suppression in the choice period or outcome period; light was applied independently with 20% probability to each period on each trial. **C**, Example session with choice period-suppression in MOs. Cyan: laser; inset: expanded timescale. **D-E**, Effect of MOs (left) or mPFC (right) suppressing during the choice period on accuracy, i.e fraction of trials choosing the Correct side (D) or probability of repeating the previous choice (E). **F-G**, Same for suppression during the outcome period. **H**, Cumulative histogram of response times on trials with optogenetic suppression in the choice period (cyan) and on no-laser trials (black), for mice with cannulas in MOs (top) and mPFC (bottom). Individual mice, thin lines, average across mice, thick lines. **I**, Left, Trial history regression weights for predicting mouse choice using past choices (perseverative weights), and choice-reward interaction (reward-seeking weights), for laser (cyan) and no laser (black) conditions. Left 4 panels show weights from trials N-n to trial N, where inactivation was delivered on trial N; right panels show weights from trial N to trial N+n, where inactivation was delivered on trial N. Right 4 panels but for PR model with different weights fit to suppression trials.

significantly stronger than that of mPFC inactivation (unpaired t-test comparing mean change in accuracy, n=12 mice, p=0.01). Furthermore, inactivation during the outcome period on random trials, either in MOs or

mPFC, did not affect repeat probability, and did not affect accuracy in MOs (accuracy in MOs paired t-test, n=6 mice, p = 0.12; repeat probability in MOs paired t-test, n=6 mice, p = 0.11; repeat probability in mPFC paired t-test, n=6 mice, p = 0.43; Figure 6 F, G). The accuracy was increased by mPFC inactivation during the outcome period, although only by -0.0065 on average (paired t-test, n=6 mice, p = 0.005, Figure 6 F, right). The magnitude of accuracy change following mPFC stimulation during the outcome period was significantly smaller than the magnitude of accuracy change during the choice period in MOs (unpaired t-test, n = 12 mice, p = 0.003). Applying trial history regression specifically to trials where MOs was inactivated yielded lower perseverative weights (i.e. weights from previous choices to the inactivated trial) and lower reward seeking weights (i.e. weights from previous choice-reward interactions) (Figure 6 I, left panels; perseverative weights: p=0.01, reward-seeking weights: p=0.007, t-test, n=6 mice), but this was not seen with mPFC inactivation (perseverative weights: p=0.41, reward-seeking weights: p=0.79, t-test, n=6 mice).

Inactivating MOs on a trial also reduced the effect of that trial on future perseveration. To show this, we refit the trial history regression, with different weights from inactivated and non-inactivated trials onto future choices. The perseverative weight from an inactivated trial to the next trial's choice was decreased, while the reward-seeking weight to the next trial's choice was increased (Figure 6I, right panels; perseverative weights: p=0.008, reward-seeking weights: p=0.01, t-test, n=6 mice). Thus, inactivating MOs during the choice period of a given trial makes the mouse less likely to perseverate in the future based on the choice it made on that inactivated trial, but does not impact the mouse's ability to learn from the reward delivered after the choice was completed.

Inactivating MOs during the choice period also caused an increase in response times (Figure 6 H). This inactivation did not prevent movement: on 84% of trials, the mice turned the wheel while the laser was still on. However, the time at which the wheel was turned was later (2.58±0.58 vs 1.91±0.15 s; median ± s.e., p=0.03, signed rank test, n=6 mice). On a further 16% of laser trials, the mice only made their choices after the laser was switched off (4.8 s after the go cue); the fraction of correct choices was similar on these trials as for the trials with responses during laser stimulation (0.66±0.02 vs 0.74±0.03, p=0.21, signrank test, n=6 mice). When mPFC was inactivated, we observed a trend toward shortened response time although it was not significant (2.96±0.45 vs 2.31±0.25 s, median ± s.e., p= 0.06, signrank test). Together, these results indicate that neural activity in MOs plays a role in guiding and rapidly executing choices, based on both perseverative and reward-seeking strategies; and that MOs activity during the choice has an impact on future choices but only for the perseverative component.

Inactivation following feedback delivery showed that mPFC, but not MOs is involved in reward-based learning. Inactivation of either region during the outcome period (1.6 s following delivery of reward or white noise feedback) on random trials did not have a large effect on subsequent choices (Figure 6F). However, because it takes mice several trials to learn about a block switch, an effect on reward learning might only be observable when multiple trials were inactivated successively. We therefore performed new experiments silencing outcome-period activity on all trials of a randomly selected half of each session (figure 7A-B). This allowed us to compare learning curves for block switches within the inactivated and non-inactivated half-sessions (Figure 7C). Inactivating mPFC significantly slowed learning about the block switch, but inactivating MOs had at most a marginal effect (Figure 7D), and the difference between the effects of mPFC and MOs inactivation were significant (figure 7E, p=0.04, t-test, n=6 mice in MOs group, n=7 mice in mPFC group). Thus, while MOs appears to be involved in perseveration and motor execution, mPFC seems to be required for reward learning in this task.

Figure 7 | Optogenetic mPFC suppression during outcome period impairs learning. A, Timing of laser pulses in this experiment. **B,** Two example sessions. For each session, light is applied randomly on either every trial in the first half session (top) or every trial in the second half (bottom). Cyan lines: laser trials. **C,** Learning curve of MOs (left) and mPFC (right) groups, for no-laser (black) and laser blocks (cyan). Thin lines, individual mice, thick lines, average. **D,** Difference in accuracy between laser and no laser blocks averaged over sessions (magenta), compared to a null distribution obtained by randomly resampling which half of trials was inactivated on each session (grey). Left: MOs suppression, right: mPFC suppression. Red Stars: timepoints for which accuracy in the inactivated half-session was worse than the null distribution (p<.01, randomization test). **E,** comparison of accuracy with MOs and mPFC inactivation. Each point shows the difference in accuracy between laser and non-laser halves, averaged over the first 15 trials following each block switch for all sessions of one mouse. The difference between mPFC and MOs was significant (p<.05; rank sum test, n=13 mice).

Discussion

We found that mouse behavior in a probabilistic reversal learning task could only be captured by computational models containing both reward seeking and perseverative mechanisms. Response times

were not correlated with model-predicted reward-seeking drive but were shorter for repeated choices and when the model predicted strong perseverative drive. Neuronal activity in many brain regions correlated with choice once movement had begun, but correlates prior to movement onset were primarily found in dorsal prefrontal cortex (MOs). This pre-movement activity was better explained by encoding an analog perseverative decision variable than the particular upcoming choice. Silencing MOs, but not mPFC activity at choice time slowed response times and made choice direction closer to random. Silencing mPFC, but not MOs following reward delivery slowed learning rates. Together, these results indicate that MOs is involved in driving perseverative behavior.

The benefits of perseveration to an animal are unclear. Most theories build from "dual-process" theories in psychology (Daw et al., 2005; Dickinson, 1985; Evans & Stanovich, 2013; Kahneman, 2011): a "system one" responds quickly and reliably but is relatively inflexible, while a "system two" is more computationally complex, slower to respond and more flexible. Some theories (Ashby et al., 2007; Bogacz, 2020; Miller et al., 2019) further propose that system one follows the perseverative law of exercise, selecting actions based in part on a running estimate of how frequently each action has been previously selected. Our findings support these predictions in four ways. First, MOs activity correlates specifically with a perseverative variable, reflecting a running average of recent actions. This is broadly consistent with fMRI recordings in human perceptual decision-making (Akaishi et al., 2014; Waldschmidt & Ashby, 2011). Second, silencing MOs reduces the influence of past choice on future choice, suggesting that these representations are causally involved in perseveration. Third, perseveration correlates with faster responses, while MOs activity both correlates with and is causally necessary for fast responses, consistent with system one processing. Finally, silencing MOs impairs perseverative learning while increasing reward-guided learning.

The neural mechanisms of perseveration are also unclear. Some models propose that perseverative information is stored in the strengths of either cortico-cortical (Ashby et al., 2007, 2010) or cortico-striatal synapses (Bogacz, 2020). These models do not require spiking correlates of perseverative information in intertrial intervals. We did however find such correlates in MOs, even before the go cue. This does not preclude a change in synaptic weights, but does raise the possibility that short-timescale perseveration might rely at least partially on persistent activity (J. X. Wang et al., 2018). Recent experiments have shown tail-striatal dopamine correlates with action prediction error, which encodes the difference between the P variable and the action made (Greenstreet et al., 2022). It is possible that MOs encoding of the P variable contributes to the computation of this signal.

A prominent theory holds that MOs is responsible for short-term memory of information relevant for guiding choice (Barthas & Kwan, 2017; Yang & Kwan, 2021), which is supported by recordings in many tasks (Barthas & Kwan, 2017; Ebitz et al., 2018; Erlich et al., 2011; Li et al., 2015; Murakami et al., 2014; Sul et al., 2011). Our finding that fixation-period MOs activity correlates with the perseverative variable P is broadly consistent with this idea: in perseverative behaviors, past choices guide future choices. However, we did not find significant correlates of the reward-seeking variable R, which also guides future choices. It remains possible that MOs encodes R but too weakly to detect here, for example MOs might represent the summed decision variable $P+R$, with the smaller size of R than P meaning that only P was individually significant. Alternatively, R might be stored elsewhere and only pass through MOs after the go cue, a possibility consistent with the facts that the previous trial's interaction variable is significantly encoded after the go cue (supplementary figure 7), and that MOs inactivation at choice time suppresses both perseverative and reward-seeking behavior. Our inactivation results contrast with those of Sul et al., 2011, who found that in a spatial reversal learning task, MOs lesions increased rats' tendency to perseverate. We hypothesize that this may be because their task encourages choice alternation rather than consistent selection of a higher-reward option. If this hypothesis is correct, then one should not take our results as

indicating a role for MOs in perseveration in all tasks; rather that MOs can be involved in both perseverative and anti-perseverative strategies, depending on the task, species, and context.

Our results are consistent with the hypothesis that perseveration enables rapid responses (Ashby et al., 2007; Gershman, 2020; Wood & Rünger, 2016). Reaction times were faster for repeated choices, and inactivating MOs at choice time increased reaction times. We therefore suggest that MOs activity serves to prime downstream circuits so that the moment the go cue sounds, a rapid response in the pre-planned direction is released. This priming might consist of depolarization in premotor units, or embodied changes such as altered muscle tension. When MOs is inactivated, the priming is removed, and the downstream circuit takes more time to produce an action in a direction selected close to randomly.

In summary, we identified area MOs as a region encoding a perseverative decision variable used in this dynamic reward learning task, and necessary for implementing the effect of this decision variable on choice direction and reaction time. Future work can establish which if any other areas also encode this variable especially subcortically, the mechanisms by which the decision variable is updated, and the mechanisms by which it affects downstream circuits involved in executing the actual action.

Methods

Initial surgery

All experimental procedures were conducted at University College London according to the UK Animals Scientific Procedures Act (1986) and under personal and project licences granted by the Home Office following appropriate ethics review.

Mice were implanted with a head-fixation plate over the interparietal bone and a 3d-printed guide, following the procedure described in detail in Steinmetz et al. (2019). During an initial surgery under isoflurane anesthesia (1-3% in O2), we implanted a steel headplate (size 15 x 3 x 0.5 mm; weight approximately 1g), followed by a custom guide. The guide was 3d printed from opaque polylactic acid, and when placed on an average mouse skull exposes an area from 3.5mm anterior to 5.5mm posterior to bregma, and 4.5mm left and right from bregma, allowing targeting multiple areas without a need for additional surgery, as well as protecting the probes from mouse paw movements that otherwise could break them. To implant the headplate, the surface of the skull was cleaned from the skin and periosteum and a green activator (Super-Bond C&B, Sun Medical) applied to the cleared skull, specifically to the area where the headplate was later implanted, to strengthen the connection between the bone and the cement. The skin was attached to the skull on the sides using Vetbond Tissue Adhesive (3M), and the skull surface was also covered by it, except for the area where the headplate was later implanted. The headplate and guide were then attached to the skull with dental cement (Super-Bond C&B, Sun Medical Co, Ltd, Japan). Two layers of UV-curing optical glue (Norland Optical Adhesives #81, Norland Products) were applied to the skull and cured for 10 seconds per layer. The layers needed to be thin to prevent excessive heat. The whole procedure took around 1 hour.

After recovery, mice were treated with carprofen added to their drinking water for three days while recovering in a home cage. After this time, they were given at least an additional two days of ad lib access to water, before beginning water restriction and habituation in preparation for behavioral training.

Task

Our task used a previously-described behavioral apparatus (Burgess et al., 2017). The mouse was head-fixed, within a half-pipe tube and covered with another half-pipe for comfort. Mouse front paws sat on a

rubber wheel (LEGO, 86652 & 32019) which could be freely turned to register choices. Wheel movements were measured using an incremental rotary encoder with a resolution of 1024 increments per turn. Rewards were delivered via a silicone lickspout placed within easy reach of the mouse's tongue. The mouse was surrounded by three video screens but no visual stimuli were displayed, just gray background.

Each trial began with a brief auditory tone (go cue, 5kHz for 100 ms), following which the mouse could turn a wheel for a liquid reward. The tone carried no information about which direction of turn would be rewarded. On each trial one of the sides was associated with 80% reward probability and 20% omission probability, and the other side was associated with 20% reward probability and 80% omission probability. The reward-probability contingencies (blocks) flipped randomly every 125-175 trials, with the duration of the first block chosen randomly between 1 and 175 trials, and the first correct side chosen at random. These block lengths are longer than what is used typically (Beron et al., 2022; Hattori et al., 2019; Le et al., 2023). Block flips were not signaled to the animal. We call choices that correspond to the 80% reward probability "correct choices", independently of whether mice are rewarded on these trials. A choice was registered when the mouse turned the wheel by 18mm in either direction. Trials did not end until this threshold was reached; there was no timeout. Upon completing a turn, the mouse received feedback (either a 1-3μL drop of 10% sucrose in water or an auditory white noise) coupled with a prolonged intertrial interval. Feedback delivery was followed by an intertrial interval of random length: uniformly distributed 1 and 2 s following reward delivery, and between 2.5 and 3.5 s following reward omission. After the inter-trial interval, the mouse had to hold the wheel completely still, to within the resolution of the rotary encoder, for a "fixation period" of 200ms before the next trial began.

Behavior training

Training began with two habituation sessions (on two consecutive days), where mice were head fixed for 15 and 30 minutes, respectively. We next trained mice 1 hour per day on a deterministic reward schedule, where the correct choice was rewarded 100% of the time and the incorrect choice was never rewarded. Once mice followed block switches for at least 3 days in a row, we changed the probabilities to 90% for the correct choice and 10% for the incorrect choice. Finally, when mice were judged to perform that version well, we changed the probabilities to 80% and 20%. Mice were trained 5 days per week, and typically became proficient in the final version of the task 4-12 weeks after the start of the training.

Behavioural models

Models used

We compared the 13 models behavioral models. These were four variants of Q-learning: standard Q-learning (Sutton & Barto, 1998), optimistic Q-learning (Lefebvre et al., 2017), differential forgetting Q-learning (Ito & Doya, 2009), and the *PR* model (Miller et al., 2021); six variants of ideal observer models: standard ideal observer, ideal observer with drifts (bayes drifts), ideal observer with flips (bayes flips), ideal observer with perseveration, ideal observer with drifts and perseveration, ideal observer with flips and perseveration; and three additional models: a local matching law (Sugrue et al., 2004), a foraging agent (Constantino & Daw, 2015), and an actor-critic model (Sutton & Barto, 1998).

Standard Q-learning

Standard Q-learning (Sutton & Barto, 1998) is a reinforcement learning algorithm that is used to learn the optimal action-value function for an agent interacting with an environment.

The two internal variables of the Q-learning model are the action values of the left and right sides (Q_L and Q_R). When a new session starts, both Q_L and Q_R are set to 0. The prediction error δ_t is calculated for trial t as a difference between the actual and the expected reward:

$$\delta_t = r_t - Q_{C,t}$$

Here C represents the choice made (L or R). This calculated reward prediction error is used to update the Q value of the chosen action with learning rate α:

$$Q_{C,t+1} = Q_{C,t} + \alpha\,\delta_t$$

The Q value associated with the unchosen side is not updated. Decisions are made according to a softmax rule, dependent on two parameters: a bias β_{bias} and an "inverse temperature" β_Q:

$$P_L = \frac{exp(\beta_Q * (Q_{L,t} + \beta_{bias}))}{sum(exp(\beta_Q * (Q_{L,t} + \beta_{bias})) + exp(\beta_Q * (Q_{R,t} - \beta_{bias})))}$$

The bounds for the parameter search are: $\alpha \in [0,1]$, other parameters $\in (-\infty, \infty)$.

Optimistic Q learning

Optimistic Q-learning (Lefebvre et al., 2017) is an extension of the standard Q-learning algorithm that allows the speed of learning to differ between positive and negative reinforcement:

$$Q_{C,t+1} = \begin{cases} Q_{C,t} + \alpha_1\delta_t & \delta_t > 0 \\ Q_{C,t} + \alpha_2\delta_t & \delta_t \le 0 \end{cases}$$

The parameter α_1 the learning rate from positive prediction error and α_2 is the learning rate from the negative prediction error. If $\alpha_1 > \alpha_2$, then the agent is implementing the optimistic Q-learning, modelling how people tend to incorporate good news into their world model faster than bad news (Lefebvre et al., 2017); and if $\alpha_1 < \alpha_2$, it is pessimistic Q-learning. If $\alpha_1 = \alpha_2$, it is standard Q-learning described in the previous section. This model has four free parameters, α_1, α_2, β_{bias}, β_Q, and the Q values are updated according to the following rules:

The bounds for the parameter search are: $\alpha_1 \in [0,1]$, $\alpha_2 \in [0,1]$, other parameters $\in (-\infty, \infty)$. Choices are made according to the softmax function with parameters β_{bias} and β_Q, as above.

Differential Forgetting Q-learning

The Differential Forgetting Q-learning rule (Ito & Doya, 2009) allows modeling of perseveration, by causing the Q-value of the unchosen side to decay toward zero. (In standard Q-learning, the Q-value of the unchosen side does not change. It uses the following rule:

$$Q_{C,t+1=} \begin{cases} (1 - \alpha_1)Q_{C,t} + \alpha_1\kappa_1 & rewarded \\ (1 - \alpha_1)Q_{C,t} + \alpha_1\kappa_2 & unrewarded \\ (1 - \alpha_2)Q_{U,t} & regardess\ of\ reward \end{cases}$$

Here, C refers to the chosen side and U to the unchosen side (both L or R). The bounds for the parameter search are: $\alpha_1 \in [0,1]$, $\alpha_2 \in [0,1]$, other parameters $\in (-\infty, \infty)$. Choices are made according to the softmax function with parameters β_{bias} and β_Q, as above.

PR model

The PR model has two decision variables: P which follows the law of exercise and mediates perseverative behavior, and R which follows the law of effect and mediates reward-seeking behavior. The perseverative decision variable P is initialized to zero at the beginning of every session, and updated according to:

$$P_{t+1} = P_t + \alpha_p(C_t\lambda_p - P_t)$$

where C_t is the choice (left or right, coded as -1 or +1) made on trial t, α_p is a learning rate parameter governing the extent to which an individual choice affects P, and λ_p is a magnitude parameter governing the largest possible magnitude of P. This rule therefore updates P towards $-\lambda_p$ following left choices and $+\lambda_p$ following right choices, so P can be thought of as a running average of previous choices (Figure 1H, yellow line).

The reward-seeking decision variable R is likewise initialized to zero at the beginning of each session, and is updated according to the following rule:

$$R_{t+1} = R_t + \alpha_r(\lambda_r C_t F_t - R_t)$$

where F_t is the feedback (unrewarded or rewarded, coded as -1 or +1) received on trial t, and α_r and λ_r are learning rate and magnitude parameters, which can be different to the perseveration parameters α_p and λ_p. The rule therefore updates R towards $-\lambda_r$ following rewarded left choices or unrewarded right choices, and towards $+\lambda_r$ following rewarded right choices or unrewarded left choices. Thus, R can be thought of as a running average of the choice-reward interaction, which provides an estimate of the correct choice (Figure 1H, orange line).

On each trial, the PR model agent chooses right over left with log odds given by the sum of the two decision variables:

$$\log\left(\frac{P(C_t = +1)}{P(C_t = -1)}\right) = P_t + R_t + b$$

where b is a parameter governing fixed left/right bias. The overall magnitudes of P and R – determined by the parameters λ_p and λ_r – therefore govern the relative influence of perseveration and reward seeking on choice. We fit the model parameters to the observed behavior of each mouse by maximum likelihoodLocal matching law

Hernstein's matching law states that the rate of responding to a particular option or stimulus is proportional to the relative rate of reinforcement associated with that option or stimulus (Herrnstein, 1970). The local matching law implements this through equations are similar to the Q-learning models discussed above (Sugrue et al., 2004). The learning rate α reflects the speed of the action value update, the sensitivity shows the strength of the connection between the reinforcement ration and the choice ratio. Then, the Q-value update for Q_L can be written as follows:

$$
\begin{aligned}
Q_{C,t+1} &= (1 - \alpha)Q_{C,t} + \alpha \quad rewarded \\
Q_{C,t+1} &= (1 - \alpha)Q_{C,t} \quad\quad\;\; unrewarded \\
Q_{U,t+1} &= (1 - \alpha)Q_{U,t} \quad\quad\;\; regardless
\end{aligned}
$$

Here, C refers to the chosen side and U to the unchosen side (both L or R). The bounds for the parameter search are: $\alpha \in [0,1]$. Choices are made according to the softmax function as above, but without bias terms in this case.

Foraging agent
The foraging agent (Constantino & Daw, 2015) explores the environment, making its decision about whether to stay with the current choice or to switch sides based on an estimate of the value of the current choice. Unlike the models above, this model only has one internal variable V, which estimates the value of the current choice, and resets when the agent switches sides. The value is updated according to

$$V_{t+1} = \begin{cases} (1 - \alpha)V_t + \alpha r_t : C_t = C_{t-1} \\ \qquad\qquad V_0 : C_t \neq C_{t-1} \end{cases}$$

Here $C_t = \pm 1$ represents the choice on trial t. The decision is made by a logistic function with a bias for switching as well as for left/right side:

$$\log\left(\frac{P(C_t = 1)}{P(C_t = -1)}\right) = \left(\beta V_t + b_{stay}\right)C_t + b_{lr}$$

The parameters α, β, b_{stay}, b_{lr} represent learning rate; strength of history, stay bias, and left/right bias, respectively. The bounds for the parameter search are: $\alpha_{local} \in [0,1]$, $V\_global \in [0,\infty)$, other parameters $\in (-\infty, \infty)$.

Actor-critic

Actor-critic algorithms are a family of algorithms that explicitly represent policy independent of value function (Sutton & Barto, 2018; Section 2.8). The policy structure (actor) selects actions, while the estimated value function (critic) criticizes those actions. The scalar signal of the critic drives learning of both actor and critic. In general, actor-critic algorithms allow both the actor and critic components to depend on sensory state; in the current task, with only a single state, the actor-critic model takes a particular simple form.

We considered a "forgetful" Actor-Critic model, which includes a decay rule that updates the policy variable towards zero on each trial, allowing the agent to produce adaptive behavior in an environment with changing reward contingencies (Miller et al., 2023). This model has three parameters: α_c, the critic learning rate; and α_l, α_f, the learning and forgetting rates for the actor. It has two state variables: V, the critic's expected reward; and H, the difference in the actor's estimated values for actions to the left and right.

The critic learns according to the following rule:

$$V_{t+1} = (1 - \alpha_c)V_t + \alpha_c * r_t$$

The actor updates its values according to the following rule:

$$H_{t+1} = \left(1 - \alpha_f\right)H_t + C_t\,\alpha_l\,(r_t - V_t)(1 - P(C_t))$$

Here $C_t = \pm 1$ represents the chosen action (L or R) on trial t. $P(C_t) = \frac{1}{1+e^{-H_t}}$ represents the probability that action C_t will be chosen, given by the logistic formula. Thus, the actor's estimate of the advantage in choosing left over right is increased when it makes a left turn and receives a larger reward than the mean value V estimated by the critic. The term $(1 - P(C_t))$ serves to boost the learning rate when it makes a rare choice. The search bounds for all parameters are [0,1].

Ideal Observer Models

An ideal observer model, also called a Bayesian filter, uses knowledge of the structure of a partially-observable dynamic environment to optimally infer the current state of that environment (Sarkka, 2013). It assumes that in each timestep the environment can be in one of several possible discrete hidden states, h_t, and that on each trial the environment first probabilistically emits one of several possible visible observations, o_t, and then probabilistically transitions into a new hidden state h_{t+1}. An ideal observer model requires three functions containing knowledge of the environment. The first is an Initialization Function, giving the model's belief distribution over hidden states on the first timestep: $p(h_0)$. The second is an Observation Function, giving the probability with which each observation will be produced in each hidden state $p(o_t|h_t)$. The third is a Dynamics Function giving the probability with which the environment will transition to each possible subsequent state $p(h_{t+1}|h_t)$.

The Bayesian filter model maintains a running belief distribution $B_t(h)$ over the possible current states of the environment. This distribution is initialized on the first trial to match the Initialization Function, and then updated after each trial in two steps. The first step makes use of the Observation Function and computes a posterior probability over h_t conditional on the observation o_t using Bayes' theorem. The second step makes use of the Dynamics Function to compute a belief over h_{t+1} marginalizing over this posterior.

Optimal Agent

We first consider the optimal agent, built around an ideal observer model with full knowledge of the true rules of the task. Specifically, it knows that the environment can be in one of 350 possible states: The current high-reward choice can be either left or right, and there can be between 1 and 175 trials remaining in the current block. We can therefore represent the agent's belief distribution as a pair of vectors of length 175, the sum of whose elements is together 1. The Initialization Function is uniform, as at the beginning of the session it is equally likely that the rig will be in any of these states. We therefore initialize the vectors uniformly at the start of each session by setting each element to 1/350.

The Observation Function is given by the reward contingencies: if the mouse selects the high-reward choice, reward is delivered with probability 80%, and if it selects the low-reward choice, reward is delivered with probability 20%. We therefore update each element of the belief distribution after each trial using Bayes' theorem and renormalize.

The Dynamics Function is given by the tasks' block switching rules: on each trial the "trials remaining in block" variable is decremented by one, until it reaches 0 in which case the "high-reward" variable is flipped and "trials remaining in block" is sampled uniformly between 125 and 175. We therefore move probability mass down the vectors by one slot, such that the belief on trial t that there are exactly n trials remaining in the block is equal to the belief on trial t+1 that there are exactly n-1 trials remaining. The belief from the final slot (belief that this was the final trial in the block) is redistributed uniformly over the first 50 slots in the alternate vector (e.g. if the model had belief 50% that this was the final trial in a left-better block, this will add 1% belief to each element between "125 trials remaining in a right-better block" and "175 trials remaining in a right-better block").

In this way, the ideal observer optimally updates a belief distribution over the true hidden states of the experiment. It can use this distribution to select actions, marginalizing over the number of remaining trials to compute an overall belief that it is currently in a left-better vs a right-better block. We convert this ideal observer into an optimal agent by having it deterministically select the action which it believes is currently high-reward.

Noisy Ideal Observer

We define the noisy ideal observer agent as containing the same ideal observer model as the optimal agent, but selecting actions using a softmax decision rule with fit parameter β_b:

$$\log\left(\frac{P(C_t = L)}{P(C_t = R)}\right) = \beta_b B_t$$

Where $B_t \in (0, 1)$ is the ideal observer's belief on trial t that the left action is currently the highly rewarding action. The noisy ideal observer agent is therefore characterized by the one free parameter β_b.

Bayesian filter assuming probabilistic reversals (Bayesian flips)

The Ideal Observer agents make use of detailed knowledge of the dynamics of our particular task that it is unlikely a mouse would be able to perfectly infer. We therefore considered a Bayesian filter agent assuming a more general task structure: that there are two discrete blocks, each with a high-reward and low-reward side delivering reward with distinct probabilities, and that these blocks change unpredictably over time.

This set of assumptions corresponds to a belief in a single hidden variable: the identity of the current high-reward side. The belief state of the agent on each trial is therefore characterized by a single number B_t giving its current belief that the left action is high-reward. We initialize this belief to 0.5 at the beginning of each session, and update it following each trial. The Observation Function involves two free parameters P_{high} and P_{low}, giving the model's assumption about the reward probabilities on the high-reward and low-

reward side. The Dynamics function involves one free parameter P_{flip} giving the models assumption about the probability with which the block will flip on each trial.

The Bayesian filter assuming probabilistic reversals therefore has four free parameters: β_b, P_{high}, P_{low}, and P_{flip}.

Bayesian filter assuming time-varying probability changes (Bayesian drifts)

The Bayesian filter assuming probabilistic reversals still contains structural knowledge about our task that might not be available to the mouse: that reward probabilities change discretely and reverse between a particular set of fixed values. We therefore also considered a Bayesian filter agent assuming an even more general task structure: that each action is associated with a separate reward probability, and that the value of each of these probabilities on each trial tends to resemble its value on recent past trials.

This set of assumptions corresponds a belief distributions over the full set of probabilities between 0 and 1 for each action. We use θ_{left} and θ_{right} to parameterize the reward probability on the left and on the right port, and B_{right, t} and B_{left, t} to indicate the observer's belief at time t over these probabilities. We approximate these distributions numerically as histograms, allowing θ to take 200 discrete values equally spaced between zero and one. This means that the first element of B_{left, t}, for example, corresponds to the observer's current belief that the reward probability of the left action is currently between 0 and 0.5%

Belief distributions are initialized to be uniform at the start of each session. After each trial, we first update the vector corresponding to the chosen action using the Observation Model given by the average probability in each bin (for example the first element is updated on the assumption that rewards are given in this state 0.25% of the time). We next update both vectors on the assumption that probabilities can change over time, by convolving them with a gaussian kernel whose standard deviation (σ) is a free parameter corresponding to the observer's belief about the plausible rate of environmental change. Probability mass which would be spread by this convolution to values less than zero is placed instead into the lowest bin (0% to 0.5%), and likewise mass which would be spread to values greater than one is placed into the highest bin (99.5% to 100%).

To make choices, the observer marginalizes these distributions, converting a belief distribution over all possible values of the generative reward probability into a single number indicating the probability with which the observer expects a reward if it selects this action:

$$V_{C,t} = \sum_{\theta'} \theta' B_{C,t}(\theta')$$

Where the index C can take the values L or R. This agent selects actions according to the difference in reward probabilities between the two actions:

$$\log\left(\frac{P(C_t = L)}{P(C_t = R)}\right) = \beta\left(V_{L,t} - V_{L,t}\right)$$

This Bayesian filter agent therefore has two free parameters: σ and β

Ideal Observers with Perseveration

For each of the three Bayesian filter agents (Noisy Ideal Observer, Bayesian Flips, Bayesian Drifts), we also define and fit "with perseveration" versions, which are mixture models comprised of the Bayesian filter alongside a perseverative agent as defined in Equation (1):

$$P_{t+1} = P_t + \alpha_p\left(C_t\lambda_p - P_t\right)$$

The decision rule of these agents is a mixture incorporating both the Bayesian filter's decision variable as well as a perseverative decision variable:

$$\log \frac{P_{t,left}}{P_{t,right}} = \beta_b B_t + P_t$$

These "with perseveration" versions therefore have all of the free parameters of the associated Bayesian filter model, as well as the two additional free parameters α_P and λ_P.

Fitting procedure

The parameters of all behavioral models were fit by maximum likelihood. Specifically we used MATLAB's fmincon function, with parameters: maxfunevals = 4000, maxiter = 2000, algorithm = 'interior-point'. For each parameter of each model, we manually specified a feasible range for the parameter search, and selected a starting value randomly in this interval. We repeated the fmincon procedure until it completed successfully three times, and recorded the parameter fits from the run that has the best likelihood of these three runs. The runs were consistent: for all but three models the difference in log-likelihood between runs was no larger than 1e-5. Among the remaining three models, two were non-perseverative (Bayesian flips and a foraging agent). For these models, we ran the fitting procedure 20 times and the log-likelihood for every run was lower than the log-likelihood of any perseverative model. For the remaining model (Bayesian flips with perseveration) we ran the fitting procedure 20 times; it had high-varying log-likelihood, but in all cases it was higher than the log-likelihood of all non-perseverative models.

For model comparison, we use a cross-validation approach that allowed for differences in strategy between mice. We fit each model's parameters on odd sessions of each mouse separately, and evaluated the model by log likelihood on the even sessions using parameters fit to the same mouse; the model parameters were thus assumed to differ between mice, but not between sessions of one mouse. The quality of fit for each mouse was obtained by dividing the total log likelihood by the total number of trials, then exponentiating to yield a geometric mean likelihood.

Trial-history regression

To gain a better understanding of the effect that MOs suppression has on choices, we compared trial history regression analyses (Lau & Glimcher, 2005) on trials with and without optogenetic suppression of MOs activity. The regression included separate terms for the choice, feedback and their interaction for past trials trying to predict the choice on the current trial. There are thus separate weights for each of the two possible choices (left, right), each of the two reward outcomes (rewarded, unrewarded), and each of the possible interactions (left rewarded or right unrewarded vs. right rewarded or left unrewarded), for each time lag considered. The pattern of trial-history regression weights provide information on the decision strategy used by the mice, without reference to explicit psychological models (Miller et al., 2022).

The full equation for trial history regression is:

$$\log\left(\frac{P_{left}(t)}{P_{right}(t)}\right) = \beta_0 + \sum_{\tau=1}^{T} \beta_C(\tau)C(t-\tau) + \sum_{\tau=1}^{T} \beta_X(\tau)C(t-\tau)R(t-\tau) + \sum_{t=1}^{T} \beta_R(\tau)R(t-\tau)$$

This equation expresses the log odds choice on trial t as a function of past choices and rewards. *C* indicates past choices, taking the value 1 for left choice trials (either rewarded or not) and -1 for right choice trials. Positive values of β_c, therefore, indicate a tendency to repeat past choices. *R* indicates past rewards, taking the value 1 for rewarded choices (either left or right) and -1 for non-rewarded choices. Positive values of β_x, therefore, indicates a tendency to repeat choices that have led to rewards and to switch away from choices that have not. Positive values of β_r would indicate a tendency to select the left action following rewards, regardless of the choice that preceded those rewards. T is a hyperparameter that governs the number of past trials that go into the model to predict the upcoming choice, set to 10 for the current analyses. The weights β were fit by maximum likelihood using MATLAB's glmfit function with default options.

Secondary surgery for electrophysiological experiments

After mice were fully trained in the task, they were retrained in the recording rig, which typically required additional 3-10 days.

On the first day of recordings, mice were anesthetized (1-3% isoflurane in O2) and a <0.5mm craniotomy was made over the desired location with a dental drill (coordinates: 1.8AP, ±2ML for targeting DMS; -1.6AP, ±0.9ML for targeting the hippocampus; 2.2AP, ±0.9ML for targeting MOs). The location of the craniotomies and the desired probe angles were determined using custom software (https://github.com/petersaj/AP_scripts_cortexlab/tree/master/allen_interface). A thin layer of Kwik-Cast (World Precision Instruments) was applied on top of the craniotomy to protect it. The animals were allowed to recover for at least 3 hours before the recording began. For each additional recording location, the procedure was repeated, with a maximum of three locations per mouse.

Electrophysiological recordings

Neuropixels probes, either 1.0 version 3B or 2.0, were held by metal rods and gradually lowered into the brain using a motorized micromanipulator (uMP-4, Sensapex Inc.), with an approximate speed of 300 micrometers per second, piercing the dura mater. To allow later track localization, prior to insertion probes were coated with a solution of DiI (ThermoFisher Vybrant V22888, V22886 or V22885) by holding 2μL in a droplet on the end of a micropipette and touching the droplet to the probe shank, letting it dry, and repeating until the droplet was gone, after which the probe appeared colored. After smooth descent to the final depth, probes were left untouched for at least 30 mins before recording. The plastic well was filled with physiological saline for the duration of the recording, prevent drying. Recordings were performed using a PXI system (NI PXIe-1071) and spikeGLX software. The recorded data were spike sorted using Kilosort 2.5 (Pachitariu et al., 2023) and then manually curated to remove noise clusters with phy (https://github.com/kwikteam/phy). Units were considered "noise" and removed from further analysis if they had non-physiological waveform shape or pattern of activity across channels. No further manual processing was performed.

Histology

Following recordings, mice were transcardially perfused following standard procedures. Briefly, they were administered a terminal dose of pentobarbital intraperitoneally, followed by a toe-pinch to ensure anesthesia. The thoracic cavity was opened, the right atrium was incised, and a solution of phosphate buffered saline (PBS) followed by 4% formaldehyde (Thermofisher 28908) in 0.1M PB pH 7.4 was perfused through the left ventricle. The brain was carefully dissected out and post-fixed in the same fixative for a minimum of 24 hours at room temperature. After washing, the tissues were stored in PBS at 4°C for no longer than three weeks before being imaged *in toto* with serial section two-photon microscopy (Economo et al., 2016; Ragan et al., 2012).

Unit alignment

The three-dimensional location of each recorded unit was aligned to the Allen common coordinate framework atlas (Q. Wang et al., 2020) using custom MATLAB code (http://github.com/petersaj/AP_histology). For the purposes of analysis, we labelled units as belonging to "MOs" if they were in Allen atlas region "Secondary motor area"; to "mPFC" if they were in "Prelimbic area", "Infralimbic area" or "Anterior cingulate area"; we did not aim for higher specificity to avoid possible mislabeling due to probe depth. All units in Allen region "Caudoputamen" were assigned a label "DMS" due to their dorsomedial location. We assign a label "Dorsal hippocampus" to any of the units in CA1, CA3 or dentate gyrus. We assign the label "Orbital areas" to any of the units in the orbital area (including lateral and medial subdivisions). We assign a label "Olfactory areas" to any of the units in the olfactory area (including main olfactory bulb, piriform area and others).

We recorded activity from more brain areas than analyzed in the paper, however the statistical methods we used require at least 6 sessions with at least 30 neurons each, therefore we did not apply these analyses to these all recorded areas (all areas recorded and number of sessions per region listed in Supplementary figure 5).

Neural data analysis

Predicting a behavioural variable from neural activity

To assess whether a behavioral variable is coded by the neural activity, we used the session permutation method (Harris, 2021). This analysis asks whether neural data better predicts behavioral variables recorded simultaneously on the same session, than behavioral variables recorded on another session. This tests the null hypothesis that the full history of neural activity of a session is independent of the full sequence of behavior of that session, but makes no assumption of independence of neural or behavioral variables between trials, or of independence between cells.

The session permutation uses a test statistic $\sum_s \rho(\boldsymbol{A}_s, \boldsymbol{B}_s)$, where $\boldsymbol{A}_s$ and $\boldsymbol{B}_s$ represent the histories of neural activity and behavior on session s encoded into arrays with N_{trials} rows, and ρ is a scalar measure of correlation to be defined below. It compares this test statistic to a null ensemble $\{\rho(\boldsymbol{A}_s, \boldsymbol{B}_{\pi(s)}): \pi \in \Pi\}$, where Π is a set of 1000 random permutations of the session labels. To use the method, all sessions must be truncated to the same length. Here we use all sessions containing at least 30 neurons in a specified area, and truncate each on to have the same number of trials as the shortest of these sessions.

The correlation measure ρ was different for different types of behavioral data considered. When correlating neural activity with a continuous behavioral variable (such as the decision variable of the PR model), we used sparse linear regression. For every behavioral session, we found the 20 neurons with largest squared correlation with the behavioral variable, predicted the behavioral variable on each trial from these neurons' activity using unpenalized linear regression, assessed performance for each session with mean squared error (MSE), and summed across sessions to get the test statistic We then generated 1000 random permutations of the sessions, and recomputed the statistic 1000 times, but now predicting the behaviour on session s using the neural activity of session $\pi(s)$. We deemed the correlation significant if the test statistic is below the first percentile of the null ensemble.

To predict a binary variable (such as left/right choice), we applied the same procedure but using logistic regression, with fit quality evaluated by log likelihood instead of MSE. We deemed the result is significant if the real likelihood was above the 99th percentile of the likelihoods of shuffles.

To verify that any differences did not arise from different numbers of neurons recorded in different areas, we ran an analysis on equally subsampled set of neurons. We selected all sessions containing at least 60 neurons in a specified area and as before kept the first L trials of every session. Then, we randomly subsampled either 10, 20, 30, 40, 50 or 60 neurons and repeated the analysis above. We deemed MOs to be significantly different from other areas for a specific neuron count if the mean difference between the real and shuffled sessions in MOs was above the 99th percentile of such differences in 5 other regions.

Accounting for other behavioural variables

Since the decision variable is correlated with choice, we tested for a partial correlation of neural activity with the decision variable given the choice: does neural activity predict the decision variable even if a common correlation with choice is accounted for?

To do so without assuming independence of trials or neurons within a session, we used an adaptation of the session permutation method that tests for partial correlation (Harris & Yuan, 2024). Again this requires all sessions to have the same length, so we took only sessions containing at least 30 neurons in the brain area of interest, and truncated each to have the same number of trials L (the length of the shortest session).

To perform partial correlation, we first construct a $L \times L$ matrix Π that projects out the choice variable from each session. To do so, we construct a matrix D of size $L \times N_{sessions}$ containing the sequence of choices for each session. We then apply singular value decomposition to obtain a matrix U of the same size as D, with orthogonal columns spanning the same subspace as the columns of D, and define $\Pi = I - UU^T$. Denoting the L-dimensional timeseries of behavioral values for session s as b_s, we next predict Πb_s from the neurons recorded in session s using a sparse method: we find the 20 neurons that have the highest correlation with Πb_s and fit an unpenalized linear regression from their fixation period activity to predict Πb_s. We compute the test statistic as the average of the MSE of this regression over all sessions. To gauge significance, we compare this test statistic to a null ensemble in which the behavioral variables, but not the neural activity, has been permuted across sessions. Because this session permutation does not affect the projection matrix Π, it yields a test for partial correlation of the behavioral variable with neural activity, allowing for a common effect of choice.

Correlation of reaction time and neural activity in MOs

For every session containing at least 30 neurons in MOs, we compute the firing rate of each neuron in the 200ms fixation period prior to the go cue, for every trial. We select the 20 neurons whose fixation-period activity best correlates with the left/right choices (assessed by Pearson correlation). We then fit an unpenalized logistic regression model to predict the left/right choices on each trial from the resulting 20-dimensional vectors. We then compare summed neural activity arising from this model (prior to passing through the logistic function) on each trial to the animals' reaction time on that trial.

Causal analysis of reward or interaction coding

We took advantage of the fact that rewards were delivered at random to assess whether reward and reward-choice interactions had a causal effect on neural activity in the subsequent fixation period. To do this, we computed a test statistic measuring the correlation of the reward (or interaction) on each trial with fixation-period activity of the next trial and averaged it across sessions; then compared this statistic to a null ensemble obtained by randomly regenerating the reward sequence using the same rules as the original session.

The test statistic was defined by sparse logistic regression. For each session, we found the 20 neurons whose activity had the highest correlation with the previous trial's reward (or interaction), and applied unpenalized logistic regression to predict the reward (or interaction) from their activity. We defined the test statistic as the average log likelihood of this prediction over trials. We compared this test statistic to a null ensemble obtained by repeating 1000 times with regenerated reward sequences, choosing the 20 best neurons and prediction weights again each time. We infer that the correlation is significant if the test statistic is above the 99th percentile of the null distribution. The analysis was run multiple times, computing activity for different time bins.

Optogenetics

To assess whether the neural activity in the MOs and mPFC plays a causal role in the task, we applied transient optogenetics inactivations to these areas. Ai32 x PV-Cre (Ai32 [Jax #012569, RRID:IMSR_JAX: 012569] x PV-Cre [Jax #008069, RRID:IMSR_JAX: 008069) mice were implanted with glass cannulas (Doric MFC_200/230-0.48_2mm_MF1.25_FLT) bilaterally, using coordinates 2.2AP, 0.9 ML (figure 6, supplementary figure 8). In these mice, blue light excites PV interneurons, thereby inhibiting the pyramidal neurons (Cardin et al., 2010; Rikhye et al., 2021). To address a possible concern about the inactivation spreading beyond the target brain region (Li et al., 2019), the dataset consists of 6 mice in which the cannulas were installed on the surface of the brain targeting MOs (MOs group, figure 6 A) and 7 mice in which the cannulas were implanted in the same AP and DV location, but 1mm deeper than the first group, and slightly angled (mPFC group, figure 6 A). This is because the goal of the experiment with the MOs group was to see if inhibiting MOs specifically affects choice behaviour. The mPFC group thus also served

as a negative control for the MOs group: an effect that is specific to the MOs group could not arise due to deeper areas being inactivated. Due to a health issue, one mouse of the mPFC group was only used in the "half session" experiment but not in the "random inactivation" experiment.

We ran two sets of experiments (figure 6 B, C, figure 7A, B) with the same population of mice. In the first experiment, light was delivered on 20% of randomly chosen trials, starting with the go cue until the animal moved a wheel or for 4.8s, whichever was faster. On a further 20% of randomly chosen trials, light was delivered starting at the time of feedback for 1.6s. Since the two stimulations are independent of one another, inhibition happens during both the choice period and the outcome period on 4% of trials. Light consisted of a train of 10ms blue (473 nm) laser pulses separated by 30ms gaps, delivered through an optic fiber (NA = 0.22, Doric Lenses) using a diode laser (Omicron LuxX 473-100). The laser power was 63-110 mW/mm^2, measured at the tip of the patch cable that was then attached to the implanted cannula before the start of every experiment.

In the second set of experiments, stimulation was not applied on random trials. Instead, we randomly chose either the first part of the session (from trial 1 until trial 250), or the second half (from trial 251 until the end of the session) and applied the stimulation on every trial of the chosen session part at the time of feedback delivery for 1.6s.

Optogenetic data analysis

Analysis of the half-session opto experiment

We constructed learning curves by computing the proportion of choices equal to the correct choice, as a function of trial number relative to the block switch, averaging over switches in all sessions. We did so separately for block switches during the laser stimulation half-session and in the other half-session. The test statistic was defined as the difference between these two curves. To assess whether the difference between the two conditions is significant, we constructed a null ensemble of 100 randomizations by recalculating this statistic after resampled the half of each session where the laser was on, according to the same rule as for the real experiment.

Reaction time and firing rate relative to block switch

To estimate mean reaction time for different trials relative to block switch (Supplementary Figure 4A), we first identified all choice switch trials (i.e. trials N where $C_N \neq C_{N-1}$). To estimate the mean reaction time n trials before the switch, we then identified the subset of these switches for which all choices from trial N-n to trial N-1 where identical; this excluded cases where choice switched twice between trial N-n and trial N. Similarly, to estimate mean reaction times n trials after the switch, we identified the subset of switches for which all choices from trial N to trial N+n were identical. For each session and each n, we then took the median reaction time of these subsets of trials (N+n or N-n for each n), and subtracted the overall median reaction time for the session. We identified statistical significance by a t-test that this difference had mean 0.

To investigate the relationship of reaction time with the perseveration or reward-seeking decision variables of the PR model (Figure 2D), we computed a mean of the binned reaction times at 8 points equally distributed from $-\lambda_p$ to λ_p and subtract the mean of these. We deem the difference significant when it's different from 0 according to Wilcoxon signed rank test with 5% significance level.

Bibliography

Aarts, H., Verplanken, B., & van Knippenberg, A. (1998). Predicting Behavior From Actions in the Past: Repeated Decision Making or a Matter of Habit? *Journal of Applied Social Psychology*, *28*(15), 1355–1374. https://doi.org/10.1111/j.1559-1816.1998.tb01681.x

Akaishi, R., Umeda, K., Nagase, A., & Sakai, K. (2014). Autonomous Mechanism of Internal Choice Estimate Underlies Decision Inertia. *Neuron*, *81*(1), 195–206. https://doi.org/10.1016/j.neuron.2013.10.018

Ashby, F. G., Ennis, J. M., & Spiering, B. J. (2007). A neurobiological theory of automaticity in perceptual categorization. *Psychological Review*, *114*(3), 632–656. https://doi.org/10.1037/0033-295X.114.3.632

Ashby, F. G., Turner, B. O., & Horvitz, J. C. (2010). Cortical and basal ganglia contributions to habit learning and automaticity. *Trends in Cognitive Sciences*, *14*(5), 208–215. https://doi.org/10.1016/j.tics.2010.02.001

Averbeck, B., & O'Doherty, J. P. (2022). Reinforcement-learning in fronto-striatal circuits. *Neuropsychopharmacology*, *47*(1), Article 1. https://doi.org/10.1038/s41386-021-01108-0

Bari, B. A., Grossman, C. D., Lubin, E. E., Rajagopalan, A. E., Cressy, J. I., & Cohen, J. Y. (2019). Stable Representations of Decision Variables for Flexible Behavior. *Neuron*, *103*(5), 922-933.e7. https://doi.org/10.1016/j.neuron.2019.06.001

Barraclough, D. J., Conroy, M. L., & Lee, D. (2004). Prefrontal cortex and decision making in a mixed-strategy game. *Nature Neuroscience*, *7*(4), Article 4. https://doi.org/10.1038/nn1209

Barthas, F., & Kwan, A. C. (2017). Secondary Motor Cortex: Where 'Sensory' Meets 'Motor' in the Rodent Frontal Cortex. *Trends in Neurosciences*, *40*(3), 181–193. https://doi.org/10.1016/j.tins.2016.11.006

Behrens, T. E. J., Woolrich, M. W., Walton, M. E., & Rushworth, M. F. S. (2007). Learning the value of information in an uncertain world. *Nature Neuroscience*, *10*(9), Article 9. https://doi.org/10.1038/nn1954

Beron, C. C., Neufeld, S. Q., Linderman, S. W., & Sabatini, B. L. (2022). Mice exhibit stochastic and efficient action switching during probabilistic decision making. *Proceedings of the National Academy of Sciences*, *119*(15), e2113961119. https://doi.org/10.1073/pnas.2113961119

Bertelson, P. (1965). Serial choice reaction-time as a function of response versus signal-and-response repetition. *Nature*, *206*(980), 217–218. https://doi.org/10.1038/206217a0

Bogacz, R. (2020). Dopamine role in learning and action inference. *eLife*, *9*, e53262. https://doi.org/10.7554/eLife.53262

Botvinick, M., Wang, J. X., Dabney, W., Miller, K. J., & Kurth-Nelson, Z. (2020). Deep Reinforcement Learning and Its Neuroscientific Implications. *Neuron*, *107*(4), 603–616. https://doi.org/10.1016/j.neuron.2020.06.014

Burgess, C. P., Lak, A., Steinmetz, N. A., Zatka-Haas, P., Reddy, C. B., Jacobs, E. A. K., Linden, J. F., Paton, J. J., Ranson, A., Schröder, S., Soares, S., Wells, M. J., Wool, L. E., Harris, K. D., & Carandini, M. (2017). High-Yield Methods for Accurate Two-Alternative Visual Psychophysics in Head-Fixed Mice. *Cell Reports*, *20*(10), 2513–2524. https://doi.org/10.1016/j.celrep.2017.08.047

Cardin, J. A., Carlén, M., Meletis, K., Knoblich, U., Zhang, F., Deisseroth, K., Tsai, L.-H., & Moore, C. I. (2010). Targeted optogenetic stimulation and recording of neurons in vivo using cell-type-specific expression of Channelrhodopsin-2. *Nature Protocols*, *5*(2), Article 2. https://doi.org/10.1038/nprot.2009.228

Cho, R. Y., Nystrom, L. E., Brown, E. T., Jones, A. D., Braver, T. S., Holmes, P. J., & Cohen, J. D. (2002). Mechanisms underlying dependencies of performance on stimulus history in a two-alternative forced-choice task. *Cognitive, Affective & Behavioral Neuroscience*, *2*(4), 283–299. https://doi.org/10.3758/cabn.2.4.283

Constantino, S. M., & Daw, N. D. (2015). Learning the opportunity cost of time in a patch-foraging task. *Cognitive, Affective, & Behavioral Neuroscience*, *15*(4), 837–853. https://doi.org/10.3758/s13415-015-0350-y

Corrado, G. S., Sugrue, L. P., Seung, H. S., & Newsome, W. T. (2005). LINEAR-NONLINEAR-POISSON MODELS OF PRIMATE CHOICE DYNAMICS. *Journal of the Experimental Analysis of Behavior*, *84*(3), 581–617. https://doi.org/10.1901/jeab.2005.23-05

Costa, V. D., & Averbeck, B. B. (2020). Primate Orbitofrontal Cortex Codes Information Relevant for Managing Explore-Exploit Tradeoffs. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *40*(12), 2553–2561. https://doi.org/10.1523/JNEUROSCI.2355-19.2020

Daw, N. D. (2011). Trial-by-trial data analysis using computational models. In *Decision Making, Affect, and Learning*. Oxford University Press. https://doi.org/10.1093/acprof:oso/9780199600434.003.0001

Daw, N. D., Niv, Y., & Dayan, P. (2005). Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nature Neuroscience*, *8*(12), Article 12. https://doi.org/10.1038/nn1560

Daw, N. D., O'Doherty, J. P., Dayan, P., Seymour, B., & Dolan, R. J. (2006). Cortical substrates for exploratory decisions in humans. *Nature*, *441*(7095), Article 7095. https://doi.org/10.1038/nature04766

Dickinson, A. (1985). Actions and habits: The development of behavioural autonomy. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, *308*(1135), 67–78. https://doi.org/10.1098/rstb.1985.0010

Ebitz, R. B., Albarran, E., & Moore, T. (2018). Exploration Disrupts Choice-Predictive Signals and Alters Dynamics in Prefrontal Cortex. *Neuron*, *97*(2), 450-461.e9. https://doi.org/10.1016/j.neuron.2017.12.007

Economo, M. N., Clack, N. G., Lavis, L. D., Gerfen, C. R., Svoboda, K., Myers, E. W., & Chandrashekar, J. (2016). A platform for brain-wide imaging and reconstruction of individual neurons. *eLife*, *5*, e10566. https://doi.org/10.7554/eLife.10566

Elber-Dorozko, L., & Loewenstein, Y. (2018). Striatal action-value neurons reconsidered. *eLife*, *7*, e34248. https://doi.org/10.7554/eLife.34248

Erlich, J. C., Bialek, M., & Brody, C. D. (2011). A cortical substrate for memory-guided orienting in the rat. *Neuron*, *72*(2), 330–343. https://doi.org/10.1016/j.neuron.2011.07.010

Evans, J. St. B. T., & Stanovich, K. E. (2013). Dual-Process Theories of Higher Cognition: Advancing the Debate. *Perspectives on Psychological Science*, *8*(3), 223–241. https://doi.org/10.1177/1745691612460685

Gershman, S. J. (2020). Origin of perseveration in the trade-off between reward and complexity. *Cognition*, *204*, 104394. https://doi.org/10.1016/j.cognition.2020.104394

Gold, J. M., Waltz, J. A., Prentice, K. J., Morris, S. E., & Heerey, E. A. (2008). Reward processing in schizophrenia: A deficit in the representation of value. *Schizophrenia Bulletin*, *34*(5), 835–847. https://doi.org/10.1093/schbul/sbn068

Gore, J. L., Dorris, M. C., & Munoz, D. P. (2002). Time course of a repetition effect on saccadic reaction time in non-human primates. *Archives Italiennes De Biologie*, *140*(3), 203–210.

Greenstreet, F., Vergara, H. M., Pati, S., Schwarz, L., Wisdom, M., Marbach, F., Johansson, Y., Rollik, L., Moskovitz, T., Clopath, C., & Stephenson-Jones, M. (2022). *Action prediction error: A value-free dopaminergic teaching signal that drives stable learning* (p. 2022.09.12.507572). bioRxiv. https://doi.org/10.1101/2022.09.12.507572

Guo, Z. V., Li, N., Huber, D., Ophir, E., Gutnisky, D., Ting, J. T., Feng, G., & Svoboda, K. (2014). Flow of Cortical Activity Underlying a Tactile Decision in Mice. *Neuron*, *81*(1), 179–194. https://doi.org/10.1016/j.neuron.2013.10.020

Hardwick, R. M., Forrence, A. D., Krakauer, J. W., & Haith, A. M. (2019). Time-dependent competition between goal-directed and habitual response preparation. *Nature Human Behaviour*, *3*(12), 1252–1262. https://doi.org/10.1038/s41562-019-0725-0

Harris, K. D. (2021). *Nonsense correlations in neuroscience* (p. 2020.11.29.402719). bioRxiv. https://doi.org/10.1101/2020.11.29.402719

Harris, K. D., & Yuan, A. E. (2024). *Tests for partial correlation between repeatedly observed nonstationary nonlinear timeseries* (arXiv:2106.07096). arXiv. https://doi.org/10.48550/arXiv.2106.07096

Hattori, R., Danskin, B., Babic, Z., Mlynaryk, N., & Komiyama, T. (2019). Area-Specificity and Plasticity of History-Dependent Value Coding During Learning. *Cell*, *177*(7), 1858-1872.e15. https://doi.org/10.1016/j.cell.2019.04.027

Herrnstein, R. J. (1970). On the law of effect. *Journal of the Experimental Analysis of Behavior*, *13*(2), 243–266. https://doi.org/10.1901/jeab.1970.13-243

Hikosaka, O., Rand, M. K., Miyachi, S., & Miyashita, K. (1995). Learning of sequential movements in the monkey: Process of learning and retention of memory. *Journal of Neurophysiology*, *74*(4), 1652–1661. https://doi.org/10.1152/jn.1995.74.4.1652

Ito, M., & Doya, K. (2009). Validation of Decision-Making Models and Analysis of Decision Variables in the Rat Basal Ganglia. *Journal of Neuroscience*, *29*(31), 9861–9874. https://doi.org/10.1523/JNEUROSCI.6157-08.2009

Ito, M., & Doya, K. (2011). Multiple representations and algorithms for reinforcement learning in the cortico-basal ganglia circuit. *Current Opinion in Neurobiology*, *21*(3), 368–373. https://doi.org/10.1016/j.conb.2011.04.001

Jun, J. J., Steinmetz, N. A., Siegle, J. H., Denman, D. J., Bauza, M., Barbarits, B., Lee, A. K., Anastassiou, C. A., Andrei, A., Aydın, Ç., Barbic, M., Blanche, T. J., Bonin, V., Couto, J., Dutta, B., Gratiy, S. L., Gutnisky, D. A., Häusser, M., Karsh, B., … Harris, T. D. (2017). Fully integrated silicon probes for high-density recording of neural activity. *Nature*, *551*(7679), Article 7679. https://doi.org/10.1038/nature24636

Kahneman, D. (2011). *Thinking, Fast and Slow*. https://www.amazon.com/Thinking-Fast-Slow-Daniel-Kahneman/dp/0374275637

Kim, H., Sul, J. H., Huh, N., Lee, D., & Jung, M. W. (2009). Role of Striatum in Updating Values of Chosen Actions. *Journal of Neuroscience*, *29*(47), 14701–14712. https://doi.org/10.1523/JNEUROSCI.2728-09.2009

Lai, L., & Gershman, S. J. (2024). Human decision making balances reward maximization and policy compression. *PLOS Computational Biology*, *20*(4), e1012057. https://doi.org/10.1371/journal.pcbi.1012057

Lau, B., & Glimcher, P. W. (2005). Dynamic Response-by-Response Models of Matching Behavior in Rhesus Monkeys. *Journal of the Experimental Analysis of Behavior*, *84*(3), 555–579. https://doi.org/10.1901/jeab.2005.110-04

Le, N. M., Yildirim, M., Wang, Y., Sugihara, H., Jazayeri, M., & Sur, M. (2023). Mixtures of strategies underlie rodent behavior during reversal learning. *PLOS Computational Biology*, *19*(9), e1011430. https://doi.org/10.1371/journal.pcbi.1011430

Lee, D., Seo, H., & Jung, M. W. (2012). Neural Basis of Reinforcement Learning and Decision Making. *Annual Review of Neuroscience*, *35*(1), 287–308. https://doi.org/10.1146/annurev-neuro-062111-150512

Lefebvre, G., Lebreton, M., Meyniel, F., Bourgeois-Gironde, S., & Palminteri, S. (2017). Behavioural and neural characterization of optimistic reinforcement learning. *Nature Human Behaviour*, *1*(4), Article 4. https://doi.org/10.1038/s41562-017-0067

Li, N., Chen, S., Guo, Z. V., Chen, H., Huo, Y., Inagaki, H. K., Chen, G., Davis, C., Hansel, D., Guo, C., & Svoboda, K. (2019). Spatiotemporal constraints on optogenetic inactivation in cortical circuits. *eLife*, *8*, e48622. https://doi.org/10.7554/eLife.48622

Li, N., Chen, T.-W., Guo, Z. V., Gerfen, C. R., & Svoboda, K. (2015). A motor cortex circuit for motor planning and movement. *Nature*, *519*(7541), Article 7541. https://doi.org/10.1038/nature14178

Miller, K. J., Botvinick, M. M., & Brody, C. D. (2021). *From predictive models to cognitive models: Separable behavioral processes underlying reward learning in the rat* (p. 461129). bioRxiv. https://doi.org/10.1101/461129

Miller, K. J., Botvinick, M. M., & Brody, C. D. (2022). Value representations in the rodent orbitofrontal cortex drive learning, not choice. *eLife*, *11*, e64575. https://doi.org/10.7554/eLife.64575

Miller, K. J., Eckstein, M., Botvinick, M. M., & Kurth-Nelson, Z. (2023). *Cognitive Model Discovery via Disentangled RNNs*. https://doi.org/10.1101/2023.06.23.546250

Miller, K. J., Shenhav, A., & Ludvig, E. A. (2019). Habits without values. *Psychological Review*, *126*(2), 292–311. https://doi.org/10.1037/rev0000120

Murakami, M., Vicente, M. I., Costa, G. M., & Mainen, Z. F. (2014). Neural antecedents of self-initiated actions in secondary motor cortex. *Nature Neuroscience*, *17*(11), 1574–1582. https://doi.org/10.1038/nn.3826

Olsen, S. R., Bortone, D. S., Adesnik, H., & Scanziani, M. (2012). Gain control by layer six in cortical circuits of vision. *Nature*, *483*(7387), 47–52. https://doi.org/10.1038/nature10835

Pachitariu, M., Sridhar, S., & Stringer, C. (2023). *Solving the spike sorting problem with Kilosort* (p. 2023.01.07.523036). bioRxiv. https://doi.org/10.1101/2023.01.07.523036

Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M., & D, H. K. (2016). *Kilosort: Realtime spike-sorting for extracellular electrophysiology with hundreds of channels* (p. 061481). bioRxiv. https://doi.org/10.1101/061481

Padoa-Schioppa, C. (2013). Neuronal origins of choice variability in economic decisions. *Neuron*, *80*(5), 1322–1336. https://doi.org/10.1016/j.neuron.2013.09.013

Palminteri, S., Wyart, V., & Koechlin, E. (2017). The Importance of Falsification in Computational Cognitive Modeling. *Trends in Cognitive Sciences*, *21*(6), 425–433. https://doi.org/10.1016/j.tics.2017.03.011

Ragan, T., Kadiri, L. R., Venkataraju, K. U., Bahlmann, K., Sutin, J., Taranda, J., Arganda-Carreras, I., Kim, Y., Seung, H. S., & Osten, P. (2012). Serial two-photon tomography for automated ex vivo mouse brain imaging. *Nature Methods*, *9*(3), 255–258. https://doi.org/10.1038/nmeth.1854

Riefer, P. S., Prior, R., Blair, N., Pavey, G., & Love, B. C. (2017). Coherency-maximizing exploration in the supermarket. *Nature Human Behaviour*, *1*(1), Article 1. https://doi.org/10.1038/s41562-016-0017

Rikhye, R. V., Yildirim, M., Hu, M., Breton-Provencher, V., & Sur, M. (2021). Reliable Sensory Processing in Mouse Visual Cortex through Cooperative Interactions between Somatostatin and Parvalbumin Interneurons. *The Journal of Neuroscience*, *41*(42), 8761–8778. https://doi.org/10.1523/JNEUROSCI.3176-20.2021

Samejima, K., Ueda, Y., Doya, K., & Kimura, M. (2005). Representation of Action-Specific Reward Values in the Striatum. *Science*, *310*(5752), 1337–1340. https://doi.org/10.1126/science.1115270

Sarkka, S. (2013). *BAYESIAN FILTERING AND SMOOTHING*.

Seo, H., Barraclough, D. J., & Lee, D. (2007). Dynamic signals related to choices and outcomes in the dorsolateral prefrontal cortex. *Cerebral Cortex (New York, N.Y.: 1991)*, *17 Suppl 1*, i110-117. https://doi.org/10.1093/cercor/bhm064

Steinmetz, N. A., Aydin, C., Lebedeva, A., Okun, M., Pachitariu, M., Bauza, M., Beau, M., Bhagat, J., Böhm, C., Broux, M., Chen, S., Colonell, J., Gardner, R. J., Karsh, B., Kloosterman, F., Kostadinov, D., Mora-Lopez, C., O'Callaghan, J., Park, J., … Harris, T. D. (2021). Neuropixels 2.0: A miniaturized high-

density probe for stable, long-term brain recordings. *Science*, *372*(6539), eabf4588. https://doi.org/10.1126/science.abf4588

Steinmetz, N. A., Zatka-Haas, P., Carandini, M., & Harris, K. D. (2019). Distributed coding of choice, action and engagement across the mouse brain. *Nature*, *576*(7786), Article 7786. https://doi.org/10.1038/s41586-019-1787-x

Sugrue et al. (2004). *Matching Behavior and the Representation of Value in the Parietal Cortex | Science*. https://www.science.org/doi/10.1126/science.1094765

Sul, J. H., Jo, S., Lee, D., & Jung, M. W. (2011). Role of rodent secondary motor cortex in value-based action selection. *Nature Neuroscience*, *14*(9), Article 9. https://doi.org/10.1038/nn.2881

Sul, J. H., Kim, H., Huh, N., Lee, D., & Jung, M. W. (2010). Distinct Roles of Rodent Orbitofrontal and Medial Prefrontal Cortex in Decision Making. *Neuron*, *66*(3), 449–460. https://doi.org/10.1016/j.neuron.2010.03.033

Sutton, R. S., & Barto, A. G. (1998). Reinforcement Learning: An Introduction. *IEEE Transactions on Neural Networks*, *9*(5), 1054–1054. https://doi.org/10.1109/TNN.1998.712192

Thorndike, E. L. (1911). Animal Intelligence: Experimental Studies. *MacMillan, New York*. https://doi.org/10.5962/bhl.title.55072

Waldschmidt, J. G., & Ashby, F. G. (2011). Cortical and striatal contributions to automaticity in information-integration categorization. *NeuroImage*, *56*(3), 1791–1802. https://doi.org/10.1016/j.neuroimage.2011.02.011

Wang, J. X., Kurth-Nelson, Z., Kumaran, D., Tirumala, D., Soyer, H., Leibo, J. Z., Hassabis, D., & Botvinick, M. (2018). Prefrontal cortex as a meta-reinforcement learning system. *Nature Neuroscience*, *21*(6), 860–868. https://doi.org/10.1038/s41593-018-0147-8

Wang, Q., Ding, S.-L., Li, Y., Royall, J., Feng, D., Lesnar, P., Graddis, N., Naeemi, M., Facer, B., Ho, A., Dolbeare, T., Blanchard, B., Dee, N., Wakeman, W., Hirokawa, K. E., Szafer, A., Sunkin, S. M., Oh, S. W., Bernard, A., … Ng, L. (2020). The Allen Mouse Brain Common Coordinate Framework: A 3D Reference Atlas. *Cell*, *181*(4), 936-953.e20. https://doi.org/10.1016/j.cell.2020.04.007

Wood, W., & Rünger, D. (2016). Psychology of Habit. *Annual Review of Psychology*, *67*(1), 289–314. https://doi.org/10.1146/annurev-psych-122414-033417

Yang, J.-H., & Kwan, A. C. (2021). Secondary motor cortex: Broadcasting and biasing animal's decisions through long-range circuits. *International Review of Neurobiology*, *158*, 443–470. https://doi.org/10.1016/bs.irn.2020.11.008

Yule, G. U. (1926). Why do we Sometimes get Nonsense-Correlations between Time-Series?—A Study in Sampling and the Nature of Time-Series. *Journal of the Royal Statistical Society*, *89*(1), 1–63. https://doi.org/10.2307/2341482

Acknowledgements

We thank Charu Reddy and Matteo Carandini for experimental help and discussions. This work was supported by Wellcome Trust grants 223144, 205093 and ERC grant 694401 to K.D.H.

Author contributions

	Lebedeva	Wang	Funnell	Terry	Oh	Miller	Harris
Conceptualization	•					•	•

Methodology	•				•	•	•
Software	•					•	
Formal Analysis	•				•	•	•
Investigation	•	•	•	•	•	•	
Resources							•
Writing – Original Draft	•					•	•
Writing – Review & Editing	•					•	•
Visualization	•					•	
Supervision						•	•
Funding Acquisition							•

Competing Interests

The authors declare no competing interests.

Supplementary figures

Supplementary figure 1. Diagnostics for all models. Two behavioural diagnostic plots are shown for each model: the learning curve (fraction of choices to the correct side as a function of time since a block switch, as in Fig. 1D); cumulative proportion of the lengths of unrewarded sequences, as in Fig. 1E. Grey lines, data for individual fits; black line, average. A, Models in which the first diagnostic is dissimilar to mice. B, Models in which the first diagnostic is dissimilar to mice. C, Models in which both diagnostics are similar to mouse diagnostics. Note, all models in C are perseverative.

Action probability heatmap

Supplementary figure 2. Similarity of choice predictions for the 5 winning models. Each panel shows a pseudocolor histogram summarizing the frequency that every pair of choice probabilities was predicted by two models, across all trials, sessions, and mice. For example, the intensity in the top left corner of the top right plot shows the frequency of trials on which the PR model predicted p(Right)=0 and the ideal observer model predicted p(Right)=1.

Supplementary figure 3 | Fit parameters: PR model. The parameter fits for the PR model for each mouse.

A

B

Supplementary figure 4 | Changes in response time and neural activity relative to behavior switches. A, median response time as a function of the number of trials relative to the choice switch, minus overall median response time. An x-axis value of n indicates median reaction times for all trials exactly n trials following a switch, without a second intervening switch. Thus the trial following the sequence LRRR would go into bin 3 only, while the last trial of sequence LRLR go into bin 1 only. Red stars indicate a significant difference from 0. **B**, Mean fixation-period activity of the 20 units that correlate best with either right or left choice, as a function of the number of trials relative to a choice switches from right to left or from left to right, on average across all MOs recordings.

Supplementary figure 5. Top, number of units recorded in each area. Bottom, number of recording sessions per area.

Supplementary figure 6. Causal effect of feedback (i.e. reward or white noise) on neural activity. To show a causal effect we took advantage of the fact that feedback delivery was randomized (80% or 20% reward probabilities for correct or incorrect choices). Black line shows log likelihood for predicting the actual feedback from neural population activity in a 200 ms time bin aligned to feedback time (left) or to the subsequent trial's go cue (right). Blue shading indicates the fixation period. Gray lines indicate null ensemble predicting randomly regenerated feedback sequences, drawn using the same behavior-dependent probabilities as the actual sequence. Red stars indicate time bins in which the likelihood for predicting the actual choice exceeds 99% of the null ensemble. Note that even if the neural activity encodes a prediction of upcoming reward magnitude, this analysis could not show a causal effect of feedback on activity before feedback delivery: because the null ensemble was randomly generated with the same probabilities as the actual feedback, there is no way that neural activity could predict the actual feedback better than null, even if it could predict whether the animal would choose the correct or incorrect side.

Supplementary figure 7. Causal effect of choice-feedback interaction on neural population activity, assessed by the same method as Figure 6. The analysis proceeds exactly as in Supp Fig. 6, but predicting the choice-feedback interaction rather than just the feedback value. Again, black lines indicate prediction of actual value, gray lines indicate the null ensemble obtained by resampling feedback values with the probabilities determined by the mouses actual choices, then recomputing the choice-feedback interaction. Blue rectangle, fixation period.

Supplementary figure 8. Estimated cannula positions for optogenetic experiments. Top two panels: experiments with inactivation on random trials. Bottom panel: experiments with inactivation in random half-experiments.