Imaging flow cytometry enables label-free cell sorting of morphological variants from

unculturable bacterial populations.

Keywords: autofluorescence, imaging flow cytometry, cell sorting, non-model system,

Pasteuria ramosa

Authors

Daniel Vocelle[1], Lindsey Thompson[2], Matthew P. Bernard[1] & Nina Wale[12,3,4,+]

1. Department of Pharmacology & Toxicology, Michigan State University, East Lansing, MI 48824, USA

2. Department of Microbiology, Genetics & Immunology, Michigan State University, East Lansing, Michigan,

USA 48824

3. Department of Integrative Biology, Michigan State University, East Lansing, Michigan, USA 48824

4. Program in Ecology, Evolution & Behavior, Michigan State University, East Lansing, Michigan, USA 48824

+corresponding author

Contributions

DV and NW conceptualized the research.

DV and LT performed the experiments.

DV designed the analysis, with consultation from MB.

DV and NW analyzed the data.

NW wrote the manuscript.

DV, LT, MB edited the manuscript.

Data availability

Description: data will be made available upon article acceptance.

Abstract

Bacterial populations often display remarkable morphological heterogeneity. Flow cytometric

cell sorting (often called FACS) is an important tool for understanding this diversity. FACs allows

researchers to obtain pure samples of each morphological variant (or morphotype) that is

present within a mixed population of cells and thus permits each morphotype to be phenotyped.

In FACS, cells are first labeled with fluorescent markers such as antibodies or transgenic

constructs, and then separated out based on their possession of these labels. However, since

the development of fluorescent labels requires *a priori* knowledge of bacterial biology, it is often

impossible to apply FACS to understudied and/or unculturable bacteria. This challenge has

limited our capacity to investigate the biology of bacterial size and shape in all but a small,

largely culturable subset of bacterial taxa. Here, we present an innovative strategy that permits

label-free cell sorting of bacterial morphotypes, using an unculturable, pleiomorphic pathogen

(*Pasteuria ramosa*) as a model bacterium. We show that imaging flow cytometry (IFC) can be

used to systematically identify light-scattering and autofluorescence "signatures" of bacterial

morphotypes, on which basis cell sorting can be conducted. Critically, our IFC-enabled cell

sorting strategy yields samples of sufficient purity (>90%) for common downstream analyses

e.g., "-omics" analyses. Our work represents an innovative application of IFC and provides an

economical, widely applicable solution to a central problem in the study of bacterial diversity.

Importance

Bacteria come in many different shapes and sizes. Why this morphological variation exists is a longstanding question in microbiology, but it remains difficult to answer. To phenotype different morphological variants (morphotypes) within a bacterial population, we need to separate them from one another. This is normally achieved using flow cytometric cell-sorting, whereby morphotypes are labelled with fluorescent antibodies and separated on the basis of their differential fluorescence. Unfortunately, it is difficult to develop fluorescent-labels specific to unculturable or poorly studied bacteria because of the limited availability of appropriate molecular tools. Here, we demonstrate that imaging flow cytometry can be used to design and validate label-free cell sorting strategies. Recently, there has been a resurgence of interest in bacterial morphological diversity and a call to expand its study across the tree of life. Our work will help microbiologists to answer this call.

Introduction

Bacterial populations often contain cells varying in size and shape, and this morphological

heterogeneity can have a profound impact on bacterial fitness (1–3). For example, different

forms of bacteria (morphotypes) may perform distinct ecological "roles" or exhibit differential

susceptibility to environmental stress, permitting population persistence in the face of such

stress (4–7). To fully understand the causes and consequences of morphological variation

within bacterial populations, we need to extend our investigations beyond well-studied taxa and

in vitro environments, which can suppress morphological diversity within a bacterial population

(2, 8–10). Here, we present imaging flow cytometry (IFC)-enabled cell sorting, a novel

procedure that allows for the purification of distinct morphotypes of unculturable bacteria living

in complex milieu.

Efforts to understand the mechanistic basis and functional significance of morphological

heterogeneity in a bacterial population often begin with isolating distinct bacterial morphotypes

from each other and/or their environmental milieu, via flow cytometric cell sorting (or,

colloquially, fluorescence activated cell sorting, FACS). However, the application of FACS to

unculturable bacteria or those found in complex samples (e.g., soil, host tissues) is a challenge.

A chicken-and-egg-scenario exists. In FACS, different cell populations are tagged with distinct

fluorescent labels (e.g., fluorescently labelled antibodies) on whose basis the populations are

separated. However, the development of morphotype-specific labels requires the

characterization of morphotype-specific antigens, which itself requires that morphotypes be

separated and studied. In the absence of morphotype specific labels, 'general' membrane

permeable labels (e.g., CFSE, Mitotracker, Hoechst), which bind to molecules present in most

cells (e.g., amines, DNA/RNA, or mitochondria), could be used to discriminate morphotypes in

certain contexts (11). However, because general labels indiscriminately permeate all cells within

a sample, they are unlikely to be helpful in distinguishing a specific morphotype if it is present in

a complex, cell-rich milieu such as host-tissue or soil.

A potential solution to the problem of sorting bacterial morphotypes in the absence of labels is to

exploit intrinsic variation in their light-scattering and -fluorescence properties (i.e., "light

signatures"). Cells differing in size and granularity differentially scatter light, and the degree of

autofluorescence may also vary among morphotypes (12). For example, viable *Bacillus*

anthracis spores exhibit a distinctive autofluorescent signature upon excitation with UV light

(13). Recent advances in imaging flow cytometry (IFC) simplify the identification of intrinsic light

signatures of morphologically distinct cells by generating images of each cell and measures of

their light-scattering and autofluorescence properties. Systematic analysis of these data allows

for the identification of morphotype-specific light signatures and has been used to identify

morphologically distinct subpopulations of mammalian and bacterial cells (14–17).

Here, using the pleiomorphic bacterial pathogen *Pasteuria ramosa* (hereafter, *Pasteuria*) as a

model, we demonstrate that IFC can be used to efficiently design and validate strategies for

label-free sorting of bacterial morphotypes. *Pasteuria* is a spore-forming pathogen of

zooplankton that exhibits striking morphological variation during its life cycle (Table 1, (18)).

Longstanding hypotheses vis-à-vis the "division of labor" among different morphotypes within

the *Pasteuria* bacterial population (19) remain untested, largely because it is difficult to separate

bacteria from host material for functional -omics and *in vitro* study. By combining state-of-the-art

IFC instrumentation, which allows us to rapidly derive thousands of images of bacterial cells,

and machine-learning algorithms, we systematically designed and validated autofluorescence-

based FACS strategies that yield high-purity samples of each of four distinct morphotypes. Our

work provides proof of principle that IFC-enabled sorting can be used to overcome the first

barrier to understanding morphological heterogeneity in bacterial species that cannot be readily

labeled.

	name (abbreviation)	Function (*hypothesized)	largest dimension (range, μm)
	Activated (A)	host attachment	-
	`cauliflower' (C)	in-host growth, host manipulation*	4-6.7
	`grape-seed' (G)	sporulation	4-6
	mature spore (S)	transmission, environmental persistence	4.2-5.4

153 **Table 1. Variation in the form and function of *Pasteuria ramosa* morphotypes.** Form and
154 function information extracted from (18, 20, 21). Illustrations are schematic representations of
155 morphotypes and are not to scale. *Indicates function is hypothesized.

Materials and methods

158 Our goal was to design and validate cell sorting strategies suitable for the purification of 4

159 distinct *Pasteuria* morphotypes from the homogenate of an infected host i.e., a sample

160 containing several *Pasteuria* morphotypes, host cells, food (algae) present in the host gut, and

161 host microbiota. Below we describe our approach in general terms. The final procedure and

162 results of our validation steps are described in the Results.

Host & Parasites

Throughout, we used *Daphnia magna* (HuHO-2 genotype) hosts and *Pasteuria* (strain C1),

originally collected in Hungary and Russia, respectively (22, 23). Both were generously donated

by D. Ebert (U. Basel, Switzerland).

To generate samples of *Pasteuria*, we experimentally infected *Daphnia magna*. Throughout, we

used a standard infection procedure. Briefly, we shook *Pasteuria* spores at 1500 rpm, overnight.

On day 0 of the experiment, we filled a vessel (e.g., 24 well-plate or beaker) with media (ADaM

or COMBO), added 1-10 *D. magna* <24 hours old and ~37,500 *Pasteuria* spores per host (24).

On the following day, we added a second equivalently sized dose of spores. We fed hosts

increasing quantities of either *Scenedesmus obliquus* or *Ankistrodesmus falcatus* as they aged,

following (25), and maintained them for 30-60 days. To diagnose *Pasteuria* infection, we

assessed hosts by eye for signs of infection, including discoloration, lack of offspring and

enlargement (26), and then confirmed it using microscopy.

Instruments

We used a Thermofisher Attune CytPix flow cytometer (cytometer, hereafter) for IFC. This

cytometer is equipped with a high-speed, brightfield camera (20x objective, 0.45 numerical

aperture, 0.3 micron/pixel resolution), four lasers and 14 filters (Table S1). Acoustic focusing

allows for the alignment of cells and particles (hereafter, events) in a narrow depth of field. As a

result, this machine can acquire in-focus images an order of magnitude faster than cytometers

with hydrodynamic-focusing (27).

For cell sorting, we used a BD Influx equipped with 5 lasers and 12 filters (Table S1). The

default measure of fluorescence on the BD Influx is "pulse height" (rather than area). Since our

goal was to use the imaging cytometer to identify fluorescence parameters that could be

employed in cell sorting, we used pulse height as our measure of fluorescence on both

machines.

Data acquisition

Immediately following diagnosis, we prepared infected animals for flow cytometric analysis.

First, each animal was washed in PBS (no Ca+ or Mg+). We transferred 1-4 animals, depending

on the degree of infection and an initial assessment of morphotype composition via microscopy,

into a 1.5mL Eppendorf tube containing PBS. We homogenized the animals briefly with an

electric pestle (~20 seconds, to avoid damaging cells) to create a slurry. We diluted the sample

in 500mL Focusing Buffer (Thermofisher Scientific, catalog no. A24904) or PBS, which have

equivalent refractive indices. We ran samples at 12.5mL/min until we acquired 30,000 events.

Between runs, we twice flushed the cytometer line to prevent sample carryover.

Pre-sort analysis: initial gating & image-processing

We first excluded ("gated out") algae on the basis of their chlorophyll autofluorescence (high

emission in channel R3, see Table S1). We then selected ("gated in") the events that (i) were

successfully photographed and processed and, furthermore, contained only one particle. This

yielded a dataset of single events, each of which was associated with an image and a set of

autofluorescence measurements.

To analyze the image data, we employed inbuilt Attune Image-Processing Software (version

6.0.1). We used the software's bead masking model (Beads_Only_Full_Resolution_v1) which is

optimized for objects 2-5mm. We chose this algorithm because (i) an alternative model

optimized for objects >5mm, excluded ~70% of our target objects and (ii) because we found that

the bead model could process a variety of *P. ramosa* morphotypes, including those that

exceeded 5mm (e.g., cauliflowers). Image-processing yielded a matrix containing the values of

22 imaging parameters including a variety of size and "texture" metrics (Table S2). In total,

including the fluorescent parameters (i.e., pulse height, area, and width) each event was

associated with 74 parameters. Before conducting further analysis, we rescaled each parameter

linearly between 0 and 1.

We visualized the data using Uniform Manifold Approximation Projection for Dimension

Reduction (UMAP) (28), implemented in FCS Express (De Novo Software). Briefly, this

algorithm allows for the visualization of high-dimensional data in a 2D plane. By inspecting

distinct regions of the UMAP, and the imaging data with which it was associated, we found that

different morphotypes clustered in distinct areas of the UMAP. Preliminary analysis of this kind

revealed that certain image parameters were particularly informative for distinguishing different

morphotypes (Table S2). We employed this subset of parameters in further analyses, described

below.

Pre-sort analysis: identifying morphotype-specific parameters for cell sorting

For each sample, we used IFC to identify gates that would enable us to sort a specific

morphotype ("target morphotype", hereafter) from the remainder of the sample.

First, we visualized the morphological complexity of the sample using a UMAP analysis of all the

scattering and fluorescent parameters, as well as several imaging parameters identified in

preliminary analyses as being most-informative (Table S2, S3). We then performed cluster

analysis to identify (i) a primary gate, on which basis we could distinguish our target morphotype

from other bacterial cells, and (ii) subsequent gate(s), on which basis we could distinguish our

target morphotype from debris (see Results). Specifically, we used FlowSOM - a neural

network-like algorithm that identifies clusters of events characterized by distinct (combinations

of) parameters (29). We set the consensus clustering parameter to between 8-20, depending on

the sample, and otherwise used default settings (Table S3).

The above-described procedure was repeated on a sample-by-sample basis. That is, the

parameters used to define the gates varied among samples (particularly in the initial stages of

our work, Table S4). Due to differences in the autofluorescence of cells among samples, the

exact position of the gates varied from sample to sample. Importantly, however, the gate -

identification and -validation strategy - which is our focus here - remained the same across

samples.

Cell sorting

Prior to cell sorting, we transferred the sample to a 5 mL FACS tube (Corning, catalog no.

352063) and filtered it through a 100mm mesh filter. We then used the gating strategy designed

using IFC (or a minor variation of the strategy, see Results, Fig. S1), to sort each of the

morphotypes in the sample that were of sufficient abundance (see Table S4 for number of

events sorted per sample).

Post-sort analysis: quantification of sample purity

We verified the purity of each sorted sample using IFC. Briefly, after running the sorted sample

on the imaging cytometer, the data was first processed as described above and then "down-

sampled" to a subset of ~500 events (with a few exceptions, see Table S4). Using the image

data, we manually classified each of the events in the downsampled dataset as an activated-,

cauliflower-, grape- or spore- morphotype of *Pasteuria* (Table 1) or as unidentified debris.

Finally, we calculated the purity of the sorted sample i.e., the proportion of total events in the

down-sampled dataset that were of the target morphotype. This purity measure is conservative

given that we include unidentified debris such as salts in our total event count.

Results

IFC enables the identification of cell sorting strategies that exploit morphotype-specific

light signatures

We established a robust procedure for identifying light signatures that can be harnessed for the

cell sorting of morphologically distinct bacterial morphotypes, in the absence of labels. To

demonstrate this approach, we provide a worked example (Fig. 1). In this example, our goal

was to sort a spore (S) population from a sample comprising host (*Daphnia*) material, host-

microbiota and non-target morphotypes of *Pasteuria* (cauliflower (C), grape (G)). Worked

examples of the procedure as applied to activated, cauliflower and grape stages are provided in

the supplement (Fig. S1).

First, we performed a dimensionality reduction (UMAP) analysis to visualize the data in 2D

space (Fig. 1A). To identify where in this 2D space each *Pasteuria* morphotype lay, we imposed

the results of a cluster analysis onto the UMAP (Fig. 1A, colors/numbers indicate clusters). This

cluster analysis was performed using all light-scattering and -fluorescence parameters and a

subset of the imaging parameters (Table S2). We manually classified each cluster as either a

morphotype-rich or debris-rich cluster using the image data (Fig. 1B). For example, since spores

appeared in most of the images associated with cluster 3, we designated cluster 3 the spore-

rich cluster (S-cluster; Fig. 1B, purple outline); similarly, clusters 1 and 2 were designated

grape- and cauliflower- rich clusters, respectively.

Next, we used the cluster analysis to identify light-scattering and fluorescence parameters that

characterized our target morphotype vs. the other morphotypes (C, G). Analysis of the median

parameter values associated with each of the morphotype clusters (Fig. 1Ci), suggested that the

different morphotypes could be broadly distinguished by the light scattering parameters FSC

and SSC (Fig.1Ci, F,S), consistent with their differences in size and granularity (Table 1). The

distribution of G-, C- and S- cluster events in FSC-SSC space confirmed this inference (Fig.

1Cii, Ciii). We thus used FSC-SSC to define our primary gate ("S1", Fig. 1Cii).

Inspection of the images associated with the events in the primary gate revealed that it included

debris as well as our target morphotype (Fig. 1Di). We therefore sought to identify a secondary

gate that could be used to distinguish the target morphotype from debris. First, we inspected the

images associated with the events in the target cluster, S1. This analysis revealed that there

was a gradient within the cluster: on the left lay a largely homogenous population of spores (Fig.

1Dii, region A, top images); whereas on the right we found a mixed population of spores and

other non-target objects (Fig. 1Dii, region B; bottom images). To identify which of the

fluorescence parameters shared by the sorter and the cytometer (Table S1) could differentiate

events in regions A and B we recolored the events in each cluster by their fluorescence

intensity in each of the shared channels (Fig. Diii, 4 parameters shown for readability; all

parameters are displayed in Fig. S2). This analysis revealed that V1 could distinguish the spore-

rich region A from the mixed region B and could therefore be used in combination with FSC to

define a secondary gate. Finally, we established the position of this secondary gate ("S2") by

plotting the events in our primary gate in bipartite FSC-V1 space (Fig. 1Div). We inspected the

images of the events in this space (Fig. 1Div, left - images of events within dotted rectangle;

right - images of events within solid rectangle) and refined the gate so that it included only

events of the target morphotype (Fig. 1Div, solid rectangle).

We then performed cell sorting using the gating strategy identified via IFC. As anticipated, the

sample exhibited the same qualitative, but not quantitative, pattern on the sorter as on the

imaging-cytometer (Fig. 1, compare. Ei vs. Ciii). We thus positioned our gates in the same

position, relative to the distribution of events, on the sorter as on the cytometer. Interestingly, we

found that UV1 – a parameter available on the cell sorter but not the cytometer – could better

discriminate spores from debris than V1 (Fig. S3). In this worked example, as well as other

analyses where spores were the target morphotype, we used UV1 rather than V1 to define our

secondary gate. With the gates set, we sorted the sample.

Finally, to validate our sorting strategy and quantify the purity of the sorted sample, we analyzed

a subset of the sorted sample via IFC. First, we confirmed that the sorted events occupied the

expected region of FSC-SSC space i.e., that the sorted sample occupied the same position in

FSC-SSC space as the S-cluster (compare Fig. 1, Ciii, 1Fi). Second, we confirmed the identity

of the sorted events using the image data (Fig. 1Fii).

IFC-enabled sorting yields high-purity populations of each morphotype

We used the above-described gate-discovery and validation strategy to conduct label-free cell

sorting on 43 occasions. These analyses involved 24 samples varying in initial composition and

targeted each of the four different *Pasteuria* morphotypes.

The sorted samples ranged in purity from 75% to 99% (Fig. 2A). Of the events that were not

identifiable as the target morphotype ("non-target events"), the majority were debris (e.g., salt

crystals, host material, Fig. 2B). Indeed, on average the sorted sample constituted only 1.8%

(range 0-8.4%) of *Pasteuria* morphotypes that were not the target (Fig. 2B).

Discussion

The first step toward understanding the physiological and ecological significance of

morphological variation in a bacterial population is to obtain a pure sample of each morphotype.

FACS is commonly employed for cell purification but its application to poorly studied or difficult-

to-culture bacteria is limited by a paucity of morphotype-specific fluorescent-labels (antibodies,

transgenics). Here, we demonstrate that IFC can be used to design and validate a label-free cell

sorting strategy that efficiently separates distinct bacterial morphotypes from each other and

from environmental debris. Critically, this procedure yields samples of a purity required for

common downstream, "-omics" analyses i.e., >=90%. High-speed image-enabled cell sorting

approaches have been innovated (30, 31) and are slowly becoming available, but their expense

and limited availability makes them out of reach for most. Our methodology represents an

immediately available, relatively economical alternative and could greatly expand our capacity to

understand morphological heterogeneity in populations of bacteria and other organisms.

Recent innovations in the acquisition and analysis of imaging flow cytometry data are critical to

the success of our approach. The first innovation is acoustic focusing. Historically, analyzing

highly heterogeneous samples using imaging cytometry has been impractical, because of the

slow speed of image acquisition. The acoustic focusing capability of the cytometer used herein

allowed us to obtain thousands of images per second. As such, even though *Pasteuria* cells

were by no means the most common events in our samples, we nonetheless obtained

thousands of pictures of them. This dataset, in turn, enabled us to use data-hungry clustering

algorithms to identify morphotype-specific parameters. In addition to increasing the speed with

which we could acquire data, acoustic focusing also reduces the noise in the imaging and

fluorescence parameters by orientating cells in a single, narrow stream (27). It is likely that this

further facilitated the discrimination of morphotype-specific light signatures. Finally, acoustic

focusing enabled us to rapidly validate our sorting protocol (as compared to e.g., a manual,

microscopy approach) by boosting the speed with which we could inspect the morphology of our

sorted events.

The second key innovation critical to our approach are machine-learning image-analysis

algorithms, which are increasingly used for the analysis of high-parameter flow cytometry data

(reviewed in (32, 33)). The multi-dimensionality of IFC data makes it difficult to analyze with

hierarchical, manual approaches alone. Here, we used two approaches to efficiently analyze

these data: the dimensionality-reduction algorithm (UMAP) allowed us to visualize the data in 2-

D space, while the clustering algorithm (FlowSOM) allowed us to rapidly locate different

morphotypes in this high-dimensional space and to extract their features. In future, a supervised

clustering approach (e.g., (14, 34, 35)) could be used for feature-detection and may improve the

efficiency of IFC-enabled gate identification.

While semi-automated computational analyses were critical to the design of our label-free cell

sorting strategy, its optimization nevertheless required user-input and experience. For example,

experimentation was required to elucidate the appropriate meta-clustering parameter for

FlowSOM: too few clusters and the target morphotype was binned with debris/other

morphotypes; too many and the target morphotype was separated into many different clusters,

making it difficult to identify morphotype-specific light signatures. Similarly, user-intervention and

observations were important for navigating the differences between the imaging cytometer and

the cell sorter. Unlike the imaging cytometer, the cell sorter used herein does not have acoustic

focusing. As a result, there was higher variability in the scattering and fluorescence data derived

using the sorter vs. the imaging cytometer (compare Fig. 1Cii, Ei). We could not, therefore,

position the gates in the same absolute position on the sorter as on the cytometer. Rather, we

needed to manually inspect the distribution of the events and then position the gate in the same

relative position on the sorter as on the imaging cytometer. Finally, prior knowledge vis-à-vis the

UV-autofluorescence of spores (13), enabled us to better discriminate the spores from debris.

These examples demonstrate that while label-free sorting is made possible by new innovations

in IFC hardware and analyses, it nevertheless requires knowledge of the bacterium's biology

and the capabilities of the hardware.

Here, we applied IFC in a novel way - to purify *known* morphotypes in a bacterial population.

However, a fortuitous byproduct of the IFC-enabled sorting procedure is that it yields a huge

amount of image data that can be used in the more traditional application of IFC: the discovery

of hitherto uncharacterized variation within a bacterial population (36). For example, in the

course of our analysis, we observed a range of grape morphologies, presumably representing

cells at various stages of the spore-development process, as well as events of unknown

provenance. This variability is difficult to observe (let alone quantify) using microscopy,

particularly when morphotypes are rare. These data may help us to address outstanding

questions vis-à-vis the life cycle of this bacterial pathogen e.g., "how does *Pasteuria*'s

morphology changes during the first week of infection?".

In principle, IFC-enabled cell sorting is readily generalizable to other study systems and

contexts. Nevertheless, we anticipate two hurdles that could narrow its widespread uptake.

First, it may not always be possible to distinguish morphotypes on the basis of intrinsic light-

scattering or autofluorescence signatures alone, particularly when hardware with limited optical

components (lasers/filters) is used. Staining the sample with 'general', membrane permeable

probes could increase the dimensionality of the data (and has proved useful in the development

of a transcriptomics-based, label-free sorting approach (11)). Second, software limitations could

limit the uptake of IFC-enabled cell sorting. To use IFC to identify gates requires that one can

simultaneously inspect the output of machine learning analyses (e.g., cluster, UMAP analyses)

and the image data. As far as we are aware, there is only one proprietary software platform

(FCS express) that permits this type of analysis using data derived with the Attune CytPix. We

encourage the development of open-source equivalents, particularly in languages commonly

used by microbial ecologists (e.g., R, Python), to overcome this potential barrier.

Cell sorting is a first step to understanding morphological heterogeneity in bacterial populations

but can only be applied to a fraction to species or contexts where that heterogeneity is

exhibited. Our work offers a novel, comparatively inexpensive method that will enable

microbiologists to purify – and hence phenotype – bacterial morphotypes from across the tree of

life and diverse environments.

Acknowledgements

The Attune CytPix used herein is located in the MSU Flow Cytometry Core and is supported by

the Equipment Grants Program, award #2022-70410-38419, from the U.S. Department of

Agriculture, National Institute of Food and Agriculture. We are grateful to members of the Wale

Lab for technical assistance and to Carrie Cizauskas PhD DVM for the illustrations in Table 1.

Bibliography

1. Yang DC, Blair KM, Salama NR. 2016. Staying in shape: the impact of cell shape on bacterial survival in diverse environments. Microbiol Mol Biol Rev 80:187–203.

2. Kysela DT, Randich AM, Caccamo PD, Brun YV. 2016. Diversity takes shape: understanding the mechanistic and adaptive basis of bacterial morphology. PLoS Biol 14:e1002565.

3. Young KD. 2006. The selective value of bacterial shape. Microbiology and Molecular Biology Reviews 70:660–703.

4. Horvath DJ, Li B, Casper T, Partida-Sanchez S, Hunstad DA, Hultgren SJ, Justice SS. 2011. Morphological plasticity promotes resistance to phagocyte killing of uropathogenic *Escherichia coli*. Microbes Infect 13:426–437.

5. Geerlings NMJ, Karman C, Trashin S, As KS, Kienhuis MVM, Hidalgo-Martinez S, Vasquez-Cardenas D, Boschker HTS, Wael KD, Middelburg JJ, Polerecky L, Meysman FJR. 2020. Division of labor and growth during electrical cooperation in multicellular cable bacteria. Proc Natl Acad Sci 117:5478–5485.

6. Zhang Z, Du C, Barsy F de, Liem M, Liakopoulos A, Wezel GP van, Choi YH, Claessen D, Rozen DE. 2020. Antibiotic production in Streptomyces is organized by a division of labor through terminal genomic differentiation. Sci Adv 6:eaay5781.

7. Prashar A, Bhatia S, Gigliozzi D, Martin T, Duncan C, Guyard C, Terebiznik MR. 2013. Filamentous morphology of bacteria delays the timing of phagosome morphogenesis in macrophages. J Cell Biol 203:1081–1097.

8. Meriläinen L, Herranen A, Schwarzbach A, Gilbert L. 2015. Morphological and biochemical features of Borrelia burgdorferi pleomorphic forms. Microbiology 161:516–527.

9. Dienes L, Weinberger HJ. 1951. The L Forms of Bacteria. Bacteriol Rev 15:245–288.

10. Katz SM, Hashemi S, Brown KR, Habib WA, Hammel JM. 1984. Pleomorphism of *Legionella pneumophila*. Ultrastruct Pathol 6:117–129.

11. Baron CS, Barve A, Muraro MJ, Linden R van der, Dharmadhikari G, Lyubimova A, Koning EJP de, Oudenaarden A van. 2019. Cell type purification by single-cell transcriptome-trained sorting. Cell 179:527-542.e19.

12. Shapiro HM. 2003. Practical Flow Cytometry, 4th ed. John Wiley & Sons, Inc, New Jersey, USA.

13. Laflamme C, Verreault D, Lavigne S, Trudel L, Ho J, Duchaine C. 2005. Autofluorescence as a viability marker for detection of bacterial spores. Front Biosci 10:1647.

14. Blasi T, Hennig H, Summers HD, Theis FJ, Cerveira J, Patterson JO, Davies D, Filby A,
Carpenter AE, Rees P. 2016. Label-free cell cycle analysis for high-throughput imaging flow
cytometry. Nat Commun 7:10256.

15. Kiepś J, Juzwa W, Dembczyński R. 2023. Imaging flow cytometry demonstrates
physiological and morphological diversity within treated probiotic bacteria groups. Int J Mol Sci
24:6841.

16. Pinto RN, Sebastian JA, Parsons MJ, Chang TC, Turner TR, Acker JP, Kolios MC. 2019.
Label-Free Analysis of Red Blood Cell Storage Lesions Using Imaging Flow Cytometry. Cytom
Part A 95:976–984.

17. Dey R, Rieger AM, Stephens C, Ashbolt NJ. 2019. Interactions of Pseudomonas aeruginosa
with Acanthamoeba polyphaga Observed by Imaging Flow Cytometry. Cytom Part A 95:555–
564.

18. Ebert D, Rainey P, Embley TM, Scholz D. 1996. Development, life cycle, ultrastructure and
phylogenetic position of *Pasteuria ramosa* Metchnikoff 1888: Rediscovery of an obligate
endoparasite of *Daphnia magna* Straus. Phil Trans R Soc B 351:1689–1701.

19. Ebert D, Duneau D, Hall MD, Luijckx P, Andras JP, Pasquier LD, Ben-Ami F. 2016. A
population biology perspective on the stepwise infection process of the bacterial pathogen
Pasteuria ramosa in *Daphnia*. Adv Parasit 91:265–310.

20. Duneau D, Luijckx P, Ben-Ami F, Laforsch C, Ebert D. 2011. Resolving the infection process
reveals striking differences in the contribution of environment, genetics and phylogeny to host-
parasite interactions. BMC biology 9:11.

21. Clerc M, Ebert D, Hall MD. 2015. Expression of parasite genetic variation changes over the
course of infection: implications of within-host dynamics for the evolution of virulence. Proc
Royal Soc B Biological Sci 282:20142820.

22. Fields PD, Reisser C, Dukić M, Haag CR, Ebert D. 2015. Genes mirror geography in
Daphnia magna. Mol Ecol 24:4521–4536.

23. Luijckx P, Ben-Ami F, Mouton L, Pasquier LD, Ebert D. 2010. Cloning of the unculturable
parasite Pasteuria ramosa and its Daphnia host reveals extreme genotype-genotype
interactions. Ecol Lett 14:125–31.

24. Klüttgen B, Dülmer U, Engels M, Ratte HT. 1994. ADaM, an artificial freshwater for the
culture of zooplankton. Water Res 28:743–746.

25. Ben-Ami F, Ebert D, Regoes RR. 2010. Pathogen Dose Infectivity Curves as a Method to
Analyze the Distribution   of Host Susceptibility: A Quantitative Assessment of Maternal
Effects   after Food Stress and Pathogen Exposure. Am Nat 175:106–115.

26. Ebert D, Zschokke-Rohringer CD, Carius HJ. 2000. Dose effects and density-dependent
regulation of two microparasites of Daphnia magna. Oecologia.

27. Ward MD, Kaduchak G. 2018. Fundamentals of Acoustic Cytometry. Curr Protoc Cytom
84:e36.

28. McInnes L, Healy J, Saul N, Großberger L. 2018. UMAP: Uniform Manifold Approximation
and Projection. J Open Source Softw 3:861.

29. Gassen SV, Callebaut B, Helden MJV, Lambrecht BN, Demeester P, Dhaene T, Saeys Y.
2015. FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry
data. Cytom Part A 87:636–645.

30. Nitta N, Sugimura T, Isozaki A, Mikami H, Hiraki K, Sakuma S, Iino T, Arai F, Endo T,
Fujiwaki Y, Fukuzawa H, Hase M, Hayakawa T, Hiramatsu K, Hoshino Y, Inaba M, Ito T,
Karakawa H, Kasai Y, Koizumi K, Lee S, Lei C, Li M, Maeno T, Matsusaka S, Murakami D,
Nakagawa A, Oguchi Y, Oikawa M, Ota T, Shiba K, Shintaku H, Shirasaki Y, Suga K, Suzuki Y,
Suzuki N, Tanaka Y, Tezuka H, Toyokawa C, Yalikun Y, Yamada M, Yamagishi M, Yamano T,
Yasumoto A, Yatomi Y, Yazawa M, Carlo DD, Hosokawa Y, Uemura S, Ozeki Y, Goda K. 2018.
Intelligent Image-Activated Cell Sorting. Cell 175:266-276.e13.

31. Schraivogel D, Kuhn TM, Rauscher B, Rodríguez-Martínez M, Paulsen M, Owsley K,
Middlebrook A, Tischer C, Ramasz B, Ordoñez-Rueda D, Dees M, Cuylen-Haering S, Diebold
E, Steinmetz LM. 2022. High-speed fluorescence image–enabled cell sorting. Science 375:315–
320.

32. Stolarek I, Samelak-Czajka A, Figlerowicz M, Jackowiak P. 2022. Dimensionality reduction
by UMAP for visualizing and aiding in classification of imaging flow cytometry data. iScience
25:105142.

33. Marsh-Wakefield FM, Mitchell AJ, Norton SE, Ashhurst TM, Leman JK, Roberts JM, Harte
JE, McGuire HM, Kemp RA. 2021. Making the most of high-dimensional cytometry data.
Immunol Cell Biol 99:680–696.

34. Hennig H, Rees P, Blasi T, Kamentsky L, Hung J, Dao D, Carpenter AE, Filby A. 2017. An
open-source solution for advanced imaging flow cytometry data analysis using machine
learning. Methods 112:201–210.

35. Haridas V, Ranjbar S, Vorobjev IA, Goldfeld AE, Barteneva NS. 2017. Imaging flow
cytometry analysis of intracellular pathogens. Methods 112:91–104.

36. Power AL, Barber DG, Groenhof SRM, Wagley S, Liu P, Parker DA, Love J. 2021. The
application of imaging flow cytometry for characterisation and quantification of bacterial
phenotypes. Front Cell Infect Microbiol 11:716592.

37. 2024. Attune Cytometric Software User Guide. MAN0026553. Thermo Fisher Scientific Ltd.,
Singapore.

A. Cluster analysis: imaging + light params.

B. Assign morphotypes to clusters using images

C. ID params. to discriminate among morphotypes (1° gate)

D. ID params. to discriminate morphotypes from debris (2° gate)

E. Use gates + FACs to sort target morphotype

F. Validate ID of sorted events via IFC

Figure 1: Design & validation of a label-free cell sorting strategy using imaging flow cytometry (IFC). Worked example of IFC-enabled cell-sorting, for the purification of *Pasteuria* spores (the "target morphotype") from a sample of host material, microbiota and (non-target) *Pasteuria* morphotypes (cauliflower, C; grapes, G). **A.** Cluster analysis of imaging and light parameters (scattering, fluorescence) is performed and imposed onto a UMAP. **B.** Each cluster is assigned to a morphotype or debris using the image data. Here, clusters 1, 2, & 3 were denoted the G (orange), C (green) and S (lilac) clusters, respectively. **C.** A primary (1°) gate (S1) that distinguishes the target morphotype from non-target morphotypes is identified using (i) a "parameter heatmap", which displays the median value of each parameter in each cluster (i; only those parameters shared among the sorter and cytometer displayed), and (ii, iii) visualization of the parameters in bipartite space (ii, events in S cluster, lilac, G cluster, orange, C cluster, green; iii, shaded from black to white to indicate increasing event density). **D.** We first inspect the images associated with the events in the 1° gate. To identify a secondary (2°) gate for separating target morphotypes from debris, regions of the target cluster that contain a pure population of the target morphotype (a) vs. a mixed population (b) are identified using the image data. Fluorescence parameters that distinguish the regions are identified by coloring the cluster by each parameter (iii, warmer colors indicate higher fluorescence intensity; note that regions A, B are differentially colored by FSC, V1 indicating the suitability of these parameters for 2° gating). Finally, events in the primary gate are plotted in bipartite space defined by the secondary parameters. The location of the secondary gate (S2) is set with respect to the image data (iv; right images, events within the gate (solid box), left images, events outside of the gate (dashed box)). **E.** The sample is analyzed using the cell sorter. It is verified that the events take approximately the same distribution on the sorter and imaging cytometer (Ei, Ciii respectively; note differences in scale); the gates are then set to the same, relative position with respect to the distribution. Here, analysis of the sample using the sorter revealed that the sorter's UV1 channel had more discriminatory power than V1 (Fig. S3) and so we used this parameter for sorting in place of V1(Eii) **F.** To verify that sorted events are of the target morphotype and to quantify the purity of the sorted sample, the sorted sample is analyzed via IFC. First, it is confirmed that sorted events lie in expected region of bipartite space (compare Fi, Cii), then the image data is inspected and the proportion of target morphotypes quantified (Fii, representative images from sorted sample)

Figure 2: IFC-enabled cell sorting yields high purity samples of four distinct *Pasteuria* morphotypes. **A.** Purity of sorted samples, determined using image data acquired via IFC (mean purity across sorts, black, large points; purity of each replicate sort, colored, small points). Note that a replicate refers to an attempt to sort a specific morphotype i.e., replicate 1 under the "activated" and "cauliflower" columns may not have involved the same sample or been performed at the same time. **B.** Composition of the non-target fraction of the sorted sample in each replicate (colored as in **A**).