Imaging flow cytometry enables label-free cell sorting of morphological variants from

unculturable bacterial populations.

Keywords: autofluorescence, flow cytometry, cell sorting, non-model, *Pasteuria ramosa*

Authors

Daniel Vocelle[1], Lindsey Thompson[2], Matthew Bernard[1] & Nina Wale[1,2,3,4,+]

1. Department of Pharmacology & Toxicology, Michigan State University, East Lansing, MI 48824, USA

2. Department of Microbiology, Genetics & Immunology, Michigan State University, East Lansing, Michigan,

USA 48824

3. Department of Integrative Biology, Michigan State University, East Lansing, Michigan, USA 48824

4. Program in Ecology, Evolution & Behavior, Michigan State University, East Lansing, Michigan, USA 48824

+corresponding author

Contributions

DV and NW conceptualized the research.

DV and LT performed the experiments.

DV designed the analysis, with consultation from MB.

DV and NW analyzed the data.

NW wrote the manuscript.

DV, LT, MB edited the manuscript.

DV, MB acquired funding for the instrumentation.

Data availability

Description: data will be made available upon article acceptance.

Abstract
Bacterial populations often display remarkable morphological heterogeneity. Fluorescence

activated cell sorting (FACS) is an important tool for understanding the biological significance of

this morphological diversity, as it permits the separation and study of distinct morphological

variants (morphotypes) from each other and their environmental milieu. In FACS, cells are first

labeled with fluorescent markers such as antibodies or transgenic constructs, and then sorted

based on their possession of such labels. However, since the development of fluorescent labels

requires *a priori* knowledge of bacterial biology, it is often impossible to apply FACS to

understudied and/or unculturable bacteria. This challenge has limited our capacity to investigate

the role of bacterial size and shape in all but a small, largely culturable subset of bacterial taxa.

Here, we present an innovative strategy that permits label-free cell sorting of bacterial

morphotypes, using an unculturable, pleiomorphic pathogen (*Pasteuria ramosa*) as a model

bacterium. We show that imaging flow cytometry (IFC) can be used to systematically identify

light-scattering and autofluorescence "signatures" of bacterial morphotypes, on which basis cell

sorting can be conducted. Critically, our IFC-enabled cell sorting strategy yields samples of

sufficient purity (>90%) for common downstream analyses e.g., "-omics" analyses. Our work

represents an innovative application of IFC and provides an economical, widely applicable

solution to a central problem in the study of bacterial diversity.

46
47
48
49

Introduction

Bacterial populations often contain cells varying in size and shape, and this morphological heterogeneity can have a profound impact on bacterial fitness (1–3). For example, different forms of bacteria (morphotypes) may perform distinct ecological "roles" or exhibit differential susceptibility to environmental stress, permitting population persistence in the face of environmental stress (4–7). To fully understand the causes and consequences of morphological variation within bacterial populations, we need to extend our investigations beyond well-studied taxa and *in vitro* environments, which can suppress morphological diversity within a bacterial population (2, 8–10). Here, we present imaging flow cytometry (IFC)-enabled cell sorting, a novel procedure that allows for the purification of distinct morphotypes from unculturable, pleiomorphic bacteria living in complex milieu.

Efforts to understand the significance of morphological heterogeneity in a bacterial population often begin with isolating distinct bacterial morphotypes from each other and/or their environmental milieu, via fluorescence activated cell sorting (FACS). However, the application of FACS to unculturable bacteria or those found in complex samples (e.g., soil, host tissues) is a challenge. A chicken-and-egg-scenario exists. In FACS, different cell populations are tagged with distinct fluorescent labels (e.g., fluorescently labelled antibodies) on whose basis the populations are separated. However, the development of morphotype-specific labels requires the characterization of morphotype-specific antigens, which itself requires that morphotypes be separated and studied. In the absence of morphotype specific labels, "general' membrane permeable labels (e.g., CFSE, Mitotracker), which bind to molecules present in most cells (e.g., amines, RNA or mitochondria), could be used to discriminate morphotypes in certain contexts (11). However, because general labels indiscriminately permeate all cells within a sample, they

are unlikely to be helpful in distinguishing a specific morphotype if it is present in a complex,

cell-rich milieu such has host-tissue or soil.

A potential solution to the problem of sorting bacterial morphotypes in the absence of labels is to

exploit intrinsic variation in their light-scattering and -fluorescence properties (i.e., "light

signatures"). Cells differing in size and granularity differentially scatter light, and the degree of

autofluorescence may also vary among morphotypes (12). For example, viable *Bacillus*

anthracis spores exhibit a distinctive autofluorescent signature upon excitation with UV light

(13). Recent advances in imaging flow cytometry (IFC) simplify the identification of intrinsic light

signatures of morphologically distinct cells, by generating images of each cell and measures of

their light-scattering and autofluorescence properties. Systematic analysis of these data allows

for the identification of morphotype-specific light signatures and has been used to identify

morphologically distinct subpopulations of mammalian and bacterial cells (14–17).

Here, using the pleiomorphic bacterial pathogen *Pasteuria ramosa* (hereafter, *Pasteuria*) as a

model, we demonstrate that IFC can be used to efficiently design and validate strategies for

label-free sorting of distinct bacterial morphotypes. *Pasteuria* is a spore-forming pathogen of

zooplankton that exhibits striking morphological variation during its life cycle (Table 1, (18)).

Longstanding hypothesis vis-à-vis the "division of labor" among different morphotypes within the

Pasteuria bacterial population (19) remain untested, largely because it is difficult to separate

bacteria from host material for functional -omics and *in vitro* study. By combining state-of-the-art

IFC instrumentation, which allows us to rapidly derive thousands of images of bacterial cells,

with machine learning algorithms we systematically designed and validated autofluorescence-

based FACS strategies that yield high-purity samples of each of four distinct morphotypes. Our

work thus provides proof of principle that IFC-enabled sorting can be used to overcome the first

barrier to understanding morphological heterogeneity in bacterial species that cannot be readily

labeled.

	name (abbreviation)	Function (*hypothesized)	largest dimension (range, µm)
	Activated (A)	host attachment	-
	Cauliflower (C)	in-host growth, host manipulation*	4-6.7
	Grape-seed (`grape', G)	sporulation	4-6
	Mature spore (`spore', S)	transmission, environmental persistence	4.2-5.4

129   **Table 1. Variation in the form and function of *Pasteuria ramosa* morphotypes. Form and**
130   function information extracted from (18, 20, 21). Schematic representations of morphotypes not
131   to scale.

Materials and methods

Our goal was to design and validate cell sorting strategies suitable for the purification 4 distinct

Pasteuria morphotypes from the homogenate of an infected host i.e., a sample containing

several *Pasteuria* morphotypes, host-cells, host food (algae) present in the gut, and host

microbiota. Below we describe our approach in general terms. The final procedure and results

of our validation steps are described in the results.

Host & Parasites

Throughout, we used *Daphnia magna* (HuHO-2 genotype) hosts and *Pasteuria ramosa* (strain

C1), originally collected in Hungary and Russia, respectively (22, 23). Both were generously

donated by D. Ebert (U. Basel, Switzerland).

To generate samples of *Pasteuria ramosa*, we experimentally infected *Daphnia magna*. The

infection procedure took the same, general format, with minor variation in husbandry details.

Briefly, *P. ramosa* spores were shaken at 1500 rpm overnight. On day 0 of the experiment, we

filled a vessel (e.g., 24 well plate or beaker) with media (ADaM or COMBO), added 1-10 *D.*

magna <24 hours old (24)**.** A second equivalently sized dose of spores was added on the

following day. Animals were fed an increasing quantity of either *Scenedesmus obliquus* or

Ankistrodesmus falcatus as they aged, up to ~30-60 days. To diagnose *Pasteuria* infection,

animals were assessed for visual signs of infection, including discoloration, lack of offspring and

enlargement (25). Infection was then confirmed by inspection using microscopy.

Instruments

We used a Thermofisher Attune CytPix flow cytometer (the cytometer, hereafter) for IFC. This

cytometer is equipped with a high-speed, brightfield camera (20x objective, 0.45 numerical

aperture, 0.3 micron/pixel resolution), four lasers and 14 filters (Table S1). Acoustic focusing

allows for the alignment of cells and particles (hereafter, events) in a narrow depth of field. This

property (along with the high-speed camera) allows for the acquisition of in-focus images at a

rate an order of magnitude faster than cytometers with hydrodynamic-focusing (26).

For cell sorting, we used a BD Influx equipped with 5 lasers and 12 filters (Table S1). The

default measure of fluorescence on the BD Influx is "pulse height" (rather than area). Since our

goal was to use the imaging cytometer to identify fluorescence parameters that could be

employed in cell sorting, we used pulse height as our measure of fluorescence on both

machines.

Data acquisition

Immediately following diagnosis, we prepared infected animals for flow cytometric analysis.

First, each animal was washed in PBS (no Ca+ or Mg+). We transferred 1-4 animals, depending

on the degree of infection and an initial assessment of morphotype composition via microscopy,

into a 1.5mL Eppendorf tube containing PBS. We homogenized the animals briefly with an

electric pestle (~20 seconds, to avoid damaging cells) to create a slurry. We diluted the sample

for analysis in 500mL Focusing Buffer (Thermofisher Scientific, catalog no. A24904) or PBS,

which have equivalent refractive indices. We ran samples at 12.5mL/min until we acquired

30,000 events. Between runs, we twice flushed the cytometer line to prevent sample carry over.

Pre-gating

We first excluded ("gated out") algae on the basis of their chlorophyll autofluorescence (high

emission in channel R3, cf. Table S1). We then selected ("gated in") the events that (i) were

successfully photographed and processed ("Is Processed" positive events) and, furthermore,

contained only one particle (Event Particle Count = 1). This yielded a dataset of single particles,

each of which had had both imaging and light-parameter data associated with it.

To analyze the image data, we employed inbuilt Attune Image-Processing Software (version

6.0.1). We used the software's bead masking model (Beads_Only_Full_Resolution_v1) which is

optimized for objects 2-5μm. We chose this algorithm because (i) an alternative model

optimized for objects >5μm, excluded ~70% of our target objects and (ii) because we found that

the bead model could process a variety of *P. ramosa* morphotypes, including those that

exceeded 5μm (e.g., cauliflowers). Image-processing yielded a matrix containing the values of

22 imaging parameters including a variety of size and "texture" metrics (Table S2). In total,

including the fluorescent parameters (and pulse parameters height, area, and width) each event

was associated with 74 parameters. Before conducting further analysis, we rescaled each

parameter linearly between 0 and 1.

We visualized the data using Uniform Manifold Approximation Projection for Dimension

Reduction (UMAP) (27), implemented in FCS Express (De Novo Software). Briefly, this

algorithm allows for the visualization of high-dimensional data in a 2D plane. By inspecting

distinct regions of the UMAP, and the imaging data with which it was associated, we found that

different morphotypes clustered in distinct areas of the UMAP. Preliminary analysis of this kind

revealed that certain image parameters were particularly informative for distinguishing different

morphotypes (Table S2). We employed this subset of parameters in further analyses, described

below.

Pre-sort analysis: identifying morphotype-specific parameters for cell sorting

For each sample, we used IFC to identify gates that would enable us to sort a specific

morphotype ("target morphotype", hereafter) from the remainder of the sample.

First, we visualized the morphological complexity of the sample using a UMAP analysis of all

scattering and fluorescent parameters, as well as several imaging parameters identified in

preliminary analyses as being most-informative (Table S2, S3). We then performed cluster

analysis to identify (i) a primary gate, on which basis we could distinguish our target morphotype

from other bacterial cells, and (ii) secondary gate(s), suitable for distinguishing our target

morphotype from debris within the primary gate (see Results). Specifically, we used FlowSOM -

a neural network-like algorithm that identifies clusters of events characterized by distinct

(combinations of) parameters (28). We set the consensus clustering parameter to between 8-

20, depending on the sample, and otherwise used default settings (Table S3).

The above-described procedure was repeated on a sample-by-sample basis. Although, over the

course of this project, we characterized distinct "signatures" of each morphotype that held

among samples, the fluorescence parameters used to define the gates varied among samples,

particularly in the initial stages of our work. Due to differences among samples in the

autofluorescence of the cells, moreover the exact position of the gates varied from sample to

sample. Importantly, however, the gate -identification and -validation strategy - which is our

focus here - remained the same for all samples.

Cell sorting

Prior to analysis via cell sorting, we transferred the sample to a 5 mL FACS tube (Corning,

catalog no. 352063) and filtered it through a 100μm mesh filter. We then used the gating

strategy designed via IFC ("pre-sort" analysis), or a minor variation of it (see Results, Fig. S1) to

sort each of the morphotypes in the sample that were of sufficient abundance.

Post-sort analysis: quantification of sample purity

We verified the purity of each sorted sample using IFC. We analyzed up to 30000 events (mean

events analyzed = 6993, range 526-30000); the exact number varied according to the relative

abundance of the morphotype present in the pre-sorted sample. We analyzed the IFC data as

before (i.e., same image processing model, and initial gating on single-processed events) and

then "down-sampled" it to 500 events (images). Using the image data, we manually classified

each of the 500 events as an identifiable bacterial morphotype (activated, cauliflower, grape or

spore) or unidentified debris. Finally, we calculated the purity of the sorted sample i.e., the

proportion of total events in the down-sampled dataset that were of the target morphotype. This

purity measure is conservative given that we include unidentified debris such as salts in our total

event count.

245   Results

IFC enables the identification of cell sorting strategies that exploit morphotype-specific

light signatures

We established a robust procedure for identifying light signatures that can be harnessed for the

cell sorting of morphologically distinct bacterial morphotypes, in the absence of labels. To

demonstrate this approach, we provide a worked example (Fig. 1). In this example, our goal

was to sort a spore (S) population from a homogenate comprising host (*Daphnia*) material, host-

microbiota and non-target morphotypes of *Pasteuria* (cauliflower (C), grape (G)). Worked

examples of the procedure as applied to activated, cauliflower and grape stages are provided in

the supplement (Fig. S1).

First, we performed a dimensionality reduction (UMAP) analysis to visualize the data in 2D

space (Fig. 1A). To identify where in this 2D space each *Pasteuria* morphotype lay, we imposed

upon this UMAP the results of a cluster analysis (Fig. 1A, colors/numbers indicate clusters).

This cluster analysis was performed using all light-scattering and -fluorescence parameters and

a subset of the imaging parameters (Table S2). We manually classified each cluster as either a

morphotype-rich or debris-rich cluster using the image data (Fig. 1B). For example, since spores

appeared in most of the images associated with cluster 3, we designated cluster 3 the spore-

rich cluster (S-cluster; Fig. 1B, purple outline); similarly, clusters 1 and 2 were designated

grape- and cauliflower- rich clusters, respectively.

Figure 1: The design & validation of a label-free cell sorting strategy using imaging flow cytometry (IFC). Worked example wherein we used IFC-enabled cell-sorting to purify *Pasteuria* spores (the "target morphotype") from a sample of host material, microbiota and (non-target) *Pasteuria* morphotypes (cauliflower, C; grapes, G). **A.** Cluster analysis of imaging and light parameters (scattering, fluorescence) is performed and imposed onto a UMAP. **B.** Each cluster is assigned to a morphotype or debris using the image data. Here, clusters 1, 2, & 3 were denoted the G (orange), C (green) and S (lilac) clusters, respectively. **C.** A primary (1°) gate that distinguishes the target morphotype from non-target morphotypes is identified using (i) a "parameter heatmap", which displays the median value of each parameter in each cluster (i; only those parameters shared among the sorter and cytometer displayed), and (ii, iii) visualization of the parameters in bipartite space (ii, events in S cluster, lilac, G cluster, orange, C cluster, green; iii, shaded from black to white to indicate increasing event density). The primary gate (S1) is thus set. **D.** We first inspect the images associated with the events in the primary gate. To identify a secondary (2°) gate for separating target morphotypes from debris, regions of the target cluster that contain a pure population of the target morphotype (a) vs. a mixed population (b) are identified using the image data. Fluorescence parameters that distinguish the regions are identified by coloring the cluster by each parameter (iii, warmer colors indicate higher fluorescence intensity; note that regions *a*, *b* are differentially colored by FSC, V1 indicating the suitability of these parameters for 2° gating). Finally, events in the primary gate are plotted in bipartite space defined by the secondary parameters. The location of the secondary gate (S2) is set with respect to the image data (iv; right images, events within the gate (solid box), left images, events outside of the gate (dashed box). **E.** The sample is analyzed using the cell sorter. It is verified that the events take approximately the same distribution on the sorter and imaging cytometer (Ei, Ciii respectively; note differences in scale); the gates are then set to the same, relative position with respect to the distribution. Here, analysis of the sample using the sorter revealed that the sorter's UV1 channel had more discriminatory power than V1 (Fig. S3) and so we used this parameter for sorting in place of V1**(**Eii**)** . F. To verify that sorted events are of the target morphotype and to quantify the purity of the sorted sample, the sorted sample is analyzed via IFC. First, it is confirmed that sorted events lie in expected region of bipartite space (compare Fi, Cii), then the image data is inspected and the proportion of target morphotypes quantified (Fii, representative images from sorted sample)

Next, we used the cluster analysis to identify light-scattering and fluorescence parameters that

characterized our target morphotype vs. the other morphotypes (C, G). Analysis of the median

parameter values associated with each of the morphotype clusters (Fig. 1Ci), suggested that the

different morphotypes could be broadly distinguished by the light scattering parameters FSC

and SSC (Fig.1Ci, F,S), consistent with their differences in size and granularity (Table 1). The

distribution of G-, C- and S- cluster events in FSC-SSC space confirmed this inference (Fig.

1Cii, Ciii). We thus used FSC-SSC to define our primary gate (S1, Fig. 1Cii).

Inspection of the images associated with the events in this primary gate revealed that it included

debris as well as our target morphotype (Fig. 1Di). We thus sought to identify a secondary gate

that could be used to distinguish the target morphotype from debris. First, we inspected the

images associated with the events in our target cluster. This analysis revealed that there was a

gradient within the cluster: on the left lay a largely homogenous population of spores (Fig. 1Dii,

region a); whereas on the right we found a mixed population of spores and other non-target

objects (Fig. 1Dii, region b). To identify which of the fluorescence parameters shared by the

sorter and the cytometer (Table S1) could differentiate events in regions a and b, we recolored

the events in each cluster by their fluorescence intensity in each of the shared channels (Fig.

Diii, 4 parameters shown for readability; all parameters are displayed in Fig. S2). This analysis

revealed that V1 could distinguish the spore-rich region (a) from the mixed region (b) and could

therefore be used in combination with FSC to define a secondary gate. Finally, we established

the position of this secondary gate ("S2") by plotting the events in our primary gate in bipartite

FSC-V1 space (Fig. 1Div). We inspected the image data (Fig. 1Div, left images, events within

dotted rectangle; right images, images from solid rectangle) and refined the gate so that it

included only events of the target morphotype (Fig. 1Div, solid rectangle).

With a gating strategy in hand, we then performed cell sorting. As anticipated, the sample

exhibited the same qualitative, but not quantitative, pattern on the sorter as on the imaging-

cytometer (compare Fig. 1Cii, Ciii vs. Ei). We thus positioned our gates in the same position,

relative to the distribution of events, on the sorter as on the cytometer. Interestingly, we found

that UV1 – a parameter available on the cell sorter but not the cytometer - could better

discriminate spores from debris than V1 (Fig. S3). In this worked example, as well as other

analyses wherein spores were the target morphotype, we thus used UV1 rather than V1 to

define our secondary gate. With the gates set, we sorted the sample.

Finally, to validate our sorting strategy and quantify the purity of the sorted sample, we analyzed

a subset of the sorted sample via IFC. First, we confirmed that the sorted events occupied the

expected region of FSC-SSC space i.e., that the sorted sample occupied the same position in

FSC-SSC space as the S-cluster (compare Figs. Ciii, 1Fi). Second, we confirmed the identity of

the sorted events using the image data (Fig. 1Fii, Fig. 2).

IFC-enabled sorting yields high-purity populations of each morphotype

We used the above-described gate-discovery and validation strategy to conduct label-free cell

sorting on 43 occasions. These analyses involved 24 samples varying in initial composition and

targeted each of the four different *Pasteuria* morphotypes.

The sorted samples ranged in purity from 75% to 99% (Fig. 2A). Of the events that were not

identifiable as the target morphotype ("non-target events"), the majority were debris (e.g., salt

crystals, host material, Fig. 2B). Indeed, on average the sorted sample constituted only 1.8%

(range 0-8.4%) of *Pasteuria* morphotypes that were not the target (Fig. 2B).

Figure 2: IFC-enabled cell sorting yields high purity samples of four distinct *Pasteuria* morphotypes. **A.** Purity of sorted samples, determined using image data acquired via IFC (mean purity across sorts, black, large points; purity of each replicate sort, colored, small points). Note that a replicate refers to an attempt to sort a specific morphotype i.e., replicate 1 under the "activated" and "cauliflower" columns may not have involved the same sample or been performed at the same time. **B.** Composition of the non-target fraction of the sorted sample in each replicate (colored as in **A**).

Discussion

The first step toward understanding the physiological and ecological significance of

morphological variation in a cellular population is to purify distinct morphotypes. FACS is

commonly employed for cell purification but its application to poorly studied or difficult-to-culture

bacteria is limited by a paucity of fluorescent-labels (antibodies, transgenics). Here, we

demonstrate that IFC can be used to design and validate a label-free cell sorting strategy that

efficiently separates distinct bacterial morphotypes from each other and from environmental

debris. Critically, this procedure yields samples of a purity required for common downstream, "-

omics" analyses i.e., >=90%. High-speed image-enabled cell sorting approaches have been

innovated (29, 30) and are slowly becoming available, but their expense and limited availability

makes them out of reach for most. Our methodology represents an immediately available,

relatively economical alternative and could greatly expand our capacity to understand

heterogeneity in cellular populations of bacteria, and indeed, other organisms or tissues.

Recent innovations in the acquisition and analysis of imaging flow cytometry data are critical to

the success of our approach. The first innovation is acoustic focusing. Historically, analyzing

highly heterogeneous samples using imaging cytometry has been impractical, because of the

slow speed of image acquisition. The acoustic focusing capability of the cytometer used herein

allowed us to obtain thousands of images per second. As such, even though *Pasteuria* cells

were by no means the most common particles in our samples, we nonetheless obtained

thousands of pictures of these cells. This dataset, in turn, enabled us to use data-hungry

clustering algorithms to identify morphotype-specific parameters. In addition to increasing the

speed with which we could acquire data, acoustic focusing also reduces the noise in the

imaging and fluorescence parameters by orientating cells in a single, narrow stream (26). It is

likely that this further facilitated the discrimination of morphotype-specific light signatures.

Finally, acoustic focusing enabled us to rapidly validate our sorting protocol (as compared to

e.g., a manual, microscopic approach) by boosting the speed with which we could inspect the

morphology of our sorted events.

The second key innovation critical to our approach are machine-learning analysis algorithms,

which are increasingly used for the analysis of high-parameter flow cytometry data (reviewed in

(31, 32)). The multi-dimensionality of IFC data makes it difficult to analyze with hierarchical,

manual approaches alone. Here, we used two approaches to efficiently analyze these data: the

dimensionality-reduction algorithm (UMAP) allowed us to visualize the data in 2-D space; while

the clustering algorithm (FlowSOM) allowed us to rapidly locate different morphotypes in this

high-dimensional space and to extract their features. In future, a supervised clustering approach

(e.g., (14, 33, 34)) could be used for feature-detection and may improve the efficiency of our

approach.

While semi-automated computational analyses were critical to the design of our label-free cell

sorting strategy, its optimization nevertheless required user-input and experience. For example,

experimentation was required to elucidate the appropriate meta-clustering parameter for

FlowSOM: too few clusters and the morphotypes were binned with debris/other morphotypes;

too many and our target morphotypes were separated into many different clusters, making it

difficult to identify morphotype-specific light signatures. Similarly, user-intervention and

observations were important for navigating the differences among the imaging-cytometer and

the cell sorter. Our cell sorter does not have acoustic focusing like the imaging-cytometer. As a

result, there is higher variability in the scattering and fluorescence parameters on the sorter than

on the cytometer (compare Fig. 1Cii, Ei), and we could not position the sorting gates in the

same absolute position on the sorter as on the cytometer. Instead, we positioned the gates in

the same *relative* position on each machine and, at least in our initial runs, confirmed that the

gates were correctly positioned by pausing sorting after a few thousand events and verifying the

composition of the sorted sample using IFC. Finally, alert to previous findings vis-à-vis the UV-

autofluorescence of spores (13), we discovered that the UV1 channel – which was available on

the sorter but not the imaging cytometer – allowed us to better discriminate the spores from

debris. Thus, while IFC-enabled sorting is arguably made possible by new innovations in IFC

hardware and analyses, it nevertheless requires user knowledge of the focal bacterium's biology

and the capabilities of the hardware.

Here, we applied IFC in a novel way - to purify *known* morphotypes in a bacterial population.

However, a fortuitous byproduct of the IFC-enabled sorting procedure is that it yields a huge

amount of image data that can be used in the more traditional application of IFC: the discovery

of hitherto uncharacterized variation within a bacterial population (35). For example, in the

course of our analysis, we observed a full range of "grape" morphologies, from those without an

"eye" to almost-spore-like stages; we also observed events of unknown provenance. Such

variability is difficult to observe (let alone quantify) using microscopy, particularly when

morphotypes are rare and may enable us to address outstanding questions in vis-à-vis the life

cycle of this pathogen e.g., what morphological stage mediates the first ~week of infection.

In principle, IFC-enabled cell sorting is readily generalizable to other study systems and

contexts. Nevertheless, we anticipate two hurdles that could narrow its widespread uptake.

First, it may not always be possible to distinguish morphotypes on the basis of intrinsic light-

scattering or autofluorescence signatures alone, particularly when hardware with limited optical

components (lasers/filters) is being used. Staining the sample with membrane permeable

probes could increase the dimensionality of the data (and proved useful in the development of a

transcriptomics based, label-free sorting approach (11)). Second, software suitable for these

analyses could limit the uptake of IFC-enabled cell sorting. This analysis requires the capacity to

simultaneously inspect the output of cluster and UMAP analyses and image data. As far as we

are aware, there is only one proprietary software platform (FCS express) that permits this type

of analysis using the imaging cytometer used herein (Attune Cytpix). We encourage the

development of open-source equivalents, particularly in languages commonly used by microbial

ecologist (e.g., R, Python).

Cell sorting is a first step to understanding morphological heterogeneity in bacterial populations

but can only be applied to a fraction to species or contexts where that heterogeneity is

exhibited. Our work offers a novel, comparatively inexpensive method that will enable

microbiologists to purify – and hence phenotype – bacterial morphotypes from across the tree of

life and from diverse environments.

407   **Acknowledgements**

408   The Attune CytPix used is located in the MSU Flow Cytometry Core and is supported by the

409   Equipment Grants Program, award #2022-70410-38419, from the U.S. Department of

410   Agriculture, National Institute of Food and Agriculture. We are grateful to members of the Wale

411   Lab for technical assistance and to Carrie Cizauskas PhD DVM for the illustrations in Table 1.

Bibliography

1. Yang DC, Blair KM, Salama NR. 2016. Staying in Shape: the Impact of Cell Shape on Bacterial Survival in Diverse Environments. Microbiol Mol Biol Rev 80:187–203.

2. Kysela DT, Randich AM, Caccamo PD, Brun YV. 2016. Diversity takes shape: understanding the mechanistic and adaptive basis of bacterial morphology. PLoS Biol 14:e1002565.

3. Young KD. 2006. The selective value of bacterial shape. Microbiology and Molecular Biology Reviews 70:660–703.

4. Horvath DJ, Li B, Casper T, Partida-Sanchez S, Hunstad DA, Hultgren SJ, Justice SS. 2011. Morphological plasticity promotes resistance to phagocyte killing of uropathogenic Escherichia coli. Microbes Infect 13:426–437.

5. Geerlings NMJ, Karman C, Trashin S, As KS, Kienhuis MVM, Hidalgo-Martinez S, Vasquez-Cardenas D, Boschker HTS, Wael KD, Middelburg JJ, Polerecky L, Meysman FJR. 2020. Division of labor and growth during electrical cooperation in multicellular cable bacteria. Proc Natl Acad Sci 117:5478–5485.

6. Zhang Z, Du C, Barsy F de, Liem M, Liakopoulos A, Wezel GP van, Choi YH, Claessen D, Rozen DE. 2020. Antibiotic production in Streptomyces is organized by a division of labor through terminal genomic differentiation. Sci Adv 6:eaay5781.

7. Prashar A, Bhatia S, Gigliozzi D, Martin T, Duncan C, Guyard C, Terebiznik MR. 2013. Filamentous morphology of bacteria delays the timing of phagosome morphogenesis in macrophages. J Cell Biol 203:1081–1097.

8. Meriläinen L, Herranen A, Schwarzbach A, Gilbert L. 2015. Morphological and biochemical features of Borrelia burgdorferi pleomorphic forms. Microbiology 161:516–527.

9. Dienes L, Weinberger HJ. 1951. The L Forms of Bacteria. Bacteriol Rev 15:245–288.

10. Katz SM, Hashemi S, Brown KR, Habib WA, Hammel JM. 1984. Pleomorphism of Legionella Pneumophila. Ultrastruct Pathol 6:117–129.

11. Baron CS, Barve A, Muraro MJ, Linden R van der, Dharmadhikari G, Lyubimova A, Koning EJP de, Oudenaarden A van. 2019. Cell Type Purification by Single-Cell Transcriptome-Trained Sorting. Cell 179:527-542.e19.

12. Shapiro HM. 2003. Practical Flow Cytometry, 4th ed. John Wiley & Sons, Inc, New Jersey, USA.

13. Laflamme C, Verreault D, Lavigne S, Trudel L, Ho J, Duchaine C. 2005. Autofluorescence as a viability marker for detection of bacterial spores. Front Biosci 10:1647.

14. Blasi T, Hennig H, Summers HD, Theis FJ, Cerveira J, Patterson JO, Davies D, Filby A, Carpenter AE, Rees P. 2016. Label-free cell cycle analysis for high-throughput imaging flow cytometry. Nat Commun 7:10256.

15. Kiepś J, Juzwa W, Dembczyński R. 2023. Imaging Flow Cytometry Demonstrates
Physiological and Morphological Diversity within Treated Probiotic Bacteria Groups. Int J Mol
Sci 24:6841.

16. Pinto RN, Sebastian JA, Parsons MJ, Chang TC, Turner TR, Acker JP, Kolios MC. 2019.
Label-Free Analysis of Red Blood Cell Storage Lesions Using Imaging Flow Cytometry. Cytom
Part A 95:976–984.

17. Dey R, Rieger AM, Stephens C, Ashbolt NJ. 2019. Interactions of Pseudomonas aeruginosa
with Acanthamoeba polyphaga Observed by Imaging Flow Cytometry. Cytom Part A 95:555–
564.

18. Ebert D, Rainey P, Embley TM, Scholz D. 1996. Development, life cycle, ultrastructure and
phylogenetic position of *Pasteuria ramosa* Metchnikoff 1888: Rediscovery of an obligate
endoparasite of *Daphnia magna* Straus. Phil Trans R Soc B 351:1689–1701.

19. Ebert D, Duneau D, Hall MD, Luijckx P, Andras JP, Pasquier LD, Ben-Ami F. 2016. A
Population Biology Perspective on the Stepwise Infection Process of the Bacterial Pathogen
Pasteuria ramosa in Daphnia. Adv Parasit 91:265–310.

20. Duneau D, Luijckx P, Ben-Ami F, Laforsch C, Ebert D. 2011. Resolving the infection process
reveals striking differences in the contribution of environment, genetics and phylogeny to host-
parasite interactions. BMC biology 9:11.

21. Clerc M, Ebert D, Hall MD. 2015. Expression of parasite genetic variation changes over the
course of infection: implications of within-host dynamics for the evolution of virulence. Proc
Royal Soc B Biological Sci 282:20142820.

22. Fields PD, Reisser C, Dukić M, Haag CR, Ebert D. 2015. Genes mirror geography in
Daphnia magna. Mol Ecol 24:4521–4536.

23. Luijckx P, Ben-Ami F, Mouton L, Pasquier LD, Ebert D. 2010. Cloning of the unculturable
parasite Pasteuria ramosa and its Daphnia host reveals extreme genotype-genotype
interactions. Ecol Lett 14:125–31.

24. Klüttgen B, Dülmer U, Engels M, Ratte HT. 1994. ADaM, an artificial freshwater for the
culture of zooplankton. Water Res 28:743–746.

25. Ebert D, Zschokke-Rohringer CD, Carius HJ. 2000. Dose effects and density-dependent
regulation of two microparasites of Daphnia magna. Oecologia.

26. Ward MD, Kaduchak G. 2018. Fundamentals of Acoustic Cytometry. Curr Protoc Cytom
84:e36.

27. McInnes L, Healy J, Saul N, Großberger L. 2018. UMAP: Uniform Manifold Approximation
and Projection. J Open Source Softw 3:861.

28. Gassen SV, Callebaut B, Helden MJV, Lambrecht BN, Demeester P, Dhaene T, Saeys Y.
2015. FlowSOM: Using self-organizing maps for visualization and interpretation of cytometry
data. Cytom Part A 87:636–645.

29. Nitta N, Sugimura T, Isozaki A, Mikami H, Hiraki K, Sakuma S, Iino T, Arai F, Endo T,
Fujiwaki Y, Fukuzawa H, Hase M, Hayakawa T, Hiramatsu K, Hoshino Y, Inaba M, Ito T,
Karakawa H, Kasai Y, Koizumi K, Lee S, Lei C, Li M, Maeno T, Matsusaka S, Murakami D,
Nakagawa A, Oguchi Y, Oikawa M, Ota T, Shiba K, Shintaku H, Shirasaki Y, Suga K, Suzuki Y,
Suzuki N, Tanaka Y, Tezuka H, Toyokawa C, Yalikun Y, Yamada M, Yamagishi M, Yamano T,
Yasumoto A, Yatomi Y, Yazawa M, Carlo DD, Hosokawa Y, Uemura S, Ozeki Y, Goda K. 2018.
Intelligent Image-Activated Cell Sorting. Cell 175:266-276.e13.

30. Schraivogel D, Kuhn TM, Rauscher B, Rodríguez-Martínez M, Paulsen M, Owsley K,
Middlebrook A, Tischer C, Ramasz B, Ordoñez-Rueda D, Dees M, Cuylen-Haering S, Diebold
E, Steinmetz LM. 2022. High-speed fluorescence image–enabled cell sorting. Science 375:315–
320.

31. Stolarek I, Samelak-Czajka A, Figlerowicz M, Jackowiak P. 2022. Dimensionality reduction
by UMAP for visualizing and aiding in classification of imaging flow cytometry data. iScience
25:105142.

32. Marsh-Wakefield FM, Mitchell AJ, Norton SE, Ashhurst TM, Leman JK, Roberts JM, Harte
JE, McGuire HM, Kemp RA. 2021. Making the most of high-dimensional cytometry data.
Immunol Cell Biol 99:680–696.

33. Hennig H, Rees P, Blasi T, Kamentsky L, Hung J, Dao D, Carpenter AE, Filby A. 2017. An
open-source solution for advanced imaging flow cytometry data analysis using machine
learning. Methods 112:201–210.

34. Haridas V, Ranjbar S, Vorobjev IA, Goldfeld AE, Barteneva NS. 2017. Imaging flow
cytometry analysis of intracellular pathogens. Methods 112:91–104.

35. Power AL, Barber DG, Groenhof SRM, Wagley S, Liu P, Parker DA, Love J. 2021. The
Application of Imaging Flow Cytometry for Characterisation and Quantification of Bacterial
Phenotypes. Front Cell Infect Microbiol 11:716592.

520