Facultative endosymbiosis between cellulolytic protists and

methanogenic archaea in the gut of the Formosan termite *Coptotermes*

formosanus

Masayuki Kaneko,[1,†] Tatsuki Omori,[1,†] Katsura Igai,[1] Takako Mabuchi,[1] Miho

Sakai-Tazawa,[1] Arisa Nishihara,[2] Kumiko Kihara,[1,3] Tsuyoshi Yoshimura,[4,‡]

Moriya Ohkuma,[2] Yuichi Hongoh[1,2]

[1]Department of Life Science and Technology, Tokyo Institute of Technology, Tokyo,

Japan.

[2] Japan Collection of Microorganisms, RIKEN BioResource Research Center, Ibaraki,

Japan

[3] National Institute of Technology, Kumamoto College, Kumamoto, Japan

[4] Research Institute for Sustainable Humanosphere, Kyoto University, Kyoto, Japan

[†]Masayuki Kaneko and Tatsuki Omori should be considered joint first author.

[‡]Passed away in May 2021.

Corresponding author: Yuichi Hongoh

Email: yhongo@bio.titech.ac.jp

Postal address: 2-12-1 Ookayama, Meguro-ku, Tokyo 152-8550, Japan.

Tel: +81-3-5734-2865

ABSTRACT

Anaerobic protists frequently harbour methanogenic archaea, which apparently contribute to the hosts' fermentative metabolism by consuming excess H_2. However, the ecological properties of endosymbiotic methanogens remain elusive in many cases. Here we investigated the ecology and genome of the endosymbiotic methanogen of the cellulolytic *Cononympha* protists in the hindgut of the Formosan termite *Coptotermes formosanus*. Microscopic and 16S rRNA amplicon sequencing analyses revealed that a single species, designated here '*Candidatus* Methanobrevibacter cononymphae', is associated with both *Cononympha leidyi* and *Cononympha koidzumii* and that its infection rate in *Cononympha* cells greatly varied from 0.0 to 99.8% among termite colonies. Fine-scale network analysis indicated that multiple 16S rRNA sequence variants coexisted within a single host cell and that identical variants were present in both *Cononympha* species and also on the gut wall; '*Ca.* Methanobrevibacter cononymphae' is facultatively associated with *Cononympha* and horizontally taken up. Indeed, transmission electron microscopy showed escape or uptake of methanogens from/by a *Cononympha* cell. The genome of '*Ca.* Methanobrevibacter cononymphae' showed features consistent with its facultative lifestyle: i.e., the genome size (2.7 Mbp) comparable to those of free-living relatives; the pseudogenization of the formate dehydrogenase gene *fdhA*, unnecessary within the non-formate-producing host cell; the dependence on acetate as an essential carbon source, which is abundant within the host cell; and the presence of a catalase gene, required for colonization on the microoxic gut wall. Our study revealed a versatile endosymbiosis between the methanogen and protists, which may be a strategy responding to changing conditions in the termite gut.

INTRODUCTION

Methanogenic archaea, or methanogens, play a key role in the final step of the decomposition of organic matter in various anaerobic environments [1]. In the guts of termites, for example, which are keystone animals in the terrestrial carbon cycle, methanogens accounted for 0–10% of the prokaryotic gut community and possibly produce 1–3% of the global methane [2,3]. Methanogens are present in the guts of diverse termite lineages [4,5] and are considered to contribute to the gut ecosystem as H_2-sinks by producing methane from H_2 and CO_2 [6,7]. Methanogens are found on the hindgut epithelium [4,8,9] or within the cells of certain gut protist species [10]. The protists in the termite gut belong to the phylum *Parabasalia* or the order *Oxymonadida* in the phylum *Preaxostyla* and play major roles in the digestion of lignocellulose [11]. These gut protists generally harbour endo- or ectosymbiotic prokaryotes, which likely contribute to the host nutrition by supplying nitrogenous compounds [11] and/or removing H_2 [12,13]. In addition to the mutualistic symbionts, commensal or parasitic endosymbiotic bacteria are occasionally present [14].

Endosymbiotic methanogens are observed, for example, in the parabasalid protist *Trichomitopsis termopsidis* in the gut of the termite *Zootermopsis angusticolis* [15,16] and the oxymonad protist *Dinenympha parva* in the gut of the termite *Reticulitermes speratus* [17,18]. It has been hypothesized that endosymbiotic methanogens promote wood decomposition of the protist hosts by consuming excess H_2 [10], but their detailed physiology and ecology remain unclear. Indeed, although there are numerous examples of endosymbiosis between anaerobic protists and methanogens in various environments, the ecological properties of endosymbiotic methanogens,

including whether they are obligate or facultative endosymbionts, are elusive in many

cases [19,20].

The Formosan subterranean termite *Coptotermes formosanus* is the most

destructive pest of wooden constructions in southern China and Japan and has invaded

Hawaii and the southern part of the United States [21]. The termite harbours a relatively

simple protistan gut community consisting of five unculturable parabasalid species:

Pseudotrichonympha grassii, *Holomastigotoides hartmanii, Holomastigotoides minor,*

Cononympha [*Spirotrichonympha*] *leidyi*, and *Cononympha koidzumii* [22–24]. Among

them, endosymbiotic methanogens are found only in *Cononympha* cells [25,26], which

are mainly localized in the posterior part of the hindgut [27,28]. *Cononympha* protists

numerically dominate the protistan gut community, occupying 2,200–10,900 out of

3,800–12,900 total protist cells per gut [28]. Transcriptomic analysis of *Cononympha*

cells suggested that *Cononympha* hydrolysed cellulose, hemicellulose, pectin, and

chitin, contributing to both lignocellulose digestion and nitrogen recycling [24].

In the present study, we examined the ecology and metabolic capacity of the

uncultured endosymbiotic methanogen of *Cononympha* species in the gut of

Coptotermes formosanus using a combination of microscopy, fine-scale phylogenetic

analysis, and genome sequence analysis. Here, we show evidence of facultative

endosymbiosis by horizontal acquisition of the methanogen and its genome features

adapted to the lifestyle. Our study sheds light on another aspect of the multi-layered

symbiotic system in the termite gut.

MATERIALS AND METHODS

Termite collection and methanogen infection rate

Coptotermes formosanus (family Rhinotermitidae) were collected from four prefectures in Japan (Table S1). The termite colonies collected in 2019 or later were kept with their nest logs in plastic containers in a laboratory until use. Colonies collected in 2018 or earlier were kept in laboratories for years fed with red pine chips. The entire guts of three worker termites per colony were removed using sterile forceps, and the gut contents were suspended in sterile solution U [29]. The ratio of methanogen-containing *Cononympha* cells to total *Cononympha* cells in individual guts was examined on the basis of F_{420} autofluorescence under an Olympus BX51 epifluorescence microscope. The procedure of CH_4 measurement emitted by termites and semi-quantitative PCR amplification of the gene for methyl coenzyme M reductase (McrA) are described in Supplemental Methods.

Collection of single *Cononympha* cells and sequencing of 18S rRNA genes

Single *Cononympha* cells were collected using a Leica AM6000 micromanipulation system and subjected to whole genome amplification (WGA) using the illustra GenomiPhi V2 kit (GE Healthcare) as described previously [30]. To identify the species of *Cononympha*, the 18S rRNA gene was amplified by PCR using *Cononympha*-specific primers designed in this study (Table S2). The PCR conditions are described in Supplemental Methods. The PCR products were purified, cloned, and sequenced using the Sanger method as described previously [30].

16S rRNA amplicon sequencing analysis

The V3–V4 region (*ca*. 400 bp) of the 16S rRNA gene was amplified by PCR with prokaryote-universal primers, Pro341F and Pro805R (Table S2), using Phusion High-Fidelity DNA Polymerase (New England Biolabs), as described previously [31]. Purification, library preparation, and paired-end sequencing (300 bp × 2) on the MiSeq platform with the MiSeq Reagent Kit v3 (Illumina) were conducted as described previously [31]. The paired-end reads were trimmed, quality-filtered, and sorted into amplicon sequence variants (ASVs) using DADA2 v1.6 [32]. The ASVs were classified using SINA v1.2.11 [33] with the SILVA v132 database [34]. ASVs classified as *Eukarya* or undetermined and ASVs of which frequency was < 0.1% of the total reads were discarded from subsequent analyses.

Phylogenetic analysis of 16S rRNA genes

Near full-length 16S rRNA genes of methanogens were amplified by PCR using Phusion Hi-Fidelity DNA Polymerase. Primers M23F and M1382R broadly targeting methanogens were used for gut wall samples, while primer CfC-M23F adjusted to the sequence of the endosymbiotic methanogen was used as the forward primer for endosymbiotic methanogens (Table S2). The PCR conditions and procedures of sequencing and alignment are described in Supplemental Methods. A maximum-likelihood tree was constructed using IQ-TREE v1.6.12 [35] with the TVM+F+I+G4 nucleotide substitution model selected by ModelFinder implemented in IQ-TREE. Network construction of sequence variants (SVs) was conducted using the PopART program depending on Qt v4.8.5 with TCS algorithms [36].

Fluorescence in situ hybridization (FISH) and transmission electron microscopy (TEM)

To discriminate between *Cononympha leidyi* and *C. koidzumii* cells, oligonucleotide

probes specific to each 18S rRNA sequence were designed using ARB [37] (Table S2).

FISH was performed as described previously [38] with hybridization at 55°C for 2 h.

Observations were conducted under an Olympus BX51 epifluorescence microscope.

TEM was performed as described previously [39] using an H-7500 transmission

electron microscope (Hitachi).

Genome sequencing

Single *Cononympha* cells were collected as described above, washed several times in

droplets of solution U, and transferred to solution U containing 0.1% Tween 20 (Nacalai

Tesque). Prokaryotic cells that leaked out from the ruptured host cell were collected

with a glass capillary attached to the Leica AM6000 micromanipulation system and

subjected to WGA as described above. Nine WGA samples of single *Cononympha* cells

were prepared. These were subjected to paired-end sequencing on the MiSeq platform.

After assembling and binning as below, one sample was selected for deeper paired-end

sequencing and mate-pair sequencing on MiSeq, and long-read sequencing on the

MinION platform (Oxford Nanopore Technologies), based on the estimated genome

completeness of the target methanogen. The procedures for library preparation and

sequencing are described in Supplemental Methods.

Genome assembly

The MiSeq paired-end reads were trimmed, quality filtered, and assembled into contigs

using SPAdes v3.9 [40]. The contigs were binned using a combination of MyCC [41]

and Contig Annotation Tool (CAT) [42]. The results were evaluated for the selection of

the best sample. The MinION reads, after being quality trimmed, were assembled

together with the quality-trimmed MiSeq paired-end and mate-pair reads into contigs

using SPAdes v3.15.2 with the 'hybrid-assembly' and 'meta' mode [40]. The assembled

contigs ≥500 bp were binned using a combination of MyCC, CAT, RAST

(http://rast.nmpdr.org/rast.cgi), and BLASTn searches. The bin affiliated with

Methanobrevibacter was further subjected to scaffolding using LINKS v2.0.1 [43] and

gap-closing using TGS-GapCloser [44]. The quality-trimmed MiSeq and MinION reads

were mapped onto the gap-closed bin using BBMap v38.96 and Minimap2 v2.24-r1122

[45]. The reads mapped onto the bin were then reassembled using SPAdes v3.15.2 with

the 'hybrid-assembly' and 'sc' mode. Reassembled contigs ≥1,000 bp were binned as

described above. Details of quality filtering are described in Supplemental Methods.

Gene annotation

Finding and functional annotation of genes were performed using a combination of

DFAST [46] and BLASTp searches of the NCBI non-redundant (nr) protein database.

CRISPR-Cas systems were identified using CRISPRCasFinder [47]. Assignment of

COG (clusters of orthologous genes) categories [48] was conducted by RPS-BLAST

v2.6.0+ searches of the NCBI Conserved Domain Database v3.16, and the results were

manually curated. Pseudogenes were manually identified as described previously [49].

Metabolic pathways were inferred using the KEGG automatic annotation server

(KAAS) [50] and KEGG Mapper [51].

Phylogenomics

To construct a phylogenomic tree, we retrieved genome sequences of

Methanobacteriaceae as references from GTDB r202 [52] and metagenome-assembled

genomes (MAGs) assigned to *Methanobacteriaceae* recently obtained from termite guts

[5,53]. Gene prediction, extraction, and alignment are described in Supplemental

Methods. A maximum-likelihood tree was constructed using IQ-TREE v1.6.12 with the

LG+F+R5 amino acid substitution model selected by ModelFinder. The robustness of

the tree topology was evaluated using 1,000 ultrafast bootstrap resamplings. Average

nucleotide identity (ANI) and average amino acid identity (AAI) were calculated using

ANI/AAI-Matrix [54].

Comparative genome analysis

The relative abundance of genes assigned to COG functional categories was calculated

for genome sequences with >80% estimated completeness (Table S3). Principal

component analysis was performed after a central log-ratio transformation using a script

(https://github.com/dkato2021/COGplot.git). To compare genome contents with

identical criterion, the identification of intact genes or pseudogenes was automatically

performed using Pseudofinder v1.1.0 [55] and the NCBI Prokaryotic Genome

Annotation Pipeline (PGAP) [48].

RESULTS

Discovery of *Cononympha* cells lacking endosymbiotic methanogens

During our preliminary observations of protists in the guts of *Coptotermes formosanus*

by epifluorescence microscopy, we unexpectedly found that, in certain *Coptotermes*

formosanus colonies, no *Cononympha* cells emitted F_{420} autofluorescence as a

characteristic signal of the presence of endosymbiotic methanogens, unlike in other

Coptotermes formosanus colonies (Fig. 1A–D) or previous reports [25,26,28]. To

confirm the absence of methanogens in *Cononympha* cells in such termite colonies, we

performed amplicon sequencing analysis of the 16S rRNA V3–V4 region for

prokaryotic microbiota associated with individual *Cononympha* cells. A total of 30

single *Cononympha* cells collected from two *Coptotermes formosanus* colonies

(O2020a and O2020b in Table S1), in which almost all *Cononympha* cells were without

F_{420} autofluorescence, were examined. For comparison, we also conducted the same

experiment using 28 single *Cononympha* cells from two colonies (K2019b and K2019c

in Table S1), in which almost all *Cononympha* cells exhibited F_{420} autofluorescence. No

16S rRNA sequences of methanogens were obtained from the former 30 *Cononympha*

cells, whereas sequences of *Methanobrevibacter* were recovered from all of the latter 28

Cononympha cells (Fig. S1).

Infection rate of methanogens in *Cononympha* cells and methane emission rate

We examined the infection rate of methanogens in *Cononympha* cells for 33 colonies of

Coptotermes formosanus collected from four prefectures in Japan (Table S1), based on

the detection of F_{420} autofluorescence. The infection rate greatly varied from 0 to 99.8%

depending on the termite colonies (Fig. 1E; Tables S1 and S4). This difference was

irrespective of natural or laboratory-reared termite colonies and prefectures of the

sampling sites (Fig. 1E; Tables S1 and S4).

To test whether the presence or absence of the endosymbiotic methanogens of

Cononympha is linked to the methane emission rate from the termite hosts, we chose

three *Coptotermes formosanus* colonies of the high and low endosymbiotic

methanogen-infection rate types, respectively (Table S1), and measured the methane

emission rates from termites (Fig. 1E; Supplementary Methods). The results showed

that termites from the colonies of the 'high infection rate' type emitted 4.5–7.8 times

more methane than those from the 'low infection rate' type colonies (Fig. 1E). Semi-

quantitative PCR targeting the *mcrA* gene in the entire gut microbiota generated results

consistent with the methane emission rates (Fig. S2; Supplementary Methods). No *mcrA*

amplification was detected in any of the four tested *Coptotermes formosanus* colonies

of the 'low infection rate' type under the PCR condition, where *mcrA* amplicons were

detected in all six tested colonies of the 'high infection rate' type (Fig. S2; Table S1).

Because the hindgut wall of *Coptotermes formosanus* can be another habitat of

methanogens [56], we examined F_{420} autofluorescence signals on gut wall fragments

from termite colonies of both 'high infection rate' and 'low infection rate' types. We

observed dense colonization of rod-shaped methanogens on fragments of the hindgut

wall from two out of seven colonies of the 'high infection rate' type, whereas almost no

signals were detected in termites from four colonies of the 'low infection rate' type (Fig.

S3; Table S1).

Fine-scale phylogenetic composition and host specificity of endosymbiotic

methanogens

In the above 16S rRNA amplicon sequencing analysis (Fig. S1), multiple ASVs

belonging to *Methanobrevibacter* were obtained from each *Cononympha* cell (Fig. S4).

Nine closely related ASVs were obtained in total, showing only 1–4 mismatches out of

386 bp. Among them, the ASV-3 sequence was identical to the corresponding region of

clone SlMeN10 (AB360373), which was derived from *Cononympha* cells in a

Coptotermes formosanus gut [26]. Since the genome of the endosymbiotic methanogen

contains only a single rRNA operon as described below, these multiple ASVs were not

derived from intragenomic variation. These results indicated that multiple strains of a

single *Methanobrevibacter* species frequently coexisted within a single *Cononympha*

cell.

Next, we examined whether both *Cononympha* species, i.e., *C. leidyi* and *C.*

koidzumii, can harbour methanogens. We obtained the 18S rRNA gene sequences of

single *Cononympha* cells used in the above 16S rRNA amplicon analysis (Fig. S1) and

found that both *Cononympha* species housed methanogens in the gut of termites from

colonies of the 'high infection rate' type (Fig. S5; Table S5). Conversely, in colonies of

the 'low infection rate' type, both *Cononympha* species lacked methanogens (Fig. S5;

Table S5). We observed that both *Cononympha* species, visually discriminated by FISH

specifically targeting 18S rRNA of each species, harboured *Methanobrevibacter*-like

rods (Fig. S6).

We further investigated whether the lineages of the endosymbiotic methanogens

differ between the two *Cononympha* host species and whether the methanogens on the

gut wall are the same species as the endosymbionts or not. To examine these with

higher resolution, we obtained near full-length 16S rRNA gene sequences of

methanogens by PCR amplification from 14 single *Cononympha leidyi* cells, three

single *C. koidzumii* cells, and seven fragments of the hindgut wall (Fig. 2; Tables S1

and S5). Three out of four 16S rRNA gene SVs of methanogens associated with *C.*

koidzumii cells were also obtained from *C. leidyi* cells. In addition, we obtained 16S

rRNA gene amplicons of methanogens from three of the seven gut wall fragments and

sequenced 24 clones in total, all of which showed 99–100% homology to SVs of the

endosymbionts (Fig. 2). These SVs from the gut wall or *Cononympha* cells formed

clusters that were delineated mainly by the host *Coptotermes formosanus* colonies (Fig.

2); thus, the sequence variations of these methanogens were strongly related to the host

termite colony and not to the *Cononympha* species.

These data together indicated that the endosymbiosis between the *Cononympha*

protists and the methanogens is facultative and that the two *Cononympha* species can

horizontally acquire the same *Methanobrevibacter* species. Furthermore, the spatial

distribution of the methanogen most likely extends to the gut wall of the termite host.

The shape and size of the methanogens on the gut wall (Fig. S3A) and within

Cononympha cells were similar. Both methanogens were straight rods, and the size was

1.1–1.8 µm by 0.4–0.5 µm (n = 50) in the former while it was 1.3–1.8 µm by 0.4–0.5

μm (n = 50) in the latter.

TEM of endosymbiotic methanogens

The facultative nature of this endosymbiosis was further corroborated by TEM. An

electron micrograph showed that electron-dense, rod-shaped prokaryotic cells,

morphologically indistinguishable, were present not only within a *Cononympha* cell but

also on its surface (Fig. 3 and S7A–D). They appeared to be in the process of phago- or

exocytosis by the *Cononympha* cell (Fig. 3 and S7A). These prokaryotic cells

morphologically resembled cells of cultured *Methanobrevibacter* species [8,9,57]. The

intracellular ones were surrounded by a host membrane (Fig. 3 and S7A–D), and some

were localized proximal to putative hydrogenosomes (Fig. 3 and S7C). Intracellular

Methanobrevibacter-like cells apparently during fission were observed (Fig. S7D); they

likely proliferate within the host *Cononympha* cell. We hereafter designated this

endosymbiotic methanogen as '*Candidatus* Methanobrevibacter cononymphae'

(abbreviated as *Mbv. cononymphae*), and the species description is presented below.

Phylogenetic position of *Mbv. cononymphae* based on 16S rRNA gene

Based on the near full-length 16S rRNA gene, *Mbv. cononymphae* was placed within a

clade exclusively comprising uncultured clones derived from termite guts (Fig. S8).

Within this clade, *Mbv. cononymphae* was closely related to clones obtained from the

protists *Microjoenia* sp. and *D. parva* in the gut of *R. speratus* [17]. This clade further

formed a monophyletic cluster with termite-gut-derived sequences, including cultured

species inhabiting the gut epithelium of the protist-dependent termite *Reticulitermes*

flavipes, i.e., *Methanobrevibacter curvatus* [8] and *Methanobrevibacter filiformis* [9].

Endosymbiotic *Methanobrevibacter* housed by an anaerobic ciliate, *Nyctotherus ovalis*,

in the hindgut of the cockroach *Blaptica dubia* [58] and that housed by an aquatic

anaerobic ciliate, *Trimyema compressum* [20], were more distantly related (Fig. S8).

General genome features of *Mbv. cononymphae*

We obtained a draft genome sequence of *Mbv. cononymphae* from a single

Cononympha cell. The genome consisted of 245 contigs with 100% and 98.6%

completeness estimated using CheckM [59] and CheckM2 [60], respectively (Table 1).

Although only a single rRNA operon was identified in this genome as in many other

Methanobrevibacter species (Table 1), we recovered four near full-length 16S rRNA

gene SVs showing 3–5 base mismatches by PCR amplification from the same DNA

sample. Thus, this genome sequence was derived from multiple, at least four,

genomovars and here designated as the composite genome 'CfCl-M3'.

The total contig length was 2,709,899 bp, and 1,834 protein-coding sequences

(CDSs) were predicted, excluding 90 CDSs truncated by contig ends and 203 CDSs

<100 amino acid sequences that were automatically predicted but with no significant

hits in BLASTp searches of the NCBI nr protein database. At least, two CRISPR-Cas

systems were identified with a total of 282 spacer sequences (Tables S6–8). Most

spacers showed no sequence similarity to those in CRISPRCasdb [61]. The number of

pseudogenes was 148 after manual inspection (Table 1), and 14 were assigned to

category [V] (defence mechanism) of COGs, including DNA restriction-modification

systems and *cas* genes (Fig. S9; Table S6). Finding of duplicated regions and the

frequency of single nucleotide polymorphisms in the CfCl-M3 composite genome are

described in Supplemental Methods, Supplemental Results, and Table S9.

Phylogenomics and comparative genome analysis

A maximum-likelihood tree based on concatenated amino acid sequences of conserved

single-copy genes (Fig. 4; Table S3) was basically congruent with the 16S rRNA gene

tree (Fig. S8). *Mbv. cononymphae* belonged to a monophyletic cluster consisting

exclusively of MAGs and two *Methanobrevibacter* isolates obtained from the guts of

protist-dependent termites. These genomes shared at least 75% ANI and 60% AAI (Fig.

S10). This clade was sister to a clade comprising genome sequences from protist-

dependent/independent termite guts or other environments (Fig. 4).

Genome sequences derived from termite guts in these two clades (Fig. 4) tended

to possess more genes assigned to the COG categories [L] (replication, recombination

and repair) and [V] (defence mechanisms) (Fig. S11). Among the genes, those related to

DNA methylation and drug resistance were especially abundant in termite-gut derived

genomes (Fig. S12). No clear differences in the number and ratio of genes assigned to

respective COG categories were observed among the five *Methanobrevibacter* species

in Table 1, except for a much smaller number and ratio of genes assigned to [V]

(defence mechanism) in *Methanobrevibacter* sp. NOE (Fig. S13), which is likely an

obligate endosymbiont [57].

The number of pseudogenes of *Mbv. cononymphae* was approximately 1.5–2.1

times larger than those of the gut-wall dwelling *Mbv. curvatus*, *Mbv. filiformis*, and

Methanobrevibacter cuticularis, but smaller (ca. 80%) than that of *Methanobrevibacter*

sp. NOE (Table 1). The distribution pattern of pseudogenes among COG categories

indicated that pseudogenes were characteristically accumulated in *Mbv. cononymphae*

in categories [L] (replication, recombination, and repair), [V] (defence mechanism), and

[X] (mobilome; prophages, transposons) (Fig. S14). No unique genes were identified in

Mbv. cononymphae among COG-assigned 911 CDSs when compared to other known

Methanobrevibacter species (Fig. S15; Table S10). Details are described in

Supplemental Results.

Predicted metabolism of *Mbv. cononymphae*

All genes for enzymes required for energy conservation through methanogenesis were

present in *Mbv. cononymphae* (Fig. 5). Similar to other *Methanobrevibacter* species, it

likely produces methane from H_2 and CO_2. On the other hand, the formate

dehydrogenase subunit alpha gene *fdhA* is pseudogenized, and the gene for formate

transporter (*fdhC*) was missing (Fig. S16). Intact *fdhA* and *fdhC* genes were not found in

sequence reads outside the assembled genome or in the other eight samples that were

not selected for deeper sequencing analysis. In addition, genes for the acetyl-CoA

decarbonylase/synthase (ACDS) multienzyme complex were absent; thus, *Mbv.*

cononymphae apparently requires acetate to produce acetyl-CoA. The cobalamin

biosynthetic pathway was absent, and instead, genes for the cobalamin/siderophore

transporter FepBCD were identified (Table S5). These characteristics, i.e., loss or

absence of genes for Fdh, ACDS, and cobalamin biosynthesis, were shared by most

MAGs derived from the guts of protist-dependent termite species, but not shared by the

gut wall-dwelling species *Mbv. curvatus*, *Mbv. filiformis*, and *Mbv. cuticularis* [8,9]

(Fig. 4).

Mbv. cononymphae possesses genes for ammonium transporter AmtB and

glutamine synthetase GlnA. The genome also encoded a major facilitator superfamily

transporter for di- and tripeptides, which may be other nitrogen sources (Fig. 5). The

amino acid and cofactor synthesis capabilities were similar to those of other

Methanobrevibacter species (Table S11). Genes for proteins involved in oxygen

tolerance, including catalase KatE, were identified (Fig. 4), and details are described in

Supplemental Results. No genes related to motility were found.

DISCUSSION

This study unveiled the ecology and genomic features of the endosymbiotic methanogen

Mbv. cononymphae harboured by two *Cononympha* species in the gut of *Coptotermes*

formosanus. A series of evidence obtained by a combination of F_{420} autofluorescence

detection, small subunit (SSU) rRNA gene sequence analysis, and TEM, clearly

indicated that their association is facultative and that the methanogens are most likely

transmitted in 'mixed mode' [62], i.e., both vertically and horizontally among

Cononympha cells across the two *Cononympha* species. In addition, our data suggest

that *Mbv. cononymphae* colonizes the hindgut wall. This facultative lifestyle is

consistent with its genomic features such as the genome size comparable to those of

free-living relatives, the number of pseudogenes intermediate between free-living

relatives and the putatively obligate endosymbiont NOE of the anaerobic ciliate *N.*

ovalis [58] (Table 1), and the presence of a catalase gene (CfCl-M3_0439, Table S6).

Catalase activity was previously observed in *Mbv. filiformis*, *Mbv. curvatus*, and

Mbv. cuticularis, all of which attach to the microoxic hindgut wall of *R. flavipes* [8,9].

The catalase gene *katE* was commonly found in the clade containing *Mbv.*

cononymphae and its sister clade; however, it is pseudogenized in *Methanobrevibacter*

sp. NOE (Fig. 4). This is possibly because NOE, unlike *Mbv. cononymphae*, has

become an obligate endosymbiont that rarely encounters a high concentration of oxygen.

Assuming that *Cononympha* cells produce H_2, CO_2, and acetate as the main

fermentation products but do not produce formate, as observed in other cellulolytic

parabasalids [63,64] and predicted based on the transcriptome of *Cononympha* [24] (see

Supplemental Methods), the loss of *fdhC* and pseudogenization of *fdhA* (Fig. S16) are

likely attributable to endosymbiosis. The amount of formate is generally limited also in

the entire hindgut of termites [7,65]. Although the *fdh* operon is apparently intact in the

genomes of the gut wall-dwelling species *Mbv. filiformis*, *Mbv. curvatus*, and *Mbv.*

cuticularis (Fig. S16), these isolates exhibited only weak or no growth on formate [8,9].

This implies that the ability to utilize formate as the sole energy and carbon source is

very poor in these isolates. It is conceivable that the reduced opportunity and/or

requirement to use formate has allowed the loss of functional *fdh* genes in *Mbv.*

cononymphae. The lack of ACDS in *Mbv. cononymphae* may also be related to

endosymbiosis. The expectedly high concentration of acetate within the host protist cell

may have allowed the endosymbiont to depend solely on acetate to produce acetyl-CoA

(Fig. 5).

The absence of the functional biosynthetic pathway of cobalamin, which is

essential in methanogenesis [66] (Fig. 5), was previously reported also in the

endosymbiotic methanogens *Methanobrevibacter* sp. NOE and *Methanocorpusculum* sp.

MCE [58]. Thus, its absence and dependence on a transporter may be a common trait in

endosymbiotic methanogens. These characteristics, i.e., the absence of functional *fdh*,

ACDS genes, cobalamin biosynthetic pathway, and the presence of the catalase gene

were common in most MAGs from the guts of protist-dependent termites (Fig. 4). This

implies that these methanogens possibly have a lifestyle similar to that of *Mbv.*

cononymphae, i.e., facultative endosymbiosis, although their localizations are

unidentified. Among them, MAG_NG87 from *Coptotermes elisae* [53] and

MAG_GCA031280375 from *Prorhinotermes canalifrons* [5] showed high ANIs, 98.8%

and 96.2%, respectively, with *Mbv. cononymphae*, and both termite genera harbour

Cononympha species [67]; these MAGs are most likely of endosymbionts of

Cononympha with similar lifestyles.

It is widely believed that hydrogenotrophic methanogens critically contribute to

the fermentative metabolism of H_2-producing anaerobic protist hosts as syntrophic

mutualists [19]. Indeed, retarded growth of anaerobic ciliates that had lost their

endosymbiotic methanogens was previously observed [68,69]. In particular, removal of

'Candidatus Methanoregula pelomyxae', an endosymbiont of *Pelomyxa schiedti*, led to

the death of the anaerobic amoeba host [70]. In the case of termite-gut protists, Messer

and Lee (1989) [16] reported that *Trichomitopsis termopsidis* treated with 2-

bromoethanesulfonate (BES) inhibiting methanogenesis was viable in the gut of *Z.*

angusticollis. However, in an experiment using cultured *Trichomitopsis termopsidis,* the

growth of the protist was severely retarded when methanogenesis was suppressed by

BES treatment [64]. Its growth was recovered when fed with a more favourable nutrient

source in addition to cellulose, i.e., autoclaved cells of a specific bacterial strain [64].

This reminded other reports that anaerobic ciliates during successive cultures under

optimal conditions tended to lose their endosymbiotic methanogens, whereas ciliates

under food-limiting conditions or at low temperatures kept housing methanogens

[71,72]. These previous results imply that the loss of *Mbv. cononymphae* may be related

to the nutritional condition of the *Cononympha* host. It is unclear whether *Mbv.*

cononymphae contributes to the nutrition of the *Cononympha* host, not only by

consuming H_2, but also by supplementing nitrogenous compounds as suggested in

several cases of obligate endosymbioses between termite-gut protists and bacteria

[12,49,73–75].

Although *Mbv. cononymphae* can also likely colonize the gut wall, the methane

emission rate shown in Fig. 1E indicated that the absence of the endosymbiont was not

compensated by methanogens outside *Cononympha* cells. Large differences in the

methane emission rate among colonies were previously observed in other termite

species, i.e., *R. speratus* [2]*, N. sugioi* [*koshunensis*] [56], and *Cryptotermes secundus*

[2,7]. Thus, the facultative association of methanogens not only with gut protists but

also with termite hosts may be not rare phenomena. Although what factors determine

the infection rate of methanogens in gut protists and their total abundance in the termite

gut remain unclear, methanogens might play a role in adjusting the fermentation process

of the protist hosts and the whole termite gut ecosystem in response to changing

nutritional conditions with their versatile localization and abundance.

Description of '*Candidatus* Methanobrevibacter cononymphae' sp. nov

Methanobrevibacter cononymphae (co.no.nym'phae. N.L. fem. n. *Cononymphae*, of

Cononympha, a genus of flagellated protist hosts). The archaea are straight rods with

dimensions of 1.3–1.8 µm (mean ± SD, 1.58 ± 0.1; *n* = 50) by 0.4–0.5 µm (0.4 ± 0.0).

The archaea are non-motile and facultatively colonise the cytoplasm of *Cononympha*

leidyi and *Cononympha koidzumii* in the gut of *Coptotermes formosanus*. The archaea

occasionally colonize the hindgut wall of *Coptotermes formosanus*. The assignment is

based on the 16S rRNA gene (LC802674). The draft genome sequence CfCl-M3

showed 96.2% average nucleotide identity to *Methanovirga procula,* which has been

named under SeqCode on the basis of a MAG (GCA_031280375) obtained from the gut

of *Prorhinotermes canalifrons* [5].

DATA AVAILABILITY

The 16S rRNA amplicon and metagenome sequence data generated in this study have

been deposited in DDBJ under the accession numbers DRR534530–534599

(BioProject: PRJDB17265). Accession numbers for the CfCl-M3 genome are

BAABUN010000001–BAABUN010000245. Representative near full-length SSU

rRNA gene sequences will appear under the accession numbers LC802720–61 (18S

rRNA) and LC802666–82 (16S rRNA).

AUTHOR CONTRIBUTIONS

Y.H. and M.O. conceived the study and provided equipment and reagents; Y.H., K.K.,

and T.Y. designed the study; Y.H. K.K., T.Y., K.M., T.O, and M.S.-T. collected

termites; T.O., M.K., M.S.-T., and K.I. performed experiments, M.K., T.O., and Y.H.

analysed and interpreted the data; T.M. performed TEM; A.N., M.K, and K.I. measured

methane; M.K., K.I., and Y.H. wrote the paper with input from the other authors.

ACKNOWLEDGEMENTS

We are grateful to Hirokazu Kuwahara for supporting the collection of protist cells and

genome sequencing, to Takumi Murakami for handling sequence data, and to Katsuhiro

Aminaka for helping with micromanipulation. We thank Yukihiro Kinjo, Osamu

Kitade, Shuji Itakura, Satoko Noda, Wakako Omura, Masao Oya, Kazuki Takahashi,

Gaku Tokuda, and Akinori Yamada for their cooperation in termite collection. We also

thank Gillian Gile for her helpful advice on the morphological discrimination of

Cononympha species, Satoshi Hattori for his advice on the metabolism of

Methanobrevibacter, and Naoya Shinzato for unpublished information on

endosymbiotic methanogens of *Trimyema compressum*. We thank Michiru Shimizu,

Jun-ichi Inoue, and Masahiro Yuki for assisting with Sanger sequencing, which was

performed in the RIKEN BSI. Sanger sequencing was also performed in the Biomaterial

Analysis Center in the Tokyo Institute of Technology. TEM was assisted by Keiko

Ikeda in the latter facility.

COMPETING INTERSTS

The authors declare no competing interests.

REFERENCES

1. Jabłoński S, Rodowicz P, Łukaszewicz M. Methanogenic archaea database

containing physiological and biochemical characteristics. *Int J Syst Evol Microbiol*

2015;65:1360–8.

2. Sugimoto A, Inoue T, Tayasu I et al. Methane and hydrogen production in a termite-

symbiont system. *Ecol Res* 1998;13:241–57.

3. Brauman A, Doré J, Eggleton P et al. Molecular phylogenetic profiling of prokaryotic

communities in guts of termites with different feeding habits. *FEMS Microbiol Ecol*

2001;35:27–36.

4. Brune A. Methanogens in the digestive tract of termites. In: Hackstein JHP (ed),

Microbiology Monographs: (Endo)symbiotic Methanogenic Archaea. Switzerland:

Springer Nature, 2018, 81–101.

5. Protasov E, Nonoh JO, Kastle Silva JM et al. Diversity and taxonomic revision of

methanogens and other archaea in the intestinal tract of terrestrial arthropods. *Front*

Microbiol 2023;14:e1281628.

6. Breznak JA, Switzer JM. Acetate synthesis from H_2 plus CO_2 by termite gut

microbes. *Appl Environ Microbiol* 1986;52:623–30.

7. Pester M, Brune A. Hydrogen is the central free intermediate during lignocellulose

degradation by termite gut symbionts. *ISME J* 2007;1:551–65.

8. Leadbetter JR, Breznak JA. Physiological ecology of *Methanobrevibacter cuticularis*

sp. nov. and *Methanobrevibacter curvatus* sp. nov., isolated from the hindgut of the

termite *Reticulitermes flavipes*. *Appl Environ Microbiol* 1996;62:3620–31.

9. Leadbetter JR, Crosby LD, Breznak JA. *Methanobrevibacter filiformis* sp. nov., a

filamentous methanogen from termite hindguts. *Arch Microbiol* 1998;169:287–92.

10. Hongoh Y, Ohkuma M. Termite gut flagellates and their methanogenic and

eubacterial symbionts. In: Hackstein JHP (ed), Microbiology Monographs:

(Endo)symbiotic Methanogenic Archaea. Switzerland: Springer Nature, 2018, 55–79.

11. Brune A. Symbiotic digestion of lignocellulose in termite guts. *Nat Rev Microbiol*

2014;12:168–80.

12. Ohkuma M, Noda S, Hattori S et al. Acetogenesis from H_2 plus CO_2 and nitrogen

fixation by an endosymbiotic spirochete of a termite-gut cellulolytic protist. *Proc Natl*

Acad Sci U S A 2015;112:10224–30.

13. Takeuchi M, Kuwahara H, Murakami T et al. Parallel reductive genome evolution in

Desulfovibrio ectosymbionts independently acquired by *Trichonympha* protists in the

termite gut. *ISME J* 2020;14:2288–301.

14. Takahashi K, Kuwahara H, Horikawa Y et al. Emergence of putative energy

parasites within *Clostridia* revealed by genome analysis of a novel endosymbiotic clade.

ISME J 2023;17:1895–906.

15. Lee MJ, Schreurs PJ, Messer AC et al. Association of methanogenic bacteria with

flagellated protozoa from a termite hindgut. *Curr Microbiol* 1987;15:337–41.

16. Messer AC, Lee MJ. Effect of chemical treatments on methane emission by the

hindgut microbiota in the termite *Zootermopsis angusticollis*. *Microb Ecol*

1989;18:275–84.

17. Tokura M, Ohkuma M, Kudo T. Molecular phylogeny of methanogens associated

with flagellated protists in the gut and with the gut epithelium of termites. *FEMS*

Microbiol Ecol 2000;33:233–40.

18. Hara K, Shinzato N, Oshima T et al. Endosymbiotic *Methanobrevibacter* species

living in symbiotic protists of the termite *Reticulitermes speratus* detected by fuorescent

in situ hybridization. *Microbes Environ* 2004;19:120–7.

19. Fenchel T, Finlay BJ. Free-living protozoa with endosymbiotic methanogens. In:

Hackstein JHP (ed), Microbiology Monographs: (Endo)symbiotic Methanogenic

Archaea. 2nd ed. Switzerland: Springer Nature, 2018, 1–11.

20. Shinzato N, Takeshita K, Kamagata Y. Methanogenic and bacterial endosymbionts

of free-living anaerobic ciliates. In: Hackstein JHP (ed), Microbiology Monographs:

(Endo)symbiotic Methanogenic Archaea. Switzerland: Springer Nature, 2018, 37–53.

21. Evans TA, Forschler BT, Grace JK. Biology of invasive termites: a worldwide

review. *Ann Rev Entomol* 2013;58:455–74.

22. Koidzumi M. Studies on the intestinal protozoa found in the termites of Japan.

Parasitol 1921;13:235–309.

23. Jasso☐Selles DE, De Martini F, Velenovsky JF et al. The complete protist symbiont

communities of *Coptotermes formosanus* and *Coptotermes gestroi*☐: morphological

and molecular characterization of five new species. *J Euk Microbiol* 2020;67:626–41.

24. Nishimura Y, Otagiri M, Yuki M et al. Division of functional roles for termite gut

protists revealed by single-cell transcriptomes. *ISME J* 2020;14:2449–60.

25. Tsunoda K, Ohmura W, Yoshimura TK et al. Methane emission by the termite,

Coptotermes formosanus Shiraki (Isoptera: Rhinotermitidae). *Wood Res* 1993;79:34–

40.

26. Inoue J, Noda S, Hongoh Y et al. Identification of endosymbiotic methanogen and

ectosymbiotic spirochetes of gut protists of the termite *Coptotermes formosanus*.

Microbes Environ 2008;23:94–7.

27. Lai P, Tamashiro M, Fujii JK. Abundance and distribution of the three species of

symbiotic protozoa in the hindgut of *Coptotermes formosanus* (Isoptera:

Rhinotermitidae). *Proc Hawaiian Entomol Soc* 1983;24:271–6.

28. Yoshimura T. Contribution of the protozoan fauna to nutritional physiology of the

lower termite, *Coptotermes formosanus* Shiraki (Isoptera: Rhinotermitidae). *Wood Res*

1995;68–129.

29. Trager W. The cultivation of a cellulose-digesting fagellate, *Trichomonas*

termopsidis, and of certain other termite protozoa. *Biol Bull* 1934;66:182–90.

30. Igai K, Kitade O, Fu J et al. Fine-scale genetic diversity and putative ecotypes of

oxymonad protists coinhabiting the hindgut of *Reticulitermes speratus*. *Mol Ecol*

2022;31:1317–31.

31. Murakami T, Segawa T, Dial R et al. Bacterial microbiota associated with the

glacier ice worm Is dominated by both worm-specific and glacier-derived facultative

lineages. *Microbes Environ* 2017;32:32–9.

32. Callahan BJ, McMurdie PJ, Rosen MJ et al. DADA2: high-resolution sample

inference from Illumina amplicon data. *Nat Methods* 2016;13:581–3.

33. Pruesse E, Peplies J, Glockner FO. SINA: accurate high-throughput multiple

sequence alignment of ribosomal RNA genes. *Bioinformatics* 2012;28:1823–9.

34. Quast C, Pruesse E, Yilmaz P et al. The SILVA ribosomal RNA gene database

project: improved data processing and web-based tools. *Nucleic Acids Res*

2013;41:D590–6.

35. Nguyen L-T, Schmidt HA, von Haeseler A et al. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*

2015;32:268–74.

36. Leigh JW, Bryant D. popart: full-feature software for haplotype network

construction. *Methods Ecol Evol* 2015;6:1110–6.

37. Ludwig W, Strunk O, Westram , et al. ARB: a software environment for sequence

data. *Nucleic Acids Res* 2004;32:1363–13671.

38. Utami YD, Kuwahara H, Murakami T et al. Phylogenetic diversity and single-cell

genome analysis of '*Melainabacteria*', a non-photosynthetic cyanobacterial group, in

the termite gut. *Microbes Environ* 2018;33:50–7.

39. Sato T, Kuwahara H, Fujita K et al. Intranuclear verrucomicrobial symbionts and

evidence of lateral gene transfer to the host protist in the termite gut. *ISME J*

2014;8:1008–19.

40. Bankevich A, Nurk S, Antipov D et al. SPAdes: a new genome assembly algorithm

and its applications to single-cell sequencing. *J Comput Biol* 2012;19:455–77.

41. Lin H-H, Liao Y-C. Accurate binning of metagenomic contigs via automated

clustering sequences using information of genomic signatures and marker genes. *Sci*

Rep 2016;6:e24175.

42. von Meijenfeldt FAB, Arkhipova K, Cambuy DD et al. Robust taxonomic

classification of uncharted microbial sequences and bins with CAT and BAT. *Genome*

Biol 2019;20:e217.

43. Warren RL, Yang C, Vandervalk BP et al. LINKS: Scalable, alignment-free

scaffolding of draft genomes with long reads. *GigaScience* 2015;4:e35.

44. Xu M, Guo L, Gu S et al. TGS-GapCloser: A fast and accurate gap closer for large

genomes with low coverage of error-prone long reads. *GigaScience* 2020;9:1–11.

45. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*

2018;34:3094–100.

46. Tanizawa Y, Fujisawa T, Nakamura Y. DFAST: a flexible prokaryotic genome

annotation pipeline for faster genome publication. *Bioinformatics* 2018;34:1037–9.

47. Couvin D, Bernheim A, Toffano-Nioche C et al. CRISPRCasFinder, an update of

CRISRFinder, includes a portable version, enhanced performance and integrates search

for Cas proteins. *Nucleic Acids Res* 2018;46:W246–51.

48. Tatusova T, DiCuccio M, Badretdin A et al. NCBI prokaryotic genome annotation

pipeline. *Nucleic Acids Res* 2016;44:6614–24.

49. Hongoh Y, Sharma VK, Prakash T et al. Complete genome of the uncultured

Termite Group 1 bacteria in a single host protist cell. *Proc Nat Acad Sci U S A*

2008;105:5555–60.

50. Moriya Y, Itoh M, Okuda S et al. KAAS: an automatic genome annotation and

pathway reconstruction server. *Nucleic Acids Res* 2007;35:W182–5.

51. Kanehisa M, Sato Y. KEGG Mapper for inferring cellular functions from protein

sequences. *Protein Sci* 2020;29:28–35.

52. Parks DH, Chuvochina M, Chaumeil PA et al. A complete domain-to-species

taxonomy for Bacteria and Archaea. *Nat Biotechnol* 2020;38:1079–86.

53. Arora J, Kinjo Y, Šobotník J et al. The functional evolution of termite gut

microbiota. *Microbiome* 2022;10:e78.

54. Rodriguez-R LM, Konstantinidis KT. The enveomics collection: a toolbox for

specialized analyses of microbial genomes and metagenomes. *PeerJ* 2016;4:e1900v1.

55. Syberg-Olsen MJ, Garber AI, Keeling PJ, et al. Pseudofinder: detection of

pseudogenes in prokaryotic genomes. *Mol Biol Evol* 2022;39:msac153.

56. Shinzato N, Yoshino H, Yara K. Methane production by microbial symbionts in the

lower and higher termites of the Ryukyu archipelago. In Endocytobiology V:

Proceedings of the Fifth International Colloquium on Endocytobiology and Symbiosis,

Kyoto, June 23–27, 1992. Sato S., Ishida M., and Ishikawa H. (eds), Kyoto: Tübingen

University Press, 1993, 161–6.

57. Miller TL. Methanobrevibacter. In: Boone DR, Castenholz RW, Carrity GM, (eds),

Bergey's Manual of Systematics Bacteriology. 2nd ed. New York: Springer Verlag,

1989, 218–26.

58. Lind AE, Lewis WH, Spang A et al. Genomes of two archaeal endosymbionts show

convergent adaptations to an intracellular lifestyle. *ISME J* 2018;12:2655–67.

59. Parks DH, Imelfort M, Skennerton CT et al. CheckM: assessing the quality of

microbial genomes recovered from isolates, single cells, and metagenomes. *Genome*

Res 2015;25:1043–55.

60. Chklovski A, Parks DH, Woodcroft BJ et al. CheckM2: a rapid, scalable and

accurate tool for assessing microbial genome quality using machine learning. *Nat*

Methods 2023;20:1203–12.

61. Pourcel C, Touchon M, Villeriot N et al. CRISPRCasdb a successor of CRISPRdb

containing CRISPR arrays and *cas* genes from complete genome sequences, and tools

to download and query lists of repeats and spacers. *Nucleic Acids Res* 2020;48:D535–

44.

62. Ebert D. The epidemiology and evolution of symbionts with mixed-mode

transmission. *Annu Rev Ecol Evol Syst* 2013;44:623–43.

63. Yamin MA. Cellulose metabolism by the flagellate *Trichonympha* from a termite Is

independent of endosymbiotic bacteria. *Science* 1981;211:58–9.

64. Odelson DA, Breznak JA. Nutrition and growth characteristics of *Trichomitopsis*

termopsidis, a cellulolytic protozoan from termites. *Appl Environ Microbiol*

1985;49:614–21.

65. Odelson DA, Breznak JA. Volatile fatty acid production by the hindgut microbiota

of xylophagous termites. *Appl Environ Microbiol* 1983;45:1602–13.

66. Weiss DS, Thauer RK. Methanogenesis and the unity of biochemistry. *Cell*

1993;72:819–22.

67. Yamin MA. Flagellates of the orders Trichomonadida Kirby, Oxymonadida Grasse,

and Hypermastigida Grassi & Foa reported from lower termites (Isoptera families

Masotermitidae, Kalotermitidae, Hodotermitidae, Termopsidae, Rhinotermitidae, and

Serritermitidae) and from the wood-feeding roach *Cryptocercus* (Dictyoptera□:

Cryptocercidae). *Sociobiology* 1979;4:5–119.

68. Fenchel T, Finlay BJ. Endosymbiotic methanogenic bacteria in anaerobic ciliates -

significance for the growth efficiency of the host. *J Protozool* 1991;38:18–22.

69. Yamada K, Kamagata Y, Nakamura K. The effect of endosymbiotic methanogens

on the growth and metabolic profile of the anaerobic free-living ciliate *Trimyema*

compressum. *FEMS Microbiol Lett* 1997;149:129–32.

70. Treitli SC, Hanousková P, Beneš V et al. Hydrogenotrophic methanogenesis is the

key process in the obligately syntrophic consortium of the anaerobic ameba *Pelomyxa*

schiedti. *ISME J* 2023;17:1884–94.

71. Wagener S, Pfennig N. Monoxenic culture of the anaerobic ciliate *Trimyema*

compressum Lackey. *Arch Microbiol* 1987;149:4–11.

72. Finlay BJ, Embley TM, Fenchel T. A new polymorphic methanogen, closely related

to *Methanocorpusculum parvum*, living in stable symbiosis within the anaerobic ciliate

Trimyema sp. *J Gen Microbiol* 1993;139:371–8.

73. Hongoh Y, Sharma VK, Prakash T et al. Genome of an endosymbiont coupling N_2

fixation to cellulolysis within protist cells in termite gut. *Science* 2008;322:1108–9.

74. Ikeda-Ohtsubo W, Strassert JF, Kohler T et al. '*Candidatus* Adiutrix intracellularis',

an endosymbiont of termite gut flagellates, is the first representative of a deep-

branching clade of *Deltaproteobacteria* and a putative homoacetogen. *Environ*

Microbiol 2016;18:2548–64.

75. Strassert JFH, Mikaelyan A, Woyke T et al. Genome analysis of '*Candidatus*

Ancillula trichonymphae', first representative of a deep-branching clade of

Bifidobacteriales, strengthens evidence for convergent evolution in flagellate

endosymbionts. *Environ Microbiol Rep* 2016;8:865–73.

LEGENDS TO FIGURES

Fig. 1. Detection of endosymbiotic methanogens of *Cononympha* protists in the hindgut

of *Coptotermes formosanus*. (A, C) Phase-contrast images of *Cononympha* cells. (B, D)

Epifluorescent images. Endosymbiotic methanogens are visible with their greenish F_{420}

autofluorescence in panel B, whereas no signal was detected in panel D. Amorphous

yellow is the autofluorescence of wood particles. Bars indicate 10 μm. (E) Ratio of

methanogen-containing *Cononympha* cells to total *Cononympha* cells in individual

termite guts across 33 termite colonies. Values indicate methane emission rates ($nmol^{-1}$

g $termite^{-1}$ h^{-1}) averaged between biological replicates. Asterisks attached to termite

colony IDs indicate laboratory-reared termite colonies.

Fig. 2. Network analysis of 16S rRNA gene sequence variants (SVs) of methanogens

obtained from *Cononympha leidyi* (green), *Cononympha koidzumii* (magenta), and gut

wall (blue). Unambiguously aligned 1,312 nucleotide sites were used. The circle size is

equivalent to the number of samples of single *Cononympha* cells or gut wall fragments,

from which the corresponding SVs were detected. Slashes on branches indicate the

number of nucleotide substitutions. The SVs were clustered by termite colonies, and

their colony IDs and *Cononympha* cell IDs are shown (see Tables S1 and S5).

Fig. 3. Transmission electron microscopy of a *Cononympha* cell. (A) Section of a

Cononympha cell. Several black lines are artefacts. (B) Magnified view of the boxed

region in panel (A). Electron-dense *Methanobrevibacter*-like cells were observed within

the host cytoplasm and also on the host cell surface. CM: cytoplasmic membrane of

Cononympha. Bars = 1 µm (A); 500 nm (B).

Fig. 4. Phylogenetic position and characteristic genetic features of '*Ca.*

Methanobrevibacter cononymphae'. Maximum likelihood tree was constructed based on

42 concatenated single-copy genes using the LG+F+R5 amino acid substitution model.

Methanobacterium veterum MK4 and *Methanobacterium bryantii* M.o.H. were used as

outgroups (Table S3). Ultrafast bootstrap values of 85–95% and ≥95% are indicated

with open and closed circles, respectively. Genomes derived from protist-dependent and

protist-independent termite species are highlighted in filled and unfilled diamonds,

respectively. The presence or absence of genes for formate dehydrogenase FdhAB,

formate transporter FdhC, acetyl-CoA decarbonylase/synthase Cdh, cobalamin

biosynthesis Cbi, and oxygen tolerance (catalase KatE, $F_{420}H_2$ oxidase FprA, and

desulfoferredoxin Dfx) are linked to the phylogenetic tree. Existing and absent genes

are indicated in blue and grey, respectively. Pseudogenes are indicated with 'φ'. The

fdhA gene of MAG_NG87 is at a contig end and truncated; therefore, it is impossible to

specify whether it is intact or pseudogenized.

Fig. 5. The predicted metabolic pathway of '*Ca.* Methanobrevibacter cononymphae'.

Genes or pathways characteristically absent in *Mbv. cononymphae* are shown in white.

The gene for formate dehydrogenase subunit alpha (*fdhA*) is pseudogenized.

Table 1. General genome features of '*Ca.* Methanobrevibacter cononymphae' and

related species.

species	'*Ca.* Mbv. cononymphae' CfCl-M3	*Mbv. curvatus* (GCA_0016392 95)	*Mbv. filiformis* (GCA_0016392 65)	*Mbv. cuticularis* (GCA_0016392 85)	NOE (GCA_0033156 55)
Habitat	Intracellular/ gut wall	gut wall	gut wall	gut wall	intracellular
Total length (Mb)	2.71 (2.67[d])	2.41	2.61	2.61	1.91
Contig	244	232	295	169	70
Completeness [a] (%)	98.6	98.2	97.1	99.7	99.3
G+C (%)	27.2	25.7	27.0	26.8	25.3
CDS [b]	2,319 (1,834)	1,883	1,985	1,869	2,175
Coding density (%) [b]	65.8 (59.7)	68.5	68.5	66.2	68.3
rRNA	3	3	3	3	3
tRNA	28	31	30	32	33
Pseudogene [c]	29/174 (148)	11/86	9/83	15/117	95/209 (187)

[a] Estimated using CheckM2 v.1.0.1.

[b] Identified using Prodigal v2.6.3. Manually curated values are shown in parentheses.

[c] Identified using Pgap/Pseudofinder with those within 1,000 bp region from both contig

ends being excluded. The number was manually curated for CfCl-M3 in this study and

previously for NOE [57] and shown in parenthesis.

[d] Total length with duplicated CDSs being excluded.