1 **Syngeneic natural killer cell therapy activates dendritic and T cells in metastatic lungs**

2 **and effectively treat low-burden metastases**

4 **Shih-Wen Huang[1,2,#], Yein-Gei Lai[1,#], Hao-Ting Liao[1,3,#], Chin-Ling Chang[1], Ruo-Yu**
5 **Ma[1], Yung-Hsiang Chen[1], Yae-Huei Liou[1], Zhen-Qi Wu[1], Yu-Chen Wu[4], Ko-Jiunn**
6 **Liu[4], Yen-Tsung Huang[5], Jen-Lung Yang[1], Ming-Shen Dai[6*], and Nan-Shih Liao[1,2*]**
7

[1]Institute of Molecular Biology, Academia Sinica, Taipei, Taiwan

[2]Molecular and Cell Biology, Taiwan International Graduate Program, Academia Sinica and

Graduate Institute of Life Science, National Defense Medical Center, Taipei, Taiwan

[3]Department of Life Sciences, National Central University, Taoyuan, Taiwan

[4]National Institute of Cancer Research, National Health Research Institutes, Miaoli, Taiwan

[5]Institute of Statistical Science, Academia Sinica, Taipei, Taiwan

[6]Department of Hematology-Oncology, Tri-Service General Hospital, Taipei, Taiwan

[#]Co-first authors, *Co-corresponding authors

ABSTRACT

Natural killer (NK) cells can control metastasis through cytotoxicity and IFN-γ production

independently of T cells in experimental metastasis mouse models. The inverse correlation

between NK activity and metastasis incidence supports a critical role for NK cells in human

metastatic surveillance. However, autologous NK cell therapy has shown limited benefit in

treating patients with metastatic solid tumors. Using a spontaneous metastasis mouse model of

MHC-I$^+$ breast cancer, we found that transfer of IL-15/IL-12-conditioned syngeneic NK cells

after primary tumor resection promoted long-term survival of mice with low metastatic burden

and induced a tumor-specific protective T cell response that is essential for the therapeutic

effect. Furthermore, NK cell transfer augments activation of conventional dendritic cells

(cDCs), Foxp3$^-$CD4$^+$ T cells and stem cell-like CD8$^+$ T cells in metastatic lungs, which requires

IFN-γ of the transferred NK cells. These results imply direct interactions between transferred

NK cells and endogenous cDCs to enhance T cell activation. We conducted an investigator-

initiated clinical trial of autologous NK cell therapy in six patients with advanced cancer and

observed that the NK cell therapy was safe and showed signs of effectiveness. These findings

indicate that autologous NK cell therapy is effective in treating established low burden

metastases of MHC-I$^+$ tumor cells by activating the cDC-T cell axis at metastatic sites.

Introduction

Metastasis is the most common cause of cancer death, and effective therapies for established metastases are currently lacking (Ganesh & Massague, 2021). Natural killer (NK) cells play a critical role in metastasis control, as the abundance of functional circulating or tumor-infiltrating NK cells is inversely correlated with metastasis incidence while being positively correlated with favorable prognoses for patients with or at risk of metastasis (Lopez-Soto, Gonzalez, Smyth, & Galluzzi, 2017). NK cells are capable of recognizing tumor cells via various activating and inhibitory NK receptors (NKRs). Tumor cells often down-regulate the expression of ligands for inhibitory NKRs, yet express stress-induced ligands for activating NKRs. Upon encountering tumor cells, the reduced inhibitory NKR engagement and enhanced activating NKR engagement facilitate NK cell activation (Jamieson et al., 2002; Karre, Ljunggren, Piontek, & Kiessling, 2005). NK cells also express receptors for activating cytokines, such as IL-15, IL-12, IL-18 and IL-21, as well as receptors for suppressive factors such as TGF-β and prostaglandin E2 present in the tumor microenvironment (Huntington, Cursons, & Rautela, 2020). The sum of signals transduced through the activating and inhibitory receptors determines NK cell activation. Activated NK cells kill engaged tumor cells via perforin, granzymes, and the death receptor ligands TNF-α, TRAIL and FasL (Smyth et al., 2005). Activated NK cells also produce IFN-γ, which directly inhibits tumor cell growth and induces type 1 T helper cell (T_H1) and cytotoxic CD8$^+$ T cell responses that are critical for effective anti-tumor immunity (Alspach, Lussier, & Schreiber, 2019; Martin-Fontecha et al., 2004). Moreover, intratumoral NK cells produce XCL1 and CCL5 to recruit type 1 conventional dendritic cells (cDC1) that cross-present cell-associated tumor antigen to CD8$^+$ T cells (Bottcher et al., 2018), and also produce Flt3L to increase the abundance of cDCs (Barry et al., 2018). However, the immunosuppressive microenvironment induced by tumor cells

elicits NK cell and T cell dysfunction (Melaiu, Lucarini, Cifaldi, & Fruci, 2019; Thommen &

Schumacher, 2018).

Two anti-metastatic mechanisms of NK cells have been revealed to date, i.e., cytotoxicity and

IFN-γ production. In the metastatic cascade, the epithelial-mesenchymal transition increases

the susceptibility of tumor cells to NK cell cytotoxicity by modulating the expression of ligands

for the activating NKRs NKG2D (Lopez-Soto et al., 2013), E-cadherin and cell adhesion

molecule 1 (Chockley et al., 2018), as well as MHC class I (MHC-I) molecules (Chen et al.,

2015). The IFN-γ produced by NK cells was shown to be essential for natural resistance to lung

metastasis in a B16 melanoma model (Takeda et al., 2011) and for the IL-12-mediated

suppression of liver metastasis in a CT-26 model (Uemura et al., 2010) using *Rag* knockout

mice that lacked T cells. Another study indicated that the anti-metastasis effect of IL-12

produced by Baft3-dependent DCs is mediated through IFN-γ production by NK cells without

the involvement of T cells (Mittal et al., 2017). NK cells also sustain the dormancy of metastatic

tumor cells seeded in distal organs through cytotoxicity and IFN-γ (Correia et al., 2021;

Malladi et al., 2016). The anti-metastatic mechanisms described above do not involve T cells.

Thus, whether T cell immunity plays a role in the anti-metastatic function of NK cells remains

to be elucidated.

Despite the clear anti-tumor function of endogenous NK cells, adoptive transfer of ex vivo-

expanded autologous NK cells has shown limited clinical benefit in treating patients with

metastatic or locally advanced solid tumors, including melanoma, renal cell carcinoma, and

digestive cancer (Parkhurst, Riley, Dudley, & Rosenberg, 2011; Sakamoto et al., 2015). Both

the immunosuppressive microenvironment induced by tumor cells (Melaiu et al., 2019) and

heterogeneity of NK cell subpopulations and their respective functions (Freud, Mundy-Bosse,

Yu, & Caligiuri, 2017) likely contribute to this therapeutic limitation. Given the vast array of

approaches used for ex vivo expansion of NK cells (Fang, Xiao, & Tian, 2017), different

methods may result in NK cells exerting different functions. We hypothesize that autologous

NK cells with anti-tumor activities are effective in treating low burden metastases. In this study,

we used syngeneic IL-15/IL-12-conditioned NK cells to treat established lung metastases after

resection of the primary MHC-I$^+$ EO771 mammary tumor. We found that the NK cell therapy

promoted long-term survival of mice with low metastatic burden in a CD8$^+$ T cell-dependent

manner, and induced tumor-specific protective immune memory. Furthermore, NK cell

transfer augmented the activation of cDC1, cDC2 and T cells in metastatic lungs, which

requires IFN-γ of the transferred NK cells. We also conducted an investigator-driven clinical

trial of autologous NK cell therapy on six patients hosting four cancer types and found that the

therapy is safe and shows signs of efficacy.

Results

Characterization of ex vivo-expanded murine NK cells

To generate NK cells with anti-tumor activity ex vivo, we cultured murine BM cells with IL-

15 and IL-12, since IL-15 is an NK cell growth and survival factor and both cytokines enhance

IFN-γ production and the cytotoxicity of NK cells (Kennedy et al., 2000; Lodolce et al., 1998;

Marcenaro et al., 2005). On average, 86% of the resulting cells were identified as TCRαβ$^-$

TCRγδ$^-$CD19$^-$NK1.1$^+$ that express NKG2D, EOMES and T-bet (Figure 1A), representing

characteristics of NK cells (Gordon et al., 2012; Raulet, 2003). Approximately 90% of the

expanded NK cells were CD27$^+$ with no-to-low levels of CD11b expression (Figure 1A). The

proportion of NK cells expressing MHC-I-interacting NKRs, including activating Ly49D/H

and inhibitory Ly49A/G2/I and NKG2A, was stable among independent cultures (Figure 1B).

In terms of activation status, on average, 87% of the expanded NK cells expressed either

intermediate or high levels of the activating receptor DNAM-1 and 82% of them displayed an

activated B220$^+$CD11c$^+$ phenotype (Figure 1C). Consistently, they expressed the anti-tumor

effectors IFN-γ, granzyme B, and TRAIL (Figure 1D). Co-culturing the NK cells with either

EO771 breast adenocarcinoma cells that express MHC-I molecules and Rae-1 or B16F10

melanoma cells that express none of those molecules in vitro (Figure 1-figure supplement 1)

resulted in dose-dependent tumor cell death (Figure 1E) and up-regulation of IFN-γ production

by the NK cells (Figure 1F). Thus, the NK cells conditioned by means of IL-15 and IL-12

possess anti-tumor activities in vitro.

Figure 1

119
120

Figure 1. Characterization of ex vivo-expanded murine NK cells.

BM cells were cultured with IL-15 and IL-12 as described in Methods. (A) Analysis of NK cells (CD19⁻TCRαβ⁻

TCRγδ⁻NK1.1⁺) among live expanded cells. Plots are representative of 20 independent cultures. (B)-(D)

Expression of MHC-I-interacting activating and inhibitory NKRs, activation markers and anti-tumor effectors by

the expanded NK cells. Representative flow histograms are presented for each molecule, and the compiled data

from independent expansion cultures is indicated by the dot plots. (E) In vitro cytotoxicity of the expanded NK

cells. CFSE-labeled EO771 or B16F10 cells (T, target) were co-cultured with expanded NK cells (E, effector) at

the indicated E:T ratios. Tumor cell viability was analyzed by PI staining using FACS. One representative of three

independent experiments is shown. Data represent mean ± SEM of triplicate wells. Statistical significance was

determined by unpaired two-tailed Student's t-test: *P<0.05, **P<0.01, ***P<0.001 relative to tumor only. (F)

Tumor cells stimulate IFN-γ production by the expanded NK cells. Sorted NK cells were cultured alone (NK cells)

or together with tumor cells (NK cells + EO771 and NK cells + B16F10) at a ratio of 1:1. Levels of intracellular

IFN-γ of NK cells were determined by staining with anti-IFN-γ or isotype control antibody, and analyzed by

FACS. The difference in mean fluorescence intensity (ΔMFI) was calculated by subtracting the MFI value of

isotype control staining from that of anti-IFN-γ antibody staining. Representative plots of two independent

experiments are shown.

Figure supplement 1. Expression of MHC-I molecules and Rae-1 by EO771 and B16F10 cells *in vitro*.

EO771 tumor-resected mice represent a model for established metastases to test NK cell therapy

Cancer recurrence occurs in a significant proportion of patients after primary tumor resection due to dissemination of tumor cells before and/or during surgery. To mimic that scenario, we established a syngeneic orthotopic breast cancer model using the MHC-I$^+$ EO771 cell line that harbors numerous mutations (Y. Yang et al., 2017). At 21 days post-inoculation and consistent with the spontaneous lung metastatic property of the EO771 line (Ewens, Mihich, & Ehrke, 2005), we observed that ~85% of mice with a primary tumor weighing $\geq$ 95 mg had metastatic foci on the lung surface. The number and area of metastatic foci was positively correlated with tumor weight on day 21 (Figure 2A). Resection of the primary tumor and its sentinel lymph node (LN) on day 21 resulted in a long-term survival rate of 18%, with mortality of all remaining mice (Figure 2B). All dead mice exhibited metastases in the lung, of which 71% displayed tumor reappearance at the surgical site (data not shown). Therefore, this tumor resection model represents a model of established metastases caused by tumor cell dissemination before and arising from surgery. To apply adoptive NK cell therapy to this model, we examined if the transferred NK cells reach the lungs of tumor-resected mice. The expanded NK cells express the chemokine receptors CXCR3, CCR5 and CXCR6 (Figure 2C), and the lung of tumor-resected mice express mRNA of the corresponding ligands CXCL9/10/11, CCL3/4/5 and CXCL16 (Figure 2D). Using eGFP$^+$ NK cells, we observed that numbers of transferred cells in the lung peaked at 4 h and then continuously declined, remaining detectable for at least 2 days (Figure 2E). A similar pattern of transferred NK cells was observed in the spleen, albeit with lower cell counts (Figure 2E).

Figure 2. EO771 tumor resection and metastasis model for NK cell treatment.

(A) Positive correlation between primary tumor weight and lung metastasis. Pearson correlations between day 21 tumor weights and the number and total area of metastatic foci on the lung surface are shown. Each dot represents a mouse. (B) Survival of mice who received day 21 tumor resection, sham surgery, or no treatment. Mouse survival was analyzed by means of the Kaplan-Meier estimator and a Log-Rank test: ****P<0.0001. (C) Expanded NK cells express the chemokine receptors CXCR3, CCR5 and CXCR6. Representative flow plots of 10 independent cultures are shown. (D) Expression of CXCR3, CCR5 and CXCR6 ligands in the lung. Specific gene expression was determined by quantitative real-time PCR using lung RNA of day 21 tumor-resected mice collected on day 24 post tumor inoculation and normalized to the expression of cyclophilin a. Each symbol represents a mouse and means ± SEM are shown. (E) Expanded eGFP+ NK cells were transferred into day 21 EO771-resected mice on day 24, and quantified in the single cell suspensions of spleen and lung at the indicated times using FACS. Numbers of eGFP+ NK cells detected in each organ are shown. Each dot represents a mouse (means ± SEM).

Syngeneic NK cell therapy is effective in treating mice with low metastatic burden in a CD8[+]-T-cell-dependent manner

We evaluated the effect of syngeneic NK cell therapy on tumor-resected mice (Figure 3A Schema), and found a negative correlation between survival time and day 21 tumor weight in the NK cell-treated group but not in the control group (Figure 3A). Consequently, we divided the mice into two groups according to their day 21 tumor weight. NK cell-treated mice displayed significantly enhanced overall survival (OS) and an improved long-term survival rate from 33% to 68% compared to control mice in the 95-400 mg tumor group, whereas the same therapy did not improve OS of the > 400 mg tumor group (Figure 3B). These results, together with the result presented in Figure 2A, indicate that the NK cell therapy promotes long-term survival of tumor-resected mice carrying low metastatic burden. We further examined if the surviving mice after NK cell treatment had acquired tumor-specific protection by re-challenging them with either EO771 or B16F10 tumor cells (Figure 3C Schema). We found that 83% of these mice were protected from EO771 re-challenge and survived beyond 180 days, whereas all mice re-challenged with B16F10 died (Figure 3C). Age-matched naïve mice inoculated for the first time with either EO771 or B16F10 as a control all died (Figure 3C). Thus, surviving mice of post-resection NK cell therapy displayed tumor-specific protective immunity.

Since the tumor re-challenge experiment was performed at 11-13 weeks following NK cell treatment (Figure 3C), the observed long-lasting protection is likely mediated by memory T cells. Accordingly, we examined the role of T cells in NK cell therapy by depleting T cell subsets two days before resecting the primary tumor (Figure 3D Schema and Figure 3-figure supplement 1). Treatment with anti-CD8 antibody alone or with anti-CD4 plus anti-CD8 antibodies, but not with anti-CD4 antibody alone, abolished the therapeutic effect of NK cell

therapy (Figure 3D), indicating the essential role of CD8$^+$ T cells. Moreover, mice depleted of

CD8$^+$ cells alone survived better than mice depleted of both CD4$^+$ and CD8$^+$ cells in response

to NK cell therapy (Figure 3D), suggesting a CD8$^+$-T-cell-independent anti-tumor effect of

CD4$^+$ T cells. Next, we examined whether the T cells in surviving mice after NK cell therapy

and tumor re-challenge possess anti-tumor activity. T cells and non-T cells were isolated from

individual NK cell-treated and EO771-rechallenged survivors or from age-matched donors,

and then transferred into naïve recipients in a 1-donor-to-1-recipient manner. The recipient

mice were inoculated with EO771 cells to evaluate the anti-tumor effects of the donor cells,

whereas naïve mice received no donor cells and served as a control for tumor growth (Figure

3E Schema). For the mice receiving cells isolated from the survivors, 5 of 6 T-cell recipients

and 1 of 6 non-T-cell recipients survived beyond 120 days post tumor inoculation (Figure 3E).

In contrast, mice receiving either T or non-T cells isolated from age-matched naïve donors all

died. Mice that received no donor cells in both experiments also all died (Figure 3E). Together,

these results indicate that the post-resection NK cell therapy induces a tumor-specific T cell

response with memory that is essential for its treatment efficacy.

Figure 3. Syngeneic NK cell therapy is effective in treating mice with low metastatic burden in a CD8[+]-T-cell-

dependent manner.

(A) Day 21 tumor weight is inversely correlated with survival duration of NK-cell-treated tumor-resected mice.

After tumor and sentinel LN resection on day 21, mice received either sorted NK cells (NK cell) or PBS (Control)

at the indicated time-points. The correlation between day 21 tumor weight and survival time of mice was evaluated

by Pearson correlation. Each dot represents a mouse. The results have been compiled from 5-8 independent

experiments. The Control group is the same dataset as presented for the Resection group in Figure 2B. (B) NK

cell therapy promotes long-term survival of mice who had light tumors on day 21. The mice described in (A) were

separated into 95-400 mg and >400 mg day 21 tumor weight groups for survival analysis by means of the Kaplan-

Meier estimator and a Log-Rank test: *P<0.05. (C) NK cell therapy induces tumor-specific protection. Surviving

mice after NK cell therapy were rechallenged with EO771 (●, compiled from four independent experiments) or

B16F10 cells (▲, compiled from two independent experiments) at 11-13 weeks post therapy, and then monitored

for survival. Age-matched naïve mice were inoculated with EO771 (○, compiled from four independent

experiments) or B16F10 cells (△, compiled from three independent experiments) as controls. Mouse survival,

using 2000 mm^3 tumor volume as a cut-off, was analyzed by means of the Kaplan-Meier estimator and a Log-

Rank test: ****P<0.001. (D) Depletion of CD8$^+$ cells abolished the efficacy of NK cell therapy. Mice were treated

with indicated antibodies at day 19 post tumor inoculation, and then received resection followed by NK cell

therapy. Mouse survival was analyzed by means of the Kaplan-Meier estimator and a Log-Rank test: ns, not

significant; *P<0.05; ***P<0.001; ****P<0.0001. The data are compiled from 2-3 experiments. (E) T cells

isolated from surviving mice post NK cell therapy and tumor re-challenge confer anti-tumor activity. Post NK

cell therapy and re-challenge survivors were collected from three independent experiments. T cells and non-T

cells isolated from each survivor were transferred separately into naïve recipient mice in a 1-donor-to-1-recipient

manner, whereas other naïve mice that received no cells served as a control (compiled from two independent

experiments). T cells and non-T cells were also isolated from age-matched naïve donors and transferred separately

into naïve recipient mice, whereas other naïve mice that received no cells served as a control (compiled from two

independent experiments). The recipient mice were then inoculated with EO771 cells one day after cell transfer.

Mouse survival, using 2000 mm^3 tumor volume as a cut-off, was analyzed by means of the Kaplan-Meier

estimator and a Log-rank test: ns, not significant; **P<0.01; ***P<0.001.

Figure supplement 1. Effectiveness of in vivo depletion of CD4$^+$ and CD8$^+$ T cells by antibody (related to Figure

3D).

Syngeneic NK cell transfer promotes cDC and T cell activation in metastatic lungs

Conventional DCs directly trigger an antigen-specific T cell response by providing ligands to antigen, costimulatory and cytokine receptors expressed on T cells (Cabeza-Cabrerizo, Cardoso, Minutti, Pereira da Costa, & Reis e Sousa, 2021). A previous in vivo study reported that A20 lymphoma expressing abnormally low levels of MHC I molecules (MHC-I) induces IL-12 production by splenic DCs ($CD11c^+CD19^-F4/80^-$) and $CD8^+$ T cell response (Mocikat et al., 2003). Therefore, we hypothesized that our NK cell therapy would modulate cDCs in metastatic lungs in favor of a T cell response. As our IL-15/IL-12-conditioned NK cells express high levels of mRNA encoding IFN-γ and IL-10 (data not shown), cytokines known to modulate cDC function, we examined whether one transfer of wild type (WT), $Il10^{-/-}$ or $Ifng^{-/-}$ NK cells affected cDCs and T cells in the metastatic lungs. Immune cells were analyzed ~18 hours after NK-cell transfer, by which time the level of transferred NK cells in the lung declined by > 60% relative to that determined at 4-hours (Fig. 2E).

We found that lung cDC1s and cDC2s consist of cells expressing either high or low level of MHC-II (MHC-II^{hi} or MHC-II^{lo}), and the MHC-II^{hi} cDCs are composed of $CCR7^+Lamp3^+$ and $CCR7^-Lamp3^-$ subsets (Figure 4A). The $CCR7^+Lamp3^+$MHC-II^{hi} subset expresses the highest level of APC function molecules, including MHC-I, MHC-II, CCR7, CD86, CD40 and PD-L1 (Figure 4A), resembling the recently identified mregDC (Cheng et al., 2021; de Saint-Vis et al., 1998; Maier et al., 2020). The $CCR7^-Lamp3^-$MHC-II^{hi} subset expresses an intermediate level of MHC-II, CD86 and CD40 among the three subsets, and similar levels of MHC-I, CCR7 and PD-L1 to that of the MHC-II^{lo} subset (Figure 4A). NK-cell transfer did not alter the proportion of total cDC1s or cDC2s in $CD45^+$ cells in the lung tissue (data not show). However, among cDC1s, transfer of WT NK cells significantly increased the proportion of the MHC-II^{hi} subsets, the levels of CD86 and CD40 expressed by all three subsets, and the levels of PD-L1,

MHC-I and MHC-II expressed by the CCR7⁻Lamp3⁻MHC-II^{hi} and MHC-II^{lo} subsets (Figure

4B). These effects require IFN-γ, but not IL-10, of the transferred NK cells, except for the PD-

L1 level in the MHC-II^{lo} subset (Figure 4B). Transfer of WT NK cells exerted many effects on

cDC2s similar to those described above for cDC1s, albeit with several differences (Figure 4B).

The differences include augmenting the MHC-II level in the CCR7⁺Lamp3⁺MHC-II^{hi} subset,

reducing the MHC-II level in the MHC^{lo} subset, and having no effect on the level of CD40 in

the CCR7⁺Lamp3⁺MHC-II^{hi} subset (Figure 4B). IFN-γ of transferred NK cells is essential for

all effects on cDC2s except for the increase of the CCR7⁺Lamp3⁺MHC-II^{hi} subset, the

increased levels of CD40 and MHC-II in the CCR7⁻Lamp3⁻MHC-II^{hi} subset, and the increased

levels of CD86 and MHC-II in the MHC-II^{lo} subset (Figure 4B).

A

B

Figure 4. Syngeneic NK cell transfer modulates APC function of cDC in the metastatic lungs.

The experimental design is the same as depicted in the Fig. 3A schema except that NK cells (WT, *Il10*$^{-/-}$ or *Ifng*$^-$

$^{/-}$) were transferred once on day 24 and lung immune cells were analyzed on day 25. (A) Gating strategies and

characterization of cDC subsets in lung tissue. The dot plots show the gating of CCR7$^+$LAMP3$^+$MHC-II^{hi}, CCR7$^-$

LAMP3$^-$MHC-II^{hi} and MHC-II^{lo} subsets of cDC1s and cDC2s. The histograms show the comparison of indicated

molecules expressed by the three subsets of cDC1s and cDC2s. Negative controls are either fluorescence-minus-

one (FMO) or single stain of another molecule. Data shown are from a representative PBS-treated mouse. (B)

NK-cell transfer affects the activation of lung cDCs. The proportion of the three cDC subsets in cDC1s and cDC2s

are shown. The expression of the indicated APC function molecules was analyzed as relative MFI, based on

normalization against the mean MFI of the PBS group in each independent experiment. Data are compiled from

two independent experiments and presented as mean ± SEM. Each symbol represents one mouse. Statistical

significance was determined by unpaired two-tailed Student's t-test: *P≤0.05, **P≤0.01, ***P≤0.001.

Next, we examined T cells in the lung tissue. Transfer of WT NK cells increased the proportion

of Foxp3$^-$CD4$^+$ and CD8$^+$ T cells without altering that of Foxp3$^+$CD4$^+$ T cells (Figure 5A and

Figure 5-figure supplement 1A). The proportion of activated Foxp3$^-$CD4$^+$ T cells, including

effector (CD62L$^-$CD44$^+$), PD-1$^+$ and Ki67$^+$ subsets, also increased (Figure 5B and Figure 5-

figure supplement 1A). These effects were not observed for mice that received *Ifng*$^{-/-}$ NK cells

(Figure 5B). CD8$^+$ T cells in a tumor microenvironment (TME) are known to display a complex

composition comprising subsets of different activation/differentiation statuses. First, we

categorized lung tissue CD8$^+$ T cells according to PD-1 and Lag3 expression (Figure 5-figure

supplement 1A), because PD-1 expression is induced by TCR stimulation (Sharpe & Pauken,

2018) and Lag3 is only expressed by exhausted or progenitor-exhausted (PEX) CD8$^+$ T cells

(Calagua et al., 2021; Z. Z. Yang et al., 2017). We reasoned that PD-1$^-$Lag3$^-$ CD8$^+$ T cells

(Figure 5-figure supplement 1A) are either naïve or at a very early stage of activation, and

found that transfer of *Il10*$^{-/-}$, but not WT, NK cells increased the proportion of CD62L$^+$CD44$^+$

and CD62L$^-$CD44$^+$ subsets, while the proportion of CD62L$^-$CD44$^+$ and Ki67$^+$ subsets were

lower in mice that received *Ifng*$^{-/-}$ NK cells compared to those proportions in mice that received

WT NK cells (Figure 5C). These results indicate that IL-10 of transferred NK cells inhibits

activation of naïve CD8$^+$ T cells, whereas IFN-γ of transferred NK cells promotes their effector

differentiation and proliferation. We further examined stem cell-like (SCL) CD8$^+$ T cells that

are Tim3$^-$ in PD-1^{lo}Lag3$^-$ CD8$^+$ cells (Figure 5-figure supplement 1A) (Castiglioni et al., 2023),

and detected an increase of total and Ki67$^+$ SCL CD8$^+$ T cells in mice that received WT or *Il10*$^{-/-}$

, but not *Ifng*$^{-/-}$, NK cells (Figure 5D). Transfer of *Il10*$^{-/-}$ NK cells resulted in a higher level of

SCL CD8$^+$ T cells than when WT NK cells were transferred (Figure 5D). In contrast, NK cell

transfer did not alter the proportion of exhausted or PEX CD8$^+$ T cells (Figure 5D and Figure

5-figure supplement 1A). The expression of PD-1, Lag3, Tim3, CD44, CD62L, Ly108 and

GrzB by SCL, exhausted, and PEX CD8$^+$ T cells in our analysis is consistent with the known

characteristics of these cell populations (Figure 5-figure supplement 1B) (Andreatta et al., 2021;

Castiglioni et al., 2023; Martinez-Usatorre et al., 2020). Thus, overall, transfer of the IL-15/IL-

12-conditioned syngeneic NK cells augmented the activation of cDC1s and cDC2s, as well as

the levels and activation of Foxp3$^-$CD4$^+$ T cells, naïve CD8$^+$ T cells and SCL CD8$^+$ T cells in

metastatic lungs. Moreover, IFN-γ of the transferred NK cells is essential for these effects.

327
328

Figure 5. Syngeneic NK cell transfer promotes T cell activation in metastatic lungs.

The experimental design is the same as Figure 4. The proportion of T cell subsets in CD45$^+$ cells in lung tissue were analyzed (Figure 5-figure supplement 1). (A) Foxp3$^+$CD4$^+$, Foxp3$^-$CD4$^+$ and CD8$^+$ T cells. (B) Foxp3$^-$CD4$^+$ T cell subsets. (C) PD1$^-$Lag3$^-$CD8$^+$ subsets. (D) Total and KI67$^+$ SCL, exhausted, and PEX CD8$^+$ T cells. Data are compiled from two independent experiments and are presented as mean ± SEM. Each symbol represents one mouse. Statistical significance was determined by unpaired two-tailed Student's t-test: *P≤0.05, **P≤0.01, ***P≤0.001.

Figure supplement 1. Analysis of T cells in lung tissue (related to Figure 5).

It is known that CCR7 mediates migration of cDC from tumor to its draining LN (dLN) (Riol-

Blanco et al., 2005), in which the migratory cDCs prime tumor-specific naive T cells (Mempel,

Henrickson, & von Andrian, 2004). As transfer of WT NK cells increased the levels of

$CCR7^+LAPM3^+MHC-II^{hi}$ cDC1s and cDC2 in metastatic lungs (Figure 4), we hypothesized

that NK cell transfer augments migration of lung cDCs to mLN and activation of T cells. We

analyzed migratory and resident cDCs in mLN (Figure 6-figure supplement 1), and found that

> 90% of migratory cDC1s and > 65% of migratory cDC2s express CCR7 and LAMP3 (Figure

6A), a phenotype similar to lung $CCR7^+LAMP3^+MHC-II^{hi}$ cDC1 and cDC2. On the other hand,

the great majority of resident cDC1s and cDC2s in mLN are $CCR7^-LAMP3^-$ (Figure 6A).

Transfer of WT NK cells increased the proportion of migratory cDC1 in cDCs and the level of

CD86 expression by both migratory cDC1s and cDC2s, which required IFN-γ of the transferred

NK cells (Figure 6B). We also examined T cells in mLN and found that NK cell transfer

affected neither the proportion nor the activation marked by $CD44^+$ or $KI67^+$ of $Foxp3^-CD4^+$

and $CD8^+$ T cells (data not show). Collectively, NK cell transfer increased the level of

migratory cDC1 and CD86 expression by migratory cDC1s and cDC2s, but did not affect T

cell activation in mLN.

353
354
Figure 6. Syngeneic NK cell transfer affects migratory cDCs in mLN.
The experimental design is the same as Fig. 4. (A) The expression of CCR7 and LAMP3 by migratory and resident
cDC1s and cDC2s. (B) NK cell transfer affected migratory cDC subsets. The proportion of migratory cDC1s and
cDC2s in cDC are shown. The expression of the indicated APC function molecules was analyzed as relative MFI,
based on normalization against the mean MFI of the PBS group in each independent experiment. Data are
compiled from two independent experiments and presented as mean ± SEM. Each symbol represents one mouse.
Statistical significance was determined by unpaired two-tailed Student's t-test: *P≤0.05, **P≤0.01, ***P≤0.001.
Figure supplement 1. Analysis of cDCs in mLN.

Clinical trial of autologous NK cell therapy on advanced cancer patients

To assess NK cell therapy in human, we expanded human NK cells from TCRβ⁻CD19⁻ PBMCs. The expanded NK cells are identifiable as CD3⁻CD19⁻CD14⁻CD56⁺EOMES⁺, with ~90% of the cells expressing HLA-DR (Figure 7A). Co-culture of the expanded HLA-DR⁺ NK cells with K562, THP-1 or U937 tumor cells resulted in dose-dependent tumor cell death (Figure 7B) and increased IFN-γ production by the NK cells (Figure 7C). We then conducted an investigator-initiated clinical trial of autologous NK cell therapy in six pre-treated advanced cancer patients enrolled between May 2016 and October 2019. Baseline characteristics of these patients are presented in Table. The patients had an ECOG performance status of 0 and a relatively small target lesion size (6.4-22.7 mm), suggesting relatively low tumor burden. The patients in cohort 1 and cohort 2 received six bi-weekly infusions of 20×10^6 or 30×10^6 autologous HLA-DR⁺ NK cells, respectively. The infused cells were total cells expanded from TCRβ⁻CD19⁻ PBMCs, of which median 92.4% (83.2-94.9%) were HLA-DR⁺ NK cells (Figure 8A and Figure 8-figure supplement 1). The composition of the infused cells differed slightly among patients, but was stable for each patient over the six infusions (Figure 8A). We examined the expression of NKRs, CD16 and CXCR3 on the HLA-DR⁺ NK cells and determined that it was stable for each patient over six batches of cell preparations, apart from some variation in NKp44 for patient 5 and CXCR3 for patient 4 (Figure 8B and Figure 8-figure supplement 2). Moreover, the HLA-DR⁺ NK cells expressed the anti-tumor effectors perforin and IFN-γ (Figure 8C and Figure 8-figure supplement 3). All patients completed the course of six infusions. The therapy proved safe and was well-tolerated by all patients, as no treatment-related adverse effect was observed throughout the entire infusion period up to 24 h after the final infusion. We assessed the preliminary impact of this therapeutic approach based on clinical responses and OS, and present the results with a median follow-up of 60.0 months (2.73-60.0). The response and survival timings were counted from the first infusion of NK cells

(Figure 8D). We adopted RECIST 1.1 to evaluate responses, except that the baseline target

lesions for patients 3, 4 and 5 (Table) were 18%, 36% and 31% smaller than measurable lesions

according to RECIST 1.1. All patients were in a stable disease (SD) state at month 1 (i.e., 1

week after the second infusion), and completed six infusions by month 2.3. Patients 1, 3 and 6

only received NK cell therapy, whereas patients 2, 4 and 5 received additional treatment(s) as

indicated. Patient 1 exhibited progressive disease (PD) at month 2.7 and left the trial. Patient 2

displayed SD for 8.2 months, during which a partial response (PR) was detected at month 4.4.

She then received maintenance metronomic Endoxan starting from month 8.4, and displayed

SD or PR for a further 50 months. Patient 3 displayed SD for 52.7 months. Patient 4 exhibited

an SD state for 13 months, received Tarceva during months 14.9-18.6, suffered PD at month

18.6 that prompted receipt of other treatments, and ultimately left the trial at month 31.5.

Patient 5 presented SD for 9.9 months, and received metronomic Endoxan during months 7.7-

13.5, 28.9-33.0, and 43.6-46.1. She underwent ultrasound imaging during months 17.6-54.2

and the results support a continuing SD state. Patient 6 retained an SD state for 36.3 months,

and subsequent ultrasound imaging in month 44.5 supports a continuing SD state. In summary,

among the four patients with metastatic solid tumors, patients 1 and 4 exhibited PD at month

2.7 and 18.6, respectively, leaving the trial at months 2.7 and 31.5, respectively. In contrast,

patients 3 and 6, who only received NK cell therapy, exhibited a SD state and both survived

for 60 months. The two follicular lymphoma patients (patients 2 and 5) who started

maintenance Endoxan therapy ~6 months after undergoing the NK-cell therapy have exhibited

a SD state and both survived for 60 months. These preliminary results suggest some level of

effectiveness for the autologous NK cell therapy, supporting the efficacy of our pre-clinical

mouse model.

411
412

Figure 7. Ex vivo-expanded human NK cells exhibit anti-tumor activities in vitro.

(A) Analysis of expanded human NK cells. Live cells among total cells obtained from human PBMC culture (as described in Methods) were analyzed for the expression of indicated molecules by FACS. Representative plots of 33 healthy donors are shown. (B) Expanded human NK cells kill tumor cells in vitro. Sorted HLA-DR$^+$ NK cells from two healthy donors were co-cultured with CFSE-labeled tumor cells at the indicated E:T ratio. Percentage of dead tumor cells represents the mean ± SEM of PI$^+$ cells among CFSE$^+$ tumor cells from triplicate wells. (C) Expanded human NK cells up-regulate IFN-γ production in response to tumor cells in vitro. Sorted HLA-DR$^+$ NK cells from two healthy donors were co-cultured with tumor cells at a 1:1 ratio for 5 h. Levels of IFN-γ in NK cells were examined by intracellular staining and are calculated as the ΔMFI ± SEM between specific antibody-stained and isotype control antibody-stained samples. Statistical significance was determined by unpaired two-tailed Student's t-test: *P<0.05, **P<0.01, ***P<0.001 relative to NK cells only.

Table. Baseline characteristics of the patients enrolled in the phase I trial.

Characteristic	Cohort 1			Cohort 2		
Patient	1	2	3	4	5	6
Age	63	42	59	66	54	65
Gender	Male	Female	Female	Female	Female	Male
Primary cancer	Colon	Follicular lymphoma	Lung	Lung	Follicular lymphoma	Bladder
Metastasis	Adrenal Lung Pelvis		Lung	Brain		Bone
Previous therapies for cancer						
Surgery	Yes	No	Yes	Yes	No	Yes
Radiotherapy	No	No	No	Yes	No	No
Lines of chemo- or targeted therapy	1	3	0	3	2	4
Immunotherapy	No	No	No	No	No	Keytruda[a]
ECOG performance status	0	0	0	0	0	0
Target lesion (mm)	Pelvic LN (16.3)	Left inguinal LN (18.4)	Mediastinal LN (12.3)	Lung left upper lobe (6.4)	Left axillary LN (10.4)	Pelvic LN (22.7)

a: Ended 7 months before NK cell therapy

Figure 8. Clinical outcomes of autologous NK cell therapy.

(A) Composition of the infused cells for each patient over six batches of cell preparation. Expression of (B) NKRs, CD16 and CXCR3 and (C) perforin and IFN-γ by the expanded HLA-DR$^+$ NK cells over six batches of cell preparation. Each dot represents a batch of cell preparation. (D) Clinical responses and survival of the six patients. The blue and yellow blocks represent the durations of NK cell treatment and subsequent follow-up, respectively. The response of patients (SD, PD or PR) was evaluated by CT imaging according to RECIST 1.1. Sono symbol marks follow-up using ultrasound imaging. Colored dotted lines indicate the timeframes for additional medications.

Figure supplement 1. Expression of CD56 and HLA-DR by CD3$^-$CD19$^-$CD14$^-$ cells after *ex vivo* expansion (related to Figure 6A).

Figure supplement 2. Expression of NKRs, CD16 and CXCR3 by the expanded HLA-DR$^+$ NK cells.

Figure supplement 3. Expression of IFN-γ and perforin in the expanded HLA-DR$^+$ NK cells.

Discussion

The limited benefit achieved with adoptive autologous NK cell therapy for metastatic solid tumors is paradoxical given the clear anti-tumor functions of endogenous NK cells. In this study, we found that transfer of syngeneic NK cells conditioned using IL-15 and IL-12 after resecting the primary tumor resulted in long-term survival of mice with low burden lung metastases. Our results indicate a requirement for CD8$^+$ T cells and the contribution of CD4$^+$ T cells to the effectiveness of this NK cell therapy (Figure 3D), which is distinct from the T-cell-independent anti-metastasis mechanisms of NK cells demonstrated previously (Correia et al., 2021; Malladi et al., 2016; Mittal et al., 2017; Takeda et al., 2011; Uemura et al., 2010). The difference in T cell dependency likely arises from differences in experimental design. These previous studies examined the effect of endogenous NK cells on experimental metastasis following intravenous or intrasplenic tumor cell injection by enumerating metastases of affected organs 14 days later. Given that the effect of NK cells on experimental metastasis is mediated by clearing intraluminal tumor cells in microvessels before they extravasate into the target tissue (Spiegel et al., 2016), these studies assessed T cell-independent control of metastasis by NK cells. In contrast, our study examined the effect of adoptively transferred ex vivo-stimulated NK cells on spontaneous metastasis by measuring long-term survival, in which lung metastases were established before primary tumor resection and NK cell administration (Figure 2A). The T cell dependency of the NK cell therapy in our model indicates that innate mechanisms are not sufficient to support the long-term survival of mice with EO771 metastases, which is consistent with the finding from experimental metastasis models that NK cells alone cannot reduce metastases after tumor cells have seeded the target tissue (Spiegel et al., 2016). Importantly, the timeframe we adopted for measuring long-term survival covers the period for the innate effect of the transferred NK cells and the effect of a full T cell response with memory.

The CD8[+] T cell dependency implies that the transferred NK cells act as an upstream controller

for an effective anti-tumor T cell response, which is in line with a previous study showing that

early intratumoral accumulation of IFN-γ-producing endogenous NK cells is required for CD8[+]

T cell-mediated eradication of *Cox*-deficient melanoma in immunocompetent wild-type mice

(Bonavita et al., 2020).

In studying the mechanism underlying T cell-dependency for the effectiveness of NK cell

therapy, our results from a mouse model reveal that transfer of WT NK cells augments

activation of cDC1s and cDC2s in the metastatic lung (Figure 4). cDC1s and cDC2s present

cell-associated and soluble antigens, respectively, to CD4[+] and CD8[+] T cells (Hildner et al.,

2008). The increase of MHC-II[hi] cDC subsets and elevated expression of MHC-I, MHC-II and

CD86 by lung cDC1s and cDC2s presumably promotes T cell activation, which is consistent

with the increase of activated lung Foxp3[-]CD4[+] and CD8[+] T cells (Figure 5). Moreover, the

elevated CD40 expression by cDC1 and CD4[+] T cell activation implies cDC1-relayed CD4[+] T

cell help for CD8[+] T cell response in the lung (Ferris et al., 2020). We also detected an increase

of SCL CD8[+] T cells in metastatic lungs upon NK cell transfer (Figure 5D). Intratumoral

tumor-specific SCL CD8[+] T cells are known to mediate the responses to PD-1/PD-L1 blockade

and adoptive T cell immunotherapies (Krishna et al., 2020; Siddiqui et al., 2019) and are

supplied from a reservoir of precursors in the tumor dLN (Connolly et al., 2021; Schenkel et

al., 2021). As cDC1s in tumor dLN maintain the SCL CD8[+] T cell precursor reservoir

(Schenkel et al., 2021), the increase of migratory cDC1 in mLN upon NK cell transfer (Figure

6B) likely contribute to the increase of lung SCL CD8[+] T cells. Moreover, a recent study

reported that the SCL CD8[+] T cell precursors are activated by tumor antigens in the tumor dLN

and acquire a stem-like state without an effector phenotype (Prokhnevska et al., 2023). After

migrating to a tumor, they differentiate from SCL to an effector state, which requires

costimulation by CD80 and CD86 of APC in the TME (Prokhnevska et al., 2023). Our findings

of augmented CD86 expression by cDCs and increased proliferating SCL CD8$^+$ T cells in

response to NK-cell transfer in metastatic lung are in line with the two-step activation of

intratumoral SCL CD8$^+$ T cells. Together, these results demonstrate a clear association between

the activation of lung cDCs and T cells in response to NK cell transfer. Furthermore, IFN-γ of

transferred NK cells is essential for the activation of both cDCs and T cells, strengthening

evidence for the scenario that NK cells activate cDCs and then cDCs activate T cells. The

requirement for IFN-γ of transferred NK cells also suggests a direct interaction between

transferred NK cells and lung cDCs. The increase of CCR7$^+$LAMP3$^+$ MHC-II^{hi} cDCs supports

this supposition, with a recent study reporting colocalization of LAMP3$^+$ DCs with NK cells

in four types of human solid tumor (Tang et al., 2023). Moreover, this result indicates that NK-

cell-derived IFN-γ drives the generation of mregDCs.

In summary, our results indicate that syngeneic IL-15/IL-12-conditioned NK cell therapy

promotes tumor-specific T cell responses, at least in part, through activation of cDC1s and

cDC2s in metastatic lung. Our findings are consistent with the critical role of endogenous NK

cells and CD8$^+$ T cells in anti-tumor immunity, and support that autologous NK cell therapy is

effective in treating established low burden MHC-I$^+$ metastases by enhancing the tumor-

specific CD8$^+$ T cell response.

Materials and Methods

Mice and cell lines

Female C57BL/6JNarl mice (National Laboratory Animal Center of Taiwan, Taipei, Taiwan),

eGFP transgenic C57BL/6JNarl mice (a gift from Chin-Yen Tsai at Academia Sinica (AS)),

Ifng[-/-] mice (strain # 002287, The Jackson Laboratory, Bar Harbor, ME, USA) and *Il10*[-/-] mice

(Strain # 002251, The Jackson Laboratory) were housed in a specific pathogen-free animal

facility at the Institute of Molecular Biology. Mice aged 8-12 weeks old were used for

experiments, unless stated otherwise. All animal protocols were approved by the IACUC of

AS. EO771 (CH3 Biosystems), K562 (a gift from Che-Kun James Shen at AS), THP-1 and

U937 (gifts from Li-Chung Hsu at National Taiwan University) cells were cultured in RPMI-

1640 (Gibco, Grand Island, NY, USA) containing 10% FBS (Hyclone, Marlborough, MA,

USA), 20 mM HEPES (Sigma-Aldrich, St. Louis, MO, USA), 100 U/ml penicillin and 100

μg/ml streptomycin (Gibco). B16F10 melanoma (a gift from Steve Roffler at AS) was cultured

in DMEM (Gibco) containing 10% FBS and 100 U/ml penicillin and 100 μg/ml streptomycin.

Flow cytometry

To stain cell surface molecules, cells were incubated with fluorophore-conjugated antibodies

for 15 min at 4 °C and washed twice with staining buffer (PBS containing 2% FBS, 5 mM

EDTA and 0.01% NaN_3). For staining of intracellular molecules, cells were fixed for 30 min

at 4 °C after surface staining with 4% paraformaldehyde, or with a Foxp3/Transcription Factor

Fixation/Permeabilization Concentrate and Diluent kit (Thermo Fisher Scientific, Waltham,

MA, USA) to stain transcription factors. The fixed cells were then washed with staining buffer,

permeabilized with 0.1% saponin, and stained with antibody for 30 min at 4 °C. Cells were

analyzed by LSRII, FACSymphony A3 or FACSCalibur (BD Biosciences, Franklin Lakes, NJ,

USA) and the data were analyzed using FlowJo (BD Biosciences). The antibodies used are

detailed in the supplementary materials.

Ex vivo expansion of murine and human NK cells

Murine bone marrow (BM) cells were depleted of red blood cells (RBCs) by means of ammonium-chloride-potassium (ACK) buffer (150 mM NH_4Cl, 10 mM $NaHCO_3$, 1 mM EDTA) and then cultured in RPMI-1640 containing 10% FBS (Corning, Corning, NY, USA), 20 mM HEPES (Sigma-Aldrich), 200 U/ml penicillin, 200 μg/ml streptomycin (Gibco), 50 μg/ml gentamycin (Sigma-Aldrich), 0.2 mg/ml L-glutamine (Sigma-Aldrich), and 50 μM 2-mercaptoethanol (Merck, Rahway, NJ, USA) in a 5% CO_2 incubator at 37 °C for 7 days with 30 ng/ml IL-15 (BioLegend, San Diego, CA, USA) and 10 ng/ml IL-12 (Peprotech, Cranbury, NJ, USA). The expanded murine NK cells ($CD19^-TCR\beta^-NK1.1^+CD11c^+B220^+$) were sorted using SORP (BD Biosciences). Human NK cells were expanded from peripheral blood mononuclear cells (PBMCs) after depleting $TCR\beta^+$ and $CD19^+$ cells with anti-TCRβ (WT31, BD Biosciences) and anti-CD19 (4G7, BD Biosciences) antibodies, and anti-mouse IgG1 microbeads (Miltenyi Biotec, Bergisch Gladbach, Germany) using an LD column with a QuadroMACS Separator (Miltenyi Biotec) under good tissue practice (GTP) conditions. The resulting $TCR\beta^-CD19^-$ PBMCs were cultured in RPMI-1640 containing autologous plasma, 20 mM HEPES (Sigma-Aldrich), 2 mM L-glutamine (Cellgro, Lincoln, NE, USA), and 50 μg/ml gentamycin (Winston Medical Supply Co. Ltd.) in a 5% CO_2 incubator at 37 °C for 7 days with 25 ng/ml IL-15 (CellGenix, Freiburg, Germany) and 10 ng/ml IL-12 (BioLegend) (Lee & Liao, 2015).

In vitro anti-tumor activity of expanded NK cells

Tumor cells were labeled using a Vybrant CFDA SE (CFSE) cell tracer kit (Invitrogen, Waltham, MA, USA). For cytotoxicity assay, sorted NK cells were co-cultured with 10^5 labeled tumor cells at the indicated ratio and then incubated for 5 h in a 5% CO_2 incubator at 37 °C. The cells were then stained with propidium iodide (PI) and the $CFSE^+$ tumor cells were

analyzed for PI⁻ cells using a LSRII system. For IFN-γ production, sorted NK cells were co-

cultured with CFSE-labeled tumor cells at a 1-to-1 ratio, with 10 μg/ml brefeldin A being

present for the last 4 h of the co-culture. Cells were then stained intracellularly with anti-IFN-

γ or isotype control antibody to detect IFN-γ production by CFSE⁻ NK cells using a LSRII

system.

Trafficking of ex vivo-expanded murine NK cells

NK cells expanded from the BM cells of *eGFP* transgenic mice were sorted and intravenously

transferred into C57BL/6JNarl mice (6 million cells/mouse) 3 days after resection of a day-21

tumor. Single cell suspensions were prepared from the lung and spleen at the indicated hours

after NK cell transfer and analyzed for eGFP⁺ donor cells using a LSRII system.

Gene expression

Lung RNA was extracted using Trizol (Invitrogen) and reverse-transcribed into cDNA using a

Reverse Transcription Kit (SMOBIO, Paramount, California, USA). Relative gene expression

was determined using a QuantStudio 12K Flex Real-Time PCR system (Thermo Fisher

Scientific) with SYBR Green and relative standard curves normalized against expression of

cyclophilin a (*Cypa*). The sequences of primers used are detailed in the supplementary

materials.

EO771 resection and metastasis model and NK cell treatment

Each mouse was inoculated with 0.3 million EO771 cells into the right fourth mammary fat

pad on day 0. Twenty-one days later, mice were anesthetized and the resulting tumor and

sentinel LN (right inguinal LN) were resected. Sham control mice received the same surgical

procedure without removing any tissue. Mice harboring a day-21 tumor weight of <95 mg were

excluded since resection alone promoted long-term survival of mice with such tumors. Mice

were grouped to have a similar mean and standard deviation of day-21 tumor weight. Each

mouse then received 0.3 million expanded NK cells or PBS (Control) via the tail vein three

times on days 24, 28 and 31, or once on day 24 as indicated. The resulting metastatic foci on

lung surfaces were visualized by staining the dissected lung with India ink under a dissecting

microscope. The area of metastatic foci was determined using ImageJ (NIH). For T cell

depletion, mice were administered intraperitoneally with 200 µg of anti-CD4 (GK1.5, BioXcell,

Lebanon, NH, USA), anti-CD8α (2.43, BioXcell), or both antibodies 19 days after EO771

inoculation, whereas control mice each received 400 µg of isotype control antibody (Rat IgG2b,

κ; BioXcell). The mice underwent tumor resection 2 days later, before receiving NK cells as

described in the previous section.

Isolation of murine T cells and non-T cells

Single cell suspensions were prepared from the BM, spleen, and LNs of NK cell-treated and

tumor-rechallenged long-term survivors and from the same tissues of age-matched naïve mice.

After lysis of RBCs with ACK buffer, T cells and non-T cells were isolated using Pan T cell

isolation Kit II (Miltenyi Biotec) via an autoMACS Pro Separator (Miltenyi Biotec) according

to the manufacturer's instructions. The T cell preparations comprised 88-93% of $CD3^+$ cells,

and the non-T cell preparations comprised 90-96% of $CD3^-$ cells.

Preparation of lung cell suspension

Each mouse received an intravenous injection of 3 µg of FITC-anti-CD45 antibody three

minutes before euthanasia to label blood $CD45^+$ cells. Lungs were harvested, minced and

washed with PBS to remove excess anti-CD45 antibody. The tissue pieces were digested with

0.2 µg/ml of collagenase IV in HBSS (Sigma H1641) containing 2% FBS and 10mM HEPES

for 30 min at 37°C with shaking at 200 rpm. RBCs in the lung cell suspension were lysed using

ACK lysis buffer.

Clinical trial design

We conducted an investigator-driven, open-label, 3+3 design trial to investigate the safety of

the autologous NK cell therapy on metastatic/refractory stage IV cancer patients at a single

center (ClinicalTrials.gov Identifier: NCT02661685). All six patients were ineligible for or

refused further systemic chemotherapy or immunotherapy, and had an Eastern Cooperative

Oncology Group (ECOG) performance status of 0, a target lesion, and adequate organ function

at the time of enrollment. The three patients in cohort 1 received six bi-weekly intravenous

infusions of 20×10^6 expanded autologous HLA-DR$^+$ NK cells, whereas the three patients in

cohort 2 were administered with 30×10^6 of the same cells. All patients underwent a

computerized tomography (CT) scan between the second and the third NK cell infusions, and

then five further CT scans at 8-week intervals starting one week after the sixth infusion.

Thereafter, CT or sonography scans were performed as indicated. Safety was evaluated by

assessing dose-limiting toxicity (DLT), defined as any treatment-related toxicity of grade 3 or

above of general or immune disorders according to the National Cancer Institute-Common

Terminology Criteria for Adverse Events (NCI-CTCAE) v4.03. Fever, chills, flu-like

symptoms, or infusion-related reactions of grade 3 or above were considered DLT only if they

remained at grade 3 or above for more than 3 days despite appropriate medication. Clinical

responses to the therapy were evaluated by CT scans according to Response Evaluation Criteria

in Solid Tumors version 1.1 (RECIST 1.1). Patient survival was followed every 8 weeks for

the first year and every 12 weeks from the second to the fifth year.

Trial oversight

The protocols for our clinical trial and follow-up were approved by the Institutional Review

Boards (IRBs) at Tri-Service General Hospital and at AS, and were conducted according to the

principles of the Declaration of Helsinki. Written informed consent was obtained from all

patients.

Statistical analysis

Results are presented as mean ± SEM. Statistical significance of in vitro cytotoxicity was

examined by unpaired two-tailed Student's t-test. Correlation was determined by Pearson

correlation coefficient. The Kaplan-Meier estimator was employed for survival analysis, and

statistical significance was determined by a Log-Rank test. Tumor volume was calculated using

the formula: length x width^2 x 0.52. Mice with a tumor volume exceeding 2000 mm^3 were

considered moribund. All statistical analyses were performed using GraphPad Prism 7

(GraphPad). ns, not significant; *P<0.05; **P<0.01; ***P<0.001; ****P<0.0001.

Acknowledgments

We thank the FACS Core and Animal Facilities at the Institute of Molecular Biology, AS, for

service and support. We thank Dr. Chee-Jen Chang at the Graduate Institute of Clinical Medical

Science, Chang Gung University, for consultation on statistical design of the clinical trial. We

thank Ms. Yi-Min Liu and colleagues at the Translation Resource Center for Genomic

Medicine, Institute of Biomedical Science, AS, for assistance with designing the case report

form, the Oracle clinical system and IRB affairs. We thank the Instrument Center at the

National Defense Medical Center for flow cytometry services. We thank Dr. John O'Brien for

manuscript editing.

References

Alspach, E., Lussier, D. M., & Schreiber, R. D. (2019). Interferon gamma and Its Important

Roles in Promoting and Inhibiting Spontaneous and Therapeutic Cancer Immunity.

Cold Spring Harb Perspect Biol, 11(3). doi:10.1101/cshperspect.a028480

Andreatta, M., Corria-Osorio, J., Muller, S., Cubas, R., Coukos, G., & Carmona, S. J. (2021).

Interpretation of T cell states from single-cell transcriptomics data using reference

atlases. *Nat Commun, 12*(1), 2965. doi:10.1038/s41467-021-23324-4

Barry, K. C., Hsu, J., Broz, M. L., Cueto, F. J., Binnewies, M., Combes, A. J., . . . Krummel,

M. F. (2018). A natural killer-dendritic cell axis defines checkpoint therapy-responsive

tumor microenvironments. *Nat Med, 24*(8), 1178-1191. doi:10.1038/s41591-018-0085-

8

Bonavita, E., Bromley, C. P., Jonsson, G., Pelly, V. S., Sahoo, S., Walwyn-Brown, K., . . .

Zelenay, S. (2020). Antagonistic inflammatory phenotypes dictate tumor fate and

response to immune checkpoint blockade. *Immunity, 53*(6), 1215-1229 e1218.

doi:10.1016/j.immuni.2020.10.020

Bottcher, J. P., Bonavita, E., Chakravarty, P., Blees, H., Cabeza-Cabrerizo, M., Sammicheli,

S., . . . Reis e Sousa, C. (2018). NK Cells Stimulate Recruitment of cDC1 into the

Tumor Microenvironment Promoting Cancer Immune Control. *Cell, 172*(5), 1022-

1037.e1014. doi:10.1016/j.cell.2018.01.004

Cabeza-Cabrerizo, M., Cardoso, A., Minutti, C. M., Pereira da Costa, M., & Reis e Sousa, C.

(2021). Dendritic Cells Revisited. *Annu Rev Immunol, 39*, 131-166.

doi:10.1146/annurev-immunol-061020-053707

Calagua, C., Ficial, M., Jansen, C. S., Hirz, T., Del Balzo, L., Wilkinson, S., . . . Einstein, D. J.

(2021). A Subset of Localized Prostate Cancer Displays an Immunogenic Phenotype

Associated with Losses of Key Tumor Suppressor Genes. *Clin Cancer Res, 27*(17),

4836-4847. doi:10.1158/1078-0432.CCR-21-0121

Castiglioni, A., Yang, Y., Williams, K., Gogineni, A., Lane, R. S., Wang, A. W., . . . Muller,

S. (2023). Combined PD-L1/TGFbeta blockade allows expansion and differentiation of

stem cell-like CD8 T cells in immune excluded tumors. *Nat Commun, 14*(1), 4703.

doi:10.1038/s41467-023-40398-4

Chen, X. H., Liu, Z. C., Zhang, G., Wei, W., Wang, X. X., Wang, H., . . . Du, J. (2015). TGF-

beta and EGF induced HLA-I downregulation is associated with epithelial-

mesenchymal transition (EMT) through upregulation of snail in prostate cancer cells.

Mol Immunol, 65(1), 34-42. doi:10.1016/j.molimm.2014.12.017

Cheng, S., Li, Z., Gao, R., Xing, B., Gao, Y., Yang, Y., . . . Zhang, Z. (2021). A pan-cancer

single-cell transcriptional atlas of tumor infiltrating myeloid cells. *Cell, 184*(3), 792-

809 e723. doi:10.1016/j.cell.2021.01.010

Chockley, P. J., Chen, J., Chen, G., Beer, D. G., Standiford, T. J., & Keshamouni, V. G. (2018).

Epithelial-mesenchymal transition leads to NK cell-mediated metastasis-specific

immunosurveillance in lung cancer. *J Clin Invest, 128*(4), 1384-1396.

doi:10.1172/JCI97611

Connolly, K. A., Kuchroo, M., Venkat, A., Khatun, A., Wang, J. W., William, I., . . . Joshi, N.

S. (2021). A reservoir of stem-like CD8 T cells in the tumor-draining lymph node

preserves the ongoing antitumor immune response. *Science Immunology, 6*(64).

doi:https://doi.org/10.1126/sciimmunol.abg7836

Correia, A. L., Guimaraes, J. C., Auf der Maur, P., De Silva, D., Trefny, M. P., Okamoto, R., . . .

Bentires-Alj, M. (2021). Hepatic stellate cells suppress NK cell-sustained breast cancer

dormancy. *Nature, 594*(7864), 566-571. doi:10.1038/s41586-021-03614-z

de Saint-Vis, B., Vincent, J., Vandenabeele, S., Vanbervliet, B., Pin, J. J., Ait-Yahia, S., . . .

Lebecque, S. (1998). A novel lysosome-associated membrane glycoprotein, DC-LAMP,

induced upon DC maturation, is transiently expressed in MHC class II compartment.

Immunity, 9(3), 325-336. doi:10.1016/s1074-7613(00)80615-9

Ewens, A., Mihich, E., & Ehrke, M. J. (2005). Distant metastasis from subcutaneously grown

E0771 medullary breast adenocarcinoma. *Anticancer Res, 25*(6b), 3905-3915.

Retrieved from http://ar.iiarjournals.org/content/25/6B/3905.full.pdf

Fang, F., Xiao, W., & Tian, Z. (2017). NK cell-based immunotherapy for cancer. *Semin*

Immunol, 31, 37-54. doi:10.1016/j.smim.2017.07.009

Ferris, S. T., Durai, V., Wu, R., Theisen, D. J., Ward, J. P., Bern, M. D., . . . Murphy, K. M.

(2020). cDC1 prime and are licensed by CD4(+) T cells to induce anti-tumour immunity.

Nature, 584(7822), 624-629. doi:10.1038/s41586-020-2611-3

Freud, A. G., Mundy-Bosse, B. L., Yu, J., & Caligiuri, M. A. (2017). The Broad Spectrum of

Human Natural Killer Cell Diversity. *Immunity, 47*(5), 820-833.

doi:10.1016/j.immuni.2017.10.008

Ganesh, K., & Massague, J. (2021). Targeting metastatic cancer. *Nat Med, 27*(1), 34-44.

doi:10.1038/s41591-020-01195-4

Gordon, S. M., Chaix, J., Rupp, L. J., Wu, J., Madera, S., Sun, J. C., . . . Reiner, S. L. (2012).

The transcription factors T-bet and Eomes control key checkpoints of natural killer cell

maturation. *Immunity, 36*(1), 55-67. doi:10.1016/j.immuni.2011.11.016

Hildner, K., Edelson, B. T., Purtha, W. E., Diamond, M., Matsushita, H., Kohyama, M., . . .

Murphy, K. M. (2008). Batf3 deficiency reveals a critical role for CD8alpha+ dendritic

cells in cytotoxic T cell immunity. *Science, 322*(5904), 1097-1100.

doi:10.1126/science.1164206

Huntington, N. D., Cursons, J., & Rautela, J. (2020). The cancer-natural killer cell immunity

cycle. *Nat Rev Cancer, 20*(8), 437-454. doi:10.1038/s41568-020-0272-z

Jamieson, A. M., Diefenbach, A., McMahon, C. W., Xiong, N., Carlyle, J. R., & Raulet, D. H.

(2002). The role of the NKG2D immunoreceptor in immune cell activation and natural

killing. *Immunity, 17*(1), 19-29. doi:10.1016/s1074-7613(02)00333-3

Karre, K., Ljunggren, H. G., Piontek, G., & Kiessling, R. (2005). Selective rejection of H-2-

deficient lymphoma variants suggests alternative immune defence strategy. 1986. *J

Immunol,   174*(11),   6566-6569.   Retrieved   from

https://www.ncbi.nlm.nih.gov/pubmed/15905492

Kennedy, M. K., Glaccum, M., Brown, S. N., Butz, E. A., Viney, J. L., Embers, M., . . . Peschon,

J. J. (2000). Reversible defects in natural killer and memory CD8 T cell lineages in

interleukin 15-deficient mice. *J Exp Med, 191*(5), 771-780. doi:10.1084/jem.191.5.771

Krishna, S., Lowery, F. J., Copeland, A. R., Bahadiroglu, E., Mukherjee, R., Jia, L., . . .

Rosenberg, S. A. (2020). Stem-like CD8 T cells mediate response of adoptive cell

immunotherapy against human cancer. *Science, 370*(6522), 1328-1334.

doi:10.1126/science.abb9847

Lee, J. M., & Liao, N. S. (2015). United States Patent No. 9157068 B2.

Lodolce, J. P., Boone, D. L., Chai, S., Swain, R. E., Dassopoulos, T., Trettin, S., & Ma, A.

(1998). IL-15 receptor maintains lymphoid homeostasis by supporting lymphocyte

homing   and   proliferation.   *Immunity,   9*(5),   669-676.   doi:10.1016/s1074-

7613(00)80664-0

Lopez-Soto, A., Gonzalez, S., Smyth, M. J., & Galluzzi, L. (2017). Control of Metastasis by

NK Cells. *Cancer Cell, 32*(2), 135-154. doi:10.1016/j.ccell.2017.06.009

Lopez-Soto, A., Huergo-Zapico, L., Galvan, J. A., Rodrigo, L., de Herreros, A. G., Astudillo,

A., & Gonzalez, S. (2013). Epithelial-mesenchymal transition induces an antitumor

immune response mediated by NKG2D receptor. *J Immunol, 190*(8), 4408-4419. doi:10.4049/jimmunol.1202950

Maier, B., Leader, A. M., Chen, S. T., Tung, N., Chang, C., LeBerichel, J., . . . Merad, M. (2020). A conserved dendritic-cell regulatory program limits antitumour immunity. *Nature, 580*(7802), 257-262. doi:10.1038/s41586-020-2134-y

Malladi, S., Macalinao, D. G., Jin, X., He, L., Basnet, H., Zou, Y., . . . Massague, J. (2016). Metastatic Latency and Immune Evasion through Autocrine Inhibition of WNT. *Cell, 165*(1), 45-60. doi:10.1016/j.cell.2016.02.025

Marcenaro, E., Della Chiesa, M., Bellora, F., Parolini, S., Millo, R., Moretta, L., & Moretta, A. (2005). IL-12 or IL-4 prime human NK cells to mediate functionally divergent interactions with dendritic cells or tumors. *J Immunol, 174*(7), 3992-3998. doi:10.4049/jimmunol.174.7.3992

Martin-Fontecha, A., Thomsen, L. L., Brett, S., Gerard, C., Lipp, M., Lanzavecchia, A., & Sallusto, F. (2004). Induced recruitment of NK cells to lymph nodes provides IFN-gamma for T(H)1 priming. *Nat Immunol, 5*(12), 1260-1265. doi:10.1038/ni1138

Martinez-Usatorre, A., Carmona, S. J., Godfroid, C., Yacoub Maroun, C., Labiano, S., & Romero, P. (2020). Enhanced Phenotype Definition for Precision Isolation of Precursor Exhausted Tumor-Infiltrating CD8 T Cells. *Front Immunol, 11*, 340. doi:10.3389/fimmu.2020.00340

Melaiu, O., Lucarini, V., Cifaldi, L., & Fruci, D. (2019). Influence of the Tumor Microenvironment on NK Cell Function in Solid Tumors. *Front Immunol, 10*, 3038. doi:10.3389/fimmu.2019.03038

Mempel, T. R., Henrickson, S. E., & von Andrian, U. H. (2004). T-cell priming by dendritic cells in lymph nodes occurs in three distinct phases. *Nature, 427*(6970), 154-159. doi:10.1038/nature02238

Mittal, D., Vijayan, D., Putz, E. M., Aguilera, A. R., Markey, K. A., Straube, J., . . . Smyth, M.

J. (2017). Interleukin-12 from CD103(+) Batf3-dependent dendritic cells required for

NK-Cell suppression of metastasis. *Cancer Immunol Res, 5*(12), 1098-1108.

doi:10.1158/2326-6066.CIR-17-0341

Mocikat, R., Braumuller, H., Gumy, A., Egeter, O., Ziegler, H., Reusch, U., . . . Rocken, M.

(2003). Natural killer cells activated by MHC class I(low) targets prime dendritic cells

to induce protective CD8 T cell responses. *Immunity, 19*(4), 561-569. Retrieved from

https://ac.els-cdn.com/S1074761303002644/1-s2.0-S1074761303002644-

main.pdf?_tid=1a97a508-b220-482d-b58b-

ff113e2c772d&acdnat=1547709680_420dfea0f64bd7bf375b970454538332

Parkhurst, M. R., Riley, J. P., Dudley, M. E., & Rosenberg, S. A. (2011). Adoptive transfer of

autologous natural killer cells leads to high levels of circulating natural killer cells but

does not mediate tumor regression. *Clin Cancer Res, 17*(19), 6287-6297.

doi:https://doi.org/10.1158/1078-0432.CCR-11-1347

Prokhnevska, N., Cardenas, M. A., Valanparambil, R. M., Sobierajska, E., Barwick, B. G.,

Jansen, C., . . . Kissick, H. (2023). CD8+T cell activation in cancer comprises an initial

activation phase in lymph nodes followed by effector differentiation within the tumor.

Immunity, 56(1), 107-124.e105. doi:10.1016/j.immuni.2022.12.002

Raulet, D. H. (2003). Roles of the NKG2D immunoreceptor and its ligands. *Nat Rev Immunol,*

3(10), 781-790. doi:10.1038/nri1199

Riol-Blanco, L., Sánchez-Sánchez, N., Torres, A., Tejedor, A., Narumiya, S., Corbí, A. L., . . .

Rodríguez-Fernández, J. L. (2005). The chemokine receptor CCR7 activates in

dendritic cells two signaling modules that independently regulate chemotaxis and

migratory speed. *Journal of Immunology, 174*(7), 4070-4080. doi:DOI

10.4049/jimmunol.174.7.4070

Sakamoto, N., Ishikawa, T., Kokura, S., Okayama, T., Oka, K., Ideno, M., . . . Yoshikawa, T.

(2015). Phase I clinical trial of autologous NK cell therapy using novel expansion

method in patients with advanced digestive cancer. *J Transl Med, 13*, 277.

doi:10.1186/s12967-015-0632-8

Schenkel, J. M., Herbst, R. H., Canner, D., Li, A., Hillman, M., Shanahan, S. L., . . . Jacks, T.

(2021). Conventional type I dendritic cells maintain a reservoir of proliferative tumor-

antigen specific TCF-1 CD8 T cells in tumor-draining lymph nodes. *Immunity, 54*(10),

2338-2353.e2336. doi:10.1016/j.immuni.2021.08.026

Sharpe, A. H., & Pauken, K. E. (2018). The diverse functions of the PD1 inhibitory pathway.

Nat Rev Immunol, 18(3), 153-167. doi:10.1038/nri.2017.108

Siddiqui, I., Schaeuble, K., Chennupati, V., Marraco, S. A. F., Calderon-Copete, S., Ferreira,

D. P., . . . Held, W. (2019). Intratumoral Tcf1 PD-1 CD8 T Cells with Stem-like

Properties Promote Tumor Control in Response to Vaccination and Checkpoint

Blockade   Immunotherapy.   *Immunity,*   *50*(1),   195-221.e110.

doi:10.1016/j.immuni.2018.12.021

Smyth, M. J., Cretney, E., Kelly, J. M., Westwood, J. A., Street, S. E., Yagita, H., . . . Hayakawa,

Y. (2005). Activation of NK cell cytotoxicity. *Mol Immunol, 42*(4), 501-510.

doi:10.1016/j.molimm.2004.07.034

Spiegel, A., Brooks, M. W., Houshyar, S., Reinhardt, F., Ardolino, M., Fessler, E., . . .

Weinberg, R. A. (2016). Neutrophils suppress intraluminal NK Cell-mediated tumor

cell clearance and enhance extravasation of disseminated carcinoma cells. *Cancer

Discov, 6*(6), 630-649. doi:10.1158/2159-8290.CD-15-1157

Takeda, K., Nakayama, M., Sakaki, M., Hayakawa, Y., Imawari, M., Ogasawara, K., . . . Smyth,

M. J. (2011). IFN-gamma production by lung NK cells is critical for the natural

835 resistance to pulmonary metastasis of B16 melanoma in mice. *J Leukoc Biol, 90*(4),

836  777-785. doi:10.1189/jlb.0411208

837 Tang, F., Li, J. H., Qi, L., Liu, D. F., Bo, Y. F., Qin, S. S., . . . Zhang, Z. M. (2023). A pan-

838  cancer single-cell panorama of human natural killer cells. *Cell, 186*(19), 4235-

839  4251.e4220. doi:10.1016/j.cell.2023.07.034

840 Thommen, D. S., & Schumacher, T. N. (2018). T Cell Dysfunction in Cancer. *Cancer Cell,*

841  *33*(4), 547-562. doi:10.1016/j.ccell.2018.03.012

842 Uemura, A., Takehara, T., Miyagi, T., Suzuki, T., Tatsumi, T., Ohkawa, K., . . . Hayashi, N.

843  (2010). Natural killer cell is a major producer of interferon gamma that is critical for

844  the IL-12-induced anti-tumor effect in mice. *Cancer Immunol Immunother, 59*(3), 453-

845  463. doi:10.1007/s00262-009-0764-x

846 Yang, Y., Yang, H. H., Hu, Y., Watson, P. H., Liu, H. Y., Geiger, T. R., . . . Wakefield, L. M.

847  (2017). Immunocompetent mouse allograft models for development of therapies to

848  target breast cancer metastasis. *Oncotarget, 8*(19), 30621-30643.

849  doi:https://doi.org/10.18632/oncotarget.15695

850 Yang, Z. Z., Kim, H. J., Villasboas, J. C., Chen, Y. P., Price-Troska, T., Jalali, S., . . . Ansell,

851  S. M. (2017). Expression of LAG-3 defines exhaustion of intratumoral PD-1(+) T cells

852  and correlates with poor outcome in follicular lymphoma. *Oncotarget, 8*(37), 61425-

853  61439. doi:10.18632/oncotarget.18251