Ezrin defines TSC1 activation at endosomal compartments through EGFR-AKT signaling

Giuliana Giamundo[1#], Daniela Intartaglia[1#], Eugenio Del Prete[2], Elena Polishchuk[2], Fabrizio

Andreone[2], Marzia Ognibene[3], Sara Buonocore[1], Francesco Giuseppe Salierno[2], Jlenia Monfregola[2],

Dario Antonini[1], Paolo Grumati[2,4], Alessandra Eva[5], Rossella de Cegli[2], Ivan Conte[1]*

1. Department of Biology, University of Naples Federico II, Strada Vicinale Cupa Cintia, 21,

Naples 80126 Italy

2. Telethon Institute of Genetics and Medicine, Via Campi Flegrei 34, Pozzuoli (Naples), 80078,

Italy.

3. U.O.C. Genetica Medica, IRCCS Istituto Giannina Gaslini, 16147, Genova, Italy

4. Clinical Medicine and Surgery, University of Naples Federico II, Via S. Pansini 5, 80131

Naples Italy.

5. Laboratory of Molecular Biology, IRCCS Istituto Giannina Gaslini, Genova, Italy.

21   #These authors contributed equally to this work

22   * Corresponding author

23   ivan.conte@unina.it

Abstract

Endosomes have emerged as major signaling hubs where different internalized ligand-receptor complexes are integrated and the outcome of signaling pathways are organized to regulate the strength and specificity of signal transduction events. Ezrin, a major membrane-actin linker that assembles and coordinates macromolecular signaling complexes at membranes, has emerged recently as an important regulator of lysosomal function. Here, we report that endosomal-localized EGFR/Ezrin complex interacts with and triggers the inhibition of the Tuberous Sclerosis Complex (TSC) in response to EGF stimuli. This is regulated through activation of the AKT signaling pathway. Loss of Ezrin was deficient in TSC repression by EGF and culminated in translocation of TSC to lysosomes triggering suppression of mTORC1 signaling. Overexpression of constitutively active EZRIN^{T567D} is sufficient to relocalize TSC to the endosomes and reactivate mTORC1. Our findings identify EZRIN as a critical regulator of autophagy via TSC in response to EGF stimuli and establish the central role of early endosomal signaling in the regulation of mTORC1. Consistently, Medaka fish deficient for Ezrin exhibit defective endo-lysosomal pathway, attributable to the compromised EGFR/AKT signaling, ultimately leading to retinal degeneration. Our data identify a pivotal mechanism of endo-lysosomal signaling involving Ezrin and its associated EGFR/TSC complex, which are essential for retinal function.

Keywords: EGFR, EZRIN, endosomal signaling, TSC1, mTOR.

Abbreviations: RPE: retinal pigment epithelium; POS: photoreceptor outer segment; ERM: Ezrin-

radix-moesin; EZR: Ezrin; NHE1: sodium-hydrogen antiporter 1; EGFR: epidermal growth factor

receptor; Vsp11: vacuolar protein sorting 11; PI3K: Phosphoinositide-3-kinase; AKT: protein kinase B;

EGF: epidermal growth factor; TSC: Tuberous sclerosis proteins; mTOR: mammalian target of

rapamycin; GTPase: Guanine nucleotide-binding proteins; MEF: Mouse Primary Embryonic

Fibroblasts; LAMP1: Lysosomal Associated Membrane Protein 1; LC3: Microtubule-

associated proteins 1A/1B light chain 3B; CTSB: Cathepsin B; HER3: human epidermal growth factor

receptor 3; HER2: human epidermal growth factor receptor 2; EEA1: early endosome antigen 1;

MAPK: mitogen-activated protein kinase; MK2: Mitogen-activated protein kinase-

activated protein kinase 2; ARPE: Retinal Pigment Epithelial Cells.

INTRODUCTION

Endosomes are intracellular membrane-bound organelles that receive, integrate, and transmit a variety

of signals to intracellular compartments. Trafficking of receptors, ion channels, lipids, and other

effector proteins within the endosomal vesicles provides a mechanism to either sustain intracellular

signaling pathways active (Palfy, Remenyi, & Korcsmaros, 2012) or to downregulate signaling

pathways through their degradation in lysosomes. Accordingly, altered endosomal maturation and

function play a key role in the pathogenesis of a wide range of human diseases including diabetes,

cancer, and neurodegenerative disorders. Therefore, new insights about endosomal signaling and

understanding the molecular components restricting signaling activity to specific pathways will

uncover new opportunities for pharmacological targeting of such disorders. In the retinal pigment

epithelium (RPE), endosomes contribute to the diurnal clearance of phagocytosed photoreceptor outer

segments (POS) that is required for RPE and photoreceptor health. This process is linked to circadian

and light phase and is initiated by the scission from the plasma membrane of phagosomes containing

POS, which undergo gradual fusion with endosomes and finally with lysosomes in a coordinated

process termed "maturation". The high demand on lysosomes for the digestion and recycling of

phagocytosed POS rather than for the clearance of mitochondria, oxidized proteins, and other cellular

components suggests the existence of a signaling pathway that can finely coordinate lysosomal

function according to needs that will not upset cellular homeostasis. A long-standing question in the

field is how functional diversity within the autophagy pathway is achieved in the RPE in the dark and

light phases. Inhibition of endosomal biogenesis, trafficking, and fusion is associated with impairment

of lysosomal biogenesis and autophagy flux. Implicit in these findings is the idea that endosomes, by

carrying signaling molecules, could serve as a signaling hub for the regulated transfer of signals to

lysosomes, acting more specifically than diffusion-based signal propagation. However, how endosomes

are essential for lysosomal function, and their relevant components regulating this process are still not

well defined.

Ezrin, a member of the ezrin-radixin-moesin (ERM) protein family, is mainly localized just beneath the

plasma membrane around cellular protrusions and villi. Ezrin acts as a scaffolding platform to cross-

link F-actin cytoskeleton with specialized membrane components (Kawaguchi & Asano, 2022) that are

implicated in the spatiotemporal dynamics of phagosomes and endosomes. Its association with both F-

actin filaments and membrane proteins is finely regulated and requires conformational activation

thorough phosphorylation at unique (Y353, Y477, and T567) residues. The central role of Ezrin in

regulating trafficking of vesicles has been described (Cha et al., 2006; Tamma et al., 2005; Zhou et al.,

2003). Indeed, maturation of endosomes and recycling/exocytosis of their components (i.e. α1β-

adrenergic receptor, NHE3, and others) require the Ezrin protein (Barroso-Gonzalez, Machado, Garcia-

Exposito, & Valenzuela-Fernandez, 2009; Cha et al., 2006; Stanasila, Abuin, Diviani, & Cotecchia,

2006; Zhao et al., 2004; Zhou et al., 2005). The phosphorylated active Ezrin is observed within early

and late endosomes (Parameswaran, Enyindah-Asonye, Bagheri, Shah, & Gupta, 2013). Moreover,

through its active and reversible interactions with actin filaments and endosomal proteins, Ezrin organizes signal transduction. Indeed, phosphorylation of the T567 residue of EZRIN leads to its colocalization in a functional complex with NHE1, EGFR, and β1-integrin in human breast tumors, suggesting its crucial role as a scaffold protein of EGFR (Antelmi et al., 2013). Accordingly, Ezrin also interacts with EGFR at membranes (Saygideger-Kont et al., 2016). In mammalian cells, depletion of an Ezrin-interacting protein, Vsp11, delays the delivery of EGFR to endosomes (Chirivino et al., 2011), thus linking the Ezrin protein network with EGFR trafficking via clathrin-coated transport vesicles. However, the mechanisms of Ezrin-EGFR interaction and its function at the endosomal compartments remain largely unexplored. Interestingly, recent findings have shown that PI3K-mediated activation of AKT upon EGF stimulation is mediated by EGFR via an early endocytic pathway (Nishimura, Takiguchi, Ito, & Itoh, 2015). These findings suggest that Ezrin may be required as a protein scaffold for coordinating EGFR/AKT signaling at endosomes. Interestingly, EZR interacts with AKT in breast cancer cells (Li et al., 2019). In addition, the Y353-phosphorylation of Ezrin is relevant for PI3K-initiated signaling through its interaction with p85, the regulatory subunit of PI3K (Gautreau, Poullet, Louvard, & Arpin, 1999). Notably, the Ezrin-p85 complex optimizes the physiological activation of AKT, supporting a central role of Ezrin in controlling intracellular pathways in response to external cell signaling (Gautreau et al., 1999).

Accumulating evidence has shown that the AKT-mediated Tuberous Sclerosis Complex (TSC) phosphorylation is a major mechanism in triggering the activity of the GTPase Rheb (Ras homolog enriched in brain), an essential activator of mTORC1 at lysosomes. We demonstrated recently that Ezrin is a key regulator of lysosomal biogenesis and functions in RPE/retina crosstalk by modulating TFEB nuclear translocation (Naso et al., 2020). Moreover, Ezrin overexpression leads to altered autophagy and an impairment of POS maturation and degradation in RPE cells (Naso et al., 2020). Thus, it is intriguing that Ezrin has been recently observed to be associated with lysosomes (Poupon,

Stewart, Gray, Piper, & Luzio, 2003). Furthermore, cancer cell proliferation and invasion through an

activated Akt/mTOR pathway was linked with activation of Ezrin (Krishnan et al., 2006). In contrast,

depletion of Ezrin was found to be associated with the repression of the mTOR pathway (Wan,

Mendoza, Khanna, & Helman, 2005). Together, these data led us to hypothesize that Ezrin-mediated

EGFR endosomal sorting and trafficking could play a central role in mTOR activation on lysosomes.

Here, we identify a previously undocumented function for Ezrin as a platform that is essential for the

endosomal signaling network involving EGFR and AKT pathways, which provides an important

insight into the spatial inactivation of the TSC complex on endosomal compartments. We show that

inactivation of Ezrin is crucial to neutralize EGF-stimulated EGFR endosomal sorting and signaling

from the plasma membrane with a reduction of AKT-mediated phosphorylation of TSC, which in turn

translocates and inhibits mTORC1 on lysosomes. These results reveal an essential layer of mTOR

regulation by Ezrin and EGFR signaling and uncover part of the paradigm of signaling from

endosomes to lysosomes to coordinate the lysosomal function in the retina and other tissues. Consistent

with a role of the Ezrin/EGFR/TSC axis in lysosomal biogenesis and function, alteration of this

molecular network alters autophagy in vivo in Medaka fish, resulting in retinal degeneration.

Derangement of this control mechanism may underpin human eye disorders and may be relevant as a

therapeutic target to restore normal vision.

RESULTS

Ezrin regulates lysosomal biogenesis.

Activated Ezrin represses the autophagy pathway in the RPE (Naso et al., 2020), but the mechanism

remains undefined. To gain insights into this, we used integrated comparative analysis by unbiased

RNA-seq and high-resolution mass spectrometry-based proteomic studies on Ezrin$^{-/-}$ mouse embryonic

fibroblasts (MEFs) (EZR^{KO}) (Ognibene et al., 2011). The comparison of the transcriptomics and

proteomics identified 572 commonly regulated genes: 317 and 213 genes are induced and inhibited in

both datasets, respectively (Figure supplement 1A). Gene Ontology (GO) and Functional Annotation

Clustering analyses were performed on these 530 commonly differentially expressed genes (DEGs),

restricting the output Cellular Compartments (CC) terms (Tables supplement 1-3). We found an

enriched overlap of these genes in cell compartments, including cell membrane and lysosome (Figure

1A, Figure supplement 1A and supplement 1B and Tables supplement 1-3). Consistent with this,

immunofluorescence analysis revealed that the EZR^{KO} displayed an increased number of lysosomes, as

assessed by quantification of lysotracker-fluorescent staining and Lamp1-LC3 colocalization (Figure

1B-F). Compared with WT MEFs, lysosomal Cathepsin B (CTSB) activity of EZR^{KO} MEFs was

significantly increased (Figure 1G). Furthermore, western blot analysis also revealed that EZR^{KO}

increased the expression of lysosomal markers (LAMP1, CTSD and LC3) as well as reducing the levels

of the autophagy substrate p62 and NBR1 (Figure 1H). To further investigate whether this autophagic

induction was Ezrin dependent, we inserted a frameshift deletion of 13 nt in the coding region (exon 2)

of the *EZRIN* gene in HeLa cells via CRISPR/Cas9-mediated genome editing (EZR$^{-/-}$) (Figure

supplement 1D). Concordantly, we found that EZR$^{-/-}$ cells have a significantly increased number of

lysosomes and increased lysosomal activity (Figure supplement 1E and F and supplement 1H),

indicative of augmented lysosomal biogenesis and function. Consistent with this, western blot analysis

showed that lysosomal markers and autophagic flux were also increased in EZR$^{-/-}$ compared to control

(Figure supplement 1G and I). Taken together, these results reveal the crucial roles of Ezrin in

lysosomal biogenesis and function. These results are consistent with our previous report showing that

the autophagy pathway is blocked by Ezrin overexpression *in vivo* (Naso et al., 2020).

Ezrin interacts with EGFR and regulates its activation.

Previous studies have implicated Ezrin in coordinating signaling complexes on membranes in cancer,

raising the question of whether the Ezrin-mediated control of autophagy may be attributed to an

alteration of signaling pathways. To identify potential signaling pathways affected by Ezrin

modulation, we performed an enrichment analysis of the 530 differentially expressed genes in EZR^{KO},

using stable isotope labeling by amino acids in cell culture (SILAC) phosphoproteomics, kinase

perturbations from GEO database, and the Proteomics drug atlas. Interestingly, SILAC

phosphoproteomics data highlighted a significant overlap with phosphorylation changes in HeLa cells

upon EGF treatment (Figure 2A and Table supplement 4). Accordingly, kinase perturbation revealed a

significant overlap with downregulated genes upon EGFR drug activation (Figure supplement 2A and

Table supplement 4), whereas Proteomics drug atlas revealed a significant enrichment in cells upon

AZ628 (a Raf Inhibitor) or MEK162 (a MEK inhibitor) (Figure supplement 2B and Table supplement

4) (B. Liu, Chen, Johns, & Neufeld, 2006; Mitchell et al., 2023; Olsen et al., 2006; Ong et al., 2002;

Warde-Farley et al., 2010) (All resources are available https://maayanlab.cloud/enrichr-kg/downloads).

Thus, we hypothesized that the EGFR needs to be selectively recognized by EZRIN to be subjected to

EZRIN-mediated endosomal trafficking and signaling. Consistent with this, gene network based on

physical interaction reveals EGFR as a possible direct EZRIN protein partner (Figure 2B and Table

supplement 5). Moreover, EGFR resulted strongly upregulated in our omics dataset (Figure 2C) and co-

immunoprecipitation (CoIP) experiments revealed a complex composed of EZRIN and EGFR (Figure

2D), consistent with human biomedical interaction repositories (Oughtred et al., 2021; Petschnigg et

al., 2014; Salokas et al., 2022). Phosphorylated Ezrin (Thr567) localizes at curved cytoplasmic

membranes and has been implicated as a membrane-cytoskeleton scaffolding protein rather than a

membrane shaper (Tsai et al., 2018). We therefore tested whether phosphorylation at Thr567 of Ezrin

was involved in interacting with EGFR in a complex at cytoplasmic membrane. As suspected,

phosphomimic active-EZR^{T567D}, but not phospho-mutant inactive-EZR^{T567A} protein (Naso et al., 2020),

when expressed in EZR$^{-/-}$ cells, co-immunoprecipitated with EGFR (Figure 2D). These data support

that active EZRIN protein interacts with EGFR. Co-immunostaining analysis confirmed that EZRIN is

localized at the plasma membrane with EGFR (Figure 2E). Moreover, a fraction of the EZRIN signal

colocalizes with EGFR, within same intracellular compartments (Figure 2E), supporting the presence

of an EZRIN/EGFR complex, in which EZRIN acts as a scaffold protein for EGFR. Thus, we

postulated that EZRIN participates in EGFR trafficking and signaling. To test this hypothesis, we

examined the expression levels and subcellular distribution of EGFR under normal and EZRIN-

depleted conditions by immunofluorescence staining and Western blot (Figure 2F and G). Interestingly,

the genetic depletion of EZRIN strongly induced the localization of EGFR at the plasma membrane and

dramatically reduced its presence within intracellular compartments (Figure 2F). Considering that the

specific EGFR signals can arise from intracellular compartments, such as the endosomal compartment

(Burke, Schooler, & Wiley, 2001), we examined whether EZRIN depletion would impair EGFR

signaling. Consistent with proteomic results, we found that the levels of total EGFR are increased in

EZR$^{-/-}$ compared to control cells (Figure 2G). Western blot analysis demonstrated that the absence of

EZRIN induced a reduction in EGFR signaling. Notably, the level of HER3 and active pY845 EGFR

were almost abolished following EZRIN depletion, which also reduced the EGFR-stimulated

phosphorylation of p38 MAPK at Threonine 180 (T180) and Tyrosine 182 (Y182), and increased

phosphorylation at T222 of its substrate, MK2 (Figure 2G). These data suggest a model by which the

EZRIN interaction with EGFR contributes to EGFR trafficking and signaling.

Ezrin regulates endocytic EGFR sorting and signaling

Activation of EGFR leads to its internalization and trafficking to early endosomes, which sustains

specific EGFR signaling and recycling (Burke et al., 2001). We asked if the increased EGFR protein

level at cell membrane and the reduction of EGFR signaling in EZR$^{-/-}$ cells could be due to an

alteration in EGFR dimerization and packaging into endosomal vesicles. To test this hypothesis, the

cellular internalization and trafficking of EGFR basis was investigated by immunofluorescence and

live-imaging studies. We found that the lack of Ezrin reduced dimerization of EGFR upon EGF

stimulation (Figure 3A and B). These results were confirmed by immunofluorescence analyses; indeed,

compared to control cells, Ezrin-depleted cells showed higher levels of EGFR on the cell surface,

which was mirrored by reduced EGFR abundance at endosomal compartments, as assessed by a

reduction in the overlap between EGFR and EEA1 signals (Figure 3C) and increased EGFR protein

levels on purified membranes and endosomes (Figure 3D and E). The reduced internalization of EGFR

to endosomes was not accompanied by a suppression of endocytosis, as indicated by the slight and

significant increase in the number of EEA1-positive early endosomes and endotracker-positive

structures in EZR$^{-/-}$ compared to WT cells (Figure 3F and G). These results support that EGFR

accumulation at the plasma membrane was not a result of an endocytosis defect. To better define

whether the lack of EZRIN alters EGFR internalization and trafficking in EZR$^{-/-}$ cells upon EGF

stimulation, we performed Total Internal Reflection Fluorescence (TIRF) time-lapse imaging at high

spatiotemporal resolution. Both EZR$^{-/-}$ and control cells, transfected with an EGFR-GFP vector, were

imaged every 0.5 s for 5 min upon EGF treatment. Notably, the EGF-induced EGFR endosomal

internalization was dramatically abolished in EZR$^{-/-}$ compared to control cells (Figure 4A and B, Figure

5A and Video 1-4). Consistent with a defective EGFR integration in the early endosome, EGFR was

localized at the plasma membrane in EZR$^{-/-}$ cells, despite EGF stimulation (Figure 4A and B, Figure

5A and Video1-4). These results support that EZRIN play a key role for the dimerization, integration,

and trafficking of EGFR in the endosomes. To strengthen these findings, we performed ultrastructural

analysis using immunoelectron microscopy (IEM) that further revealed the reduced number of EGFR-

positive endosomal compartments and the increased presence of EGFR at the plasma membrane in

EGF-stimulated EZR$^{-/-}$ compared with EGF-stimulated control cells (Figure 5B). Consistent with this,

pY845 EGFR and pT180/pY182 p38 MAPK were reduced upon EGF stimulation in EZR$^{-/-}$ cells

(Figure 5C). Moreover, the increased EGFR internalization from membranes to endosomes by EGF

stimulation was significantly inhibited in EZR$^{-/-}$ cells compared to WT (Figure 5D). As expected, we

found that the endosomal EGFR internalization was further repressed in MEF-EZR^{KO} (Figure

supplement 2C-E) and in HeLa cells upon *NSC668394* treatment (Figure supplement 2F and G), a

specific Ezrin inhibitor (Naso et al., 2020). Taken together, these data strongly support a primary role

of EZRIN in mediating the internalization and trafficking of EGFR from plasma membrane to

endosomes.

The endosomal Ezrin-EGFR complex targets TSC1 protein

We next sought to identify the molecular networks by which Ezrin/EGFR axis controls lysosomal

biogenesis and function. Interestingly, EGFR stimulates several downstream effectors, including

PI3K/AKT signaling in response to multiple stimuli (Wee & Wang, 2017). This led us to investigate

the role of Ezrin/EGFR axis in the control of AKT signaling. AKT binds, phosphorylates, and inhibits

hamartin (TSC1) and tuberin (TSC2) complex. TSC complex is essential to turn off the activity of

Rheb, a crucial activator of mTORC1 at lysosomal surface (Dibble & Cantley, 2015). This raised the

possibility of Ezrin-mediated activation of EGFR signaling would be required for AKT activation and

thus stimulation of the mTORC1 pathway via TSC repression. To test this hypothesis, we analyzed the

interaction between endogenous Ezrin/EGFR with AKT and TSC1 to define an endosomal signaling

platform. In agreement with previously presented data (Haddad et al., 2002), TSC1 and AKT co-

immunoprecipitated with EZRIN (Figure 6A). Consistently, immunoprecipitation of EGFR was able to

pull down both TSC1 and AKT (Figure 6A), suggesting that EGFR, AKT, TSC1 and EZRIN are

present in a complex. The latter result led us to investigate whether EGFR could interact with AKT and

TSC1 indirectly through EZRIN. Co-immunoprecipitation experiments confirmed this possibility given

that *EZRIN* depletion abolished the interactions of EGFR with TSC1 and AKT, thereby point out the

key role of Ezrin as a scaffold protein for the formation and activation of the EGFR/AKT/TSC1

signaling (Figure 6A). In agreement with this hypothesis, lack of Ezrin reduced pS473 AKT activation

and in turn suppressed AKT-mediated phosphorylation of pS939 TSC2 (Figure 6B). As expected,

inactivation of AKT promoted activation of TSC1 and TSC2, which localized on the lysosomes in

EZR$^{-/-}$ cells (Figure 6C and D). Consistently, translocation of the TSC complex on the lysosomes led to

inhibition of mTORC1 pathway, as demonstrated by reduction of pT389 P70 S6 Kinase and pS65 4E-

BP1 levels (Figure 6B). These data were confirmed also in Ezrin-inhibited HeLa (Figure supplement

3A), ARPE-19 cells (Figure supplement 3B and C) and MEF EZR^{KO} cells (Figure supplement 3D and

E). Together, these results establish an Ezrin-dependent molecular machinery coordinating EGFR

sorting and signaling at the endosome to a well-regulated signals transfer to lysosomes via AKT/TSC

axis. Consistent with this idea, phosphorylation of pS473 AKT was significantly abolished in response

to EGF treatment in EZR$^{-/-}$ compared to control cells (Figure 6E). Accordingly, using confocal

Airyscan high-resolution microscopy, we found that the majority of the TSC complex was present in

early endosomes of HeLa WT cells upon EGF treatment, as shown by co-localization with the

endosomal marker EEA1 (Figure 6F). Notably, the endosomal TSC localization was abolished in EGF-

treated EZR$^{-/-}$ cells (Figure 6F), which indicates that EGFR-mediated repression of TSC by AKT

activation could occur in a stable endosomal complex dependent by Ezrin. Consistently, the ectopic

expression of a constitutively active EZR^{T567D} protein, but not a constitutively inactive Ezr^{T567A} protein,

rescued EGFR sorting and signaling activation at the endosome in EZR$^{-/-}$ cells upon EGF treatment

(Figure supplement 4A). Additionally, EZR^{T567D}, but not EZR^{T567A}, rescued the physiological

localization of TSC complex on the cytoplasm in EZR$^{-/-}$ cells (Figure supplement 4B). Moreover, full

translocation of TSC on endosomes was restored in EZR$^{-/-}$ cells expressing EZR^{T567D} protein after EGF

treatment, as shown by co-localization between TSC1 and EEA1 proteins (Figure supplement 4B).

Aberrant EGFR signaling induces retinal degeneration in EZR[-/-] medaka fish

To further investigate on the role of EZRIN/EGFR axis, which is conserved among vertebrates, in daily modulation of lysosomal biogenesis and function in retinal cells, we carried out *in vivo* experiments. Accordingly, we found that the EGFR expression pattern in the rodents' retina diminished in response to light and increased after light off (Figure supplement 5A and B), coinciding with Ezrin expression and diurnal lysosomal biogenesis in the RPE/retina (Naso et al., 2020). Consistently, we found an inhibition of TSC2 and an increase of AKT/mTOR pathway in the mice retina in response to dark condition, when active Ezrin (Naso et al., 2020) and EGFR are highly expressed (Figure supplement 5C). Moreover, TSC2 resulted activated and AKT/mTOR signaling decreased in response to light in the retina, when inactive Ezrin (Naso et al., 2020) and EGFR are weakly expressed (Figure supplement 5C), which suggests that activation of Ezrin underlies the requirement endosomal EGFR signaling to assemble the EGFR/AKT/TSC complex and represses lysosomal biogenesis. This data supported that EGFR signaling in retinal cells could be regulated by Ezrin for finely control lysosomal biogenesis and function in mTOR-dependent manner. Thus, we used the highly effective CRISPR-Cas9 mediated mutagenesis to create stable Ezrin mutant lines in Medaka fish (*Oryzias latipes*, Ol) as *in vivo* model system. Targeting two sgRNAs (sgRNA1 and sgRNA2) in the exon 1 of *Ezrin* gene, we generated a 386 bp deletion and established founder lines for this deletion. This mutation eliminates the first 129 amino acids containing ATG (olEzrin[Δ386]), generating a severely truncated Ezrin protein (Figure 7A), which was not detectable by western blot analyses (Figure 7B). This indicated that the Δ386 Ezrin allele is likely functionally null, and mutants will hereafter be called Ezrin[-/-] medaka line. Larval homozygous Ezrin[-/-] medaka line appeared almost visually indistinguishable from wild-type siblings, and the Δ386 allele was inherited in Mendelian ratios (Figure 7C). Interestingly, the Ezrin[-/-] medaka larvae recapitulated in part a previously reported phenotype characterized in postnatal ezrin knockout

mice (Bonilha, Rayborn, Saotome, McClatchey, & Hollyfield, 2006). Consistent with our *in vitro* data,

we observed increased levels of the EGFR protein accompanied by a significant reduction of active

pY845 EGFR. This was accompanied by a significant reduction of the phosphorylation of pS473 AKT

in Ezrin$^{-/-}$ medaka line compared to control larvae (Figure 7D). Consistently, we observed reduced

AKT-mediated phosphorylation of pT1462 of TSC2 and an increased autophagy, as demonstrated by

higher levels of LC3-II and Lamp1 (Figure 7D and E). Notably, endosomal internalization of EGFR

was significantly repressed in the RPE of Ezrin$^{-/-}$ medaka line. Consistent with defective EGFR

internalization and trafficking, whole mount immunofluorescence analysis showed that EGFR

accumulated at plasma membrane of RPE of Ezrin$^{-/-}$ medaka line compared with control fish (Figure

7F). Considering the role of endosomal sorting and signaling in the health of retinal cells (Toops, Tan,

& Lakkaraju, 2014), we addressed the consequences of aberrant EGFR signaling pathway in retina of

Ezrin$^{-/-}$ medaka line. Notably, defective endosomal EGFR signaling was sufficient to induce deleterious

consequences for the health of photoreceptor cells, which showed reduction in photoreceptor outer

segments compared with native rods, similarly to the pathogenesis of macular degeneration (Borrelli et

al., 2020; Kaur & Lakkaraju, 2018). Altogether, these data support the dynamic regulation of EGFR

signaling at endosomal compartments in response to Ezrin activation, which assembles and activates an

EGFR/AKT/TSC signalosome at endosomes to finely regulate the lysosomal signaling by mTOR

pathway, required for the correct autophagy and retinal cell health (Figure 8).

DISCUSSION

Canonical EGFR signaling begins at the plasma membrane with the engagement of the EGF ligand

(Tanaka et al., 2018). Emerging studies have indicated that sorting of the EGF-EGFR complex to

endosomal vesicles requires spatiotemporally defined encounters with distinct cytoskeleton platforms

resulting in internalization, activation, maintenance or termination of EGFR signaling (Ceresa, 2012;

Wang, Pennock, Chen, & Wang, 2002). Consistent with this notion, it is now increasingly recognized

that many molecules participating in signal transduction are central sorting hubs that coordinate

signaling from and to different intracellular compartments, including early endosomes, late endosomes,

phagosomes, and lysosomes (Sorkin & von Zastrow, 2009). However, molecular networks determining

selective signal transduction from endosomes to lysosomes are not well defined. In this study, we

demonstrated that EZRIN is a cytoskeleton scaffold protein aligned along internal membranes and that

this localization is essential for endosomal EGFR signal transduction to the TSC complex. Endosomal

EGFR sorting and activation occurs mainly due to binding with EZRIN that facilitates dimerization and

activation of the EGF-EGFR receptor complex, resulting in their recruitment to endosomes, followed

by AKT activation that targets and inhibits the TSC complex. Indeed, time-lapse confocal imaging

revealed that EGFR fails to be recruited to endosomal compartments upon EGF stimulation in the

absence of Ezrin or in the presence of its inactive form (EZRIN^{T567A}). The loss of Ezrin function

compromises EGFR-mediated AKT activation, which in turn reduces TSC complex inhibition resulting

in TSC complex translocation to lysosomes where it constrains mTOR activity. Consistent with this,

not only does loss of Ezrin impact on lysosomal TSC complex translocation, but we also documented

that dephosphorylation of Ezrin in its inactive form is required for EGFR inactivation and TSC1 and

TSC2 release from an EGFR/EZRIN complex, possibly to support the lysosomal biogenesis and

function. Indeed, autophagy appears to be highly sensitive to pharmacological Ezrin inhibition via the

EGFR/AKT axis. Moreover, overexpression of EZRIN^{T567A}, but not EZRIN^{T567D}, fails to restore EGFR

endosomal signaling and lysosomal function in Ezrin-defective cells, indicating that phosphorylation of

Ezrin is indispensable for this activity. Supporting this possibility, loss of βA3/A1-crystallin affects

PITPβ/PLC signaling axis associated with an age-related loss of PLC-mediated Ezrin phosphorylation

and subsequent compromised RPE cell polarity and EGFR signaling. Notably, the lysosome-mediated

POS clearance was disrupted in the Cryba1 cKO RPE (Shang et al., 2021). Beyond this, our findings

also showed an interaction of EGFR with the TSC complex and their co-localization with endosomes,

opening to future work on mechanisms of how endosomal system connects extracellular signals with

lysosomes under different physiological and pathological conditions.

Upregulation of Ezrin has been shown to induce an age-related macular degeneration-like phenotype in

miR-211[-/-] mice (Naso et al., 2020), where light-mediated cell clearance is completely abolished. We

can speculate that recruitment of EGFR on endosomal compartments by Ezrin orchestrates a local

signal between endosomes and lysosomes to drive tight control on the lysosomal cargo demands,

although future studies are needed in this regard. Interestingly, inhibition of EGFR activity, by

silencing Rubicon (RUBCN), switches lysosomal cargo degradation from POS associated to LC3-

associated phagocytosis to autophagy process in the RPE cells (Muniz-Feliciano, Doggett, Zhou, &

Ferguson, 2017). However, the molecular mechanisms are completely unknown. Therefore, Ezrin may

represent a nodal point in endosomal compartments where EGFR signaling and AKT converge and

integrate to directly control the TSC/mTOR pathway and lysosomal cargo demands and degradation.

Notably, mutations affecting TSC1 and TSC2 alter lysosomal function with retinal manifestations in

40–50% of individuals (Rosset, Netto, & Ashton-Prolla, 2017). This phenotype is also showed in RPE-

specific deletion of TSC1 profoundly leading to an age-related impairment in lysosomal function

associated with RPE degeneration *in vivo* (Huang et al., 2019).

The findings of the cellular mechanisms governing endosomal EGFR sorting and signaling might be of

therapeutic relevance. Indeed, alteration of endosomal biogenesis and signaling have been shown to

participate in the age-related, progressive neurodegeneration such as in age-related macular

degeneration and Alzheimer's disease (REF Kaur et al). Thus, the identification of the mechanisms that

control Ezrin/EGFR/mTOR molecular network might be exploited for the treatment of diseases in

which defective endo-lysosomes play a part.

MATERIALS AND METHODS

3' mRNA sequencing library preparation

The transcriptional response of four biological replicates for both MEF^{WT} and MEF^{Ezr} KO cell lines

was analyzed using QuantSeq 3' mRNA sequencing. RNA extraction, quality control and preparation

of RNA-seq libraries and sequencing on an NovaSeq6000 platform were carried out in collaboration

with the Next Generation Sequencing (NGS) Facility at TIGEM following standard procedures as

previously described (Carotenuto et al., 2022). An average yield of ~4.5 Mb was obtained per sample.

Computational analysis of deep sequencing data

Data analysis was performed using the pipeline already established at the Bioinformatics and Statistics

Core Facility at TIGEM (Pinelli et al., 2016). Briefly, the reads were trimmed to remove adapter

sequences and low-quality ends and reads mapping to contaminating sequences (e.g. ribosomal RNA,

phIX control) were filtered-out. Alignment was performed with STAR 2.6.0a3 (Dobin et al., 2013) on

mm10 reference assembly obtained from cellRanger website4 (Ensembl assembly release 93). The

expression levels of genes were determined with htseq-count 0.9.15 using mm10 Ensembl assembly

(release 93) downloaded from the cellRanger website4. We filtered out all genes having < 1 cpm in less

than n_min samples and Perc MM reads > 20% simultaneously. Differential expression analysis was

performed using edgeR6 (S. Liu et al., 2021).

Mass Spectrometry

Protein extraction and preparation of MS samples were carried out in accordance with standard

procedures currently utilized in the Mass Spectrometry Facility at TIGEM. About 30 mg of cell lysate

was used. Peptides were purified using the iST Kit (Preomics) following the company instructions.

Peptide separation and LC MS/MS analysis was carried out accordingly to standard procedures as

detailed in (Di Malta et al., 2023)

Data analysis of mass spectrometry

At least three independent biological replicates were performed for all experiments. For mass

spectrometry, all acquired raw files were processed using MaxQuant (1.6.2.10) and the implemented

Andromeda search engine. For protein assignment, spectra were correlated with the UniProt Homo

Sapiens including a list of common contaminants. Searches were performed with tryptic specifications

and default settings for mass tolerances for MS and MS/MS spectra. Carbamidomethyl at cysteine

residues was set as a fixed modification, while oxidations at methionine and acetylation at the N-

terminus were defined as variable modifications. The minimal peptide length was set to seven amino

acids, and the false discovery rate for proteins and peptide-spectrum matches to 1%. The match-

between-run feature with a time window of 0.7 min was used. For further analysis, the Perseus

software was used and first filtered for contaminants and reverse entries as well as proteins that were

only identified by a modified peptide. For full proteomes and IP-interactomes, the LFQ Ratios were

logarithmized, grouped and filtered for min. valid number (min. 3 in at least one group). Missing values

were replaced by random numbers that are drawn from a normal distribution. Finally, the intensities

were normalized by subtracting the median intensity of each sample. Significantly regulated proteins

between conditions were determined by student t-test using FDR < 0.05 as threshold.

Functional analysis on transcriptomics and proteomics data

The threshold for the statistical significance of gene expression was FDR < 0.05. The threshold for the

statistical significance of the proteomics analysis was -log10 > 1,3 and -Log2 > =1. GOEA and KEGG

Pathway were performed on induced and inhibited genes, separately, both in the transcriptome and in

the proteome experiments using the DAVID Bioinformatic tool (Huang da, Sherman, & Lempicki,

2009a, 2009b) restricting the output to Biological Process (BP), Cellular Compartments (CC) terms.

The threshold for statistical significance of GOEA was FDR <0.1 and the Enrichment Score $\geq$ 1.5,

while for the KEGG Pathway analyses it was FDR < 0.1. The comparison of the transcriptomics and

proteomics identified 572 commonly regulated genes: 317 and 213 genes were induced and inhibited in

both datasets, respectively.

Data visualization

Heatmap and Venn diagram were generated using custom annotation scripts.

Accession code

The transcriptomics data have been deposited in the NCBI Gene Expression Omnibus (GEO) (Edgar,

Domrachev, & Lash, 2002) and are accessible through GEO Series accession number GSE195983**.** The

title of the dataset is: "Transcriptome profile of EZR_KO cells"**.** For this dataset, a secure token has

been created to allow a review of the record: private token. The proteome data was deposited in PRIDE

repository and are available via ProteomeXchange with identifier PXD045157.

Western blot analysis

After transfection and/or treatments, cells were collected to extract total protein, endosomal proteins

(Invent, biotechnologies, ED-028), or membrane proteins (Thermo Fisher, 89842), while mouse eyes

were enucleated, and the retina was separated from the RPE. Both mice and cell samples were lysed

using RIPA buffer (150 mM sodium chloride, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1%

sodium dodecyl sulfate, 50 mM Tris, pH 8.0) with an inhibitor cocktail (Thermo Fisher Scientific,

78420). The protein concentration was determined by Bradford analysis and quantified using a Thermo

Fisher Helios γ spectrophotometer. Proteins were fractionated by sodium dodecyl sulfate-

polyacrylamide gel electrophoresis (SDS-PAGE) and transferred to PVDF membranes (EMD

Millipore, IPVH00010), then blocked in Tween 0.1% Tris-buffered saline containing 5% bovine serum

albumin (Tocris 5217) for at least 1h at room temperature and subsequently incubated overnight at 4°C

with primary antibodies. For Western blot analysis, the following antibodies were used: mouse anti-

NBR1 (1:1000, Abnova MO1), rabbit anti-LAMP1 (1:500, Sigma L1418), mouse anti-Ezrin (1:1000,

Novex 357300), mouse anti-SQSTM1/P62 (1:1000, Abcam ab56416), rabbit anti-Cathepsin D (1:1000,

Cell Signaling 2284), rabbit anti-LC3 (1:1000, Novus NB100-2220), mouse anti-GAPDH (1:1000,

Santa Cruz SC-32233), rabbit anti-HER2/ErbB2 (1:1000, Cell Signaling 2165), rabbit anti-

HER3/ErbB3 (1:1000, Cell Signaling 12708), rabbit anti-phospho-EGF receptor (Tyr845) (1:1000, Cell

Signaling 6963), rabbit anti-EGF receptor (1:1000, Cell Signaling 4267), rabbit anti-MAPKAPK-2

(1:1000, Cell Signaling 3042), rabbit anti-phospho-MAPKAPK-2 (Thr222) (1:1000, Cell Signaling

3316), rabbit anti-p38 MAPK (1:1000, Cell Signaling 8690), rabbit anti-phospho-p38 MAPK

(Thr180/Tyr182) (1:1000, Cell Signaling 4511), rabbit anti-ZO1 (1:1000, Abcam ab216880), mouse

anti-EEA1 (1:1000, BD 610457), rabbit anti-Tuberin/TSC2 (1:1000, Cell Signaling 4308), rabbit anti-

phospho-Tuberin/TSC2 (Ser939) (1:1000, Cell Signaling 3615), rabbit anti-phospho-Tuberin/TSC2

(Thr1462) (1:1000, Cell Signaling 3617), rabbit anti-p70 S6 Kinase (1:1000, Cell Signaling 9202),

mouse anti-phospho-p70 S6 Kinase (Thr389) (1:1000, Cell Signaling 9206), rabbit anti-Akt (1:1000,

Cell Signaling 9272), rabbit anti-phospho-Akt (Ser473) (1:1000, Cell Signaling 4060), rabbit anti-4E-

BP1 (1:1000, Cell Signaling 9644), rabbit anti-phospho-4E-BP1 (Ser65) (1:1000, Cell Signaling 9456),

rabbit anti-phospho-4E-BP1 (Thr37/46) (1:1000, Cell Signaling 2855), rabbit anti-Hamartin/TSC1

(1:1000, Cell Signaling 6935), mouse anti-EGFR (1:500, Santa Cruz sc-120), mouse anti-p-EGFR

(1:500, Santa Cruz sc-57542). After washing 3 times with Tween 0.1% Tris-buffered saline (TBS-T),

the membranes were incubated for 1 h at room temperature with the following secondary antibodies:

goat anti-rabbit IgG antibody, HPR conjugate, and goat anti-mouse IgG antibody HPR conjugate

(1:10,000 EMD Millipore, 12-348; 12-349). Western blot detection was done with ChemiDoc XRS+

System-Bio-Rad and quantified using ImageJ software.

Immunofluorescence

Mouse eyes were fixed overnight in 4% paraformaldehyde in PBS at 4°C and then cryopreserved by

treatment first with 5% and then with 30% sucrose in phosphate-buffered saline and embedded in OCT.

Twenty-micrometer cryosections were collected on slides (Superfrost Plus; Fisher Scientific,

Pittsburgh, PA). Cells were fixed with 4% paraformaldehyde (Chem Cruz sc-281692) for 15 min at

room temperature followed by washing with 1% PBS. After fixation, the cells were permeated with

blocking buffer (0.5% BSA, 0.005% saponin, 0.02% NaN$_3$) for 1 h at room temperature. Medaka fish

at stage 40 were subjected to anesthesia and then fixed by incubation in 4% PFA for 4 h at room

temperature (RT). Samples were rinsed three times with PTW 1X (1X PBS, 0.1% Tween, pH 7.3) and

then incubated overnight in 15% sucrose/PTW 1X at 4°C, and then again incubated overnight in 30%

sucrose/PTW 1X at 4°C and embedded. Sixteen-micrometer cryosection were collected on slides. The

following primary antibodies were used: rat anti-LAMP-1 (1:400, Santa Cruz sc-19992), mouse anti-

LAMP-1 (1:1000, DSHB H4A3), rabbit anti-LAMP1 (1:100, Abcam ab24170), rabbit anti-LC3B

(1:200, Novus NB100-2220), rabbit anti-EGF receptor (1:50, Cell Signaling 4267), mouse anti-EEA1

(1:100, BD 610457), rabbit anti-Tuberin/TSC2 (1:100, Cell Signaling 4308), rabbit anti-

Hamartin/TSC1 (1:1000, Cell Signaling 6935), mouse anti-EGFR (1:50, Santa Cruz sc-120), chicken

anti-GFP (1:500, Abcam ab13970), LysoTracker Red (Invitrogen L7528), CellLight Early Endosomes-

RFP (Invitrogen C10587). All incubations were performed overnight at 4°C. After washing with 1%

PBS, slides were incubated with the following secondary antibodies: Alexa 488 goat anti-

rabbit/mouse/Chicken (1:1000, Invitrogen A-11008 rabbit, A-11032 mouse, A-11039), Alexa 594 goat

anti-mouse/rat (1:1000, Invitrogen A-11032 mouse, A-11007 rat) and DAPI (1:500, Vector

Laboratories H-1200) for 1 h at room temperature; then, the slides were washed with 1% PBS and

mounted with PBS/glycerol and imaged with a Zeiss LSM800 microscope. Three dimensional images

were imaged with a Zeiss LSM880 confocal microscope equipped with Airyscan super-resolution

imaging module, using x63/1.40 NA Plan Apochromat Oil DIC M27 objective lens (Zeiss

MicroImaging, Jena, Germany).

Live cell imaging

HeLa cells were transiently transfected with EGFR-GFP and treated as indicated in the Fig.s. Time

lapse video were acquired for 5 min. One frame was acquired roughly every 0.5 s with lasers set at

30% power or below. Total Internal Reflection Fluorescence (TIRF) time-lapse imaging was performed

with a 60 x Plan Apo oil immersion lens using a Nikon Eclipse Ti Spinning Disk microscope, and

images were annotated, and the video was reconstitute using ImageJ software.

Image analysis

Lysotracker and Endotracker quantification. Fluorescent images of the cells were captured at 40 X

magnification using a LSM700 Zeiss Confocal Microscopy system, converted to grey-scale and

normalized to background staining, using ImageJ. Quantification of lysotracker and endotracker

reactivity was measured as mean values to define fluorescence signal intensity (IntDen/Area) and as the

area occupied by fluorescent-labeling in each region of interest.

LC3-LAMP-1, TSC1-LAMP-1 and TSC2-LAMP-1 colocalization. The colocalization of LC3 (green)

and LAMP-1 (red) and TSC1/TSC2 (green) and LAMP-1 (red) were evaluated using a LSM700 Zeiss

Confocal Microscopy after immunostaining of endogenous proteins. Average values were calculated

over 10 images, each containing a mean of 10 cells per image, and collected from at least three independent experiments. Exposure settings were unchanged throughout acquisition. Images were analysed using the JaCoP plugin (Bolte & Cordelieres, 2006) in ImageJ software.

EGFR-positive endosome quantification. Morphometric analysis of the distribution of gold particles (EGFR-labeled) at endosomal structures was performed using iTEM software (Olympus SYS, Germany). In detail, we counted the number of EGFR-positive endosomes on almost 10 26500 x magnification images. In the absence of specific staining, early endosome identification relied on morphological characteristics described in the literature (Vogel et al., 2015).

Cathepsin B assay

Cathepsin B activity was measured by a fluorometric assay kit (AB65300; Abcam, Cambridge, MA, USA) following the manufacturer's instructions. The reaction and fluorescence were read at 400 nm (excitation) and 505 nm (emission) on Promega GloMax discover.

Cross-linking assay

HeLa cells were washed twice with PBS and then cross-linked with DSP solution (Lomant's Reagent, Thermo Fisher 22585) at a final concentration of 1 mM for 30 min at RT. The reaction was stopped by adding stop solution (Tris-HCl 1 M, pH 7.5) at a final concentration of 10 mM for 15 min at RT. Each sample was analyzed by Western blot assay.

Immunoprecipitation assay

Cells were washed three times with ice-cold PBS and then homogenized with ice-cold lysis buffer (20 mM Tris-HCl pH 7.4, 150 mM NaCl, 1 mM EDTA, 0.5% NP40). At least 1 mg of proteins were immunoprecipitated with mouse anti-Ezrin (Novex, 357300) and rabbit anti-EGF receptor (Cell

Signaling 4267) in rotation at 4° C overnight. Then, the immunoprecipitates were conjugated with

protein G Beads (Dynabeads Protein G, Thermo Fisher Scientific 10004D), eluted in Laemmli buffer,

and subjected to immunoblot analyses.

Cell culture and treatments

ARPE-19, HeLa, and MEF cell lines were obtained from American Type Culture Collection (ATCC).

ARPE-19 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM)/F-12, while HeLa and

MEF cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Gibco) supplemented with

10% (v/v) FBS and 5% penicillin-streptomycin. All cell lines were maintained at 37°C, 5% CO_2 in a

humified incubator according to the guidelines provided by the vendors. MEF EZR^{KO} cells were kindly

donated by Alessandra Eva of Istituto G. Gaslini, Genova, Italy. To analyze the autophagic flux, cells

were treated with 200 nM Bafilomycin A1 (Sigma-Aldrich, B1793) for 3h in an incubator and

maintained in starvation for 30 min in HBSS medium (Thermo Fischer Scientific, 14025092)

supplemented with 10 mM HEPES (Thermo Fischer Scientific, 156330080). To evaluate EGFR

localization in immunoelectron microscopy, HeLa cells were treated with 100 μg/ml cycloheximide

(CHX) (Sigma-Aldrich, C4859). Drug treatment was performed for 6 h with 10 μM of *NSC668394* or

DMSO as previously reported (Bulut et al., 2012). EGF stimulation was obtained with 10 ng/ml of

animal-free recombinant human EGF (Peprotech AF-100-15) for 3h. We used a sub-confluent cell

culture (i.e. 80% of confluence) for each *in vitro* experiment.

Generation of an EZR$^{-/-}$ HeLa cell line

HeLa (ATCC CCL-2) full knock-out of the *EZRIN* gene was generated using the CRISPr/Cas9 system.

The   gRNA   sequence   CAATGTCCGAGTTACCACCA was   selected   using

the http://crispor.tefor.net/crispor.py online tool. HeLa cells were electroporated using the Amaxa

system with the nucleofection kit Cat No VCA-1003 from Lonza. Cells were FACS-sorted into 96-well

plates to obtain single-cell derived colonies carrying the INDEL mutations. Upon genomic DNA

extraction, the genomic sequence containing the targeted region were amplified by PCR reaction with

the   specific   primers: hEZRNup   TGCCGTCGCCACACTGAGGA, hEZRNlow

TCCTTTGCTTCCATGCCTGG. PCR products were analyzed by DNA Sanger sequencing and the cell

clone carrying the homozygous deletion *c.23 DEL AGTTACCACCATG* was selected and expanded.

Plasmids and Transfections

Cells were transfected at 80% confluence using Lipofectamine 2000 (Invitrogen, 12566014), following

the manufacturer's protocol. The plasmids used were Ezrin^{T567D} and Ezrin^{T567A}-mCherry, modified

from vectors described by Coscoy et al, provided by the S.Coscoy lab (Insitute Curie, Paris) (Coscoy et

al., 2002), and EGFR-GFP (Addgene, 32751).

Immunoelectron microscopy analysis

HeLa cells were fixed with a mixture of 4% paraformaldehyde (PFA) and 0.05% glutaraldehyde (GA)

for 10 min at RT, then washed with 4% PFA once to remove the residual GA and fixed again with 4%

PFA for 30 min at RT. Next, the cells were incubated with a blocking/permeabilizing mixture (0.5%

BSA, 0.1% saponin, 50-mM NH$_4$Cl) for 30 min and subsequently with the primary monoclonal

antibody anti-GFP, diluted 1:500 in blocking/permeabilizing solution. The following day, the cells

were washed and incubated with the secondary antibody, an anti-rabbit Fab fragment coupled to 1.4 nm

gold particles (diluted 1:50 in blocking/permeabilizing solution) for 2 h at RT. The cells were then

post-fixed as described in Polishchuk and Polishchuk(Polishchuk & Polishchuk, 2019). After

dehydration, the specimens were embedded in epoxy resin and polymerized at 60°C for 72 h. Thin 60

nm sections were cut on a Leica EM UC7 microtome. The EM images were acquired from thin sections

using a FEI Tecnai-12 electron microscope equipped with a VELETTA CCD digital camera (FEI,

Eindhoven, the Netherlands).

Animals and Ethics Approval Statement

All studies on animals were conducted in strict accordance with the institutional guidelines for animal

research and approved by the Italian Ministry of Health, Department of Public Health, Animal Health,

Nutrition and Food Safety in accordance with the law on animal experimentation (article 31; D.L.

26/2014; protocol number: 0016304-21/07/2020-DGSAF-MDS-P.

RPE and retina dissection

To analyze protein expression levels in RPE individually, mouse eyes were dissected to remove optic

nerve, cornea, lens, and retina in ice-cold PBS 1X under stereomicroscopy (Leica). The RPE was

peeled from the eyecup and transferred to a tube containing 100 µl of RIPA buffer. RPE cells were

pelleted by centrifugation at 12,000 x g for 15 min at 4°C.

Light/Dark adaptation of mice for tissue isolation

Mice were maintained in dark conditions with a maximum of 0.4 lux from 19:00 pm to 7:00 am. Then,

animals were kept in a room with the light phase (450 lux) from 7:00 am to 19:00 pm. For light/dark

transition studies, some animals were transferred after 3 h from light conditions to dark conditions and

sacrificed. Eyes from dark mice were isolated under dim red light.

Medaka stocks

The cab strain of wild-type and Ezrin$^{-/-}$ medaka (*Oryzias latipes*) lines were maintained following

standard conditions (i.e., 12h/ 12h dark/light conditions at 27°C). Embryos were staged according to

the method proposed by Iwamatsu (Iwamatsu, 2004). All studies on fish were conducted in strict

accordance with the Institutional Guidelines for animal research and approved by the Italian Ministry

of Health, Department of Public Health, Animal Health, Nutrition and Food Safety in accordance with

the law on animal experimentation (D. Lgs.26/2014). Furthermore, all fish treatments were reviewed

and approved in advance by the Ethics Committee at the TIGEM institute (Pozzuoli (NA), Italy).

Ezrin[-/-] medaka generation by CRISPR/Cas9 system

The genomic sequence of medaka *Ezrin* was obtained the medaka genome database at the Ensembl

Genome Database Project (http://www.ensembl.org/Oryzias_latipes; ENSORLG00000012128). Design

and construction of *OlEzrin*-sgRNA was committed to SYNTHEGO. The sequences of *OlEzrin*-

sgRNA oligonucleotides are listed in Table 1. Instead of Cas9 mRNA, the commercial reagent of

pCS2-nCas9n (Addgene, #4729) was used in this study. After pCS2-nCas9n was digested by NotI

treatment, this linearized vector was used as the template for synthesizing capped Cas9 mRNA with a

mMessage mMachine SP6 Kit (Life Technologies). Microinjection of the medaka embryos followed a

method described preciously by Kinoshita et al. (Kinoshita, Kani, Ozato, & Wakamatsu, 2000). A

mixture containing 200ng/uL of Cas9 mRNA and 20ng/ul of *OlEzrin*-sgRNA was prepared and

injected into the fertilized eggs at the one-cell stage. After hatching, the larvae were raised to sexual

maturity and used as "founder" fish (F0). To observe the genomic DNA mutations induced by Cas9

and *OlEzrin*-sgRNA in CRISPR/Cas9-mediated Ezrin-mutated medaka, a small piece of the caudal fin

from individual F0 fish was collected and subjected to genomic DNA analysis, using the primer set

indicating in Table 1. After the above screening had confirmed the occurrence of CRISPR/Cas9-

mediated Ezrin mutation in the F0 generation, these founder fish were crossed with each other, and

their offspring (F1) were checked for Ezrin mutations in the same way. Two of the F1 progeny with the

same mutation patterns were mated to produce the F2 generation. The F2 generation were crossed with

each other to produce F3 progeny, that was screened as described above to confirm that the same

mutation patterns were successfully inherited.

	Name	Sequence (5'-3')	Usage
EZRIN- gRNA	*olEzrin* Sense	ACAATGGATGAGCCTATTAG	CRISPR/Cas 9-LoxP target site
	olEzrin Antisense	AGACTGATGCTGCCTCACTG	
Primers	*olEzrin*_Forward	GAACTCCTTCTAGCACCC	PCR for olEzrin genotype
	*olEzrin*_Reverse	CCGCCTCCCTCCTCAATC	

Table 1. Primer sequences.

Whole mount immunostaining

Medaka larvae were fixed in 4% PFA, 2X PBS, and 0.1% Tween-20. The fixed larvae were washed

with PTW 1X and digested for 20 min with 10 µg/ml proteinase K and washed two-fold with 2 mg/ml

glycine/PTW 1x. The samples were fixed for 20 min in 4% PFA, 2X PBS, and 0.1% Tween-20,

washed with PTW 1X, and then incubated for 2 h in FBS 1%/PTW 1X, at room temperature. The

larvae were incubated with mouse anti-EGFR (1:50, Santa Cruz sc-120) overnight at 4°C. The samples

were washed with PTW 1X, incubated with the secondary antibody, Alexa-488 goat anti-mouse IgG

(ThermoFisher), then with DAPI. Finally, the larvae were placed in glycerol 100%.

Statistical analysis

<u>T-test, Welch's t-test, Mann-Whitney test.</u> For the analysis of the statistically significant differences

between two conditions, we performed the Shapiro-Wilk test to check if each condition had followed

the normal distribution (null hypothesis): we performed the non-parametric Mann-Whitney test in case

of rejection of the null hypothesis (p-value < 0.05), and we performed the parametric unpaired t-test in

case of non-rejection of the null hypothesis (p-value ≥ 0.05). In the second case, we also performed the

F-test to check the homoscedasticity between the compared conditions (null hypothesis): we applied

the parametric Welch's t-test in case of rejection of the null hypothesis (p-value < 0.05). All the tests

were performed with GraphPad Prism 10.0.0, GraphPad Software, Boston, Massachusetts USA.

<u>ANOVA, Welch's ANOVA, Kruskal-Wallis test (with multiple comparisons post hoc tests).</u> For the

analysis of the statistically significant differences among multiple conditions, we performed the

Shapiro-Wilk test to check if each condition had followed the normal distribution (null hypothesis): we

performed the non-parametric Kruskal-Wallis test in case of rejection of the null hypothesis (p-value $<$

0.05), and we performed the parametric one way ANOVA in case of non-rejection of the null

hypothesis (p-value ≥ 0.05). In the second case, we also performed the Brown-Forsythe test to check

the homoscedasticity between the compared conditions (null hypothesis): we applied the parametric

Welch's one way ANOVA in case of rejection of the null hypothesis (p-value < 0.05). For

completeness, we computed the p-values with post hoc tests for the pairwise multiple comparisons:

Tukey's test for one way ANOVA, Dunnett's test for Welch's one way ANOVA, and Dunn's test for

Kruskal-Wallis test. All the tests were performed with GraphPad Prism 10.0.0, GraphPad Software,

Boston, Massachusetts USA.

<u>Poisson Regression.</u> For the analysis of the statically significant differences between two conditions

with discrete values (i.e., counts), we performed the Poisson Regression over data, considering a

Generalized Linear Model with Likelihood Ratio test. No correction for multiple comparisons was

necessary. Poisson Regression with Generalized Linear Model and Likelihood Ratio test were

performed with the package 'car' (version 3.1-2) in the R environment (version 4.2.3).

ACKNOWLEDGEMENTS

681 We are grateful to Dr. Cathal Wilson for critical reading and English editing of the manuscript. We are

682 grateful to Edoardo Nusco for mice technical support. Acknowledgment is made to BioRender

683 (www.biorender.com) for model images.

AUTHOR CONTRIBUTIONS

GG and DI performed the experiments and analyzed the data. DI and GG contributed to the

experimental design, implementation, and interpretation. SB and FGS contributed to the technical part

of the work. EDP and RDC performed bioinformatics analysis. EDP performed and supervised

statistical evaluations. RDC analyzed transcriptomics and proteomics data. FA and JM generated HeLa

$EZR^{-/-}$ cells. MO and AE donated MEF EZR^{KO} cells. PG performed and analyzed mass spectrometry

analysis. GG, DI and IC conceived the experiments, analyzed the data, and wrote the manuscript. All

authors read and approved the final manuscript.

FUNDING

693 Work in the Conte group was supported by grants from the Million Dollar Bike Ride Grant Program

694 MDBR-21-103-CHM, International Retinal Research Foundation, MIUR FISR2020IP_03551, MIUR –

695 PRIN, and Sanfilippo Children's Foundations and National MPS Society.

REFERENCES
697 Antelmi, E., Cardone, R. A., Greco, M. R., Rubino, R., Di Sole, F., Martino, N. A., . . . Reshkin, S. J.
698   (2013). ss1 integrin binding phosphorylates ezrin at T567 to activate a lipid raft signalsome
699   driving invadopodia activity and invasion. *PLoS One, 8*(9), e75113.
700   doi:10.1371/journal.pone.0075113

Barroso-Gonzalez, J., Machado, J. D., Garcia-Exposito, L., & Valenzuela-Fernandez, A. (2009). Moesin regulates the trafficking of nascent clathrin-coated vesicles. *J Biol Chem, 284*(4), 2419-2434. doi:10.1074/jbc.M805311200

Bolte, S., & Cordelieres, F. P. (2006). A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc, 224*(Pt 3), 213-232. doi:10.1111/j.1365-2818.2006.01706.x

Bonilha, V. L., Rayborn, M. E., Saotome, I., McClatchey, A. I., & Hollyfield, J. G. (2006). Microvilli defects in retinas of ezrin knockout mice. *Exp Eye Res, 82*(4), 720-729. doi:10.1016/j.exer.2005.09.013

Borrelli, E., Sacconi, R., Zuccaro, B., Cavalleri, M., Bordato, A., Zucchiatti, I., . . . Querques, G. (2020). Photoreceptor alteration in intermediate age-related macular degeneration. *Sci Rep, 10*(1), 21036. doi:10.1038/s41598-020-78201-9

Bulut, G., Hong, S. H., Chen, K., Beauchamp, E. M., Rahim, S., Kosturko, G. W., . . . Uren, A. (2012). Small molecule inhibitors of ezrin inhibit the invasive phenotype of osteosarcoma cells. *Oncogene, 31*(3), 269-281. doi:10.1038/onc.2011.245

Burke, P., Schooler, K., & Wiley, H. S. (2001). Regulation of epidermal growth factor receptor signaling by endocytosis and intracellular trafficking. *Mol Biol Cell, 12*(6), 1897-1910. doi:10.1091/mbc.12.6.1897

Carotenuto, P., Romano, A., Barbato, A., Quadrano, P., Brillante, S., Volpe, M., . . . Franco, B. (2022). Targeting the MITF/APAF-1 axis as salvage therapy for MAPK inhibitors in resistant melanoma. *Cell Rep, 41*(6), 111601. doi:10.1016/j.celrep.2022.111601

Ceresa, B. P. (2012). Spatial regulation of epidermal growth factor receptor signaling by endocytosis. *Int J Mol Sci, 14*(1), 72-87. doi:10.3390/ijms14010072

Cha, B., Tse, M., Yun, C., Kovbasnjuk, O., Mohan, S., Hubbard, A., . . . Donowitz, M. (2006). The NHE3 juxtamembrane cytoplasmic domain directly binds ezrin: dual role in NHE3 trafficking and mobility in the brush border. *Mol Biol Cell, 17*(6), 2661-2673. doi:10.1091/mbc.e05-09-0843

Chirivino, D., Del Maestro, L., Formstecher, E., Hupe, P., Raposo, G., Louvard, D., & Arpin, M. (2011). The ERM proteins interact with the HOPS complex to regulate the maturation of endosomes. *Mol Biol Cell, 22*(3), 375-385. doi:10.1091/mbc.E10-09-0796

Coscoy, S., Waharte, F., Gautreau, A., Martin, M., Louvard, D., Mangeat, P., . . . Amblard, F. (2002). Molecular analysis of microscopic ezrin dynamics by two-photon FRAP. *Proc Natl Acad Sci U S A, 99*(20), 12813-12818. doi:10.1073/pnas.192084599

Di Malta, C., Zampelli, A., Granieri, L., Vilardo, C., De Cegli, R., Cinque, L., . . . Ballabio, A. (2023). TFEB and TFE3 drive kidney cystogenesis and tumorigenesis. *EMBO Mol Med, 15*(5), e16877. doi:10.15252/emmm.202216877

Dibble, C. C., & Cantley, L. C. (2015). Regulation of mTORC1 by PI3K signaling. *Trends Cell Biol, 25*(9), 545-555. doi:10.1016/j.tcb.2015.06.002

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R. (2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21. doi:10.1093/bioinformatics/bts635

Edgar, R., Domrachev, M., & Lash, A. E. (2002). Gene Expression Omnibus: NCBI gene expression and hybridization array data repository. *Nucleic Acids Res, 30*(1), 207-210. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/11752295

Gautreau, A., Poullet, P., Louvard, D., & Arpin, M. (1999). Ezrin, a plasma membrane-microfilament linker, signals cell survival through the phosphatidylinositol 3-kinase/Akt pathway. *Proc Natl Acad Sci U S A, 96*(13), 7300-7305. doi:10.1073/pnas.96.13.7300

Haddad, L. A., Smith, N., Bowser, M., Niida, Y., Murthy, V., Gonzalez-Agosti, C., & Ramesh, V. (2002). The TSC1 tumor suppressor hamartin interacts with neurofilament-L and possibly functions as a novel integrator of the neuronal cytoskeleton. *J Biol Chem, 277*(46), 44180-44186. doi:10.1074/jbc.M207211200

Huang da, W., Sherman, B. T., & Lempicki, R. A. (2009a). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Research, 37*(1), 1-13. doi:10.1093/nar/gkn923

Huang da, W., Sherman, B. T., & Lempicki, R. A. (2009b). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nature Protocols, 4*(1), 44-57. doi:10.1038/nprot.2008.211

Huang, J., Gu, S., Chen, M., Zhang, S. J., Jiang, Z., Chen, X., . . . Zhao, C. (2019). Abnormal mTORC1 signaling leads to retinal pigment epithelium degeneration. *Theranostics, 9*(4), 1170-1180. doi:10.7150/thno.26281

Iwamatsu, T. (2004). Stages of normal development in the medaka Oryzias latipes. *Mech Dev, 121*(7-8), 605-618. doi:10.1016/j.mod.2004.03.012

Kaur, G., & Lakkaraju, A. (2018). Early Endosome Morphology in Health and Disease. *Adv Exp Med Biol, 1074*, 335-343. doi:10.1007/978-3-319-75402-4_41

Kawaguchi, K., & Asano, S. (2022). Pathophysiological Roles of Actin-Binding Scaffold Protein, Ezrin. *Int J Mol Sci, 23*(6). doi:10.3390/ijms23063246

Kinoshita, M., Kani, S., Ozato, K., & Wakamatsu, Y. (2000). Activity of the medaka translation elongation factor 1alpha-A promoter examined using the GFP gene as a reporter. *Dev Growth Differ, 42*(5), 469-478. doi:10.1046/j.1440-169x.2000.00530.x

Krishnan, K., Bruce, B., Hewitt, S., Thomas, D., Khanna, C., & Helman, L. J. (2006). Ezrin mediates growth and survival in Ewing's sarcoma through the AKT/mTOR, but not the MAPK, signaling pathway. *Clin Exp Metastasis, 23*(3-4), 227-236. doi:10.1007/s10585-006-9033-y

Li, N., Kong, J., Lin, Z., Yang, Y., Jin, T., Xu, M., . . . Chen, L. (2019). Ezrin promotes breast cancer progression by modulating AKT signals. *Br J Cancer, 120*(7), 703-713. doi:10.1038/s41416-019-0383-z

Liu, B., Chen, H., Johns, T. G., & Neufeld, A. H. (2006). Epidermal growth factor receptor activation: an upstream signal for transition of quiescent astrocytes into reactive astrocytes after neural injury. *J Neurosci, 26*(28), 7532-7540. doi:10.1523/JNEUROSCI.1004-06.2006

Liu, S., Wang, Z., Zhu, R., Wang, F., Cheng, Y., & Liu, Y. (2021). Three Differential Expression Analysis Methods for RNA Sequencing: limma, EdgeR, DESeq2. *J Vis Exp*(175). doi:10.3791/62528

Mitchell, D. C., Kuljanin, M., Li, J., Van Vranken, J. G., Bulloch, N., Schweppe, D. K., . . . Gygi, S. P. (2023). A proteome-wide atlas of drug mechanism of action. *Nat Biotechnol, 41*(6), 845-857. doi:10.1038/s41587-022-01539-0

Muniz-Feliciano, L., Doggett, T. A., Zhou, Z., & Ferguson, T. A. (2017). RUBCN/rubicon and EGFR regulate lysosomal degradative processes in the retinal pigment epithelium (RPE) of the eye. *Autophagy, 13*(12), 2072-2085. doi:10.1080/15548627.2017.1380124

Naso, F., Intartaglia, D., Falanga, D., Soldati, C., Polishchuk, E., Giamundo, G., . . . Conte, I. (2020). Light-responsive microRNA miR-211 targets Ezrin to modulate lysosomal biogenesis and retinal cell clearance. *EMBO J, 39*(8), e102468. doi:10.15252/embj.2019102468

Nishimura, Y., Takiguchi, S., Ito, S., & Itoh, K. (2015). EGF-stimulated AKT activation is mediated by EGFR recycling via an early endocytic pathway in a gefitinib-resistant human lung cancer cell line. *Int J Oncol, 46*(4), 1721-1729. doi:10.3892/ijo.2015.2871

Ognibene, M., Vanni, C., Segalerba, D., Mancini, P., Merello, E., Torrisi, M. R., . . . Eva, A. (2011). The tumor suppressor hamartin enhances Dbl protein transforming activity through interaction with ezrin. *J Biol Chem, 286*(34), 29973-29983. doi:10.1074/jbc.M111.270785

Olsen, J. V., Blagoev, B., Gnad, F., Macek, B., Kumar, C., Mortensen, P., & Mann, M. (2006). Global, in vivo, and site-specific phosphorylation dynamics in signaling networks. *Cell, 127*(3), 635-648. doi:10.1016/j.cell.2006.09.026

Ong, S. E., Blagoev, B., Kratchmarova, I., Kristensen, D. B., Steen, H., Pandey, A., & Mann, M. (2002). Stable isotope labeling by amino acids in cell culture, SILAC, as a simple and accurate approach to expression proteomics. *Mol Cell Proteomics, 1*(5), 376-386. doi:10.1074/mcp.m200025-mcp200

Oughtred, R., Rust, J., Chang, C., Breitkreutz, B. J., Stark, C., Willems, A., . . . Tyers, M. (2021). The BioGRID database: A comprehensive biomedical resource of curated protein, genetic, and chemical interactions. *Protein Sci, 30*(1), 187-200. doi:10.1002/pro.3978

Palfy, M., Remenyi, A., & Korcsmaros, T. (2012). Endosomal crosstalk: meeting points for signaling pathways. *Trends Cell Biol, 22*(9), 447-456. doi:10.1016/j.tcb.2012.06.004

Parameswaran, N., Enyindah-Asonye, G., Bagheri, N., Shah, N. B., & Gupta, N. (2013). Spatial coupling of JNK activation to the B cell antigen receptor by tyrosine-phosphorylated ezrin. *J Immunol, 190*(5), 2017-2026. doi:10.4049/jimmunol.1201292

Petschnigg, J., Groisman, B., Kotlyar, M., Taipale, M., Zheng, Y., Kurat, C. F., . . . Stagljar, I. (2014). The mammalian-membrane two-hybrid assay (MaMTH) for probing membrane-protein interactions in human cells. *Nat Methods, 11*(5), 585-592. doi:10.1038/nmeth.2895

Pinelli, M., Carissimo, A., Cutillo, L., Lai, C. H., Mutarelli, M., Moretti, M. N., . . . di Bernardo, D. (2016). An atlas of gene expression and gene co-regulation in the human retina. *Nucleic Acids Res, 44*(12), 5773-5784. doi:10.1093/nar/gkw486

Polishchuk, R. S., & Polishchuk, E. V. (2019). From and to the Golgi - defining the Wilson disease protein road map. *FEBS Lett, 593*(17), 2341-2350. doi:10.1002/1873-3468.13575

Poupon, V., Stewart, A., Gray, S. R., Piper, R. C., & Luzio, J. P. (2003). The role of mVps18p in clustering, fusion, and intracellular localization of late endocytic organelles. *Mol Biol Cell, 14*(10), 4015-4027. doi:10.1091/mbc.e03-01-0040

Rosset, C., Netto, C. B. O., & Ashton-Prolla, P. (2017). TSC1 and TSC2 gene mutations and their implications for treatment in Tuberous Sclerosis Complex: a review. *Genet Mol Biol, 40*(1), 69-79. doi:10.1590/1678-4685-GMB-2015-0321

Salokas, K., Liu, X., Ohman, T., Chowdhury, I., Gawriyski, L., Keskitalo, S., & Varjosalo, M. (2022). Physical and functional interactome atlas of human receptor tyrosine kinases. *EMBO Rep, 23*(6), e54041. doi:10.15252/embr.202154041

Saygideger-Kont, Y., Minas, T. Z., Jones, H., Hour, S., Celik, H., Temel, I., . . . Uren, A. (2016). Ezrin Enhances EGFR Signaling and Modulates Erlotinib Sensitivity in Non-Small Cell Lung Cancer Cells. *Neoplasia, 18*(2), 111-120. doi:10.1016/j.neo.2016.01.002

Shang, P., Stepicheva, N., Teel, K., McCauley, A., Fitting, C. S., Hose, S., . . . Sinha, D. (2021). betaA3/A1-crystallin regulates apical polarity and EGFR endocytosis in retinal pigmented epithelial cells. *Commun Biol, 4*(1), 850. doi:10.1038/s42003-021-02386-6

Sorkin, A., & von Zastrow, M. (2009). Endocytosis and signalling: intertwining molecular networks. *Nat Rev Mol Cell Biol, 10*(9), 609-622. doi:10.1038/nrm2748

Stanasila, L., Abuin, L., Diviani, D., & Cotecchia, S. (2006). Ezrin directly interacts with the alpha1b-adrenergic receptor and plays a role in receptor recycling. *J Biol Chem, 281*(7), 4354-4363. doi:10.1074/jbc.M511989200

Tamma, G., Klussmann, E., Oehlke, J., Krause, E., Rosenthal, W., Svelto, M., & Valenti, G. (2005).
Actin remodeling requires ERM function to facilitate AQP2 apical targeting. *J Cell Sci, 118*(Pt
16), 3623-3630. doi:10.1242/jcs.02495
Tanaka, T., Zhou, Y., Ozawa, T., Okizono, R., Banba, A., Yamamura, T., . . . Sakurai, H. (2018).
Ligand-activated epidermal growth factor receptor (EGFR) signaling governs endocytic
trafficking of unliganded receptor monomers by non-canonical phosphorylation. *J Biol Chem,*
293(7), 2288-2301. doi:10.1074/jbc.M117.811299
Toops, K. A., Tan, L. X., & Lakkaraju, A. (2014). A detailed three-step protocol for live imaging of
intracellular traffic in polarized primary porcine RPE monolayers. *Exp Eye Res, 124*, 74-85.
doi:10.1016/j.exer.2014.05.003
Tsai, F. C., Bertin, A., Bousquet, H., Manzi, J., Senju, Y., Tsai, M. C., . . . Bassereau, P. (2018). Ezrin
enrichment on curved membranes requires a specific conformation or interaction with a
curvature-sensitive partner. *Elife, 7*. doi:10.7554/eLife.37262
Vogel, G. F., Ebner, H. L., de Araujo, M. E., Schmiedinger, T., Eiter, O., Pircher, H., . . . Hess, M. W.
(2015). Ultrastructural Morphometry Points to a New Role for LAMTOR2 in Regulating the
Endo/Lysosomal System. *Traffic, 16*(6), 617-634. doi:10.1111/tra.12271
Wan, X., Mendoza, A., Khanna, C., & Helman, L. J. (2005). Rapamycin inhibits ezrin-mediated
metastatic behavior in a murine model of osteosarcoma. *Cancer Res, 65*(6), 2406-2411.
doi:10.1158/0008-5472.CAN-04-3135
Wang, Y., Pennock, S., Chen, X., & Wang, Z. (2002). Endosomal signaling of epidermal growth factor
receptor stimulates signal transduction pathways leading to cell survival. *Mol Cell Biol, 22*(20),
7279-7290. doi:10.1128/MCB.22.20.7279-7290.2002
Warde-Farley, D., Donaldson, S. L., Comes, O., Zuberi, K., Badrawi, R., Chao, P., . . . Morris, Q.
(2010). The GeneMANIA prediction server: biological network integration for gene
prioritization and predicting gene function. *Nucleic Acids Res, 38*(Web Server issue), W214-
220. doi:10.1093/nar/gkq537
Wee, P., & Wang, Z. (2017). Epidermal Growth Factor Receptor Cell Proliferation Signaling
Pathways. *Cancers (Basel), 9*(5). doi:10.3390/cancers9050052
Zhao, H., Shiue, H., Palkon, S., Wang, Y., Cullinan, P., Burkhardt, J. K., . . . Turner, J. R. (2004). Ezrin
regulates NHE3 translocation and activation after Na+-glucose cotransport. *Proc Natl Acad Sci*
U S A, 101(25), 9485-9490. doi:10.1073/pnas.0308400101
Zhou, R., Cao, X., Watson, C., Miao, Y., Guo, Z., Forte, J. G., & Yao, X. (2003). Characterization of
protein kinase A-mediated phosphorylation of ezrin in gastric parietal cell activation. *J Biol*
Chem, 278(37), 35651-35659. doi:10.1074/jbc.M303416200
Zhou, R., Zhu, L., Kodani, A., Hauser, P., Yao, X., & Forte, J. G. (2005). Phosphorylation of ezrin on
threonine 567 produces a change in secretory phenotype and repolarizes the gastric parietal cell.
J Cell Sci, 118(Pt 19), 4381-4391. doi:10.1242/jcs.02559

Figure legends

Figure 1. **Deletion of Ezrin increases the lysosomal pathway. A,** Gene Ontology of 530 genes

differentially expressed at mRNA and protein levels (EZR[KO] versus WT). Bubble plot representing

some of the most enriched GO terms regarding cellular components. Color and x axis represent minus

logarithms of p Value. The size represents numbers of genes enriched in a GO term. **B**, WT and

EZR^{KO} MEF cells were cultured in 6-cm cell plates for 24 hours, then fixed and immunostained with

lysotracker and DAPI. Scale bar 10 μm. **C**, Data represent mean of lysotracker-positive cells ± SEM

(n=3 experiments at least). Statistical test: unpaired t-test. **D**, MEF cells WT and Ezr^{KO} were cultured in

6-well plates for 24 hours, then fixed and immunostained with LAMP1 and LC3 antibodies and DAPI.

Scale bar 10 μm (magnification 1 μm). **E**, Data represent mean of LAMP1-LC3 colocalization spots ±

SEM (n=3 experiments at least). Statistical test: unpaired t-test. **F,** Model showing autophagic flux

induction in EZR^{-/-} cells. **G**, MEF Ezr^{KO} showed CTSB enzymatic activity increase compared to control

cells. **H**, MEF cells WT and Ezr^{KO} were lysed and immunoblotted with NBR1, LAMP1, EZR, P62,

LC3 antibodies or GAPDH antibodies as a loading control. The graphs show the mean NBR1, LAMP1,

EZR, P62, LC3 levels relative to GAPDH ± SEM (n=3 experiments at least). Statistical test: unpaired t-

test for NBR1, P62, LC3-I; Welch's t-test for CTSD, LC3-II; Mann-Whitney test for LAMP1.

Figure 2. Ezrin binds EGFR and regulate its activation. A, Bubble plot representing the enrichment

analysis of 530 DEGs performed in SILAC Phosphoproteomics data. Color and x axis represent minus

logarithms of p Value. The size represents numbers of genes enriched in the indicated data. **B**, Physical

interactions, obtained by GeneMANIA, highlight Ezrin and EGFR binding. **C**, Volcano plot of DEGs,

with up-regulated EGFR and down-regulated MAP2 and ERBB2 (no threshold on Log2FC and 0.05

threshold on -Log10FDR). Legend: red dot, up-regulated gene; blue dot, down-regulated gene; grey

dot, not significant gene. **D,** Co-IP data for Ezrin-EGFR interaction. For the co-IP analyses, was used

Ezrin antibody, conjugated with beads, and immunoblotted with EGFR antibody for WT and EZR^{-/-}

(left) and HeLa EZR^{T567D} and EZR^{T567A} (right) HeLa cells, respectively. Schematic representation of

HeLa EZR^{T567D} and EZR^{T567A} co-IP (bottom). **E**, Confocal microscopy images showing EGFR (green)

and EZR (red) co-localization on the membrane in HeLa WT cells (left) and magnified views of the

regions are provided (right). Scale bar 10 μm (magnification 1 μm). Representative plots of co-

localization profiles on the membrane between EGFR (green) and EZR (red). **F**, Immunofluorescent labelling images of EGFR in HeLa WT and EZR$^{-/-}$ cells, observed by confocal microscopy. Scale bar 10 μm. **G**, Immunoblots and calculated levels (bottom) of HER3, pY845 EGFR, EGFR, pT222 MK2, MK2, pT180/pY182 p38 MAPK and P38 MAPK in HeLa WT and EZR$^{-/-}$ cells. Data are expressed as mean of pY845EGFR/EGFR, pT222 MK2/MK2 and pT180/pY182 p38 MAPK/P38 MAPK ratio ± SEM (n=3 experiments at least). GAPDH was used as loading control. Statistical test: unpaired t-test for pY845 EGFR; Mann-Whitney test for HER3, EGFR, pT222 MK2, pT180/pY182 p38 MAPK.

Figure 3. Ezrin controls EGFR localization. A, Western blot analysis of chemical crosslinked EGFR in HeLa WT and EZR$^{-/-}$ with (+) and without (-) EGF stimulation. Arrowheads indicate detected signals of dimeric and monomeric form of EGFR. **B**, Model showing the crosslinking effect of EGFR dimer formation in HeLa WT and EZR$^{-/-}$. **C**, Immunofluorescence images of EGFR (green) and EEA1 (red) in HeLa WT (top) and EZR$^{-/-}$ (bottom) observed by confocal microscopy. Scale bar 10 μm (magnification 1 μm). Representative plots of co-localization profiles of EGFR in early endosome. HeLa EZR$^{-/-}$ cells do not show EGFR and EEA1 co-localization compared to control cells. **D**, Representative immunoblots of EGFR in membrane (top) and endosomes (bottom) proteins in HeLa WT and EZR$^{-/-}$. ZO-1 and EEA1 are used as membrane and endosomes extraction control, respectively. GAPDH are used as loading control. **E**, Schematic translocation of EGFR in the endosomes in HeLa WT compared to HeLa EZR$^{-/-}$. **F**, HeLa cells were fixed and immunostained with endotracker and EEA1 (red) and DAPI (blue). Scale bar 10 μm (magnification 1 μm). **G**, Graph shows mean of endotracker-positive cells ± SEM (n=3 experiments at least). Statistical test: unpaired t-test.

Figure 4. EGFR migrates on the endosomes depending on Ezrin. A, Immunofluorescence labelling images of EGFR-GFP (green), EEA1 (red) and DAPI (blue) after 3h of EGF stimulation (right) in

HeLa WT (top) and EZR$^{-/-}$ (bottom). Magnified views of the regions in the boxes are provided in both

Airyscan high-resolution microscopy and 3D-confocal microscopy. **B**, EGFR and EEA1 co-localization

is expressed as a representative plot in HeLa WT (top) and EZR$^{-/-}$ (bottom). Scale bar 10 µm

(magnification 1 µm).

Figure 5. EGF stimulation does not affect on EGFR in absence of Ezrin. A, Live cell imaging and

model for EGFR (green) translocation from the membrane to the endosomes in HeLa WT (top) and

EZR$^{-/-}$ (bottom) cells without EGF stimulation (T0) and with a progressive EGF stimulation (from

T10'' to T60''). White boxes are magnifications that depict EGFR protein migration. Scale bar 1 µm).

Please refer to Video 1. **B**, IEM (anti-GFP immunolabelling) of cycloheximide treated HeLa WT, WT

+ EGF, EZR$^{-/-}$ and EZR$^{-/-}$ + EGF cells expressing EGFR-GFP. Endosomes containing EGFR is shown

in green. Scale bar 200 nm. Quantitative analysis (right) of EGFR positive endosomes expressed as

mean ± SEM. Statistical test: generalized Linear Model with Likelihood Ratio (Poisson Regression). **C**,

immunoblots and calculated levels (bottom) of HER2, pY845 EGFR, EGFR, pT180/pY182 p38 MAPK

and P38 MAPK in HeLa WT and EZR$^{-/-}$ cells with (+) and without (-) EGF stimulation. Data are

expressed as mean of pY845EGFR/EGFR and pT180/pY182 p38 MAPK/P38 MAPK ratio ± SEM

(n=3 experiments at least). GAPDH was used as loading control. Statistical test: Unpaired t-test for

HER2 WT, HER2 EZR$^{-/-}$, pY845 EGFR EZR$^{-/-}$, EGFR WT, EGFR EZR$^{-/-}$, pT180/pY182 p38 MAPK

WT, pT180/pY182 p38 MAPK EZR$^{-/-}$; unpaired t-test with Welch's correction for pY845 EGFR WT.

D, Representative immunoblots of EGFR in membrane (top) and endosomes (bottom) proteins in HeLa

WT and EZR$^{-/-}$ with (+) and without (-) EGF stimulation. ZO-1 and EEA1 are used as membrane and

endosomes extraction control, respectively. GAPDH are used as loading control.

Figure 6. EGFR-Ezrin complex interacts with TSC1. A, co-IP analysis for Ezr-TSC1 (left) and EGFR-TSC1 (right) interaction. For co-IP analyses, Ezrin (left) and EGFR (right) antibodies were used. The proteins immunoprecipitated were blotted for TSC1 and AKT antibodies in HeLa WT and EZR[-/-]. **B**, HeLa WT and EZR[-/-] cells were lysed and immunoblotted with pS939 TSC2, TSC2, PT389 P70 S6 Kinase, P70 S6 Kinase, pS473 AKT, AKT, pS65 4E-BP1, 4E-BP1 and GAPDH as a loading control. Data represent the mean of pS939 TSC2/TSC2, T389 P70 S6 Kinase/P70 S6 Kinase, pS473 AKT/AKT and pS65 4E-BP1/4E-BP1 ratio ± SEM (n=3 experiments at least). Statistical test: unpaired t-test for pT389 P70 S6 Kinase, pS473 AKT; Unpaired t-test with Welch's correction for pS939 TSC2; Mann-Whitney test for pS939 TSC2. **C**, Representative confocal images of LAMP1 and TSC1 (left) and LAMP1 and TSC2 (right) immunofluorescence in HeLa WT and EZR[-/-] cells. Magnified insets of TSC1/2 localization are shown. Scale bar 10 μm (magnification 1 μm). **D**, Data represent mean of LAMP1-TSC1 (left) and LAMP1-TSC2 (right) colocalization spots ± SEM (n=3 experiments at least). Statistical test: unpaired t-test for LAMP1-TSC1; Unpaired t-test with Welch's correction for LAMP1-TSC2. **E**, pS473 AKT western blotting with EGF time course in HeLa WT (left) and EZR[-/-] (right) cells. Graph shows the mean of pS473 AKT/AKT ratio ± SEM (n=3 experiments at least). Statistical test: One Way ANOVA with Dunnett's post-hoc test for WT curve; Kruskal-Wallis test with Dunn's post-hoc test for KO curve (pairwise comparisons with reference T0). NS: not significant. **F**, HeLa WT, WT + EGF, EZR[-/-] and EZR[-/-] + EGF cells were immunostained with EGFR (green), TSC1 (red) and EEA1 (gray). Representative magnifications are shown. Scale bar 10 μm (magnification 1 μm).

Figure 7. *Ezrin* depletion induces EGFR-mediated retinal degeneration. A, Schematic representation of used CRISPR/Cas9 strategy to generate Ezrin[-/-] medaka lines. The red box highlighted the deleted nucleotides in the *Ezrin* exon 1 gene. **B**, WT and Ezrin[-/-] medaka proteins were immunoblotted with Ezrin antibody and Actin as a loading control. **C**, Stereo-microscopic

representative images of WT and Ezrin[-/-] medaka at stage 40. Scale bar 1 mm. **D**, Immunoblots and

calculated levels (right) of pT1462 Tsc2, pS473 Akt, LC3-I, LC3-II, pY845 Egfr and Egfr in WT and

Ezrin[-/-] medaka fish. Data are expressed as mean of pT1462 Tsc2/Tsc2, pS473 Akt/Akt and

pY845EGFR/EGFR ratio ± SEM (n=3 experiments at least). Actin was used as loading control.

Statistical test: Unpaired t-test. **E**, Representative confocal images of LAMP1 immunofluorescence in

WT and Ezrin[-/-] medaka fish. Magnified insets of RPE LAMP1 localization are shown. Scale bar 10

µm. RPE: retinal pigment epithelium; ONL: outer nuclear layer; INL: inner nuclear layer. **F**, Medaka

WT and Ezrin[-/-] fish were immunostained with EGFR. Scale bar 10 µm. **G**, Immunofluorescence

labelling images of RHO (left) and ZPR1 (right) in WT and Ezrin[-/-] fish. Magnified views of the

regions in the boxes are provided in bottom. Scale bar 10 µm.

Figure 8. EGFR/Ezrin/TSC molecular pathway

Diurnal inactivation of Ezrin lead to incapacity of EGFR to dimerize. The absence of active EGFR on

the endosome causes the migration of TSC complex on the lysosome, where it inhibits mTORC1. On

the contrary, the nocturnal activation of Ezrin favorites the phosphorylation and dimerization of EGFR,

that translocate from the plasma membrane on the early endosome. Ezrin localized with EGFR, on the

endosomal membrane, binds TSC complex, preventing mTORC1 inactivation on the lysosome.

Figure supplement 1. **Ezrin knockout cells show lysosomal enhancement. A,** VENN diagram

showing the comparison of transcriptomics (GSE195983) and proteomics (PRIDE ID: PXD045157)

performed in EZR[KO] MEF cells. 572 genes are commonly regulated: 317 and 213 genes are induced

and inhibited in both datasets, respectively; the remaining 42 are regulated in opposite manner.

Heatmap on 22 Lysosomal genes (out of 317) induced in the transcriptome **B** and in the proteome **C**. **D,**

Schematic representation of *Ezrin* gene and the corresponding protein. The position of the mutation in

1001 exon 2 and the relative changes in the coding sequence is highlighted. **E,** WT and EZR$^{-/-}$ HeLa cells

1002 were cultured in 6-cm cell plates for 24 hours, then fixed and immunostained with lysotracker and

1003 DAPI. Scale bar 10 μm. **F,** Data represent mean of lysotracker-positive cells ± SEM (n=3 experiments

1004 at least). Statistical test: unpaired t-test. **G,** HeLa WT and EZR$^{-/-}$ cells were lysed and immunoblotted

1005 with NBR1, LAMP1, EZR, p62, LC3 antibodies or GAPDH antibody as a loading control. Data

1006 represent the mean relative NBR1, LAMP1, p62 and LC3 levels relative to GAPDH ± SEM (n=3

1007 experiments at least). Statistical test: unpaired t-test for NBR1, p62, LC3-I and LC3-II; Mann-Whitney

1008 test for LAMP1. **H,** HeLa EZR$^{-/-}$ showed CTSB enzymatic activity increase compared to control cells.

1009 Statistical test: unpaired t-test. **I,** HeLa WT and EZR$^{-/-}$ cells were cultured in normal medium (stv-, baf-

1010 ). Starved HBSS medium (stv+), or starved medium supplemented with bafilomycin (baf+) or without

1011 bafilomycin (baf-), were lysed and immunoblotted with LC3 antibodies or GAPDH antibody as a

1012 loading control. Data represent the mean relative LC3 levels relative to GAPDH ± SEM (n=3

1013 experiments at least). Statistical test: unpaired t-test for all the comparisons against WT; Mann-

1014 Whitney test for all the comparisons against EZR$^{-/-}$.

1015

1016 **Figure supplement 2**. **E**zrin genetic and pharmacological depletion causes EGFR signaling

1017 **alteration. A** and **B,** Bubble plots representing the enrichment analysis of 530 DEGs performed in

1018 kinase perturbation from GEO database and from the Proteomics Drug Atlas. Color and x axis

1019 represent minus logarithms of p Value. The size represents numbers of genes enriched in the indicated

1020 data. **C,** MEF cells were fixed and immunostained with endotracker (red) and DAPI (blue). Scale bar

1021 10 μm. **D,** Graph shows mean of endotracker-positive cells ± SEM (n=3 experiments at least).

1022 Statistical test: unpaired t-test. **E,** Immunoblots and calculated levels (bottom) of HER2, HER3,

1023 EGFR, pT222 MK2, MK2, pT180/pY182 p38 MAPK and P38 MAPK in MEF WT and EZR^{KO} cells.

1024 Data are expressed as mean of HER2, HER3, EGFR, pT222 MK2/MK2 and pT180/pY182 p38 MAPK/

p38 MAPK ratio ± SEM (n=3 experiments at least). GAPDH was used as loading control. Statistical

test: unpaired t-test for HER3 and pT180/pY182 p38 MAPK; unpaired t-test with Welch's correction

for EGFR; Mann-Whitney test for HER2 and pT222 MK2. **F,** Immunoblots and calculated levels

(bottom) of HER2, HER3, pY845 EGFR, EGFR, pT222 MK2, MK2, pT180/pY182 p38 MAPK and

P38 MAPK in MEF WT and EZR^{KO} cells. Data are expressed as mean of HER2, HER3, pY845 EGFR,

EGFR, pT222 MK2 and pT180/pY182 p38 MAPK ratio ± SEM (n=3 experiments at least). GAPDH

was used as loading control. Data are expressed as mean of HER2, HER3, pY845EGFR/EGFR, EGFR,

pT222 MK2/MK2 and pT180/pY182 p38 MAPK/P38 MAPK ratio ± SEM (n=3 experiments at least).

Statistical test: One Way ANOVA with Tukey's post-hoc test for HER2, EGFR, pT222 MK2 and

pT180/pY182 p38 MAPK; Kruskal-Wallis test with Dunn's post-hoc for pY845 EGFR. **G,**

Immunofluorescence labelling images of EGFR-GFP (green), EEA1 (red) and DAPI (blue) after 3h of

EGF stimulation (bottom) in HeLa WT (left) and *NSC668394*-treated cells (right). Scale bar 10 μm

(magnification 1 μm).

1038

Figure supplement 3. **Ezrin inhibition induces mTORC1 pathway inhibition**. **A,** Immunoblots and

calculated levels (bottom) of pS939 TSC2, TSC2, pT389 P70 S6 Kinase, P70 S6 Kinase, pS473 AKT,

AKT, pS65 4E-BP1, 4E-BP1 in HeLa WT, DMSO-treated and *NSC668394*-treated cells. Data are

expressed as mean of pS939 TSC2/TSC2, pT389 P70 S6 Kinase/P70 S6 Kinase, pS473 AKT/ AKT

and pS65 4E-BP1/4E-BP1 ratio ± SEM (n=3 experiments at least). GAPDH was used as loading

control. Statistical test: One Way ANOVA with Tukey's post-hoc test for pS939 TSC2, pS473 AKT,

pS65 4E-BP1; Kruskal-Wallis test with Dunn's post-hoc test for pT389 P70 S6 Kinase. **B,**

Immunoblots and calculated levels (bottom) of pT1462 TSC2, TSC2, pT389 P70 S6 Kinase, P70 S6

Kinase, pS473 AKT, AKT, pS65 4E-BP1 and 4E-BP1 in HeLa WT, DMSO-treated and *NSC668394-*

treated cells. Data are expressed as mean of pT1462 TSC2/TSC2, pT389 P70 S6 Kinase/P70 S6

Kinase, pS473 AKT/ AKT and pS65 4E-BP1/4E-BP1 ratio ± SEM (n=3 experiments at least). GAPDH

was used as loading control. Statistical test: One Way ANOVA with Tukey's post-hoc test for pT389

P70 S6 Kinase, pS473 AKT, pS65 4E-BP1; Kruskal-Wallis test with Dunn's post-hoc test for pT1462

TSC2. **C,** Immunofluorescence labelling images of LAMP1 (red), TSC1/TSC2 (green) and DAPI

(blue) in MEF WT (left) and *NSC668394*-treated cells. Scale bar 10 μm (magnification 1 μm). **D,**

Immunoblots and calculated levels (bottom) of pS939 TSC2, TSC2, pT389 P70 S6 Kinase, P70 S6

Kinase, pS473 AKT and AKT in MEF WT and EZR^{KO} cells. Data are expressed as mean of pS939

TSC2/TSC2, pT389 P70 S6 Kinase/P70 S6 Kinase and pS473 AKT/ AKT ratio ± SEM (n=3

experiments at least). GAPDH was used as loading control. Statistical test: unpaired t-test. **E,**

Immunofluorescence labelling images of LAMP1 (red), TSC1 (green) and DAPI (blue) in MEF WT

and EZR^{KO} cells. Scale bar 10 μm (magnification 1 μm).

1060

Figure supplement 4. Ezrin overexpression rescue EGFR and TSC1 localization. A,

Immunofluorescence labelling images of EGFR-GFP (green), EZR^{T567D}-CHE (top), EZR^{T567A}-CHE

(bottom), EEA1 (gray) and DAPI (blue) in HeLa EZR$^{-/-}$. EZR^{T567D}, but not EZR^{T567A}, with EGF

stimulation rescued EGFR translocation on the endosomes in EZR$^{-/-}$ cells. Magnified views of EGFR

and Ezrin localization are provided in both Airyscan high-resolution microscopy and 3D-confocal

microscopy. Scale bar 10 μm (magnification 1 μm). **B**, Representative confocal images of EGFR-GFP

(green), EZR^{T567D}-CHE (top), EZR^{T567A}-CHE (bottom), TSC1 (gray) and DAPI (blue) in HeLa EZR$^{-/-}$.

TSC1 localization misplaced from lysosomes to endosomes in HeLa EZR$^{-/-}$ cells, after

EZR^{T567D} transfection and upon EGF stimulation. Boxes show magnifications of TSC1 position,

displayed both as Airyscan high-resolution microscopy and 3D-confocal microscopy. Scale bar 10 μm

(magnification 1 μm).

1072

Figure supplement 5. **Light/dark transitions regulate EGFR and mTOR1 signaling in mice RPE.**

A, WT mice were exposed 3h in light on (left) and dark (right) condition and after 3 h the retina were

analyzed by EGFR immunofluorescence. Representative images are shown. Scale bar 50 μm.

Schematic summary of EGFR and Ezrin trend in mice retina. **B**, WT mice RPE were analyzed by

western blot for EGFR pathway in light and dark conditions. Data are expressed as mean of pY845

EGFR/EGFR and pT180/pY182 p38 MAPK/P38 MAPK ratio ± SEM (n=3 experiments at least).

GAPDH was used as loading control. Statistical test: unpaired t-test for EGFR, pY845 EGFR, and

pT180/pY182 p38 MAPK. **C,** Immunoblots and calculated levels (right) of pS939 TSC2, TSC2, pT389

P70 S6 Kinase, P70 S6 Kinase, pS473 AKT, AKT, pT37/46 4E-BP1 and 4E-BP1 in light and dark-

exposed WT mice. Data are expressed as mean of pS939 TSC2/TSC2, pT389 P70 S6 Kinase/P70 S6

Kinase, pS473 AKT/ AKT and pT37/46 4E-BP1/4E-BP1 ratio ± SEM (n=3 experiments at least).

GAPDH was used as loading control. Statistical test: unpaired t-test for pS939 TSC2, pS473 AKT;

Unpaired t-test with Welch's correction for pT389 P70 S6 Kinase; Mann-Whitney test for pT37/46 4E-

BP1.

1087

Video 1

WT HeLa cells expressing EGFR-GFP were imaged by TIRF super-resolution microscopy every were

imaged every 0.5 s for 5 min after EGF stimulation (related to Figure 5A).

Video 2

Magnification from WT HeLa cells expressing EGFR-GFP, imaged by TIRF super-resolution

microscopy every were imaged every 0.5 s for 5 min after EGF stimulation (related to Figure 5A).

Video 3

EZR$^{-/-}$ HeLa cells expressing EGFR-GFP were imaged by TIRF super-resolution microscopy every

were imaged every 0.5 s for 5 min after EGF stimulation (related to Figure 5A).

Video 4

Magnification from EZR$^{-/-}$ HeLa cells expressing EGFR-GFP, imaged by TIRF super-resolution

microscopy every were imaged every 0.5 s for 5 min after EGF stimulation (related to Figure 5A).

a

Ex.1

sgRNA1 ΔEx1 sgRNA2

Wild type

Ex.1

TTCTTTTGTGTTGCATG - - - 370bp - - - GGCGCGCGGTAGCCGCTTT

Ezrin⁻/⁻

TTCTTTTGTGAGCCGCTT

b

	WT	Ezrin⁻/⁻	kDa
Ezr			80
Act			45

c

Wild type

Ezrin⁻/⁻

d

	WT	Ezrin⁻/⁻	kDa
pT1462 Tsc2			200
Tsc2			200
pS473 Akt			65
Akt			60
LC3-I			19
LC3-II			17
pY845 Egfr			175
Egfr			175
Act			45

pT1462 Tsc2 P=0.00041

pS473 Akt P=0.023

LC3-I P=0.0040

LC3-II P=0.0019

pY845 Egfr P=0.0018

Egfr P=0.0040

Relative levels

e

WT LAMP1

Ezrin⁻/⁻ LAMP1

RPE ONL INL

RPE

f

WT EGFR

Ezrin⁻/⁻ EGFR

g

WT RHO

Ezrin⁻/⁻ RHO

WT ZPR1

Ezrin⁻/⁻ ZPR1