Protection of savings by reducing the salience of opposing errors

Mousa Javidialsaadi[1], Scott T. Albert[2], Badr Moufarrej S Al Mutairi[1], Jinsung Wang[1*]

[1]Kinesiology programs, Zilber College of Public Health, University of Wisconsin–Milwaukee,

Milwaukee, WI, 53201

[2]Neuroscience Center, University of North Carolina at Chapel Hill – Chapel Hill, NC 27514

Correspondence: Jinsung Wang, 492 Enderis Hall, University of Wisconsin – Milwaukee, WI

53201, USA. Email wang34@uwm.edu. Phone: 414-229-2645

Key words: Gradual adaptation; Visual rotation; Washout; Unlearning; Implicit learning;

Readaptation

Abstract

When humans encounter the same disturbance twice, they adapt to it faster during the second

exposure. To examine how subconscious learning systems contribute to this savings process,

previous studies have suppressed explicit awareness of the perturbation by gradually increasing

its magnitude during initial learning. This has produced mixed effects, with some studies

demonstrating faster relearning, and others observing no acceleration during relearning. Here

we examined whether these differences might be due to the nature of a de-adaptation period

that separates two learning periods. To test this idea, we manipulated the magnitude of washout

errors by de-adapting participants abruptly, gradually, or by removing feedback entirely.

Empirical analyses indicated that the different classes of washout errors had a profound effect

on savings: large washout errors nullified the ability to save, whereas small errors or the absence

of error protected savings. Model-based analyses suggested that changes in learning rates were

mediated by an increase in sensitivity to error that could be reversed by experience with

oppositely-oriented washout errors. This suggests that the experience of error produces both a

facilitation of learning for similar errors and a reduction in learning for dissimilar errors. The latter

can abolish the expression of savings following gradual adaptation.

Introduction

When an unexpected perturbation to our motor system occurs, the resulting errors in our actions and predictions[1–5] update our motor commands, predictively countering future disturbances. The removal of a perturbation causes behavior to rapidly revert to its initial state, but a lingering memory of our initial encounter is stored within the nervous system, whereby re-exposure to the perturbation results in more rapid adaptation. This learning hallmark, termed savings[6], can occur on the same day, overnight[7], or even a year after the initial disturbance[8], and is ubiquitous across effectors including saccade adaptation[9], locomotion[10,11], visuomotor adaptation[7,12,13], force field adaptation[14–16], and conditioning paradigms[17]. What components of adaptation experiments are necessary to produce savings during re-learning phases?

Our ability to flexibly increase our learning rate is known to rely on two parallel adaptive systems: an explicit process directed by our conscious strategies[18–20], and a subconscious implicit process that operates without our awareness[2,4]. The implicit and explicit contributions to savings have been typically studied using visuomotor rotations, a paradigm that can provide an opportunity to parse adapted responses into a re-aiming direction (*i.e.*, the explicit strategy) and an additional uncontrolled adjustment (*i.e.*, the implicit correction)[18].

In visuomotor adaptation, improvements in strategy use are salient in protocols where a rotation is abruptly initiated during two separate exposures. Explicit savings can be documented by quantifying strategy use via reaction time manipulations[7,21], or by directly querying subjects[12]. When conditions are applied to suppress explicit strategies, such as limiting response time[22], or minimizing errors experienced during initial learning[23], implicit savings can be isolated. This latter minimization approach is performed by gradually introducing a perturbation, thus maintaining a

small discrepancy between the disturbance and motor action that produces smaller error sizes

that go undetected, limiting strategy use[23].

Intriguingly, however, studies that have employed gradual perturbations have observed

mixed effects on the ability of *gradual* learning to facilitate more rapid *abrupt* learning. Absence

of savings following gradual learning[24] is consistent with a memory of errors model[24,25], in which

gradual adaptation cannot improve abrupt adaptation due to a mismatch in error size: errors are

large in the abrupt case, but small in the gradual case. Thus, to the brain, an abrupt perturbation

following gradual adaptation may appear to be a novel context, because large errors it

experiences never occurred in the past. While this prediction is consistent with at least one prior

study[24], more recent work has exhibited opposing results where gradual adaptations improve

abrupt learning rates[23,26], particularly through an acceleration in implicit learning[23]. How is it that

gradual perturbations result in savings in some experiments, but not in others?

Here we consider the possibility that typical sensorimotor learning paradigms may fail to

produce savings not because of differences in the way individuals learn, but rather by the way

they forget. That is, savings protocols require not only two learning periods, but also the decay

of the adapted behavior in between the two learning phases. This forgetting is achieved in a

washout period in which the perturbation is removed to return the adapted state to its baseline

level. The conditions used to achieve this washout, however, vary across studies. In some cases,

washout consists of abrupt cessation of the perturbation, and lasts the same number of trials as

the initial exposure period[24]. On the other hand, some savings experiments employ much shorter

washout phases[23,26] that are preceded by other conditions such as invariant error-clamp[26] or no-

feedback[23] trials.

Critically, these differences in the washout period alter the size of errors experienced by

the subject. Could it be that *larger* washout errors experienced over *longer* durations reduce the

ability to save? This idea seems plausible, given that washout produces errors that are opposite

in sign to the errors experienced during adaptation phases, thus necessitating a learning response

in the opposite direction. Here we examine the overlooked possibility that the nature of washout

plays a central role in the expression of savings. Across three groups, we manipulate the types of

washout errors by (1) removing errors entirely by eliminating visual feedback, (2) limiting the size

of errors by de-adapting subjects gradually, or (3) facilitating large errors by abruptly disengaging

a visuomotor rotation. We then apply empirical and model-based approaches to assess whether

some washout conditions more easily produce savings during abrupt rotation periods that follow

initial gradual learning phases. These experiments seek to not only clarify previous discrepancies

in the literature, but also to expand our current models of savings in the implicit learning system.

Materials & Methods

Participants

40 healthy volunteers (18-35 years old; 18 females) participated in this study. All subjects were

right-handed as assessed by the Edinburgh handedness inventory (Oldfield, 1971). Informed

consent was given by each participant, and consent forms were approved by the University of

Wisconsin-Milwaukee Institutional Review Board. Subjects were randomly assigned to 1 of 4

groups: 3 gradual rotation groups and 1 control group.

Apparatus

Participants were situated in a robotic exoskeleton (KIMARM, BKIN Technologies Ltd, Kingston,

ON, Canada) that provided gravitational support to the right arm (all subjects used this arm in

the present study). The exoskeleton was positioned so that the arm was hidden underneath a

horizontal display (Fig. 1A). To track the hand's position, a small cursor was projected onto the

display, over the subject's index fingertip. During each trial, the KINARM projected visual stimuli

onto the display, so that they appeared in the same plane as the arm. The visual stimuli consisted

of a centrally located start circle (2 cm in diameter) and 1 of 4 target circles (2 cm in diameter)

located 10 cm away from the start target (Fig. 1A). We sampled the hand's position in the x-y

plane at 1000 Hz. Position data were low-pass filtered at 15 Hz, and then differentiated to

calculate velocity. Post-processing, analysis, and modeling were conducted in MATLAB R2018a

(The MathWorks Inc., Natick, MA).

Experimental Design

Participants were assigned to 1 of 3 experimental groups (n=10/group). Each experimental group

experienced 4 experiment phases: (1) a baseline reaching period, (2) a gradual adaptation period,

(3) a washout period, and (4) an abrupt readaptation period (Fig. 1B). During the gradual learning

period, a rotation was introduced that slowly increased in magnitude (up to 24°). In the abrupt

readaptation period, the rotation immediately attained its maximum value. To determine how

gradual learning and washout type altered reaching behavior during the final abrupt learning

phase, we compared behavior to a control group (n=10) that completed the same 4-phase

adaptation paradigm, with the only exception being that the perturbation was introduced

abruptly, not gradually, in the initial adaptation period.

In the baseline period (Fig. 1B, baseline), participants moved their arm to each of the four

targets over a 10-cycle reaching period. On each trial, continuous visual feedback was provided

via a cursor indicating the location of the index fingertip. Participants were instructed to move

rapidly and accurately to the target location. The trial ended 1.5 sec after target presentation. To

begin the next trial, the participants brought their hand back to the central start position.

Throughout the entire experiment, the four targets were presented in a pseudorandom order in

4-trial cycles (i.e., all four targets presented once in a cycle).

During the gradual adaptation period, participants were exposed to a visuomotor rotation

that systematically increased in size throughout the session. It started at 0° for 9 cycles, increased

by 1° (i.e., visual display of the cursor rotated 1° counterclockwise (CCW) about the start circle)

for next 23 cycles, and remained at 24° for last 8 cycles (Fig. 2A, gradual rotation). We provided

visual feedback on all trials. Following this session, participants underwent a washout period. This

washout period was the critical manipulation in our study and differed across all 3 gradual

learning groups. One group experienced a no-feedback condition in which the cursor was

removed during movement execution. Another group experienced abrupt washout in which

visual feedback was given and the rotation was immediately set to 0° (*i.e.*, the baseline

condition). Last, a third group experienced gradual de-adaptation: the rotation schedule was

identical to the initial gradual adaptation sequence, but in the reverse order (9 cycles 24°, 23

cycles in which the rotation decreased by 1°, 8 cycles with 0° rotation). Note that the control

group, that initially experienced an abrupt rotation condition, de-adapted in the no-feedback

condition in between rotation exposures.

The experiment ended with an abrupt readaptation period that lasted 40 cycles (160 trials

total). During this session, the cursor was immediately rotated 24° CCW relative to the hand path

(Fig. 1A, readaptation period). This last abrupt adaptation was used to examine whether gradual

or abrupt learning, followed by varying washout conditions, would lead to savings during a

second rotation period.

Empirical Data Analysis

Data analysis was conducted in MATLAB (MathWorks, Natick, MA). To quantify performance, we

calculated the reaching movement angle at peak velocity. This angle was computed as the hand's

angular position relative to a line segment connecting the start and target positions. We removed

outlier reach angles, by first removing angles with an absolute value greater than 60° (0.567% of

trials), and then removing angles that deviated by more than 5 median absolute deviations within

all trials in a given cycle, collapsed across participants in a specific experimental group (1.798% of

trials). Next, data were averaged within each 4-trial cycle.

To empirically compare adaptation between groups, we assessed the learning rate during

each abrupt rotation phase via an exponential function. We focused on rate measures, to prevent

our evaluation from being biased by the initial state of the learner (*i.e.*, subjects may demonstrate

greater total adaptation not because they learn more rapidly, but because their initial reach angle

is greater during the second abrupt rotation due to incomplete washout from the initial rotation).

Thus, we fitted an exponential curve to participant hand angles, and used the rate parameter as

our measure of learning rate. For this, we employed a 2-parameter exponential function designed

to enforce that each curve begins at the initial pre-rotation reach angle. For the initial

perturbation in the control group, this hand angle was computed as the average angle over the

last 4 baseline cycles. For the abrupt rotations during the second exposure period (across all

groups), this angle was computed as the average angle over the last 4 washout cycles. The

exponential function took the mathematical form:

$$y(t) = \alpha - (\alpha - y_0)e^{-\beta t} \qquad\qquad (1)$$

Here, α is parameter that encodes the exponential curve's asymptote, y_0 is the initial reach angle

described above, β is the rate parameter of interest, $y(t)$ is the reach angle on cycle t (starting at

$t = 0$). The function was fit to 41 cycles of data at a time (40 cycles of the abrupt rotation plus the

initial reach angle cycle on $t = 0$) using a bounded least-squares approach (MATLAB *fmincon*). The

parameter search was repeated 100 times, each time varying the initial parameter estimate that

seeds the algorithm. The α parameter was bounded between -45° and 45°. The β parameter was

bounded between 0 cycle^{-1} and 0.5 cycle^{-1}. The model parameters that minimized squared error

across all 100 repetitions were selected.

State-space learning model

To interpret our empirical results using an error-based learning framework, we employed a state-

space model[15,22,25,28,29,31]. The model describes adaptation as a trial-by-trial learning process

whereby the adapted state of the individual is updated by (1) learning from the error experienced

on each trial and (2) a small amount of decay that reflects temporal instability of short-term

adaptation[30,48,49]. In the model, the total update in the motor memory due to an error is

controlled by one's sensitivity to error (b) and the strength of memory retention between trials

is encoded by a retention factor (a). Together, learning and forgetting collectively determine how

the subject's internal state (x) evolves in response to an error (e) experienced on trial t in the

presence of Gaussian state noise (ε_x, normally distributed with zero mean, and standard

deviation of σ_x) according to:

$$x^{(t+1)} = ax^{(t)} + be^{(t)} + \varepsilon_x^{(t)} \tag{2}$$

194   Eq. (2) allows us to ascribe performance differences during the adaptation period to interpretable

195   quantities: retention (a) and error sensitivity (b).

196   Note, however, that the internal state (x) is not a measurable. Instead, on each cycle, the

197   motor output (reach angle) is measured. The reach angle (y) directly reflects the subject's internal

198   state, but is altered by execution noise (ε_y, normal with zero mean, std. dev. = σ_y) according to:

$$y^{(t)} = x^{(t)} + \varepsilon_y^{(t)} \tag{3}$$

Together, Eqs. (2) and (3) represent a single module state-space model. We fit this model to each

participant's reach angles during the adaptation period using the Expectation-Maximization (EM)

algorithm[28]. For each adaptation period, all 40 abrupt rotation cycles and 3 preceding baseline or

washout cycles were included. EM is an algorithm that conducts maximum likelihood estimation

in an iterative process. We used EM to identify the parameters {a, b, x_1, σ_x, σ_y, σ_1} that maximized

the likelihood of observing the data (x_1 and σ_1 represent the subject's initial state and variance,

respectively). We conducted 100 iterations, each time modifying the initial parameter guess that

seeded the algorithm. The retention factor (a) was bounded between 0.85 and 1. The initial state

(x_1) was bounded between -10 and 20°. Error sensitivity (b) was bounded between 0 and 1. Last,

all 3 variance parameters were bounded between $1.0e^{-3}$ and 100. We selected the parameter set

that maximized the likelihood function across all 100 iterations.

Statistics

For statistical comparisons between 2 groups or conditions, paired or independent two-sample

t-tests were employed. When more than 2 groups were compared, we applied one-way ANOVAs.

For post-hoc testing, we used Dunnett's test to measure differences between each experimental

group and the control group (we reached analogous conclusions when Tukey's test was used for

post-hoc comparisons across all groups without specifying a control condition).

Results

Participants executed point-to-point movements between a start location and a target (Fig. 1A).

In each group, a visual rotation was applied to the cursor motion in two separate exposure

periods: a condition which generally leads to an increase in learning rate called savings. Here, we

altered the size of errors experienced in the washout period between each exposure (Fig. 1B) to

examine which conditions protect or abolish visuomotor savings.

Visuomotor adaptation during point-to-point movements is accelerated by visuomotor savings

Adaptation to a novel visuomotor rotation is often studied in the context of "slicing" movements

in which participants rapidly move a cursor through a target location without online feedback

corrections. Here, we trained subjects to execute point-to-point reaches that require accurate

endpoint control. To confirm that our point-to-point design also produces visuomotor savings as

we recently reported[27], we tested participants in our control group. After a short baseline period,

a 24° rotation was abruptly introduced. Over time subjects adapted their reach angles to partially

counter the disturbance (Fig. 1C, abrupt rotation). To washout this initial learning, we removed

visual feedback, causing reach angles to gradually decay towards their baseline state (Fig. 1C, no

feedback). To determine whether the learning system was improved by prior experience, we

abruptly introduced a second 24° abrupt rotation, causing a rapid change in reach angle (Fig. 1C,

abrupt rotation). How did learning in this second exposure period compare to naïve adaptation?

To answer this question, we isolated the two learning periods and fit an exponential curve

to each participant's reach angles during both adaptation phases (Fig. 1D). As expected, the prior

exposure to the rotation accelerated adaptation, almost doubling the subjects' learning rates

(Fig. 1E; paired t-test, $t(9)=6.8$, $p<0.001$, Cohen's $d=2.15$). In sum, we find that point-to-point

adaptation to a 24° rotation produces savings and is amenable to further study.

Figure 1. Savings in a point-to-point visuomotor rotation task. **A.** Experiment apparatus. Subjects were

seated in a KINARM exoskeleton and executed point-to-point reaching movements to 1 of 4 targets. **B.**

Perturbation schedule. The control experiment consisted of 4 experimental periods: (1) 10-cycle baseline

period, (2) abrupt 24°, 40-cycle rotation (exposure 1), (3) 40-cycle no visual feedback washout, (4) 40-cycle re-exposure to the 24° rotation (exposure 2). **C.** Learning curves. The reach angle at maximum velocity was calculated on each trial and averaged within each 4-trial cycle. **D.** Savings visualization. The learning curves during the first and second exposure to the rotation are shown. We fit a 2-parameter exponential curve to individual participants to quantify learning rates. The empirical model fit (empirical fit) is also shown overlaid on the measured learning curves (exposure 1 or exposure 2). **E.** Quantification of savings. We compared the learning rate parameters across the first and second exposures quantified via the exponential model fit in **D**. Errors bars indicated mean ± 1 standard error of the mean. Single points in **E** represent individual participants. Statistics: *** indicates p<0.001.

Abrupt and gradual perturbation exposures similarly enhance the rate of learning

Past studies have yielded conflicting evidence as to the necessary conditions for savings. In some cases, savings appears to require the experience of similarly sized errors during the initial and the re-exposure phases of the experiment[24]. This is achieved by creating large errors in each adaptation period via an abrupt perturbation. More recently, however, other studies have observed savings without the experience of large errors[23,26]. In this case, the rotation is introduced in a gradual manner whereby its magnitude slowly increases during the initial exposure. Do gradual rotations lead to savings in point-to-point reaching movements?

To answer this question, we tested a second group of participants in a gradual adaptation paradigm (Fig. 2A). Whereas the abrupt group in Fig. 1 experienced a 24° perturbation all at once, subjects in the gradual group experienced a time-varying rotation that grew 1° over 24 cycles to reach its terminal value. This dramatically slowed the rate of initial adaptation (Fig. 2B, compare blue: gradual and black: abrupt). After this initial exposure, participants in the gradual and abrupt groups experienced identical trial conditions: a no-feedback washout period in which the adapted hand angles returned to their baseline state (Fig. 2B, no feedback), followed by a second rotation that was introduced abruptly in both groups.

To determine how the initial exposure to the rotation modulated savings, we isolated the

adapted reach angles during the second exposure period (Fig. 2C) and fit an exponential curve to

each participant's performance across both groups. Intriguingly, we did not detect a statistically

significant difference in learning rates (Fig. 2D; $t(18)=0.328$, $p=0.747$, Cohen's $d=0.147$),

suggesting that abrupt and gradual exposures both lead to visuomotor savings.

Figure 2. Abrupt and gradual adaptation lead to similar expression of savings in a point-to-point reaching

task. **A.** Perturbation schedule. Two groups were tested in an experiment consisting of 4 phases: baseline,

initial perturbation, washout, secondary perturbation. Groups differed only based on the visuomotor

rotation encountered during the initial exposure: the control group from Fig. 1 experienced a 24° rotation

that immediately reached its terminal value (labeled abrupt here), and the gradual group experienced a

rotation that slowly increased in magnitude across 24 cycles to reach the same terminal value. **B.** Learning

curves. The reach angle at maximum velocity was calculated on each trial and averaged within each 4-trial

cycle. The control group (black during exposure 1, and brown during exposure 2) behavior is compared to

the gradual group (cyan) behavior. The perturbation schedules are also shown overlaid with the data. **C.**

Savings visualization. The learning curves during the second exposure period are shown; both groups

adapted to an abrupt rotation during this period. We fit a 2-parameter exponential curve to individual

participants to quantify learning rates. The empirical model fit (empirical fit) is shown overlaid on the

measured learning curves (abrupt or gradual). **D.** Quantification of savings. We compared the learning

rate parameters during the second exposure period quantified via the exponential model fit in **C**. Errors

bars indicated mean ± 1 standard error of the mean. Single points in **D** represent individual participants.

Statistics: n.s. indicates $p > 0.05$.

Size of error during washout modulates the strength of visuomotor savings

Our results in Fig. 2 add to a growing body of evidence that savings can be achieved via a gradual

exposure to a perturbation, suggesting that large errors are not explicitly required to facilitate an

increase in learning rate. Why have some prior studies not observed the same result? Is it possible

that disagreement across studies is caused not by perturbation conditions, but rather by the type

of error experienced during the intervening period of washout that separates two adaptation

phases?

In Figs. 1 and 2, abrupt and gradual group subjects were de-adapted in a no-feedback

condition, preventing the experience of all visual errors. The absence of error led to a gradual

decay in reach angles, reproduced in Fig. 3B (cyan). To assess whether this de-adaptation

condition can alter the expression of savings, we tested two additional gradual groups (Fig. 3A)

that differed only by the errors experienced during washout. In the abrupt washout group, the

perturbation immediately ceased at the end of the initial adaptation period, creating large

oppositely-signed errors that induced rapid unlearning in the subsequent washout phase (Fig. 3B,

green). In the gradual washout group, the rotation was slowly decreased from cycle-to-cycle

during the washout period. This prevented the occurrence of large oppositely-signed errors,

leading to gradual unlearning of the initial adapted reach direction (Fig. 3B, magenta).

Next, all 3 gradual learning groups were exposed to an abrupt 24° rotation. To determine

how preceding washout modulated subsequent learning rates, we compared reach angles during

this abrupt readaptation period to those measured in naïve participants that had not previously

experienced a rotation (*i.e.*, the initial adaptation period in the control group examined in Fig. 1).

As before, we fit exponential curves to participant reach angles to empirically quantify individual

subject learning rates (Fig. 3C). Intriguingly, we observed that washout conditions had a dramatic

effect on re-learning rates (Fig. 3D, one-way ANOVA, $F_{(3,36)}=4.97$, $p=0.006$). Namely, savings was

observed only after no-feedback (post-hoc Dunnett's test, $p=0.019$) and gradual washout (post-

hoc Dunnett's test, $p=0.019$), but not after abrupt washout (post-hoc Dunnett test, $p=0.987$).

In sum, the ability to express savings after gradual learning is modulated by the errors the

participant experiences during de-adaptation. When large errors occur, as in our abrupt washout

group, savings is eliminated. When errors are small (*i.e.*, gradual washout) or are removed during

washout (*i.e.*, no feedback), savings is preserved.

Figure 3. Savings after gradual learning is attenuated by washout error magnitude. **A.** Perturbation

schedule. Three groups were tested in an experiment consisting of 4 phases: baseline, initial perturbation,

washout, secondary perturbation. All groups experienced a gradual perturbation during the initial learning

period, and an abrupt rotation during the second exposure. Groups differed only based on the errors

experienced during the washout (de-adaptation) phase. One group de-adapted in the absence of visual

feedback (cyan, no feedback). A second group de-adapted in the presence of large errors due to the

abrupt removal of the rotation (abrupt, green). A third group de-adapted in the presence of small errors

due to the gradual removal of the rotation (gradual, magenta). **B.** Learning curves. The reach angle at

maximum velocity was calculated on each trial and averaged within each 4-trial cycle. The abrupt group

(black and brown) behavior is compared to all 3 gradual group from described in inset **A**. The perturbation

schedules are overlaid with the data. **C.** Savings visualization. The learning curves during the second abrupt

exposure period are shown. Each gradual group is shown in a separate inset (left: abrupt de-adaptation,

middle: gradual de-adaptation, right: no-feedback de-adaptation). All learning curves are overlaid with

the naïve performance of the control group that experienced the initial rotation abruptly. We fit a 2-

parameter exponential curve to individual participants to quantify learning rates. The empirical model fit

(empirical fit) is shown overlaid on the measured learning curves (naïve, abrupt, gradual, or no feedback).

D. Quantification of savings. We compared the learning rate parameters during the second exposure

period in each gradual group to the learning rate parameters in the initial abrupt exposure of the control

group. Errors bars indicated mean ± 1 standard error of the mean. Single points in **D** represent individual

participants. Statistics: * indicates $p < 0.05$.

Savings following gradual learning is promoted by an increase in sensitivity to error

Short-term sensorimotor adaptation is due to the combined effects of two processes: (1) trial-to-

trial error-based learning, and (2) trial-to-trial forgetting[15,22,25,28–31]. In principle, savings could be

caused by either an increase in the amount of learning that follows the experience of an error,

an increase in the amount of learning retained with the passage of time, or a combination of both

sources. These two processes can be captured by a state-space model that represents adaptation

as a recursive process controlled by two parameters: an error sensitivity (controls the amount of

learning from error) and a retention factor (controls the amount of decay).

To determine if the savings we observed following gradual adaptation was due to changes

in error-based learning or forgetting, we fit state-space models to individual participant learning

curves during the second rotation periods in all 3 gradual adaptation groups (Fig. 4A, green, blue,

and magenta). To obtain naïve learning parameter estimates, we fit the state-space model to the

initial abrupt adaptation period in the control group in Fig. 1 (Fig. 4A, black).

359   To determine if the initial gradual adaptation and subsequent washout period altered the

360   stability of the motor memory during the readaptation period, we compared retention factors

361   across all 4 groups (Fig. 4B). We did not observe a statistically significant change in retention after

362   exposure to the gradual rotation (one-way ANOVA, $F_{(3,36)}=0.2$, $p=0.895$).

363   To determine if instead, a change in error sensitivity boosted adaptation rates during the

second exposure period, we compared the state-space model error sensitivity parameters across

all 4 groups (Fig. 4C). As in past studies[11,16,22], prior adaptation did alter the sensitivity to error

during the second exposure (one-way ANOVA, $F_{(3,36)}=3.81$, $p=0.0181$), accelerating re-learning

rate (Fig. 3D). Interestingly, we found a statistically significant increase in error sensitivity over

naïve, in the no-feedback condition (post-hoc Dunnett's test, $p=0.0387$), but not in the abrupt

condition (post-hoc Dunnett's test, $p=0.897$). In the gradual washout condition, we observed a

trend towards an increase in error sensitivity over the naïve condition, though this trend did not

reach statistical significance (post-hoc Dunnett's test, $p=0.548$).

Figure 4. Savings after gradual learning is mediated by an increase in error sensitivity. **A.** Learning curves

and visualization of savings. The reach angle at maximum velocity was calculated on each trial and

averaged within each 4-trial cycle. Each inset compares naïve adaptation (control, black) to 1 of 3 groups

that experienced an initial gradual rotation and secondary abrupt rotation, separated by washout (left:

gradual group that de-adapted with abrupt removal of the rotation, middle: gradual group that de-

adapted with gradual removal of the rotation, right: gradual group that de-adapted with any visual error).

We fit a state-space learning model (model) to individual participant reach angles that describes learning

as a recursive trial-to-trial process of error-based adaptation and decay. **B.** Comparison of retention

factors. Retention factors estimated by state-space model fits are compared across the secondary abrupt

learning periods of the gradual group and the initial naïve adaptation period in the control group. **C.**

Comparison of error sensitivity. The sensitivity to error parameters estimated by state-space model fits

are compared across the secondary abrupt learning periods of the gradual group and the initial naïve

adaptation period in the control group. Errors bars indicated mean ± 1 standard error of the mean. Single

points in **B** and **C** represent individual participants. Statistics: n.s. indicates $p > 0.05$, ** indicates $p < 0.01$.

Taken together, this indicates that initial gradual adaptation elevates sensitivity to error, leading

to savings. However, errors experienced during the washout phase can reverse this gain in error

sensitivity. Enhancement in error sensitivity is optimally preserved when visual errors are totally

removed in the washout phase via a no-feedback condition.

Discussion

The conditions necessary to increase the learning rate of the sensorimotor system are not

completely known. This is clearly illustrated in gradual adaptation paradigms where participants

adapt to a perturbation slowly and are then re-tested in an abrupt context following a period of

washout. Whereas some studies have observed more rapid learning in this context[23,26], other

accounts have found that gradual learning is unable to improve abrupt adaptation[24]. Here we

considered whether this contradiction might be due to variations in the de-adaptation phase that

separates two learning periods in typical savings paradigms. Specifically, does the salience of

oppositely-signed errors during washout inhibit savings after gradual learning?

Here we examined this idea by altering the type of error experienced during the washout

phase after a gradual adaptation period. We considered no-feedback, gradual, and abrupt de-

adaptation conditions that differ along the following error size gradient. (1) Removing visual

feedback eliminates visuomotor error completely, and so adaptation decays only due to a

forgetting process[32]. (2) With gradual de-adaptation, errors are present, but are small because

the motor action and perturbation are reduced in tandem preventing the occurrence of large

errors. (3) Similar to abrupt adaptation, abrupt de-adaptation results in the largest errors due to

the immediate removal of the perturbation when subjects still remain in the fully adapted state.

Our empirical analyses revealed accelerated learning after subjects de-adapted gradually or

without feedback, but not after abrupt removal of the initial perturbation (Fig. 3D). Interpreted

in an error-based framework, this suggests that the size of oppositely-signed errors during the

washout phase is inversely related to the propensity to save: larger errors appear to nullify

potential improvements in learning following initial gradual adaptation.

This conclusion qualitatively aligns with previous work by Kojima et al. (2004)[9], which

demonstrated that savings in the oculomotor system is sensitive to the duration of washout:

when the duration of the washout phase is increased, the learning rate during the re-exposure

phase reverts to its baseline state, abolishing savings. Thus, both the duration of error

presentation and the error magnitude during washout seem to negatively interact with the

expression of savings. This hypothesis may help to explain the mixed results noted above in

visuomotor studies of gradual learning. Both Coltman et al. (2021)[26] and Yin & Wei (2020)[23]

employed shorter washout periods, and observed savings following gradual adaptation, whereas

Herzfeld et al. (2014)[24] applied a longer washout period and did not find savings after gradual

learning. This possible explanation could easily be tested in a follow-up study that varies the

length of the washout period after an initial gradual learning period.

Along these lines, Kitago et al. (2013)[32] anecdotally noted that increasing the number of

washout trials appeared to dampen the magnitude of savings relative to earlier studies that

applied much fewer de-adaptation trials[33,34]. With that said, their observation was made in the

context of an abrupt-then-abrupt savings protocol, instead of the mixed gradual-then-abrupt

design we used. The expression of savings following abrupt or gradual learning periods may have

critical differences. That is, whereas removing feedback after gradual learning resulted in rapid

adaptation in our study, it abolished savings after abrupt learning in the experimental conditions

used by Kitago et al. (2013)[32]. We think that this stark discrepancy is likely caused by variations

in the utilization of implicit and explicit learning processes, which were not directly measured by

Kitago et al., nor in our current study. That is, gradual learning paradigms reduce the size of error,

which in turn limits cognitive awareness of the perturbation and associated strategic re-aiming

strategies[22,23,35]. We also anticipate larger strategy use in the Kitago et al.'s study due to the

utilization of a single target (as opposed to the 4 targets used herein), which also appears to

attenuate procedural aftereffects[36,37]. Accordingly, savings in gradual-then-abrupt learning

paradigms have been linked to increases in implicit learning rates[23], as opposed to overt

strategies that tend to dominate savings responses in abrupt-then-abrupt protocols[7,12]. Thus, we

hypothesize that washout has divergent effects on the ability of implicit and explicit systems to

save: removing feedback during washout may protect savings in the implicit system (which might

otherwise be suppressed by "negatively-signed" errors during the washout period[38]), but

negatively impact savings in the explicit system. This fascinating possibility could be tested in

follow-up studies that assay implicit and explicit contributions to learning following abrupt and

gradual adaptations with differing washout conditions.

To understand what learning property links washout to the capacity to save, we applied

a state-space model that casts short-term adaptation as a cumulative process of trial-to-trial

error-based learning and decay. This model indicated that gradual savings appears to involve an

upregulation in sensitivity to error, that is reversed by exposure to large errors in the washout

phase (Fig. 4C). This outcome comports with past work, suggesting that increases in error

sensitivity are the primary source of accelerated re-learning after abrupt, gradual, or random

perturbations[26], and the adaptation of disparate motor systems such as reaching[16,22,26] and

walking[11]. Thus, a growing body of evidence suggests that exposure to a perturbation causes an

increase in error sensitivity that can improve performance in subsequent exposures to similar

perturbations[22,24]. Our results suggest that increases in error sensitivity can be undermined by

inconsistently signed errors experienced in between learning periods.

What specific properties of error suppress previous gains in error sensitivity? We found

that error sensitivity was elevated specifically in individuals that de-adapted without feedback,

suggesting that improvements in error sensitivity can be protected by the absence of error. With

that said, given that the subjects' re-learning rates were similarly increased in both no-feedback

and gradual de-adaptation conditions (Fig. 3D), it is possible that error sensitivity was also

elevated in the gradual de-adaptation group (consistent with the trend in Fig. 4C), but that we

did not have sufficient statistical power to detect this effect. Thus, it is not currently clear

whether error sensitivity is reduced by the experience of *any* opposing error, or whether the size

of the opposing error further modulates the rate of error sensitivity decline.

Our findings have critical implications for the learning rules suggested by an influential

error sensitivity model defined by a memory of errors[13,24,25]. The memory of errors model that

was initially proposed by Herzfeld et al. (2014)[24] accounts for changes in error sensitivity based

on the statistical properties of error. Namely, when errors have consistent signs, this indicates

that the learning system could have adapted more to the initial experience of error to better

eliminate performance error. A critical feature of this model is its locality: changes in error

sensitivity are restricted to errors of a given size. In other words, the experience of consistently

small or large errors modulates error sensitivity specific to errors of that magnitude. Thus,

contrary to our findings, a memory of errors model predicts that a gradual adaptation paradigm

will not lead to savings during an abrupt perturbation, because abrupt errors are much larger

than those experienced during a gradual perturbation. Thus, our results add to a growing body

of evidence[23,26] suggesting a need to revisit properties of the error sensitivity landscape; it may

be that the experience of consistent errors generalizes more broadly to unexperienced errors of

disparate sizes than previously appreciated.

The possibility that changes in error sensitivity extend beyond the experienced range of

errors could help not only to understand patterns of adaptation in gradual-then-abrupt learning

paradigms, but also suppression of learning rates in anterograde interference paradigms[22,31,39–

41]. That is, whereas increases in error sensitivity have been linked to improvements in learning

when two similar perturbations are experienced in sequence, decreases in error sensitivity have

been found to produce a slowing of adaptation when transitioning between two contrasting

perturbations[31]. To date, no memory of errors model has attempted to integrate these opposing

error sensitivity-driven effects on learning rate. Limited application of the memory of errors

model to interference paradigms arises, because learning in response to "positively-signed"

errors during the first exposure does not produce error sensitivity changes that generalize to

"negatively-signed" errors during the second exposure (as they have not been experienced

before). Our findings, that large oppositely-signed errors during washout reverse prior

enhancements in error sensitivity (Fig. 4C), suggest that the experience of a sensorimotor error

produces two distinct effects: (1) an increase in sensitivity to similar errors, and (2) a suppression

in error sensitivity to contrasting errors. This second suppressive learning property may be a

critical process missing from current memory of errors models that would aid in interpreting both

the dissolution of savings following certain types of washout, as well as interference effects.

We speculate that this antagonistic interaction between the response to "positive" and

"negative" errors may be encoded within cerebellar learning circuits that mediate error-based

learning processes, a structure known to encode error directionality based on its anatomical

connections with the inferior olive[42–44]. In particular, errors of a given direction (*i.e.*, sign) lead to

complex spiking in some Purkinje-cell (P-cell) subpopulations in the cerebellar cortex, and

suppression in others. Complex spiking events in turn lead to long term potentiation or

depression at P-cell synapses with parallel fibers[45]. While it is not clear precisely how error

sensitivity is encoded by groups of P-cells, our findings suggest the possibility that prolonged

absence of complex spikes (when errors are opposite of the preferred direction) during washout

periods may produce cellular or synaptic-level changes that reduce the plastic learning potential

of these neurons in the future (*i.e.*, during the re-exposure period). This idea, and the source of

cerebellar error sensitivity more generally, remain to be examined.

It is important to distinguish this memory of errors framework from that of Bayesian

inference models that incorporate both error-based learning processes and inference across

multiple sensorimotor contexts[46,47]. In single-context memory of errors models, the same

adaptive states apply uniformly across different experimental periods, such that any change in

learning rate must be due to a variation in learning parameters: sensitivity to error or memory

retention. But with multiple contexts, as in the Bayesian COIN model by Heald et al. (2021)[46], the

apparent learning rate is influenced by inferring which context is most likely given sensory

information. For the COIN model, visuomotor adaptation involves two contexts: a baseline

context and a rotation context. Faster learning during the second exposure is caused not by a

change in the response to error, but by an increase in the propensity to expect the rotation given

prior training during the initial exposure period. Thus, our findings could have important

implications for how this inference occurs in the sensorimotor system. Namely, the errors (*i.e.*,

sensory information) experienced during the washout period alters the inference process: with

large "negatively-signed" errors during the washout period, the nervous system decreases its

expectation that the rotation state (with its "positively-signed" errors) will occur again in the

future. This idea, and a broader comparison between the memory of errors and Bayesian

frameworks on the effects of washout on future learning, remain to be tested.

Competing interests

The authors declare no competing interests.

Data availability

All data and analysis code will be made accessible via OSF upon acceptance.

Author contributions

MJ, BM and JW designed the experiments and methodology. MJ and BM collected data. MJ and

STA analyzed data and created figures. MJ and STA wrote manuscript. MJ, STA, and JW edited

that manuscript.

References

1. Tsay, J. S., Haith, A. M., Ivry, R. B. & Kim, H. E. Interactions between sensory prediction

error and task error during implicit motor learning. *PLoS Comput Biol* **18**, e1010005-

(2022).

2. Taylor, J. A. & Ivry, R. B. Flexible cognitive strategies during motor learning. *PLoS Comput*

Biol **7**, e1001096 (2011).

3. Leow, L.-A., Marinovic, W., de Rugy, A. & Carroll, T. J. Task errors drive memories that

improve sensorimotor adaptation. *Journal of Neuroscience* (2020)

doi:10.1523/JNEUROSCI.1506-19.2020.

4. Mazzoni, P. & Krakauer, J. W. An implicit plan overrides an explicit strategy during

visuomotor adaptation. *Journal of Neuroscience* **26**, 3642–3645 (2006).

5. Albert, S. T., Blaum, E. C. & Blustein, D. H. Sensory prediction error drives subconscious

motor learning outside of the laboratory. *J Neurophysiol* **130**, 427–435 (2023).

6. Ebbinghaus, H. *Uber Das Gedachtnis*. (Dunacker and Humblot, Leipzig, Germany, 1885).

7. Haith, A. M., Huberdeau, D. M. & Krakauer, J. W. The Influence of Movement Preparation

Time on the Expression of Visuomotor Learning and Savings. *Journal of Neuroscience* **35**,

5109–5117 (2015).

8. Landi, S. M., Baguear, F. & Della-Maggiore, V. One week of motor adaptation induces

structural changes in primary motor cortex that predict long-term memory one year

later. *J Neurosci* **31**, 11808–11813 (2011).

9. Kojima, Y., Iwamoto, Y. & Yoshida, K. Memory of Learning Facilitates Saccadic Adaptation

in the Monkey. *Journal of Neuroscience* **24**, 7531–7539 (2004).

10.   Malone, L. A., Vasudevan, E. V. L. & Bastian, A. J. Motor Adaptation Training for Faster

Relearning. *Journal of Neuroscience* **31**, 15136–15143 (2011).

11.   Mawase, F., Shmuelof, L., Bar-Haim, S. & Karniel, A. Savings in locomotor adaptation

explained by changes in learning parameters following initial adaptation. *J Neurophysiol*

111, 1444–1454 (2014).

12.   Morehead, J. R., Qasim, S. E., Crossley, M. J. & Ivry, R. Savings upon Re-Aiming in

Visuomotor Adaptation. *Journal of Neuroscience* **35**, 14386–14396 (2015).

13.   Leow, L.-A., de Rugy, A., Marinovic, W., Riek, S. & Carroll, T. J. Savings for visuomotor

adaptation require prior history of error, not prior repetition of successful actions. *J

Neurophysiol* **116**, 1603–1614 (2016).

14.   Brashers-Krug, T., Shadmehr, R. & Bizzi, E. Consolidation in human motor memory.

Nature **382**, 252–255 (1996).

15.   Smith, M. A., Ghazizadeh, A. & Shadmehr, R. Interacting adaptive processes with

different timescales underlie short-term motor learning. *PLoS Biol* **4**, e179 (2006).

16.   Coltman, S. K., Cashaback, J. G. A. & Gribble, P. L. Both fast and slow learning processes

contribute to savings following sensorimotor adaptation. *J Neurophysiol* **121**, 1575–1583

(2019).

17.   Medina, J. F., Garcia, K. S. & Mauk, M. D. A mechanism for savings in the cerebellum.

Journal of Neuroscience **21**, 4081–4089 (2001).

18.   Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning in

a Sensorimotor Adaptation Task. *Journal of Neuroscience* **34**, 3023–3032 (2014).

19. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning.

Nat Commun **10**, 40 (2019).

20. McDougle, S. D., Bond, K. M. & Taylor, J. A. Explicit and Implicit Processes Constitute the

Fast and Slow Processes of Sensorimotor Learning. *Journal of Neuroscience* **35**, 9568–

9579 (2015).

21. Huberdeau, D. M., Krakauer, J. W. & Haith, A. M. Practice induces a qualitative change in

the memory representation for visuomotor learning. *J Neurophysiol* **122**, 1050–1059

(2019).

22. Albert, S. T. *et al.* Competition between parallel sensorimotor learning systems. *Elife* **11**,

e65361 (2022).

23. Yin, C. & Wei, K. Savings in sensorimotor adaptation without an explicit strategy. *J*

Neurophysiol **123**, 1180–1192 (2020).

24. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in

sensorimotor learning. *Science (1979)* **345**, 1349–1353 (2014).

25. Albert, S. T. *et al.* An implicit memory of errors limits human sensorimotor adaptation.

Nat Hum Behav (2021) doi:10.1038/s41562-020-01036-x.

26. Coltman, S. K., van Beers, R. J., Medendorp, W. P. & Gribble, P. L. Sensitivity to error

during visuomotor adaptation is similarly modulated by abrupt, gradual, and random

perturbation schedules. *J Neurophysiol* **126**, 934–945 (2021).

27. James, R., Bao, S., D'Amato, A. & Wang, J. The nature of savings associated with a

visuomotor adaptation task that involves one arm or both arms. *Hum Mov Sci* **81**, 102896

(2022).

28.   Albert, S. T. & Shadmehr, R. Estimating properties of the fast and slow adaptive

604   processes during sensorimotor adaptation. *J Neurophysiol* **119**, 1367–1393 (2018).

605   29.   Orozco, S. P., Albert, S. T. & Shadmehr, R. Adaptive control of movement deceleration

606   during saccades. *PLoS Comput Biol* **17**, e1009176–e1009176 (2021).

607   30.   Brennan, A. E. & Smith, M. A. The Decay of Motor Memories Is Independent of Context

608   Change Detection. *PLoS Comput Biol* **11**, e1004278 (2015).

609   31.   Lerner, G. *et al.* The Origins of Anterograde Interference in Visuomotor Adaptation.

610   *Cerebral Cortex* **30**, 4000–4010 (2020).

611   32.   Kitago, T., Ryan, S. L., Mazzoni, P., Krakauer, J. W. & Haith, A. M. Unlearning versus

612   savings in visuomotor adaptation: comparing effects of washout, passage of time, and

613   removal of errors on motor memory. *Front Hum Neurosci* **7**, 307 (2013).

614   33.   Zarahn, E., Weston, G. D., Liang, J., Mazzoni, P. & Krakauer, J. W. Explaining Savings for

615   Visuomotor Adaptation: Linear Time-Invariant State-Space Models Are Not Sufficient. *J*

616   *Neurophysiol* **100**, 2537–2548 (2008).

617   34.   Huang, V. S., Haith, A., Mazzoni, P. & Krakauer, J. W. Rethinking motor learning and

618   savings in adaptation paradigms: model-free memory for successful actions combines

619   with internal models. *Neuron* **70**, 787–801 (2011).

620   35.   Neville, K.-M. & Cressman, E. K. The influence of awareness on explicit and implicit

621   contributions to visuomotor adaptation over time. *Exp Brain Res* **236**, 2047–2059 (2018).

622   36.   Tsay, J. S. *et al.* Large-scale citizen science reveals predictors of sensorimotor adaptation.

623   *Nat Hum Behav* (2024) doi:10.1038/s41562-023-01798-0.

624   37.   Wang, J., Bao, S. & Tays, G. D. Lack of generalization between explicit and implicit

625   visuomotor learning. *PLoS One* **14**, e0224099 (2019).

626   38.   Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor

627   perturbation produces opposite effects on explicit and implicit learning processes. *PLoS*

628   *Biol* **19**, e3001147 (2021).

629   39.   Sing, G. C. & Smith, M. A. Reduction in learning rates associated with anterograde

630   interference results from interactions between different timescales in motor adaptation.

631   *PLoS Comput Biol* **6**, e1000893 (2010).

632   40.   Caithness, G. *et al.* Failure to consolidate the consolidation theory of learning for

633   sensorimotor adaptation tasks. *Journal of Neuroscience* **24**, 8662–8671 (2004).

634   41.   Miall, R. C., Jenkinson, N. & Kulkarni, K. Adaptation to rotated visual feedback: a re-

635   examination of motor interference. *Exp Brain Res* **154**, 201–210 (2004).

636   42.   Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of error and learning to

637   correct that error by the Purkinje cells of the cerebellum. *Nat Neurosci* **21**, 736–743

638   (2018).

639   43.   Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of action by the Purkinje

640   cells of the cerebellum. *Nature* **526**, 439–442 (2015).

641   44.   Soetedjo, R., Kojima, Y. & Fuchs, A. F. Complex spike activity in the oculomotor vermis of

642   the cerebellum: a vectorial error signal for saccade motor learning? *J Neurophysiol* **100**,

643   1949–1966 (2008).

644   45.   Suvrathan, A., Payne, H. L. & Raymond, J. L. Timing Rules for Synaptic Plasticity Matched

645   to Behavioral Function. *Neuron* **92**, 959–967 (2016).

646   46.   Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference underlies the learning of

647   sensorimotor repertoires. *Nature* **600**, 489–493 (2021).

648   47.   Haruno, M., Wolpert, D. M. & Kawato, M. MOSAIC Model for Sensorimotor Learning and

649   Control. *Neural Comput* **13**, 2201–2220 (2001).

650   48.   Joiner, W. M. & Smith, M. A. Long-term retention explained by a model of short-term

651   learning in the adaptive control of reaching. *J Neurophysiol* **100**, 2948–2955 (2008).

652   49.   Alhussein, L., Hosseini, E. A., Nguyen, K. P., Smith, M. A. & Joiner, W. M. Dissociating

653   effects of error size, training duration, and amount of adaptation on the ability to retain

654   motor memories. *J Neurophysiol* **122**, 2027–2042 (2019).