Protection of savings by reducing the salience of opposing errors

Mousa Javidialsaadi[1†], Scott T. Albert[2†], Badr Moufarrej S Al Mutairi[1], Jinsung Wang[1*]

[1]Kinesiology programs, Zilber College of Public Health, University of Wisconsin–Milwaukee,

Milwaukee, WI, 53201, USA

[2]Neuroscience Center, University of North Carolina at Chapel Hill, Chapel Hill, NC 27514, USA

[†]These authors contributed equally to this work.

*Correspondence: Jinsung Wang, 492 Enderis Hall, University of Wisconsin – Milwaukee,

Milwaukee, WI 53201, USA. Email wang34@uwm.edu. Phone: 414-229-2645

Key words: Gradual adaptation; Visual rotation; Washout; Unlearning; Implicit learning;

Readaptation

Abstract

When humans encounter the same perturbation twice, they typically adapt faster the second

time, a phenomenon called savings. Studies have examined savings following adaptation to a

gradually introduced perturbation, with mixed results. These inconsistencies might be caused by

differences in how behavior returns to its baseline state during the 'washout' phase in between

learning periods. To test this, participants controlled a cursor that was subject to a visual rotation

in its motion direction. The rotation was applied during two learning periods, separated by a

washout period where the rotation was removed abruptly, gradually, or without error feedback.

We found that the type of error experienced during washout affected savings: abrupt washout

with large errors eliminated savings, whereas gradual or no-feedback washout preserved it.

Model-based analyses indicated these effects were driven by changes in error sensitivity

suggesting that salient, opposing errors experienced during washout downregulate the response

to error, nullifying savings.

Introduction

When an unexpected perturbation to our motor system occurs, the resulting errors in our actions and predictions[1–5] update our motor commands, predictively countering future disturbances. For example, when we walk on slippery ice, a sudden fall results in an unwanted error that alters how we traverse the icy patch in the future. Even when the icy patch (i.e., the perturbation) is removed and we reestablish our normal walking patterns, a lingering memory is stored within the nervous system, whereby re-exposure to the perturbation results in more rapid re-learning. This learning hallmark, termed savings[6], can occur on the same day, overnight[7], or even a year after the initial perturbation[8], and is ubiquitous across movements executed with our eyes (saccade adaptation[9]), legs (locomotion[10,11]), arms (visuomotor[7,12,13] and force field adaptation[14–16]), as well as classical conditioning paradigms[17].

Our ability to flexibly increase our learning rate is known to rely on two parallel adaptive systems: an explicit process directed by our conscious strategies[18–20], and a subconscious implicit process that operates without our awareness[2,4]. The implicit and explicit contributions to savings in motor tasks are typically studied using visuomotor rotations. In this task, subjects control some type of cursor with their hand or arm and move it to a target location. Following the baseline period of task familiarization, a visuomotor perturbation is introduced that rotates the cursor's trajectory relative to the hand's actual reaching direction. This is akin to altering the orientation of your laptop so that moving your cursor up results in an unexpected diagonal trajectory. The visual rotation thus induces an error, which individuals use to alter their behavior over time[21]. This altered behavior, or in this case, a preemptive change in the direction you move your hand to counteract the cursor's rotation, is supported by intentional changes in the aiming direction

you select (i.e., explicit strategy) along with an additional uncontrolled adjustment (*i.e.*, the

implicit correction)[18].

In visuomotor rotation tasks, improvements in strategy use are salient in cases where a

large rotation is introduced abruptly. The abrupt introduction of a large rotation causes large

errors in the anticipated cursor motion, which in turn encourages individuals to adopt an

intentional re-aiming strategy[22–24]. If the rotation is reintroduced in the future, participants recall

their past strategy, resulting in a greatly accelerated re-learning rate (termed "explicit

savings")[7,12,25]. This form of explicit savings can be suppressed either by limiting response time[22]

(which causes people to respond reflexively to the presentation of a target, thereby preventing

them from using a re-aiming strategy) or by reducing the error magnitude during initial learning[26]

(which reduces awareness of the rotation). The latter is typically done by gradually introducing

the rotation. When the rotation size increases gradually, our slow-adapting, subconscious (i.e.,

implicit) learning system is better able to mitigate the cursor errors, resulting in smaller

discrepancies between intended and actual cursor movements. This can help prime the implicit

system to exhibit savings that are detectable during subsequent encounters with the rotation[22,26].

Intriguingly, however, studies that employed gradual rotations have observed mixed

effects on the ability of *gradual* learning to facilitate more rapid *abrupt* re-learning. Absence of

savings following gradual learning[27] is consistent with a standard memory of errors model[27,28], in

which gradual learning cannot improve abrupt learning due to a mismatch in error size: errors

are small in the gradual case, and thus gradual learning fails to generalize to abrupt learning in

which large errors are experienced. Thus, to the brain, an abrupt perturbation following gradual

learning may appear to be a novel context (even though the rotation size at the end of learning

is the same in both conditions), because it has never experienced the large errors all at once in

the past. Although this prediction is consistent with at least one prior study[27], more recent work

has exhibited opposing results where gradual learning can improve abrupt learning rates[26,29,30],

possibly by accelerating implicit learning[26]. This raises a key question: why does gradual learning

lead to savings in some experiments, but not in others?

Here we consider the possibility that typical sensorimotor learning paradigms may fail to

produce savings not due to differences in how individuals learn, but rather because of how the

learned behavior decays between the learning and re-learning phases. Specifically, sensorimotor

savings is typically studied in paradigms where individuals adapt to a perturbation during an

initial learning period, their performance returns to baseline levels during a washout period, and

they are exposed to the same perturbation during a re-learning period (in psychological terms,

this 'washout' period is analogous to 'extinction'). For visuomotor rotation tasks, conditions

employed for washout of initial learning vary across studies. In some cases, a washout period

consists of abrupt cessation of the perturbation and includes the same number of trials as the

initial learning period[27]. Other studies use a much shorter washout phase[26,29] consisting of

augmented feedback that constrains the cursor motion to an exact straight line (i.e., zero visual

error, termed invariant error-clamp[29]) or trials where visual feedback is removed entirely (termed

no-feedback trials[26]). This no-feedback condition is akin to moving a hidden cursor on a screen.

Critically, these differences in washout conditions alter the size or quality of errors that

induce performance decay. Here we propose that *larger, or more salient,* errors experienced

during a washout period might reduce the nervous system's ability to save what was learned

from the initial learning period. In the present study, thus, we examine the overlooked possibility

in motor learning, that the nature of washout plays a central role in the expression of savings

following gradual learning. Across three groups we manipulate the types of washout errors by (1)

removing errors entirely by eliminating visual feedback, (2) limiting the size of errors by removing

a visual rotation gradually, or (3) facilitating large errors by abruptly disengaging the rotation. We

then apply empirical and model-based approaches to assess whether some washout conditions

more easily produce savings during abrupt learning that follows initial gradual learning.

Our current study seeks to not only clarify previous discrepancies in the literature, but

also to extend past studies by identifying sources that interfere with the expression of savings.

This has important implications for understanding how to preserve increases in our sensitivity to

errors[22,26,28] (thus resulting in greater savings), which in turn may inform new ways to account for

motor plasticity in current models of motor adaptation[22,28,31]. Over the long term, this knowledge

will facilitate the design of strategies to boost learning rates in neurorehabilitation paradigms.

Results

Adaptation to a novel visuomotor rotation is often studied in the context of "slicing" movements,

in which individuals briskly move their hand (or cursor) *through* a visual target, or point-to-point

movements, in which they move their hand to a target and stop on the target. Here, we employed

the latter movements that required accurate endpoint control. Our subjects grasped a robotic

exoskeleton that controlled the motion of a cursor on a horizontal screen (their arm was hidden

from view), and executed point-to-point movements between a start location and one of four

targets (Fig. 1a). Initially, the cursor followed the hand's position in a continuous stroke (baseline),

but during the learning periods (exposures 1 and 2), a 24° visual rotation was applied to the

cursor's motion (Fig. 1b). A washout period was placed in between the two learning periods in

which the rotation was removed through three different strategies (abrupt removal, gradual

removal, or completely removing visual feedback during each movement). Our primary goal was

to examine how errors experienced during the washout phase protect or abolish visuomotor

savings differently depending on the washout conditions.

Figure 1. Savings in a point-to-point visuomotor rotation task. **a**. Experiment apparatus. Subjects were seated in a
KINARM exoskeleton and executed point-to-point reaching movements to one of four targets. **b**. Perturbation
schedule for abrupt adaptation group. This control experiment consisted of 4 experimental periods: (1) 10-cycle
baseline period, (2) 40-cycle abrupt 24° rotation (exposure 1), (3) 40-cycle no-visual feedback washout, (4) 40-cycle
re-exposure to abrupt 24° rotation (exposure 2). **c**. Learning curves. Solid black line indicates size of rotation during
each period. Reach angle at maximum velocity was calculated on each trial and averaged within each 4-trial cycle. **d**.
Savings visualization. Learning curves during first and second exposure to the rotation are shown. We fit a 2-
parameter exponential curve to individual participants to quantify learning rates. The empirical model fit (empirical
fit) is also shown overlaid on the measured learning curves (exposure 1 or exposure 2). **e**. Quantification of savings.
We compared the learning rate parameters across exposures 1 and 2 quantified via the exponential model fit in d.
Errors bars indicated mean ± SE. Single points in e represent individual participants. Statistics: *** indicates p<0.001.

Visuomotor adaptation during point-to-point movements is accelerated by visuomotor savings

To confirm that our point-to-point movement design leads to savings (as we recently reported[32]),

we first tested a group of subjects (control group) who experienced a visuomotor rotation

abruptly during the initial learning period. After a short baseline period, a 24° rotation was

abruptly introduced. Over time subjects adapted their reach angles to partially counter the

rotation (Fig. 1c, exposure 1). To washout the initial learning, we removed visual feedback,

causing reach angles to gradually decay towards their baseline state (Fig. 1c, washout period). To

determine whether subsequent learning was improved by prior experience, we abruptly

introduced a 24° rotation again, causing a rapid change in reach angle (Fig. 1c, exposure 2). How

did learning during the second exposure period compare to naïve adaptation?

To answer this question, we isolated the two learning periods and fit an exponential curve

to each participant's reach angles during both adaptation phases (Fig. 1d). The exponential curve

produced an excellent fit to behavior, yielding a root-mean-squared-error (RMSE) within 0.57° of

baseline cycle-to-cycle variability (i.e., the theoretical floor for model error: see Methods:

Evaluating model error; RMSE is shown in Supplementary Fig. 1). As expected, prior exposure to

the rotation accelerated adaptation, almost doubling subject learning rates (Fig. 1e; paired t-test,

$t(14)=5.16$, $p<0.0001$, Cohen's $d=1.33$). A power law model for learning, commonly applied for

psychological learning[33–35], also showed an increased learning rate during the second exposure

(see Supplementary Fig. 2; Fig. 2a shows model fits; Fig. 2b provides power parameter; paired t-

test, $t(14)=4.13$, $p=0.001$, Cohen's $d=1.07$; model RMSE provided in Supplementary Fig. 1). In sum,

we find that point-to-point adaptation to a 24° rotation produces savings and is amenable to

further study.

Measuring re-learning rates after initial abrupt and gradual adaptations

Past studies have yielded conflicting evidence as to the necessary conditions for savings. In some

cases, savings appears to require the experience of similarly sized errors during both the initial

learning and re-learning periods[27]. This is achieved by creating large errors in each learning period

via an abrupt perturbation. Other studies, however, have observed savings without initial

experience of large errors[26,29]. In this case, the rotation is introduced gradually whereby its

magnitude slowly increases during initial exposure to the rotation.

To examine how gradual learning influences subsequent abrupt learning, we tested a

second group of participants with a gradual adaptation paradigm (Fig. 2a). Whereas the abrupt

adaptation group (controls) in Fig. 1 experienced a 24° rotation all at once during exposure 1,

participants in this gradual adaptation group experienced a time-varying rotation that grew 1°

per cycle over 24 cycles to reach its terminal value. Linear models fit to the learning curves of the

abrupt and gradual adaptation groups (Supplementary Fig. 3) confirmed that gradually

introducing the rotation slowed the rate of adaptation (additional participant groups described

in the following Results section were used in this analysis; see Methods: *Comparing gradual and*

abrupt learning rates; statistics provided in Supplementary Table 1).

Figure 2. Abrupt and gradual adaptation lead to similar expression of savings in a point-to-point reaching task. **a**. Perturbation schedule for abrupt or gradual adaptation group. Groups differed only based on the visuomotor rotation encountered during the initial exposure: control group from Fig. 1 experienced a 24° rotation that immediately reached its terminal value (labeled abrupt here); gradual group experienced a rotation that slowly increased in magnitude across 24 cycles to reach the same terminal value. **b**. Learning curves. Solid brown line indicates size of rotation during each period. Reach angle at maximum velocity was calculated on each trial and averaged within each 4-trial cycle. Control group (black during exposure 1, and brown during exposure 2) behavior is compared to gradual group (cyan) behavior. Perturbation schedules are overlaid with the data. **c**. Savings visualization. Learning curves during the second exposure period are shown; both groups adapted to an abrupt rotation during this period. We fit a 2-parameter exponential curve to individual participants to quantify learning rates. The empirical model fit (empirical fit) is shown overlaid on the measured learning curves (abrupt or gradual). **d**. Quantification of savings. We compared the learning rate parameters during exposure 2 quantified via the exponential model fit in c. Errors bars indicated mean ± SE. Single points in **e** represent individual participants. Statistics: n.s. indicates $p > 0.05$. Abbreviations: "gradual" indicates gradual adaptation group; "abrupt" indicates abrupt adaptation group.

After the initial, gradual adaptation (exposure 1), participants in this group experienced a no-feedback washout period, followed by abrupt adaptation (exposure 2). To compare between the gradual and abrupt adaptation groups, we isolated the learning curves during exposure 2 (Fig. 2c) and fit both exponential and power law models to assess their rates of learning. We did not detect a statistically significant difference in re-learning rate between the groups that

experienced the rotation gradually or abruptly during initial learning (Fig. 2d shows exponential

model; t(28)=1.315, p=0.199, Cohen's d=0.48; Supplementary Fig. 2d show the power law model;

t(28)=0.90, p=0.377, Cohen's d = 0.33), indicating the presence of savings following gradual

adaptation, a finding consistent with prior studies[29]. That being said, this null result may reflect

insufficient statistical power. The critical way to confirm whether the initial gradual rotation led

to savings is to compare re-learning rates to a naïve control condition. This is pursued in greater

detail in the following Results section.

Size of error during washout modulates the strength of visuomotor savings

It is unclear what causes gradual adaptation to induce savings in some conditions,[26,29,30] but not

others[27]. We considered the idea that savings between initial learning and re-learning phases

might be shaped by the washout period that separates the two learning phases.

As shown in Figs. 1 and 2, abrupt and gradual adaptation groups experienced a no-

feedback washout condition, preventing the experience of any visual errors during the washout

period (Fig. 3b, cyan). To assess whether the washout condition alters the expression of savings,

we tested two additional gradual adaptation groups (Fig. 3a) that differed based on the type of

error experienced during washout. In the abrupt washout group, the rotation immediately ended

at the end of the initial learning period, creating large, oppositely signed errors that induced a

rapid decline (Fig. 3b, green). In the gradual washout group, the rotation gradually decreased

cycle-to-cycle during the washout period. This prevented the occurrence of large errors in the

opposite direction, leading to a gradual decline in performance (Fig. 3b, magenta).

Figure 3. Savings after gradual learning is attenuated by washout error magnitude. **a**. Perturbation schedule for gradual adaptation groups in 3 washout conditions (no feedback, abrupt, gradual). All groups experienced a gradual rotation during exposure 1 and an abrupt rotation during exposure 2. Groups differed only based on errors experienced during washout. One group deadapted in the absence of visual feedback (cyan, no feedback). Second group deadapted in the presence of large errors due to abrupt removal of the rotation (abrupt, green). Third group deadapted in the presence of small errors due to gradual removal of the rotation (gradual, magenta). **b**. Learning curves. Solid lines in different colors indicate size of rotation for corresponding group during each period. Reach angle at maximum velocity was calculated on each trial and averaged within each 4-trial cycle. Abrupt adaptation group (black and brown) behavior is compared to all 3 gradual adaptation groups from described in inset a. The perturbation schedules are overlaid with the data. **c**. Savings visualization. Learning curves during exposure 2 (abrupt adaptation) are shown. Each gradual group is shown in a separate inset (left: abrupt washout; middle: gradual washout; right: no-feedback washout). All learning curves are overlaid with naïve performance of the control group that experienced initial rotation abruptly. We fit a 2-parameter exponential curve to individual participants to quantify learning rates. The empirical model fit (empirical fit) is shown overlaid on the measured learning curves (naïve, abrupt, gradual, or no feedback). **d**. Quantification of savings. We compared the learning rate parameters during exposure 2 in each gradual group to the learning rate parameters in the initial abrupt exposure of the control group. Errors bars indicated mean ± SE. Single points in **d** represent individual participants. Statistics: * indicates p < 0.05.

Following that, all three washout groups were exposed to an abrupt 24° rotation (i.e.,

exposure 2). To determine how washout influenced subsequent learning rates, we compared

reach angles during this abrupt adaptation period to those of naïve participants who had not

previously experienced a rotation (*i.e.*, the initial learning period of the control group shown in

Fig. 1). Comparing the rates of learning to a naïve control group is a standard approach to detect

acceleration in learning rates in conditions where initial adaptation occurred in a gradual

fashion[13,26,27,29,36–38]. As before, we fit both exponential and power law curves to participant reach

angles to empirically quantify individual subject learning rates (Fig. 3c for exponential model;

Supplementary Fig. 2e for power law model). Both the exponential and power models exhibited

excellent matches to behavior and possessed an average RMSE within 0.59° of the theoretical

minimum error level estimated based on intrinsic cycle-to-cycle reach variability during non-

adaptation periods (RMSE distributions are shown in Supplementary Fig. 1; we did not detect a

statistically significant difference between the power law and exponential model errors, paired

t-test, $t(59)=0.71$, $p=0.478$).

Critically, both empirical model types showed that altering washout errors had a dramatic

effect on re-learning rates (Fig. 3d, 1-way ANOVA for exponential model, $F(3,56)=5.49$, $p=0.002$;

Supplementary Fig. 2f for power law, $F(3,56)=5.82$, $p=0.002$). Namely, savings was observed only

after no-feedback washout (post-hoc Dunnett's test, $p=0.004$ for exponential model and $p=0.002$

for power law) and gradual washout (Dunnett's test, $p=0.04$ for exponential model and $p=0.045$

for power law), but not after abrupt washout (post-hoc Dunnett test, $p=0.996$ for exponential

model and $p=0.985$ for power law). In sum, initial gradual adaptation did induce savings, and the

ability to express savings was modulated by errors subjects experienced during washout. When

large errors occurred in our abrupt washout condition, savings was eliminated. When errors were

masked (either by making them small with gradual washout, or removing them via no feedback),

savings was preserved.

Improved rates of re-learning are not due to a bias in initial performance state

Improved re-learning in exposure 2, as shown in Fig. 3, could be driven by two different sources:

(1) a difference in the starting point for re-learning (i.e., the adapted state each subject possesses

at the end of washout; we will refer to this as the 'terminal washout state') or (2) an increased

rate of re-learning irrespective of the starting point for re-learning. This distinction is critical given

that the no-feedback washout group experienced less performance decay during washout (Fig.

4c provides the terminal washout state; 1-way ANOVA, $F(3,56)=10.86$, $p<0.0001$) than other

experimental groups (post-hoc statistics in Supplementary Table 2), meaning that they started

their re-learning period at an advantaged (i.e., larger) learning state.

282
Figure 4. Improved re-learning is not caused by a biased starting point. Exponential and power
law models in Fig. 3 were fit to the entire re-learning period in exposure 2. No-feedback washout
group started this period at a biased state. Inset **a** ('unadjusted') provides the terminal washout
state (average angle on last 4 washout cycles (control shows average on last 4 baseline cycles).
We refit the exponential model to participant data after removing the initial cycles in the re-
learning phase required for Control, Abrupt washout, and Gradual washout groups to 'catch up'
to no-feedback washout starting point. Inset **d** shows the initial reaching angle on the adjusted
starting cycle. The exponential model fits to the adjusted data are shown in Inset **a.** Learning
curves in a are overlaid with control group for ease of comparison. The exponential model fit is
shown overlaid on the measured learning curve. Inset **b** shows the associated rate parameters
for the exponential model. Inset **e**, provides a comparison between re-learning rates (or the naïve
learning rate in control group) and the adaptation starting point (i.e., the terminal washout state).
X-axis provides the terminal washout state for each participant (the collapsed individual points
in **C**). Y-axis provides the learning rates shown in **b**. Both axes were zero-centered at the group
level (i.e., the group mean was subtracted from each data point prior to visualization and
analysis). Statistics in **e** refer to a linear regression. Errors bars in all insets indicate mean ± SEM.
Single dots in **b-e** are individual subjects. Statistics: * indicates $p < 0.05$, ** indicates $p < 0.01$, and
*** indicates $p < 0.001$.

Did no-feedback washout produce faster re-learning because other conditions needed to

'catch up' to their initial adapted state? To answer this question, we conducted a control analysis

where we matched the initial state across groups to that of the no-feedback group (see Methods:

Analyzing relationships between initial state and re-learning rates). That is, we fit the exponential

and power law models to only a portion of the re-learning curve. The starting point for the model

fit was chosen to exclude the initial catch-up period where subjects in the control, abrupt, and

gradual washout groups fell below the initial state of the no-feedback group. The adjusted curves

were defined by the starting points in Fig. 4d; this process removed the initial state bias between

groups (1-way ANOVA, $F_{(3,56)}=1.94$, $p=0.134$). Importantly, we obtained similar results as before;

our model (exponential fit shown in Fig. 4a; power fit shown in Supplementary Fig. 2g) suggested

that learning rates during re-learning periods differed between groups ($F_{(3,56)}=3.9$, $p=0.013$ for

exponential model; $F_{(3,56)} = 4.9$, $p=0.004$ for power model), with no-feedback washout

exhibiting faster re-learning (post-hoc Dunnett's test against control, $p=0.022$ for exponential and

$p=0.007$ for power), a marginal effect in the gradual washout group ($p=0.137$ for exponential and

$p=0.079$ for power), and no effect in the abrupt washout group ($p=1$ for exponential and $p=0.999$

for power).

To corroborate these group-level results, we analyzed the relationship between terminal

washout state and re-learning rates at the individual participant level. We collapsed data across

all four groups and de-trended datasets (see Methods: *Analyzing relationships between initial*

state and re-learning rates) to prevent correlations from being caused by group-level effects.

Linearly regressing re-learning rates (Fig. 4e, y-axis) onto the terminal washout states (Fig. 4e, x-

axis) did not yield a statistically significant correlation ($R=-0.206$, $p=0.114$, $R^2=0.043$), and instead

trended toward an inverse relationship such that the subjects with greater terminal washout

states adapted *more slowly* (counter to the idea that better retention of initial learning speeds

re-learning).

In sum, we did not observe clear evidence for the idea that less performance decay in the

washout phase would lead to faster re-learning. The improved learning rates we observed in Figs.

3 and 4 appeared to reflect an enhanced capacity for learning as opposed to a bias in the starting

point for learning.

Savings following gradual learning is promoted by an increase in sensitivity to error

Short-term sensorimotor adaptation is due to the combined effects of two processes: (1) trial-to-

trial error-based learning, and (2) trial-to-trial performance decay (generally called 'retention' or

'forgetting' in the sensorimotor learning literature)[15,22,28,39–42]. In principle, savings could be

caused by either an increase in the amount of learning that follows the experience of an error,

an increase in the amount of learning retained after time passage, or a combination of both

sources. These two processes can be captured within a state-space model that represents

adaptation as a recursive process controlled by two parameters: an error sensitivity (controls the

amount of learning from error) and a retention factor (controls the amount of decay)[15,20,22,28,39,42].

To determine if the savings we observed following gradual adaptation was due to changes

in error-based learning or performance decay, we fit state-space models to individual subject

learning curves during the second rotation period in all three gradual adaptation groups (Fig. 5a,

green, magenta, blue). To estimate learning and performance decay measures in naïve

participants, we fit a state-space model to the initial abrupt adaptation period in the control

group shown in Fig. 1 (Fig. 5a, black). To ensure that our state-space model required both a

retention parameter and sensitivity to error term, we compared the 'complete' model to a

simpler one where retention was set to a value of 1 (no performance decay). A model comparison

with Akaike's information criterion (AIC) indicated that the complete model was superior in 55 of

60 participants (see Supplementary Fig. 4a to visualize fits for the reduced model; model

comparison is shown in Supplementary Fig. 4b). As such, only the complete parameter model

was used in all subsequent analyses. As with power law and exponential model fits, the state-

space model yielded a close match to individual subject behavior, with the average error falling

within 0.65° of the theoretical minimum value estimated during periods without any adaptation

(see Methods: *Evaluating model error* and Supplementary Fig. 1).

Figure 5. Savings after gradual learning is mediated by an increase in error sensitivity. **a**. Learning
curves and visualization of savings. Reach angle at maximum velocity was calculated on each trial
and averaged within each 4-trial cycle. Each inset compares naïve adaptation (control, black) to
1 of 3 gradual adaptation groups that experienced different washout conditions (left: abrupt
washout; middle: gradual washout; right: no-feedback washout). We fit a state-space learning
model (model) to individual participant reach angles that describes learning as a recursive trial-
to-trial process of error-based adaptation and decay. **b**. Comparison of retention factors.
Retention factors estimated by state-space model fits are compared across the control group
during exposure 1 and 3 gradual adaptation groups during exposure 2. **c**. Comparison of error
sensitivity. Sensitivity-to-error parameters estimated by state-space model fits are compared
across the control group during exposure 1 and 3 gradual adaptation groups during exposure 2.
Errors bars indicated mean ± SE. Single points in **b** and **c** represent individual participants.
Statistics: n.s. indicates $p > 0.05$, ** indicates $p < 0.01$.
To determine if the initial gradual adaptation and subsequent washout period altered the

stability of the motor memory during the readaptation period, we compared retention factors

across the four groups (Fig. 5b). We did not observe a statistically significant change in retention

after exposure to the gradual rotation (one-way ANOVA, $F(3,56)=1.43$, $p=0.244$).

To determine if instead, a change in error sensitivity boosted adaptation rates during the

second exposure period, we compared the state-space model error sensitivity parameters across

the four groups (Fig. 5c). As in past studies[11,16,22], prior adaptation did alter the sensitivity to error

during the second exposure (1-way ANOVA, $F(3,56)=6.99$, $p=0.0004$), reflecting the accelerated

re-learning rates (Fig. 3d). Interestingly, we found a statistically significant increase in error

sensitivity over naïve, in the no-feedback condition (post-hoc Dunnett's test, $p=0.0011$), but not

in the abrupt condition (post-hoc Dunnett's test, $p=0.994$). In the gradual washout condition, we

observed a trend towards an increase in error sensitivity over the naïve condition, though this

trend did not reach statistical significance (post-hoc Dunnett's test, $p=0.123$).

Taken together, this indicates that initial gradual adaptation elevates sensitivity to error,

leading to savings. However, errors experienced during the washout phase can reverse this gain

in error sensitivity. Enhancement in error sensitivity is optimally preserved when visual errors are

totally removed in the washout phase via a no-feedback condition.

Discussion

The conditions required to increase sensorimotor learning rates are not completely known. This

is clearly illustrated in gradual adaptation paradigms where participants adapt to a perturbation

slowly and are then re-tested in an abrupt context following a period of washout. Whereas some

studies have observed more rapid learning in this situation[26,29], other accounts have found that

gradual learning cannot improve abrupt learning[27]. We considered whether this contradiction is

due to the washout phase that separates the two learning phases in savings paradigms.

Specifically, does salience of oppositely signed errors during washout inhibit savings after gradual

learning?

Here we examined this idea by altering the type of error experienced during the washout

phase after gradual learning. We examined no-feedback, gradual, and abrupt washout conditions

that differ along an error size gradient. (1) Removing visual feedback eliminates visuomotor error

completely, causing the adapted behavior to decay in performance in the absence of overt errors

[43]. (2) With gradual washout, errors are present, but are reduced in magnitude because both the

motor action and the perturbation decrease in tandem. (3) Similar to abrupt learning, abrupt

washout results in the largest errors due to the immediate removal of the perturbation when

subjects still remain at a highly adapted performance level. Our findings suggest that the savings

produced by gradual adaptation is protected when washout occurs gradually or in the absence

of error, but not abruptly (Fig. 3d).

The idea that washout impacts savings qualitatively aligns with prior work by Kojima et

al. (2004),[9] which demonstrated that savings in the oculomotor system is sensitive to the

duration of washout: when the duration of the washout phase is increased, the learning rate

during the re-exposure phase reverts to its baseline state, abolishing savings. Thus, both the

duration of error presentation and the error magnitude during washout (as shown in our current

study) seem to negatively interact with the expression of savings. This hypothesis may explain

the mixed results noted above in visuomotor studies of gradual learning. Both Coltman et al.

(2021)[29] and Yin & Wei (2020)[26] employed short washout periods and observed savings following

gradual learning, whereas Herzfeld et al. (2014)[27] applied longer washout periods and did not

find savings after gradual learning. This possibility could easily be studied in future experiments

that vary the length of the washout, along with varying types of washout conditions after an

initial gradual learning period.

Along these lines, Kitago et al. (2013)[43] remarked that increasing the number of washout

trials appeared to dampen the magnitude of savings relative to other studies that used much

fewer washout trials[44,45]. That being said, their observation was made in the context of an abrupt-

then-abrupt savings protocol, instead of the gradual-then-abrupt design we used. The savings

that follows abrupt or gradual learning may have critical differences. For example, we observed

that removing errors in the washout phase after gradual learning protected savings, but the same

washout condition abolished savings after abrupt learning in Kitago et al. (2013)[43] (see Bao et

al.[46] for the case of interlimb transfer).

This discrepancy may be due to variations in the utilization of implicit and explicit learning

processes. Gradual perturbations reduce the error size, in turn limiting one's cognitive awareness

of the perturbation. In visuomotor rotation tasks, reducing the awareness impairs cognitive

strategies that can be used to counter the rotation, allowing subconscious learning (i.e., implicit)

systems to increase their contribution to the adaptation process[22,23,26]. Thus, during the initial

gradual learning period, we expect greater reliance on implicit adaptation, and reduced

engagement of explicit strategies in our tasks, relative to the tasks employed by Kitago et al.

(2013). Further, both Kitago et al. and Bao et al. used a single reaching target, whereas our

current study incorporated four targets. Increasing the target count enhances task complexity,

reducing explicit strategy use and boosting implicit learning (as measured via procedural

aftereffect trials[47,48]). Thus, washout may influence savings differently in our study, where the

initial learning that is being washed out is largely implicit, as compared to the studies by Kitago

et al. and Bao et al., where learning may have relied more heavily on explicit strategies. This idea

matches the observation that savings in gradual-then-abrupt learning paradigms like ours have

been linked to an enhancement in implicit learning rates[26,30], whereas savings in abrupt-then-

abrupt protocols is due to accelerated explicit responses[7,12]. Thus, we hypothesize that washout

has divergent effects on the ability of implicit and explicit systems to save: removing feedback

during washout protects savings in the implicit system, but impairs explicit savings.

That being said, it is critical to note that implicit and explicit processes were not measured

in our study, and thus, our hypothesis remains both speculative and limited. To examine implicit

and explicit contributions to savings in our paradigm, future experiments could assay each system

at regular intervals during learning, washout, and re-learning. One approach would be to include

implicit probe trials (trials where participants are instructed to abandon any explicit strategy that

they are using) to measure the subconscious contribution to their adapted state. Without studies

like this, we cannot definitively attribute the observed savings effect to either the implicit or

explicit process.

To understand what learning property links washout to the capacity to save, we applied

a state-space model that casts short-term adaptation as a cumulative process of trial-to-trial,

error-based learning and decay. This model showed that savings relied on an upregulation in

error sensitivity, which was reversed by large errors in the washout phase (Fig. 5c). This outcome

comports with past work, suggesting that increases in error sensitivity are the primary source of

accelerated re-learning after abrupt, gradual, or random perturbations[29], and the adaptation of

disparate motor systems including reaching[16,22,29] and walking[11]. Intriguingly, our results suggest

that increases in error sensitivity are undermined by opposite errors experienced in between

learning periods.

We found that error sensitivity was higher in the subjects who washed out without

feedback, suggesting that improvements in error sensitivity is protected in the absence of error.

We found a trend towards an increase in error sensitivity in the gradual washout group as well

(Fig. 4c), but this did not reach statistical significance. Because our exponential model (Fig. 3d)

and power law model (Supplementary Fig. 3f) suggested the presence of savings in the gradual

washout group, we suspect that our study lacked the sufficient power to detect an enhancement

in error sensitivity in the gradual washout condition (power was analyzed using learning rates,

see Methods: *Power analysis*). Thus, it remains to be determined whether error sensitivity is

reduced by the experience of *any* opposing error, or whether the size of the opposing error

amplifies the decline in error sensitivity.

Our findings have critical implications for an influential error sensitivity model defined by

a memory of errors[13,27,28]. The memory of errors model that was initially proposed by Herzfeld et

al. (2014)[27] describes changes in error sensitivity based on statistical properties of error. Namely,

when two errors have consistent signs, the nervous system could have adapted more to the initial

error. A critical model property is locality: consistently small errors or large errors modulate error

sensitivity specific to errors of that magnitude. Thus, contrary to our findings, a memory of errors

model predicts that a gradual perturbation will not lead to savings during an abrupt perturbation,

because abrupt errors are larger than those experienced during a gradual perturbation. Thus, our

results add to a growing body of evidence[26,29] suggesting a need to revisit properties of the error

sensitivity landscape; it may be that the experience of consistent errors generalizes more broadly

to unexperienced errors of disparate sizes than previously appreciated.

The possibility that changes in error sensitivity extend beyond similarly sized errors could

also help to understand declines in learning rate in anterograde interference paradigms[22,42,49–51].

That is, whereas increases in error sensitivity have been linked to improvements in learning when

two similar perturbations occur in sequence, a decline in error sensitivity appears to slow learning

when transitioning between two opposing perturbations (e.g., clockwise and counterclockwise

rotations)[42]. The memory of errors model has not been altered to describe this interference

effect. The limited application of the memory of errors model to interference paradigms arises,

because learning in response to "positively signed" errors during the first exposure does not

produce error sensitivity changes that generalize to "negatively signed" errors during the second

exposure. Our findings, that large, oppositely signed errors during washout reverse prior

enhancements in error sensitivity (Fig. 5c), suggest that experience of a sensorimotor error

produces two distinct effects: (1) an increase in sensitivity to similar errors, and (2) a suppression

in error sensitivity to opposite errors. This second suppressive learning property may be a critical

process missing from current memory of errors models that would aid in interpreting both the

dissolution of savings following certain types of washout, as well as interference effects.

We speculate that antagonistic interaction between the response to positive and negative

errors may arise within cerebellar learning circuits that mediate error-based learning, a structure

known to encode error directionality based on anatomical connections with the inferior olive[52–

54]. That is, an error of a given direction (*i.e.*, sign) leads to complex spiking in certain Purkinje-cell

(P-cell) subpopulations in the cerebellar cortex, and suppression in others. Complex spike events

in turn lead to long term potentiation or depression at P-cell synapses with parallel fibers[55]. While

it is unknown how error sensitivity is encoded by P-cells, our findings suggest the possibility that

prolonged absence of complex spikes (when errors are opposite the neuron's preferred direction)

during washout periods may produce synaptic changes that reduce the plastic learning potential

of these cells in the future (*i.e.*, during re-exposure). This idea, and the source of cerebellar error

sensitivity more generally, remain to be examined.

Note that the state-space model posits that a memory is created during the initial learning

period and then decays during washout. Savings reflects the rate at which the original memory

can be reacquired. This idea, that savings is related more to a memory of how to respond to

errors than to a memory of past actions, is supported by various motor learning studies[13,27,28].

However, the psychology of cognitive decision making as well as subconscious classical

conditioning mechanisms, point towards a more nuanced view of learning, namely, as a

composition of multiple memories that compete with one another to be expressed[56]. In this view,

savings depends on inferring the correct memory to draw upon in the current context.

In general, learning involves predictions. Adaptation is achieved by predicting an outcome

and updating one's actions and beliefs based on errors in this prediction. Broadly speaking, errors

encoded in the cerebellum[52,53,57] are also represented within dopaminergic systems[58,59] and can

be modeled via reinforcement learning algorithms[60,61]. This process is distilled to its fundamental

elements in conditioning paradigms, where a stimulus is provided that precedes an appetitive or

aversive outcome with some probability. Such experiments have been used extensively to study

savings: cases where an initial response is acquired, extinguished, and reacquired within different

systems such as eyeblink conditioning, nictitating membrane responses, and salivation in various

species including rabbits, mice, and dogs[62–68]. In Pavlovian tasks, faster reacquisition depends on

two variables: (1) how strongly the memory created during initial exposure generalizes to the re-

exposure period, and (2) how deeply the memory created in extinction (i.e., washout) interferes

with the recall process[56]. That is, memories created during the initial conditioning period are not

abolished by extinction, but instead remain latent and require the appropriate contexts and cues

to promote their re-expression[69,70].

Many factors determine how well conditioned memories can be recalled, such as

similarity between the contexts surrounding the initial acquisition and re-acquisition periods[71,72],

biological significance of the initial learning events[73,74], and the ability to couple the initial context

and the associated memory[75]. Such conditioning rules may help to understand why savings in the

abrupt and gradual contexts explored here can be differentially expressed. The large errors

experienced during an abrupt perturbation are salient, increasing their biological significance.

Further, in tasks with an abrupt-washout-abrupt design, the similarity between the initial

learning and re-learning phases is clearly high, promoting savings. This may be why savings in the

abrupt-washout-abrupt design can withstand an abrupt washout period with large opposing

errors[7,9,22,27], while less salient gradual errors that are less similar to abrupt errors create

memories with increased susceptibility to disruption during an abrupt washout period as we

observed in our study (Fig. 3d).

This same framework can be applied to interpret why washing out without error feedback

best facilitated savings from the gradual adaptation phase to the subsequent abrupt adaptation

phase. That is, washout creates a new pairing between the stimulus (i.e., target presentation in

our study) and the appropriate response[76–78] (e.g., predicting presence of a visual rotation). When

the washout phase is experienced in a similar context to the initial exposure, interference occurs.

For example, Bouton (1986)[79] observed that when extinction persists longer than the initial

acquisition, savings is completely abolished; with even longer washout durations, reacquisition

can further slow and produce a type of retrograde interference[14,50,80,81]. Intriguingly, altering the

context of the learner during the extinction phase in ABA paradigms[76,82] allowed renewal of the

initial memory, because the extinction context formed a separate memory that did not exert

interference. For our study, the no-feedback condition (in which no cursor or outcome is present

on the screen) was the most dissimilar context from the initial learning and re-learning periods

(in which full feedback was provided). For this reason, the washout phase creates a memory in a

separable context, protecting the initial conditioned response from potential interference.

A related, yet distinct, viewpoint suggests that creating a new context during the washout

period is triggered by the salience of prediction errors. When errors are high during washout, this

may be a trigger to create a new memory state instead of updating the original state. For example,

in Gershman et al.'s study (2013),[83] subjects in a fear conditioning paradigm more fully

extinguished their fear memory, when the extinction (i.e., washout) phase was executed

gradually by reducing the probability of the aversive stimulus in a slow continuous manner, as

opposed to all at once. The critical idea was that this gradual extinction would encourage the

animal to associate the shock with their initial memory, as opposed to an extinction-specific

memory. This hypothesis comports with our findings, namely, that when the learned behavior

decays gradually or in the absence of feedback, no washout-specific memory state is formed to

interfere with the initially obtained memory. This would promote more rapid expression of the

initial memory, hence savings.

Multiple memories and statistical inference are used in Bayesian models of sensorimotor

learning. These models incorporate both error-based learning and state inference across multiple

sensorimotor contexts[84,85]. In single-context memory of errors models, the same adaptive states

apply uniformly across different experimental periods, such that any change in learning rate must

be due to variation in constitutive learning parameters: sensitivity to error or memory retention.

But with multiple contexts, as in the COIN model described by Heald et al. (2021)[84], the apparent

learning rate is influenced by inferring which context is most likely given sensory information. For

the COIN model, visuomotor adaptation involves two contexts: a baseline context and a rotation

context. Faster learning during the second exposure is caused not by a change in the response to

error, but by a greater propensity to expect the rotation given prior training. Our findings have

important implications for this inference step, namely, that the errors (*i.e.*, sensory information)

experienced during washout alter the inference process: with large opposing errors during the

washout period, the nervous system decreases its expectation that the initial rotation state (with

its positively-signed errors) will occur again in the future. A broader comparison between

memory of errors models and Bayesian frameworks on the effects of washout on future learning

remains to be conducted.

Methods

Participants

Initially, 10 healthy young adults per group were tested. We then conducted a power analysis to

determine the necessary group size to achieve statistical significance with 80% power. The data

from these 40 participants (10 per group) were used to estimate effect sizes for this analysis. Our

power analysis indicated that a minimum of 13 participants per group were required to achieve

80% power (see *Power Analysis* below for details*)*. Based on this, we collected an additional 5

participants per group, resulting in n=15 per group (60 participants in total). All subjects were

right-handed as assessed by the Edinburgh handedness inventory (Oldfield, 1971). The subjects

were randomly assigned to the following groups: one abrupt adaptation (i.e., control) group and

three gradual adaptation groups. All experimental protocols were approved by the University of

Wisconsin-Milwaukee (UWM) Institutional Review Board (IRB). Subjects gave written informed

consent prior to participation, which was approved by the UWM IRB in accordance with the

Declaration of Helsinki (IRB #16.163).

Apparatus

Participants were situated in a robotic exoskeleton (KIMARM, BKIN Technologies Ltd, Kingston,

ON, Canada) that provided gravitational support to the right arm (all subjects used this arm in

the present study). The exoskeleton was positioned so that the arm was hidden underneath a

horizontal display (Fig. 1a). To track the hand's position, a small cursor was projected onto the

display, over the subject's index fingertip. During each trial, the KINARM projected visual stimuli

onto the display, so that they appeared in the same plane as the arm. The visual stimuli consisted

of a centrally located start circle (2 cm in diameter) and one of four target circles (2 cm in

diameter) located 10 cm away from the start target (Fig. 1a). We sampled the hand's position in

the x-y plane at 1000 Hz. Position data were low-pass filtered at 15 Hz, and then differentiated

to calculate velocity. Post-processing, analysis, and modeling were conducted in MATLAB R2018a

(The MathWorks Inc., Natick, MA).

Experimental Design

All subjects experienced 4 experiment phases: (1) a baseline period, (2) an initial learning period

(exposure 1), (3) a washout period, and (4) a re-learning period (exposure 2) (Fig. 1b). In the

baseline period (Fig. 1b, baseline), participants moved their arm to each of the four targets over

a 10-cycle period (all four targets were presented once in a cycle). On each trial, continuous visual

feedback was provided via a cursor indicating the location of the index fingertip. Participants

were instructed to move rapidly and accurately to the target location. The trial ended 1.5 sec

after target presentation. To begin the next trial, the participants brought their hand back to the

central start position. Throughout the entire experiment, the four targets were presented in a

pseudorandom order in 4-trial cycles.

During the initial learning period (exposure 1), a 24° rotation was introduced abruptly for

the abrupt adaptation (control) group (see Fig. 1), and gradually for three gradual adaptation

groups (see Figs. 2-5). More specifically, the visual display of the cursor in the control group was

rotated 24° counterclockwise (CCW) about the start circle from trial 1 and remained constant

throughout the entire exposure period. In the gradual adaptation groups, the rotation size

started at 0° for the first 9 cycles, then increased by 1° (CCW about the start circle) per cycle for

next 23 cycles, and remained at 24° for last 8 cycles. During the re-learning period (exposure 2),

all participants experienced the same abrupt 24° rotation condition as that experienced by the

control group in the initial learning period. All participants performed reaching movements over

40 cycles (i.e., 160 trials) in each of the two exposure periods.

In the washout period, the control group performed reaching movements without visual

feedback. The three gradual adaptation groups experienced one of three washout conditions: (1)

abrupt washout, in which the rotation was completely removed from trial 1 (i.e., the same as the

baseline period); (2) gradual washout, in which the rotation size was decreased gradually from

24° down to 0°); and (3) no-feedback washout, in which subjects performed reaching movements

without visual feedback. All participants performed reaching movements over a 40-cycle period.

In sum, all groups re-learned in an abrupt context. Thus, the four groups were separated by their

initial learning and washout conditions. The abrupt adaptation group (controls) is the group that

initially learned abruptly and washed out without feedback. The gradual adaptation/gradual

washout group is the group that initially learned gradually and then washed out gradually. The

gradual adaptation/abrupt washout group is the group that initially learned gradually and

washed out abruptly. The gradual adaptation/no-feedback washout group is the group that

initially learned gradually and washed out without feedback.

Behavioral data processing

Data analysis was conducted in MATLAB (MathWorks, Natick, MA). To quantify performance, we

calculated the reaching movement angle at peak velocity (c.f., direction error[37]). This angle was

computed as the hand's angular position relative to a line segment connecting the start and target

positions. Outlying reach angles were detected and removed as detailed in *Outlier Detection* prior

to additional analysis. Next, data were averaged within each 4-trial cycle.

Comparing gradual and abrupt learning rates

To compare learning rates between gradual and abrupt adaptations, we analyzed learning curves

in the exposure 1 period with a linear model. For gradual learning, the linear model incorporated

a delay parameter that allowed the starting point at which learning begins to vary between each

participant (the parameter was allowed to vary between 9 and 20 cycles: note that 9 cycles were

chosen because the rotation was held at 0° for 9 cycles). The model was fit to individual subject

behavior and the delay parameter that minimized squared error was selected. Because abrupt

learning is not well-captured by a linear model (see Fig. 1c) only the initial 8 cycles were used to

quantify an 'early' learning rate to compare to the gradual condition. No delay was allowed in

the linear fit. Model fits are shown in Supplementary Fig. 3a, and learning rates (i.e., linear slope)

are provided in Supplementary Fig. 3b. Abrupt exposure to the rotation more than doubled

learning rates (1-way ANOVA, $F(3,56)=17.54$, $p < 1e^{-7}$; all post-hoc comparisons between abrupt

group and each of the 3 gradual learning groups via Tukey's test yielded $p<0.0001$; other

comparisons $p > 0.97$). See Supplementary Table 1 for post-hoc t-test statistics.

Exponential and power law models

To empirically compare adaptation between groups, we assessed the learning rate during each

abrupt rotation period via an exponential function. We focused on rate measures, to prevent our

evaluation from being biased by the initial state of the learner (*i.e.*, subjects may demonstrate

greater total adaptation not because they learn more rapidly, but because their initial reach angle

is greater during the second abrupt rotation due to incomplete washout from the initial rotation).

Thus, we fitted an exponential curve to participant hand angles, and used the rate parameter as

our measure of learning rate. For this, we employed a 2-parameter exponential function designed

to enforce that each curve begin at the initial pre-rotation reach angle. For the initial rotation in

the control group, this hand angle was computed as the average angle over the last 4 baseline

cycles. For the abrupt rotations during the second exposure period (across all groups), this angle

was computed as the average angle over the last 4 washout cycles. The exponential function took

the mathematical form:

$$y(t) = \alpha - (\alpha - y_0)e^{-\beta t}$$ (1)

Here, α is parameter that encodes the exponential curve's asymptote, y_0 is the initial reach angle

described above, β is the rate parameter of interest, $y(t)$ is the reach angle on cycle t (starting at

$t = 0$). The function was fit to 41 cycles of data at a time (40 cycles of the abrupt rotation plus the

initial reach angle cycle on $t = 0$) using a bounded least-squares approach (MATLAB *fmincon*). The

parameter search was repeated 100 times, each time varying the initial parameter estimate that

seeds the algorithm. The α parameter was bounded between -45° and 45°. The β parameter was

bounded between 0 cycle^{-1} and 0.5 cycle^{-1}. The model parameters that minimized squared error

across all 100 repetitions were selected.

Exponential models are commonly used in sensorimotor learning studies to assess

adaptation rates. For learning in the psychological domain, models with a positive hazard such as

a power law are more common[33–35]. Thus, to corroborate our exponential model, we repeated

our primary analyses using a 2-parameter power law given by:

$$y(t) = a - (a - y_0)t^{-b}$$ (2)

Similar to our exponential model, the power function was constrained to begin at the terminal

washout state (y_0) by starting with $t = 1$. In this way a sets asymptotic learning and b is the power

parameter that we used to quantify and compare learning rates. As with the exponential model,

fmincon was used to fit the model to individual participant data in a space where *a* was bounded

between 0 and 45° and *b* was bounded between 0 and 5.

Analyzing relationships between initial state and re-learning rates

We conducted a set of control analyses (results shown in Fig. 4) to determine how the initial state

at the start of the re-exposure period altered learning rate. Note that this initial state at the start

of exposure 2 is synonymous with the terminal washout state (as the washout directly precedes

exposure 2). Two analyses were conducted: one at the group-level and one at the subject-level.

708   For our group-level analyses we repeated our exponential model fit to a smaller segment

of the re-exposure period (or the initial exposure period in the control group). We selected this

period based on the no-feedback group. This group exhibited greater lingering adaptation at the

end of the washout phase (see Fig. 4c), meaning that they would begin adaptation at the start of

exposure 2 at a higher level than the other groups. To control for this, we identified new starting

points for exponential model fits that more closely matched the mean starting point exhibited in

the no-feedback group (6.203°). To do this, we used a 2-step approach. First, we used exponential

model fits to the entire exposure period (as described in *Exponential and power law models*) and

identified when these models crossed the threshold value of 6.203° for each participant. We then

trimmed the data to remove all data points prior to this threshold value, such that the beginning

of the adjusted learning curves corresponded to the cycle just prior to the threshold crossing (the

resulting initial states are shown in Fig. 4d). We then refit the exponential model to the truncated

data, using the exact same process as described in *Exponential and power law models*. Resulting

learning rate parameters are reported in Fig. 4b). This process was repeated (using the same start

points) for the power law model, and results are shown in Supplementary Figs. 2g and 2h.

Finally, we compared learning rates during the re-exposure period (or naïve exposure for

the control group) to their terminal reach angle at the end of washout (or the end of the baseline

period for the naïve control group). This was done to test the hypothesis that the subjects who

had an initial bias towards the adapted state might exhibit an unfair advantage in their learning

rate. For this analysis, the exponential rate parameters were compared to the average reach

angle at the end of washout (the 4-cycle period described in *Exponential and power law models*).

For this, data were collapsed across the four groups (n=60 participants in total). Data were de-

trended at the group level by zero-centering; the group mean was subtracted within each group

prior to analysis. The resulting data (Fig. 4e) were then compared across participants via linear

regression.

State-space learning model

To interpret our empirical results using an error-based learning framework, we employed a state-

space model[15,22,28,39,40,42]. The model describes adaptation as a trial-by-trial learning process

whereby the adapted state of the individual is updated by (1) learning from the error experienced

on each trial and (2) a small amount of decay that reflects temporal instability of short-term

adaptation[41,86,87]. In the model, the total update in the motor memory due to error is controlled

by one's sensitivity to error (b) and the strength of memory retention between trials is encoded

by a retention factor (a). Together, learning and performance decay (this is termed 'forgetting'

or 'retention' in the sensorimotor adaptation literature) collectively determine how the subject's

internal state (x) evolves in response to error (e) experienced on trial t in the presence of Gaussian

state noise (ε_x, normally distributed with zero mean, and standard deviation of σ_x) according to:

$$x^{(t+1)} = ax^{(t)} + be^{(t)} + \varepsilon_x^{(t)} \qquad (2)$$

Eq. (2) allows us to ascribe performance differences during the adaptation period to interpretable

quantities: retention (a) and error sensitivity (b).

Note, however, that the internal state (x) is not observable. Instead, on each cycle, the

motor output (reach angle) is measured. The reach angle (y) directly reflects the subject's internal

state, but is altered by execution noise (ε_y, normal with zero mean, std. dev. = σ_y) according to:

$$y^{(t)} = x^{(t)} + \varepsilon_y^{(t)} \qquad (3)$$

Together, Eqs. (2) and (3) represent a single module state-space model. We fit this model to each

participant's reach angles during the adaptation period using the Expectation-Maximization (EM)

algorithm[39]. For each adaptation period, all 40 abrupt rotation cycles and 3 preceding baseline or

washout cycles were included. EM is an algorithm that conducts maximum likelihood estimation

in an iterative process. We used EM to identify the parameters $\{a, b, x_1, \sigma_x, \sigma_y, \sigma_1\}$ that maximized

the likelihood of observing the data (x_1 and σ_1 represent the subject's initial state and variance,

respectively). We conducted 100 iterations, each time modifying the initial parameter guess that

seeded the algorithm. The retention factor (a) was bounded between 0.85 and 1. The initial state

(x_1) was bounded between -10 and 20°. Error sensitivity (b) was bounded between 0 and 1. Last,

all 3 variance parameters were bounded between $1.0e^{-3}$ and 100. We selected the parameter set

that maximized the likelihood function across all 100 iterations.

Last, we conducted a control analysis to assess if a reduced model without performance

decay (i.e., complete memory retention; $a = 1$) was sufficient to explain behavior. The same fitting

procedure described above was used to fit the state-space model, but the retention parameter

was constrained to a value of 1, thereby reducing model complexity by one parameter. The

resulting model fits are provided in Supplementary Fig. 4a. Akaike's Information Criterion (AIC)

was used to compare the complete model to the reduced model in Supplementary Fig. 4b.

Outlier detection

As noted above in *Behavioral data processing*, outlier reach angles were removed prior to model

fits, empirical analyses, and statistical testing. To remove outliers, we used a 2-step process which

started by eliminating trials where the absolute reach angle exceeded 60° (0.567% of trials). Then,

in the second pass, we removed additional outlier trials in each cycle. For this, we collapsed reach

angles across all subjects and 4 trials in each cycle, and removed trials that deviated by more than

5 median absolute deviations from the cycle median (1.798% of trials). This was done separately

within each experimental group, such that each cycle removal process used 60 trials (4 trials in a

cycle, multiplied by 15 participants in the group).

In a control analysis, we employed an alternative way to detect outliers within individual

participants without considering other subjects in the group. For this, we binned reach angles in

consecutive 10-trial bins. In each bin we calculated the median reach angle. Outliers were taken

as trials that deviated from the median by more than a threshold level. We varied this threshold

to explore more liberal and conservate outlier detection; 4 levels were used (from most to least

aggressive: 15°, 20°, 25°, and 30°) leading to removal of 0.68% to 2.6% of trials. Resulting analysis

was done based on our exponential model learning rates (Supplementary Fig. 5). We found that

all thresholds produced the same qualitative result (statistics are given in Supplementary Tables

3 and 4), which also matched the conclusions reported in our Results section.

Evaluating model error

The goodness-of-fit for each model (exponential, power, and state-space model) was quantified

by calculating the root-mean-squared-error (RMSE) for each individual participant learning curve

during the abrupt learning period (exposure 2 was used for the initial gradual learning groups but

exposure 1 was used for the control group). All RMSE measures are shown in Supplementary Fig.

1. For comparison, we also calculated the intrinsic variability in cycle-to-cycle reach angle (shown

as a horizontal line in Supplementary Fig. 1). This intrinsic variability provides a theoretical lower

bound for the RMSE that could be achieved by any model, as it represents the inherent random

movement noise that corrupts movement angles during baseline periods without any adaptation.

To calculate this value, we calculated the standard deviation in angle during the baseline period

as well as the last 10 cycles of the washout period, and averaged these two values together (note

that in the no-feedback group, we used only the baseline standard deviation because during the

washout period the rotation had not been removed long enough to achieve a stable steady-state

at which to assess motor noise in the absence of learning).

Statistical analysis

For statistical comparisons between 2 groups or conditions, paired or independent two-sample

t-tests were employed. When more than 2 groups were compared, we applied one-way ANOVAs.

In post-hoc tests on savings-related measures (see *Measuring savings* below), we used Dunnett's

test to measure differences between each experimental group and the control. For cases where

initial learning rates and terminal washout states were compared across groups, Tukey's test was

used to evaluate all possible pair-wise comparisons.

Measuring savings

Our principal analyses concern whether experiencing a perturbation gradually during initial

learning produces a benefit for re-learning in the future. For sensorimotor adaptation, savings

refers to an increase in the rate or extent of re-learning. However, in many cases, including our

current study, initial learning rates during gradual learning cannot be observed due to a mismatch

between perturbation types in exposure 1 and 2. In exposure 1, participants adapt to a gradual

rotation. In exposure 2, the rotation is abrupt. The gradual and abrupt learning rates are not

comparable: by design, gradual learning is necessarily slower than abrupt learning due to the

presence of smaller errors[28,88,89] throughout the learning process which are known to drive

smaller trial-to-trial updates in reach angle[28,88,89]. In our case, abrupt learning was more than

twice as rapid as gradual learning (see *Comparing gradual and abrupt learning rates* and

Supplementary Fig. 3).

In cases such as ours where initial learning is gradual and re-learning is abrupt, savings is

assessed by comparing learning rates during the abrupt re-learning period to that of a separate

control group that only experiences an abrupt rotation (this control condition allows us to obtain

a proxy estimate for the naïve abrupt learning rate)[13,26,27,29,36–38]. This is the approach we utilized

herein, and it informed our choice of statistical tests described in the previous section.

Power analysis

To determine our group size, we conducted a power analysis. As noted before, we recruited 10

participants in each group and used these preliminary results in our analysis (n=40 subjects total).

For our analysis, we selected our principal investigation in Fig. 3. Exponential models were fit for

exposure 2 in three experimental groups (groups that experienced gradual adaptation during

exposure 1) and exposure 1 in the naïve control condition. The fitting procedure is described

above in *Exponential and power law models*. We then extracted the learning rate for each

participant and calculated the mean and variance for each group. For each group, we drew

simulated learning rates from a normal distribution, and then conducted a 1-way ANOVA. Power

was measured based on the finding that $p < 0.05$ for a given simulated experiment. The resulting

power is reported in Supplementary Fig. 6 as a function of group size. Note that 13 subjects were

required in our simulations to reach 80% power.

Competing interests

The authors declare no competing interests.

Data availability

The datasets used and/or analyzed during the current study will be made available upon

request by contacting the corresponding author.

Code availability

The MATLAB scripts used for empirical analyses, model fitting, model comparison, and

statistical testing will be made available upon request by contacting the corresponding author.

Author contributions

MJ, BM and JW designed the experiments and methodology. MJ and BM collected data. MJ and

STA analyzed data and created figures. MJ and STA wrote manuscript. MJ, STA, and JW edited

the manuscript.

Acknowledgements

This work was supported in part by the National Institute of Child Health and Human

Development (T32HD040127 to STA), the W.M. Keck Foundation (to STA) and a Burroughs

Wellcome Fund Careers at the Scientific Interface Award (to STA).

References

1. Tsay, J. S., Haith, A. M., Ivry, R. B. & Kim, H. E. Interactions between sensory prediction error and task error during implicit motor learning. *PLoS Comput Biol* **18**, e1010005- (2022).
2. Taylor, J. A. & Ivry, R. B. Flexible cognitive strategies during motor learning. *PLoS Comput Biol* **7**, e1001096 (2011).
3. Leow, L.-A., Marinovic, W., de Rugy, A. & Carroll, T. J. Task errors drive memories that improve sensorimotor adaptation. *Journal of Neuroscience* (2020) doi:10.1523/JNEUROSCI.1506-19.2020.
4. Mazzoni, P. & Krakauer, J. W. An implicit plan overrides an explicit strategy during visuomotor adaptation. *Journal of Neuroscience* **26**, 3642–3645 (2006).
5. Albert, S. T., Blaum, E. C. & Blustein, D. H. Sensory prediction error drives subconscious motor learning outside of the laboratory. *J Neurophysiol* **130**, 427–435 (2023).
6. Ebbinghaus, H. *Uber Das Gedachtnis*. (Dunacker and Humblot, Leipzig, Germany, 1885).
7. Haith, A. M., Huberdeau, D. M. & Krakauer, J. W. The Influence of Movement Preparation Time on the Expression of Visuomotor Learning and Savings. *Journal of Neuroscience* **35**, 5109–5117 (2015).
8. Landi, S. M., Baguear, F. & Della-Maggiore, V. One week of motor adaptation induces structural changes in primary motor cortex that predict long-term memory one year later. *J Neurosci* **31**, 11808–11813 (2011).
9. Kojima, Y., Iwamoto, Y. & Yoshida, K. Memory of Learning Facilitates Saccadic Adaptation in the Monkey. *Journal of Neuroscience* **24**, 7531–7539 (2004).
10. Malone, L. A., Vasudevan, E. V. L. & Bastian, A. J. Motor Adaptation Training for Faster Relearning. *Journal of Neuroscience* **31**, 15136–15143 (2011).
11. Mawase, F., Shmuelof, L., Bar-Haim, S. & Karniel, A. Savings in locomotor adaptation explained by changes in learning parameters following initial adaptation. *J Neurophysiol* **111**, 1444–1454 (2014).
12. Morehead, J. R., Qasim, S. E., Crossley, M. J. & Ivry, R. Savings upon Re-Aiming in Visuomotor Adaptation. *Journal of Neuroscience* **35**, 14386–14396 (2015).
13. Leow, L.-A., de Rugy, A., Marinovic, W., Riek, S. & Carroll, T. J. Savings for visuomotor adaptation require prior history of error, not prior repetition of successful actions. *J Neurophysiol* **116**, 1603–1614 (2016).
14. Brashers-Krug, T., Shadmehr, R. & Bizzi, E. Consolidation in human motor memory. *Nature* **382**, 252–255 (1996).
15. Smith, M. A., Ghazizadeh, A. & Shadmehr, R. Interacting adaptive processes with different timescales underlie short-term motor learning. *PLoS Biol* **4**, e179 (2006).
16. Coltman, S. K., Cashaback, J. G. A. & Gribble, P. L. Both fast and slow learning processes contribute to savings following sensorimotor adaptation. *J Neurophysiol* **121**, 1575–1583 (2019).
17. Medina, J. F., Garcia, K. S. & Mauk, M. D. A mechanism for savings in the cerebellum. *Journal of Neuroscience* **21**, 4081–4089 (2001).
18. Taylor, J. A., Krakauer, J. W. & Ivry, R. B. Explicit and Implicit Contributions to Learning in a Sensorimotor Adaptation Task. *Journal of Neuroscience* **34**, 3023–3032 (2014).

19. McDougle, S. D. & Taylor, J. A. Dissociable cognitive strategies for sensorimotor learning.
Nat Commun **10**, 40 (2019).
20. McDougle, S. D., Bond, K. M. & Taylor, J. A. Explicit and Implicit Processes Constitute the
Fast and Slow Processes of Sensorimotor Learning. *Journal of Neuroscience* **35**, 9568–9579
(2015).
21. Shadmehr, R., Smith, M. A. & Krakauer, J. W. Error correction, sensory prediction, and
adaptation in motor control. *Annu Rev Neurosci* **33**, 89–108 (2010).
22. Albert, S. T. *et al.* Competition between parallel sensorimotor learning systems. *Elife* **11**,
e65361 (2022).
23. Neville, K.-M. & Cressman, E. K. The influence of awareness on explicit and implicit
contributions to visuomotor adaptation over time. *Exp Brain Res* **236**, 2047–2059 (2018).
24. Salomonczyk, D., Cressman, E. K. & Henriques, D. Y. P. Proprioceptive recalibration following
prolonged training and increasing distortions in visuomotor adaptation. *Neuropsychologia*
49, 3053–3062 (2011).
25. Huberdeau, D. M., Krakauer, J. W. & Haith, A. M. Practice induces a qualitative change in
the memory representation for visuomotor learning. *J Neurophysiol* **122**, 1050–1059 (2019).
26. Yin, C. & Wei, K. Savings in sensorimotor adaptation without an explicit strategy. *J*
Neurophysiol **123**, 1180–1192 (2020).
27. Herzfeld, D. J., Vaswani, P. A., Marko, M. K. & Shadmehr, R. A memory of errors in
sensorimotor learning. *Science (1979)* **345**, 1349–1353 (2014).
28. Albert, S. T. *et al.* An implicit memory of errors limits human sensorimotor adaptation. *Nat*
Hum Behav (2021) doi:10.1038/s41562-020-01036-x.
29. Coltman, S. K., van Beers, R. J., Medendorp, W. P. & Gribble, P. L. Sensitivity to error during
visuomotor adaptation is similarly modulated by abrupt, gradual, and random perturbation
schedules. *J Neurophysiol* **126**, 934–945 (2021).
30. Wang, J., Joshi, M. & Lei, Y. The extent of interlimb transfer following adaptation to a novel
visuomotor condition does not depend on awareness of the condition. *J Neurophysiol* **106**,
259–64 (2011).
31. Avraham, G., Morehead, J. R., Kim, H. E. & Ivry, R. B. Reexposure to a sensorimotor
perturbation produces opposite effects on explicit and implicit learning processes. *PLoS Biol*
19, e3001147 (2021).
32. James, R., Bao, S., D'Amato, A. & Wang, J. The nature of savings associated with a
visuomotor adaptation task that involves one arm or both arms. *Hum Mov Sci* **81**, 102896
(2022).
33. Newell, A. & Rosenbloom, P. S. Mechanisms of skill acquisition and the law of practice. in
Cognitive skills and their acquisition (ed. Anderson, J. R.) 1–55 (Erlbaum, Hillsdale, NJ, 1981).
34. Donner, Y. & Hardy, J. L. Piecewise power laws in individual learning curves. *Psychon Bull*
Rev **22**, 1308–19 (2015).
35. Anderson, J. R. *et al.* An Integrated Theory of the Mind. *Psychol Rev* **111**, 1036–1060 (2004).
36. Roemmich, R. T. & Bastian, A. J. Two ways to save a newly learned motor pattern. *J*
Neurophysiol **113**, 3519–3530 (2015).
37. Orban de Xivry, J.-J. & Lefèvre, P. Formation of model-free motor memories during motor
adaptation depends on perturbation schedule. *J Neurophysiol* **113**, 2733–2741 (2015).

38. Turnham, E. J. A., Braun, D. A. & Wolpert, D. M. Facilitation of learning induced by both
random and gradual visuomotor task variation. *J Neurophysiol* **107**, 1111–22 (2012).

39. Albert, S. T. & Shadmehr, R. Estimating properties of the fast and slow adaptive processes
during sensorimotor adaptation. *J Neurophysiol* **119**, 1367–1393 (2018).

40. Orozco, S. P., Albert, S. T. & Shadmehr, R. Adaptive control of movement deceleration
during saccades. *PLoS Comput Biol* **17**, e1009176–e1009176 (2021).

41. Brennan, A. E. & Smith, M. A. The Decay of Motor Memories Is Independent of Context
Change Detection. *PLoS Comput Biol* **11**, e1004278 (2015).

42. Lerner, G. *et al.* The Origins of Anterograde Interference in Visuomotor Adaptation. *Cerebral*
Cortex **30**, 4000–4010 (2020).

43. Kitago, T., Ryan, S. L., Mazzoni, P., Krakauer, J. W. & Haith, A. M. Unlearning versus savings
in visuomotor adaptation: comparing effects of washout, passage of time, and removal of
errors on motor memory. *Front Hum Neurosci* **7**, 307 (2013).

44. Zarahn, E., Weston, G. D., Liang, J., Mazzoni, P. & Krakauer, J. W. Explaining Savings for
Visuomotor Adaptation: Linear Time-Invariant State-Space Models Are Not Sufficient. *J*
Neurophysiol **100**, 2537–2548 (2008).

45. Huang, V. S., Haith, A., Mazzoni, P. & Krakauer, J. W. Rethinking motor learning and savings
in adaptation paradigms: model-free memory for successful actions combines with internal
models. *Neuron* **70**, 787–801 (2011).

46. Bao, S., Wang, J., Wright, D. L., Buchanan, J. J. & Lei, Y. The decay and consolidation of
effector-independent motor memories. *Sci Rep* **12**, 3131 (2022).

47. Tsay, J. S. *et al.* Large-scale citizen science reveals predictors of sensorimotor adaptation.
Nat Hum Behav (2024) doi:10.1038/s41562-023-01798-0.

48. Wang, J., Bao, S. & Tays, G. D. Lack of generalization between explicit and implicit
visuomotor learning. *PLoS One* **14**, e0224099 (2019).

49. Sing, G. C. & Smith, M. A. Reduction in learning rates associated with anterograde
interference results from interactions between different timescales in motor adaptation.
PLoS Comput Biol **6**, e1000893 (2010).

50. Caithness, G. *et al.* Failure to consolidate the consolidation theory of learning for
sensorimotor adaptation tasks. *Journal of Neuroscience* **24**, 8662–8671 (2004).

51. Miall, R. C., Jenkinson, N. & Kulkarni, K. Adaptation to rotated visual feedback: a re-
examination of motor interference. *Exp Brain Res* **154**, 201–210 (2004).

52. Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of error and learning to
correct that error by the Purkinje cells of the cerebellum. *Nat Neurosci* **21**, 736–743 (2018).

53. Herzfeld, D. J., Kojima, Y., Soetedjo, R. & Shadmehr, R. Encoding of action by the Purkinje
cells of the cerebellum. *Nature* **526**, 439–442 (2015).

54. Soetedjo, R., Kojima, Y. & Fuchs, A. F. Complex spike activity in the oculomotor vermis of the
cerebellum: a vectorial error signal for saccade motor learning? *J Neurophysiol* **100**, 1949–
1966 (2008).

55. Suvrathan, A., Payne, H. L. & Raymond, J. L. Timing Rules for Synaptic Plasticity Matched to
Behavioral Function. *Neuron* **92**, 959–967 (2016).

56. Bouton, M. E. Context, time, and memory retrieval in the interference paradigms of
Pavlovian learning. *Psychol Bull* **114**, 80–99 (1993).

57. Ohmae, S. & Medina, J. F. Climbing fibers encode a temporal-difference prediction error
during cerebellar learning in mice. *Nat Neurosci* **18**, 1798–1803 (2015).
58. Schultz, W., Dayan, P. & Montague, P. R. A Neural Substrate of Prediction and Reward.
Science (1979) **275**, 1593–1599 (1997).
59. Gershman, S. J. *et al.* Explaining dopamine through prediction errors and beyond. *Nat
Neurosci* **27**, 1645–1655 (2024).
60. Rescorla, R. A. & Wagner, A. A theory of Pavlovian conditioning: Variations in the
effectiveness of reinforcement and nonreinforcement. in *Classical Conditioning II: Current
Research and Theory* vol. Vol. 2 (1972).
61. Soto, F. A. & Wasserman, E. A. Error-driven learning in visual categorization and object
recognition: a common-elements model. *Psychol Rev* **117**, 349–81 (2010).
62. Konorski, J. & Szwejkowska, G. Chronic extinction and restoration of conditioned reflexes. I.
Extinction against the excitatory background. *Acta Biol Exp (Warsz)* **15**, 155–170 (1950).
63. Konorski, J. & Szwejkowska, G. Chronic extinction and restoration of conditioned reflexes:
III. Defensive motor reflexes. *Acta Biol Exp (Warsz)* **16**, 91–94 (1952).
64. Shurrager, P. S. & Culler, E. Conditioning in the spinal dog. *J Exp Psychol* **26**, 133–159 (1940).
65. Frey, P. W. & Butler, C. S. Extinction after aversive conditioning: An associative or
nonassociative process? *Learn Motiv* **8**, 1–17 (1977).
66. Frey, P. W. & Ross, L. E. Classical conditioning of the rabbit eyelid response as a function of
interstimulus interval. *J Comp Physiol Psychol* **65**, 246–250 (1968).
67. Hoehler, F. K., Kirschenbaum, D. S. & Leonard, D. W. The effects of overtraining and
successive extinctions upon nictitating membrane conditioning in the rabbit. *Learn Motiv* **4**,
91–101 (1973).
68. Napier, R. M., Macrae, M. & Kehoe, E. J. Rapid reacquisition in conditioning of the rabbit's
nictitating membrane response. *J Exp Psychol Anim Behav Process* **18**, 182–192 (1992).
69. Spear, N. E. Retrieval of memories: A psychobiological approach. in *Handbook of learning
and cognitive processes: Vol 4. Attention and memory* (ed. Estes, W. K.) 17–90 (Erlbaum,
Hillsdale, NJ, 1976).
70. Spear, N. E. *The Processing of Memories: Forgetting and Retention*. (Erlbaum, Hillsdale, NJ,
1978).
71. McGeoch, J. A. Forgetting and the law of disuse. *Psychol Rev* **39**, 352–370 (1932).
72. Tulving, E. Cue-dependent forgetting. *Am Sci* **62**, 74–82 (1974).
73. Devenport, L. Sampling behavior and contextual change. *Learn Motiv* **20**, 97–114 (1989).
74. Hendersen, R. W. Fearful memories: The motivational significance of forgetting. in *Affect,
conditioning, and cognition: Essays on the determinants of behavior* (eds. Brush, F. R. &
Overmier, J. B.) 43–54 (Erlbaum, Hillsdale, NJ, 1985).
75. Baddeley, A. D. Domains of recollection. *Psychol Rev* **89**, 708–729 (1982).
76. Bouton, M. E., Todd, T. P., Vurbic, D. & Winterbauer, N. E. Renewal after the extinction of
free operant behavior. *Learn Behav* **39**, 57–67 (2011).
77. Bouton, M. E. Context, ambiguity, and unlearning: sources of relapse after behavioral
extinction. *Biol Psychiatry* **52**, 976–986 (2002).
78. Bouton, M. E. Context and Behavioral Processes in Extinction: Table 1. *Learning & Memory*
11, 485–494 (2004).

79. Bouton, M. E. Slow reacquisition following the extinction of conditioned suppression. *Learn*
Motiv **17**, 1–15 (1986).
80. Krakauer, J. W., Ghez, C. & Ghilardi, M. F. Adaptation to visuomotor transformations:
consolidation, interference, and forgetting. *J Neurosci* **25**, 473–478 (2005).
81. Bouton, M. E. & Swartzentruber, D. Slow reacquisition following extinction: Context,
encoding, and retrieval mechanisms. *J Exp Psychol Anim Behav Process* **15**, 43–53 (1989).
82. Todd, T. P., Winterbauer, N. E. & Bouton, M. E. Effects of the amount of acquisition and
contextual generalization on the renewal of instrumental behavior after extinction. *Learn*
Behav **40**, 145–157 (2012).
83. Gershman, S. J., Jones, C. E., Norman, K. A., Monfils, M.-H. & Niv, Y. Gradual extinction
prevents the return of fear: implications for the discovery of state. *Front Behav Neurosci* **7**,
(2013).
84. Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference underlies the learning of
sensorimotor repertoires. *Nature* **600**, 489–493 (2021).
85. Haruno, M., Wolpert, D. M. & Kawato, M. MOSAIC Model for Sensorimotor Learning and
Control. *Neural Comput* **13**, 2201–2220 (2001).
86. Joiner, W. M. & Smith, M. A. Long-term retention explained by a model of short-term
learning in the adaptive control of reaching. *J Neurophysiol* **100**, 2948–2955 (2008).
87. Alhussein, L., Hosseini, E. A., Nguyen, K. P., Smith, M. A. & Joiner, W. M. Dissociating effects
of error size, training duration, and amount of adaptation on the ability to retain motor
memories. *J Neurophysiol* **122**, 2027–2042 (2019).
88. Kim, H. E., Morehead, J. R., Parvin, D. E., Moazzezi, R. & Ivry, R. B. Invariant errors reveal
limitations in motor correction rather than constraints on error sensitivity. *Commun Biol* **1**,
19 (2018).
89. Wei, K. & Kording, K. Relevance of error: what drives motor adaptation? *J Neurophysiol* **101**,
655–664 (2009).

1060

1061 **Supplementary Figures**

1062   Supplementary Figure 1

1063

1064

1065

1066

1067

Supplementary Figure 2

1068

1069

1070

Supplementary Figure 3

1071

1072

1073

Supplementary Figure 4

1074

1075

1076

Supplementary Figure 5

1077

1078

1079

Supplementary Figure 6

1080