Size-Dependent Mechanoadaptation
Enables Migration with Oversized Parasitic Cargo

Authors: Mauricio J.A. Ruiz-Fernandez[1,10], Jianfei Jiang[2,3,10], Alice Battistella[4,5], Armina Mortazavi[2,3,10], Sterre van Wierst[6], Bingzhi Wang[1], Artur Kuznetcov[1], Rüya Aslan[1], Jack Merrin[7], Daan Vorselen[6], Jochen Guck[4,5,8], Markus Meissner[9], Javier Periz[9]*, Benedikt Sabass[2,3,10]*, Jörg Renkawitz[1]*

Affiliations:
[1] Biomedical Center, Walter Brendel Center of Experimental Medicine, Institute of Cardiovascular Physiology and Pathophysiology, Klinikum der Universität, Ludwig-Maximilians-University Munich; Munich, Germany.
[2] Department for Veterinary Sciences, Institute for Infectious Diseases and Zoonoses, Ludwig-Maximilians-University Munich; Munich, Germany
[3] Faculty of Physics and Center for NanoScience, Ludwig-Maximilians-University Munich; Munich, Germany
[4] Max Planck Institute for the Science of Light, Erlangen, Germany
[5] Max-Planck-Zentrum für Physik und Medizin, Erlangen, Germany
[6] Department of Cell Biology and Immunology, Wageningen University & Research, Wageningen, The Netherlands
[7] Institute of Science and Technology (IST) Austria; Klosterneuburg, Austria
[8] Friedrich-Alexander-Universität Erlangen-Nürnberg, Physics Department and Medical Faculty, Erlangen, Germany
[9] Experimental Parasitology, Department for Veterinary Sciences, Ludwig-Maximilians-University Munich; Munich, Germany
[10] Department of Physics, Technische Universität Dortmund, Otto-Hahn-Str. 6, 44221 Dortmund, Germany

*Correspondence: Javier.Periz@para.vetmed.uni-muenchen.de, Benedikt.Sabass@tu-dortmund.de, and Joerg.Renkawitz@med.uni-muenchen.de.

Abstract:

Cell migration is fundamental to development, homeostasis, and immunity, and requires cells to traverse space-limited microenvironments by deforming intracellular organelles such as the nucleus. Although nuclear mechanics are established as a rate-limiting factor during confined migration, how cells transport bulky foreign cargo remains unclear. Here, using *Toxoplasma gondii*, a disease-causing parasite that hijacks motile immune cells for dissemination while replicating into large intracellular assemblies, we discover that parasitic cargoes exceed the size and stiffness of host nuclei, and therefore constitute a primary bottleneck for migration. We uncover a mechanoadaptive response to parasite size, characterized by size-dependent intracellular repositioning, myosin-II contractility, and bleb-based protrusions. This reprogramming of forces and subcellular architecture enables the transport of oversized, mechanically rigid cargo through space-limited microenvironments. These findings establish a mechanobiological principle of how cells size-dependently adapt forces and their subcellular architecture to accommodate intracellular objects that challenge the physical limits of cell migration.

Keywords: cell trafficking, organelle mechanobiology, actomyosin cytoskeleton, cell-matrix interplay, leukocytes, parasitophorous vacuole, Apicomplexa parasites, dendritic cells, macrophages.

Introduction

Cell movement is essential in biology, impacting processes such as organismic development, tissue maintenance, and immune responses. Within multicellular organisms, motile cells face the challenge of navigating complex local microenvironments with spaces substantially smaller than the cells themselves[1]. As a result, motile cells must continuously deform their cell body, as well as their intracellular organelles, such as the nucleus[2,3]. Since the nucleus is typically the stiffest and largest organelle within a cell, its deformation during movement through space-limiting microenvironments is rate-limiting for transmigration[4,5]. To overcome this challenge, motile cells rely on forces generated by their cytoskeleton, which propel themselves and their organelle cargo through these constraining environments[6–8].

Immune cells face the challenge of constantly transmigrating through space-limiting microenvironments, as they move along long migratory tracks during their surveillance and effector responses[9,10]. To alleviate the physical burden of their nuclear cargo along these trafficking routes, immune cells frequently possess nuclei with lower stiffness than non-immune cells[11]. Still, their nucleus is stiffer when compared to other subcellular organelles, and its deformation represents a major bottleneck[5]. To enable their rapid movement, immune cells employ dedicated cytoskeleton networks that accurately position their nucleus intracellularly[12] and squeeze it through narrow pores[6,9]. However, it is not understood how immune cells transport foreign cargoes along their challenging trafficking routes, such as phagocytosed cell corpses, microplastics, and pathogens[13–15], particularly when these cargoes exceed the size of their own organelles.

One example of such a cargo is the human pathogen *Toxoplasma gondii*, an obligate intracellular parasite that infects a large part of the human population and causes toxoplasmosis, which can lead to encephalitis in immunocompromised patients and fetal abnormalities during pregnancy[16]. Upon uptake, the parasite efficiently infects immune cells in the host gut and arrives in lymph nodes within a few days, which represent the homing destination of many immune cells, as well as in distant organs like the heart and brain[17,18]. This fast dissemination of the parasite relies on two complementary behaviors: (1) the extracellular motility of the parasite itself, and (2) the hitchhiking by motile host cells for its transport as an intracellular cargo. The extracellular motility of the parasite is essential for invading into and egressing from host cells, leading to local dissemination[19,20]. In addition, hitchhiking motile host cells is a second, complementary mechanism for long-range parasitic dissemination. Notably, *T. gondii* efficiently infects host immune cells, thereby hijacking their ability to rapidly move between distant tissue locations[15,21]. Moreover, the parasite induces a highly motile mode of immune cells upon infection[22–26], which further facilitates parasite dissemination[27,28]. Thus, the parasite not only hides within immune cells to avoid immune detection, but also turns these cells into its own transport shuttles[15]. However, parasite replication within the parasitophorous vacuole[29] generates large intracellular parasitic objects. This raises the unresolved paradox of how rapidly migrating immune cells efficiently traverse constraining microenvironments while carrying foreign, bulky cargoes.

Here, we discover that the parasitic cargo represents a novel and primary bottleneck for cell migration of infected immune cells, whose transport is even more challenging than the bulky host nucleus. Using custom-made bioengineering, advanced imaging, mechanical force measurements, computer simulations, and genetic engineering of parasites and immune cells, we establish the underlying mechanical and cellular principles that enable efficient immune cell migration despite the physical challenges posed by the foreign cargo. Our findings thereby reveal a new size-dependent mechano-adaptation strategy that redistributes forces and the subcellular architecture.

Results

Efficient Transport of Large Assemblies of Intracellular Parasites Through Dense Environments.

To address the paradox of how motile immune cells transport large intracellular parasitic cargoes through constrained environments, we employed *Toxoplasma gondii* as a well-established eukaryotic parasite model that grows intracellularly[19,22,23,30,31], and mouse dendritic cells (DCs) as a well-established model for motile immune cells that are infected by *Toxoplasma* and move with an amoeboid migration strategy[32–34]. Upon timed infection, we loaded parasite-infected DCs into straight tunnel-like microchannels with a defined width of 8 micrometers and height of 5 micrometers (**Figure 1A**), providing a constrained microenvironment that mimics confining tissues and narrow capillaries[35–38]. Using simultaneous imaging of hundreds of microchannels[39] allowed us to precisely quantify the migratory velocity of parasite-infected DCs within these constraining microenvironments. Importantly, by using genetically engineered *Toxoplasma* parasites that stably encode a Halo-tagged version of the major surface antigen SAG1 for fluorescent visualition[40,41] (**Figure 1A**), we were able to track the velocity of the infected host cell while mapping the parasitic replication stage and thus the intracellular size of the parasitic cargo (**Figures 1B** and **C**).

Dendritic cells infected with small 1-stage parasites rather showed faster migration velocities than uninfected bystander cells (**Figure 1C**), consistent with previous findings that identified a high mobility of *Toxoplasma*-infected DCs[22,27,42]. Cells carrying 2-stage and even 4-stage parasites, of which the latter have comparable sizes to the host nucleus and fill the entire width of the microchannel (**Figures 1A** and **1B**), still showed rapid migratory velocities of around 10 micrometers per minute, comparable to the velocity of non-infected bystander DCs (**Figure 1C** and **Supplemental Movie S1**). When the parasites further replicated to 8-, 16-, and 32-stages, the velocity of the host cell decreased but still reached substantial speeds of 3-8 micrometers per minute (**Figure 1C** and **Supplemental Movie S1**). Yet, notably, at these stages, the parasitic cargo is larger than the host nucleus, the bulkiest organelle in most cell types (**Figure 1B**). Thus, even cells that carried extremely large 16- and 32-stage parasites, which reside within a parasitophorous vacuole (PV) that is (i) up to 5 times longer than the host nucleus, (ii) entirely filled the width of the microchannels, and (iii) occupied large parts of the host cytoplasm (**Figures 1A, 1B** and **Supplemental Movie S1**), still moved in the range of micrometers per minute and thus a magnitude faster than motile mesenchymal cells.

To generalize these findings, we investigated the transport of large parasitic cargoes through constrained microenvironments by macrophages, an immune cell type that, in contrast to DCs, uses a slower adhesive mesenchymal migration strategy[10]. Notably, macrophages are preferentially infected by *Toxoplasma gondii*[43] and are modulated by the parasite into a low-adhesive and highly motile state[25,26]. When we loaded macrophages upon their timed infection with *Toxoplasma gondii* parasites into linear microchannels (**Figure S1A**), quantification of their velocity revealed an extremely rapid motility of around 16 to 18 micrometers per minute of cells infected with 1-, 2-, and 4-stage parasites (**Figures 1D** and **S1A,** and **Supplemental Movie S2**). Comparable to infected dendritic cells, macrophages loaded with large 8-, 16- and 32-stage parasites carried parasitic cargoes that were up to 3 times larger than the macrophage's nucleus (**Figures S1A and S1B**), slowed down but migrated still with substantial speeds of 3 to 10 micrometers per minute (**Figure 1D** and **S1A,** and **Supplemental Movie S2**). Importantly, these velocities represent rapid speeds for macrophages since they typically only migrate with speeds of around 1 micrometer per minute in their uninfected state[44]. Overall, these findings demonstrate that the transport of extremely large parasitic cargoes by immune cells through constrained microenvironments is remarkably efficient.

To identify the limits of parasitic transport by motile immune cells, we next introduced a
defined pore of 4 micrometers in width into the migratory track (**Figure S1C**). Although larger
parasitic cargoes took longer to squeeze through the pore within dendritic cells, they still passed
surprisingly quickly (**Figures 1F** and **S1C**). To provide a smaller pore that is even smaller than the
diameter of an individual parasite within the parasitophorous vacuole, we challenged parasite-infected
DCs with pores of only 2 micrometers in width (**Figure 1E**). To our surprise, we still observed the
rapid translocation of the host cell and its parasitic cargo through these narrow microenvironmental
barriers (**Figures 1E** and **1G,** and **Supplemental Movie S3**). Dendritic cells infected with 1-, 2-, and
4-stage parasites took the same time to pass through the 2 micrometers pore as uninfected cells, and
even large parasitic 16-stage cargoes only slowed down two-fold (**Figure 1G**). Similarly, parasite-
infected macrophages crossed these narrow microenvironmental barriers with only a 2- to 3-fold delay
at 4 to 8 and 16 to 32 parasite stages, respectively, compared to 1- to 2-staged parasites (**Figure S1D**
and **Supplemental Movie S4**). Thus, *Toxoplasma* parasites efficiently overcome microenvironmental
constraints and cross local barriers within infected dendritic cells and macrophages, even when the
parasitic cargo reaches sizes substantially larger than any host cell's organelle.
154
Intracellular Parasites Cause Migration Bottlenecks for Infected Immune Cell Hosts.
One solution to the paradox of how large parasitic cargoes might be transported efficiently through
space-constrained microenvironments is that the parasite may be mechanically soft, enabling large
deformations. To test this hypothesis, we performed atomic force microscopy (AFM) on *Toxoplasma*-
carrying DCs after fixation in removable microenvironments that confine cells between two
surfaces[45] (**Figure 2A**). This approach allowed us (i) to measure the physical properties of the
parasites and the host cells from the outside when they are flatly spread, and (ii) to uncouple the
measurement of stiffness from the motility of the host cell. Quantification of the Young's modulus, a
mechanical property that increases with increasing stiffness, showed a higher Young's modulus of
each individual parasite in comparison to the local intra-vacuolar matrix between the individual
parasites within a parasitophorous vacuole (PV) (**Figures 2A** and **2B**). Surprisingly, the stiffness of
the parasites was even slightly higher than that of the host nucleus (**Figure 2A** and **2B**), which
typically represents the major deformation bottleneck for motile cells crossing constrained
microenvironments[2,4,5]. When we analyzed the dissipated viscous energy measured by AFM as a
mechanical parameter for viscosity[46,47] , we detected a slightly lower dissipated viscous energy of the
parasites compared to the host nucleus (**Figure 2A** and **2C**), indicating a more pronounced elastic
response of the parasite. These data suggest that *Toxoplasma* -infected immune cells harbour a novel,
stiff yet elastic, intracellular obstacle in addition to the bulky host nucleus.
To corroborate this finding, we measured the internal mechanical properties of the parasites
within infected immune cells using Brillouin microscopy[48] . Combining this approach with confocal
imaging of the host cell nucleus and SAG1-Halo-labeled *Toxoplasma* parasites (**Figure 2D**) revealed
a higher Brillouin shift (**Figure 2D** and **2E**) and a higher Brillouin peak linewidth (**Figure 2D** and **2F**)
of the entire parasitic cargo in comparison to the host cell's nucleus. The increased Brillouin peak
linewidth, a proxy for the viscosity of the sample[48] , suggests that rather the low viscosity of the
individual parasites measured with AFM is relevant during squeezing than the internal property of the
entire cargo measured by Brillouin microscopy. The increased Brillouin shift, a proxy for the
longitudinal modules and thus stiffness, further validates that the entire parasitic cargo is stiffer than
the host nucleus. Together, these data support the finding that *Toxoplasma* parasites within motile
immune cells have physical properties that make their deformation challenging.
Indeed, when we measured the shape of the parasite and the host nucleus when they were in
direct neighborhood, we observed strong deformations of the host nucleus by individual parasites
(**Figures 2G** and **S2A**), confirming a lower deformability of the parasite in comparison to the stiffest

host organelle. We further validated this finding with live-cell imaging, which revealed dynamic and
frequent nuclear deformation by the parasitic cargo when they were in close proximity (**Figure S2B**).
Thus, intracellular *Toxoplasma* parasites are not only a large but also a physically challenging cargo.
To directly test whether this physically challenging cargo represents a new primary bottleneck
for infected cells to move through constrained microenvironments, we next performed fast time-lapse
and high-resolution imaging of labeled parasites translocating within either dendritic cells or
macrophages through 2 micrometer pores (**Figures 2H** and **2I,** and **Supplemental Movie S3**). This
revealed that large parasitic cargoes are compressed once they encounter the pore (**Figure 2J** and
Supplemental Movie S3), followed by the sequential passage of individual parasites through the pore
(**Figures 2H, 2I, S2D,** and **S2E,** and **Supplemental Movies S3 and S4**). During this sequential
passage of individual parasites, their ellipsoid shape was substantially deformed (**Figures 2H, 2I,**
S2D, and **S2E**). Behind the pore, the parasites rapidly returned to their original shape and reorganized
within their parasitophorous vacuole (**Figures 2H-J**). Notably, not only were large parasitic cargoes
compressed in front of the pore (**Figure 2J**) but also the individual parasites frequently resided for
some time at the entrance of the pore (**Figure S2C-E**), which was followed by a quick translocation
through the pore once the individual parasite was deformed (**Figures 2H** and **2J**), both underscoring
the physical challenge of the parasitic cargo for passage through constrained microenvironments.
Indeed, when we measured the time for the parasitic cargo to pass through the pore, compared with
the passage times of the host nucleus and the entire host cell, we observed that the major translocation
time is determined by large parasitic cargoes (**Figures S2F-I**), identifying that parasite-infected
immune cells carry a novel intracellular bottleneck that is even more challenging than the bulky host
nucleus. In line with these findings, we often observed an increased distance between the host nucleus
and the parasitic cargo after the host nucleus passed through a constriction, while the parasitic cargo
remained at the pore entrance (**Figure 2H and S2D**). Together, these results demonstrate that
intracellular *Toxoplasma* parasites within motile immune cells are large, bulky, and rigid cargoes that
locally unfold and re-fold to allow single parasites to translocate sequentially. Overall, these findings
identify the parasitic cargo as the primary bottleneck during the migration of infected immune cells
through constrained microenvironments.
215

Intracellular Positioning Adapts to the Size of the Parasitic Cargo.
To identify the mechanism by which infected immune cells migrate in constrained environments
while carrying the physical burden of their parasitic cargo, we first quantified the intracellular position
of the parasite relative to the host nucleus, as the position of organelles serves as an indicator of
underlying cell polarity and thus cytoskeletal forces[49,50] . To our surprise, we observed that parasitic
cargoes are non-randomly positioned within the cytoplasm of motile DCs, depending on the parasitic
cargo stage (**Figures 3A** and **3B,** and **Supplemental Movie S1**). While small 1-stage parasitic cargoes
are mostly positioned in the back of the host nucleus, growing parasitic cargoes are positioned with
increasing frequency towards the cellular leading edge in front of the host nucleus, until very large 32-
stage PV cargoes are predominantly positioned towards the host cell front (**Figure 3B**). To confirm
this re-localization of the parasitic cargo with increasing size, we analyzed parasite-infected DCs in
microchannels of larger width (**Figure S3A** and **S3B**) as well as during their navigation through
maze-like microenvironments (**Figure S3C**). In all microenvironments, large parasitic cargoes were
particularly frequently positioned in front of the host nucleus (**Figures 3B, S3A-C**). This even more
forward positioning of large parasitic cargoes compared to the host nucleus is particularly unexpected,
since DCs belong to the class of migrating cells that employ an amoeboid migration mode that is
characterized by low-adhesiveness to the substrate and forward positioning of the nucleus towards the
cellular protrusion[49,51]. Overall, these data reveal that the *Toxoplasma* parasitic cargo is positioned in a
manner that adapts to its size.

In agreement with size-dependent cargo positioning, we frequently observe the repositioning
of large parasitic cargoes from the rear to the front during our imaging intervals but rarely vice versa
(**Figures 3C, 3D**, **S3D** and **S3E,** and **Supplemental Movie S5**). Since these repositioning events of
the large parasitic cargo require overtaking of the nucleus and thus substantial deformation of both
cargoes (**Figure 3C, S3D** and **Supplemental Movie S5**), these results demonstrate that parasitic
cargoes actively adapt their intracellular localization.

To confirm that repositioning is an active process that is specific to the parasitic cargo, we
loaded parasite-infected DCs with rigid polystyrene beads. With a diameter of 6 micrometers, these
beads mimic the size of a 4-stage parasitic cargo. However, we observed the beads mostly positioned
behind the parasitic cargo (**Figures S4A** and **S4B**), suggesting that a passive bead cargo differs from a
parasitic cargo. To further corroborate this finding, we loaded deformable beads (acrylamide co-
acrylic acid microparticles[52,53]) with a size of 6 micrometers and a stiffness of 6.5 kPa into motile
DCs. This approach has the advantages of mimicking (i) the size and (ii) the stiffness of individual
parasites, also when compared to 1-5 kPa stiffness range of immune cell nuclei[54] . Indeed, as for
parasitic cargoes, these bead cargoes dynamically deform the host cell nucleus (**Figure S4C** and
Supplemental Movie S6). Moreover, DCs frequently took up multiple beads that clustered
intracellularly, allowing quantification of cells carrying a range of bead assemblies and thus cargo
sizes (**Figure 3E**), thereby closely mimicking the intracellular sizes of parasitic cargoes. However, in
contrast to parasitic cargoes, these bead cargoes often localized to the rear half of the host cell body
and randomly in relation to the host nucleus (**Figure 3F** and **3G**). Moreover, bead cargoes did not
show a correlation between their positioning and cargo size (**Figure 3F**). Additionally, in contrast to
parasitic cargoes, the intracellular position of the bead cargo remained stable along long migration
tracks due to rare reposition events of the bead cargoes to the front of the host nucleus (**Figures 3G**
and **S4D,** and **Supplemental Movie S6**). Together, these findings reveal that the cellular organization
and behavior differ depending on whether the cells carry large bead- or parasitic-cargoes.

To characterize the adaptation of cellular organization to the size of the parasitic cargo, we
next mapped the position of the parasite in relation to the microtubule-organizing center (MTOC), a
key marker of cellular organization[55] . To do so, we infected EB3-mcherry expressing DCs, which
serves as a microtubule-plus end and microtubule-organizing center (MTOC) marker within the host
cell[12,56,57] . This revealed that large parasitic cargoes are exclusively positioned forward of the MTOC
(**Figures 3H** and **S3E,** and **Supplemental Movie S7**), which typically localizes approximately to the
cell center[49,55] . Moreover, in most cases, the large parasitic cargo is even positioned together with the
nucleus in front of the MTOC (**Figure S3F,** and **Supplemental Movie S7**). These data show that both
the parasitic cargo and the nucleus can simultaneously acquire an amoeboid-like positioning in front
of the MTOC, and that the parasitic cargo preferentially locates forward of these key host organelles,
particularly when it is large.

To test whether large parasitic *Toxoplasma* cargoes generally localize towards the cell front in
infected immune cells, we infected macrophages and again observed parasitic cargoes actively
positioned in front of the host nucleus (**Figures 3I**) in a size-dependent manner (**Figures 3J**),
comparable to parasitic cargoes in motile DCs. Together, these results reveal that *Toxoplasma*
parasitic cargoes actively position towards the front of their immune host cells despite their large,
bulky nature. Overall, these findings reveal a mechanical adaptation of forces and intracellular
positioning in response to the size of the parasitic cargo.

Myosin-driven Host Cell Contractility Adapts to the Parasite Size.
To identify the underlying mechanism of size-adaptive parasite positioning within infected immune
cells, we developed a particle-based model to simulate the arrangement of the parasitic cargo relative
to the host nucleus during immune cell migration. By combining dissipative particle dynamics[58] , with

a representation of the host nucleus, the host cortex, and the parasitic cargo as elastic shells, this
simulation captures the effect of (i) mechanical coupling between the host nucleus, the host cortex and
the extracellular microenvironment, (ii) cytoplasmic flows, (iii) cell contractility, and (iv) parasite
cargo size onto the localization of intracellular cargoes in a spatiotemporal manner (see
Supplementary Information for details on the simulation). The position of the parasitic cargo is
determined by a balance of pressure at the rear, representing actomyosin activity, and visco-elastic
forces that result from a coupling between the cortex and the cargo. The simulation predicts that the
pushing and pulling forces on the cargo depend on its size in constricted environments. Moreover,
they revealed that increasing the host contractility, simulated by the host cortical tension, results in a
higher likelihood of forward localization of the parasitic cargo, in particular when the parasitic cargo
reaches sizes larger than the host nucleus (**Figures 4A** and **4B,** and **Supplemental Movie S8**).

To directly test the prediction of the model experimentally, we imaged the localization of host
myosin-IIA, since myosin-IIA is a key driver of cellular contractility in immune cells[36] , and since the
myosin-II upstream regulators Rho/ROCK are critical for the tissue dissemination of the parasite by
monocytes and macrophages[30] . Live cell imaging of non-muscle myosin heavy chain IIA (myosin-
IIA, encoded by the MYH9 gene) using DCs encoding MYH9-GFP and infected with *Toxoplasma*
parasites revealed a strong enrichment of myosin-IIA in the rear of the host cell behind the parasitic
cargo (**Figure 4C** and **Supplemental Movie S9**). While uninfected immature DCs also often exhibit a
rearward myosin accumulation[59] , the intracellular distribution pattern was more evenly distributed in
uninfected bystander DCs when compared to parasite-infected DCs (**Figure 4C** and **D**). Notably, the
more the parasite size increased, the more prominently myosin-IIA was enriched to the rear (**Figure
4D**). This finding demonstrates that large parasitic cargoes cause a shift in the distribution of host
myosin-IIA from the front to the rear, which is indicative of rapid immune cell motility[59] . In contrast,
DCs loaded with bead cargoes of different sizes did not show a correlation between cargo size and
myosin-IIA enrichment at the host cell rear (**Figures 4C** and **4E**). Thus, the distribution of host
myosin-IIA adapts to the size of the parasitic cargo.

To test whether redistribution of host myosin-IIA alters cellular contractility, we first
examined the morphology of DCs infected with parasites. While motile DCs are morphologically
characterized by veil-like protrusions and many macropinosome-like vesicles in the front part of the
cell body[60,61] , *Toxoplasma gondii*-infected DCs frequently showed roundish, smooth, and vesicle-free
cell fronts that can even have bleb-like morphologies (**Figure 4F** and **Supplemental Movie S10**),
both indicators of a high degree of intracellular contractility. Notably, when we quantified the
frequency of this smooth cell front phenotype in relation to the size of the parasitic cargo, we
observed that large parasitic cargoes almost always caused the appearance of a non-veiled, smooth, or
blebbing cell front (**Figure 4G**). A key defining feature of a bleb during bleb-based migration
strategies is a high contractility, leading to the expansion of the bleb faster than the polymerization of
F-actin[62,63]. By infecting DCs that express Lifeact-GFP as a marker for F-actin, we indeed observed
rapidly expanding blebs that were devoid of F-actin (**Figure 4H** and **S5A,** and **Supplemental Movie
S10**). Thus, large parasitic cargoes induce high contractility within their motile host immune cells. To
further corroborate this finding, we depolymerized the host microtubule cytoskeleton with low doses
of nocodazole, well-known to increase cellular contractility by the release of microtubule-bound
myosin activators like RhoGEFH1/Lfc1[57,64] , and observed that the blebbing phenotype already
appeared even with smaller parasitic cargoes (**Figure S5B** and **S5C**). Together, these data identify an
enhanced myosin-based contractility in cells carrying large parasitic cargoes.

To directly test whether this enhanced myosin-based contractility causes the forward
positioning of large parasitic cargoes, as predicted by our simulations, we used the myosin inhibitor
para-nitroblebbistatin and measured the localization of the parasitic cargo during DC migration. While
control cells positioned their large parasite cargoes forward, myosin-inhibited cells displayed a

localization of the parasitic cargo independent of its size (**Figure 4I**). This finding was substantiated
when we quantified the frequency of repositioning of the parasitic cargo and the host nucleus along
long migration tracks, showing that myosin inhibition strongly impaired the ability of parasitic
cargoes to move towards the host cell front (**Figure 4J**). Thus, myosin-based contractility within the
host cell drives the intracellular positioning of the parasite as an adaptation to its size. Overall, these
findings reveal a shift in the force balance of motile immune cells hijacked by *Toxoplasma* parasite,
depending on the pathogen's size.
338
Adaptive Myosin-driven Contractility Facilitates Parasitic Transport Across Local Barriers.
These findings suggest a model in which myosin-based contractility at the host cell's rear generates a
contractile force that propels the parasitic cargo through space-limiting environments in a size-
dependent manner. To experimentally test this model, we infected DCs and measured the force they
generate in their local microenvironment using quantitative traction force microscopy (TFM). Since
motile immune cells, and particularly *Toxoplasma*-infected immune cells, use an amoeboid migration
strategy that is non-adhesive and therefore prevents their analysis on flat two-dimensional
substrates[10,21], we measured the traction forces generated by cells confined between two substrates,
one of which had embedded fluorescent beads (**Figure 5A** and **Supplemental Movie S11**). Mapping
and quantification of the displacement of the beads during the motility of infected DCs (**Figure S5D**)
showed higher bead displacements (**Figure S5E**) and, therefore, higher traction magnitudes (**Figure
5B**) of cells carrying large parasitic cargoes. These results show that immune cells can adjust their
force-regime to the size of the parasitic cargo.
Next, to test whether this adaptive force generation is myosin-dependent, we infected DCs
with a genetic knockout of myosin-IIA. This revealed that the overall average traction magnitude was
lower in *Toxoplasma*-infected myosin-IIA DCs than in *Toxoplasma*-infected wild-type DCs, and
increased less with the size of the parasitic cargo (**Figure 5B** and **S5E**). Together, these data
demonstrate that immune cells adapt their force-generation to the size of the parasitic cargo in a
myosin-IIA-dependent manner.
To test whether these myosin-IIA-dependent forces are required for parasitic cargo transport,
we next quantified the velocity of parasite-loaded motile DCs upon myosin inhibition in linear 8-
micrometer-wide microchannels. This demonstrated that myosin inhibition by para-nitroblebbistatin
impaired the migration of DCs carrying larger parasitic cargoes (**Figure 5C**). In contrast, uninfected
bystander cells and parasite-infected cells with small 1-stage PVs reached the same velocities upon
myosin inhibition as control cells (**Figure 5C**). These data show that the migration of hijacked
immune cells through confined microenvironments is facilitated by adaptive myosin contractility to
the size of the parasite cargo.
To investigate the consequences for migration across local barriers, we next investigated how
frequently parasite-infected DCs passed through 2 micrometers pores following myosin inhibition.
While almost all control cells efficiently translocated through these small pores, myosin inhibition
strongly impaired the translocation of larger PVs (**Figures 5D** and **5E,** and **Supplemental Movie
S12**). While small 1- and 2-stage parasites successfully migrated through the pores in approximately
60% of cases, large 4-, 8-, and 16-stage parasites almost always got stuck at the 2 micrometer-sized
pores (**Figures 5D** and **5E**). This behavior was reminiscent of cells loaded with assemblies of
deformable beads, which rarely achieved the successful movement of the cargo across 2 micrometers
pores (**Figures 5F-H,** and **Supplemental Movie S12**). Notably, when we imaged the behavior of
parasite-loaded DCs that become stuck at the pore entrance, we observed parasite egress from the host
(**Figure 5I** and **Supplemental Movie S12**). Quantification relative to the cargo size revealed that
extracellular egress was rare in control cells but occurred frequently with large parasitic cargoes upon
the impairment of myosin activity (**Figure 5J**). Together, these findings reveal that contractility

adaptive to the size of the parasitic cargo is critical for the successful transport of *Toxoplasma*
parasites within host cells through space-limited environments and across local barriers.
Overall, our results establish that the parasite *Toxoplasma gondii* alters the cellular force
balance of its motile host cells by hijacking and adapting their myosin-based contractility. Thereby,
we reveal a novel hijacking strategy in which the parasite modifies its host's force regime and
subcellular architecture to accommodate the size of its parasitic cargo. This mechano-adaptation to
size facilitates the transport of large intracellular parasite assemblies across locally complex and
confined microenvironments, and resolves the paradox of how parasite-infected immune cells can
effectively migrate through such environments despite their bulky parasitic cargo.

Discussion

Cell migration through confining tissue environments requires continuous coordination between cytoskeletal force generation, subcellular architecture, and the mechanical properties of intracellular organelles[1,2,9,36,49]. While the nucleus has emerged as a dominant mechanical bottleneck in confined migration[2,4,5], cells like phagocytes can carry additional intracellular cargoes. However, whether such cargoes and their physical properties challenge the migratory capacity, and how cells respond to this challenge, has remained unknown. Here, we identify large intracellular parasites as previously unrecognised mechanical constraints for immune cell migration and uncover a mechanism by which host cells adaptively rewire cytoskeletal forces and intracellular organization to enable the transport of oversized, rigid cargoes through confining microenvironments.

Our findings reveal a novel principle of subcellular architecture dynamics and mechanoadaptation during cell migration: motile immune cells dynamically reorganize the spatial organization between force-generating cytoskeletal networks, the nuclear organelle bottleneck, and a mechanically demanding intracellular cargo. In uninfected cells, the nucleus typically represents the stiffest and largest intracellular object, having a defined intracellular position[50] and acting as a dominant intracellular obstacle for migration through space-limiting environments[2,4,5,34,65,66]. In contrast, we show that intracellular *Toxoplasma gondii* parasites can exceed the nucleus in both stiffness and size, effectively overriding it as the primary mechanical obstacle. Remarkably, instead of preventing migration, infected cells reposition the parasitic cargo to the front of the host cell body as the size of the cargo burden increases, mediated by increasingly rearward-localized contractile forces. This adaptive positioning highlights the plasticity of subcellular architecture during migration. Notably, inert cargoes of comparable size and physical properties do not elicit this adaptive contractile response, in contrast to the previously discovered recruitment of the actin cytoskeleton to beads within confined migrating DCs[6], suggesting that immune cells engage a cargo-selective or cargo-induced contractile program. More broadly, these findings establish that foreign cargoes influence immune cell trafficking, either by impeding passage through micron-scale pores, as observed for inert particles, or by inducing force and architecture rewiring, as observed for active parasitic cargoes.

From a mechanical perspective, our data demonstrate that immune cells adapt cytoskeletal force generation to the size of the parasitic cargo. We observe a progressive rearward localization of myosin-II and an increase in contractile, bleb-inducing forces as the parasite size increases, suggesting a scaling relationship between cargo dimensions and force output. The spatial distribution of myosin-II, the independence from microtubule-based pulling forces, the bleb-based migration mode, and the loss of efficient parasite translocation upon myosin inhibition together support a model in which rearward contractility predominantly generates pushing forces that drive the parasitic cargo through confinement. Notably, these mechanistic and mechanical findings are fully consistent with previous work showing that Rho/ROCK signaling, which is a key upstream regulator of myosin-II activity, is critical for *in vivo* tissue dissemination of *Toxoplasma* by monocytes and macrophages[30].

In addition to force scaling, we uncover a mechanical shape-adaptation mechanism in which the parasitic cargo transiently deforms and unfolds into smaller substructures during the passage through narrow constrictions. This dynamic reduction in the effective cross-sectional area indicates that confined migration with oversized cargo is not determined solely by static size or stiffness, but also by the capacity for reversible cargo remodeling under force. This behavior mirrors principles observed at the cellular and nuclear levels, where cells exploit shape plasticity to traverse restrictive microenvironments[1], and where neutrophil granulocytes use lobulated nuclear shapes to efficiently translocate through micron-scale constrictions[11,67]. Our findings extend these principles to intracellular cargo, highlighting that force-induced cargo remodeling represents an important layer of mechanical adaptation during migration.

Beyond cell biology and mechanics, these findings have important implications for parasite dissemination within host organisms. Intracellular pathogens are known to hijack host cytoskeletal forces to facilitate uptake, intracellular motility, stability, and evasion of degradation. For instance, bacterial pathogens such as *Shigella flexneri* exploit actin-driven macropinocytosis for entry[68], *Chlamydia* relies on host actin scaffolds for vacuole stability[69], and *Listeria* and *Rickettsia* hijack actin polymerisation for comet-like propulsion[70,71]. In contrast, how large eukaryotic intracellular parasites exploit host cytoskeletal forces has remained poorly understood. Our findings demonstrate that *Toxoplasma* parasites, unlike non-living bead cargoes of comparable sizes and stiffness, elicit a size-dependent reorganization of host contractility, indicating that the parasite is not a passive mechanical burden but an active modulator of host cell mechanics. This size-dependent force-rewiring of motile immune cells complements the known ability of *Toxoplasma* parasites to induce a rapid migratory, low-adhesive state in their immune cell hosts, together likely providing an evolutionary advantage by enabling a robust dissemination strategy across diverse tissues and physical environments.

In summary, our study uncovers a previously unrecognized intersection between subcellular architecture, adaptive force generation, and host-pathogen interactions. By revealing how migrating immune cells reorganize intracellular mechanics to accommodate oversized parasitic cargoes, we establish new principles of mechanoadaptation with implications for cell migration, organelle positioning, immune responses, and pathogen dissemination across tissues.

References

1. Yamada, K.M., and Sixt, M. (2019). Mechanisms of 3D cell migration. Nat. Rev. Mol. Cell Biol. *20*, 738–752. https://doi.org/10.1038/s41580-019-0172-9.

2. Calero-Cuenca, F.J., Janota, C.S., and Gomes, E.R. (2018). Dealing with the nucleus during cell migration. Curr. Opin. Cell Biol. *50*, 35–41. https://doi.org/10.1016/j.ceb.2018.01.014.

3. Nader, G.P. de F., Williart, A., and Piel, M. (2021). Nuclear deformations, from signaling to perturbation and damage. Curr Opin Cell Biol *72*, 137–145. https://doi.org/10.1016/j.ceb.2021.07.008.

4. Kalukula, Y., Stephens, A.D., Lammerding, J., and Gabriele, S. (2022). Mechanics and functional consequences of nuclear deformations. Nat Rev Mol Cell Bio, 1–20. https://doi.org/10.1038/s41580-022-00480-z.

5. Wolf, K., Lindert, M. te, Krause, M., Alexander, S., Riet, J. te, Willis, A.L., Hoffman, R.M., Figdor, C.G., Weiss, S.J., and Friedl, P. (2013). Physical limits of cell migration: Control by ECM space and nuclear deformation and tuning by proteolysis and traction force. J Cell Biol *201*, 1069–1084. https://doi.org/10.1083/jcb.201210152.

6. Thiam, H.-R., Vargas, P., Carpi, N., Crespo, C.L., Raab, M., Terriac, E., King, M.C., Jacobelli, J., Alberts, A.S., Stradal, T., et al. (2016). Perinuclear Arp2/3-driven actin polymerization enables nuclear deformation to facilitate cell migration through complex environments. Nat. Commun. *7*, 10997. https://doi.org/10.1038/ncomms10997.

7. Blanchoin, L., Boujemaa-Paterski, R., Sykes, C., and Plastino, J. (2014). Actin Dynamics, Architecture, and Mechanics in Cell Motility. Physiol. Rev. *94*, 235–263. https://doi.org/10.1152/physrev.00018.2013.

8. Shah, E.A., and Keren, K. (2013). Mechanical forces and feedbacks in cell motility. Current Opinion in Cell Biology. https://doi.org/10.1016/j.ceb.2013.06.009.

9. Calmettes, V., Quintanilla, M.A., Mariano, L.L., Piel, M., Moreau, H.D., and Lennon□Duménil, A. (2026). Mechanosensing in Dendritic Cells. Immunol. Rev. *337*. https://doi.org/10.1111/imr.70086.

10. Kameritsch, P., and Renkawitz, J. (2020). Principles of Leukocyte Migration Strategies. Trends Cell Biol. *30*, 818–832. https://doi.org/10.1016/j.tcb.2020.06.007.

11. Renkawitz, J., Yesin, A., Kroll, J., Cabral, A.T., D'Annunzio, S., and Thiam, H.R. (2026). Nuclear mechanobiology rules immune cells' functions: from differentiation to cell trafficking and pathogen killing. Nucleus *17*, 2590843. https://doi.org/10.1080/19491034.2025.2590843.

12. Kroll, J., Hauschild, R., Kuznetcov, A., Stefanowski, K., Hermann, M.D., Merrin, J., Shafeek, L., Müller□Taubenberger, A., and Renkawitz, J. (2023). Adaptive pathfinding by nucleokinesis during amoeboid migration. EMBO J., e114557. https://doi.org/10.15252/embj.2023114557.

13. Weavers, H., Evans, I.R., Martin, P., and Wood, W. (2016). Corpse Engulfment Generates a Molecular Memory that Primes the Macrophage Inflammatory Response. Cell *165*, 1658–1671. https://doi.org/10.1016/j.cell.2016.04.049.

14. Vanetti, C., Broggiato, M., Pezzana, S., Clerici, M., and Fenizia, C. (2025). Effects of
microplastics on the immune system: How much should we worry? Immunol. Lett. *272*, 106976.
https://doi.org/10.1016/j.imlet.2025.106976.

15. Drewry, L.L., and Sibley, L.D. (2019). The hitchhiker's guide to parasite dissemination. Cell
Microbiol *21*, e13070. https://doi.org/10.1111/cmi.13070.

16. Montoya, J., and Liesenfeld, O. (2004). Toxoplasmosis. Lancet *363*, 1965–1976.
https://doi.org/10.1016/s0140-6736(04)16412-x.

17. Barragan, A., and Sibley, L.D. (2003). Migration of Toxoplasma gondii across biological barriers.
Trends Microbiol *11*, 426–430. https://doi.org/10.1016/s0966-842x(03)00205-1.

18. Matta, S.K., Rinkenberger, N., Dunay, I.R., and Sibley, L.D. (2021). Toxoplasma gondii infection
and its implications within the central nervous system. Nat. Rev. Microbiol. *19*, 467–480.
https://doi.org/10.1038/s41579-021-00518-7.

19. Frénal, K., Dubremetz, J.-F., Lebrun, M., and Soldati-Favre, D. (2017). Gliding motility powers
invasion and egress in Apicomplexa. Nat Rev Microbiol *15*, 645–660.
https://doi.org/10.1038/nrmicro.2017.86.

20. Tardieux, I., and Baum, J. (2016). Reassessing the mechanics of parasite motility and host-cell
invasion. J Cell Biol *214*, 507–515. https://doi.org/10.1083/jcb.201605100.

21. Ólafsson, E.B., and Barragan, A. (2020). The unicellular eukaryotic parasite Toxoplasma gondii
hijacks the migration machinery of mononuclear phagocytes to promote its dissemination. Biol Cell
112, 239–250. https://doi.org/10.1111/boc.202000005.

22. Lambert, H., Hitziger, N., Dellacasa, I., Svensson, M., and Barragan, A. (2006). Induction of
dendritic cell migration upon Toxoplasma gondii infection potentiates parasite dissemination. Cellular
microbiology *8*, 1611–1623. https://doi.org/10.1111/j.1462-5822.2006.00735.x.

23. Weidner, J.M., Kanatani, S., Castañeda, M.A.H., Fuks, J.M., Rethi, B., Wallin, R.P.A., and
Barragan, A. (2013). Rapid cytoskeleton remodelling in dendritic cells following invasion by
Toxoplasma gondii coincides with the onset of a hypermigratory phenotype. Cellular microbiology
15, 1735–1752. https://doi.org/10.1111/cmi.12145.

24. Cook, J.H., Ueno, N., and Lodoen, M.B. (2018). Toxoplasma gondii disrupts β1 integrin signaling
and focal adhesion formation during monocyte hypermotility. J Biol Chem *293*, 3374–3385.
https://doi.org/10.1074/jbc.m117.793281.

25. Hoeve, A.L. ten, Rodriguez, M.E., Säflund, M., Michel, V., Magimel, L., Ripoll, A., Yu, T.,
Hakimi, M.-A., Saeij, J.P.J., Ozata, D.M., et al. (2024). Hypermigration of macrophages through the
concerted action of GRA effectors on NF-κB/p38 signaling and host chromatin accessibility
potentiates Toxoplasma dissemination. mBio *15*, e02140-24. https://doi.org/10.1128/mbio.02140-24.

26. Hoeve, A.L. ten, Braun, L., Rodriguez, M.E., Olivera, G.C., Bougdour, A., Belmudes, L., Couté,
Y., Saeij, J.P.J., Hakimi, M.-A., and Barragan, A. (2022). The Toxoplasma effector GRA28 promotes
parasite dissemination by inducing dendritic cell-like migratory properties in infected macrophages.
Cell Host Microbe. https://doi.org/10.1016/j.chom.2022.10.001.

27. Lambert, H., Vutova, P.P., Adams, W.C., Loré, K., and Barragan, A. (2009). The Toxoplasma
gondii-Shuttling Function of Dendritic Cells Is Linked to the Parasite Genotype. Infection and
Immunity *77*, 1679–1688. https://doi.org/10.1128/iai.01289-08.

28. Sangaré, L.O., Ólafsson, E.B., Wang, Y., Yang, N., Julien, L., Camejo, A., Pesavento, P., Sidik,
S.M., Lourido, S., Barragan, A., et al. (2019). In Vivo CRISPR Screen Identifies TgWIP as a
Toxoplasma Modulator of Dendritic Cell Migration. Cell Host Microbe *26*, 478-492.e8.
https://doi.org/10.1016/j.chom.2019.09.008.

29. Clough, B., and Frickel, E.-M. (2017). The Toxoplasma Parasitophorous Vacuole: An Evolving
Host–Parasite Frontier. Trends Parasitol *33*, 473–488. https://doi.org/10.1016/j.pt.2017.02.007.

30. Drewry, L.L., Jones, N.G., Wang, Q., Onken, M.D., Miller, M.J., and Sibley, L.D. (2019). The
secreted kinase ROP17 promotes Toxoplasma gondii dissemination by hijacking monocyte tissue
migration. Nat Microbiol *4*, 1951–1963. https://doi.org/10.1038/s41564-019-0504-8.

31. Li, W., Grech, J., Stortz, J.F., Gow, M., Periz, J., Meissner, M., and Jimenez-Ruiz, E. (2022). A
splitCas9 phenotypic screen in Toxoplasma gondii identifies proteins involved in host cell egress and
invasion. Nat. Microbiol. *7*, 882–895. https://doi.org/10.1038/s41564-022-01114-y.

32. Alraies, Z., Rivera, C.A., Delgado, M.-G., Sanséau, D., Maurin, M., Amadio, R., Piperno, G.M.,
Dunsmore, G., Yatim, A., Mariano, L.L., et al. (2024). Cell shape sensing licenses dendritic cells for
homeostatic migration to lymph nodes. Nat. Immunol., 1–14. https://doi.org/10.1038/s41590-024-
01856-3.

33. Alanko, J., Uçar, M.C., Canigova, N., Stopp, J., Schwarz, J., Merrin, J., Hannezo, E., and Sixt, M.
(2023). CCR7 acts as both a sensor and a sink for CCL19 to coordinate collective leukocyte
migration. Sci. Immunol. *8*, eadc9584. https://doi.org/10.1126/sciimmunol.adc9584.

34. Raab, M., Gentili, M., Belly, H. de, Thiam, H.-R., Vargas, P., Jimenez, A.J., Lautenschlaeger, F.,
Voituriez, R., Lennon-Duménil, A.-M., Manel, N., et al. (2016). ESCRT III repairs nuclear envelope
ruptures during cell migration to limit DNA damage and cell death. Science *352*, 359–362.
https://doi.org/10.1126/science.aad7611.

35. Deygas, M., Bernard, M., Nivoit, P., Barbier, L., Wang, L., Maurin, M., Tejerina, E., Jakuszeit, T.,
Deslys, A., Garcia-Gomez, M., et al. (2025). Adaptive motility enables neutrophils to rapidly navigate
confined capillaries. bioRxiv, 2025.01.08.631737. https://doi.org/10.1101/2025.01.08.631737.

36. Moreau, H.D., Piel, M., Voituriez, R., and Lennon-Duménil, A.-M. (2018). Integrating Physical
and Molecular Insights on Immune Cell Migration. Trends Immunol. *39*, 632–643.
https://doi.org/10.1016/j.it.2018.04.007.

37. Garcia-Arcos, J.M., Chabrier, R., Deygas, M., Nader, G., Barbier, L., Sáez, P.J., Mathur, A.,
Vargas, P., and Piel, M. (2019). Reconstitution of cell migration at a glance. Journal of Cell Science
132, jcs225565. https://doi.org/10.1242/jcs.225565.

38. Rodriguez, M.E., Hassan, A., Linaroudis, N., Harryson-Oliveberg, F., Hoeve, A.L. ten, and
Barragan, A. (2025). ICAM-1/CD18-mediated sequestration of parasitized phagocytes in cortical
capillaries promotes neuronal colonization by Toxoplasma gondii. Nat. Commun. *16*, 3529.
https://doi.org/10.1038/s41467-025-58655-z.

39. Kroll, J., Ruiz□Fernandez, M.J.A., Braun, M.B., Merrin, J., and Renkawitz, J. (2022).
Quantifying the Probing and Selection of Microenvironmental Pores by Motile Immune Cells. Curr
Protoc 2, e407. https://doi.org/10.1002/cpz1.407.

40. Periz, J., Rosario, M.D., McStea, A., Gras, S., Loney, C., Wang, L., Martin-Fernandez, M.L., and
Meissner, M. (2019). A highly dynamic F-actin network regulates transport and recycling of
micronemes in Toxoplasma gondii vacuoles. Nature Communications 10, 290.
https://doi.org/10.1038/s41467-019-12136-2.

41. Koreny, L., Mercado-Saavedra, B.N., Klinger, C.M., Barylyuk, K., Butterworth, S., Hirst, J.,
Rivera-Cuevas, Y., Zaccai, N.R., Holzer, V.J.C., Klingl, A., et al. (2023). Stable endocytic structures
navigate the complex pellicle of apicomplexan parasites. Nat. Commun. 14, 2167.
https://doi.org/10.1038/s41467-023-37431-x.

42. Kanatani, S., Uhlén, P., and Barragan, A. (2015). Infection by Toxoplasma gondii Induces
Amoeboid-Like Migration of Dendritic Cells in a Three-Dimensional Collagen Matrix. PLoS ONE
10, e0139104. https://doi.org/10.1371/journal.pone.0139104.

43. Wang, Y., Sangaré, L.O., Paredes-Santos, T.C., Hassan, M.A., Krishnamurthy, S., Furuta, A.M.,
Markus, B.M., Lourido, S., and Saeij, J.P.J. (2020). Genome-wide screens identify Toxoplasma
gondii determinants of parasite fitness in IFNγ-activated murine macrophages. Nat Commun 11,
5258. https://doi.org/10.1038/s41467-020-18991-8.

44. Paterson, N., and Lämmermann, T. (2021). Macrophage network dynamics depend on
haptokinesis for optimal local surveillance. eLife 11, e75354. https://doi.org/10.7554/elife.75354.

45. Clausen, B.E., Amon, L., Backer, R.A., Berod, L., Bopp, T., Brand, A., Burgdorf, S., Chen, L.,
Da, M., Distler, U., et al. (2023). Guidelines for mouse and human DC functional assays. Eur. J.
Immunol. 53, e2249925. https://doi.org/10.1002/eji.202249925.

46. Battistella, A., Andolfi, L., Zanetti, M., Zilio, S.D., Stebel, M., Ricci, G., and Lazzarino, M.
(2022). Atomic force spectroscopy□based essay to evaluate oocyte postovulatory aging. Bioeng.
Transl. Med. 7, e10294. https://doi.org/10.1002/btm2.10294.

47. Yun, X., Tang, M., Yang, Z., Wilksch, J.J., Xiu, P., Gao, H., Zhang, F., and Wang, H. (2017).
Interrogation of drug effects on HeLa cells by exploiting new AFM mechanical biomarkers. RSC
Adv. 7, 43764–43771. https://doi.org/10.1039/c7ra06233h.

48. Schlüßler, R., Kim, K., Nötzel, M., Taubenberger, A., Abuhattum, S., Beck, T., Müller, P.,
Maharana, S., Cojoc, G., Girardo, S., et al. (2022). Correlative all-optical quantification of mass
density and mechanics of subcellular compartments with fluorescence specificity. Elife 11, e68490.
https://doi.org/10.7554/elife.68490.

49. Kroll, J., and Renkawitz, J. (2024). Principles of organelle positioning in motile and non-motile
cells. EMBO Rep. 25, 2172–2187. https://doi.org/10.1038/s44319-024-00135-4.

50. Gundersen, G.G., and Worman, H.J. (2013). Nuclear positioning. Cell 152, 1376–1389.
https://doi.org/10.1016/j.cell.2013.02.031.

51. Renkawitz, J., Kopf, A., Stopp, J., Vries, I. de, Driscoll, M.K., Merrin, J., Hauschild, R., Welf,
E.S., Danuser, G., Fiolka, R., et al. (2019). Nuclear positioning facilitates amoeboid migration along
the path of least resistance. Nature 568, 546–550. https://doi.org/10.1038/s41586-019-1087-5.

52. Vorselen, D., Wang, Y., Jesus, M.M. de, Shah, P.K., Footer, M.J., Huse, M., Cai, W., and Theriot,
J.A. (2020). Microparticle traction force microscopy reveals subcellular force exertion patterns in
immune cell–target interactions. Nature Communications *11*, 1–14. https://doi.org/10.1038/s41467-
019-13804-z.

53. Mali, A., Peeters, Y., Mercado, R.R. de, Settle, A.H., Footer, M.J., Srinivas, M., Theriot, J.A., and
Vorselen, D. (2025). Using tunable hydrogel microparticles to measure cellular forces. Nat. Protoc.,
1–27. https://doi.org/10.1038/s41596-025-01281-2.

54. Guilak, F., Tedrow, J.R., and Burgkart, R. (2000). Viscoelastic Properties of the Cell Nucleus.
Biochem. Biophys. Res. Commun. *269*, 781–786. https://doi.org/10.1006/bbrc.2000.2360.

55. Bornens, M. (2008). Organelle positioning and cell polarity. Nature Reviews Molecular Cell
Biology *9*, 874–886. https://doi.org/10.1038/nrm2524.

56. Schmitt, M.T., Kroll, J., Ruiz-Fernandez, M.J.A., Hauschild, R., Ghosh, S., Kameritsch, P.,
Merrin, J., Schmid, J., Stefanowski, K., Thomae, A.W., et al. (2025). Protecting centrosomes from
fracturing enables efficient cell navigation. Sci. Adv. *11*, eadx4047.
https://doi.org/10.1126/sciadv.adx4047.

57. Kopf, A., Renkawitz, J., Hauschild, R., Girkontaite, I., Tedford, K., Merrin, J., Thorn-Seshold, O.,
Trauner, D., Häcker, H., Fischer, K.-D., et al. (2020). Microtubules control cellular shape and
coherence in amoeboid migrating cells. J. Cell Biol. *219*, e201907154.
https://doi.org/10.1083/jcb.201907154.

58. Groot, R.D., and Warren, P.B. (1997). Dissipative particle dynamics: Bridging the gap between
atomistic and mesoscopic simulation. J. Chem. Phys. *107*, 4423–4435.
https://doi.org/10.1063/1.474784.

59. Chabaud, M., Heuzé, M.L., Bretou, M., Vargas, P., Maiuri, P., Solanes, P., Maurin, M., Terriac,
E., Berre, M.L., Lankar, D., et al. (2015). Cell migration and antigen capture are antagonistic
processes coupled by myosin II in dendritic cells. Nature Communications *6*, 7526.
https://doi.org/10.1038/ncomms8526.

60. Moreau, H.D., Blanch-Mercader, C., Attia, R., Maurin, M., Alraies, Z., Sanséau, D., Malbec, O.,
Delgado, M.-G., Bousso, P., Joanny, J.-F., et al. (2019). Macropinocytosis Overcomes Directional
Bias in Dendritic Cells Due to Hydraulic Resistance and Facilitates Space Exploration. Dev. Cell *49*,
171-188.e5. https://doi.org/10.1016/j.devcel.2019.03.024.

61. Leithner, A., Eichner, A., Müller, J., Reversat, A., Brown, M., Schwarz, J., Merrin, J., Gorter,
D.J.J. de, Schur, F., Bayerl, J., et al. (2016). Diversified actin protrusions promote environmental
exploration but are dispensable for locomotion of leukocytes. Nat. Cell Biol. *18*, 1253–1259.
https://doi.org/10.1038/ncb3426.

62. García-Arcos, J.M., Jha, A., Waterman, C.M., and Piel, M. (2024). Blebology: principles of bleb-
based migration. Trends Cell Biol. https://doi.org/10.1016/j.tcb.2024.02.009.

63. Paluch, E.K., and Raz, E. (2013). The role and regulation of blebs in cell migration. Current
Opinion in Cell Biology *25*, 582–590. https://doi.org/10.1016/j.ceb.2013.05.005.

64. Krendel, M., Zenke, F.T., and Bokoch, G.M. (2002). Nucleotide exchange factor GEF-H1
mediates cross-talk between microtubules and the actin cytoskeleton. Nat Cell Biol *4*, 294–301.
https://doi.org/10.1038/ncb773.

65. Denais, C.M., Gilbert, R.M., Isermann, P., McGregor, A.L., Lindert, M. te, Weigelin, B.,
Davidson, P.M., Friedl, P., Wolf, K., and Lammerding, J. (2016). Nuclear envelope rupture and repair
during cancer cell migration. Science *352*, 353–358. https://doi.org/10.1126/science.aad7297.

66. Harada, T., Swift, J., Irianto, J., Shin, J.-W., Spinler, K.R., Athirasala, A., Diegmiller, R., Dingal,
P.C.D.P., Ivanovska, I.L., and Discher, D.E. (2014). Nuclear lamin stiffness is a barrier to 3D
migration, but softness can limit survival. J. Cell Biol. *204*, 669–682.
https://doi.org/10.1083/jcb.201308029.

67. Grinsven, E. van, Mukherjee, A.K., and Udalova, I.A. (2025). Shaping neutrophil morphology and
function: the importance of a segmented nucleus. Nat. Rev. Immunol., 1–15.
https://doi.org/10.1038/s41577-025-01237-3.

68. Weiner, A., Mellouk, N., Lopez-Montero, N., Chang, Y.-Y., Souque, C., Schmitt, C., and
Enninga, J. (2016). Macropinosomes are Key Players in Early Shigella Invasion and Vacuolar Escape
in Epithelial Cells. PLoS Pathog. *12*, e1005602. https://doi.org/10.1371/journal.ppat.1005602.

69. Kumar, Y., and Valdivia, R.H. (2008). Actin and Intermediate Filaments Stabilize the Chlamydia
trachomatis Vacuole by Forming Dynamic Structural Scaffolds. Cell Host Microbe *4*, 159–169.
https://doi.org/10.1016/j.chom.2008.05.018.

70. Welch, M.D., Rosenblatt, J., Skoble, J., Portnoy, D.A., and Mitchison, T.J. (1998). Interaction of
Human Arp2/3 Complex and the Listeria monocytogenes ActA Protein in Actin Filament Nucleation.
Science *281*, 105–108. https://doi.org/10.1126/science.281.5373.105.

71. Haglund, C.M., Choe, J.E., Skau, C.T., Kovar, D.R., and Welch, M.D. (2010). Rickettsia Sca2 is a
bacterial formin-like mediator of actin-based motility. Nat. Cell Biol. *12*, 1057–1063.
https://doi.org/10.1038/ncb2109.

72. Redecke, V., Wu, R., Zhou, J., Finkelstein, D., Chaturvedi, V., High, A.A., and Häcker, H. (2013).
Hematopoietic progenitor cell lines with myeloid and lymphoid potential. Nature methods *10*, 795–
803. https://doi.org/10.1038/nmeth.2510.

73. Sixt, M., and Lämmermann, T. (2011). In vitro analysis of chemotactic leukocyte migration in 3D
environments. Methods in molecular biology (Clifton, N.J.) *769*, 149–165.
https://doi.org/10.1007/978-1-61779-207-6_11.

74. Leithner, A., Renkawitz, J., Vries, I.D., Hauschild, R., Häcker, H., and Sixt, M. (2018). Fast and
efficient genetic engineering of hematopoietic precursor cells for the study of dendritic cell migration.
Eur. J. Immunol. *48*, 1074–1077. https://doi.org/10.1002/eji.201747358.

75. Braun, M., Reza, R., Kameritsch, P., Ruiz-Fernandez, M., Schmitt, M., Shnipova, M., Krug, A.,
Merrin, J., Polleux, J., and Renkawitz, J. (2024). Microenvironmental Regulation of Macropinocytosis
Facilitates Extracellular Fluid Sampling. https://doi.org/10.21203/rs.3.rs-4223901/v1.

76. Rosario, M.D., Periz, J., Pavlou, G., Lyth, O., Latorre□Barragan, F., Das, S., Pall, G.S., Stortz,
J.F., Lemgruber, L., Whitelaw, J.A., et al. (2019). Apicomplexan F□actin is required for efficient
nuclear entry during host cell invasion. EMBO Rep. *20*, e48896.
https://doi.org/10.15252/embr.201948896.

77. Huang, Y., Schell, C., Huber, T.B., Şimşek, A.N., Hersch, N., Merkel, R., Gompper, G., and
Sabass, B. (2019). Traction force microscopy with optimized regularization and automated Bayesian

parameter selection for comparing cells. Sci. Rep. *9*, 539. https://doi.org/10.1038/s41598-018-36896-
x.

78. Schlüßler, R., Möllmert, S., Abuhattum, S., Cojoc, G., Müller, P., Kim, K., Möckel, C.,
Zimmermann, C., Czarske, J., and Guck, J. (2018). Mechanical Mapping of Spinal Cord Growth and
Repair in Living Zebrafish Larvae by Brillouin Imaging. Biophys. J. *115*, 911–923.
https://doi.org/10.1016/j.bpj.2018.07.027.

79. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S.,
Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image
analysis. Nature methods *9*, 676–682. https://doi.org/10.1038/nmeth.2019.

705
706

Figure legends

Figure 1. Efficient transport of large assemblies of intracellular parasites by immune cells through dense environments. (A) Representative *Toxoplasma gondii*-infected dendritic cells (DCs) migrating in linear microchannels are shown as kymographs over 6.5 minutes. *Toxoplasma gondii* is shown in red, the host cell shape in yellow, and the DC nucleus in cyan (Hoechst). Note the different replication stages (1- to 32-stage) and respective sizes of the parasitic cargo within the motile host cell (see also the zoom-ins in the yellow boxes). **(B)** Quantification of the length of the parasitophorous vacuole in comparison to the host cell nucleus during DC migration along linear microchannels. Note that a 4-stage PV equals the length of the host nucleus, while 8-, 16, and 32-staged PVs exceed the size of the host nucleus. N = 8 cells (1-stage PV), 10 cells (2-stage PV), 24 cells (4-stage PV), 28 cells (8-stage PV), 37 cells (16-stage PV), and 9 cells (32-stage PV) from 3 independent biological replicates. Unpaired t-tests. Data are the mean±95% CI. **(C)** DC velocity in relation to the replication stage of the parasite cargo. N = 86 cells (non-infected bystander DCs), 22 cells (1-stage PV), 29 cells (2-stage PV), 50 cells (4-stage PV), 93 cells (8-stage PV), 67 cells (16-stage PV), and 11 cells (32-stage PV) from 3 independent biological replicates. Kruskal-Wallis multiple comparison test on bystander cells. Data are the mean±95% CI. **(D)** Macrophage velocity in relation to the replication stage of the parasite cargo. N = 28 cells (1-stage PV), 28 cells (2-stage PV), 34 cells (4-stage PV), 54 cells (8-stage PV), 39 cells (16-stage PV), and 17 cells (32-stage PV) from 4 independent biological replicates. Kruskal-Wallis multiple comparison test on 1-stage infected cells. Data are the mean±95% CI. See Figure S1A for representative image examples, Figure S1B for parasitic sizes within macrophages, and Figure S1D for passing times through 2 micron pores within macrophages. **(E)** Representative dendritic cell carrying a 16-stage *Toxoplasma* cargo across a 2-micron-wide pore within a microchannel. *Toxoplasma gondii* is shown in red, the host cell shape in yellow, and the DC nucleus in cyan (Hoechst). **(F)** Passing times of parasitic cargoes of different sizes during DC migration through 4 micrometer-sized pores. See Figure S1C for representative images. One-way ANOVA with multiple comparisons on bystander cells. Data are the mean±SD, N = 3 biological replicates. **(G)** Passing times of parasitic cargoes of different sizes during DC migration through 2 micrometer-sized pores, as shown in Figure 1E. One-way ANOVA with multiple comparisons on bystander cells. Data are mean±SD, N = 3 biological replicates.

Figure 2. Intracellular parasites cause migration bottlenecks for infected immune cell
hosts. (A) Representative fluorescent microscopy (left panel, T. gondii in red, host nucleus in
blue) and atomic force microscopy (Young's modulus, middle panel; dissipated viscous
energy, right panel) of a migrating dendritic cell in confinement and upon PFA-based
fixation. **(B)** Quantification of A), quantifying the Young's modulus of the dendritic cell's
nucleus (host cell), the individual parasites, and the regions within the parasitophorous
vacuole (PV) between the individual parasites ("PV matrix"). N = 23 cells from 4
independent biological replicates. Data are the mean±SEM. Statistics: Friedman ANOVA.
(C) Quantification of A), quantifying the dissipated viscous energy of the dendritic cell's
nucleus (host cell), the individual parasites, and the PV matrix. N = 23 cells from 4
independent biological replicates. Data are the mean±SEM. Statistics: Friedman ANOVA.
(D) Representative simultaneous fluorescent (left panel) and Brillouin microscopy (Brillouin
shift, middle panel; Brillouin linewidth, right panel) images of a *Toxoplasma*-infected
dendritic cell migrating in confinement and upon PFA-based fixation. **(E)** Quantification of
D), quantifying the Brillouin shift of intracellular *Toxoplasma* parasites in comparison to the
dendritic cell's nucleus. N = 26 cells from 4 independent biological replicates. Data are
mean±95% CI. Statistics: Wilcoxon matched-pairs signed rank test. **(F)** Quantification of D),
quantifying the Brillouin peak of intracellular *Toxoplasma* parasites in comparison to the
dendritic cell's nucleus. N = 26 cells from 4 independent biological replicates. Data are the
mean±95% CI. Statistics: Wilcoxon matched-pairs signed rank test. **(G)** Representative
indentation of the host nucleus by *Toxoplasma* parasites. See Figure S2A for quantification.
(H) Representative example of the squeezing of individual parasites in a 2-stage
parasitophorous vacuole (PV) within a dendritic cell during passage through a 2 micrometer-
sized pore. The yellow dashed square is shown in a zoom. **(I)** Representative 16-stage
example showing unfolding of the parasitophorous vacuole (PV) during passage within a
dendritic cell through a 2 micrometer-sized pore. The yellow dashed square is shown in a
zoom. **(J)** Quantification of the change in length of 16-stage parasites during passage through
2 micrometer-sized pores. N= 19 cells from 3 independent biological replicates. Data are the
mean±SEM. Statistics: Friedman ANOVA.

Figure 3. Intracellular positioning adapts to the size of the parasitic cargo. (A) Representative examples of the intracellular positioning of parasites forward or rearward relative to the host cell nucleus during dendritic cell migration in a straight 8 micrometer-wide microchannel. *Toxoplasma gondii* in red, host cell nucleus in cyan, host cell shape in yellow. **(B)** Quantification of the positioning of the parasite cargo in relation to the dendritic cell's host nucleus, as shown in B). Note the increasing frequency of parasitic frontward positioning with its size. Data are the mean±SD. Unpaired t-tests. N=3 individual biological replicates (23 cells 1-stage PV, 29 cells 2-stage PV, 52 cells 4-stage PV, 99 cells 8-stage PV, 76 cells 16-stage PV, and 12 cells 32-stage PV). **(C)** Representative examples of dynamic repositioning of PV cargoes in front of the host cell nucleus towards the host cell front. **(D)** Quantification of dynamic repositioning events in the parasite to host nucleus axis, as shown in C). Note the increasing frequency of parasitic forward positioning with its size. Data are the mean±SD. Unpaired t-tests. N=3 individual biological replicates (23 cells 1-stage PV, 29 cells 2-stage PV, 52 cells 4-stage PV, 99 cells 8-stage PV, 76 cells 16-stage PV, and 12 cells 32-stage PV). **(E)** Representative dendritic cells with deformable (6.5 kPa) beads as cargoes, either positioned forward or rearward of the dendritic cell's nucleus. Note how multiple beads cluster, comparable to the parasitic assemblies. Deformable beads in green, dendritic cell nucleus in cyan, dendritic cell shape in yellow. **(F)** Quantification of the intracellular positioning of deformable beads relative to the host cell nucleus, as shown in E). Data are the mean±SD. Unpaired t-tests. N=3 individual biological replicates (44 cells w/o beads, 25 cells 1 bead, 19 cells 2 beads, 31 cells 3-4 beads, 21 cells 5-8 beads, 22 cells 9-40 beads). **(G)** Quantification of dynamic repositioning events in the deformable bead to the dendritic cell's nucleus axis. Data are the mean±SEM. Unpaired t-tests. N=3 individual biological replicates (44 cells w/o beads, 25 cells 1 bead, 19 cells 2 beads, 31 cells 3-4 beads, 21 cells 5-8 beads, 22 cells 9-40 beads). **(H)** Quantification of the positioning of the parasite in relation to the host cell's microtubule-organizing center (MTOC) marked by EB3-mCherry, as shown in Figure S4E. Data are the mean±95% CI. N=3 individual biological replicates (19 cells 1-stage PV, 26 cells 2-stage PV, 22 cells 4-stage PV, 44 cells 8-stage PV, 18 cells 16-stage PV, and 5 cells 32-stage PV). See also the quantification of the parasite positioning in relation to the nucleus and the MTOC, as shown in Figure S4F. **(I)** Representative macrophage carrying an 8-stage parasite, in which the parasitic cargo dynamically repositions forward of the macrophage's nucleus. **(J)** Quantification of the positioning of the parasite cargo in relation to the macrophage's host nucleus. See also representative examples in Figure S1A. Data are the mean±SD. N=3 individual biological replicates (109 cells).

Figure 4. Myosin-driven host cell contractility adapts to parasite sizes. (A) Representative snapshots of simulating the intracellular positioning of parasites and the host nucleus in moving cells. *Toxoplasma gondii* in red, host cell nucleus in cyan, the host cytoplasm in gray (upper panel), the cell cortex in orange (upper panel), and the intracellular fluid particle density of the host in fire-color (lower panel). Note the repositioning of the parasite frontward of the host nucleus. **(B)** Simulation of parasite positioning in dependence of parasite size relative to the host nucleus, and the tension in the host cell rear that leads to an intracellular pressure gradient. **(C)** Representative examples of *Toxoplasma*-infected (upper panel) or bead-loaded (lower panel) Myh9-GFP (Myosin-II)-expressing dendritic cells migrating along linear paths. *Toxoplasma gondii* in red, host cell nucleus in cyan, Myh9-GFP in white or as fire-color. **(D)** Quantification of the Myh9-GFP signal over the cell length in *Toxoplasma*-infected dendritic cells. N=4 individual biological replicates (50 bystander cells, 11 cells 1-stage PV, 15 cells 2-stage PV, 38 cells 4-stage PV, 57 cells 8-stage PV, 30 cells 16-stage PV, and 4 cells 32-stage PV). **(E)** Quantification of the Myh9-GFP signal over the cell length in dendritic cells loaded with deformable beads. N=3 individual biological replicates (83 cells w/o beads, 73 cells 1 bead, 40 cells 2 beads, 37 cells 3-4 beads, 26 cells 5-10 beads). **(F)** Representative examples of protrusion shapes in uninfected or *Toxoplasma*-infected dendritic cells. Note the frequent appearance of smoother or even blebbing protrusions in infected dendritic cells, while uninfected cells typically show classical veil-like protrusions. *Toxoplasma gondii* in red and host cell nucleus in cyan. **(G)** Quantification of the frequency of veiled, smooth, or blebbing protrusions in uninfected or *Toxoplasma*-infected dendritic cells, as shown in F). Data are the mean±SEM. One-way ANOVA with Geisser-Greenhouse correction on the fraction of cells with veils. N=3 individual biological replicates (28 bystander cells, 12 cells 1-stage PV, 14 cells 2-stage PV, 14 cells 4-stage PV, 20 cells 8-stage PV, and 23 cells 16-stage PV). **(H)** Representative example of Lifeact-GFP-expressing, *Toxoplasma*-infected dendritic cell. Zoom-ins in the yellow boxes highlight a reduced actin signal within bleb-like protrusions, which is a typical feature of blebs. See also Figure S5A for comparison of uninfected bystander cells. **(I, J)** Positioning (I) and repositioning (J) of the parasitic cargo in the presence of the myosin II inhibitor para-nitro-blebbistatin (pNB) or DMSO controls. Note that the control cells are the same as in Fig. 3b. N=3 individual biological replicates (I: 22 cells 1-stage PV DMSO, 16 cells 1-stage PV pNB, 29 cells 2-stage PV DMSO, 34 cells 2-stage PV pNB, 50 cells 4-stage PV DMSO, 55 cells 4-stage PV pNB, 93 cells 8-stage PV DMSO, 62 cells 8-stage PV pNB, 67 cells 16-stage PV DMSO, 20 cells 16-stage PV pNB; J: 17 cells 1-stage PV DMSO, 8 cells 1-stage PV pNB, 18 cells 2-stage PV DMSO, 22 cells 2-stage PV pNB, 34 cells 4-stage PV DMSO, 38 cells 4-stage PV pNB, 69 cells 8-stage PV DMSO, 35 cells 8-stage PV pNB, 16 cells 16-stage PV DMSO, 16 cells 16-stage PV pNB). Unpaired t-tests. Data are the mean±SEM.

Figure 5. Adaptive myosin-driven contractility facilitates transport across local barriers. (A) Representative traction force measurement of a dendritic cell infected with *Toxoplasma gondii* (16-stage PV) migrating in confinement. Traction vectors are shown in orange, the DC nucleus in cyan, and the parasite in red. See Supplemental Figure 5C for representative examples of bead displacements that are the basis for the traction vector calculation, and Supplemental Figure 5D for the quantification of bead displacements. **(B)** Quantification of A), showing the average traction magnitude (Pa) of WT control cells and myosin-IIA knockout cells infected with *Toxoplasma gondii*. Note the increase in traction magnitudes in WT cells with increasing parasitic sizes, and the reduced traction magnitudes in myosin-IIA knockout cells. N = 29 cells (1-stage PV, Control), 23 cells (2-stage PV, Control), 23 cells (4-stage PV, Control), 29 cells (8-stage PV, Control), 36 cells (16-stage PV, Control), and 14 cells (32-stage PV, Control), and 6 cells (1-stage PV, Myosin-IIA KO), 5 cells (2-stage PV, Myosin-IIA KO), 8 cells (4-stage PV, Myosin-IIA KO), 19 cells (8-stage PV, Myosin-IIA KO), 22 cells (16-stage PV, Myosin-IIA KO), and 10 cells (32-stage PV, Myosin-IIA KO) from 5 independent biological replicates. Data are the mean±SEM. **(C)** Dendritic cell migration velocity through 8 micrometer-wide microchannels upon *Toxoplasma gondii* infection, in the presence of the myosin II inhibitor para-nitroblebbistatin (pNB) or DMSO controls. N=3 individual biological replicates (86 cells (bystander, Control), 23 cells (1-stage PV, Control), 29 cells (2-stage PV, Control), 52 cells (4-stage PV, Control), 99 cells (8-stage PV, Control), and 76 cells (16-stage PV, Control), and 103 cells (bystander, pNB), 16 cells (1-stage PV, pNB), 34 cells (2-stage PV, pNB), 55 cells (4-stage PV, pNB), 62 cells (8-stage PV, pNB), and 20 cells (16-stage PV, pNB)). Data are the mean±SD. **(D)** Representative kymographs of *Toxoplasma gondii*-infected motile dendritic cells encountering 4 (left) and 2 (right) micrometer-wide microchannel pores in the presence of the myosin II inhibitor para-nitro-blebbistatin. *Toxoplasma gondii* is shown in red, the host cell shape in yellow, and the DC nucleus in cyan (Hoechst). **(E)** Representative kymographs of bead(deformable)-loaded motile dendritic cells encountering 2 micrometer-wide microchannel pores. Deformable beads in green, dendritic cell nucleus in cyan, dendritic cell shape in yellow. **(F)** Quantification of successful passage of dendritic cells through 4 and 2 micrometer-wide microchannel pores upon *Toxoplasma gondii* infection, in the presence of the myosin II inhibitor para-nitroblebbistatin (pNB) or DMSO controls. N=3 individual biological replicates (84 cells (bystander, 2 µm, Control), 15 cells (1-2-stage PV, 2 µm, Control), 10 cells (4-stage PV, 2 µm, Control), 16 cells (8-stage PV, 2 µm, Control), and 11 cells (16-32-stage PV, 2 µm, Control), and 48 cells (bystander, 2 µm, pNB), 21 cells (1-2-stage PV, 2 µm, pNB), 14 cells (4-stage PV, 2 µm, pNB), 19 cells (8-stage PV, 2 µm, pNB), and 15 cells (16-32-stage PV, 2 µm, pNB), and 74 cells (bystander, 4 µm, Control), 63 cells (1-2-stage PV, 4 µm, Control), 28 cells (4-stage PV, 4 µm, Control), 14 cells (8-stage PV, 4 µm, Control), and 26 cells (16-32-stage PV, 4 µm, Control), and 37 cells (bystander, 4 µm, pNB), 28 cells (1-2-stage PV, 4 µm, pNB), 17 cells (4-stage PV, 4 µm, pNB), 9 cells (8-stage PV, 4 µm, pNB), and 5 cells (16-32-stage PV, 4 µm, pNB)). Data are the fraction of the total±95% CI. **(G)** Quantification of successful passage of bead(deformable)-loaded dendritic cells through 2 micrometers-wide microchannel pores. N=3 individual biological replicates (28 cells w/o beads, 11 cells 1 bead, 12 cells 2 beads, 14 cells 3-4 beads, 13 cells 5-10 beads). Data are the fraction of the total±95% CI. **(H)** Quantification of the frequency of subcategories, if deformable-bead loaded dendritic cells do not successfully pass the 2 micron-sized pore. N=3 individual biological replicates (9 cells 1 bead, 9 cells 2 beads, 11 cells 3-4 beads, 13 cells 5-8 beads, 7 cells 9-40 beads). Data are the fraction of the total±95% CI. **(I)** Representative parasitic egress from a *Toxoplasma*-infected dendritic cell through a 2 micron-pore. **(J)** Quantification of parasitic egress upon migration or attempt migration

through 4 and 2 micrometer-wide microchannel pores upon *Toxoplasma gondii* infection, in
the presence of the myosin II inhibitor para-nitroblebbistatin (pNB) or DMSO controls. N=3
individual biological replicates (15 cells (1-2-stage PV, 2 µm, Control), 10 cells (4-stage PV,
2 µm, Control), 16 cells (8-stage PV, 2 µm, Control), and 11 cells (16-32-stage PV, 2 µm,
Control), and 20 cells (1-2-stage PV, 2 µm, pNB), 13 cells (4-stage PV, 2 µm, pNB), 19 cells
(8-stage PV, 2 µm, pNB), and 15 cells (16-32-stage PV, 2 µm, pNB), and 63 cells (1-2-stage
PV, 4 µm, Control), 27 cells (4-stage PV, 4 µm, Control), 14 cells (8-stage PV, 4 µm,
Control), and 26 cells (16-32-stage PV, 4 µm, Control), and 28 cells (1-2-stage PV, 4 µm,
pNB), 17 cells (4-stage PV, 4 µm, pNB), 9 cells (8-stage PV, 4 µm, pNB), and 5 cells (16-32-
stage PV, 4 µm, pNB)). Data are the fraction of the total±95% CI.
917
918

Methods

Cell culture.

Dendritic cells, macrophages, and HoxB8-FL cells: Murine dendritic cells (DCs), macrophages, and immortalised hematopoietic precursor Hoxb8-FL cells[72], were cultured at 37°C in a humidified incubator with 5% CO_2. For cell culture, a basis medium composed of RPMI-1640 medium (Invitrogen, 21875034), 10% fetal bovine serum (Gibco, 10437-028), 1% penicillin-streptomycin (Sigma-Aldrich, P0781, 100 U/ml penicillin and 100 µg/ml streptomycin), and 50 µM 2-mercaptoethanol (Gibco, 31350-010), was supplemented with cell culture supernatant of growth-factor-expressing cells. DC medium was supplemented with 10% GM-CSF supernatant, macrophage medium was supplemented with 5% Flt3l supernatant and 20% M-CSF supernatant, and Hoxb8-FL medium was supplemented with 5% Flt3l supernatant and 1 µM estradiol (Sigma-Aldrich, E2758-1G). All centrifugations for washing or pelleting murine cells were performed at 300 g for 5 min. During experiments, all cells were maintained in imaging medium, phenol-free basis medium supplemented with 50 µM L-ascorbic acid (MilliporeSigma, W210901) to limit photobleaching and phototoxicity.

Bone marrow isolation from mice. All animal experiments were performed in accordance with the German Animal Welfare Act. The Core Facility Animal Models of the Biomedical Center (BMC) at LMU Munich housed the used mice. Bone marrow from male and female C57Bl6/J mice, sacrificed at 8 to 12 weeks of age, was used to generate bone marrow-derived dendritic cells (BMDCs). For this, mice were euthanised by cervical dislocation, femurs and tibiae were isolated, cut open and flushed with basis medium as previously described[73].

Differentiation of dendritic cells and macrophages. DCs were differentiated from hematopoietic progenitor cells, either from murine bone marrow, obtained from male or female 8- to 12-week-old C57Bl6/J wild-type or MyoIIA-Flox*CD11c-Cre mice[12], or from Lifeact-GFP, EB3-mCherry[57], or Myh9-GFP Hoxb8-FL progenitor cell lines, as previously described[74]. In short, progenitor cells were seeded in DC medium. On day 3, 20% GM-CSF supernatant DC medium was added to the cells at a 1:1 ratio and on day 6, half the culture medium was replaced with 20% GM-CSF supernatant DC medium without pipetting the remaining cells. DCs were frozen on day 8, except for those with the MyoIIA-Flox*CD11c-Cre genetic modification, which were differentiated until day 10, with an extra medium change on day 8, and then frozen on day 10. DCs were frozen in fetal bovine serum (FBS, Gibco, 10437-028) supplemented with 10% DMSO, first for 3 days at -72 °C in a suitable freezing container, then stored in liquid nitrogen storage and used for experiments up to 1 year later.
Macrophages were differentiated from wild-type Hoxb8-FL progenitor cells, as described before[75]. Hoxb8-FL progenitor cells were cultured in macrophage medium, with medium changes on days 3 and 5, and macrophages were used for experiments on day 6 of the differentiation.

Culturing of Toxoplasma gondii parasites. Toxoplasma gondii RH Δku80DiCre SAG1 gene was tagged with Halo following standard protocols established in the lab [31,41]. Parasites were maintained on HFFs (Human foreskin fibroblasts (HFFs) (RRID: CVCL_3285, ATCC) in DMEM (Dulbecco's modified Eagle's medium), 10% fetal bovine serum, 2 mM L glutamine and 25 mg/ml gentamicin, and maintained at 37°C and 5% CO_2. Cultured cells and parasites were screened against mycoplasma contamination with LookOut Mycoplasma detection kit (Sigma) and treated with Mycoplasma Removal Agent (Bio Rad), if needed.

Infection of dendritic cells with Toxoplasma gondii parasites. HFFs 4 cm dish cell cultures
were infected with 5×10^6 parasites and left to replicate for 48 h until replicating vacuoles
were ready to egress. Following the established protocols in the lab[76], parasites were released
by scratching and syringing, and then filtered to remove cell debris. Next, *Toxoplasma* cells
were pelleted, resuspended in prewarmed PBS, washed again, and resuspended in prewarmed
culture medium for the respective host cell type. For the infection with *Toxoplasma gondii,*
differentiated DCs were thawed 2-4 h before infection in a water bath at 37 °C, washed with
basis medium, resuspended in DC medium and incubated in a 6-well plate at 1×10^6 DCs in 3
ml medium per well. For their infection, macrophages were grown close to confluency and
used on day 6 of their differentiation after washing in PBS and adding 3 ml new macrophage
medium per well of 6-well plates. DCs and macrophages were challenged with 0.5×10^6
parasites per well for 20 h, constituting a ~1:2 ratio of parasites to immune cells. Then, the
cells were stained for 50 min with Janelia dye 646 HaloTag ligand (Promega, GA1120) at a
1:15000 dilution and two drops of NucBlue (Hoechst 33342, Invitrogen, R37605) were added
per well without medium change. Infected macrophages lost adhesion and were harvested by
flushing the culture dish surface with a 1 ml pipette. Stained DCs and MCs were pelleted and
resuspended in imaging medium, optionally supplemented with inhibitors or DMSO for
inhibitor experiments.

Migration assays.

PDMS microchannel assays: Microchannels were generated as previously described[39].
Polydimethylsiloxane (PDMS, Sylgard 184, Biesterfield) was mixed at a 1:10 elastomer-to-
curing agent ratio, poured over a silicon wafer with negative microchannel structures
imprinted by photolithography, degassed under vacuum, and cured at 80°C overnight. The
PDMS was cut into blocks of approximately 7x12 mm. Two parallel holes, each 2x7 mm,
were punched with a custom-made puncher at a 1 mm spacing in between. The devices and
glass coverslips, cleaned with isopropanol and ethanol in an ultrasonic cleaning bath, were
then cleaned in a plasma cleaner and bonded at 120°C. The PDMS devices were glued to a 6-
well plate with 18 mm holes. After 2 min of treatment in the plasma cleaner, the
microchannels were flooded with imaging medium and equilibrated in the cell culture
incubator. Imaging medium supplemented with 0.625 µg/ml CCL19 was loaded into one
hole, and 5×10^4 *T. gondii*-infected and stained DCs were loaded into the second hole.
Imaging was started after about 0.5-1 h, when the first cells entered the microchannels.
Linear microchannels that were 8 µm wide and 5 µm high were used for migration analysis.
For analysis of pore passage dynamics, the linear microchannels were constricted at a single
point to 2 or 4 µm over a 5 µm length. For the analysis of rigid bead positioning within DCs,
a pillar forest PDMS design was used, with a height of 5 µm, 7 µm-diameter pillars, and a 10
µm spacing between pillars.

1009

Under-agarose migration assay. For cell migration under confinement in a 2D plane, DCs
were allowed to migrate under 1% agarose towards the chemokine CCL19, as described
before[45] . Briefly, agarose solution was prepared by mixing 1 part 2×HBSS, 2 parts phenol-
free basis medium with 20% fetal bovine serum and 1 part high-molecular weight agarose
(Biozym Gold Agarose, 850152) dissolved in pure water at 4% (w/v), adjusted to 55 °C, and
the resulting 1% agarose mix was poured into 3.5 cm glass bottom dishes or per well of an 8-
well slide, 3 ml or 300 µl, respectively. The agarose was solidified at room temperature for 1

h, then equilibrated in a cell culture incubator and in a petri dish containing a wet paper towel
to provide additional humidity.
1019
1020   *Inhibitor experiments.* Inhibitors were added to the cells after the last washing step, directly
1021   after staining and before adding them to the experimental setup. Inhibitors were also added to
1022   the experimental environment and to the CCL19 medium. Para-Nitroblebbistatin was used at
1023   25 µM, with the same 1:2000 DMSO dilution in the medium as a control. Nocodazole was
1024   used at 1 µM, with the same 1:3333 DMSO dilution as a control.
1025
Deformable acrylamide co-acrylic acid beads. Deformable acrylamide co-acrylic acid beads
with a diameter of 6.2 µm and a stiffness of 3.9 kPa were generated as previously
described[52,53]. DAAM-particles (5% v/v) were covalently conjugated with 20 mg/mL BSA
and 0.1 mM TMR-cadaverine (Invitrogen, A1318). First, they were washed twice (3000 g
centrifugation for 1 min, and subsequent resuspension and mixing by vortexing) in activation
buffer (100 mM MES pH 6, 200 mM NaCl). Particles were incubated in activation buffer
with 0.1% (v/v) Tween 20, 40 mg/mL N-(3-Dimethylaminopropyl)-N'-ethylcarbodiimide
hydrochloride, and 20 mg/mL N-hydroxysuccimide for 15 min while mixing. Particles were
subsequently washed three times with PBS pH 6 with 0.2% (v/v) Tween 20. After the final
wash, particles were resuspended in PBS pH 6 with 0.2% (v/v) Tween 20 at 10% (v/v) solids
and mixed thoroughly. Then, 2x PBS pH 8.5 with 0.2% (v/v) Tween 20 was added and
particles were directly incubated with BSA. After 1h, 0.1 mM TMR-cadaverine was added,
and after another 0.5 h, blocking buffer consisting of 100 mM TRIS (pH 9), 100 mM NaCl
and 100 mM ethanolamine was added. After 0.5 h, particles were washed three times in PBS
pH 7.4 + 0.1% Tween 20. Beads were previously determined to have increased Young's
moduli of 6.5 kPa after BSA functionalization[52]. Beads were further opsonized with 0.03
mg/ml anti-BSA IgG antibody (MP Biomedicals, 08651111) in PBS with 0.1% Tween for 1
h under rotation, washed three times with PBS (pH 7.4) with 10% Tween with centrifugation
steps at 3000 g for 1 min, and stored in PBS supplemented with 5 mM sodium azide. The
beads were washed with PBS and resuspended in DC medium before adding them to DCs.
Differentiated DCs were thawed and 3.3×10^5 DCs were challenged with 1.5×10^6 deformable
beads for 20 h in 1 ml DC medium in wells of a 24-well plate. DCs were stained with one
drop of NucBlue (Hoechst 33342, Invitrogen, R37605) per well for 20 min, and the medium
was changed to imaging medium. Imaging during migration in linear microchannels or
microchannels with 2 µm constrictions was performed at 1 frame per minute.
1051
1052
1053
1054   **Imaging.**
1055
Live-cell imaging: Live-cell imaging was performed using a Leica DMi8 inverted wide-field
epifluorescence microscope with a motorised stage, under cell culture conditions of 37°C and
5% CO_2 in a humidified atmosphere (Pecon). Time-lapse recordings were performed using a
20x (Leica HC PL APO 20x/0.80, 11506529) or 40x (Leica HC PL APO 40x/0.95 CORR,
11506414) lens. The Leica sCMOS Camera DFC9000 GT was used as the camera, with
LED5 fluorescence light source. *Toxoplasma* signal (Janelia dye 646) was imaged with a far-
red filter (Excitation: 638/31, Emission: 695/58) and 640 nm excitation, EB3-mcherry signal
with an orange filter (Excitation: 554/24, Emission: 594/32) and 550 nm excitation, and the
nuclear signal (Hoechst 33342 or DAPI) with a blue filter (Excitation: 391/32, Emission:
435/30) and 395 nm excitation. Exposure to 395 nm light was kept to a minimum, just

enough to clearly localize the nucleus position and size, to avoid phototoxicity affecting
experimental results.
1068
STED imaging: Fixation of cells was performed as described in Del Rosario et al.[76] . Briefly,
cells were fixed in cytoskeleton buffer composed of two components: cytoskeleton buffer
(CB1) (MES pH 6.1 10 mM, KCL 138 mM, MgCl 3 mM, EGTA 2 mM, 5% PFA) and
cytoskeleton buffer (CB2) (MES pH6.1 10 mM, KCL 163.53 mM, MgCl 3.555 mM, EGTA
2.37 mM, sucrose 292 mM). These buffers were mixed in a 4:1 ratio, respectively, and
subsequently used for fixation for 10 min. The samples were quenched (NH_4CL 50 mM) for
10 min and washed three times in PBS. To visualise microtubules, rat anti-human alpha
tubulin (BioRad) was used at 1:300 in 2% BSA 0.2% triton x100 in PBS for 1 h, washed
three times in PBS, followed by an incubation with 1:1000 secondary antibody Alexa 488
(Life Technologies).
STED and confocal imaging were performed on an Abberior Expert Line STED
microscope (Abberior Instruments GmbH, Göttingen, Germany) built around an Olympus
IX83 inverted microscope body, equipped with an Olympus 100x / NA 1.4 Oil UPLSAPO
objective and the easy3D STED module (spatial light modulator for 2D and 3D beam
shaping). The system was controlled by Imspector software (Abberior Instruments). Samples
were excited with 405 nm, and pulsed 485 nm, or 640 nm laser diodes at 1%, 2%, 5% of
maximum power. Stimulated emission depletion was performed with a pulsed 595 nm STED
laser at 5% of maximum power (repetition rate 40 MHz) using a 2D doughnut depletion
pattern. The 405 nm channel was acquired only in confocal mode. Fluorescence was detected
on single-photon-counting avalanche photodiodes (APDs) with the following spectral
detection windows/ Filter Cubes: channel 1: 500–550 nm (485 nm excitation); channel 2:
650–720 nm (640 nm excitation). The pinhole was set to 1.0 Airy unit. Images were acquired
with 40 nm pixel size, 4 µs pixel dwell time, and 10 line accumulations.
1092
1093
1094 **Force and mechanical measurements.**
1095
Traction force microscopy. Glass coverslips (Marienfeld, 60 × 24 mm, #1.5H) were cleaned
in two stages. Initially, they were sonicated in a 5% v/v Hellmanex III solution for 10
minutes, followed by sonication in 100% ethanol. At each step, the coverslips were washed
rigorously in double-distilled water for 5 minutes. Finally, the coverslips were dried using
compressed nitrogen.
Silicone substrates were fabricated by combining a 1.2:1 weight ratio of two
components, Cy-A and Cy-B (Dow DOWSIL™ CY 52-276 Kit), resulting in a substrate
stiffness of approximately 2-3 kPa. The gel solution was thoroughly mixed, centrifuged, and
degassed before 400 µl were transferred onto a glass coverslip. The solution was spin-coated
for 90 s using a homemade spinner, with acceleration and deceleration every 30 seconds to
create a smooth, uniform surface. The gels were then cured at 80 °C for 3 h, after which they
were stored at RT. To functionalize the silicone surface, the gels were treated with a 7% (v/v)
solution of 3-aminopropyl trimethoxysilane (APTES) in pure ethanol for 2 hours. The gels
were washed with pure ethanol, a 1:1 ethanol:PBS mixture, and PBS alone, each for three
cycles.
As an imaging chamber, an 8-well bottomless sticky-slide (Ibidi, 80828) was mounted
on top of each PDMS substrate. Following three washes with PBS (Sigma-Aldrich, D8537),
100 µL of a 100 nm, red-orange fluorescent (540/560) FluoSpheres carboxylate-modified
microspheres (Invitrogen, F8800) solution at a dilution of 1:1,000 in PBS was mixed with a
final concentration of 100 µg/mL EDC and pipetted onto the top surface of the gel containing

200 µL PBS. The solution was incubated for 20 min at RT and then washed three times with
PBS.
To measure traction forces in the under-agarose assay, 300 µl of 1%-agarose was used
per well of an 8-well slide with the bottoms coated with the PDMS-TFM substrate described
above. At two sites per well, 5×10^3 *T.gondii*-infected DCs in 1 µl were injected under the
agarose layer and allowed to migrate towards a chemokine source of 7 µl 2.5 µg/ml CCL19
in a hole punched with a 3 mm puncher. Live imaging was started 30 min after the injection
of DCs and recorded at 30 s per frame.
Image processing was carried out in customized MATLAB scripts. Some crucial
functions employed in the workflow are written in *italics* below. First, raw time sequences of
fluorescent beads are normalized by *mat2gray* to reduce the brightness differences. We
correct the spatial drift resulting from fluctuations of the stage and the microscope by
normxcorr2 for each frame relative to the first frame. Then, aligned image sequences are
used for further processing. We select a frame as the reference frame when no cell is in the
field of view. We apply *detectMinEigenFeatures* to localize fluorescent beads with subpixel
precision at the reference frame. Next, the detected points are fed to a built-in optical flow
tracker in MATLAB *vision.PointTracker* to obtain the displacements at each frame relative to
the reference. Subsequently, tracking results are structured as input for traction
reconstruction. We calculate the traction fields using published software[77], in which we
employ regularized Fourier transform cytometry with optimal estimation of regularization
parameters by the Bayesian approach.
1137
Atomic force microscopy and Brillouin microscopy. To test the stiffness of the *T. gondii*
parasitophorous vacuole, we employed atomic force microscopy (AFM) and Brillouin
microscopy (BM). *T. gondii*-infected DCs were let migrate in an under-agarose assay. For
this, two parallel holes were punched in the 1%-agarose using a custom-made 2×7 mm
puncher at a 2 mm spacing. One hole was loaded with 25 µl of imaging medium
supplemented with 0.625 µg/ml CCL19 as chemoattractant, and the other holes were loaded
with 25 µl of imaging medium containing 1×10^5 *T. gondii*-infected DCs stained with only
Janelia dye 646 HaloTag ligand (Promega, GA1120). DCs migrated toward the chemokine
source, where they squeezed themselves under the 1% agarose gel. After 4 h of migration, the
DCs were fixed while still under the agarose layer with 3.7% paraformaldehyde in PBS for 1
h at 37°C, then washed three times with PBS. Then, the agarose was removed to make the
cells accessible for AFM and BM, the nuclei were stained with 0.1 µg/ml DAPI (4',6-
Diamidine-2'-phenylindole, #62248, Thermo Scientific), and the dishes were washed three
times with PBS.
For AFM measurements, a Nanowizard IV was used (JPK BioAFM, Bruker Nano
GmbH) with SNL-10 triangular pyramidal cantilevers (Bruker) with a tip radius of 2-12 nm
and a nominal spring constant of 0.35 N/m. Cantilevers were calibrated by the thermal noise
method directly inside a glass-bottom petri dish. Quantitative imaging (QI) mode (JPK,
Berlin) was used. Cells were scanned with a force load of 3 nN, at 150 µm/s, with a
resolution of 2.56 px/µm. The dissipated viscous energy was extracted by calculating the area
between the approach and retraction curves in the positive force range for each pixel of the
image using a custom-made Python script. Extraction of Young's modulus and height was
performed using the JPKSPM Data Processing software. Here, the Poisson's ratio was set to
0.5, and the Sneddon model for a triangular pyramidal indenter was applied for the fitting,
with a limitation of the indentation curve to 200 nm in the contact region.
Brillouin frequency shift images were obtained, as described before[78], by employing
a Brillouin spectrometer with a two-stage virtually imaged phase array VIPA etalon (Light
Machinery, cat. nr. OP-6721-6743-4), a diode laser beam (λ = 780.24 nm, DLC TA PRO 780,

Toptica), a Fabry-Perot interferometer in two-pass configuration and a monochromatic grating (Toptica), a molecular absorption cell filled with rubidium 85Rb (Precision Glassblowing cat nr. TG-ABRB-I85-Q), and a sCMOS camera for the Brillouin spectrum (Teledyne cat nr. Prime BSI). The system was integrated with an inverted microscope stand with motorised stage and focus (AxioObserver 7, Zeiss). For the measurements, a 40x objective lens was used and the exposure time was set to 0.2 s per measurement point. Confocal fluorescence images were acquired in the same region as the Brillouin microscopy with a rescan confocal microscopy module (RCM2, Confocal.nl) with a sCMOS camera (Prime BSI Express, Teledyne), fluorescence excitation with a multi-line laser unit (Skyra, Cobolt, excitation wavelength $\lambda = 405$ nm and 637 nm used), and a 50 µm pinhole.

For analysis of the AFM and BM data, epifluorescence (AFM) and confocal microscopy (BM) images were semi-manually aligned with the AFM and BM images, respectively, using a custom-made ImageJ script. Host nuclei and parasitic cargoes were segmented manually based on their respective fluorescence signals, excluding overlapping regions of both the parasite and the host nucleus. Mean BM and AFM values of the host nucleus and parasitic cargo were extracted using a custom-made ImageJ script.

Image Analysis.

Software: Fiji (version 2.14.0) [79] and LAS X (Version 3.7.6) were used for image processing. Only cells not interacting with other cells, and with single 1-32 stage PVs or uninfected bystanders, were analyzed. For each cell, individual parasite cells inside the PV were counted to assess the PV stage. All data was grouped accordingly. Cell positions were tracked using the ROI manager of Fiji and quantified using custom-made ImageJ scripts. Single-cell data were further analyzed and summarized with R (R version: 4.3.3, R-studio version 2023.12.1.402, tidyverse package version 2.0.0) and exported to GraphPad Prism (version 8.0.1) for statistical analysis.

Analysis of migration in straight channels: PV-host nucleus axis, migration speed, PV size. For the orientation of the *Toxoplasma* PV–host nucleus axis, DCs or macrophages were analyzed as they migrated in linear PDMS microchannels of varying widths (8, 10, or 16 µm). The same analysis was performed for DCs carrying beads as well. PV/beads-nucleus orientation and position were documented at the beginning and end of cells migrating ~500 µm through the microchannels. From this, the frequency of cells with PV-nucleus or bead-nucleus orientations and the frequency of repositioning changes were determined, and the average migration speed over the ~500 µm distance travelled was calculated for each cell. PV or nucleus size was measured as the length of the longer axis of the PV and the nucleus. Cells were imaged at a constant time interval, depending on the experiment, ranging from 45 to 75 seconds.

Analysis of migration in 2 and 4 µm constriction channels. Pore passing dynamics were quantified for DCs or macrophages challenged with *T. gondii,* or for DCs challenged with beads, migrating in linear microchannels of 8 µm width and 5 µm height with a 2 or 4 µm constriction. The passage time was determined by measuring the time between the first time point at which the cell, the nucleus, or the PV entered the constriction on one side and the first time point at which the constriction was fully excited. Further, cells that entered but did not exit the constriction, or that egressed during the 5 h recording period, were documented to calculate the frequency of DC passing, DC being stuck in the constriction, or *T. gondii* egressing. Cells were imaged at 2 min time interval for the fraction of cells passing analysis

and at 10 s for the PV-unfolding analysis and analysis of pore passage of the nucleus and PV
compared to the whole host cell. For the PV-unfolding analysis, the length of the PV was
measured just before the host DC touched the constriction, when the PV touched but did not
yet enter the constriction, just after the PV left the constriction, and when the entire host DC
left the constriction. Further, the passaging time of single parasites within the PV was
measured by determining the time between PV touching both sides of the constriction for the
first time, the rear of the first parasite cell passing the middle point of the constriction, and
then each subsequent parasite cell passing the middle point of the constriction with its rear.
1224
Analysis of EB3-mCherry and Myosin-GFP live fluorescence. DCs derived from EB3-
mcherry- or Myosin-GFP-expressing Hoxb8 cells were imaged while migrating through 8
µm-wide, 5 µm-high microchannels. For the microtubule-organizing center (MTOC)
positioning analysis, cells with a clearly defined MTOC, as a single brightest spot in the EB3-
mCherry fluorescence signal that was consistently present throughout the recording, were
analyzed. PV-nucleus-MTOC axis orientation was documented after cells migrated 300-500
µm in the microchannels. The localization of Myosin-GFP was measured using the Fiji line
profile over an 8 µm-wide region from the rear to the front of the DC. This was performed by
using a custom-made ImageJ script and R script to subtract background signal determined for
each replicate, normalize the fluorescence intensity to the average of the entire cell for each
cell, resulting in a mean intensity of 1 for each cell, and then the length of each cell was
normalized to 0 (rear) to 100 (front).
1237
Analysis of blebbing and smooth phenotype. Bone marrow derived *T. gondii*-infected DCs
were imaged while migrating in 8 µm wide and 5 µm high microchannels and categorized
manually in (1) veiled, for cells with constantly veiled protrusions containing many vesicles,
(2) smooth, for cells with a non-veiled cell front not containing vesicles most of the time, or
(3) blebbing cells, for cells showing clear signs of frequent quickly developing blebs in some
of the frames.
1244
1245
1246 **Computational model.**
1247 See the extended supplemental file for details.
1248
Statistics.
All experiments were performed at least as three independent biological replicates, and all
replicates were validated independently. Statistical analysis was conducted using GraphPad
Prism (version 8.0.1) with the appropriate tests for a normal or non-normal distribution, as
stated in the figure legend. Fractions of cell subpopulations were calculated from pooled cell
numbers of at least three independent biological replicates, and the 95% confidence intervals
of the fractions were calculated using the Wilson/Brown hybrid method in Prism. Error bars
are defined in the figure legends.
1257
1258

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contacts, Jörg Renkawitz (Joerg.Renkawitz@med.uni-muenchen.de), Benedikt Sabass (Benedikt.Sabass@TU-dortmund.de), and Javier Periz (Javier.Periz@para.vetmed.uni-muenchen.de).

Materials availability

All newly generated materials associated with the paper are available upon request from the lead contacts.

Data and code availability

All data supporting the findings of this study are available within the paper. The particle-based model presented in the paper is described in detail in the Supplementary Information and is publicly available: https://github.com/CellMicroMechanics/ToxoDCMigrationSimulationModel. Any additional information required to reanalyze the data reported in this work is available from the lead contacts upon request.

ACKNOWLEDGEMENTS

We thank Kasia Stefanowski for excellent technical assistance and Janina Kroll for critical reading of the manuscript, Ana-Maria Lennon-Dumenil and Aline Yatim for bone marrow from MyoIIA-Flox*CD11c-Cre mice, Michael Sixt for Myh9-GFP expressing HoxB8 cells, and the Core Facility Bioimaging, the Core Facility Flow Cytometry, and the Animal Core Facility of the Biomedical Center (BMC) for excellent support. This study was supported by the Deutsche Forschungsgemeinschaft (DFG; German Research Foundation; Priority Programme SPP2332, project 492014049, to JR and BS; and DFG project ME 2675/6-2 to JP), the Peter Hans Hofschneider Professorship of the foundation "Stiftung Experimentelle Biomedizin" (to JR), the LMU Institutional Strategy LMU-Excellent within the framework of the German Excellence Initiative (to JR), and the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (BacForce, G.A.No. 852585, to BS).

AUTHOR CONTRIBUTIONS

Mauricio J.A. Ruiz-Fernandez: Conceptualization; investigation; methodology; writing – review and editing.
Jianfei Jiang: Software and simulation; methodology; writing – review and editing.
Alice Battistella: Investigation; writing – review and editing
Armina Mortazavi: Investigation; methodology; writing – review and editing.
Sterre van Wierst: Methodology; writing – review and editing.
Bingzhi Wang: Investigation; writing – review and editing.
Artur Kuznetcov: Investigation; writing – review and editing.
Rüya Aslan: Investigation; writing – review and editing.
Jack Merrin: Methodology; writing – review and editing.
Daan Vorselen: Supervision; methodology; writing – review and editing.
Jochen Guck: Supervision; methodology; writing – review and editing.
Markus Meissner: Supervision; methodology; writing – review and editing.
Javier Periz: Conceptualization; supervision; investigation; writing – review and editing.

1308 Benedikt Sabass: Conceptualization; supervision; investigation; funding acquisition; project
1309 administration; writing – review and editing.
1310 Jörg Renkawitz: Conceptualization; supervision; investigation; funding acquisition; project
1311 administration; writing - original draft; writing – review and editing.

1312

DECLARATION OF INTEREST
1314 The authors declare no competing interests.

1315

Declaration of generative AI and AI-assisted technologies in the writing process
1317 The manuscript was edited for grammatical correctness and readability using
1318 Grammarly.com and ChatGPT. The AI-assisted modifications were restricted to improving
1319 readability, with all scientific content and conclusions remaining the sole responsibility of the
1320 authors.

1321

1322

SUPPLEMENTAL INFORMATION
1324 Supplemental Figures S1 to S5
1325 Supplemental Movies S1 to S11
1326 Supplemental information to numerical model

1327

1328

 SUPPLEMENTAL INFORMATION

1330
1331
1332
1333 **Supplemental Figure Legends**

1334

Supplemental Figure 1. Migration and crossing of microenvironmental barriers by
***Toxoplasma*-infected macrophages carrying large parasitic cargoes. (A)** Representative
Toxoplasma gondii-infected macrophages migrating in linear microchannels are shown as
kymographs over time. *Toxoplasma gondii* SAG1-Halo, stained with Janelia dye 646
HaloTag ligand, is shown in red, the host cell shape in yellow, and the macrophage's nucleus
in cyan (Hoechst). Note the different replication stages (1- to 32-stage) and sizes of the
parasite within the motile host macrophage. See Figure 1D for quantification of migration
speed. **(B)** Quantification of the length of the parasitophorous vacuole in comparison to the
host cell nucleus during macrophage migration along linear microchannels, as shown in A).
Note that a 4-stage PV equals the length of the host nucleus, while 8-, 16, and 32-staged PVs
exceed the size of the macrophage's nucleus. N = 14 cells (1-stage PV), 19 cells (2-stage
PV), 21 cells (4-stage PV), 32 cells (8-stage PV), 15 cells (16-stage PV), and 7 cells (32-stage
PV) from 4 independent biological replicates. Paired t-tests. Data are the mean±95% CI. **(C)**
Representative dendritic cell carrying a high-stage *Toxoplasma* cargo across a 4-micron-wide
pore within a microchannel. *Toxoplasma gondii* is shown in red and the DC nucleus in cyan
(Hoechst). See Figure 1F for quantification. **(D)** Passing times of parasitic cargoes of
different sizes during macrophage (MC) migration through 2 micrometer-sized pores. Data
are the mean±95% CI. Unpaired, non-parametric data: Mann-Whitney test. N = 26 (1- & 2-
stage PVs), 34 (4- & 8-stage PVs), 22 (16- & 32-stage PVs) from 4 independent biological
replicates.

1355
1356

1357

Supplemental Figure 2. Intracellular *Toxoplasma* parasites are a major physical
bottleneck within motile dendritic cells and macrophages. (A) Quantification of the
frequency of indentation of the dendritic cell (host) nucleus in *Toxoplasma*-infected dendritic
cells in the direction of the parasitophorous vacuole (PV), categorized in PVs that are either
directly next to the host nucleus, close by (up to 1 micrometer) or distant (more than 1
micrometer). See representative image in Figure 2G. Data are mean±SD, N = 52 cells. **(B)**
Live-cell imaging of a representative *Toxoplasma*-infected dendritic cell migrating through a
maze-like microenvironment composed of pillars, which connect the lower and upper
surfaces and are spaced at 10 micrometer distances. *Toxoplasma gondii* is shown in red, the
host cell shape in yellow, and the DC's nucleus in cyan (Hoechst). Note the dynamic
deformations of the host nucleus by the parasitic cargo (see zoom-ins in the yellow boxes).
(C) Time of passage through 2 micrometer-sized pores of the 1st and 2nd translocating
parasite within a parasitophorous vacuole (PV) within a motile dendritic cell. Longer passage
times of the 1st parasite in comparison to the 2nd parasite are shown as red connecting lines,
whereas the same or shorter passage times are shown in grey and black, respectively. Data
are the mean±95%. CI Paired, non-parametric data: Wilcoxon matched-pairs signed rank test.
N = 14 (2-stage PVs), 14 (4-stage PVs), 20 (8-stage PVs), 23 (16-stage PVs) from 3
independent biological replicates. **(D)** Representative example of the squeezing of an
individual parasite (1-stage parasitophorous vacuole (PV)) within a macrophage during
passage through a 2 micrometer-sized pore. *Toxoplasma gondii* is shown in red, the host cell
shape in yellow, and the DC's nucleus in cyan (Hoechst). The yellow dashed square is shown
in a zoom. **(E)** Representative 16-stage example showing unfolding of the parasitophorous
vacuole (PV) during passage within a macrophage through a 2 micrometer-sized pore.
Toxoplasma gondii is shown in red, the host cell shape in yellow, and the DC's nucleus in
cyan (Hoechst). The yellow dashed square is shown in a zoom. **(F-H)** Passing times of the
host cell body (F), the host nucleus (G), and the parasitophorous vacuole (H) while
Toxoplasma-infected dendritic cells cross a 2-micrometer pore. Data are the mean±95% CI.
Non-paired, non-parametric data: Kruskal-Wallis multiple-comparison test. N = 28 (non-
infected bystander DCs), 12 (1-stage PVs), 14 (2-stage PVs), 14 (4-stage PVs), 20 (8-stage
PVs), 23 (16-stage PVs) from 3 independent biological replicates. **(I)** Relative passing times
of the parasitic cargo and the host nucleus in relation to the host cell passing time. Data are
the mean±95% CI. Paired, non-parametric data: Multiple Wilcoxon tests. N = 28 (non-
infected bystander DCs), 12 (1-stage PVs), 14 (2-stage PVs), 14 (4-stage PVs), 20 (8-stage
PVs), 23 (16-stage PVs) from 3 independent biological replicates.

1392
1393

1394

Supplemental Figure 3. The intracellular position of the parasitic cargo and its
repositioning dynamics change with parasite size. (A) Representative kymographs of
Toxoplasma-infected dendritic cells migrating in 10- and 16-micrometer-wide linear channels
(1.7 minutes between each frame in the kymograph). *Toxoplasma gondii* is shown in red and
the DC's nucleus in cyan (Hoechst). **(B)** Quantification of the parasitophorous vacuole (PV)
to nucleus axis configuration during dendritic cell migration in 10- and 16-micrometer-wide
linear channels, as shown in A). N=3 individual biological replicates (20 cells (1-stage PV,
10 µm), 33 cells (2-stage PV, 10 µm), 55 cells (4-stage PV, 10 µm), 38 cells (8-stage PV, 10
µm), 9 cells (16-stage PV, 10 µm) and 11 cells (1-stage PV, 16 µm), 25 cells (2-stage PV, 16
µm), 39 cells (4-stage PV, 16 µm), 49 cells (8-stage PV, 16 µm), 14 cells (16-stage PV, 16
µm)). See also Figure 3B for comparison to 8-micrometer-wide linear channels. **(C)**
Positioning of the parasitophorous vacuole (PV) in relation to the host cell nucleus during
migration of *Toxoplasma*-infected dendritic cells (DCs) through a maze of pillars (10
micrometer spacing between pillars). See also representative images in Figure S2B. N= 22
cells (1-stage PV), 23 cells (2-stage PV), 17 cells (4-stage PV), 17 cells (8-stage PV), 2 cells
(16-stage PV) from 3 independent biological replicates. Data are mean±95% CI. **(D)**
Representative kymographs of parasitic cargo repositioning in 10- and 16-micrometer-wide
linear channels (3.4 minutes for the 10-micrometer-wide channel, and 1.7 minutes for the 16-
micrometer-wide channel between each frame in the kymograph). *Toxoplasma gondii* is
shown in red and the DC's nucleus in cyan (Hoechst). **(E)** Quantification of the dynamic
repositioning of PV cargoes towards the host cell front, frontward of the host nucleus, as
shown in D). N=3 individual biological replicates (20 cells (1-stage PV, 10 µm), 33 cells (2-
stage PV, 10 µm), 56 cells (4-stage PV, 10 µm), 39 cells (8-stage PV, 10 µm), 9 cells (16-
stage PV, 10 µm), and 12 cells (1-stage PV, 16 µm), 27 cells (2-stage PV, 16 µm), 40 cells
(4-stage PV, 16 µm), 50 cells (8-stage PV, 16 µm), 14 cells (16-stage PV, 16 µm)).

1420
1421

Supplemental Figure 4. Immune cell motility with non-deformable and deformable bead cargoes. (A) Representative dendritic cell (DC; nucleus in cyan) carrying a 2-stage parasitic cargo (red) and a 6 micron-sized rigid bead (green) migrating through the maze of a 3D pillar forest. **(B)** Quantification of the PV to nucleus axis configuration in relation to rigid latex beads, as shown in A). N=21 cells from 3 individual biological replicates. Data are shown as violin plots with the median. **(C)** Representative dendritic cell (DC) with a cargo composed of two deformable (6.5 kPa) beads, migrating in a linear microchannel with a two-micrometer pore. Deformable beads in green and dendritic cell nucleus in cyan. Note the deformation of the DC nucleus by the deformable bead. **(D)** Transport and positioning of deformable beads (as in C) within motile DCs along linear microchannels. These representative cells show examples of forward and rearward positioning of the beads relative to the host nucleus within cells carrying small (1 bead), intermediate (7 beads), or large (11 beads) cargoes of deformable beads. See Figure 3F for quantification. **(E)** Representative example of EB3-mcherry expressing dendritic cells (white; label for the MTOC) infected with *Toxoplasma gondii* shown in red. The host nucleus is shown in cyan, and the position of the MTOC is marked with a purple arrow. **(F)** Quantification of the intracellular configuration of the DC MTOC, DC nucleus, and parasite axis, showing that most cells position the parasite together with the host nucleus forward of the host MTOC. N=3 individual biological replicates (19 cells (1-stage PV, 10 µm), 26 cells (2-stage PV, 10 µm), 22 cells (4-stage PV, 10 µm), 44 cells (8-stage PV, 10 µm), 18 cells (16-stage PV, 10 µm), and 5 cells (32-stage PV, 16 µm).

1446

Supplemental Figure 5. Adaptation of Host Cell Forces to the Size of the Parasitic Cargo. (A) Representative examples of Lifeact-GFP-expressing dendritic cells carrying different stages of *Toxoplasma gondii* (small 2-stage and large 8-stage cargoes, as well as uninfected bystander cell). Zoom-ins in the yellow boxes highlight the different distributions of actin within the protrusions. Note the reduced actin signal within bleb-like protrusions in the dendritic cells carrying a large 8-stage parasite. **(B)** Representative images of alpha-tubulin immunofluorescence stainings (white) of control (DMSO) or microtubule-inhibited (nocodazole) dendritic cells (nucleus in blue; DAPI) infected with *Toxoplasma gondii* (red; SAG1-Halo). **(C)** Quantification of the blebbing and smooth cell front phenotype in the presence of the microtubule inhibitor nocodazole or DMSO controls. N=5 individual biological replicates (16 cells (1-stage PV, DMSO), 17 cells (2-stage PV, DMSO), 13 cells (4-stage PV, DMSO), 33 cells (8-stage PV, DMSO), 42 cells (16-stage PV, DMSO), and 4 cells (32-stage PV, DMSO), and 13 cells (1-stage PV, nocodazole), 11 cells (2-stage PV, nocodazole), 13 cells (4-stage PV, nocodazole), 26 cells (8-stage PV, nocodazole), 31 cells (16-stage PV, nocodazole), and 3 cells (32-stage PV, nocodazole)). **(D)** Representative image (left) and heat map (right) of TFM-beads and their displacement during the migration of *Toxoplasma*-infected dendritic cells. **(E)** Quantification of D, comparing beads displacements in wild-type (WT) and myosin-IIA knockout dendritic cells. N = 29 cells (1-stage PV, Control), 23 cells (2-stage PV, Control), 23 cells (4-stage PV, Control), 29 cells (8-stage PV, Control), 36 cells (16-stage PV, Control), and 14 cells (32-stage PV, Control), and 6 cells (1-stage PV, Myosin-IIA KO), 5 cells (2-stage PV, Myosin-IIA KO), 8 cells (4-stage PV, Myosin-IIA KO), 19 cells (8-stage PV, Myosin-IIA KO), 22 cells (16-stage PV, Myosin-IIA KO), and 10 cells (32-stage PV, Myosin-IIA KO) from 5 independent biological replicates. Data are the mean±SEM.

1471
1472
1473

Supplemental Movies

Supplemental Movie S1. Transport of *Toxoplasma*-cargoes by infected dendritic cells through confining microenvironments. Representative dendritic cells (DCs) infected with *Toxoplasma gondii* were imaged while migrating along confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Note the different replication stages (1, 2, 4, 8, 16, and 32-stages) and thus sizes of the parasitic cargo and that the parasitic cargoes can position in front and in the rear of the host. *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the DC nuclei are shown in cyan (Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S2. Transport of *Toxoplasma*-cargoes by infected macrophages through confining microenvironments. Representative macrophages infected with *Toxoplasma gondii* were imaged while migrating along confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Note the different replication stages (1, 2, 4, 8, 16, and 32-stages) and thus sizes of the parasitic cargo. *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the macrophages nuclei are shown in cyan (Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S3. *Toxoplasma*-cargoes passing 2 micrometer-wide pores during their transport in infected dendritic cells. Representative dendritic cells (DCs) infected with *Toxoplasma gondii* were imaged while migrating through a two micrometer-wide pore within confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Note the different replication stages (1, 2, 4, 8, and 16-stages) and thus sizes of the parasitic cargo. Individual parasites squeeze and deform during the crossing of this microenvironmental barrier. The representative 2- to 16-stage parasitic cargoes show unfolding and refolding during the pore passage, allowing individual parasites to pass one-by-one. *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the DC nuclei are shown in cyan (Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S4. *Toxoplasma*-cargoes passing 2 micrometer-wide pores during their transport in infected macrophages. Representative macrophages infected with *Toxoplasma gondii* were imaged while migrating through a two micrometer-wide pore within confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Note the different replication stages (1, 2, 4, 8, and 16-stages) and thus sizes of the parasitic cargo. Individual parasites squeeze and deform during the crossing of this microenvironmental barrier. The representative 2- to 16-stage parasitic cargoes show unfolding and refolding during the pore passage, allowing individual parasites to pass one-by-one. *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the macrophage nuclei are shown in cyan (Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S5. Active repositioning of *Toxoplasma*-cargoes towards the cell fronts of infected, motile dendritic cells. Representative dendritic cells (DCs) infected with *Toxoplasma gondii* were imaged while migrating along confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Note that the parasitic cargoes can reposition from the cell front to behind the host nucleus (1-stage) or overtake the host nucleus to reposition towards the cell front (4- to 32-stage). *Toxoplasma*

gondii parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the DC nuclei are shown in cyan (NucBlue, Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S6. Migration of dendritic cells carrying deformable bead cargoes. Representative dendritic cells (DCs) migrating along confining, linear microchannels (8 micron wide and 5 microns high) while carrying different numbers of deformable beads (1, 2, 4, 7, and 11 beads; green). The DC nucleus (Hoechst) is shown in blue. The representative examples show the two possible configurations of beads and the DC nucleus along the cell axis (beads in front or rear of the DC nucleus). Time is indicated as minutes (min).

Supplemental Movie S7. Positioning of the parasitic cargo relative to the microtubule-organising center. Representative dendritic cells (DCs) expressing EB3-mCherry, as a microtubule-organising center (MTOC) marker, infected with *Toxoplasma gondii* were imaged while migrating along confining, linear microchannels (8 micron wide and 5 microns high) towards a CCL19 chemokine gradient. Representative uninfected bystander DCs and DCs carrying 1- or 4-stage parasitic cargoes are shown in the most prominent MTOC-nucleus (uninfected) or parasite-MTOC-nucleus conformations. Lastly, a DCs carrying a 16-stage parasitic cargo is shown, demonstrating how the parasite can overtake the host MTOC while still behind the host nucleus, notably also incurring some deformation of the parasitic cargo for it to fit next to the MTOC. EB3 microtubules tips and MTOC visible in grey (bottom videos, EB3-mCherry), *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand) and the DC nuclei are shown in cyan (NucBlue, Hoechst 33342). The positioning of the parasite to the host MTOC and nucleus is indicated at the top and the time is indicated in minutes (min).

Supplemental Movie S8. Simulating the intracellular localization of the parasitic cargo and the host nucleus. Representative movie from the computational simulation, showing the parasitic cargo in red, the host cytoplasm in gray, the host cell cortex in orange and the host nucleus in blue (first movie part). The second movie part shows additionally the intracellular fluid particle density. See main text, main figures, and the Supplemental Information for further details.

Supplemental Movie S9. Localization of myosin during the migration of dendritic cells carrying parasitic or deformable bead cargoes. Representative myosin-GFP (white) expressing dendritic cells (DC) migrating along confining, linear microchannels while carrying different parasitic stages (red; 1- and 8-staged) or different numbers of deformable beads (green; 1 and 9 beads). The DC nucleus (Hoechst) is shown in blue. Note the rearward localized myosin accumulation in the DC carrying a large, 8-staged parasitic cargo. Time is indicated as minutes (min).

Supplemental Movie S10. Blebbing cell protrusions in dendritic cells carrying large parasitic cargoes. First part: Representative dendritic cells (DC) migrating along confining, linear microchannels while carrying different sizes of parasitic cargo (red). Note different protrusion shapes, including bleb-like protrusions. Second part: Representative Lifeact-GFP (green) expressing dendritic cell (DC). migrating along confining, linear microchannels while carrying a large parasitic cargo (red). Note the quick extension of the bleb that is initially devoid of a Lifeact-GFP signal. Time is indicated as minutes (min).

Supplemental Movie S11. Forces onto the environment by *Toxoplasma gondii*-infected dendritic cells. Representative example of traction force microscopy, showing dendritic cells (DC) carrying either a small, 2-stage or a large, 16-stage parasitic cargo. Traction vectors are shown in orange, *Toxoplasma gondii* parasites expressing SAG1-Halo are shown in red (Janelia dye 646 Halo ligand), and the DC nuclei are shown in cyan (NucBlue, Hoechst 33342). Time is indicated as minutes (min).

Supplemental Movie S12. Impaired translocation of *Toxoplasma*-cargoes through narrow pores upon myosin inhibition, and when carrying deformable beads. First part: Representative dendritic cell (DC) infected with *Toxoplasma gondii*, and migrating through a 4 micrometer-sized pore in the presence of the myosin inhibitor para-nitroblebbistatin. Second part: Representative dendritic cell (DC) infected with *Toxoplasma gondii*, trying to migrate through a 2 micrometer-sized pore in the presence of the myosin inhibitor para-nitroblebbistatin. Third part: Representative dendritic cell (DC) carrying deformable beads as a cargo, and trying to migrate through a 2 micrometer-sized pore. Fourth part: Representative parasitic egress at a 2 micrometer-sized pore. The movies show representative cells from at least three independent biological replicates. Time is indicated as minutes (min).

Figure 1. Efficient Transport of Large Assemblies of Intracellular Parasites by Immune Cells Through Dense Environments.

Figure 2. Intracellular Parasites Cause Migration Bottlenecks for Infected Immune Cell Hosts.

Figure 3. Intracellular Positioning Adapts to the Size of the Parasitic Cargo.

Figure 4. Myosin-driven Host Cell Contractility Adapts to Parasite Sizes.

Figure 5. Adaptive Myosin-driven Contractility Facilitates Transport Across Local Barriers.