Differences in size and number of embryonic type-II neuroblast lineages correlate with

divergent timing of central complex development between beetle and fly

Authors: Simon Rethemeier[1,2], Sonja Fritzsche[1], Dominik Mühlen[1], Gregor Bucher[1] and Vera

S. Hunnekuhl[1]*

* Author for correspondence

[1] University of Göttingen, Johann-Friedrich-Blumenbach Institute, GZMB, Department of

Evolutionary Developmental Genetics, Germany

[2] University Medical Center Göttingen (UMG), Germany

Abstract

Despite its conserved basic structure, the morphology of the insect brain and the timing of

its development underwent evolutionary adaptations. However, little is known about the

developmental processes that create this diversity. The central complex is a brain centre

required for multimodal information processing and an excellent model to understand

neural development and divergence. It is produced in large parts by type-II neuroblasts,

which produce intermediate progenitors, another type of cycling precursor, to increase their

neural progeny. These neural stem cells are believed to be conserved among insects, but

little is known on their molecular characteristics in insects other than flies. *Tribolium*

castaneum has emerged as a valuable new insect model for brain development and

evolution. However, type-II neuroblast lineages and their role in central complex

development have so far not been studied in this beetle.

Using CRISPR-Cas9 we created a fluorescent enhancer trap marking expression of *Tribolium*

fez/earmuff, a key marker for type-II neuroblast derived intermediate progenitors. Using

combinatorial labelling of further markers including *Tc-pointed*, *Tc-deadpan*, *Tc-asense* and

Tc-prospero we characterized the type-II neuroblast lineages present in the *Tribolium*

embryo and their sub-cell-types. Intriguingly, we found 9 type-II neuroblast lineages per

hemisphere in the *Tribolium* embryo while *Drosophila* produces only 8 per brain

hemisphere. In addition, these lineages are significantly larger at the embryonic stage of

Tribolium than they are in *Drosophila* and contain more intermediate progenitors. Finally,

we mapped these lineages to the domains of early expressed head patterning genes.

Notably, *Tc-otd* is absent from all type-II neuroblasts and intermediate progenitors, whereas

Tc-six3 marks an anterior subset of the type-II-lineages. The placodal marker *Tc-six4*

specifically marks the territory where anterior medial type-II neuroblasts differentiate.

In conclusion, we identified a conserved pattern of gene expression in holometabolan

central complex forming type-II neuroblast lineages, and conserved head patterning genes

emerged as new candidates for conferring spatial identity to individual lineages. The higher

number and greater lineage size of the embryonic type-II neuroblasts in the beetle correlate

with a previously described embryonic phase of central complex formation which is not

found in the fly. These findings stipulate further research on the causal link between timing

of stem cell activity and temporal and structural differences in central complex

development.

47
Introduction

Neural development of insects allows to study molecular and cellular principles in easy to

manipulate invertebrate models. The fruit fly *Drosophila melanogaster* (*Drosophila*

hereafter) has mostly been used for this purpose, making it one of the best understood

models for neurogenesis [1,2]. However, it has remained a puzzle how the huge ecological

diversity of insects and the divergent neural anatomies that are adapted to different niches

[3,4] evolved. Furthermore, *Drosophila* may in many instances not be a good representation

of insect development and some processes are derived in the fly lineage [5–7]. For these

reasons the beetle *Tribolium castaneum* (*Tribolium* hereafter) has been introduced as an

additional model for insect neural development [8–11]. *Tribolium* is a grain storage pest and

all stages live in flour, hence there is no major transition of habitat or lifestyle between the

larval and adult stage as in flies. However, when compared to *Drosophila* the larvae are

more mobile through the use of walking legs which allows them to navigate within or

between food sources [3,6,10]. In this beetle many molecular genetic manipulation and

labelling methods have been established [12–16]. Specifically, labelling of neural cell types

in *Tribolium* has been facilitated by the advent of CRISPR-Cas9 [12,15,17]. Another factor

that makes *Tribolium* an informative model for the understanding of insect brain

development is the more conserved development of the head neuroectoderm facilitating

the identification of specific neurogenic domains and allowing cross-species comparisons

[11,18,19].

The insect central complex is an anterior, midline spanning neuropile and constitutes an

important brain centre for the processing of sensory input, coordination of movement and

navigation [20,21]. The highly conserved basic structure of the central complex, along with

diverse specifications in response to ecological requirements among different insect species,

make this brain structure a very interesting model for evolutionary developmental studies

[3,10,20]. Between the two insect models *Drosophila* and *Tribolium* a temporal shift in the

emergence of central complex neuropile was observed with *Tribolium* developing a

functional larval central complex during embryogenesis whereas it develops only at the

onset of adult life in *Drosophila* [3,10]. It is however not known how these temporal

differences are established during development on a cellular and molecular level.

The entire insect central nervous system including the brain is produced by neural stem

cells, the neuroblasts (NBs) [22]. While there are evolutionary modifications in the relative

position of trunk NBs and their gene expression profiles between fly and beetle, the core

determinants specifying their role as neural progenitors are conserved [8]. NBs undergo

repeated divisions producing rows of ganglion mother cells (GMCs) which divide one more

time to produce neurons and/or glia (see Fig. 1, top panel). This leads to neural cell lineages

that include the neuroblast itself and its progeny [23,24]. Several NBs of the anterior-most

part of the neuroectoderm contribute to the central complex, which has been characterized

as the most complex brain structure (apart from the optic lobe) [25,26]. Intriguingly, in

Drosophila and in the grasshopper *Schistocerca gregaria* a neuroblast subtype, the type-II

neuroblasts (type-II NBs), were found to prominently contribute to the formation of the

central complex [27,28]. These specific neuroblasts generate more offspring by producing

another class of neural precursors, the intermediate progenitors (INPs), which also divide in

a stem cell like fashion [29,30] (Fig. 1, lower panel). Type-II NBs and INPs have attracted a lot

of attention because intermediate, cycling progenitors have also been described in

vertebrates, the radial glia cells, which produce a variety of cell types [31,32]. In addition, in

Drosophila brain tumours have been induced from type-II NB lineages [33], opening up the

possibility of modelling tumorigenesis in an invertebrate brain, thus making these lineages

one of the most intriguing stem cell model in invertebrates [34,35].

Figure 1) Neuroblasts of the _Drosophila_ nervous system. Type-I neuroblasts (type-I NBs)

(top panel) and type-II neuroblasts (type-II NBs) (lower panel). Type-I NBs undergo stem-

cell-like divisions with each round producing a ganglion mother cell (GMC) that divides one

more time to produce two neurons or glia cells. In _Drosophila_, all type-I NBs express _dpn_ and

ase whereas GMCs express only _ase_. Type-II NBs express _dpn_ and _pnt_ but not _ase_. They also

undergo repeated divisions, with each round producing an intermediate progenitor cell

(INP) which also is a proliferating progenitor and expresses _ase_. INPs undergo a maturation

process during which _pnt_ is still expressed initially, but not _dpn_. Mature INPs express _dpn_,

ase and _fez/erm_ whereas GMCs express _ase_ and _fez/erm_. Type-II NB lineages are only found

in the anterior brain, they produce central complex neurons and glia [30,36].

In the developing insect brain the number of neurons that is produced by each type-II NB is

significantly larger than the offspring of a type-I NB. Medial _Drosophila_ type-II NBs produce

an average of 450 neurons whereas type-I NBs of the brain only generate 100-200 neurons

each [37–39]. This increased number of neural cells that are produced by individual type-II

lineages along with the fact that neuroblast lineages can produce different types of neurons,

leads to the generation of extensive neural complexity within the anterior insect brain

[27,30].

Based on descriptions of large proliferative lineages in the grasshopper _Schistocerca_

gregaria, a hemimetabolous insect distantly related to flies, it is widely believed that type-II

NBs and INPs are conserved within insects [25,40]. However, a molecular characterisation of

such lineages that on the one hand will allow to test whether the cell type markers

characterized in _Drosophila_ are more widely conserved, and on the other hand will facilitate

an identification of sub-cell types, has not been performed in an insect other than

Drosophila.

Whereas it is widely believed that _Tribolium_ does use type-II NB in central complex

formation [3,17,41] their presence has so far not been unequivocally shown and a thorough

comparison of type-II NB lineages and their sub-cell-types between the fly and the beetle

model has been lacking.

The characterization of type-II NBs in _Drosophila_ has in large parts been based on a specific

reporter line in which eGFP expression is driven by an _earmuff (fez/erm)_ enhancer element

and marks INP-lineages [30,42,43]. _Tribolium earmuff_ has been first described as _Tc-fez_,

referring to the vertebrate *earmuff* ortholog *fez* [19]. We will use *Tc-fez/erm* in the

following. A defined sequence of gene expression has been described in *Drosophila* type-II

NBs, INPs and GMCs. The ETS-transcription factor *pointed* (*pnt*) marks type-II NBs [44,45],

which do not express the type-I NB marker *asense* (*ase*) but the neural gene deadpan (*dpn*)

(Fig. 1). Intermediate progenitors express *fez/erm* and *ase* and at a more mature stage also

dpn, whereas *prospero* (*pros*) marks mature INPs and GMCs [30,46]. *Six4* is another marker

for *Drosophila* type-II NBs and INPs [47].

In the present work we characterize type-II neuroblasts in *Tribolium* with respect to their

number, their location and the conservation of the molecular markers known from

Drosophila. We also wanted to test, in how far the earlier emergence of the central complex

in beetles would be reflected by a change of division activity of type-II NBs or intermediate

progenitors. To that end, we used CRISPR-Cas9 induced genome editing for creating a *Tc-*

fez/erm enhancer trap line to characterize embryonic *Tc-fez/erm* -expressing cells including

INPs and GMCs of type-II NB lineages and combined this line with other labelling methods

including multicolour *hybridisation chain reaction* (HCR) [48,49]. We found that the

Tribolium embryo produces 9 *Tc-pnt*-expressing type-II NBs compared to only 8 type-II NB

lineages in the *Drosophila* embryo [50]. We show that these lineages produce central

complex cells and confirm a largely conserved molecular code that differentiates INPs of

distinct maturation stages and GMCs. Intriguingly, we found that lineage sizes in *Tribolium*

embryos are considerably larger and include more mature INPs than in the *Drosophila*

embryo [50]. We also show that the placodal marker gene *Tc-six4* characterizes the

embryonic tissue that gives rise to the larger anterior group of type-II NBs, whereas *Tc-six3*

is marking only an anterior subset of these lineages. Interestingly there is a part of the

central complex precursors expressing neither *Tc-six3* nor *Tc-otd,* which is absent from all

type-II NB lineages.

Results

A CRISPR-Cas9-NHEJ generated enhancer trap lines marks *Tc-fez/erm*-expressing cells

at the embryonic and larval stage

We created an enhancer trap line driving eGFP that reflects *Tc-fez/erm* gene expression in

the embryo. We named the line *fez magic mushrooms* (fez-mm-eGFP) because it marks the

mushroom bodies of the larval brain (Fig. 2 A-C). The reporter construct was inserted 160 bp

upstream of the *fez/erm*-transcription start site using the *non-homologous end joining*
(NHEJ) repair mechanism (see supplementary Fig. S1 for scheme). Enhancer traps
sometimes only reflect parts of a gene expression pattern, therefore we carefully evaluated
eGFP co-expression with *Tc-fez/erm*-RNA. We found that at the embryonic stage fez-mm-
eGFP co-localized with most *Tc-fez/erm*-expressing cells in an area that will give rise to the
protocerebral structures mushroom bodies, larval eyes and central complex [11,19] (Fig. 2,
panel D). The fez-mm-eGFP line marked several progenitor cell types including cells that we
characterize as type-II NB-derived intermediate progenitors (see below). Based on these
analyses, we use the fez-mm-eGFP expression as a reporter for *Tc-fez/erm* expression.

Figure 2) Magic mushrooms (fez-mm-eGFP) reporter line. A-C) eGFP-expression in a larval
instar brain, D) EGFP antibody labelling in combination with *Tc-fez/erm* RNA *in situ* in the
embryonic head, A-D) Dapi labelling of nuclei. Scalebars in A and D are 50 μm. **A)** Full
projection of whole brain scan (4th instar larvae): fez-mm-eGPF expression marks many cells
of the mushroom bodies (white arrowheads) and of the larval optic lobe (purple
arrowheads). Note that we use the terms dorsal and ventral with respect to the neuraxis of
the larva. See [10] for orientation. **B)** Central part of the whole brain scan (projection of
substack, same brain as in A): mm-eGFP marks the peduncles of the mushroom bodies
(white arrowheads) and some central complex ensheathing cells (probably glia cells; yellow
arrowhead). **C)** Projection of the dorsal part of same brain as in A shows calyces and
peduncles of mushroom bodies (white arrowheads), and optic neuropile (purple

arrowheads). **Panel D)** Extensive co-localization of fez-mm-eGFP with *Tc-fez/erm*-RNA in

white encircled area which gives rise to protocerebral structures. Projection of all planes

where expression was detected, embryonic stage NS11.

***Tc-pointed* marks a population of nine type-II NBs that are associated with fez-mm-**

eGFP marked INP-lineages

To identify type-II NBs, we searched for cell groups that express the markers known from

flies. The transcription factor *pointed* (*pnt*) marks type-II NBs in *Drosophila* and is required

for their correct differentiation [44,45]. In both *Tribolium* and *Drosophila*, *pnt* is also

expressed in other areas (this work/[51,52]). Therefore, we looked for *Tc-pnt* expression in

large cells with large nuclei (typical NB morphology) that were closely associated with *Tc-*

fez/erm-expressing cells (marking putative INPs). We found a total of 9 such clusters instead

of the 8 expected from flies (Fig. 3 A-C). We also found that the large *Tc-pnt* expressing cells

as well as fez-mm-eGFP expressing cells are mitotically active (Fig. 3 C-I). Further molecular

and cell size analysis corroborated our interpretation that these cells constitute type-II NBs,

intermediate progenitors (INPs) at different maturation stages and ganglion mother cells

(GMCs) (see details below).

Looking at a series of embryonic stages we determined the emergence and further

development of *Tc-pnt* and fez-mm-eGFP expressing lineages as well as their arrangement

with respect to the embryonic head and to one another. We first detected conspicuous

adjacent expression of *Tc-pnt* and fez-mm-eGFP at stage NS7 (not shown). At stage NS11

seven *Tc-pnt*-clusters are present in a horse-shoe-shaped formation surrounding fez-mm-

eGFP positive cells (white arrows in Fig. 3 A, star marks one cluster out of focus). At stage

NS13 these groups are found in a position at the medial border of the head lobes and their

number has increased to nine. The seven clusters that were already present at stage NS11

are arranged in an anterior group (white arrowheads in Fig. 3B) at stage NS13 while two

additional clusters have emerged more posteriorly (yellow arrows in Fig. 3B-I). The clusters

are in different depths (reflected by the three optical sections shown in Fig. 3 B-I to B-III,

clusters out of focus are visualized by grey circles in Fig. 3 B'I-II, respectively). Fez-mm-eGFP

expression not related to type-II NB offspring was found in the lateral head lobe from NS13

onwards (blue arrowheads in Fig. 3). At stage NS14 six of the anterior-medial clusters are

aligned in one plane along the medial rim of the developing brain (Fig. 3 C-I), whereas the

most posterior cluster of the anterior group is located at a separate position and in a deeper

plane (Fig. 3C-II). The posterior group is found at about the same focal plane as well (Fig. 3

C-II). The aligned six clusters produce their *Tc-fez/erm*-positive offspring towards laterally

whereas the *Tc-fez/erm*-positive offspring of the more posterior cluster is found medially of

it (Fig. 3 C-II). The two most posterior clusters produce *Tc-fez/erm*-cells in an anterior-lateral

direction (Fig. 3 C-II). The apparent re-arrangement of the *Tc-pnt*-clusters from the anterior

rim of the embryonic head to a medial position reflects not active migration but follows the

overall tissue movements during head morphogenesis that was previously described in

Tribolium [19] (Fig. 3 D). Based on their position by the end of embryogenesis (stage NS14)

we assign the nine type-II NB clusters into two groups: one anterior median group consisting

of seven type-II NBs and one posterior group consisting of two type-II NBs (see Fig. 3 C and

D).

Figure 3) Developmental occurrence and progression of *Tc-pnt*-clusters and associated fez-mm-eGFP cells (type-II NB lineages). A-C) Anti-GFP antibody staining reflecting fez-mm-eGFP expression in combination with *Tc-pnt* RNA *in situ* hybridisation and dapi labelling of nuclei (A, C) or FM1-43 membrane staining (B), single planes from confocal z-stacks of right head lobes. Scalebars are 25 μm. A'-C') Schematic drawing of the head lobe at the respective stages; clusters out of focus depicted by grey circles. D) Schematic depiction of the localization of type-II NBs during *Tribolium* head morphogenesis. **A)** At stage NS11 seven *Tc-pnt*-clusters (white arrows) are found in an anterior-medial horseshoe-like arrangement with fez-mm-eGFP cells in the centre. Asterisk indicates the position of one *Tc-pnt*-cluster which is outside the focal plane of this image. **B)** At stage NS13 the seven anterior-medial clusters have been moved to the medial margin of the developing brain by

morphogenetic movements of the head lobes (white arrows). The clusters are situated in
different dorso-ventral levels shown in the different images B-I to B-III. Two additional
posterior *Tc-pnt*-clusters have appeared (yellow arrows). Blue arrows point to fez-mm-eGFP
positive cells that are not part of the type-II NB lineages. **B-I**: ventral level **B-II** mid-level, **B-
III**: dorsal level. **C)** At stage NS14 a group of six clusters are arranged in one plane anterior-
medially in the brain lobe (white arrows) (**C-I**). Anti-PH3 labelling marks mitotic cells within
the lineages (white signal) and clusters out of the focal plane are indicated by grey circles in
C'-I. **C-II** One cluster of the anterior median group (white arrow) is in a deeper plane, as well
as the two posterior clusters (yellow arrows). Blue arrowheads point to lateral fez-mm-eGFP
expression not associated with type-II NBs. A'-C') AMG =anterior medial group (blue), PG
=posterior group (pink). Grey circles in B'-C' indicate position of type-II NBs in different focal
planes. **D)** *Tc-pnt*-expressing cells identified as type-II NBs first appear bilaterally in the
anterior most part of the embryonic head (stage NS11). During head morphogenesis this
anterior tissue folds over so that the type-II NBs end up in a more medial position. Two
more type-II NBs emerge posteriorly (stage 13). At the end of embryogenesis (stage NS14)
the type-II NBs can be divided in an anterior medial group (AMG (blue); including type-II NBs
1-7, numbering starting from anterior) and a posterior group (PG (pink), including type-II
NBs 8-9).

Characterization of cell- and nuclear size of *pnt*-positive type II NBs
We wanted to confirm that each of the *Tc-pnt* and *Tc-fez/erm* expressing lineages contains a
neuroblast. Neuroblasts, including type-II NBs, are larger than other cells, have larger nuclei
and are mitotically active [40]. Therefore, we determined the cell size of the largest cells of
the *Tc-pnt* cluster and compared it to a random sample of cells of the embryonic head. Both
samples were taken from stage NS13 and NS14 embryos. We found that the cells that we
had assigned as Type-II NBs had a significantly larger diameter (av.: 12.98 µm; n=43) than
cells of the reference sample (av.: 6.75 µm; n=1579) (Fig. 4). We also found that the
diameters of nuclei of type-II NBs were significantly larger (av.: 8.73 µm; n=17) than the
ones of a control sample (av.: 5.21 µm; n=1080) (Fig. 4). Although type-II NBs were clearly
larger than the average cell of the head lobes the control sample also contained cells of an
equal size (figure 4) which may constitute type-I NBs.

Figure 4) Cell- and nuclear diameter of type-II NBs compared to a control group. Stages NS13-14. Left: diameter of *Tc-pnt*-pos. type-II NBs (average diameter=12.98 µm; n=43) compared to control group (average diameter=6.75, n=1579). Right: average nuclear diameter of type-II NBs (=8,73 µm, n=17) compared to average nuclear diameter of control group (=5.21, n=1080). Differences between groups are significant (t-test; *** ≅ P < 0.001).

Conserved patterns of gene expression mark *Tribolium* type-II NBs, different stages of INPs and GMCs

Drosophila type-II NBs, INPs and GMCs express a specific sequence of genes including *asense* (*ase*), *deadpan* (*dpn*) and *prospero* (*pros*) (Fig. 1) [30]. However, these markers have up to now not been tested for expression in type-II lineages in other organisms. We tested if the respective lineages express these markers in a comparable sequence in the beetle and if they mark distinctive cell types within the lineages (Fig. 5 panel A-D and Fig. panel 6 A, B). We found that the large *Tc-pnt*-positive but fez-mm-eGFP negative cells (i.e. type-II NBs) express the gene *Tc-dpn*, in line with expression of this gene in *Drosophila* neural precursors

(Fig. 5, panel A). Like *Drosophila* type-II NBs, these cells do not express the type-I NB marker

Tc-ase (yellow arrowhead in Fig. 5, panel D). We further found additional cells directly

adjacent of the type-II NBs itself which we believe are recently born immature INPs. They

are *Tc-pnt*-positive but fez-mm-eGFP-negative and neither express *Tc-ase* (Fig. panel D, pink

arrowheads). Expression of *Tc-ase* starts in the following in immature *Tc-pnt*-positive INPs

(Fig. 5 panel D; blue arrowhead) and is maintained in INPs that are *Tc-pnt*-negative but are

marked by fez-mm-eGFP (Fig. 5, panel D and E). *Tc-dpn* is absent from immature *Tc-pnt*-

positive INPs (Fig. 5D, pink arrowhead) but is expressed in mature INPs (orange arrowheads

in Fig. 5, panel B). Cells that are located at the distal end of the lineages that do not express

Tc-dpn but are positive for fez-mm-eGFP and *Tc-ase* are classified as GMCs (Fig. 5, panels B,

C and E). We found the gene *Tc-pros* expressed in most fez-mm-eGFP expressing cells of a

lineage (in Fig. 6, panel A-B) and infer from the extent of the expression that it is expressed

in mature INPs (*Tc-dpn*+) and GMCs (*Tc-dpn*-). Fez-mm-eGFP positive cells at the base of the

lineage that do not express *Tc-pros* are immature INPs (Fig. 6, panel A, B; blue arrowheads).

In summary, the expression dynamics found in the *Tribolium* type-II NB lineages are very

similar to the one found in *Drosophila.* Therefore these neural markers can be used for a

classification of cell types within the lineages into type-II NBs (*Tc-pnt+*, *Tc-fez/erm-*, *Tc-ase-*,

Tc-dpn+, *Tc-pros-*), immature-I INPs (*Tc-pnt+*, *Tc-fez/erm-*, *Tc-ase-*, *Tc-dpn-*, *Tc-pros-*),

immature-II INPs (*Tc-pnt+*, *Tc-fez/erm+*, *Tc-ase+*, *Tc-dpn-*, *Tc-pros-*), mature INPs (*Tc-pnt-*, *Tc-*

fez/erm+, *Tc-ase+*, *Tc-dpn+*, *Tc-pros+*), and GMCs (*Tc-pnt-*, *Tc-fez/erm+*, *Tc-ase+*, *Tc-dpn-*, *Tc-*

pros+). This classification is summarized in Fig. 7 A-B. Image stacks from specimens co-

stained with the anti-PH3 mitosis marker (data shown in fig. 3C and listed in table 1 and 2,

available at https://doi.org/10.25625/8IVICL) indicate that immature-I and -II INPs are not

dividing whereas mature INPs are.

Figure 5: Differential expression of the neural markers *Tc-pnt, Tc-fez/erm, Tc-dpn* and *Tc-ase* in type-II NBs, INPs and GMCs. A, B, D) Anti-GFP staining visualizing fez-mm-eGFP expression in combination with HCR-labelling of further factors. C, E) GFP antibody staining in combination with RNA *in situ* hybridisation. A-E) Dapi staining of nuclei. All panels are single planes from confocal z-stacks. **Panel A)** Stage NS12, expression of *Tc-pnt* and *Tc-dpn* in type-II NBs (yellow arrowhead) with adjacent fez-mm-eGFP positive INPs. **Panel B)** Stage NS13, white line marks one lineage. Expression of *Tc-dpn* is absent in the immature-I *Tc-pnt+* INPs (pink arrowhead) but present in mature INPs (orange arrowheads), white line marks one lineage. Type-II NB not in focus of this image. **Panel C)** Stage NS14, expression of

Tc-dpn in type-II NBs (yellow arrowheads), and mature INPs (orange arrowheads). Asterisk

marks mitosis in a fez-mm-eGFP cell visible through the condensation of chromatin stained

by dapi. **Panel D)** Stage NS13, *Tc-ase-* and *Tc-pnt*-expression in type-II NBs and INPs, white

line marks one lineage. Yellow arrowhead: type-II NBs (*Tc-pnt+*), pink arrowheads:

immature-I INP (*Tc-pnt+*), blue arrowhead: immature-II INP (*Tc-fez+/ Tc-ase+/ Tc-pnt+*).

Panel E) Stage NS13, expression of *Tc-ase* in fez-mm-eGFP positive cells, location of type-II

NBs and immature-I INPs is outside the focus.

332
333

Figure 6) Expression of *Tc-pros* in fez-mm-eGFP cells. A-B) Anti-GFP antibody staining in

combination with *Tc-pros* RNA *in situ* hybridization. Membranes stained with FM 1-43.

Single planes from confocal z-stacks. **Panel A)** Stage NS13, two type-II NB lineages producing

INPs and GMCs in opposing directions. Expression of *Tc-pros* in mature INPs and GMCs.

Cluster of type-II NBs (yellow arrowheads) and immature-I INPs (pink arrowheads).

Immature-II INPs express *Tc-fez/erm* but not *Tc-pros* (blue arrowhead). **Panel B)** Stage NS13,

detail of *Tc-pros*-negative type-II NBs (yellow arrowhead) and immature-I INPs (pink

arrowheads) and immature-II INP (blue arrowhead). Note that distinction between type-II

NB and immature INPs in **A)** and **B)** is only made based on position.

343
344
345
346

348
Figure 7) Schematic drawing of expression of different markers in a type-II NB lineage. A)

and B) show the same lineage with *Tc-fez/erm* and *Tc-pnt* expression but with different

additional markers mapped on top. **A)** Subclassification into type-II NBs (*Tc-pnt+*),

immature-I INPs (*Tc-pnt+*), immature-II INPs (*Tc-pnt+, Tc-fez/erm+, Tc-ase+*) and mature

INPs and GMCs (*Tc-fez/erm+, Tc-ase+*) (based on stainings shown in Fig. 5). **B)** Type-II NB (*Tc-*

pnt+, Tc-dpn+), immature-I (*Tc-pnt+*) and immature-II (Tc-pnt+, *Tc-fez/erm+*). Mature INPs

express *Tc-dpn*, *Tc-pros* and *Tc-fez/erm* whereas GMCs express only *Tc-fez/erm* and *Tc-pros*

(based on staining shown in Fig. 5 and 6). Note that of the selected markers immature-I INPs

express *Tc-pnt* only.

358
Type-II NB lineages produce central complex cells that are marked by shaking hands (skh)

To test if the *Tribolium* embryonic type-II NB lineages contribute to the beetle central

complex like in the fly, we used the shaking hands- (skh-) reporter line that marks central

complex neurons and their postmitotic embryonic precursors [41]. By co-staining for *Tc-pnt*

and *Tc-fez/erm* RNA in the background of the skh-line we found that the skh-positive cells

are located directly distally of the type-II NB lineages in a position where the type-II NBs

derived neural cells are expected (Fig. 8 panels A-B, E,F). Most skh-cells have switched off

fez/erm-expression, but at the transition between *Tc-fez/erm* and skh cells we found some

cells that express both markers (Fig. 8 panel D, E), supporting the view that many skh-

positive central complex cells stem from the *fez*-expressing GMCs of the type-II NB lineages

(Fig. 8 E). We found skh cells at the end of lineages of the anterior-median group (Fig. 8,

panel A) and at the end the two posterior lineages (Fig. 8, panel B). We therefore conclude

that both groups contribute to central complex neuropile (Fig. 8 F). We cannot say whether

the lineage of type-II NB seven, which we assigned to the anterior-median group, but which

is located with some distance to the other type-II NB of that group (Fig. 3 C, D; grey circle in

Fig. 8 F), produces skh+ cells, but we assume that it does.

Figure 8) Expression of the transgenic central complex reporter skh/eGFP in relation to

type-II NB lineages. A-D) Antibody labelling of eGFP, which is expressed in the pattern of

skh [41] in combination with HCR visualizing *Tc-fez/erm* and *Tc-pnt*, dapi labelling of nuclei, single planes of confocal z-stacks, all stage NS14. **A)** Left head lobe, anterior group of *Tc-pnt*-positive type-II NBs, *Tc-fez/erm*-expressing INPs/GMCs and skh/eGFP positive postmitotic cells. **B)** Left head lobe, posterior group of type-II NB lineages. *Tc-pnt+* type-II NBs or immature-I/-II INPs (white arrows), *Tc-fez/erm+* INPs/GMCs (orange arrowheads) and skh/eGFPpostmitotic cells (yellow arrowheads). **C)** Detail of one lineage including type-II NB cluster (*Tc-pnt+*, white arrows), INPs/GMCs (*fez+*, orange arrowheads) and skh/eGFPcells at the end of the lineage (yellow arrowheads). **D)** Detail of transition from *Tc-fez/erm+* cells (orange arrowheads) to skh/eGFPcells reveals a small area of co-expression of both factors (yellow arrowhead). **E)** Schematic drawing of type-II NB lineage showing relative positions and markers of type-II NBs, INPs/GMCs and postmitotic central complex forming cells. **F)** Schematic overview of the arrangement of the type-II NB lineages including skh/eGFP+ central complex forming cells in the head lobe of stage NS14 (immature-I/-II INPs not shown for simplicity). No data on skh-GFP expression in lineage of type-II NB 7 (marked in grey) is available.

The *Tribolium* embryonic lineages of type-II NBs are larger and contain more mature INPs than those of *Drosophila*

In beetles, a single-unit central body (most likely constituting the fan-shaped body [10]) develops during embryogenesis and has gained functionality at the onset of larval life [15]. By contrast, in *Drosophila* central complex commissural tracts originating from type-II NBs form in the embryo but the type-II NB become quiescent and only resume divisions in the late larval stage [29,50]. In stark contrast to *Tribolium* a functional central complex is only observed in the adult fly [3]. In addition, the volume of the L1 central brain including the central complex is about four times larger in *Tribolium* than it is in *Drosophila* [53], while there are only small differences in the relative neuropilar volumes of the adult beetle and fly central bodies [53,54]. As type-II NBs contribute to central complex development we asked in how far the embryonic division pattern of these lineages in the beetle would reflect this heterochronic development. To assess the size of the embryonic type-II NB lineages in beetles we counted the *Tc-fez/erm* positive (fez-mm-eGFP) cells (INPs and GMCs) of the anterior medial group (type-II NB lineages 1-7). As we were not always able to distinguish between cells belonging to neighbouring INP-lineages we quantified and averaged the cell

411   number of all lineages of the anterior medial group. We also evaluated the proportion of

412   dividing cells in that group based on anti-PH3 staining (table 1).

414   **Table 1: *Tc-fez/erm* positive cell number (type-II NBs, INPs and GMCs)**

Embryo#/stage	Total number of cells of 7 anterior lineages	% dividing cells (PH3 antibody)	Cells per lineage (average)
1/NS 13	146	8.2%	20.9
2/NS 13	111	6.3%	15.9
3/NS 14	139	5.0%	19.9
4/NS 14	96	10.5%	13.7

We found that at stage NS13 (early retracting embryo) each lineage consisted on average of

18,4 progenitor cells and at stage NS14 (retracted embryo) of 16.8 progenitors. Between 5-

10 % of the cells were marked by the PH3 antibody. As PH3 marks dividing cells in a specific

phase, the portion of dividing cells may be higher. We further quantified the number of

mature *Tc-dpn*-positive INPs (table 2).

Table 2: Mature INPs (*Tc-fez/erm* and *Tc-dpn* positive)

Embryo #/ stage	Total number of mature INPs of 7 anterior lineages	% dividing cells (PH3 antibody)	Number of mature INPs per lineage (average)
1/NS 13	24	n.d.	3,4
2/NS 13	12	n.d.	1,7
3/NS 13	23	n.d.	3,3
4/NS 14	27	3,7%	3,9
5/NS 14	30	13,3%	4,3

We used the numbers from *Tribolium* (table 1, 2) for comparison with type-II NB lineages

from *Drosophila* embryos (data from [29]). At fly embryonic stage 13, which is comparable

to *Tribolium* stage NS14 in that is represents a retracted germband, lineages only contained

one or more INPs but no GMCs [29]. Therefore, we compared our data for the anterior

lineages of *Tribolium* stages NS13 and NS14 to the anterior and median cluster in the

Drosophila stage 15 and 16 embryos. In both species these stages are the two penultimate

stages of embryogenesis when all type-II NB lineages are present, and importantly, in

Drosophila the maximum lineage sizes and numbers of INPs are reached at these stages. We

found that the beetle lineages were significantly larger than the corresponding ones in

Drosophila (Fig. 9; *Drosophila* data taken from [29]). Of note, cell counts in *Drosophila* also

included neurons [29]. Our counts are based on *Tc-pnt* and fez-mm-eGFP expression and

therefore mostly included progenitors. However, based on the small area of co-expression

of *Tc-fez/erm* and skh-expression (see above/ figure 8) they may also include some newly

born neurons, but the counts in *Drosophila* included more mature neurons which further

increases the difference in the number of progenitor cells between the two species and the

observed difference in lineage size represents a minimum estimate.

Due to a lower quality of the image data from the posterior group which is in a deeper layer

(see Fig. 3) we were not able to quantify cells of this cluster in *Tribolium* . In *Drosophila,* the

posterior lineages are larger (av. 10 cells [29]) than the anterior and median ones, but they

are still smaller than the anterior lineages of *Tribolium* at the stages that were investigated.

We also found that the embryonic type-II NB lineages in *Tribolium* comprised significantly

more mature INPs (*Tc-dpn+/ Tc-fez+*) than the corresponding lineages in *Drosophila* (Fig. 9).

The described observations on the higher numbers of progenitor cells in *Tribolium* when

compared to *Drosophila* correlate with a previously described earlier (i.e. embryonic)

formation of a functional central complex in beetles, which is present in the first larval stage

[3].

Figure 9) Size comparison of *Tribolium* and *Drosophila* type-II NB lineages. Anteriormedian lineages from *Tribolium* stages 13 and 14 (quantified in this work, see table 1 and 2) compared to *Drosophila* stages 15 and 16 (pooled data from [29]). Left: Lineage sizes comprising type-II NBs, INPs and GMCs (*Tribolium* n=4; *Drosophila* n=8). Right: Number of *Tc-dpn*-positive mature INPs (*Tribolium* n=5; *Drosophila* n=8). T-test p-value significance level (t-test; *** ≅ P < 0.001, ** ≅ P < 0.01).

Type-II NBs and their lineages are differentially marked by the head patterning transcription factors *Tc-six4* and *Tc-six3* but do not express *Tc-otd*

The anterior-most embryonic insect head is divided into different territories by highly conserved developmental transcription factors that give rise to different parts of the protocerebrum [11]. Hence, we wondered in which of these molecular territories subsets of type-II NB lineages are located, as neuroectoderm regionalization genes are candidates for giving specific spatial identities to individual lineages. We tested co-expression with the previously characterized head patterning transcription factors *Tc-otd*, *Tc-six4*, *Tc-six3*, known to be active in the developing protocerebrum (*Tc-otd*) and in the anterior medial region of the neuroectoderm, where type-II NBs emerge (*Tc-six3*, *Tc-six4*) [11,19,55,56]. We found no co-expression of fez-mm-eGFP marked cells of both the anterior medial and posterior lineages with *Tc-otd*. Rather, *Tc-otd* was expressed in the surrounding embryonic head tissue. *Tc-otd* was also absent from the type-II NBs themselves (Fig. 10 panel A-B). However, *Tc-otd* might be expressed in neural cells derived from these lineages as we can detect expression in the central area of the brain lobes where for instance skh cells are located (Fig. 8 panel A, F; Fig. 10 panel A-B). Interestingly, we found that *Tc-six4* is expressed specifically in that area of the head lobe in which the type-II NBs clusters 1-6 of the anterior medial group first differentiate. At stage NS13 *Tc-six4* marks both, type-II NBs and fez-mm-eGFP expressing INPs and some surrounding cells (Fig. 11, panel A). At the following differentiation stage (NS14) *Tc-six4* still marks the type-II NBs clusters of the anterior medial group but is not expressed in the fez-mm-eGFP INPs (Fig. 11 panel B). It is also not expressed in the posterior group of type-II NBs but only marks type-II NBs of the anterior medial group. Finally, we found that *Tc-six3* marks, and is restricted to, the anterior-most lineages 1-4 including both the type-II NBs and the fez-mm-eGFP marked INPs/GMCs (Fig. 11 panel C). These results reveal candidates for distinguishing type-II identity from type-I NBs in

the brain. Only the latter is possibly marked by *Tc-otd* as *Drosophila otd* is expressed in type-I brain neuroblasts [57]. By contrast, *Tc-six4* only marks type-II NBs and appears to be an early determinant of the anterior medial group. *Tc-six4* is also a candidate for distinguishing the anterior medial from the posterior cluster. The results also reveal a subdivision of the anterior cluster by *Tc-six3*, which marks the four most anterior type-II NBs.

Figure 10) Expression of *Tc-otd* in relation to the fez-mm-eGFP lineages. A/B) RNA *in situ* hybridisation for *Tc-otd* in combination with GFP antibody staining and dapi labelling of nuclei, A and B show projections of different levels of NS14 embryos, right head lobe. **A)** *Tc-otd* is neither expressed in type-II NBs nor the in the fez-mm-eGFP positive INPs of the anterior medial group (both positioned within encircled area) but it is expressed in the directly surrounding tissue. **B)** Likewise, *Tc-otd* is not expressed in type-II NBs or fez-mm-eGFP cells of the posterior group (encircled area). A lateral area of *Tc-fez/erm* and *Tc-otd* co-expression (**A/B**, blue arrow) likely is part of the eye anlagen [19].

Figure 11) Expression of *Tc-six4* and *Tc-six3* in type-II NBs and INPs. A-C) RNA *in situ*

hybridisations of *Tc-six4* and *Tc-six3* in combination with GFP antibody staining reflecting

fez-mm-eGFP expression, dapi labelling of nuclei, left head lobes shown. **Panel A)** Stage

NS13; *Tc-six4* is expressed in an area encompassing type-II NBs (positioned in encircled area)

and mm-eGFP expressing INPs. **Panel B)** Stage NS14; *Tc-six4* is now restricted to type-II NBs

and their youngest progeny that do not express fez-mm-eGFP (yellow arrowheads). **Panel C)**

Stage NS14; *Tc-six3* specifically marks the four anterior-most lineages. It is expressed in the

type-II NBs at the base of the lineages (yellow arrowheads) and in the fez-mm-eGFP positive

INPs and possibly also in GMCs. It is absent from the remaining lineages of the anterior

medial group and is also not seen in the posterior group (not shown).

Discussion

Evolutionary divergence of number and grouping of embryonic type-II NB lineages between beetle, fly and grasshopper

We have identified a total of 9 type-II NB lineages on each side of the *Tribolium* embryonic head. This represents a major evolutionary divergence as in both, the hemimetabolan grasshopper *Schistocerca gregaria* and the fruit fly *Drosophila melanogaster* only 8 embryonic type-II NBs were identified [29,38]. Our finding of an additional type-II NB lineage at the embryonic stage shows that the number of type-II NBs is not fixed across insects and suggests that evolutionary divergence in brain development between species may be driven by differences in the number of progenitor lineages. To test this, further research is required to determine where the additional lineage in *Tribolium* is located and if it is associated with morphological differences of the brain.

Although in the frame of this work we did not perform lineage tracing we can speculate about which NB might be lacking in fly embryos based on our data. In the hemimetabolan *Schistocerca* there is no obvious clustering of type-II NBs, and they are all arranged in one row at the medial rim of the head lobes. In *Tribolium* we have observed an arrangement of the type-II NBs into two groups per side, one large anterior medial group containing seven type-II NB clusters and one posterior group of two type-II NBs. The anterior six of the anterior medial group are arranged in one row like in the grasshopper. In *Drosophila* there are three described groups referred to as the anterior, the middle and the posterior cluster [29]. The posterior cluster consists of two type-II NBs and therefore most likely corresponds to the posterior two type-II NBs in *Tribolium*. The *Drosophila* middle and anterior cluster are most likely equivalent to the anterior medial group of *Tribolium.* However, the type-II NB 7, which we assigned to the anterior medial group, but which is a bit separated and produces offspring into the opposite direction (see Fig. 2 C-II, white arrow) might be the one that does not have a homologue in the fly embryo. The identification of more specific spatial markers for type-II NBs (in addition to *Tc-six3* and *Tc-six4*) or lineage tracing tools are required to identify the neuroblast, which is not present in fly embryos. Subsequently, it would be especially interesting to see what the role of the additional type-II NBs in the beetle is and which neuropile it contributes to.

Summing up, there is a tendency towards grouping of type-II NBs into subsets in the holometabolan models that was not observed in the only hemimetabolan studied so far. A ninth type-II NB was only found in the *Tribolium* embryo represents a striking

developmentally divergent feature, but its role and the homologisation of individual type-II

NBs between the different insects requires further studies.

Gene expression identifies homologous cell types and suggests conservation of gene

function between fly and beetle type-II NB lineages

The genes defining the different stages of differentiation in type-II NB lineages have been

identified and intensively studied in *Drosophila,* but it had remained unclear, in how far the

respective patterns and mechanisms were conserved. Our results show a large degree of

conservation of expression and probably also function – at least in holometabola. We

identified type-II NBs by their larger than average size [40] and the expression of the

signature marker *Tc-pnt* adjacent to a group of *Tc-fez/erm* expressing cells (INPs and GMCs).

In *Drosophila* both these factors have key functions in defining the developmental potential

of type-II NBs and INPs: Pnt suppresses Ase in type-II NBs and promotes the formation of

INPs [45]. Loss of *pnt* expression leads to a de-differentiation of INPs as *fez/erm* is no longer

repressed in young INPs [44]. Fez/Erm controls proliferation of INPs by activating Pros and

prevents dedifferentiation of INPs into type-II NBs [43]. We show that also in *Tribolium* INPs

undergo a maturation process, during which expression of *Tc-pnt* stalls and expression of

Tc-fez/erm is switched on. Despite both factors characterising different stages of INP

maturation they are not completely mutual exclusive and we observed a small window of

overlap in immature-II INPs, highly suggestive of a conserved role of Pnt in the activation of

fez/erm [44] (Fig. 12). *Drosophila dpn* is a neural marker [58] expressed in type-II NBs and

again in mature INPs, leaving a gap of expression in immature-I and II INPs. *Drosophila ase* is

repressed in type-II NBs but expressed together with *fez*/*erm* in INPs and GMCs where its

expression overlaps with *pros* [59]. We found identical dynamics of gene expression in

Tribolium (see Fig. 12) suggesting conservation of gene function within the lineages and a

conserved process of INP maturation.

In conclusion, expression of the genes examined in this study in type-II NBs and their

lineages is highly conserved in fly and beetle, which is implicit of conserved gene function

and a conserved process of central complex formation.

Figure 12) Conserved aspects of gene expression in type-II NB lineages of the
holometabolan models *Drosophila* and *Tribolium*. Overview on the different cell types
within one lineage (type-II NBs, INPs, GMCs and central complex neurons), their mitotic
activity, and summary of the expression of key factors in the respective cell types. **I**NPs are
divided into immature-I, immature-II, and mature INPs with each subtype having a unique
expression profile. Note that each lineage can have several mature INPs. We strongly
assume that glia are also produced from these lineages in *Tribolium*, as they are in
Drosophila [30]. * In the *Tribolium* embryo *Tc-six4* is only expressed in the anterior medial
lineages, and its expression is overlapping with *Tc-fez/erm* at an earlier stage (NS12)
whereas it is restricted to type-II NBs and immature-I INPs, mutually exclusive with *Tc-*
fez/erm, at a later embryonic stage (NS14). *Drosophila six4* was found in all 8 larval lineages
[47]. The scheme is based on this work and [30,41,43,44,47]. NB= neuroblast; imm.-I/-II INP
= immature-I/-II intermediate progenitor; mat. INP= mature intermediate progenitor;
N=neuron; skh=shaking hands (enhancer trap) [41]*.

Divergent timing of type-II NB activity and heterochronic development of the central
complex
Previous work described a heterochronic shift in central complex formation between fly and
beetle [3,10]. In *Tribolium* parts of the central complex develop during embryogenesis. It
becomes functional at the onset of the first larval stage, and is presumably required for the

599	more complex movements of the beetle larvae using legs [3]. In both *Drosophila* and
600	*Tribolium* the anterior groups of type-II NBs (anterior medial group of *Tribolium* and anterior
601	and middle cluster of *Drosophila*) are fully present in the last 3rd of embryogenesis, with the
602	beginning of germ band retraction [8,29,60]. However, in the *Drosophila* embryo only a
603	limited number of INPs and GMCs are produced before they enter a resting stage prior to
604	hatching [29]. By contrast, we show that *Tribolium* type-II NB lineages are much larger at the
605	embryonic stage compared to *Drosophila* [29]. They also include more mature, *Tc-dpn*
606	expressing cycling INPs which have the capability of driving the increase in lineage size. We
607	believe that this increased activity may contribute to the embryonic development of a
608	functional central body and protocerebral bridge [3]. This hypothesis does however require
609	further confirmation by lineage tracing experiments and by testing how manipulating
610	central genes acting within the lineages will affect the timing of central complex
611	development. In addition, differences in cell number in the individual central complex
612	neuropiles between *Tribolium* and *Drosophila* have to be assessed, as the higher number of
613	progenitors may also reflect a higher number of neurons present in the adult beetle fan
614	shaped body or may be due to a slower rate of division and a resulting accumulation of
615	progenitor cells. Lastly, we do not yet have any information whether *Tribolium* type-II NBs,
616	or *Tribolium* neuroblasts in general, also enter a stage of quiescence at the end of
617	embryogenesis as they do in *Drosophila* [29] and what role neurons born in the 1st larva play
618	in *Tribolium*.
619	Another emerging question is if and when *Tribolium* type-II NB lineages are active during
620	larval development and how that relates to the described steps of central complex
621	development. After preformation in the embryo *Drosophila* type-II NBs have their main
622	period of activity in the 3rd larva and the lineages are much larger and contain more INPs
623	than the *Tribolium* embryonic lineages [30]. Therefore, it would be interesting to see if type-
624	II NB lineages are present and active in the late larval brain of *Tribolium*.
625	Taken together, we have for the first time identified and molecularly characterized
626	embryonic type-II NBs and INPs as the major central complex progenitor pool in an insect
627	outside *Drosophila*. We have observed quantitative differences within these lineages when
628	compared between *Tribolium* and *Drosophila*. Based on these findings we hypothesize that
629	increased division behaviour of type-II NB lineages is required for the embryonic central
630	complex development as found in beetles. Our findings set the basis for future work on the

631 relationship of the identified type-II NB lineages and the timing of central complex

632 development.

The placodal marker *Tc-six4* may be responsible for the differentiation and the spatial

identity of the anterior medial group of type-II NBs

The different type-II NB lineages contribute to different parts of the brain. For instance, only

the anterior four type-II NB lineages form the w, x, y, and z tracts of the central complex in

flies and grasshoppers. Hence, there must be signals that make them different despite the

conserved and well-studied sequence of neural gene expression and functional interactions

in *Drosophila* (Fig. 12) [30,43,44,59]. In the ventral nerve cord, the different identities of the

NBs are specified by the combination of spatial patterning genes expressed in the

neuroectoderm at the time of delamination [24]. However, little is known about these

signals for type-II NBs. In *Tribolium* the domains of head patterning genes in the brain

neuroectoderm are well characterized and some highly conserved territories like the

anterior-medial *six3*-positive domain were described and suggested to give respective NBs

different spatial identities [11,19,55]. The *Tc-six4* expression domain is also largely

coincident with an embryonic structure termed insect head placode, a neurogenic,

invaginating tissue [56,61] potentially homologous to the vertebrate adenohypophyseal

placode which is also marked by *six4* [11,56]. A placodal origin of *Drosophila* type-II NBs has

been shown for one type-II NB, and it is assumed that the other type-II NBs also stem from

placodal tissue [29,50,62].

Tc-six4 is also a very interesting factor with regards to the specification of type-II NBs

because it is expressed only in the anterior group of type-II NBs. Hence, it could contribute

to distinguishing their developmental fate from the posterior group. This is however

different in the *Drosophila* larva*.* Here Six4 is expressed in all eight lineages where it

prevents the formation of supernumerary type-II NBs and a premature differentiation of

INPs [47]. In the *Tribolium* embryo Six4 may have a similar rate limiting role within the

anterior group of type-II NBs. However, its early expression in a small part of the

neuroectoderm [56] and the delamination of the anterior-medial type-II NB group within

this domain also hints at an instructive role of *Tc-six4* in the formation of the anterior medial

group.

Tc-six3 marks a subset of type-II NBs whereas _Tc-otd_ is absent from all lineages

Importantly, we found that in _Tribolium_ late embryogenesis, _Tc-six3_ is expressed specifically

within the lineages of type-II NBs 1-4 of the anterior medial group, but not in the other

linages. In grasshopper and fly these anterior 4 lineages (DM1-4) give rise to the z, y, x and

w tracts [25,38] and contribute crucially to the development of protocerebral bridge and

central body. These tracts form a major midline crossing neuropile of the central body which

serves as a scaffold in the development of this structure [3,25]. We conclude that in

Tribolium the evolutionary ancient _six3_ territory gives rise to the neuropile of the z, y, x and

w tracts. This might well be the same in other insects such as flies and grasshoppers but has

to our knowledge not been looked at in detail in these models. In the beetle the central

body is also missing in weak _Tribolium Tc-six3_-RNAi phenotypes, suggesting that _Tribolium_

six3 is required for the formation of this structure [19]. Given that central complex neuropile

is marked by additional factors such as _Tc_-foxQ2 (_Drosophila_ fd102c [63]) and _Tc-rx_- [3,17], it

would be interesting to see the relationship of additional spatial patterning genes with

regards to the type-II NB lineages in future studies.

The gene _otd_ is a marker of the posterior protocerebrum and _six3_ and _otd_ are believed to

subdivide the embryonic anterior brain into two major domains [55]. This subdivision is also

part of the recently revisited concept of an ancestral division of the insect protocerebrum

into archicerebrum (_six3_-positive) and prosocerebrum (_otd_-positive) [11]. As discussed

above, _Tc-six3_ is expressed in the anterior-most four type-II NB lineages, but unexpectedly,

we found that _Tc-otd_ is specifically absent from all type-II NB lineages, suggesting that its

inhibition is required for the development of all these lineages. Interestingly, the lineages of

type-II NB 5 and 6 (and presumably of type-II NB 7) of the anterior cluster and the posterior

type-II NB lineages do neither express _Tc-otd_ nor _Tc-six3_ showing that there are

protocerebral structures expressing none of these conserved markers in development,

questioning the assigned generalized role of these two factors in subdividing the

protocerebrum [11,55] .

In summary, our findings are just the beginning of the quest for the genes required for the

identity specification of type-II NB lineages. The genes known to be involved in head

patterning are an excellent starting point for this purpose [11,18].

Methods

Creation of transgenic reporter line

Using CRISPR-Cas9 we have inserted a transgene into the *Tc-fez/erm* locus that includes *egfp* behind the *Tribolium* basal heat shock promoter (bhsp68), which is not active by itself but can be activated by nearby enhancer elements [64,65]. The transgene also contained the coding sequence of Cre-recombinase (not used in this work) transcribed from the same promoter, and the eye pigmentation gene *Tc-vermilion* behind the eye-specific 3xP3 enhancer element [66] (see Fig. S1 A for map of the transgene). On the repair plasmid the transgene was flanked by CRISPR target sequences derived from *Dm-yellow* and *Dm-ebony*, which were used for *in vivo* linearisation of the transgene.

Using CRISPR Optimal Target Finder [67] we designed three guide RNAs (gRNAs) targeting the genomic region 2.5 kb upstream of the *Tc-fez/erm* transcription start site (see table S1). We produced three plasmids where single gRNAs were transcribed from the *Tribolium* pU6b RNA promoter, as described in [12]. We co-injected the three gRNAs plasmids at 125µg/µl each, gRNAs targeting the *Dm-yellow* and *Dm-ebony* sequence at 125µg/µl each, Cas9 helper plasmid (see [12]) at 500 µg/µl and the repair plasmid at 500 µg/µl. Embryos of the white eyed *Tribolium vermilion-white* (*vw*) strain (G0) were injected, raised and then bred to *vw*-beetles in individual crosses. The offspring of these crosses (F1) were screened for black eyed heterozygous carriers of the transgene. These were again outcrossed to *vw*-beetles and the heterozygous offspring (F2) were further analysed genetically (see table S2) and with respect to fluorescent signal. Siblings of the F2 generation were then crossed to one another to generate the homozygous line (fez-mm-eGFP). Details of the insertion can be found in Fig. S1 A.

We also used the shaking hands (skh) enhancer trap line (G10011-GFP) which marks central complex cells [41].

Larval brain dissection, fixation, and staining

Larval brains of the newly generated *Tc-fez/erm* enhancer trap line fez-mm-eGFP were dissected and stained as described in [16]. Brains were mounted in *Vectashield* for microscopic inspection. See table S4 for a list of antibodies and staining reagents used.

Embryonic RNA *in situ* hybridisation, hybridisation chain reaction and antibody staining

Gene identifiers and primer sequences that were used for amplification can be found in

table S3. The fragments were inserted into a pJet1.2 cloning vector. Standard RNA *in situ*

probes were synthesised using the 5X Megascript T7 kit (Ambion) according to the

manufacturers protocol. Embryo fixation and single colour RNA *in situ* chain reaction in

combination with antibody staining to eGFP or phospo-histone H3 staining to mark mitoses

was conducted as described in [68]. Embryos were also stained for DNA (dapi) or cell

membranes (FM1-43). A full list of antibodies and staining reagents can be found in table S4.

Probes for multicolour *hybridisation chain reaction (HCR)* binding *Tc-fez/erm*, *Tc-dpn*, *Tc-ase*

and *Tc-pnt* were produced by Molecular Instruments (see table S5). Labelling reactions were

performed as described in [69]. Antibody staining to eGFP was performed following the

completion of the HCR staining (see [68]). All embryos were mounted in *Vectashield* for

microscopic inspection (see table S4).

Image acquisition and analysis

Multichannel image stacks were recorded using a Zeiss LSM 980 confocal laser scanning

microscope. Image stacks consisted of 100-300 slices, depending on the specimen. The

resolution ranged from 1024 x 1024 to 2048 x 2048 pixels. Plane thickness was optimized

according to width of the pinhole and ranged from 0.1 – 1 µm. We used FIJI [70] for 3

dimensional inspection, to export individual planes or to generate maximum intensity

projections. Cropping and adjustment of brightness and contrast was done with GIMP

(version 2.10.32) or Adobe Photoshop CS5.

Evaluation and of cell size and cell number

Cell and nuclear diameters of type-II NBs and a control group were measured based on the

image stacks using FIJI [70] across 7 embryos. Lineage size and numbers of mitoses (based

on *Tc-pnt* and *fez*-mm-eGFP expression/ anti-PH3 staining) was evaluated by manual

counting based on image stacks derived from 4 head lobes of 4 embryos ($\cong$ 28 anterior-

medial lineages). The number of *Tc-dpn* expressing intermediate progenitors was evaluated

in the same way across 5 embryos ($\cong$ 35 anterior-medial lineages). Reference data on

Drosophila lineage size (type-II NBs, INPs, GMCs and neurons) and numbers of *dpn*-

expressing intermediate INPs were extracted from [29]. Statistics on all numerical data

(standard deviations and two-tailed t-tests) were performed using Microsoft Excel 2016.

References

1. Konstantinides N, Kapuralin K, Fadil C, Barboza L, Satija R, Desplan C. Phenotypic
Convergence: Distinct Transcription Factors Regulate Common Terminal Features. Cell.
Elsevier Inc.; 2018;174:622-635.e13. https://doi.org/10.1016/j.cell.2018.05.021
2. Li X, Chen Z, Desplan C. Temporal Patterning of Neural Progenitors in Drosophila. Curr Top
Dev Biol. 2014;8:1385–95. https://doi.org/10.1016/B978-0-12-396968-2.00003-8
3. Farnworth MS, Eckermann KN, Bucher G. Sequence heterochrony led to a gain of
functionality in an immature stage of the central complex: A fly–beetle insight [Internet].
PLoS Biol. 2020. http://dx.doi.org/10.1371/journal.pbio.3000881
4. Couto A, Wainwright JB, Morris BJ, Montgomery SH. Butterfly brain evolution. Curr Opin
Insect Sci. Elsevier Inc; 2020;42:55–60. https://doi.org/10.1016/j.cois.2020.09.002
5. Hakeemi MS, Ansari S, Teuscher M, Weißkopf M, Großmann D, Kessel T, et al. Screens in
fly and beetle reveal vastly divergent gene sets required for developmental processes. BMC
Biol. BMC Biology; 2022;20:1–13. https://doi.org/10.1186/s12915-022-01231-4
6. Klingler M, Bucher G. The red flour beetle T. castaneum: elaborate genetic toolkit and
unbiased large scale RNAi screening to study insect biology and evolution. Evodevo. BioMed
Central; 2022;13:1–11. https://doi.org/10.1186/s13227-022-00201-9
7. Tautz D, Friedrich M, Schröder R. Insect embryogenesis - What is ancestral and what is
derived? Development. 1994;120:193–9.
https://doi.org/10.1242/dev.1994.Supplement.193
8. Biffar L, Stollewerk A. Conservation and evolutionary modifications of neuroblast
expression patterns in insects. Dev Biol. Elsevier; 2014;388:103–16.
http://dx.doi.org/10.1016/j.ydbio.2014.01.028
9. Biffar L, Stollewerk A. Evolutionary variations in the expression of dorso-ventral
patterning genes and the conservation of pioneer neurons in Tribolium castaneum. Dev Biol.
Elsevier; 2015;400:159–67. http://dx.doi.org/10.1016/j.ydbio.2015.01.025
10. Koniszewski N, Kollmann M, Bigham M, Farnworth M, He B, Büscher M, et al. The insect
central complex as model for heterochronic brain development—background, concepts, and
tools. Dev Genes Evol. Development Genes and Evolution; 2016;226:209–19.
http://dx.doi.org/10.1007/s00427-016-0542-7
11. Posnien N, Hunnekuhl VS, Bucher G. Gene expression mapping of the neuroectoderm
across phyla – conservation and divergence of early brain anlagen between insects and
vertebrates. Elife. 2023;12:1–30. https://doi.org/10.7554/eLife.92242
12. Gilles AF, Schinko JB, Averof M. Efficient CRISPR-mediated gene targeting and transgene
replacement in the beetle Tribolium castaneum. Development. 2015;142:2832–9.
http://dev.biologists.org/cgi/doi/10.1242/dev.125054
13. Gilles AF, Schinko JB, Schacht MI, Enjolras C, Averof M. Clonal analysis by tunable
CRISPR-mediated excision. Dev. 2019;146:1–8. 10.1242/dev.170969
14. Posnien N, Schinko J, Grossmann D, Shippy TD, Konopova B, Bucher G. RNAi in the red
flour beetle (Tribolium). Cold Spring Harb Protoc. 2009;4:1–7. doi:10.1101/pdb.prot5256
15. Farnworth MS, Eckermann KN, Ahmed HM, Mühlen DS, He B, Bucher G. The red flour
beetle as model for comparative neural development: Genome editing to mark neural cells
in Tribolium brain development. Methods Brain Dev. 2020. https://doi.org/10.1007/978-1-
4939-9732-9_11

16. Hunnekuhl VS, Siemanowski J, Farnworth MS, He B, Bucher G. Immunohistochemistry
and Fluorescent Whole Mount RNA In Situ Hybridization in Larval and Adult Brains of
Tribolium. Methods Mol Biol. 2020. p. 233–51. https://doi.org/10.1007/978-1-4939-9732-
9_13
17. He B, Buescher M, Farnworth MS, Strobl F, Stelzer E, Koniszewski NDB, et al. An
ancestral apical brain region contributes to the central complex under the control of foxQ2
in the beetle tribolium. Elife. 2019;8:1–29. https://doi.org/10.7554/eLife.49065
18. Posnien N, Schinko JB, Kittelmann S, Bucher G. Genetics, development and composition
of the insect head - A beetle's view. Arthropod Struct Dev. Elsevier Ltd; 2010;39:399–410.
http://dx.doi.org/10.1016/j.asd.2010.08.002
19. Posnien N, Koniszewski N, Hein H, Bucher G. Candidate gene screen in the red flour
beetle tribolium reveals six3 as ancient regulator of anterior median head and central
complex development. PLoS Genet. 2011;7. https://doi.org/10.1371/journal.pgen.1002416
20. Honkanen A, Adden A, Da Silva Freitas J, Heinze S. The insect central complex and the
neural basis of navigational strategies. J Exp Biol. 2019;222.
https://doi.org/10.1242/jeb.188854
21. Pfeiffer K, Homberg U. Organization and functional roles of the central complex in the
insect brain. Annu Rev Entomol. 2014;59:165–84. https://doi.org/10.1146/annurev-ento-
011613-162031
22. Hartenstein V, Stollewerk A. The evolution of early neurogenesis. Dev Cell. Elsevier Inc.;
2015;32:390–407http://dx.doi.org/10.1016/j.devcel.2015.02.004
23. Doe CQ, Skeath JB. Neurogenesis in the insect central nervous system. Curr Opin
Neurobiol. 1996;6:18–24. https://doi.org/10.1016/S0959-4388(96)80004-3
24. Stollewerk A. A flexible genetic toolkit for arthropod neurogenesis. Philos Trans R Soc B
Biol Sci. 2016;371. https://doi.org/10.1098/rstb.2015.0044
25. Boyan GS, Reichert H. Mechanisms for complexity in the brain: Generating the insect
central complex. Trends Neurosci. Elsevier Ltd; 2011. p. 247–57. 10.1016/j.tins.2011.02.002
26. Strausfeld NJ. Arthropod Brains: Evolution, functional elegance, and historical
significance. The Belknap Press of Harvard University Press; 2012.
27. Boyan GS, Reichert H. Mechanisms for complexity in the brain: Generating the insect
central complex. Trends Neurosci. Elsevier Ltd; 2011;34:247–57.
http://dx.doi.org/10.1016/j.tins.2011.02.002
28. Boyan G, Liu Y, Khalsa SK, Hartenstein V. A conserved plan for wiring up the fan-shaped
body in the grasshopper and Drosophila. Dev Genes Evol. Development Genes and
Evolution; 2017;227:253–69. https://doi.org/10.1007/s00427-017-0587-2
29. Walsh KT, Doe CQ. Drosophila embryonic type II neuroblasts: Origin, temporal
patterning, and contribution to the adult central complex. Dev. 2017;144:4552–62.
https://doi.org/10.1242/dev.157826
30. Bayraktar OA, Doe CQ. Combinatorial temporal patterning in progenitors expands neural
diversity. Nature. Nature Publishing Group; 2013;498:449–55.
https://doi.org/10.1038/nature12266
31. Malatesta P, Appolloni I, Calzolari F. Radial glia and neural stem cells. Cell Tissue Res.
2008;331:165–78. https://doi.org/10.1007/s00441-007-0481-8
32. Barry DS, Pakan JMP, McDermott KW. Radial glial cells: Key organisers in cns
development. Int J Biochem Cell Biol. Elsevier Ltd; 2014;46:76–9.
http://dx.doi.org/10.1016/j.biocel.2013.11.013
33. Caussinus E, Gonzalez C. Induction of tumor growth by altered stem-cell asymmetric

division in Drosophila melanogaster. Nat Genet. 2005;37:1125–9.

https://doi.org/10.1038/ng1632

34. Homem CCF, Knoblich JA. Drosophila neuroblasts: A model for stem cell biology. Dev.

2012;139:4297–310. https://doi.org/10.1242/dev.080515

35. Bowman SK, Rolland V, Betschinger J, Kinsey KA, Emery G, Knoblich JA. The Tumor

Suppressors Brat and Numb Regulate Transit-Amplifying Neuroblast Lineages in Drosophila.

Dev Cell. 2008;14:535–46. 10.1016/j.devcel.2008.03.004

36. Southall TD, Brand AH. Neural stem cell transcriptional networks highlight genes

essential for nervous system development. EMBO J. 2009;28:3799–807.

https://doi.org/10.1038/emboj.2009.309

37. Bello BC, Izergina N, Caussinus E, Reichert H. Amplification of neural stem cell

proliferation by intermediate progenitor cells in Drosophila brain development. Neural Dev.

2008;3. https://doi.org/10.1186/1749-8104-3-5

38. Boyan G, Williams L. Embryonic development of the insect central complex: Insights

from lineages in the grasshopper and Drosophila. Arthropod Struct Dev. Elsevier Ltd;

2011;40:334–48. http://dx.doi.org/10.1016/j.asd.2011.02.005

39. Izergina N, Balmer J, Bello B, Reichert H. Postembryonic development of transit

amplifying neuroblast lineages in the Drosophila brain. Neural Dev. 2009;4:1–13.

https://doi.org/10.1186/1749-8104-4-44

40. Boyan G, Williams L, Legl A, Herbert Z. Proliferative cell types in embryonic lineages of

the central complex of the grasshopper Schistocerca gregaria. Cell Tissue Res.

2010;341:259–77. https://doi.org/10.1007/s00441-010-0992-6

41. Garcia-Perez NC, Bucher G, Büscher M. Shaking hands is a homeodomain transcription

factor that controls axon outgrowth of central complex neurons in the insect model

Tribolium. Development. 2021. https://doi.org/10.1242/dev.199368

42. Pfeiffer BD, Jenett A, Hammonds AS, Ngo TTB, Misra S, Murphy C, et al. Tools for

neuroanatomy and neurogenetics in Drosophila. Proc Natl Acad Sci U S A. 2008;105:9715–

20. https://doi.org/10.1073/pnas.080369710

43. Weng M, Golden KL, Lee CY. dFezf/Earmuff Maintains the Restricted Developmental

Potential of Intermediate Neural Progenitors in Drosophila. Dev Cell. Elsevier; 2010;18:126–

35. http://dx.doi.org/10.1016/j.devcel.2009.12.007

44. Xie Y, Li X, Deng X, Hou Y, O'Hara K, Urso A, et al. The Ets protein Pointed prevents both

premature differentiation and dedifferentiation of Drosophila intermediate neural

progenitors. Development. 2016;143:3109–18.

http://dev.biologists.org/lookup/doi/10.1242/dev.137281

45. Zhu S, Barshow S, Wildonger J, Yeh L, Jan Y. Ets transcription factor Pointed promotes

the generation of intermediate neural progenitors in Drosophila larval brains.

2011;108:20615–20. https://doi.org/10.1073/pnas.111859510

46. Bayraktar OA, Boone JQ, Drummond ML, Doe CQ. Drosophila type II neuroblast lineages

keep Prospero levels low to generate large clones that contribute to the adult brain central

complex. Neural Dev. 2010;5. https://doi.org/10.1186/1749-8104-5-26

47. Chen R, Hou Y, Connell M, Zhu S. Homeodomain protein Six4 prevents the generation of

supernumerary Drosophila type II neuroblasts and premature differentiation of

intermediate neural progenitors. PLoS Genet. 2021.

http://dx.doi.org/10.1371/journal.pgen.1009371

48. Choi HMT, Schwarzkopf M, Fornace ME, Acharya A, Artavanis G, Stegmaier J, et al. Third-

generation in situ hybridization chain reaction: Multiplexed, quantitative, sensitive,

versatile, robust. Dev. 2018;145:1–10. https://doi.org/10.1242/dev.165753

49. Choi HMT, Chang JY, Trinh LA, Padilla JE, Fraser SE, Pierce NA. Programmable in situ
amplification for multiplexed imaging of mRNA expression. Nat Biotechnol. Nature
Publishing Group; 2010;28:1208–12. https://doi.org/10.1038/nbt.1692

50. Álvarez JA, Díaz-Benjumea FJ. Origin and specification of type II neuroblasts in the
Drosophila embryo. Dev. 2018;145:1–10. https://doi.org/10.1242/dev.158394

51. Brunner D, Dücker K, Oellers N, Hafen E, Scholzi H, Klambt C. The ETS domain protein
Pointed-P2 is a target of MAP kinase in the Sevenless signal transduction pathway. Nature.
1994;370:386–9. https://doi.org/10.1038/370386a0

52. Klambt C. The Drosophila gene pointed encodes two ETS-like proteins which are
involved in the development of the midline glial cells. Development. 1993;117:163–76.
https://doi.org/10.1242/dev.117.1.163

53. Bullinger G. Conservation and divergence of the retinal homeobox genetic neural lineage
between Drosophila melanogaster and Tribolium castaneum. University of Göttingen; 2024.
http://dx.doi.org/10.53846/goediss-10513

54. Dreyer D, Vitt H, Dippel S, Goetz B, el Jundi B, Kollmann M, et al. 3D standard brain of
the red flour beetle Tribolium castaneum: a tool to study metamorphic development and
adult plasticity. Front Syst Neurosci. 2010;4:1–13.
http://journal.frontiersin.org/article/10.3389/neuro.06.003.2010/abstract

55. Steinmetz PRH, Urbach R, Posnien N, Eriksson J, Kostyuchenko RP, Brena C, et al. Six3
demarcates the anterior-most developing brain region in bilaterian animals. Evodevo.
BioMed Central Ltd; 2010;1:14. http://www.evodevojournal.com/content/1/1/14

56. Posnien N, Koniszewski N, Bucher G. Insect Tc-six4 marks a unit with similarity to
vertebrate placodes. Dev Biol. Elsevier Inc.; 2011;350:208–16.
http://dx.doi.org/10.1016/j.ydbio.2010.10.024

57. Urbach R, Technau GM. Neuroblast formation and patterning during early brain
development in Drosophila. BioEssays. 2004;26:739–51. https://doi.org/10.1002/bies.20062

58. Bier E, Vaessin H, Younger-shepherd S, Jan LY. deadpan, an essential pan-neural gene in
Drosophila ,encodes a helix-loop-helix protein similar to the hairy gene product. Genes Dev.
1992;2137–51. 10.1101/gad.6.11.2137

59. Li X, Xie Y, Zhu S. Notch maintains Drosophila type II neuroblasts by suppressing
expression of the Fez transcription factor Earmuff. Development. 2016;143:2511–21.
http://dev.biologists.org/lookup/doi/10.1242/dev.136184

60. Campos-Ortega JA, Hartenstein V. The embryonic development of Drosophila
melanogaster. 1st ed. Berlin, Heidelberg: Springer; 1985.

61. de Velasco B, Erclik T, Shy D, Sclafani J, Lipshitz H, McInnes R, et al. Specification and
development of the pars intercerebralis and pars lateralis, neuroendocrine command
centers in the Drosophila brain. Dev Biol. 2007;302:309–23.
https://doi.org/10.1016/j.ydbio.2006.09.035

62. Hwang HJ, Rulifson E. Serial specification of diverse neuroblast identities from a
neurogenic placode by Notch and Egfr signaling. Development. 2011;138:2883–93.
https://doi.org/10.1242/dev.055681

63. Lee HH, Frasch M. Survey of Forkhead Domain Encoding Genes in the Drosophila
Genome: Classification and Embryonic Expression Patterns. Dev Dyn. 2004;229:357–66.
https://doi.org/10.1002/dvdy.10443

64. Schinko JB, Weber M, Viktorinova I, Kiupakis A, Averof M, Klingler M, et al. Functionality
of the GAL4/UAS system in Tribolium requires the use of endogenous core promoters. BMC

Dev Biol. 2010;10. https://doi.org/10.1186/1471-213X-10-53

65. Schinko JB, Hillebrand K, Bucher G. Heat shock-Mediated misexpression of genes in the
beetle Tribolium castaneum. Dev Genes Evol. 2012;222:287–98.
https://doi.org/10.1007/s00427-012-0412-x

66. Berghammer AJ, Klingler M, Wimmer E a. A universal marker for transgenic insects.
Nature. 1999;402:370–1. http://www.ncbi.nlm.nih.gov/pubmed/10586872

67. Gratz SJ, Ukken FP, Rubinstein CD, Thiede G, Donohue LK, Cummings AM, et al. Highly
specific and efficient CRISPR/Cas9-catalyzed homology-directed repair in Drosophila.
Genetics. 2014;196:961–71. https://doi.org/10.1534/genetics.113.160713

68. Buescher M, Oberhofer G, Garcia-Perez NC, Bucher G. A Protocol for Double Fluorescent
In Situ Hybridization and Immunohistochemistry for the Study of Embryonic Brain
Development in Tribolium castaneum. Methods Brain Dev. 2020. p. 219–32.
https://link.springer.com/protocol/10.1007/978-1-4939-9732-9_12

69. Tidswell ORA, Benton MA, Akam M. The neuroblast timer gene nubbin exhibits
functional redundancy with gap genes to regulate segment identity in Tribolium. Dev.
2021;148. https://doi.org/10.1242/dev.199719

70. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: An
open source platform for biological image analysis. Nat Methods. 2012;9:676–82.
https://doi.org/10.1038/nmeth.2019

Acknowledgements

We would like to thank Claudia Hinners for technical support with molecular biology methods, and Elke Küster for help with screening and beetle stock keeping. We also thank Christoph Viehbahn for valuable feedback on the project. Author Simon Rethemeier was supported by a scholarship of the *Göttingen Promotionskolleg für Medizinstudierende*, funded by the *Jacob-Henle-Programm*.