Epigenetic targeting of PGBD5-dependent DNA damage in SMARCB1-deficient sarcomas

Yaniv Kazansky[1,2], Helen S. Mueller[1,2], Daniel Cameron[1,2], Phillip Demarest[1,2], Nadia Zaffaroni[3], Noemi Arrighetti[3], Valentina Zuco[3], Prabhjot S. Mundi[4], Yasumichi Kuwahara[5], Romel Somwar[6], Rui Qu[7], Andrea Califano[8], Elisa de Stanchina[7], Filemon S. Dela Cruz[8], Andrew L. Kung[8], Mrinal M. Gounder[9], Alex Kentsis[1,2,10]*

[1] Molecular Pharmacology Program, Sloan Kettering Institute, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[2] Tow Center for Developmental Oncology, Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[3] Molecular Pharmacology Unit, Department of Experimental Oncology, Fondazione IRCCS Istituto Nazionale dei Tumori di Milano, Milan, Italy
[4] Department of Systems Biology, Columbia University Irving Medical Center, New York, NY, USA
[5] Department of Biochemistry and Molecular Biology, Kyoto Prefectural University of Medicine
[6] Department of Pathology, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[7] Antitumor Assessment Core, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[8] Department of Pediatrics, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[9] Department of Medicine, Memorial Sloan Kettering Cancer Center, New York, NY, USA
[10] Departments of Pediatrics, Pharmacology, and Physiology & Biophysics, Weill Medical College of Cornell University, New York, NY, USA

* Correspondence: Alex Kentsis, MD, PhD, email kentsisresearchgroup@gmail.com, telephone +1-646-888-2593, 1275 York Avenue, ZRC-1863, New York, NY 10021

Running title: Epigenetic targeting of DNA damage
Keywords: epigenetics, EZH2 inhibition, BAF, combination therapy, PRC2

Conflict of Interest

The authors declare no potential conflicts of interest. AK is a consultant for Novartis, Rgenta, Blueprint, and Syndax.

Abstract

Despite the potential of targeted epigenetic therapies, most cancers do not respond to current epigenetic drugs. The Polycomb repressive complex EZH2 inhibitor tazemetostat was recently approved for the treatment of *SMARCB1*-deficient epithelioid sarcomas, based on the functional antagonism between PRC2 and loss of SMARCB1. Through the analysis of tazemetostat-treated patient tumors, we recently defined key principles of their response and resistance to EZH2 epigenetic therapy. Here, using transcriptomic inference from *SMARCB1*-deficient tumor cells, we nominate the DNA damage repair kinase ATR as a target for rational combination EZH2 epigenetic therapy. We show that EZH2 inhibition promotes DNA damage in epithelioid and rhabdoid tumor cells, at least in part via its induction of the transposase-derived PGBD5. We leverage this collateral synthetic lethal dependency to target PGBD5-dependent DNA damage by inhibition of ATR but not CHK1 using elimusertib. Consequently, combined EZH2 and ATR inhibition improves therapeutic responses in diverse patient-derived epithelioid and rhabdoid tumors *in vivo*. This advances a combination epigenetic therapy based on EZH2-PGBD5 synthetic lethal dependency suitable for immediate translation to clinical trials for patients.

Introduction

Targeted epigenetic therapies offer potential improvements over conventional cytotoxic chemotherapies through superior clinical efficacy and reduced toxicity. In cancers caused by genetic mutations of transcriptional and epigenetic regulators, specific inhibition of epigenetic effectors can directly block dysregulated gene expression by leveraging cancer-specific dependencies. One promising example of this therapeutic approach is in cancers caused by mutation of the chromatin remodeling SWI/SNF/BAF (Brg/Brahma-associated factors) complex, a ubiquitous epigenetic regulator that is mutated in at least 20% of human cancers (1). In particular, the highly lethal solid tumors caused by loss of the core BAF subunit *SMARCB1* (2-4), which include malignant rhabdoid tumors (MRT) and epithelioid sarcomas (ES), are known to be dependent on the methyltransferase EZH2, a component of the Polycomb Repressive Complex 2 (PRC2). This dependency is thought to result from epigenetic antagonism between BAF and PRC2, in which normal BAF activity evicts PRC2 from tumor suppressor loci (5, 6). In a recent clinical trial, this dependency was targeted using the EZH2 methyltransferase inhibitor tazemetostat (TAZ), leading to its FDA approval (7). Yet despite the promise of this therapeutic approach, EZH2 inhibition as a monotherapy exhibited efficacy in only a subset of patients, with most patient tumors having either primary resistance or acquiring resistance after treatment (7, 8). Thus, there is a critical need to develop improved epigenetic combination therapies, which can be achieved at least in part through improved understanding of the effects of EZH2 inhibition in epithelioid and rhabdoid tumors.

We recently leveraged functional genomics of epithelioid and rhabdoid tumors to elucidate the mechanisms of resistance to EZH2 inhibition, including frequent disruption of the RB1/E2F cell cycle control axis causing TAZ resistance in patients. This discovery led to the identification of the AURKB inhibitor barasertib combination therapy that can overcome *RB1*-mediated TAZ resistance *in vitro* and improve TAZ responses in preclinical epithelioid and rhabdoid tumor xenografts *in vivo* (8). This approach bypasses primary and acquired RB1/E2F-mediated checkpoint defects to maintain TAZ-induced reprogramming of oncogenic gene expression. However, RB1/E2F pathway alterations were observed in less than half of epithelioid and rhabdoid tumors, raising the question of additional combinations that may achieve effective epigenetic therapy for patients, including those with primary or acquired TAZ resistance.

Epithelioid and rhabdoid tumors occur due to characteristic biallelic deletions and inactivating
mutations of *SMARCB1*, either as a result of germline rhabdoid tumor predisposition syndrome,
or due to somatic mutations (2, 9, 10). For rhabdoid tumors, differences in the types of mutations,
e.g. large versus small chromosomal deletions or missense and nonsense mutations of
SMARCB1 correspond with distinct molecular subtypes and clinical features (11). Genome
sequencing studies have identified distinct sequence-specific mutational features in rhabdoid
tumors, including sequence-specific deletions of *SMARCB1* and recurrent mutations of other
genes, associated at least in part with the mutagenic activity of the domesticated transposase-
derived Piggybac Transposable Element Derived 5 (PGBD5) (9).

PGBD5 is among the most evolutionarily conserved domesticated transposase-derived genes in
vertebrates and can mediate sequence-specific DNA rearrangements dependent on its putative
nuclease activity (9, 12). In particular, Pgbd5 promotes sequence-specific somatic mutagenesis
and tumor development in mouse models of medulloblastoma, one of the most common childhood
brain tumors (13). PGBD5-expressing cells, including rhabdoid tumor cells, require a distinct form
of non-homologous end-joining (NHEJ) DNA repair, leading to their hypersensitivity to inhibition
of DNA repair signaling, specifically ATR inhibition (14). We have now found that TAZ inhibition
of EZH2 can increase PGBD5 expression in epithelioid and rhabdoid tumor cells, potentiating
PGBD5-dependent DNA damage, and conferring synergy with the ATR-selective DNA repair
inhibitor elimusertib. Accordingly, combined treatment of patient-derived rhabdoid and epithelioid
tumors engrafted into immunodeficient mice, including from patients with multiply relapsed
metastatic disease, demonstrates synergistic activity *in vivo*.

Results

To define potential therapeutic targets for TAZ combination therapy in *SMARCB1*-deficient
tumors, we leveraged a recently developed method for transcriptomic inference of oncogenic
protein activity, termed metaVIPER (Virtual Inference of Protein activity by Enriched Regulon), as
applied to gene expression profiles of 68 patient rhabdoid tumors (15-17). We used the
OncoTarget annotation of high-affinity pharmacologic inhibitors of putative master gene
expression regulators (MRs) to order them by their mean activity scores (**Figure 1A**). As
expected, EZH2 was among the most activated MR proteins in rhabdoid tumors, as was the
previously identified TAZ combination therapy target, AURKB (**Figure 1A**). This analysis also
prioritized CDK4, CDK2, and AURKA (**Figure 1A**), consistent with the previously established

activity of cell cycle checkpoints downstream of G1/S in rhabdoid tumors (8), validating this
ranking approach.

In addition to these known dependencies, this analysis also identified the NHEJ DNA repair
structural factor XRCC6 (Ku70), and its mediator kinase ATR among the most activated MR
genes among all 6,000 MR proteins in metaVIPER (**Figure 1B**), both of which have been shown
to be required for the survival of cells expressing active PGBD5 in prior studies (14). Indeed,
XRCC6 ranked higher in this analysis than the well-validated target EZH2 (**Figure 1B**).

To ascertain the robustness of this prediction, we next analyzed an independent dataset from
Kuster and colleagues (18), which assessed the responses of sarcoma cell lines to a panel of 151
pharmacologic inhibitors, including G401 and A204 rhabdoid tumor and VA-ES-BJ epithelioid
sarcoma cell lines. We observed that the ATR-selective kinase inhibitor elimusertib was among
the most active drugs against epithelioid and rhabdoid tumor cells, even more active than both
TAZ and barasertib (**Figure 1C**). Thus, *SMARCB1*-deficient epithelioid and rhabdoid tumors are
highly sensitive to ATR inhibition (14).

Since both modulation of gene expression by TAZ and DNA repair by ATR inhibitor elimusertib
appears to have prominent activity in rhabdoid tumor cells, we inquired whether TAZ may regulate
PGBD5 itself. We found that in addition to the expected upregulation of Polycomb gene sets,
EZH2 inhibition in G401 rhabdoid tumor cells with TAZ for 11 days also caused a significant
increase in the expression of *PGBD5* (mean fold-increase of 6.6 and Student's t-test p = 1.2E-7;
Figure 1D). Using CRISPR gene editing, we generated isogenic *RB1* wild-type and *RB1^{del}* mutant
G401 cells and confirmed correct biallelic *RB1* inactivating mutations and consequent loss of RB1
protein expression, using the *AAVS1* safe harbor as a negative control (8). In prior work, we found
that defects in the RB1/E2F axis, including mutations of *RB1*, cause TAZ resistance (8).
Consistent with the independence of this TAZ resistance mechanism from ATR inhibitor
susceptibility, two independent G401 *RB1^{del}* mutant cell lines also exhibited significant TAZ-
mediated induction of *PGBD5* expression (mean fold-increase of 333 and 8.6 and t-test p = 1.3E-
6 and 1.6E-6 for E1 and F2 clones, respectively; **Figure 1D**). Additionally, both *RB1^{WT}* and *RB1^{del}*
mutant G401 cells exhibited nanomolar susceptibility to the ATR-selective inhibitor elimusertib
(half-maximal effective concentration of 18 ± 1.6 nM, 19 ± 3.8 nM, and 27 ± 3.2 nM for *RB1^{WT}*,
RB1^{del} E1, and F2 clones, respectively; **Figure 1E**).

Elimusertib is currently undergoing clinical trials in patients with refractory or relapsed solid
tumors, including patients with PGBD5-expressing tumors such as MRT and ES (Clinical Trials
Identifier NCT05071209). We therefore investigated the activity of combination treatment with
TAZ and elimusertib, reasoning that TAZ-induced upregulation of *PGBD5* expression may
potentiate the anti-tumor effects of ATR inhibition. We observed greater antitumor effects from
the combination of TAZ and elimusertib than either drug alone against diverse MRT and ES cell
lines, including TAZ-resistant ES1 and VAESBJ cells (mean fold-decrease in normalized cell
viability of 0.64, 0.63, 0.78, 0.84, and 0.84 compared with most effective monotherapy for
TTC642, KP-MRT-NS, KP-MRT-RY, G401 and ES1 respectively, t-test 1.0E-3, 3.0E-5, 3.3E-3,
2.6E-4, and 6.1E-5 ;**Figure 2A**). Strikingly, the combination of TAZ and elimusertib was highly
synergistic in G401 rhabdoid and ES1 epithelioid sarcomas cells (ZIP synergy = 3.4 and 2.0 and
p = 9.8E-1 and 7.3E-1, respectively; **Figure 2B-C**).

If the improved anti-tumor activity of TAZ and elimusertib is due to the induction of PGBD5-
dependent DNA damage, then this treatment combination should be associated with the induction
of DNA damage repair signaling. To test this prediction, we used confocal immunofluorescence
microscopy to quantify γH2AX as a specific marker of DNA damage (19). In agreement with prior
studies (14), vehicle-treated G401 cells showed measurable γH2AX staining associated with
baseline *PGBD5* expression (**Figure 2D-E**). Consistent with TAZ-mediated induction of *PGBD5*
expression (**Figure 1D**), we found that TAZ treatment alone significantly increased nuclear γH2AX
(mean normalized intensity = 0.42 versus 0.29 for TAZ and DMSO, respectively; *t*-test *p* = 9.3E-
3; **Figures 2D-E**). The combination of TAZ and elimusertib induced a significant increase in
γH2AX as compared to either drug alone (mean normalized intensity = 1.8 for the combination
versus 0.42 and 0.85 for TAZ and elimusertib, respectively; *t*-test *p* = 6.2E-14 and 1.5E-29 for
combination versus elimusertib and TAZ, respectively; **Figures 2D-E**).

Recently, EZH2 suppression was shown to induce replication stress through upregulation of
MYCN expression in T-ALL cells, which in turn sensitized cells to inhibition of CHK1 (19), a
downstream mediator of ATR signaling (20). To test this possibility, we measured *MYCN*
expression in TAZ-treated G401 rhabdoid tumor cells, which was significantly increased (**Figure**
3A). However, this induction of *MYCN* gene expression was not associated with the accumulation
of MYCN protein, as measured by Western blotting with *MYCN*-amplified IMR5 neuroblastoma
cells as a positive control (**Figure 3B**). Consistent with this, EZH2 inhibition either alone or in

combination with CHK1 inhibition did not induce apparent replication stress, as measured by RPA
phosphorylation, using the treatment of cells with the DNA topoisomerase inhibitor camptothecin
as positive control for replication stress. This was despite the effective inhibition of CHK1 auto-
phosphorylation by the CHK1-selective SRA737 inhibitor (**Figure 3C**). Concordantly, the CHK1
inhibitor SRA737 showed poor activity against G401 cells, regardless of *RB1* status (**Figure 3D**).
Thus, rhabdoid tumor cells exhibit a specific dependency on ATR-dependent but CHK1-
independent DNA damage repair signaling.

TAZ-mediated induction of *PGBD5* and DNA damage would indicate that this form of DNA
damage should be dependent on *PGBD5* expression. To test this prediction, we engineered short
hairpin RNA-mediated depletion of *PGBD5* in G401 cells using lentiviral transduction of two
independent *PGBD5*-specific shRNAs (shPGBD5), as compared to a control GFP-targeting
shRNA (shGFP). We confirmed that shPGBD5-expressing cells were significantly depleted of
PGBD5 as compared to control shGFP cells (mean fold-depletion = 0.59 and 0.54; t-test p = 4.7E-
3 and 1.2E-3, respectively; **Figure 4A**). We then measured DNA damage using quantitative
confocal immunofluorescence microscopy of γH2AX, and found that while the combination of TAZ
and elimusertib induced DNA damage in shPGBD5 cells, this effect was significantly reduced
compared to control shGFP cells (mean fold-reduction = 0.34 and 0.39; t-test p = 3.9E-8 and
4.4E-8 for shGFP versus shPGBD5-1 and shPGBD5-3, respectively; **Figure 4B-C**). This PGBD5-
dependent reduction in DNA damage was observed both when examining all nuclei with γH2AX
staining (**Figure 4B**), as well as only nuclei with punctate γH2AX staining, corresponding to more
localized DNA damage (mean fold-reduction = 0.59 and 0.61; t-test p = 2.4E-5 and 2.7E-5 for
shGFP versus shPGBD5-1 and shPGBD5-3, respectively ;**Figure 4D**), as opposed to pan-nuclear
γH2AX staining due to genome-wide unrepaired DNA damage and cellular apoptosis (mean fold-
reduction = 0.31 and 0.55; t-test p = 9.0E-4 and 1.6E-2 for shGFP versus shPGBD5-1 and
shPGBD5-3, respectively; **Figure 4E**). Thus, PGBD5 is at least in part necessary for TAZ-
mediated induction of DNA damage and its potentiation by the combination with elimusertib.

This nominates therapeutic targeting of the EZH2-PGBD5 synthetic lethal dependency as an
improved combination strategy for epithelioid and rhabdoid tumors. To test this idea, we
assembled a phase 2-like cohort of diverse MRT and ES tumors derived from patients with
relapsed and metastatic disease, including tumors with numerous additional acquired mutations
(**Supplementary Table S1**). We engrafted these tumors into immunodeficient *NOD-scid*
IL2Rgamma^{null} mice, and randomized tumor-bearing animals to treatment with TAZ or elimusertib

or the combination of both (**Figure 5A**). The combination of TAZ and elimusertib exceeded the effect of treatment with either drug alone when assessed by tumor growth measurements (Vardi *U*-test p = 2.33E-2 and 0.20 for combination versus elimusertib or TAZ, respectively; **Figure 5A**) and significantly extended tumor-free survival from 51 days (95% CI = 42-60 days) for elimusertib and 68 days (95% CI = 53-82 days) for TAZ to 100 days (95% CI = 76-124 days) for the combination (log-rank test p = 5.8E-4 and 3.8E-2 for combination versus elimusertib or TAZ, respectively; **Figure 5B**).

This activity was most pronounced for the HYMAD_EPIS_X0004aS1 and SOMWR_EPIS_X00013aS1 PDX models (**Figures 5C-H**), despite the former exhibiting a relatively poor response to TAZ monotherapy, when assessed by tumor growth measurements (Vardi *U*-test p = 2.0E-4 for combination versus elimusertib or TAZ for HYMAD_EPIS_X0003aS1; **Figure 5C**; p = 1.0E-3 and 6.0E-2 for combination versus elimusertib and TAZ, respectively for SOMWR_EPIS_X00013aS1; **Figure 5E**) and tumor-free survival (log-rank test p = 6.2E-3 and 6.3E-5 for combination versus elimusertib or TAZ, respectively for HYMAD_EPIS_X0003aS1; **Figure 5D**; p = 9.5E-5 and 6.0E-2 for combination versus elimusertib or TAZ, respectively for SOMWR_EPIS_X00013aS1; **Figure 5F**). This effect leverages the EZH2-PGBD5 collateral synthetic lethal dependency, targeting PGBD5-dependent DNA damage to improve TAZ clinical response and overcome resistance in epithelioid and rhabdoid tumors.

Discussion

Prior studies showed that inhibition of EZH2 methyltransferase activity using tazemetostat is insufficient to induce durable tumor regressions in most patients with epithelioid and rhabdoid tumors (8). Our recent functional genetic studies of patient tumors before and after clinical TAZ therapy led to a specific model of effective epigenetic therapy, including rational combinations to overcome RB1/E2F pathway defects that were observed in 43% of tumors with primary or acquired TAZ resistance (8). Here, we further advanced this approach by identifying a collateral synthetic lethal dependency between EZH2 and PGBD5 in rhabdoid and epithelioid sarcomas that confers a susceptibility to combined epigenetic therapy using EZH2 and ATR inhibition. We found that EZH2 inhibition can upregulate the transposase-derived PGBD5 nuclease, which induces DNA damage, requiring ATR but not CHK1-mediated DNA damage repair signaling in *SMARCB1*-deficient rhabdoid and epithelioid tumors regardless of their *RB1* mutational status. As a result, combined EZH2 and ATR inhibition exerts synergistic anti-tumor effects, as measured

by DNA damage induction *in vitro* and tumor growth reduction and improvements in tumor-free

survival *in vivo*.

The combination of EZH2 and ATR inhibition offers both a promising therapeutic approach that

can be rapidly translated to clinical trials for patients with rhabdoid and epithelioid sarcomas,

and a compelling example of drug-induced synthetic lethality. Originally described as genetic

interactions (21), synthetic lethal targeting has proven to be a powerful therapeutic approach for

the treatment of cancers. For example, breast and ovarian carcinomas with *BRCA1/2* mutations

exhibit increased dependence on poly(adenosine diphosphate-ribose) polymerase (PARP)-

mediated DNA repair, conferring a susceptibility to PARP1/2 inhibitors such as olaparib (22, 23).

Synthetic dependencies in DNA damage signaling, chromatin remodeling, and metabolic

functions have recently been defined to develop improved targeted therapies (24-28). In

particular, acquired defects in DNA repair in tumor cells, such as inactivating mutations or

functional defects in ATM signaling, can confer susceptibility to ATR inhibition due to the

specific requirements of concurrent ATM and ATR signaling for efficient NHEJ DNA repair and

tumor cell survival (28-32).

ATR-selective inhibitors also synergize with genotoxic chemotherapies, such as DNA cross-

linking platinum drugs, due to the specific requirements of ATR-dependent DNA repair during

DNA replication (28). However, this approach has limited therapeutic efficacy due to its effects

on healthy cells. Thus, effective targeting of DNA damage repair must leverage tumor-specific

mutational and repair processes, providing a rationale for combining epigenetic and DNA repair

inhibitors. In the case of rhabdoid and epithelioid sarcomas with *SMARCB1* deficiency, the

epigenetic antagonism between the chromatin remodeling activities of BAF and PRC2

contributes to the dependency of tumor cells on EZH2 activity (5), whose inhibition promotes the

expression of tumor suppressors otherwise repressed by PRC2 (8, 33). This epigenetic

reprogramming also appears to increase the expression of PGBD5, with the associated DNA

damage and requirement for its ATR-dependent repair. It remains to be defined whether this

effect is due to direct repression through PRC2-mediated histone 3 lysine 27 trimethylation of

the *PGBD5* locus and/or indirect transcriptional or post-transcriptional regulation of *PGBD5*

expression.

Though the cell of origin of rhabdoid and epithelioid sarcomas is currently unknown, these tumors

exhibit epigenetic and transcriptional features of neuronal and neural crest cells (34-36). EZH2 is

known to regulate neuronal differentiation (37-40), and therefore it will also be important to determine whether EZH2 inhibition of other PGBD5-expressing tumors, such as neuroblastomas, medulloblastomas, and Ewing and other fusion sarcomas, all of which also share features of neuronal lineages, can also induce PGBD5-dependent DNA damage, thereby conferring a susceptibility to collateral synthetic lethal targeting with ATR inhibitors. Importantly, susceptibility to ATR inhibition can also result from PGBD5-independent sources of intrinsic DNA damage, such as alternative lengthening of telomeres (ALT) (41), replication stress due to transcriptional-replication interference (42-44), and functional ATM defects (30, 45), all of which may occur concurrently in tumor cells undergoing PGBD5-dependent DNA damage. Similarly, EZH2 may also contribute to other mechanisms of DNA damage repair (46-53). Thus, future biochemical and genetic studies will be needed to define specific mechanisms of EZH2 and ATR-dependent DNA damage repair signaling in tumor and healthy cells to identify improved targets to develop exclusively tumor-selective therapies. This direction is particularly compelling because PGBD5-dependent DNA damage repair appears to require ATR but not CHK1 kinase signaling in epithelioid and rhabdoid tumor cells, in contrast to ATR/CHK1-dependent canonical signaling observed during replication stress in healthy tissues.

Further definition of the EZH2-PGBD5 collateral synthetic lethal mechanisms should also aid in therapy stratification of combined EZH2 and ATR inhibition. For example, apparent variation in response to the tazemetostat-elimusertib combination between patient-derived epithelioid and rhabdoid tumors may result from biological differences in DNA damage repair signaling, PGBD5 activity, or other sources of intrinsic DNA damage, which in turn may be associated with the recently described distinct molecular subtypes of ES and MRT tumors (54). In addition, both EZH2 and ATR inhibition can be immunogenic (8, 55-57), and future studies will be needed to define immunologic effects of this combination therapy that may contribute to therapeutic effects in patients. In all, this work emphasizes how improved understanding of collateral dependencies of intrinsic mutators and epigenetic dysregulation responsible for causing childhood and young-onset cancers can be leveraged for rational combination therapies.

Acknowledgements

This work is dedicated to Maggie Schmidt and her family, and to the many other patients and their advocates who inspire and support our research. We thank Richard Koche, Nicholas Socci, and Mithat Gönen for technical advice, Marc Ladanyi for patient-derived xenografts,

members of our labs for critical advice and manuscript comments, and Epizyme (now Ipsen) and Bayer for supplying tazemetostat and elimusertib, respectively. This work was supported by the MSK Integrated Genomics Operation Core, Anti-Tumor Assessment Core, Bioinformatics Core, Molecular Diagnostics Service and the Department of Pathology, the Marie-Josée and Henry R. Kravis Center for Molecular Oncology, NIH R01 CA214812, P30 CA08748, T32 GM007739, Burroughs Wellcome Fund, Rita Allen Foundation, Pershing Square Sohn Cancer Research Alliance and the G. Harold and Leila Y. Mathers Foundation, Cycle for Survival, MSK Sarcoma Center, the Starr Cancer Consortium, and Maggie's Mission. Yaniv Kazansky was supported by a Medical Scientist Training Program grant from the National Institute of General Medical Sciences of the National Institutes of Health under award number T32 GM007739 to the Weill Cornell/Rockefeller/Sloan Kettering Tri-Institutional MD-PhD Program. AK is a Scholar of the Leukemia & Lymphoma Society.

Methods

Cell culture

G401, TTC642, VAESBJ, and TM8716 cell lines were obtained from the American Type Culture Collection. The ES1 cell line was generated and kindly provided by Nadia Zaffaroni. Rhabdoid tumor cell lines KP-MRT-NS and KP-MRT-RY were kindly provided by Yasumichi Kuwahara and Hajime Hosoi. The identity of all cell lines was verified by STR analysis. Absence of *Mycoplasma* contamination was determined at every plating using the MycoAlert kit according to manufacturer's instructions (Lonza). Cell lines were cultured in 5% CO_2 in a humidified atmosphere in 37°C. All media were obtained from Corning and supplemented with 10% fetal bovine serum (FBS), 1% L-glutamine, and 100 U/mL penicillin and 100 µg/mL streptomycin (Gibco). G401, ES1, and VAESBJ cells were cultured in Dulbecco's Modified Eagle Medium (DMEM). TTC642, TM8716, KP-MRT-NS, and KP-MRT-RY cells were cultured in Roswell Park Memorial Institute (RPMI) medium. All experiments were performed using cell lines kept in culture for no more than 10 passages. Generation of G401 with *RB1* deletion was described previously (8) using the gRNA sequence in Table 1 below.

Western blotting

To assess protein expression by Western immunoblotting, pellets of 1 million cells were prepared and washed once in cold PBS. Cells were resuspended in 100-130 µL of RIPA lysis buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1.0% NP-40, 0.5% sodium deoxycholate, 0.1% sodium

dodecyl sulfate) and incubated on ice for 10 minutes. Cell suspensions were then disrupted using
a Covaris S220 adaptive focused sonicator for 5 minutes (peak incident power: 35W, duty factor:
10%, 200 cycles/burst) at 4 °C. Lysates were cleared by centrifugation at 18,000 g for 15 min at
4 °C. Protein concentration was assayed using the DC Protein Assay (Bio-Rad) and 15-35 µg
whole cell extract was used per sample. Samples were boiled at 95 °C in Laemmli buffer (Bio-
Rad) with 40 mM DTT and resolved using sodium dodecyl sulfate-polyacrylamide gel
electrophoresis. Proteins were transferred to Immobilon FL PVDF membranes (Millipore), and
membranes were blocked using Intercept Blocking buffer (Li-Cor). Primary antibodies used were:
anti-EZH2 (Cell Signaling Technology, 5246) at 1:1,000, anti-RPA32 pT21 (abcam, ab109394) at
1:2,000, anti-RPA32 pS4/pS8 (ThermoFisher, A300-245A) at 1:2,000, anti-pCHK1 S296 (Cell
Signaling Technology, 90178) at 1:250, anti-MYCN (Cell Signaling Technology, 9405) at 1:250,
anti-Actin (Cell Signaling Technology, 4970 and 3700) at 1:5,000. Blotted membranes were
visualized using goat secondary antibodies conjugated to IRDye 680RD or IRDye 800CW (Li-
Cor, 926-68071 and 926-32210) at 1:15,000 and the Odyssey CLx fluorescence scanner,
according to manufacturer's instructions (Li-Cor). Image analysis was done using the Li-Cor
Image Studio software (version 4).

Transcriptomic data
Expression levels of *PGBD5* and *MYCN* in G401 cells were determined from our previously
published dataset (8). Transcriptomic data and metaVIPER protein activity inference was
performed as previously described (15, 16).

Generating shPGBD5 cells
For shRNA cells, pLKO.1 shRNA vectors targeting *PGBD5* (TRCN0000138412,
TRCN0000135121) and control shGFP were obtained from the RNAi Consortium (Broad
Institute). Lentivirus production was carried out as described previously (8). G401 cells were
transduced at an MOI ~1.5 and selected with puromycin at 2 µg/mL for 72 hours. Knockdown was
confirmed by quantitative RT-PCR as previously described (9), using primers specified in the
Table 1.

Combination drug treatment
Drugs used for *in vitro* treatment were supplied by Selleckchem (TAZ, S7128; Elimusertib, S9864;
camptothecin, S1288; SRA737, S8253).

For combination treatment with TAZ and elimusertib, we used a two-dimensional dose matrix
design, treating the cells for 9 days. After the addition of cells, drugs were added using a pin tool
(stainless steel pins with 50 nL slots, V&P Scientific) mounted onto a liquid handling robot (CyBio
Well vario, Analytik Jena). On Day 9, CellTiter-Glo reagent was freshly reconstituted and added
in a 1:1 proportion to cell media, according to the manufacturer's instructions. A similar protocol
was used for monotherapy dose response curves for elimusertib and SRA737, although drug
solutions were added manually, and specific treatment times are indicated in the corresponding
figure legends. For analysis of synergy, we used the synergyFinder package (58). Outliers due to
pinning errors were excluded after manual examination.
403
<u>Immunofluorescence</u>
Immunofluorescence for γH2AX was performed on cells plated on Millicell EZ Slide glass slides
(EMD Millipore), coated for 45 minutes with bovine plasma fibronectin (Millipore Sigma). After
drug treatment, cells were washed once with PBS and fixed in 4% formaldehyde for 10 minutes
at room temperature. Slides were then washed three times in PBS for 5 minutes, permeabilized
for 15 minutes in 0.3% Triton X-100, washed again in PBS three times, and blocked with 5%
goat serum (Millipore Sigma, G9023) in PBS for 1 hour at room temperature. Slides were
incubated with mouse anti-γH2A.X primary antibody (Sigma-Aldrich, 05-636) at 1:500 in
blocking buffer for 1 hour, washed three times in PBS, and incubated with goat anti-mouse
secondary antibody conjugated to AlexaFluor555 (Invitrogen, A-21422) at 1:1,000. Cells were
then counterstained with DAPI at 1:1,000 for 10 minutes and treated with ProLong Diamond
Antifade Mountant with DAPI (Invitrogen, P36962) for 48 hours.
416
Images were acquired on a Zeiss LSM880 confocal microscope at 63x magnification. Images
were then processed using a custom pipeline in CellProfiler (59). Per-cell integrated γH2A.X
intensity was normalized against per-cell integrated DAPI intensity. All image analysis used the
same pipeline settings, with the exception of the RescaleIntensity module for the AF555
channel, which used the settings 0.009-0.09 for the images in Figure 2 and 0.005-0.09 for
Figure 4. CellProfiler analysis pipeline files and raw image files are available on Zenodo (DOI:
10.5281/zenodo.10982946). Overlaid images were prepared using Fiji (60).
424
<u>Xenografts</u>
All mouse experiments were carried out in accordance with institutionally approved animal use
protocols (8). To generate PDXs, tumor specimens were collected under approved IRB protocol

14-091, immediately minced and mixed (50:50) with Matrigel (Corning, New York, NY) and

implanted subcutaneously in the flank of 6-8 weeks-old female NOD.Cg-*Prkdc*^{scid} *Il2rg*^{tm1Wjl}/*Szj*

(NSG) mice (Jackson Laboratory, Bar Harbor, ME), as described previously (61). Mice were

monitored daily and PDX samples were serially transplanted three times before being deemed

established. PDX tumor histology was confirmed by review of H&E slides and direct comparison

to the corresponding patient tumor slides. PDX identity was further confirmed by MSK-IMPACT

sequencing analysis.

Therapeutic studies used female and male NSG mice obtained from the Jackson Laboratory.

Xenografts were prepared as single-cell suspensions, resuspended in Matrigel, and implanted

subcutaneously into the right flank of 6-10 week old mice. 100 µL of tumor cell suspension was

used for each mouse. Tumors were allowed to grow until they reached a volume of 100 mm^3, at

which point they were randomized into treatment groups without blinding. Drugs were prepared

using the following formulations: Tazemetostat was dissolved at 25 mg/mL in 5% DMSO, 40%

PEG 300, 5% Tween 80, and 50% water. Elimusertib was dissolved at 5 mg/mL in 10% DMSO,

40% PEG 300, 5% Tween 80, and 45% water using a sonicator. Drugs were reconstituted daily.

TAZ was dosed at 250 mg/kg twice daily by oral gavage, 7 days per week. Elimusertib was

dosed at 40 mg/kg twice daily by oral gavage using 2 days on and 12 days off cycle. Caliper

tumor measurements were taken twice weekly. Tumor volumes were calculated using the

formula Volume = (π/6) x length x width^2. Tumor growth analysis was performed using the Vardi

U-test (62), as implemented in the clinfun R package using the aucVardiTest function. Tumor-

free survival analysis was calculated using OriginPro (Microcal) by the Kaplan-Meier method,

using the log-rank test. Raw data and R scripts used for data analysis are available on Zenodo

(DOI: 10.5281/zenodo.10398544).

Supplementary Table S1: List of PDX models used in this study, with clinical characteristics of
the original tumor specimens, followed by a list of mutations found in all PDX models, as
determined by targeted MSK-IMPACT sequencing.

References

1. Kadoch C, Crabtree GR. Mammalian SWI/SNF chromatin remodeling complexes and cancer: Mechanistic insights gained from human genomics. Sci Adv. 2015;1(5):e1500447.
2. Versteege I, Sévenet N, Lange J, Rousseau-Merck MF, Ambros P, Handgretinger R, et al. Truncating mutations of hSNF5/INI1 in aggressive paediatric cancer. Nature. 1998;394(6689):203-6.
3. Kim KH, Kim W, Howard TP, Vazquez F, Tsherniak A, Wu JN, et al. SWI/SNF-mutant cancers depend on catalytic and non-catalytic activity of EZH2. Nat Med. 2015;21(12):1491-6.
4. McKenna ES, Sansam CG, Cho YJ, Greulich H, Evans JA, Thom CS, et al. Loss of the epigenetic tumor suppressor SNF5 leads to cancer without genomic instability. Mol Cell Biol. 2008;28(20):6223-33.
5. Wilson BG, Wang X, Shen X, McKenna ES, Lemieux ME, Cho YJ, et al. Epigenetic antagonism between polycomb and SWI/SNF complexes during oncogenic transformation. Cancer Cell. 2010;18(4):316-28.
6. Kadoch C, Williams RT, Calarco JP, Miller EL, Weber CM, Braun SM, et al. Dynamics of BAF-Polycomb complex opposition on heterochromatin in normal and oncogenic states. Nat Genet. 2017;49(2):213-22.
7. Gounder M, Schöffski P, Jones RL, Agulnik M, Cote GM, Villalobos VM, et al. Tazemetostat in advanced epithelioid sarcoma with loss of INI1/SMARCB1: an international, open-label, phase 2 basket study. Lancet Oncol. 2020;21(11):1423-32.
8. Kazansky Y, Cameron D, Mueller HS, Demarest P, Zaffaroni N, Arrighetti N, et al. Overcoming clinical resistance to EZH2 inhibition using rational epigenetic combination therapy. Cancer Discov. 2024.
9. Henssen AG, Koche R, Zhuang J, Jiang E, Reed C, Eisenberg A, et al. PGBD5 promotes site-specific oncogenic mutations in human tumors. Nat Genet. 2017;49(7):1005-14.
10. Biegel JA, Zhou JY, Rorke LB, Stenstrom C, Wainwright LM, Fogelgren B. Germ-line and acquired mutations of INI1 in atypical teratoid and rhabdoid tumors. Cancer Res. 1999;59(1):74-9.
11. Chun HE, Johann PD, Milne K, Zapatka M, Buellesbach A, Ishaque N, et al. Identification and Analyses of Extra-Cranial and Cranial Rhabdoid Tumor Molecular Subgroups Reveal Tumors with Cytotoxic T Cell Infiltration. Cell Rep. 2019;29(8):2338-54.e7.
12. Henssen AG, Henaff E, Jiang E, Eisenberg AR, Carson JR, Villasante CM, et al. Genomic DNA transposition induced by human PGBD5. Elife. 2015;4.
13. Yamada M, Keller RR, Gutierrez RL, Cameron D, Suzuki H, Sanghrajka R, et al. Childhood cancer mutagenesis caused by transposase-derived PGBD5. Sci Adv. 2024;10(12):eadn4649.
14. Henssen AG, Reed C, Jiang E, Garcia HD, von Stebut J, MacArthur IC, et al. Therapeutic targeting of PGBD5-induced DNA repair dependency in pediatric solid tumors. Sci Transl Med. 2017;9(414).
15. Ding H, Douglass EF, Jr., Sonabend AM, Mela A, Bose S, Gonzalez C, et al. Quantitative assessment of protein activity in orphan tissues and single cells using the metaVIPER algorithm. Nat Commun. 2018;9(1):1471.
16. Coutinho DF, Mundi PS, Marks LJ, Burke C, Ortiz MV, Diolaiti D, et al. Validation of a non-oncogene encoded vulnerability to exportin 1 inhibition in pediatric renal tumors. Med. 2022;3(11):774-91.e7.
17. Mundi PS, Dela Cruz FS, Grunn A, Diolaiti D, Mauguen A, Rainey AR, et al. A Transcriptome-Based Precision Oncology Platform for Patient–Therapy Alignment in a Diverse Set of Treatment-Resistant Malignancies. Cancer Discovery. 2023;13(6):1386-407.
18. Lee CY, The M, Meng C, Bayer FP, Putzker K, Müller J, et al. Illuminating phenotypic drug responses of sarcoma cells to kinase inhibitors by phosphoproteomics. Mol Syst Biol. 2024;20(1):28-55.

19. León TE, Rapoz-D'Silva T, Bertoli C, Rahman S, Magnussen M, Philip B, et al. EZH2-Deficient T-cell Acute Lymphoblastic Leukemia Is Sensitized to CHK1 Inhibition through Enhanced Replication Stress. Cancer Discov. 2020;10(7):998-1017.

20. Ngoi NYL, Pilié PG, McGrail DJ, Zimmermann M, Schlacher K, Yap TA. Targeting ATR in patients with cancer. Nature Reviews Clinical Oncology. 2024;21(4):278-93.

21. Dobzhansky T. Genetics of natural populations; recombination and variability in populations of Drosophila pseudoobscura. Genetics. 1946;31(3):269-90.

22. Farmer H, McCabe N, Lord CJ, Tutt AN, Johnson DA, Richardson TB, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. Nature. 2005;434(7035):917-21.

23. Bryant HE, Schultz N, Thomas HD, Parker KM, Flower D, Lopez E, et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. Nature. 2005;434(7035):913-7.

24. Muller FL, Colla S, Aquilanti E, Manzo VE, Genovese G, Lee J, et al. Passenger deletions generate therapeutic vulnerabilities in cancer. Nature. 2012;488(7411):337-42.

25. Zhao D, Lu X, Wang G, Lan Z, Liao W, Li J, et al. Synthetic essentiality of chromatin remodelling factor CHD1 in PTEN-deficient cancer. Nature. 2017;542(7642):484-8.

26. Kwok M, Davies N, Agathanggelou A, Smith E, Petermann E, Yates E, et al. Synthetic lethality in chronic lymphocytic leukaemia with DNA damage response defects by targeting the ATR pathway. Lancet. 2015;385 Suppl 1:S58.

27. Toledo LI, Murga M, Zur R, Soria R, Rodriguez A, Martinez S, et al. A cell-based screen identifies ATR inhibitors with synthetic lethal properties for cancer-associated mutations. Nat Struct Mol Biol. 2011;18(6):721-7.

28. Reaper PM, Griffiths MR, Long JM, Charrier JD, Maccormick S, Charlton PA, et al. Selective killing of ATM- or p53-deficient cancer cells through inhibition of ATR. Nat Chem Biol. 2011;7(7):428-30.

29. Middleton FK, Patterson MJ, Elstob CJ, Fordham S, Herriott A, Wade MA, et al. Common cancer-associated imbalances in the DNA damage response confer sensitivity to single agent ATR inhibition. Oncotarget. 2015;6(32):32396-409.

30. Shruti M, Marcus RB, Yone Phar L, Hannah A, Shruthi P, Ann H, et al. FET fusion oncoproteins disrupt physiologic DNA repair networks in cancer. bioRxiv. 2023:2023.04.30.538578.

31. Min A, Im SA, Jang H, Kim S, Lee M, Kim DK, et al. AZD6738, A Novel Oral Inhibitor of ATR, Induces Synthetic Lethality with ATM Deficiency in Gastric Cancer Cells. Mol Cancer Ther. 2017;16(4):566-77.

32. Menezes DL, Holt J, Tang Y, Feng J, Barsanti P, Pan Y, et al. A synthetic lethal screen reveals enhanced sensitivity to ATR inhibitor treatment in mantle cell lymphoma with ATM loss-of-function. Mol Cancer Res. 2015;13(1):120-9.

33. Knutson SK, Warholic NM, Wigle TJ, Klaus CR, Allain CJ, Raimondi A, et al. Durable tumor regression in genetically altered malignant rhabdoid tumors by inhibition of methyltransferase EZH2. Proc Natl Acad Sci U S A. 2013;110(19):7922-7.

34. Gadd S, Sredni ST, Huang C-C, Perlman EJ. Rhabdoid tumor: gene expression clues to pathogenesis and potential therapeutic targets. Laboratory Investigation. 2010;90(5):724-38.

35. Custers L, Khabirova E, Coorens THH, Oliver TRW, Calandrini C, Young MD, et al. Somatic mutations and single-cell transcriptomes reveal the root of malignant rhabdoid tumours. Nat Commun. 2021;12(1):1407.

36. Chun HE, Lim EL, Heravi-Moussavi A, Saberi S, Mungall KL, Bilenky M, et al. Genome-Wide Profiles of Extra-cranial Malignant Rhabdoid Tumors Reveal Heterogeneity and Dysregulated Developmental Pathways. Cancer Cell. 2016;29(3):394-406.

37. Pereira JD, Sansom SN, Smith J, Dobenecker MW, Tarakhovsky A, Livesey FJ. Ezh2, the histone methyltransferase of PRC2, regulates the balance between self-renewal and differentiation in the cerebral cortex. Proc Natl Acad Sci U S A. 2010;107(36):15957-62.

38. Zhang J, Ji F, Liu Y, Lei X, Li H, Ji G, et al. Ezh2 regulates adult hippocampal neurogenesis and memory. J Neurosci. 2014;34(15):5184-99.

39. Di Meglio T, Kratochwil CF, Vilain N, Loche A, Vitobello A, Yonehara K, et al. Ezh2 orchestrates topographic migration and connectivity of mouse precerebellar neurons. Science. 2013;339(6116):204-7.

40. Zhang M, Zhang Y, Xu Q, Crawford J, Qian C, Wang GH, et al. Neuronal Histone Methyltransferase EZH2 Regulates Neuronal Morphogenesis, Synaptic Plasticity, and Cognitive Behavior in Mice. Neurosci Bull. 2023;39(10):1512-32.

41. Flynn RL, Cox KE, Jeitany M, Wakimoto H, Bryll AR, Ganem NJ, et al. Alternative lengthening of telomeres renders cancer cells hypersensitive to ATR inhibitors. Science. 2015;347(6219):273-7.

42. Dorado García H, Pusch F, Bei Y, von Stebut J, Ibáñez G, Guillan K, et al. Therapeutic targeting of ATR in alveolar rhabdomyosarcoma. Nat Commun. 2022;13(1):4297.

43. Zeman MK, Cimprich KA. Causes and consequences of replication stress. Nat Cell Biol. 2014;16(1):2-9.

44. Schoppy DW, Ragland RL, Gilad O, Shastri N, Peters AA, Murga M, et al. Oncogenic stress sensitizes murine cancers to hypomorphic suppression of ATR. J Clin Invest. 2012;122(1):241-52.

45. Kwok M, Davies N, Agathanggelou A, Smith E, Oldreive C, Petermann E, et al. ATR inhibition induces synthetic lethality and overcomes chemoresistance in TP53- or ATM-defective chronic lymphocytic leukemia cells. Blood. 2016;127(5):582-95.

46. Karakashev S, Fukumoto T, Zhao B, Lin J, Wu S, Fatkhutdinov N, et al. EZH2 Inhibition Sensitizes CARM1-High, Homologous Recombination Proficient Ovarian Cancers to PARP Inhibition. Cancer Cell. 2020;37(2):157-67.e6.

47. Zhang X, Huo X, Guo H, Xue L. Combined inhibition of PARP and EZH2 for cancer treatment: Current status, opportunities, and challenges. Front Pharmacol. 2022;13:965244.

48. Ratz L, Brambillasca C, Bartke L, Huetzen MA, Goergens J, Leidecker O, et al. Combined inhibition of EZH2 and ATM is synthetic lethal in BRCA1-deficient breast cancer. Breast Cancer Res. 2022;24(1):41.

49. Campbell S, Ismail IH, Young LC, Poirier GG, Hendzel MJ. Polycomb repressive complex 2 contributes to DNA double-strand break repair. Cell Cycle. 2013;12(16):2675-83.

50. Chou DM, Adamson B, Dephoure NE, Tan X, Nottke AC, Hurov KE, et al. A chromatin localization screen reveals poly (ADP ribose)-regulated recruitment of the repressive polycomb and NuRD complexes to sites of DNA damage. Proc Natl Acad Sci U S A. 2010;107(43):18475-80.

51. Ito T, Teo YV, Evans SA, Neretti N, Sedivy JM. Regulation of Cellular Senescence by Polycomb Chromatin Modifiers through Distinct DNA Damage- and Histone Methylation-Dependent Pathways. Cell Rep. 2018;22(13):3480-92.

52. Piunti A, Rossi A, Cerutti A, Albert M, Jammula S, Scelfo A, et al. Polycomb proteins control proliferation and transformation independently of cell cycle checkpoints by regulating DNA replication. Nat Commun. 2014;5:3649.

53. Rondinelli B, Gogola E, Yücel H, Duarte AA, van de Ven M, van der Sluijs R, et al. EZH2 promotes degradation of stalled replication forks by recruiting MUS81 through histone H3 trimethylation. Nat Cell Biol. 2017;19(11):1371-8.

54. Pinto EM, Hamideh D, Bahrami A, Orr BA, Lin T, Pounds S, et al. Malignant rhabdoid tumors originating within and outside the central nervous system are clinically and molecularly heterogeneous. Acta Neuropathol. 2018;136(2):315-26.

55. Burr ML, Sparbier CE, Chan KL, Chan YC, Kersbergen A, Lam EYN, et al. An Evolutionarily
Conserved Function of Polycomb Silences the MHC Class I Antigen Presentation Pathway and
Enables Immune Evasion in Cancer. Cancer Cell. 2019;36(4):385-401.e8.
56. Mehdipour P, Marhon SA, Ettayebi I, Chakravarthy A, Hosseini A, Wang Y, et al. Epigenetic
therapy induces transcription of inverted SINEs and ADAR1 dependency. Nature.
2020;588(7836):169-73.
57. Ishak CA, Marshall AE, Passos DT, White CR, Kim SJ, Cecchini MJ, et al. An RB-EZH2
Complex Mediates Silencing of Repetitive DNA Sequences. Mol Cell. 2016;64(6):1074-87.
58. Ianevski A, Giri AK, Aittokallio T. SynergyFinder 2.0: visual analytics of multi-drug
combination synergies. Nucleic Acids Res. 2020;48(W1):W488-w93.
59. Stirling DR, Swain-Bowden MJ, Lucas AM, Carpenter AE, Cimini BA, Goodman A.
CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics. 2021;22(1):433.
60. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an
open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.
61. Mattar M, McCarthy CR, Kulick AR, Qeriqi B, Guzman S, de Stanchina E. Establishing and
Maintaining an Extensive Library of Patient-Derived Xenograft Models. Front Oncol. 2018;8:19.
62. Vardi Y, Ying Z, Zhang CH. Two-sample tests for growth curves under dependent right
censoring. Biometrika. 2001;88(4):949-60.

651 **Table 1:** List of oligonucleotides

gRNAs	
RB1 crRNA	5'- AGAGAGAGCUUGGUUAACUU -3'
Primers for amplification	
PGBD5 forward	5'- CTCTGGGTCAGACAATGTTCTTC -3'
PGBD5 reverse	5'- GCTTATTCTTCAGCGCATCCA -3'
GAPDH forward	5'- GATCATCAGCAATGCCTCCT -3'
GAPDH reverse	5'- GTCATGAGTCCTTCCACGATAC -3'

665
666
Figure 1: ATR inhibition is a compelling therapeutic target and overcomes TAZ
resistance in rhabdoid tumors: (**A-B**) metaVIPER analysis of rhabdoid tumor transcriptomes
for proteins within OncoTarget (**A**) and the complete metaVIPER protein set (**B**). (**C**) Inhibitors
rank-ordered by IC50 from Kuster et al. (**D**) DESeq2-normalized read counts of cells treated
with 10 µM TAZ versus equivalent volume of DMSO for 11 days. n=3 biological replicates per
condition. *p = 1.6E-6, **p = 1.3E-6, ***p = 1.2E-7 by two-sided Student's t-test. (**E**) G401 cells
treated with elimusertib for 4 days.

692
Figure 2: Combination EZH2 and ATR inhibition improves response in vitro: (**A**) Panel of
MRT and ES cell lines ordered left to right by decreasing response to TAZ monotherapy. Cells
were treated with the indicated monotherapy or combination for 11 days. Drug concentrations
used were: TAZ: 200 nM, elimusertib: 8 nM. We selected an elimusertib dose below its
monotherapy IC_{50} (for G401 cells) in order to visualize any additive effects upon combination with
TAZ. *p = 0.14, **p = 3.3E-3, ***p = 1.0E-3, ****p = 2.6E-4, #p = 6.1E-5, ##p = 3.0E-5 by two-
sided Student's t-test. All comparisons refer to TAZ + elimusertib vs. elimusertib conditions,
except for G401 cells, in which comparison is for TAZ + elimusertib vs. TAZ. n = 5 replicates per
condition. (**B-C**) Synergy plots for combination treatment with TAZ (left axis) and elimusertib (right
axis) for (**B**) G401 and (**C**) ES1 cells. Cells were treated at the indicated doses for 9 days and
analyzed for synergy using the Zero Interaction Potency (ZIP) model. (**D**) Representative images
of G401 cells treated with the indicated treatment for 7 days. Elimusertib was added on Day 5.

Doses used were 500 mM TAZ and 100 nM elimusertib. (**E**) Quantification of γH2AX fluorescence
relative to DAPI fluorescence. *p = 9.3E-3, **p = 6.2E-14, ***p = 1.5E-29 by two-sided Student's
t-test. n = 332 nuclei for DMSO, 404 for TAZ, 400 for elimusertib, 257 for TAZ + elimusertib.

728
729
Figure 3: CHK1 inhibition does not induce replication stress or synergize with TAZ: (**A**)
DESeq2-normalized read counts of cells treated with 10 µM TAZ versus equivalent volume of
DMSO for 11 days. n=3 biological replicates per condition. (**B**) Dose-response curves of G401
cells treated with the CHK1 inhibitor SRA737 for 9 days. (**C**) Western blot assaying replication
stress as measured by RPA phosphorylation at S4/8 and T21. Camptothecin treatment (1.5 µM)
for 2 h was used as a positive control for replication stress. Autophosphorylation of CHK1 at S296
was used to confirm CHK1 inhibition. Cells were pre-treated with 10 µM TAZ or DMSO for 9 days.
Cells were then split and additionally treated with SRA737 (3 µM) or equivalent volume of DMSO
for 2 days. (**D**) Cells treated with 10 µM TAZ or DMSO for 11 days do not express MYCN protein.
MYCN-amplified neuroblastoma cell line IMR5 was used as a positive control for MYCN
expression.

741
742
743

744
Figure 4: TAZ-induced DNA damage is PGBD5-dependent: (**A**) RT-qPCR showing *PGBD5*
745
expression vs. *GAPDH* in G401 cells with the indicated shRNA. *p = 4.7E-3, **p = 1.2E-3. (**B**)
746

Quantification of γH2AX fluorescence relative to DAPI fluorescence in all nuclei. $*p = 8.6E-3$, $**p$

$= 3.9E-8$, $***p = 4.4E-8$, $****p = 1.1E-12$, $\# p = 4.9E-34$, $\#\# p = 4.0E-47$ by two-sided Student's t-

test. (**C**) Representative images of cells quantified in (B). (**D**) Quantification of γH2AX

fluorescence relative to DAPI fluorescence in nuclei with punctate γH2AX staining. $*p = 3.6E-2$,

$**p = 2.7E-5$, $***p = 2.4E-5$, $****p = 1.1E-15$, $\#p = 5.6E-51$, $\#\#p = 1.7E-81$. For B-C, n for shGFP

cells is 539 for DMSO, 321 for TAZ, 491 for elimusertib, 418 for TAZ + elimusertib. n for shPGBD5-

1 is 497 for DMSO, 390 for TAZ, 354 for elimusertib, 277 for TAZ + elimusertib. n for shPGBD5-

3 is 703 for DMSO, 418 for TAZ, 554 for elimusertib, 317 for TAZ + elimusertib. (**E**) Proportion of

nuclei with pan-nuclear γH2AX staining per field. Each dot represents one field. $*p = 9.2E-2$, $**p$

$= 1.6E-2$, $***p = 9.0E-4$ n for shGFP cells is 22 fields for DMSO, 22 for TAZ, 22 for elimusertib,

28 for TAZ + elimusertib. n for shPGBD5-1 is 22 for DMSO, 22 for TAZ, 22 for elimusertib, 20 for

TAZ + elimusertib. n for shPGBD5-3 is 22 for DMSO, 22 for TAZ, 20 for elimusertib, 22 for TAZ +

elimusertib.

Figure 5: TAZ plus elimusertib improves therapeutic response in vivo: (**A**) Tumor growth curves for 5 mouse PDXs treated with the indicated drug regimen. n = 20 mice for vehicle and elimusertib-treated groups, n = 21 for TAZ and TAZ + elimusertib-treated groups. Vardi *U*-test *p* = 2.3E-2 and 0.20 for combination vs. elimusertib or TAZ, respectively. (**B**) Kaplan-Meier curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs in panel C. Mean survival is 51 days (95% CI: 42-60 days) for elimusertib, 68 days (95% CI: 53-82 days) for TAZ to 100 days (95% CI: 74-124 days) for the combination. Log-rank test *p* = 5.8E-4 and 3.8E-2 for combination versus elimusertib or TAZ, respectively. (**C**) Tumor growth curves for the HYMAD_EPIS_X0004aS1 PDX model treated with the indicated drug regimen. Vardi *U*-test *p* = 2.0E-4 for combination versus elimusertib or TAZ. n = 14 mice per treatment group. (**D**) Kaplan-Meier curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the PDXs in panel C. Log-rank test *p* = 6.2E-3 and 6.3E-5 for combination versus elimusertib or TAZ, respectively. (**E**) Tumor growth curves for the SOMWR_EPIS_X00013aS1 PDX model. Vardi *U*-test *p* = 1.0E-3 and 6.0E-2 for combination versus elimusertib and TAZ, respectively. *n* = 8 mice for all groups. (**F**) Kaplan-Meier curves showing tumor-free survival (defined as tumor volume ≤ 1,000 mm^3) for the SOMWR_EPIS_X00013aS1 PDX model; Mean survival is 90 days (95% CI: 83-97 days) for combination versus 80 days (95% CI: 70-90 days) for TAZ and 45 days (95% CI: 37-52 days) for elimusertib. Log-rank test *p* = 9.5E-5 and 6.0E-2 for combination versus elimusertib and TAZ, respectively. (**G**) Image of representative tumors extracted from mice in C-D and on Day 52 of treatment (**H**) their corresponding weights. *p = 6.7E-3 by two-sided Student's t-test.