Postnatal hearing development limits early parent-to-offspring vocal communication and gates sensorimotor learning in the zebra finch

Tommi Anttonen, Jakob Christensen-Dalsgaard, Coen P. H. Elemans

Affiliation

Sound Communication and Behaviour Group, Department of Biology, University of Southern Denmark, Odense, Denmark

Correspondence

Tommi Anttonen, t.anttonen@outlook.com; Coen P.H. Elemans, coen@biology.sdu.dk

ABSTRACT

Hearing is critical for vocal communication, and in birds its onset is linked to egg hatching. Hearing develops after hatching in early hatching (altricial) species, while already functional in the egg in later hatching (precocial) species. Recent work suggests that altricial zebra finches exhibit parent-to-embryo acoustic communication, requiring functional, high-frequency hearing already in the egg. Yet, the early postnatal development of their auditory sensitivity remains unknown. By measuring hearing and vibration-detection capacity during postnatal development in zebra finches, we show that zebra finch hatchlings are deaf even to loud, broadband sounds. Auditory nerve sensitivity develops gradually in a low-to-high frequency manner. Auditory nerve responses show adult-like properties only after 20 days from hatching. Our data provide evidence against the plausibility of early parent-to-offspring acoustic communication in zebra finches and suggest that hearing functionality may set the onset of vocal learning in zebra finches.

INTRODUCTION

The ability to detect sound is critical for vocal communication, vocal learning, and maintenance of learned vocalizations (Konishi, 1965; Marler *et al.*, 1972; Nordeen and Nordeen, 1992; Heaton, Dooling and Farabaugh, 1999; Zevin, Seidenberg and Bottjer, 2004). In terrestrial vertebrates hearing improves during development through growth and maturation of the inner and middle ear. In birds, the onset of hearing capacity is often linked to the timing of egg hatching, with many precocial bird species hatching later than altricial birds. For example, the average incubation time of precocial chickens and ducks is 21 and 28 days while for altricial zebra finches and budgerigars it is 14 and 18 days, respectively (Hamburger and Hamilton, 1951; Montgomery, Burke and Byers, 1978; Ar and Tazawa, 1999; Murray *et al.*, 2013).

The hearing of altricial birds is widely believed to mature after hatching with little to no functional hearing during the embryonic, pre-hatching stages (Woolley, 2017). In contrast, precocial bird species show auditory function already while still being inside the egg (Gos, 1933; Gottlieb, 1965; Konishi, 1973; Saunders, Coles and Gates, 1973). The embryonic onset of hearing function in precocial birds enables embryo-to-embryo (Vince, 1966; Woolf, Bixby and Capranica, 1976; Schwagmeyer *et al.*, 1991; Noguera and Velando, 2019), and parent-to-embryo vibroacoustic communication where it plays a crucial role in imprinting on species-specific sounds (Grier, Counter and Shearer, 1967; Hess, 1972; Gottlieb, 1979). Embryonic sensitivity to vibroacoustic cues mediates synchronization and induction of hatching in other precocial animals, including frogs (Warkentin, Caldwell and McDaniel, 2006), snakes (Aubret *et al.*, 2016), turtles (Doody *et al.*, 2012; McGlashan *et al.*, 2015), and crocodiles (Vergne and Mathevon, 2008).

Surprisingly, recent research suggests that parent-to-embryo vibroacoustic communication may also occur in altricial birds (Colombelli-Négrel *et al.*, 2012, 2016; Colombelli-Négrel, Hauber and Kleindorfer, 2014; Kleindorfer *et al.*, 2024). This challenges the previous distinction between precocial and altricial birds in terms of functional embryonic hearing. In additional support of a less rigid distinction, a recent series of studies suggest that altricial zebra finch embryos can detect sound while still being inside the egg. When in a hot environment, zebra finch parents emit "heat calls" with a fundamental frequency of 7-10 kHz that have been proposed to epigenetically guide their unhatched offspring to adapt to high

temperatures (Mariette and Buchanan, 2016, 2019; Katsis *et al.*, 2018, 2021, 2023; Mariette *et*
al., 2018; Pessato *et al.*, 2020; Udino *et al.*, 2021; Udino and Mariette, 2022). The presence of
such parent-to-embryo communication via the auditory modality crucially requires that zebra
finch embryos can detect these sound stimuli in the egg. Furthermore, under the assumption that
no sudden loss of hearing function occurs around hatching, these studies also imply that zebra
finches must be able to hear upon hatching.

The above predictions contradict the prevailing understanding that zebra finch hearing
develops predominantly after hatching. The earliest measurements of hearing sensitivity in zebra
finches are at 10 days-post-hatch (DPH) and indicate a reduced hearing sensitivity of 25 dB
sound pressure level (SPL) compared to 20 DPH when their hearing sensitivity is comparable to
adult levels (Amin, Doupe and Theunissen, 2007). The adult frequency sensitivity ranges from
0.25-to-8 kHz with the highest sensitivity around 3 kHz (Okanoya and Dooling, 1987; Amin,
Doupe and Theunissen, 2007; Yeh, Rivera and Woolley, 2023). If embryonic detection of heat
calls relies on the auditory system, either through air-borne sound pressure waves on the
tympanum or via bone conduction (Sakabe, Arayama and Suzuki, 1969; Mudry and Tjellström,
2011), then the onset of zebra finch hearing must occur earlier than currently assumed. As an
alternative route of detection, heat call acoustic signals could induce egg vibrations that embryos
sense as vibrotactile signals. However, both hearing capacity via air or bone conduction remain
unknown during the early development in zebra finches. To test the hypothesis that zebra finches
have functional hearing prior to hatching and possibly even before, it is imperative to conduct
measurements of hearing sensitivity at early postnatal ages.

Hearing plays a crucial role in another key behaviour in songbirds, vocal learning which
is the ability to learn song by imitation. Juvenile male zebra finches listen to the song of adult
tutors and form a memory template of it during a restricted sensory learning phase (Immelmann,
1967). This phase is followed by a sensorimotor phase, where the birds practice their own
vocalizations through trial-and-error learning until they match their internal song template. In the
zebra finch, the dominant model for studying neural circuitry and behaviour underlying this
complex motor learning process, the sensory and sensorimotor phase partly overlap and start on
25 DPH (Immelmann, 1967; Böhner, 1990; Roper and Zann, 2006; Braaten, 2010; Gobes,
Jennings and Maeda, 2019). Experimentally priming early sensorimotor learning can push the

sensory phase to start around 20 DPH (Leitão and Gahr, 2024). The onset of the sensory phase

thus closely coincides with hearing sensitivity maturation, which raises the question if hearing

maturation constrains the onset of sensory and sensorimotor learning in zebra finches. However,

our current understanding is limited by the lack of data on the maturation of hearing in songbirds

(Woolley, 2017).

Here we test the hypothesis that zebra fiches possess auditory function at early post-hatch

ages, by measuring air conduction- (AC-) and bone conduction-induced (BC-) auditory

brainstem responses (ABRs). ABR recordings allow hearing threshold estimation when

behavioural training is challenging (Heffner, Koay and Heffner, 2008). Additionally, we

measured the vibrations that air conducted sounds can induce on zebra finch eggs. We show that

the earliest AC-ABRs evoked by intense, broadband sounds could not be detected on 2 DPH, but

earliest at 4 DPH. Responses to frequency specific sounds develop later and gradually in a low-

to-high frequency manner. AC-ABR thresholds remain high throughout development for

frequencies associated with heat calls. AC-ABR sensitivity for clicks matures to adult-like levels

at 10 DPH, but the response amplitude continues to increase beyond 20 DPH. Head vibrations

surprisingly do not elicit BC-ABRs in zebra finches. Airborne sounds induce low amplitude

vibrations in eggs that are far below avian or mammalian vibrotactile detection thresholds.

Together, our findings provide evidence against the hypothesis that zebra finch embryos and

early hatchlings can detect vibroacoustic stimuli via air and bone conduction hearing or via

vibrotactile senses. Our results also suggest that neural responses to sounds at the level of the

auditory nerve continue to mature past 20 DPH, which suggests that hearing functionality may

set the onset of vocal learning in zebra finches.

RESULTS

AC-ABRs to loud clicks appear during the first postnatal week

To assess the development of zebra finch hearing function we quantified the responses of auditory pathway neurons to auditory stimuli by recording AC-ABRs (Fig. 1a) throughout postnatal development. First, to evaluate overall hearing sensitivity we presented loud yet physiologically relevant (95 dB SPL) clicks that stimulate the auditory system across a wide frequency spectrum (Fig. 1b). Clicks activate most auditory neurons nearly simultaneously and result in prominent AC-ABRs (Jones, Jones and Shukla, 1997), thereby providing maximal sensitivity for threshold detection.

None of the tested animals displayed click-evoked AC-ABRs at 2 DPH (both with detection at signal-to-noise ratio of 2 (S/N > 2) and with detailed visual inspection; n=0/15; Fig. 1c,d). However, during the first postnatal week, click-evoked AC-ABRs became observable in some animals (S/N > 2; 4 DPH, n = 5/13; 6 DPH, n = 9/10; Fig. 1d). Subsequently, from the second postnatal week onwards, all tested animals consistently demonstrated click-evoked AC-ABRs (S/N > 2; 8 DPH, n = 11; 10 DPH, n = 11; 20 DPH, n = 7; 25 DPH, n = 8; Adult, n = 8; Fig. 1d). Thus, sounds at loud yet physiologically relevant SPLs do not evoke AC-ABRs in young zebra finches in the days after hatching.

Detection of click-evoked AC-ABRs is adult-like at 10 DPH

To estimate thresholds of hearing over postnatal age, we recorded series of click-evoked AC-ABRs at 5 dB downward steps from 95 dB SPL (Fig. 1e). Hearing thresholds are strongly affected by the age of the animal ($F(6)=25.22$, $p=4.93^{-14}$; Fig. 1f, Supplementary Table 1). Click-evoked AC-ABR thresholds are significantly higher than adult-like levels (40.6 ± 4.0 dB SPL) in animals younger than 10 DPH, but not in older animals (4 DPH vs adult, $t = 8.16$, $p = 3.27*10^{-10}$; 6 DPH vs adult, $t = 5.91$; $p = 1.38*10^{-6}$; DPH 8 vs adult, $t = 3.37$, $p = 0.008$; 10 DPH vs adult, $t = 1.23$, $p = 1$; 20 DPH vs adult, $t = -0.13$, $p = 1$; 25 DPH vs adult, $t = -1.29$, $p = 1$; p-values are Bonferroni-corrected; Fig. 1f, Supplementary Table 1). Click thresholds are as high as 80.6 ± 1.6 dB SPL on 4 DPH. Therefore, zebra finch hearing sensitivity rapidly increases during the first two postnatal weeks.

Anttonen et al., 2024, Figure 1

Figure 1. Click-evoked AC-ABRs appear and subsequently mature after hatching in zebra finches. a ABR

recording setup. **b** Used sound stimuli. *Red*, phase inversed signal. **c** Representative AC-ABRs to 95 dB SPL clicks

at tested DPH timepoints. The first negative peak (*I*) and the following trough are designated as wave I. Noise RMS

is determined from the *red area* and signal RMS from the *blue area*. **d** Percentage of animals that exhibited a

detectable (S/N > 2) click-induced AC-ABR. **e** Detection thresholds were detected with a set of clicks with

decreasing SPL (5 dB steps). *Arrow* indicates the determined threshold for this 8 DPH example. **f** Average click

thresholds at different DPH ages. Orange box indicates heat call playback levels used by Katsis et al., (2018). **g**

Click-evoked AC-ABR wave I amplitude and **h** latency over SPL. Age groups are color coded in c,d,e,g,h. *Gray*

areas in f-h = s.e.m.

AC-ABR waveforms change past the sensory learning onset

The click-evoked AC-ABR consists of multiple waves that reflect the subsequent activations of various neural populations of the auditory pathway. The first wave (wave I in Fig. 1c) corresponds to the activity of the primary auditory neurons (Buchwald and Huang, 1975). To study the postnatal maturation of this synchronized neural output, we measured changes in both amplitude and latency of wave I across developmental age.

In adults, the amplitude of wave I increases with the SPL of the click (Fig. 1c,g) as it has been reported for humans, mice, and budgerigars (Hecox and Galambos, 1974; Henry and Haythorn, 1978; Brittan-Powell and Dooling, 2004). We found that the age of the animal significantly affects both wave I amplitude directly and the relation between SPL and wave I amplitude (Age-SPL interaction compared to only additive effects of age and SPL: $\chi^2(6)=164.23$, $p=7.52*10^{-33}$, additive effects of age and SPL compared to only SPL: $\chi^2(6)=79.43$, $p=4.69*10^{-15}$, SPL only compared to null model: $\chi^2(1)=438.12$, $p=2.78*10^{-97}$, Fig. 1g, Supplementary Table 1). The steepness of the amplitude-SPL slope is significantly different between adults and younger animals but not between adult and those aged 25 DPH (4 DPH vs adult, $t = -5.40$, $p = 1.50*10^{-6}$; 6 DPH vs adult, $t = -5.65$, $p = 1.8910^{-6}$; DPH 8 vs adult, $t = -6.30$, $p = 3.79*10^{-7}$; 10 DPH vs adult, $t = -5.53$, $p = 6.53*10^{-6}$; 20 DPH vs adult, $t = -3.00$, $p = 0.03$; 25 DPH vs adult, $t = -0.69$, $p = 1$; p-values are Bonferroni-corrected; Fig.- 1g, Supplementary Table 1).

The latency of wave I decreases with increasing click SPL (Fig. 1c,h) consistent with findings in humans, mice, and budgerigars (Hecox and Galambos, 1974; Henry and Haythorn, 1978; Brittan-Powell and Dooling, 2004). Similar to amplitude, the age of the animal significantly affects the relation between dB SPL and wave I latency (Age-SPL interaction compared to only additive effects of age and SPL: $\chi^2(6) = 116$, $p = 1.31*10^{-22}$; additive effects of age and SPL compared to only SPL, $\chi^2(6) = 109$, $p = 2.76*10^{-21}$; SPL only compared to null model: $\chi^2(1) = 754$, $p = 4.81*10^{-166}$; Fig. 1h, Supplementary Table 1). The observed latencies are significantly longer during early postnatal development (4-to-10 DPH) when compared to latencies observed in adults. No significant differences are observed between the latencies of adults and animals older than 10 DPH (4 DPH vs adult, $t = 12.56$, $p = 3.41*10^{-19}$; 6 DPH vs adult, $t = 10.06$, $p = 1.49*10^{-13}$; 8 DPH vs adult, $t = 7.24$, $p = 1.16*10^{-8}$; 10 DPH vs adult, $t = 6.18$, $p = 5.95*10^{-7}$; 20 DPH vs adult, $t = 2.23$, $p = 0.18$; 25 DPH vs adult, $t = 1.67$, $p = 0.61$; p-values are Bonferroni-corrected; Fig. 1h, Supplementary Table 1).

The observed changes in the synchronized output of the auditory nerve indicate that peripheral

encoding of sounds in zebra finches mature during the first three weeks post hatching and

resemble the responses seen in adults at 25 DPH.

Hearing sensitivity to high frequency sounds develops last

While clicks provide general sensitivity thresholds, they do not inform about frequency

sensitivity of the auditory system. To study frequency-specific hearing sensitivity we tested

whether loud (95 dB SPL) tone bursts centred at specific frequencies between 0.25-to-8 kHz

elicit detectable AC-ABRs. Each stimulus was 25 ms in length so that the lowest studied

frequency stimulus would undergo at least 6 cycles during this period. The rise/fall time of the

stimulus was set to 5 ms to minimize spectral splatter that compromises the frequency specificity

of the stimulus (Lauridsen, Brandt and Christensen-Dalsgaard, 2021).

Tone burst-evoked AC-ABRs are detectable for the first time at 8 DPH at 1 kHz (Fig. 2a

and Supplementary Figure 1a). As observed with click stimuli, the amplitude of the tone burst-

induced AC-ABRs appears to increase (Fig. 2a and Supplemental Fig. 1b; Age as a predictor of

amplitude compared to null model: $\chi^2(3) = 6.11$, p = 0.11, Supplementary Table 2) while the

latency significantly decreases during postnatal development (Fig. 2a and Supplementary Figure

1c; Age as a predictor of latency compared to null model: $\chi^2(3) = 33.72$, p = $2.27*10^{-7}$,

Supplementary Table 2). Latencies are significantly longer at 8-to-10 DPH (Latencies at 1 kHz:

8 DPH, 6.60 ± 0.77 ms; 10 DPH, 5.01 ± 0.40 ms) and are adult-like at 20-to-25 DPH (Latencies

at 1 kHz: 20-to-25 DPH, 3.34 ± 0.18 ms; adult, 3.67± 0.30 ms; 8 DPH vs adult, t = 5.95, p =

$1.87*10^{-6}$; 10 DPH vs adult, t = 4.52, p = 0.0004; 20-to-25 DPH vs adult, t = 1.16, p = 0.81; p-

values are Bonferroni-corrected; Supplementary Table 2). During the following weeks of

postnatal development, tone burst-induced AC-ABRs became detectable at 0.5-to-4 kHz

(Supplementary Fig. 1a). No responses were detected at higher frequencies (Fig. 2b and

Supplementary Fig. 1a). Taken together, we observed that maturation of the hearing sensitivity in

zebra finches progresses in a low-to-high frequency order, as is also observed in mice and cats

(Ehret, 1976; Ehret and Romand, 1981) and in other birds (Brittan-Powell and Dooling, 2004;

Kraemer *et al.*, 2017).

Figure 2. Frequency dependent hearing maturation over postnatal development. a Representative AC-ABRs to

1 kHz and **b** 8 kHz tone burst stimuli over postnatal development. **c** Zebra finch ABR thresholds obtained with tone

bursts (*solid lines*, this study) with tone pips (*dashed line*., this study). Behavioural hearing thresholds recorded by

Okanoya and Dooling (1987) are plotted with a *dotted line*. *Orange box* displays the playback level of heat calls

used by Katsis et al. (2018). Shaded areas = s.e.m.

Limited hearing sensitivity at heat call frequencies

To test how readily the adult zebra finch auditory system detects heat calls, we analysed their

hearing sensitivity for higher frequency sound stimuli in more detail. Importantly, tone bursts

evoked AC-ABRs only in about half of the adult zebra finches (Supplementary Fig. 1a) and the

detection thresholds of > 60 dB SPL were high. This indicates that while being more frequency

specific, tone bursts failed to recruit zebra finch auditory neurons to fire concurrently enough to

generate detectable AC-ABRs. A significant effect of age on thresholds could not be observed

(Age as a predictor of threshold compared to null model: $\chi^2(3) = 1.64$, p = 0.65, Supplementary

Table 2), likely due to the low statistical power caused by the limited amount of observed tone

burst-evoked AC-ABRs.

Next, we designed sound stimuli that aim to recruit more auditory neurons to fire

simultaneously. We shortened the stimulus rise/fall time to 1 ms and the plateau time to 3 ms.

The obtained 5 ms-long tone pips (Fig. 1b) have a broader spectral distribution of acoustic

energy when compared to tone bursts (Lauridsen, Brandt and Christensen-Dalsgaard, 2021).

Compared to tone burst-induced AC-ABRs, the SPL matched tone pips produced much more

prominent AC-ABRs (Fig. 2c and Supplementary Fig. 2a). Tone pip-evoked AC-ABRs were

readily detected in all tested adult zebra finches around their most sensitive hearing range of 1-

to-4 kHz (Okanoya and Dooling, 1987) (Supplementary Fig. 2b). Detection thresholds for tone

pips are ~20 dB SPL more sensitive than those of tone bursts (Additive effects of stimulus type

and frequency as predictors of threshold compared to stimulus type alone $\chi^2(1) = 6.84$, p = 0.009;

stimulus type as a predictor of threshold compared to null model: $\chi^2(1) = 10.27$, p = 0.001,

Supplementary Table 2). Importantly, the tone pip-based audiogram resides ~10-to-20 dB SPL

above the known behavioural thresholds (Okanoya and Dooling, 1987) (Fig. 2c) as one would

expect based on previous studies (Stapells, 2000). Based on these audiograms, we can conclude

that heat calls at playback levels of 62-to-67 dB SPL at the ear (Katsis *et al.*, 2018) are just on or

below the limit of what adult zebra finches can readily detect.

Bone conduction does not mediate hearing in zebra finches

After evaluating the development of air conduction-based hearing, we next studied whether bone

conduction can mediate hearing during early postnatal development in zebra finches. By using

the same electrode configuration as for AC-ABRs, we tested whether head vibrations can evoke

BC-ABRs.

In adult zebra finches, tone pip- and tone burst-like head vibrations at high acceleration

levels gave rise to large recordable potentials at frequencies below 2 kHz (Figs. 3a and 4a) and to

smaller, ABR-like potentials at 2-to-3 kHz (Figs. 3b and 4a). Aiming first to resolve the

unknown source of these responses, we tested whether a simultaneous air conduction white noise

sound stimulus at 95 dB SPL masks the vibration-induced signals. Interestingly, responses below

2 kHz were not sensitive to noise masking (Fig. 3c), which strongly suggests that they did not

originate from the auditory system.

Figure 3. Characteristics of potentials generated by head vibrations in zebra finches. *Red insets* in a-f show the start of the response at a higher magnification. **a** Tone pip vibrational stimuli induce large recordable potentials at frequencies below 2 kHz. The visualised example is at 0.5 kHz. **b** At 2-to-3 kHz, small ABR-like potentials can be recorded. This example is at 3 kHz. **c** Concurrent 95 dB SPL white noise fails to mask the large potentials but **d** eliminates the small ABR-like potentials. **e** Detaching the vibration probe from the head eliminates the large potentials while the ABR-like response remains unaffected. **g** Both response types disappear after the animal dies **h** the vibration exciter produces audible sounds at high acceleration levels.

In contrast, the head vibration-induced ABR-like responses were effectively masked by noise (Fig. 4d, n = 3) as expected for BC-ABRs. To test if head vibrations are required for generating these potentials, we lifted the vibration exciter a few millimeters to prevent direct head contact. While this adjustment eliminated the potentials with stimuli below 2 kHz (Fig. 3e), the smaller ABR-like potentials persisted (Fig. 3f, n = 3). Furthermore, the observed potentials require the animal to be alive to occur, indicating that they originate from the animal and are not artefacts of e.g. electrode vibration (Fig. 3g; n = 3). Taken together, we conclude that the ABR-like

responses at frequencies above 2 kHz are driven by something else than head vibrations, while
responses below 2 kHz are driven by head vibrations.

While searching for the origin of noise-sensitive responses described above, we noticed
that the vibration exciter used by us and others (Chhan, McKinnon and Rosowski, 2017)
produces audible sounds in addition to vibration at high stimulus intensities. When driven for
example at 3 kHz and above 85 dB re 0.1 mm s^{-2}, the exciter produces sounds above 40 dB SPL
at the animals' ear that are of sufficient amplitude to evoke AC-ABRs (Fig. 3h). Therefore, we
suggest that the observed ABR-like responses during vibration excitation are AC-ABRs evoked
by exciter-generated sounds and not head vibration-evoked BC-ABRs.

Responses to head vibrations mature postnatally and require high acceleration

To study the postnatal development of head vibration-induced potentials, we induced head
vibrations with tone bursts in zebra finches at frequencies between 0.25-to-8 kHz. We used tone
bursts as they have a lower capability to induce AC-ABRs than tone pips, minimizing the AC-
ABRs evoked by the audible sounds originating from the exciter. Potentials induced by head
vibrations can be detected only with stimuli below 2 kHz (Fig. 4a,c). These responses typically
consist of two negative deflections (n_I and n_{II}) and of a single positive one between them (p_I).
The response waveform changed during postnatal development (Fig. 4b) with the number of
animals displaying these responses increasing as they matured (Fig. 4c). Such responses are
detected with 0.25 kHz stimulation as early as 2 DPH, followed with 0.5 kHz at 6 DPH, and
finally at 1 kHz in three-week-old animals (Fig. 4c). A significant effect of age on thresholds
could be observed (Age as a predictor of threshold compared to null model: $\chi^2(6) = 23.94$, p =
0.0005, Supplementary Table 3). The acceleration threshold levels reduce over postnatal
development (2 DPH vs adult, t = 2.74, p = 0.051; 4 DPH vs adult, t = 3.60, p = 0.005; 6 DPH vs
adult, t = 2.74, p = 0.051; 8 DPH vs adult, t = 4.13, p = 0.003; 10 DPH vs adult, t = 3.68, p =
0.006; 20-to-25 DPH vs adult, t = 3.73, p = 0.012; p-values are Bonferroni-corrected;
Supplementary Table 3) and are around 80 dB re 0.1 mm s^{-2} in adults (Fig. 4d). Taken together,
head vibration-induced potentials mature during postnatal development in a low-to-high
frequency manner.

Figure 4. Postnatal maturation of head vibration-induced potentials (HVIPs). a-c relate to tone burst vibration stimuli delivered on the head at 100 dB re 0.1 mm/s² **a** Representative, adult HVIPs evoked by intense 0.25-to-8 kHz tone bursts. **b** Representative HVIPs induced by intense 0.5 kHz tone bursts at various DPH timepoints. **c** Percentage of animals displaying detectable HVIPs (S/N > 2) at various ages (DPH). Number of tested animals is displayed above the bars. *Gray bars* indicate animals without signal. **d** Acceleration thresholds for HVIPs at various DPH timepoints. Shaded area = s.e.m.

Sound-induced egg vibrations are below detectable levels

Our results do not support the hypothesis that embryonic zebra finches can use their immature auditory system to detect high frequency heat calls. We therefore tested the hypothesis that airborne sounds can induce zebra finch eggs to vibrate at high enough velocities to be detected via vibrotactile senses (Mountcastle, LaMotte and Carli, 1972; Hörster, 1990). We played sound frequency sweeps to zebra finch eggs while simultaneously recording the sound-induced vibrations of the egg with a laser vibrometer (Fig. 5a). The vibration velocity transfer function shows that zebra finch eggs vibrate at velocities around -85 to -55 dB re 1 mm s^{-1} when exposed to a 94 dB SPL frequency sweep between 0.25-to-10 kHz (Fig. 5b). Within the frequency ranges associated with most sensitive hearing of zebra finches (1-to-4 kHz, *green*) and with heat calls (7-to-10 kHz, *red*), sound exposures at 94 dB SPL induce average vibration velocities of -72.4 ± 0.8 and -69.2 ± 0.9 dB re 1 mm s^{-1}, respectively (standard deviation, Fig. 5c). We calculated the corresponding vibration velocity levels induced by SPLs used during experimental heat call playbacks to zebra finch eggs (67 dB SPL) (Katsis *et al.*, 2018) to be between -99.4 and -96.2 dB re 1 mm s^{-1}, respectively (Fig. 5d). The obtained vibration velocities are approximately 10-to-40 dB lower than typical vertebrate bone conduction thresholds (Håkansson, Tjellström and Rosenhall, 1985; Sakai, Karino and Kaga, 2006; Chhan, McKinnon and Rosowski, 2017), 80 dB

327 lower than what human (Mountcastle, LaMotte and Carli, 1972) and bird (Hörster, 1990)

328 vibrotactile senses typically can detect, and 80 dB lower than the threshold levels to generate

329 HVIPs in zebra finches (Fig. 5d). In conclusion, air-borne sounds at physiologically relevant

330 SPLs are not able to induce zebra finch eggs to vibrate at high enough velocities to be detected

331 by avian (or mammalian) vibrotactile senses or by bone conduction-mediated hearing.

Anttonen et al., 2024; Figure 5

Figure 5. Intense sound stimulation induces zebra finch eggs to vibrate at very low velocities. a Setup used to
measure sound induced vibrations of eggs. **b** Average (N=5) vibration velocity transfer function of a zebra finch egg
during a 94 dB SPL sound sweep between 0.25-to-10 kHz. *Gray area* marks the standard deviation. *Green* and *red*
areas display the frequency windows related to the best sensitivity of zebra finch hearing (*song*) and to heat calls,
respectively. **c** The average egg vibration velocities induced by a 94 dB SPL sounds at frequencies associated with
song (*green*) and with heat calls (*red*). Error bars = s.d. **d** The average egg vibration velocities caused by a 67 dB
SPL sounds plotted together with thresholds for human BC-ABR (Håkansson, Tjellström and Rosenhall, 1985),
mouse BC-ABR (Chhan, McKinnon and Rosowski, 2017) and zebra finch HVIPs.

DISCUSSION

Our data challenge previous findings that zebra finch parents can use sound to communicate with their hatchlings and embryos through calls with high fundamental frequencies of 7-to-10 kHz. The lack of ABRs even to loud stimuli show that zebra finch hatchlings are deaf right after hatching (Fig. 1). Furthermore, the progressively increasing hearing sensitivity over postnatal development (Fig. 1) strongly suggests that zebra finch embryos should have even lower hearing sensitivity than hatchlings. Even in precocial bird species that show auditory function already during embryonic development, auditory thresholds remain very high until a few days before hatching (Saunders, Coles and Gates, 1973; Saunders, Gates and Coles, 1974; Rebillard and Rubel, 1981; Lippe, 1994). For example, evoked responses to very loud sounds (>100 dB SPL) have been obtained at around E12 in precocial chickens (21 embryonic days in total in chickens) (Saunders, Coles and Gates, 1973). However, ambient sound levels evoke responses much later around E16 (Jones, Jones and Paggett, 2006). Importantly, the above discussed measurements of embryonic hearing are often overestimations of the natural hearing sensitivity since they involve opening the egg and clearing the auditory meatus from fluids, and even drainage of fluids from the middle ear (Rebillard and Rubel, 1981). Future experiments with non-invasive techniques are required to estimate embryonic hearing capabilities of zebra finch embryos.

Our data demonstrates that hearing functionality in zebra finches gradually increases over postnatal development. The gradual increase of ABR sensitivity over postnatal development is consistent with previous observations in zebra finches, and another songbird, the pied flycatcher (Aleksandrov and Dmitrieva, 1992) as well as in other altricial bird species such as budgerigars (Brittan-Powell and Dooling, 2004), and barn owls (Köppl and Nickel, 2007; Kraemer *et al.*, 2017). It is important to note that ABR recordings are minimally invasive but crude measurements of hearing sensitivity and do not inform about the underlying processes that are behind the observed development of hearing sensitivity. These processes have been studied well in chickens. For example, the onset of evoked responses to very loud sounds coincides with synapse formation on hair cells of the chicken inner ear (Saunders, Coles and Gates, 1973; Cohen and Fermin, 1978; Caus Capdevila, Sienknecht and Köppl, 2021). The fluid unloading and maturation of the middle ear structures have been suggested to cause some of the significant improvements in auditory sensitivity during chicken development (Saunders, Coles and Gates,

1973; Cohen, Hernandez and Saunders, 1992).When these hallmark events occur in zebra finch

development remains unknown. Future work on the development of the outer, middle, and inner

ear of the zebra finch is thus needed to offer insights into the physiological basis for the observed

postnatal improvement of hearing sensitivity.

During early postnatal development prior to 10 DPH, the hearing sensitivity of zebra

finches is limited to frequencies around 1 kHz, and sensitivity for higher frequencies develops

progressively afterwards. The progressive development of high frequency hearing is consistent

with observations on other altricial bird species, such as budgerigars (Brittan-Powell and

Dooling, 2004) and precocial bird species such as chickens, ducks, and geese (Konishi, 1973;

Saunders, Coles and Gates, 1973; Saunders, Gates and Coles, 1974; Rebillard and Rubel, 1981;

Cohen, Hernandez and Saunders, 1992; Dmitrieva and Gottlieb, 1992). Therefore, the functional

maturation of hearing in zebra finches follows a gradual, progressive pattern consistent with

other bird species. The late maturation of high frequency hearing sensitivity combined with the

fact that heat calls consist of high fundamental frequencies (7-to-10 kHz) make heat calls

particularly challenging to be detected by developing zebra finches - even so by adults (Fig. 2).

This difficulty in detection is not unique to zebra finches; even precocial species that display

hearing function before hatching have high response thresholds for sounds above 6 kHz as

neonates (chicken: >120 dB SPL; mallard duck: >90 dB SPL) (Saunders, Coles and Gates, 1973;

Dmitrieva and Gottlieb, 1992). Consistent with limited high frequency hearing, the established

parent-embryo communication of precocial mallard ducks depends specifically on the low-

frequency auditory sensitivity of the embryo to detect maternal calls with frequency components

below 3 kHz (Gottlieb, 1975). Taken together, our results provide evidence against the

hypothesis that zebra finch embryos can hear heat calls, and we consider it thus unlikely that

adult zebra finches can communicate with their embryos via air-borne sound.

Sound-induced vibrations of the egg and bone conduction hearing could provide an

alternative explanation for the detection of heat calls by zebra finch embryos (Mariette, 2024).

However, the apparent lack of BC-ABRs suggest that bone conduction hearing experienced by

zebra finches is limited or non-existent (Figs 3, 4). This finding is surprising as bone conduction

hearing functions in multiple other species (Håkansson, Tjellström and Rosenhall, 1985; Sakai,

Karino and Kaga, 2006; Bergin et al., 2015; Chhan, McKinnon and Rosowski, 2017). The lack

of masking of the head vibration-induced potentials by intense noise (Fig. 4), suggest that these

responses may be vestibular compound action potentials (Jones and Jones, 1996; Jones, Jones

and Shukla, 1997; Brown, Pastras and Curthoys, 2017). This hypothesis is further supported by

the notion that vestibular responses can be recorded prior to responses of the auditory system

(Freeman *et al.*, 1999). Even if sound-induced vibrations of the egg would translate without any

loss into vibrations of the zebra finch embryo, the level of the generated vibrations is orders of

magnitudes too small to evoke responses of the vestibular system or to be detected by typical

vertebrate vibrotactile senses (Fig. 5). Taken together, i) the lack of cochlear responses to bone

conduction stimulation throughout postnatal development and at adulthood and ii) the absence of

sufficiently strong vibrotactile stimuli to the egg generated by airborne sounds make it very

unlikely that developing zebra finch embryos can perceive heat calls via detection of sound-

induced vibrations.

The role of heat calls as a communication signal has been debated (McDiarmid, Naguib

and Griffith, 2018; Mariette and Buchanan, 2019). They have instead been proposed to be a side

product of thermally induced panting to increase ventilation (McDiarmid, Naguib and Griffith,

2018; Mariette and Buchanan, 2019; Pessato *et al.*, 2020). While heat calls have been recorded in

several studies, the source level of the heat calls is currently unknown. It is therefore hard to

evaluate the biological relevance of the used experimental playbacks because heat calls may be

much softer *in vivo*. How the heat call playbacks have affected the embryos in the earlier studies

remains to be further tested and reproduced.

While the sensitivity of the zebra finch auditory system to click stimuli is similar at 10

DPH and at adulthood, other aspects of the ABR continue to change. For example, the response

latency of the auditory nerve (wave I) shortens past 20 DPH. Furthermore, the responses

generated by the more upstream neural populations of the auditory system (wave II and above)

appear to go through major changes after 10 DPH (Fig. 1c). Based on our findings, we believe

that the peripheral structures and sound encoding circuits of the zebra finch auditory system

continue to develop for several weeks after hatching. Similar observations have been made on

the well-studied precocial chicken. While chicken auditory nerve sensitivity at hatching is close

to what is observed at adulthood, the temporal coding ability of the auditory nuclei - the ability to

respond to high stimulus rates - increases post-hatch (Saunders, Coles and Gates, 1973;

Saunders, Gates and Coles, 1974; Tucci and Rubel, 1990). Furthermore, the response

characteristics of the auditory nerve fibers and the latencies of the ABR show postnatal changes

(Katayama, 1985; Manley *et al.*, 1991; Jones, Jones and Paggett, 2006). The periphery of the

chicken auditory system also continues to structurally mature for several weeks after hatching

(Ryals, Creech and Rubel, 1984; Cotanche and Corwin, 1991; Cohen, Hernandez and Saunders,

1992; Jones, Jones and Paggett, 2006). Surprisingly, we know little about these changes in zebra

finches. Experimental work indicates that zebra finches learn song only after 25 DPH, the onset

of sensory learning (Arnold, 1975; Roper and Zann, 2006; Braaten, 2010; Gobes, Jennings and

Maeda, 2019). The striking co-occurrence of this onset with the maturation of the auditory nerve

responses (Fig. 1) suggests that the onset of song learning in zebra finches may be gated by the

developmental maturation of the auditory system. It would be interesting to investigate this

possibility in other songbird species where auditory system development has also been studied,

and sensory and sensorimotor phases are more separated, such as e.g. budgerigars, and the

characterization of motor learning is on the way (Zhao *et al.*, 2023).

Interestingly, all currently known vocal learners (songbirds, parrots, hummingbirds, bats,

seals, humans, whales) seem to be altricial species. This raises the question if altriciality and

particularly parental care could have been the key that unlocked the ability for vocal learning

from ancestral precocial birds.

METHODS

Animals

Local, captivity-bred zebra finches (*Taeniopygia guttata,* order *Passeriformes*) were used in this study. Zebra finches were kept and bred indoors in temperature- and moisture-controlled group aviaries (100-to-200 individuals) or pairwise in breeding cages supplemented with nesting boxes. Lights in the aviaries were on a 12/12 hour light/dark photoperiod. Zebra finches were given water, food, cuttlefish bones, and nesting material *ad libitum*. Nesting boxes were monitored daily for accurate aging of the hatched chicks. The day when a given zebra finch chick was observed to have hatched in the morning was defined to be post-hatch day 0 (0 DPH). Hatched chicks were recognized from each other by making individualized, non-invasive down feather cuts until the usage of leg bands was feasible at DPH 10. Animal experiments were approved by the Danish National Animal Experimentation Board (Dyreforsøgstilsynet) under license number 2019-15-0201-00284.

Anesthesia

Preparation: 30-to-60 minutes prior to recording, 20 DPH-to-adult zebra finches were removed from their breeding cage or aviary, brought to the test room in a transport cage, and were fasted to minimize the reflux of fluid/food during anesthesia. Younger zebra finches were moved to the test room in a warmed transport box and were not fasted. Zebra finches were weighed prior to the induction of anesthesia.

Dosing: Anesthesia was induced with an intrapectoral injection of ketamine (Ketaminol Vet., 100 mg/ml, MSD Animal Health, NJ, USA) in combination with xylazine (Rompun vet., 20 mg/ml, Bayer Animal Health GmbH, Leverkusen, Germany). 20 DPH-to-adult zebra finches were administered 50 µg of ketamine and 10 µg of xylazine per gram of body weight. 2-to-10 zebra finches were given 50 µg of ketamine and 5 µg of xylazine per gram of body weight. After one hour, a supplementary dose (75% of the induction dose) was administered. In some zebra finches, a clear liquid - possibly due to fluid reflux or due to increased production of oral

secretions - was noticed in their mouths within 10-to-15 minutes from anesthesia induction. To

avoid airways from being obstructed, a small piece of paper tissue was inserted to the side of the

mouth to drain the liquid.

Air conduction -induced auditory brainstem responses (AC-ABRs)

Recording room: All recordings were performed by placing the zebra finch inside an audiometric

room (T-room, C-A TEGNÉR AB, Bromma, Sweden) equipped with a self-built vibration

isolation table with a padded surface. The walls of the room were padded with sound absorbing

wedges (Classic Wedge 30, EQ Acoustics, UK).

Maintenance of body temperature: The body temperature of the animals was maintained by

placing a heating pad warmed by circulating water (TP702, Gaymar Instruments Inc., Orchard

Park, NY, USA) underneath them and placing a small cloth on top of their body. Body

temperature monitoring was performed with a J-type temperature probe connected to a

thermocouple measurement device (NI USB-TC01, National Instruments). The tip of the probe

was coated with Sylgard (184 Silicone Elastomer Kit, Dow Europe GMBH, Wiesbaden,

Germany), covered with vaseline, and was placed withing the cloaca. With 2-to-10 DPH zebra

finches where the cloaca was too small for probe insertion the probe was placed under the wing.

Head placement: The head of the zebra finch was slightly elevated by placing it on top of wax

mound (Surgident Periphery Wax, Kulzer, IN, US). The ends of a wedge-shaped notch on the

wax were slightly pressed together so that the wax held the lower mandible of the zebra finch in

place. The auricular feathers covering the left external auditory meatus in mature zebra finches

were cut with scissors to allow unobstructed stimulation of the auditory system. Prior to

electrode placement, the appropriate depth of anaesthesia was confirmed by the lack of

responsiveness to a toe-pinch.

Electrodes: Three subdermal needle electrodes were placed subcutaneously on the head of the

animal in the following configuration (Fig. 1A): (1) Active electrode: Dorsal to the left auditory

meatus, pointing towards to the top of the head. (2) Inverting electrode: On top of the head along

the midline, pointing towards the bill; (3) Ground electrode: Dorsal to the right auditory meatus,

pointing towards to the top of the head. Of note, to compare these recordings to studies where the

positions of the active and inverting electrodes are the opposite one must invert the recorded

signal. The impedances of the electrodes were measured (RA4LI, Tucker-Davis Technologies,

FL, USA) to be below 1 kOhm in 2-to-10 DPH zebra finches and below 3 kOhm in 20 DPH-to-

adult zebra finches.

Recording devices: Recorded evoked potentials were passed from the electrodes to a headstage

(RA4LI) in series with a preamplifier (RA4PA) and with a digital signal processor (RM2, all

from Tucker-Davis Technologies, FL, USA). RM2 was controlled with custom-written software

QuickABR (Brandt, Brande-Lavridsen and Christensen-Dalsgaard, 2018). Recorded responses

were digitized with a sampling rate of 25 kHz (16 bits).

Sound stimulation: Sound stimulation was generated in the RM2 controlled by QuickABR,

amplified (Cambridge Audio, Azur 740A Integrated Amplifier, London, UK), and emitted from

a loudspeaker (Wharfedale Diamond 220, Wharfedale Ltd., Huntingdon, UK). The loudspeaker

was string-suspended 25 cm away from the animal, directly facing the left auditory meatus.

Sound reflections were minimized by placing the loudspeaker at an ~45-degree angle relative to

the closest wall of the recording room. To calibrate the loudspeaker, a ½" microphone (G.R.A.S.

microphone type 26AK, G.R.A.S., Holte, Denmark) was suspended at the position of the head of

the animal and connected to a microphone amplifier (G.R.A.S. Power Module Type 12AA, Gain

0, Linear filtering, Direct mode select = Ampl.). The microphone itself was calibrated using an

acoustical calibrator (type 4321, Output: 1 kHz at 94 dB re. 20 µPa, Brüel & Kjær). Calibration

of the loudspeaker was controlled by QuickABR.

Stimuli: AC-ABRs were elicited with broadband clicks (20 µs) and 25 ms-long tonebursts at 0.25

kHz, 0.5 kHz, 1 kHz, 2 kHz, 3 kHz, 4 kHz, 6 kHz, and 8 kHz. To minimize "spectral splatter"

caused by abrupt on/offsets of stimuli, the rise/fall time (5 ms, cos^2-gated) of the tonebursts was

selected to accommodate at least one full cycle of the stimulus with the lowest frequency. To

further increase the frequency specificity of the stimulus, the plateau time (15 ms) was selected

to allow at least 3 full cycles of the stimulus with the lowest frequency. Additionally, 5 ms-long

tonebursts (1 ms rise/fall time, cos^2-gated) were collected from adult zebra finches. Stimuli were

sampled at 25 kHz and repeated at a rate of 25 Hz to collect average responses consisting of 400

individual responses. Every second presented individual stimulus was inverted to reduce

microphonic effects. The maximal sound pressure level (SPL) used in this study was selected to

be 95 dB SPL as it can be viewed as being physiologically relevant. Click amplitudes are stated

as the root mean square (RMS) dB value of a sinusoid with the same peak amplitude as the click

(peak equivalent dB).

Recording paradigm: The stability of the recorded responses was monitored throughout the

recording session by recording a 95 dB SPL click-evoked AC-ABR periodically between each

stimulus type. A specific stimulus recording was accepted into further analysis if the amplitude

of the click-evoked AC-ABR wave I had changed less than 25% from the most temporally stable

amplitude of that recording session. To minimize test order bias, the recording order of stimuli

with varying frequencies was randomized. Furthermore, BC-ABRs were collected to prior to

AC-ABRs in some animals.

Head vibration-induced potentials and bone conduction-induced ABRs (BC-ABRs)

Setup and calibration: A vibrometer was attached to a piston (4501 A 30424, Brüel & Kjær,

Copenhagen) and connected to a preamplifier (1704-A-002, Gain 100x, Lin filter, Brüel & Kjær,

Copenhagen). The piston was then attached to an electrodynamic vibration exciter (Mini Shaker

Type 4810, Brüel & Kjær, Copenhagen) surrounded by a grounded wire mesh. The system was

calibrated by using a vibration calibrator (TYPE 4294 Portable Vibration Calibrator, 10 m s^{-2}

standard acceleration level at 159.15 Hz, Brüel & Kjær, Copenhagen). Calibration was controlled

by QuickABR. Maximal tested acceleration level was selected to be 10 m s^{-2} (100 dB re 0.1 mm

s^{-2}) to ensure animal safety during recordings and as the value is close to the physiologically

relevant gravitational acceleration (1 g-force).

Procedure: The vibrational exciter was lowered on a holder so that the attached piston made

physical contact with the proximal end of the upper mandible of the anesthetised zebra finch.

The piston was perpendicular to the culmen of the upper mandible. A small drop of tissue

adhesive (VetBond, 3M) was placed between the piston and the mandible to secure their physical

contact. Head vibration-induced potentials were elicited with the vibration exciter with same

stimulus types and recorded with the same electrodes/devices as AC-ABRs (see above).

Control experiments: After a recording session, used vibrational stimuli were re-recorded in the

absence of probe-to-head contact by slightly lifting the probe off from the bill. To test whether

an intense AC-stimulus could mask the head vibration-induced potentials, white noise was

played at 95 dB SPL simultaneously while recording the vibrational responses. A subset of

animals (N = 3) was used to determine whether the recorded responses are dependent on the

animal being alive. At the end of the specific recording session the animal was killed with a

lethal overdose of ketamine-xylazine without removing it from the recording setup. *Ex vivo*

responses were recorded after the animal visibly had stopped breathing. The SPL of the airborne

sounds generated by the vibration exciter were recorded with the same devices as the AC-ABR

stimuli were calibrated with. The used microphone was placed next to the vibration exciter at the

location where the ear of the tested zebra finch would reside.

Egg vibration measurements with laser doppler vibrometry

Zebra finch eggs were removed from their nest and placed in an upright position on a dental wax

covered platform at the centre of an anechoic room. The anechoic room was custom-made with

its walls and floor covered with 30 cm rockwool wedges. The sample platform was mounted on a

heavy tripod which also held a laser Doppler vibrometer (OFV-5000 with OFV-505 sensor,

Polytec, Waldbronn, Garmany) attached on a manipulator. The laser was pointed at the center of

the side of the egg. Strong reflections were directly obtained from the egg surface so adding

reflectors was not necessary. Sound-induced vibrations of the egg were induced with a speaker

(JBL IG, Northridge, CA, USA) placed next to the laser vibrometer. A microphone (½ inch

G.R.A.S., Brüel & Kjær) was placed few millimeters above the egg and was used for speaker

calibration. The egg was stimulated with sound frequency sweeps (0.25-to-10 kHz) at 84 dB

SPL. The sweeps were generated with Tucker-Davis system 2 hardware (Tucker-Davis

Technologies, TDT, Alachua, FL, USA) controlled by custom-written software DragonQuest

(Christensen-Dalsgaard and Manley, 2008). The signal was deconvoluted by dividing the

spectrum of the sweep with the measured transfer function of the speaker. The analog laser

signal of the laser vibrometer was digitized using an A–D converter (AD2, TDT) and averaged

over 100 presentations. During recordings, the laser signal was monitored using an oscilloscope

and a vibrometer controller (Polytec OFV-5000). Stimulation and recording were controlled by

DragonQuest.

Data analysis and visualization

Data was visualized and analysed with custom scripts in MATLAB R2022B (Mathworks).
Figures were edited with Illustrator CS6 (Adobe).

Threshold determination: For AC-ABR and head vibration-induced potential threshold detection,
the stimulus level was lowered from the maximal value in 5 dB steps. For the reported thresholds
to reflect the average sensitivity of zebra finches capable of detecting the given stimulus, only
zebra finches that displayed a detectable (both visually and above the signal-to-noise ratio of 2)
AC-ABR or a head vibration-induced potential at the maximal used stimulation level are
included in the analysis. The response threshold was designated to be 2.5 dB below the last
observable response-generating stimulus level (*arrow* in Fig. 1f).

Signal-to-noise ratio calculation: The recording prior to stimulus onset was used to calculate the
noise RMS. Signal-to-noise ratio (S/N) was determined by dividing the RMS value of the signal
(*blue window*) by the RMS of the recording prior to stimulus onset (*red window*, Figs. 1c., 2a,
2b, 4a, 4b, and Supplementary Fig. 2a). The analysis window to determine the signal RMS for
head vibration-induced potentials was positioned so that the AC-ABR responses evoked by the
sounds generated by the vibration exciter would not contaminate the results (Fig. 4a).

Response latency and amplitude calculation: Response latency was calculated as the time
difference between the first negative deflection of the AC-ABR (designated as wave I) and the
arrival time of the sound at the ear (Fig. 1c). The amplitude of the wave I (reflecting the activity
of the auditory nerve) was calculated as the potential difference between the first negative peak
and the following positive trough that together formed the most prominent component of the
response (Fig. 1c). Of note, a second negative peak/deflection between the first peak and the
trough existed. This peak appeared to merge with the first negative peak at soft SPLs as reported
for AC-ABRs of bald eagles and red-tailed hawks (McGee *et al.*, 2019).

Statistics

To test for main effects, we used linear mixed effects models. We sequentially compared models
by adding hypothesised effects as fixed effects one-by-one, starting from a null model and
ending with full interaction models. For all age comparisons, age was treated as an ordinal rather

than a continuous variable to account for discrete timepoints of measurements and to allow for

post-hoc pairwise comparisons between ages. All models had the individual animals as random

effect. We used chi-square tests, Akaike Information Criterion (AIC), and Bayes Information

Criterion (BIC) as criteria for model comparison. For models with significant main effects, we

then used estimated marginal means for post-hoc comparisons. We used Bonferroni p-value

correction to account for multiple comparisons. In all figures, we denote significance levels as

follows: *, p(Bonf.) = 0.05; **, p(Bonf.) = 0.01; ***, p(Bonf.) = 0.001. All statistical testing was

carried out in R 4.2.1 (R Core Team, 2024) We used the lme4 package (Bates *et al.*, 2015) for

mixed effects modelling and emmeans package (Lenth, 2024) for post-hoc comparisons.

References

Aleksandrov, L.I. and Dmitrieva, L.P. (1992) 'Development of auditory sensitivity of altricial
birds: absolute thresholds of the generation of evoked potentials', *Neuroscience and Behavioral*
Physiology, 22(2), pp. 132–137. Available at: https://doi.org/10.1007/BF01192385.

Amin, N., Doupe, A. and Theunissen, F.E. (2007) 'Development of Selectivity for Natural
Sounds in the Songbird Auditory Forebrain', *Journal of Neurophysiology*, 97(5), pp. 3517–3531.
Available at: https://doi.org/10.1152/jn.01066.2006.

Ar, A. and Tazawa, H. (1999) 'Analysis of heart rate in developing bird embryos: effects of
developmental mode and mass', *Comparative Biochemistry and Physiology Part A: Molecular &*
Integrative Physiology, 124(4), pp. 491–500. Available at: https://doi.org/10.1016/S1095-
6433(99)00142-7.

Arnold, A.P. (1975) 'The effects of castration on song development in zebra finches (Poephila
guttata)', *The Journal of Experimental Zoology*, 191(2), pp. 261–278. Available at:
https://doi.org/10.1002/jez.1401910212.

Aubret, F. *et al.* (2016) 'Heartbeat, embryo communication and hatching synchrony in snake
eggs', *Scientific Reports*, 6(1), p. 23519. Available at: https://doi.org/10.1038/srep23519.

Bates, D. *et al.* (2015) 'Fitting Linear Mixed-Effects Models Using lme4', *Journal of Statistical*
Software, 67, pp. 1–48. Available at: https://doi.org/10.18637/jss.v067.i01.

Bergin, M.J. *et al.* (2015) 'High frequency bone conduction auditory evoked potentials in the
guinea pig: Assessing cochlear injury after ossicular chain manipulation', *Hearing Research*,
330(Pt A), pp. 147–154. Available at: https://doi.org/10.1016/j.heares.2015.10.009.

Böhner, J. (1990) 'Early acquisition of song in the zebra finch, *Taeniopygia guttata*', *Animal*
Behaviour, 39(2), pp. 369–374. Available at: https://doi.org/10.1016/S0003-3472(05)80883-8.

Braaten, R.F. (2010) 'Song recognition in zebra finches: Are there sensitive periods for song
memorization?', *Learning and Motivation*, 41(3), pp. 202–212. Available at:
https://doi.org/10.1016/j.lmot.2010.04.005.

Brandt, C., Brande-Lavridsen, N. and Christensen-Dalsgaard, J. (2018) 'The Masked ABR
(mABR): a New Measurement Method for the Auditory Brainstem Response', *Journal of the*
Association for Research in Otolaryngology, 19(6), pp. 753–761. Available at:
https://doi.org/10.1007/s10162-018-00696-x.

Brittan-Powell, E.F. and Dooling, R.J. (2004) 'Development of auditory sensitivity in
budgerigars', *J. Acoust. Soc. Am.*, 115(6).

Brown, D.J., Pastras, C.J. and Curthoys, I.S. (2017) 'Electrophysiological Measurements of
Peripheral Vestibular Function—A Review of Electrovestibulography', *Frontiers in Systems*
Neuroscience, 11. Available at: https://doi.org/10.3389/fnsys.2017.00034.

Buchwald, J.S. and Huang, C. (1975) 'Far-field acoustic response: origins in the cat', *Science*
(New York, N.Y.), 189(4200), pp. 382–384. Available at:
https://doi.org/10.1126/science.1145206.

Caus Capdevila, M.Q., Sienknecht, U.J. and Köppl, C. (2021) 'Developmental maturation of
presynaptic ribbon numbers in chicken basilar-papilla hair cells and its perturbation by long-term
overexpression of Wnt9a', *Developmental Neurobiology*, 81(6), pp. 817–832. Available at:
https://doi.org/10.1002/dneu.22845.

Chhan, D., McKinnon, M.L. and Rosowski, J.J. (2017) 'Identification of induced and naturally
occurring conductive hearing loss in mice using bone conduction', *Hearing Research*, 346, pp.
45–54. Available at: https://doi.org/10.1016/j.heares.2017.02.001.

Christensen-Dalsgaard, J. and Manley, G.A. (2008) 'Acoustical Coupling of Lizard Eardrums',
JARO: Journal of the Association for Research in Otolaryngology, 9(4), pp. 407–416. Available
at: https://doi.org/10.1007/s10162-008-0130-2.

Cohen, G.M. and Fermin, C.D. (1978) 'The Development of Hair Cells in the Embryonic
Chick's Basilar Papilla', *Acta Oto-Laryngologica*, 86(1–6), pp. 342–358. Available at:
https://doi.org/10.3109/00016487809124756.

Cohen, Y.E., Hernandez, H.N. and Saunders, J.C. (1992) 'Middle-ear development: II.
Morphometric changes in the conducting apparatus of the chick', *Journal of Morphology*,
212(3), pp. 257–267. Available at: https://doi.org/10.1002/jmor.1052120305.

Colombelli-Négrel, D. *et al.* (2012) 'Embryonic learning of vocal passwords in superb fairy-
wrens reveals intruder cuckoo nestlings', *Current biology: CB*, 22(22), pp. 2155–2160. Available
at: https://doi.org/10.1016/j.cub.2012.09.025.

Colombelli-Négrel, D. *et al.* (2016) 'Vocal imitation of mother's calls by begging Red-backed
Fairywren nestlings increases parental provisioning', *The Auk*, 133(2), pp. 273–285. Available
at: https://doi.org/10.1642/AUK-15-162.1.

Colombelli-Négrel, D., Hauber, M.E. and Kleindorfer, S. (2014) 'Prenatal learning in an
Australian songbird: habituation and individual discrimination in superb fairy-wren embryos',
Proceedings. Biological Sciences, 281(1797), p. 20141154. Available at:
https://doi.org/10.1098/rspb.2014.1154.

Cotanche, D.A. and Corwin, J.T. (1991) 'Stereociliary bundles reorient during hair cell
development and regeneration in the chick cochlea', *Hearing Research*, 52(2), pp. 379–402.
Available at: https://doi.org/10.1016/0378-5955(91)90027-7.

Dmitrieva, L.P. and Gottlieb, G. (1992) 'Development of brainstem auditory pathway in mallard
duck embryos and hatchlings', *Journal of Comparative Physiology. A, Sensory, Neural, and
Behavioral Physiology*, 171(5), pp. 665–671. Available at: https://doi.org/10.1007/BF00194114.

Doody, J.S. *et al.* (2012) 'Good vibrations? Sibling embryos expedite hatching in a turtle',
Animal Behaviour, 83(3), pp. 645–651. Available at:
https://doi.org/10.1016/j.anbehav.2011.12.006.

Ehret, G. (1976) 'Development of absolute auditory thresholds in the house mouse (Mus
musculus)', *Journal of the American Audiology Society*, 1(5), pp. 179–184.

Ehret, G. and Romand, R. (1981) 'Postnatal development of absolute auditory thresholds in
kittens', *Journal of Comparative and Physiological Psychology*, 95(2), pp. 304–311. Available
at: https://doi.org/10.1037/h0077770.

Freeman, S. *et al.* (1999) 'Development of short latency vestibular evoked potentials in the
neonatal rat', *Hearing Research*, 137(1), pp. 51–58. Available at: https://doi.org/10.1016/S0378-
5955(99)00137-9.

Gobes, S.M.H., Jennings, R.B. and Maeda, R.K. (2019) 'The sensitive period for auditory-vocal
learning in the zebra finch: consequences of limited-model availability and multiple-tutor
paradigms on song imitation', *Behavioural processes*, 163, pp. 5–12. Available at:
https://doi.org/10.1016/j.beproc.2017.07.007.

Gos, M. (1933) 'Les reflexes conditionnels chez l'embryon d'oiseau.', *Bull. Soc. Roy. Sci. Liége*,
4(194–199), pp. 246–250.

Gottlieb, G. (1965) 'PRENATAL AUDITORY SENSITIVITY IN CHICKENS AND DUCKS',
Science (New York, N.Y.), 147(3665), pp. 1596–1598. Available at:
https://doi.org/10.1126/science.147.3665.1596.

Gottlieb, G. (1975) 'Development of species identification in ducklings: I. Nature of perceptual
deficit caused by embryonic auditory deprivation', *Journal of Comparative and Physiological
Psychology*, 89(5), pp. 387–399. Available at: https://doi.org/10.1037/h0077068.

Gottlieb, G. (1979) 'Development of species identification in ducklings: V. Perceptual
differentiation in the embryo', *Journal of Comparative and Physiological Psychology*, 93(5), pp.
831–854. Available at: https://doi.org/10.1037/h0077614.

Grier, J.B., Counter, S.A. and Shearer, W.M. (1967) 'Prenatal auditory imprinting in chickens', *Science (New York, N.Y.)*, 155(3770), pp. 1692–1693. Available at: https://doi.org/10.1126/science.155.3770.1692.

Håkansson, B., Tjellström, A. and Rosenhall, U. (1985) 'Acceleration levels at hearing threshold with direct bone conduction versus conventional bone conduction', *Acta Oto-Laryngologica*, 100(3–4), pp. 240–252. Available at: https://doi.org/10.3109/00016488509104786.

Hamburger, V. and Hamilton, H.L. (1951) 'A series of normal stages in the development of the chick embryo', *Journal of Morphology*, 88(1), pp. 49–92.

Heaton, J.T., Dooling, R.J. and Farabaugh, S.M. (1999) 'Effects of deafening on the calls and warble song of adult budgerigars (Melopsittacus undulatus)', *The Journal of the Acoustical Society of America*, 105(3), pp. 2010–2019. Available at: https://doi.org/10.1121/1.426734.

Hecox, K. and Galambos, R. (1974) 'Brain stem auditory evoked responses in human infants and adults', *Archives of Otolaryngology (Chicago, Ill.: 1960)*, 99(1), pp. 30–33. Available at: https://doi.org/10.1001/archotol.1974.00780030034006.

Heffner, H.E., Koay, G. and Heffner, R.S. (2008) 'Comparison of behavioral and auditory brainstem response measures of threshold shift in rats exposed to loud sound', *The Journal of the Acoustical Society of America*, 124(2), pp. 1093–1104. Available at: https://doi.org/10.1121/1.2949518.

Henry, K.R. and Haythorn, M.M. (1978) 'Effects of age and stimulus intensity of the far field auditory brain stem potentials in the laboratory mouse', *Developmental Psychobiology*, 11(2), pp. 161–168. Available at: https://doi.org/10.1002/dev.420110208.

Hess, E.H. (1972) '"Imprinting" in a natural laboratory', *Scientific American*, 227(2), pp. 24–31. Available at: https://doi.org/10.1038/scientificamerican0872-24.

Hörster, W. (1990) 'Vibrational sensitivity of the wing of the pigeon (Columba livia) — a study using heart rate conditioning', *Journal of Comparative Physiology A*, 167(4), pp. 545–549. Available at: https://doi.org/10.1007/BF00190825.

Immelmann, K. (1967) 'Zur ontogenetischen Gesangsentwicklung bei Prachtfinken.', pp. 320–332.

Jones, S.M. and Jones, T.A. (1996) 'Short latency vestibular evoked potentials in the chicken embryo', *Journal of Vestibular Research: Equilibrium & Orientation*, 6(2), pp. 71–83.

Jones, S.M., Jones, T.A. and Shukla, R. (1997) 'Short latency vestibular evoked potentials in the Japanese quail (Coturnix coturnix japonica)', *Journal of Comparative Physiology A*, 180(6), pp. 631–638. Available at: https://doi.org/10.1007/s003590050079.

Jones, T.A., Jones, S.M. and Paggett, K.C. (2006) 'Emergence of Hearing in the Chicken Embryo', *Journal of Neurophysiology*, 96(1), pp. 128–141. Available at: https://doi.org/10.1152/jn.00599.2005.

Katayama, A. (1985) 'Postnatal development of auditory function in the chicken revealed by auditory brain-stem responses (ABRs)', *Electroencephalography and Clinical Neurophysiology*, 62(5), pp. 388–398. Available at: https://doi.org/10.1016/0168-5597(85)90048-6.

Katsis, A.C. *et al.* (2018) 'Prenatal exposure to incubation calls affects song learning in the zebra finch', *Scientific Reports*, 8(1), p. 15232. Available at: https://doi.org/10.1038/s41598-018-33301-5.

Katsis, A.C. *et al.* (2021) 'Long-term effects of prenatal sound experience on songbird behavior and their relation to song learning', *Behavioral Ecology and Sociobiology*, 75(1), p. 18. Available at: https://doi.org/10.1007/s00265-020-02939-5.

Katsis, A.C. *et al.* (2023) 'Prenatal sound experience affects song preferences in male zebra finches', *Animal Behaviour*, 199, pp. 1–9. Available at: https://doi.org/10.1016/j.anbehav.2023.02.008.

Kleindorfer, S. *et al.* (2024) 'Nestling Begging Calls Resemble Maternal Vocal Signatures When Mothers Call Slowly to Embryos', *The American Naturalist* [Preprint]. Available at: https://doi.org/10.1086/728105.

Konishi, M. (1965) 'The Role of Auditory Feedback in the Control of Vocalization in the White-Crowned Sparrow', *Zeitschrift für Tierpsychologie*, 22(7), pp. 770–783. Available at: https://doi.org/10.1111/j.1439-0310.1965.tb01688.x.

Konishi, M. (1973) 'Development of Auditory Neuronal Responses in Avian Embryos', *Proceedings of the National Academy of Sciences of the United States of America*, 70(6), pp. 1795–1798.

Köppl, C. and Nickel, R. (2007) 'Prolonged maturation of cochlear function in the barn owl after hatching.', *Journal of comparative physiology. A, Neuroethology, sensory, neural, and behavioral physiology*, 193(6), pp. 613–624.

Kraemer, A. *et al.* (2017) 'Development of auditory sensitivity in the barn owl', *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural, and Behavioral Physiology*, 203(10), pp. 843–853. Available at: https://doi.org/10.1007/s00359-017-1197-1.

Lauridsen, T.B., Brandt, C. and Christensen-Dalsgaard, J. (2021) 'Three auditory brainstem response (ABR) methods tested and compared in two anuran species', *The Journal of Experimental Biology*, 224(Pt 2), p. jeb237313. Available at: https://doi.org/10.1242/jeb.237313.

Leitão, A. and Gahr, M. (2024) 'Babbling opens the sensory phase for imitative vocal learning', *Proceedings of the National Academy of Sciences of the United States of America*, 121(18), p. e2312323121. Available at: https://doi.org/10.1073/pnas.2312323121.

Lenth, R. (2024) '_emmeans: Estimated marginal means, aka Least-Squares Means_. R package version 1.10.0.' Available at: https://CRAN.R-project.org/package=emmeans.

Lippe, W. (1994) 'Rhythmic spontaneous activity in the developing avian auditory system', *The Journal of Neuroscience*, 14(3), pp. 1486–1495. Available at: https://doi.org/10.1523/JNEUROSCI.14-03-01486.1994.

Manley, G.A. *et al.* (1991) 'Activity patterns of primary auditory-nerve fibres in chickens: development of fundamental properties', *Hearing Research*, 57(1), pp. 1–15. Available at: https://doi.org/10.1016/0378-5955(91)90068-k.

Mariette, M.M. *et al.* (2018) 'Parent-embryo acoustic communication: a specialised heat vocalisation allowing embryonic eavesdropping', *Scientific Reports*, 8(1), p. 17721. Available at: https://doi.org/10.1038/s41598-018-35853-y.

Mariette, M.M. (2024) 'Developmental programming by prenatal sounds: insights into possible mechanisms', *The Journal of Experimental Biology*, 227(Suppl_1), p. jeb246696. Available at: https://doi.org/10.1242/jeb.246696.

Mariette, M.M. and Buchanan, K.L. (2016) 'Prenatal acoustic communication programs offspring for high posthatching temperatures in a songbird', *Science (New York, N.Y.)*, 353(6301), pp. 812–814. Available at: https://doi.org/10.1126/science.aaf7049.

Mariette, M.M. and Buchanan, K.L. (2019) 'Calling in the heat: the zebra finch incubation call depends on heat AND reproductive stage—a comment on McDiarmid et al. 2018', *Behavioral Ecology*, 30(3), pp. e1–e3. Available at: https://doi.org/10.1093/beheco/arz045.

Marler, P. *et al.* (1972) 'Effects of acoustical stimulation and deprivation on song development in red-winged blackbirds (Agelaius phoeniceus)', *Animal Behaviour*, 20(3), pp. 586–606. Available at: https://doi.org/10.1016/s0003-3472(72)80024-1.

McDiarmid, C.S., Naguib, M. and Griffith, S.C. (2018) 'Calling in the heat: the zebra finch "incubation call" depends on heat but not reproductive stage', *Behavioral Ecology*, 29(6), pp. 1245–1254. Available at: https://doi.org/10.1093/beheco/ary123.

McGee, J. *et al.* (2019) 'Auditory performance in bald eagles and red-tailed hawks: a comparative study of hearing in diurnal raptors', *Journal of Comparative Physiology. A, Neuroethology, Sensory, Neural, and Behavioral Physiology*, 205(6), pp. 793–811. Available at: https://doi.org/10.1007/s00359-019-01367-9.

McGlashan, J.K. *et al.* (2015) 'Hatching behavior of eastern long-necked turtles (*Chelodina longicollis*): The influence of asynchronous environments on embryonic heart rate and phenotype', *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*, 188, pp. 58–64. Available at: https://doi.org/10.1016/j.cbpa.2015.06.018.

Montgomery, R.A., Burke, C.J. and Byers, S.M. (1978) 'A Field Guide to the Aging of Wood Duck Embryos', *The Journal of Wildlife Management*, 42(2), pp. 432–437. Available at: https://doi.org/10.2307/3800287.

Mountcastle, V.B., LaMotte, R.H. and Carli, G. (1972) 'Detection thresholds for stimuli in humans and monkeys: comparison with threshold events in mechanoreceptive afferent nerve

fibers innervating the monkey hand', *Journal of Neurophysiology*, 35(1), pp. 122–136. Available
at: https://doi.org/10.1152/jn.1972.35.1.122.

Mudry, A. and Tjellström, A. (2011) 'Historical background of bone conduction hearing devices
and bone conduction hearing aids', *Advances in Oto-Rhino-Laryngology*, 71, pp. 1–9. Available
at: https://doi.org/10.1159/000323569.

Murray, J.R. *et al.* (2013) 'Embryological Staging of the Zebra Finch, Taeniopygia guttata',
Journal of Morphology, 274(10), pp. 1090–1110. Available at:
https://doi.org/10.1002/jmor.20165.

Noguera, J.C. and Velando, A. (2019) 'Bird embryos perceive vibratory cues of predation risk
from clutch mates', *Nature Ecology and Evolution*, 3(8), pp. 1225–1232. Available at:
https://doi.org/10.1038/s41559-019-0929-8.

Nordeen, K.W. and Nordeen, E.J. (1992) 'Auditory feedback is necessary for the maintenance of
stereotyped song in adult zebra finches', *Behavioral and Neural Biology*, 57(1), pp. 58–66.
Available at: https://doi.org/10.1016/0163-1047(92)90757-u.

Okanoya, K. and Dooling, R.J. (1987) 'Hearing in passerine and psittacine birds: a comparative
study of absolute and masked auditory thresholds', *Journal of Comparative Psychology
(Washington, D.C.: 1983)*, 101(1), pp. 7–15.

Pessato, A. *et al.* (2020) 'Vocal panting: a novel thermoregulatory mechanism for enhancing heat
tolerance in a desert-adapted bird', *Scientific Reports*, 10(1), p. 18914. Available at:
https://doi.org/10.1038/s41598-020-75909-6.

R Core Team (2024) *R: The R Project for Statistical Computing. R Foundation for Statistical
Computing, Vienna, Austria.* Available at: https://www.r-project.org/ (Accessed: 26 April 2024).

Rebillard, G. and Rubel, E.W. (1981) 'Electrophysiological study of the maturation of auditory
responses from the inner ear of the chick', *Brain Research*, 229(1), pp. 15–23. Available at:
https://doi.org/10.1016/0006-8993(81)90741-1.

Roper, A. and Zann, R. (2006) 'The Onset of Song Learning and Song Tutor Selection in
Fledgling Zebra Finches', *Ethology*, 112(5), pp. 458–470. Available at:
https://doi.org/10.1111/j.1439-0310.2005.01169.x.

Ryals, B.M., Creech, H.B. and Rubel, E.W. (1984) 'Postnatal changes in the size of the avian
cochlear duct', *Acta Oto-Laryngologica*, 98(1–2), pp. 93–97. Available at:
https://doi.org/10.3109/00016488409107539.

Sakabe, N., Arayama, T. and Suzuki, T. (1969) 'Human fetal evoked response to acoustic
stimulation', *Acta Oto-Laryngologica. Supplementum*, 252, pp. 29–36. Available at:
https://doi.org/10.3109/00016486909120512.

Sakai, Y., Karino, S. and Kaga, K. (2006) 'Bone-conducted auditory brainstem-evoked responses
and skull vibratory velocity measurement in rats at frequencies of 0.5–30 kHz with a new giant

magnetostrictive bone conduction transducer', *Acta Oto-Laryngologica*, 126(9), pp. 926–933.
Available at: https://doi.org/10.1080/00016480500536871.

Saunders, J.C., Coles, R.B. and Gates, G.R. (1973) 'The development of auditory evoked
responses in the cochlea and cochlear nuclei of the chick', *Brain Research*, 63, pp. 59–74.
Available at: https://doi.org/10.1016/0006-8993(73)90076-0.

Saunders, J.C., Gates, G.R. and Coles, R.B. (1974) 'Brain-stem evoked responses as an index of
hearing thresholds in one-day-chicks and ducklings', *Journal of Comparative and Physiological*
Psychology, 86(3), pp. 426–431. Available at: https://doi.org/10.1037/h0036132.

Schwagmeyer, P.L. *et al.* (1991) 'Effects of sibling contact on hatch timing in an asynchronously
hatching bird', *Animal Behaviour*, 41(5), pp. 887–894. Available at:
https://doi.org/10.1016/S0003-3472(05)80355-0.

Stapells, D. (2000) 'Threshold Estimation by the Tone-Evoked Auditory Brainstem Response: A
Literature Meta-Analysis', *Journal of speech-language pathology and audiology*, 24, pp. 74–83.

Tucci, D.L. and Rubel, E.W. (1990) 'Physiologic Status of Regenerated Hair Cells in the Avian
Inner Ear following Aminoglycoside Ototoxicity', *Otolaryngology–Head and Neck Surgery*,
103(3), pp. 443–450. Available at: https://doi.org/10.1177/019459989010300317.

Udino, E. *et al.* (2021) 'Prenatal acoustic programming of mitochondrial function for high
temperatures in an arid-adapted bird', *Proceedings. Biological Sciences*, 288(1964), p.
20211893. Available at: https://doi.org/10.1098/rspb.2021.1893.

Udino, E. and Mariette, M.M. (2022) 'How to Stay Cool: Early Acoustic and Thermal
Experience Alters Individual Behavioural Thermoregulation in the Heat', *Frontiers in Ecology*
and Evolution, 10. Available at: https://www.frontiersin.org/articles/10.3389/fevo.2022.818278
(Accessed: 1 March 2024).

Vergne, A.L. and Mathevon, N. (2008) 'Crocodile egg sounds signal hatching time', *Current*
Biology, 18(12), pp. R513–R514. Available at: https://doi.org/10.1016/j.cub.2008.04.011.

Vince, M.A. (1966) 'Artificial acceleration of hatching in quail embryos', *Animal Behaviour*,
14(4), pp. 389–394. Available at: https://doi.org/10.1016/s0003-3472(66)80034-9.

Warkentin, K.M., Caldwell, M.S. and McDaniel, J.G. (2006) 'Temporal pattern cues in
vibrational risk assessment by embryos of the red-eyed treefrog, Agalychnis callidryas', *The*
Journal of Experimental Biology, 209(Pt 8), pp. 1376–1384. Available at:
https://doi.org/10.1242/jeb.02150.

Woolf, N.K., Bixby, J.L. and Capranica, R.R. (1976) 'Prenatal Experience and Avian
Development: Brief Auditory Stimulation Accelerates the Hatching of Japanese Quail', *Science*,
194(4268), pp. 959–960.

Woolley, S.M.N. (2017) 'Early Experience and Auditory Development in Songbirds', in K.S.
Cramer et al. (eds) *Auditory Development and Plasticity: In Honor of Edwin W Rubel*. Cham:

Springer International Publishing, pp. 193–217. Available at: https://doi.org/10.1007/978-3-319-21530-3_8.

Yeh, Y.-T., Rivera, M. and Woolley, S.M.N. (2023) 'Auditory sensitivity and vocal acoustics in five species of estrildid songbirds', *Animal Behaviour*, 195, pp. 107–116. Available at: https://doi.org/10.1016/j.anbehav.2022.11.002.

Zevin, J.D., Seidenberg, M.S. and Bottjer, S.W. (2004) 'Limits on Reacquisition of Song in Adult Zebra Finches Exposed to White Noise', *The Journal of Neuroscience*, 24(26), pp. 5849–5862. Available at: https://doi.org/10.1523/JNEUROSCI.1891-04.2004.

Zhao, Z. *et al.* (2023) 'Anterior forebrain pathway in parrots is necessary for producing learned vocalizations with individual signatures', *Current biology: CB*, 33(24), pp. 5415-5426.e4. Available at: https://doi.org/10.1016/j.cub.2023.11.014.

END NOTES

Acknowledgements

We thank Catherine Carr and Albertine Leitão for their feedback on the manuscript. We also thank Sonja Jacobsen, Emilie Radoor, Emilie Jensen, Dina Stær Arengoth, and Niels Peter Jørgensen for animal care and maintenance.

Author contributions statement

T.A., J.C.D., and C.P.H.E. conceptualized experiments; T.A. collected and analyzed the data for figures 1-to-4. T.A. and J.C.D. collected and analyzed the data for figure 5. T.A. and C.P.H.E. wrote the draft of the manuscript; T.A., J.C.D., and C.P.H.E. wrote the manuscript. T.A. and C.P.H.E. acquired funding.

Funding

This grant was supported by Lundbeck Foundation (Lundbeckfonden) grant to T.A. and Novo Nordisk Fonden grant to C.P.H.E.

Competing interests

The authors declare no competing interests.