Estropausal gut microbiota transplant improves measures of ovarian function in adult mice

Minhoo Kim[1], Justin Wang[1,2,3], Steven E. Pilley[4], Ryan J. Lu[1,5], Alan Xu[1,6], Younggyun Kim[1,7], Minying Liu[8], Xueyan Fu[8], Sarah L. Booth[8], Peter J. Mullen[1,4,9], and Bérénice A. Benayoun[1,9, 10,11,12]

1. Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA 90089, USA.

2. Department of Quantitative and Computational Biology, University of Southern California, Los Angeles, CA 90089, USA.

3. Department of Chemistry, University of Southern California, Los Angeles, CA 90089, USA.

4. Department of Molecular Microbiology and Immunology, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

5. Graduate Program in the Biology of Aging, University of Southern California, Los Angeles, CA 90089, USA.

6. Thomas Lord Department of Computer Science, USC Viterbi School of Engineering, Los Angeles, CA 90089, USA.

7. Alfred E. Mann Department of Biomedical Engineering, University of Southern California, Los Angeles, CA 90089, USA.

8. Jean Mayer USDA Human Nutrition Research Center on Aging at Tufts University, Boston, MA 02111, USA

9. Norris Comprehensive Cancer Center, Keck School of Medicine, University of Southern California, Los Angeles, CA, USA

10. Molecular and Computational Biology Department, USC Dornsife College of Letters, Arts and Sciences, Los Angeles, CA 90089, USA.

11. Biochemistry and Molecular Medicine Department, USC Keck School of Medicine, Los Angeles, CA 90089, USA.

12. USC Stem Cell Initiative, Los Angeles, CA 90089, USA.

*Correspondence: berenice.benayoun@usc.edu

Abstract

Decline in ovarian function with age not only affects fertility but is also linked to a higher risk of

age-related diseases in women (*e.g.* osteoporosis, dementia). Intriguingly, earlier menopause is linked to

shorter lifespan; however, the underlying molecular mechanisms of ovarian aging are not well understood.

Recent evidence suggests the gut microbiota may influence ovarian health. In this study, we

characterized ovarian aging associated microbial profiles in mice and investigated the effect of the gut

microbiome from young and estropausal female mice on ovarian health through fecal microbiota

transplantation. We demonstrate that the ovarian transcriptome can be broadly remodeled after

heterochronic microbiota transplantation, with a reduction in inflammation-related gene expression and

trends consistent with transcriptional rejuvenation. Consistently, these mice exhibited enhanced ovarian

health and increased fertility. Using metagenomics-based causal mediation analyses and serum

untargeted metabolomics, we identified candidate microbial species and metabolites that may contribute

to the observed effects of fecal microbiota transplantation. Our findings reveal a direct link between the

gut microbiota and ovarian health.

Keywords

Microbiome, ovarian aging, menopause, 16S amplicon sequencing, ovarian RNA-seq

44 **Main**

45   Women are born with a definitive number of ovarian follicles and the gradual decrease in the

46 number of the follicles ultimately leads to menopause[1-3]. The onset of menopause has been linked with an

47 elevated risk of numerous age-related conditions, including osteoporosis and dementia[4-6]. Moreover, a

48 later onset of menopause has been identified as a potential predictor of extended lifespan[7,8]. While the

49 profound effects of menopause on the health- and lifespan of women are well-documented, the factors

50 that contribute to ovarian aging and the strategies for its prevention remain poorly understood.

51   During the reproductive years, the hypothalamus-pituitary-ovarian hormone axis ensures the

52 proper production and release of hormones necessary for ovulation and menstruation[9]. However, as

53 menopause progresses, ovaries become less responsive due to reduced follicle numbers, ultimately

54 leading to the hormonal imbalances characteristic of menopause[10]. Specifically, the serum concentrations

55 of follicle-stimulating hormone (FSH) and luteinizing hormone (LH) increase, whereas anti-Müllerian

56 hormone (AMH) and Inhibin A (INHBA) decrease with menopause progression[11-13]. These hormonal

57 adjustments are hallmarks of the menopausal transition and highlight the diminished function of the

58 ovaries[10]. Importantly, this disrupted feedback mechanism has been attributed to the wide array of

59 menopausal symptoms and health risks associated with this transition phase[14,15]. For instance, recent

60 studies have shown FSH receptor activation by FSH in the brain increases susceptibility for Alzheimer's

61 disease pathogenesis in women and blockade of FSH improves cognition in a mouse model of

62 Alzheimer's disease[14,16].

63   The adult human gut microbiome is composed of approximately 10^{13} to 10^{14} micro-organisms[17,18].

64 The microbiome consists of bacteria, viruses and fungi, with bacteria representing the largest proportion

65 of the microbial population[19]. Perturbations in the gut microbiota configuration are reported to lead to

66 development and/or progression of serious disease, including diabetes and depression[20,21]. Interestingly,

67 reported health effects of fecal microbiota derived from aged individuals have been inconsistent[22,23]. For

68 example, one study demonstrated that fecal microbiota transplantation (FMT) from 24-month-old mice to

69 3-month-old mice led to the acceleration of age-associated phenotypes in the brain, retina, and gut of the

70 recipient mice[22]. In contrast, another study found that grafting of fecal microbiota from 24-month-old mice

71 to 5- to 6-week-old germ-free mice enhanced neurogenesis in the hippocampus and induced intestinal

growth[23]. These findings underscore the microbiome's complex, multifaceted effects on various organ

systems throughout life, significantly affecting host health.

A recent study discovered that women with premature ovarian insufficiency exhibited significantly

altered gut microbial profiles compared to healthy women[24]. Notably, alternations in the gut microbiome in

premature ovarian insufficiency patients were associated with changes in serum hormone levels of key

ovarian endocrine markers, including FSH, LH and AMH[24]. Similarly, the composition of the gut

microbiome has been implicated in the development of polycystic ovary syndrome (PCOS), a disorder

characterized by hyperandrogenism and polycystic ovaries[25]. Specifically, FMT of samples from PCOS

patients to mice resulted in ovarian dysfunction in otherwise healthy recipient mice[25]. Consistent with the

importance of a healthy microbiome for ovarian function, another study has shown that 17β-estradiol

levels are significantly lowered in the germ-free female mice[26]. Moreover, transcriptome analysis of sexual

development marker genes of the liver and histological studies of ovarian follicle development in germ-

free female mice indicated that sexual maturation is perturbed in microbiota-depleted mice[26]. Importantly,

the gut microbiota has been shown to modulate serum estrogen levels (through the "estrobolome"),

further highlighting the potential role of the gut microbiota in female reproductive health[27]. The gut

microbiota regulates action of estrogens through the secretion of β-glucuronidase, an enzyme that

deconjugates estrogens into their active forms[27-29]. Dysbiosis of the gut microbiota can impair this process,

leading to a reduction in circulating estrogens and potentially contributing to the development of estrogen-

modulated conditions[27]. Collectively, these findings highlight a significant causal relationship between the

gut microbiota and ovarian function and health, suggesting a pivotal role for microbial communities in

modulating female reproductive conditions.

In this study, we analyzed the gut microbial profiles of young and estropausal (post-reproductive

phase in rodents, analogous to human menopause) female mice via 16S rRNA amplicon sequencing and

explored how the distinct microbial communities affect ovarian health through FMT experiments in young

female mice. Unbiased RNA-seq analyses revealed that modifying the gut microbiota leads to profound

changes in the ovarian transcriptomic landscape. Surprisingly, heterochronic gut microbiota transplant

from estropausal mice to young female recipients led to a reduction in inflammation-related gene

expression in the ovaries, and dampened expression of genes that tend to be upregulated in aging

ovaries. Consistently, we observed improved ovarian function and health in the estropausal FMT

recipients, supporting a surprising finding whereby old gut microbiota has beneficial effects on ovarian

health. Lastly, we identified specific microbial genera and species, through 16S rRNA amplicon

sequencing and whole genome shotgun sequencing, respectively, that might drive the observed

transcriptomic changes in the ovaries via causal mediation analyses. Overall, our study shows that

manipulating the gut microbiota can directly influence ovarian function and health.

Results

Young and estropausal female mice show distinct fecal microbial profiles

To investigate the potential link between changes in the gut microbiome and ovarian aging in

mice, we assessed (1) the ovarian health state and (2) microbial profiles of young (YF, 4-months) and

naturally aged estropausal (EF, 20-months) C57BL/6JNia female mice[30] (Fig. 1A). To quantify the overall

ovarian health state of the mice, we developed a composite "ovarian health index", modeled after the

concept of frailty index[31] (Extended Data Fig. 1A). Our goal in constructing this index was to create a

standardized, integrative tool that would enable comparative analysis of ovarian function across different

datasets. The index combines ovarian follicle counts and serum levels of AMH, FSH, and Inhibin A, using

a 3-tier scoring system based on the median values of the young and estropausal groups as reference

points. Specifically, values beyond the young median are scored as 3, values between the medians of the

young and estropausal groups are scored as 2, and values beyond the estropausal median are scored as

1 (Extended Data Fig. 1A). Given that ovarian follicle count and serum levels of key pituitary and ovarian

hormones—AMH, FSH, and Inhibin A—are widely recognized as reliable indicators of ovarian health[32-35],

we combined these parameters to form a comprehensive and objective evaluation of ovarian function

(Fig.1A-D; see Methods). Additionally, to facilitate accessibility and broader use of the ovarian health

index, we developed an R shiny application, allowing the public to analyze their own datasets using this

tool (https://alanxu-usc.shinyapps.io/Ovarian_Health_Index_Calculator/).

Consistent with previous reports[30,36], we observed a significant reduction in ovarian follicle counts

and serum levels of AMH and Inhibin A, as well as increased FSH levels in the estropausal group

(Fig.1B,C and Extended Data Fig. 1B-D). Furthermore, the estropausal group exhibited a significantly

lower ovarian health index score compared to the young group (p-value ~0.0117; Fig.1D), validating the

subsequent use of our index as a measure of ovarian health and function.

We then collected fecal samples from four independent cohorts of mice and performed 16S rRNA

V3-V4 amplicon sequencing to compare the gut microbial profiles of the young and estropausal female

mice. 16S rRNA amplicon sequencing allows identification of bacteria at the genus level by amplifying

and sequencing variable regions of the 16S rRNA gene (*e.g.* V3-V4 region), which contains unique

sequences specific to different species[37]. Quantification and analysis of the sequences from the variable

regions, or amplicon sequence variants (ASVs), have been shown to provide important insights of the

microbial compositions and environment[38].

Our analysis focused on both beta diversity, which assesses differences between microbial

communities, and alpha diversity, which looks at diversity within a community[39,40]. Beta diversity was

evaluated using the Bray-Curtis dissimilarity and Jaccard index (Extended Data Fig. 2A). The analysis of

Bray-Curtis dissimilarity and Jaccard index, conducted by cohort, revealed a distinct separation between

the young and estropausal female groups (Extended Data Fig. 2A). Moreover, the results of the principal

component analysis, applied to center log-ratio (CLR)-transformed and batch-corrected ASV counts,

further highlighted clear clustering between the young and estropausal females across 4 independent

cohorts of animals (Fig. 1E). Batch correction was performed to account for technical variability across

cohorts[41]. Alpha diversity measures the diversity within a single microbial community or ecosystem,

focusing on the variety and abundance of ASVs in a specific environment[36]. Importantly, estropausal

females exhibited significantly higher scores in both observed features and Shannon entropy compared to

the young female group (Fig. 1F). Our results suggest that the estropausal females have a higher

biodiversity and increased species richness and evenness. This pattern of increased alpha diversity with

age in female mice aligns with findings from previous studies[42]. However, in humans, alpha diversity has

been shown variable trends with age, with both increases and decreases reported in different studies[43,44].

Additionally, a recent study reported elevated alpha diversity indices in PCOS model mice[45].

We next asked which specific microbial genera changed in abundance between young and

estropausal female mice recurrently across our 4 independent cohorts (Fig. 1G and Extended Data Fig.

2B, 3). Our analysis identified three genera that were more abundant ("UP in EF") and six that were less

abundant ("DOWN in EF") in the microbiota of estropausal females (Fig. 1G and Extended Data Fig. 2B, 3;

absolute effect size > 1, adjusted p-value < 0.05). Most of the differentially abundant genera belonged to

the *Lachnospiraceae* family, offering a potential lead for future research on their impact on ovarian health.

Previous studies have reported a relative decrease in abundance of microbial species from the

Lachnospiraceae family with age in humans[46,47]. Further studies will be required to elucidate the

molecular mechanisms through which these genera affect ovarian function and health. Additionally, we

conducted a predictive analysis of functional consequences of these changes in microbial abundances

using PICRUSt2[48] (Fig. 1H). This analysis suggested an increase in metabolic pathways such as

ketogluconate metabolism and the superpathway of glycerol degradation to 1,3-propanediol, while

pathways including the superpathway of fatty acid biosynthesis initiation and phospholipases were

predicted to decrease in the estropausal group (Fig. 1H). These predicted metabolic changes, including

an increase in ketoglutarate and alterations in lipid metabolism, have been previously associated with

ovarian cancers, underscoring the potential relevance of our findings to understanding ovarian health[49,50].

Our results collectively underscore the pronounced differences in the fecal microbial profiles between

young and estropausal females.

Chemically-induced premature ovarian failure model mice show distinct fecal microbial profiles

The 4-VinylCyclohexene Diepoxide (VCD)-injected mouse model is a follicle-deplete, ovary-intact

animal model that is known to recapitulate many key aspects of human ovarian aging[51,52]. VCD

accelerates atresia by causing selective loss of primary and primordial follicles in the ovary, while leaving

the rest of the ovarian structure intact for residual hormone production[51,52]. We assessed the ovarian

health index and fecal microbial profiles of vehicle- and VCD-exposed mice (CTL and VCD, respectively;

Extended Data Fig. 4A). Briefly, 4-month-old C57BL/6NTac female and male mice were subjected to daily

intraperitoneal injections of either vehicle (safflower oil) or VCD (at a dosage of 160 mg/kg/day) for 15

consecutive days. Then, the mice were maintained for an additional 100 days to ensure the complete

development of phenotypes associated with VCD exposure[53] (Extended Data Fig. 4A). To determine

whether the observed effects on microbial profiles in females were a direct consequence of VCD-induced

ovarian dysfunction rather than off-target effects of VCD administration, we also generated and analyzed

data from male mice.

For female mice, ovarian follicle depletion and alterations in crucial ovarian hormone levels (*i.e.*

AMH, FSH, and Inhibin A) were confirmed through ovarian histology analysis and serum hormone

concentration measurements, respectively (Extended Data Fig. 4B-E). In line with these findings, the

VCD-treated group exhibited a markedly lower ovarian health index (p-value ~0.0104; Extended Data Fig.

4F), consistent with overall depleted ovarian function.

Subsequently, we conducted 16S rRNA V3-V4 amplicon sequencing on vehicle- and VCD-treated

animals. Analysis through principal component analysis of CLR-transformed ASV counts, along with Bray-

Curtis dissimilarity and Jaccard index, demonstrated a distinct separation between the vehicle-treated

(CTL) and VCD-exposed groups (Extended Data Fig. 5A-B). Alpha diversity indices, on the other hand,

revealed no significant differences between these two groups (Extended Data Fig. 5C). These findings

suggest that the overall diversity within the group's communities may appear similar, but they may be

different in the species present, their relative abundances and the functions the species perform. Notably,

beta diversity analysis of microbial profiles in male mice showed no notable distinctions between vehicle-

and VCD-treated groups (Extended Data Fig. 5A, B). Additionally, there were no differences in alpha

diversity between these groups (Extended Data Fig. 5D). These findings indicate that the observed

effects on microbial profiles in females are causally induced by ovarian dysfunction and not off-target

effects of VCD administration.

Our analysis revealed differentially abundant genera from CTL vs. VCD groups (Extended Data

Fig. 5E and 6; absolute effect size > 1, adjusted p-value < 0.05). We detected several genera from the

Lachnospiraceae family, yet none of the unique ASV IDs corresponded with those identified as

differentially abundant in the estropausal female mice. Thus, further research to accurately identify the

species significant to ovarian aging and health will be important. Then, we utilized PICRUSt2 to predict

functional abundance between the CTL and VCD groups (Extended Data Fig. 5F). Notably, we detected

down-regulation of the phospholipases pathway in the VCD group, a finding consistent with observations

in the estropausal female group (Fig. 1H and Extended Data Fig. 5F). Our findings suggest that

physiological ovarian aging and chemically-induced ovarian failure lead to substantially different

outcomes on the gut microbiota.

Young and old male mice show distinct fecal microbial profiles

Next, as an important comparison point, we also characterized the fecal microbial profiles of

young (4-months) and naturally aged old (20-months) C57BL/JNia male mice (Extended Data Fig. 7A).

This comparative study among female and male aging mice, as well as the VCD-exposed mice, aimed to

identify differences in microbial profiles attributable to general aging and those specific to ovarian aging.

We collected fecal samples from three independent cohorts of mice and performed 16S rRNA V3-V4

amplicon sequencing to compare the gut microbial profiles of the young and old male mice. Analysis

using principal component analysis of CLR-transformed ASV counts, coupled with Bray-Curtis

dissimilarity and Jaccard index, showed a distinct divergence between the young and old male groups

(Extended Data Fig. 7B-C). However, this separation was less pronounced compared to what was

observed in the female cohorts (Fig. 1E). Alpha diversity indices indicated no significant disparities

between the two age groups of males (Extended Data Fig. 7D). Similarly, a previous study examining

microbial profiles in male mice aged 3-4 months and 19-20 months found comparable alpha diversity

index values between the two age groups[54]. Through differential abundance and functional abundance

prediction analyses, we identified several genera and functional terms that varied significantly between

young and old male mice (Extended Data Fig. 7E-F and 8). Importantly, we observed a few consistent

functional terms across female and male aging cohorts. In particular, the L-lysine biosynthesis I pathway

was found to be up-regulated, while the pyridoxal 5-phosphate biosynthesis I pathway and the

superpathway of pyridoxal 5-phosphate biosynthesis were down-regulated with age in both sexes (Fig.

1H and Extended Data Fig. 7F). These results suggest that these pathways may play roles in the general

aging processes, rather than being uniquely related to ovarian aging. Further research will be necessary

to elucidate the influence of these functional pathways on the aging process.

Gut microbiota manipulation can induce changes in the ovarian transcriptome

To investigate the potential effects of gut microbiota on ovarian health, we conducted FMT

experiments to directly assess how alterations in the gut microbiota after estropause may influence

ovarian function. Specifically, young adult female mice (4-months) were subjected to treatment with anti-

biotics/anti-mycotic (Ab/Am) treatment to eliminate their initial microbial environment, and then received

fecal microbiota grafts from either young (FMT-YF) or estropausal (FMT-EF) female mice (Fig. 2A).

Subsequently, the ovaries were harvested for bulk RNA-seq analysis (Fig. 2A). Based on previous

research that has shown adverse effects of gut microbiota from donors with ovarian dysfunction, such as

PCOS[25], our initial hypothesis was that transplanting fecal matter from estropausal mice could potentially

induce estropause-like conditions in the recipient mice.

Multi-dimensional scaling (MDS) analysis of the ovarian RNA-seq dataset revealed a clear

separation between the FMT-YF and FMT-EF groups (Fig. 2B). Additionally, we detected 2,131

differentially expressed genes between the FMT-YF and FMT-EF groups (FDR < 5%; Extended Data Fig.

9A). Then, we investigated the pathways that were distinctively regulated between the FMT-YF and FMT-

EF groups. Our gene set enrichment analysis (GSEA)[55] results highlighted that pathways related to

immune regulation were significantly down-regulated in the FMT-EF groups (Fig. 2C-E; highlighted in red).

To evaluate whether these changes were the results of substantial changes in immune cells infiltration in

the tissue, we leveraged a publicly available ovarian single-cell RNA-seq dataset[56] to conduct

deconvolution analysis, thus allowing us to deduce alterations in immune cell proportions within our

dataset. Using two distinct tools, CSCDRNA[57] and Granulator[58], we determined that the proportions of

immune cells did not significantly differ between the FMT-YF and FMT-EF groups (Extended Data Fig.

9B), consistent with the notion that the transcriptional changes that we observed may reflect cell

autonomous changes in gene expression profiles. In addition, our GSEA results revealed terms

associated with chromatin structure and dynamics, including kinetochore, chromatin silencing complex,

chromosome, centromeric region, and spindle pole (Fig. 2E). These findings suggest that transcriptional

changes in the ovaries may reflect alterations in chromatin organization and chromosome dynamics,

processes critical for maintaining oocyte quality[59,60]. Together, these results indicate that pathways

associated with both immune regulation and chromatin dynamics distinguish the FMT-YF and FMT-EF

groups.

To identify transcription factors potentially driving the transcriptomic alterations observed between

the FMT-YF and FMT-EF groups, we utilized Gene Transcription Regulation Database (GTRD)[61] with the

GSEA paradigm. While our analysis did not identify any transcription factors meeting the usual stringent

criteria of FDR < 5%, we found two transcription factors showing trends of up-regulation and eight

displaying down-regulation trends from the FMT-EF group, with an adjusted FDR < 10% threshold (Fig.

2F). Then, we identified publicly available ChIP-seq datasets for the trending transcription factors via the

Cistrome Data Browser[62,63], using corresponding *bona fide* peaks to determine potential enrichment by

GSEA. Our analysis showed significant negative enrichment scores of Ncoa1 and Usp7-regulated genes

within the FMT-EF group (p-value ~0.0138 and ~1e-04, respectively; Extended Data Fig. 9C). Moreover,

we noted that the expression levels of *Ncoa1* and *Usp7* remained consistent across FMT-YF and FMT-EF

groups (Extended Data Fig. 9D), suggesting potential post-transcriptional or post-translational regulation.

While research on the roles of Ncoa1 and Usp7 in ovarian function is sparse, one study highlighted that

Ncoa1 function is important in egg production traits in Shaobo hens[64]. Additionally, an inhibitor of Usp7,

P5091, was found to inhibit the proliferation of ovarian cancer cells in another study[65]. Together with our

findings, this suggests that Ncoa1 and Usp7 represent potential transcriptional regulators of ovarian

function.

Heterochronic FMT may have rejuvenating effects on the ovarian transcriptome of the recipients

Cumulative studies have indicated that aging leads to a pro-inflammatory shift in the ovarian

microenvironment, adversely affecting ovarian function and oocyte quality[66,67]. Based on our GSEA

results which indicated down-regulation of inflammatory pathways in the ovaries of the FMT-EF group, we

next asked whether changes in the ovarian transcriptome upon FMT-EF could represent a more general

trend of transcriptional rejuvenation. For this purpose, we analyzed two publicly available bulk RNA-seq

datasets profiling ovaries from young (2-3-months) and aged/estropausal (12-months) wild-type female

mice[68,69] to identify gene sets that are either up- or down-regulated with ovarian aging (Fig. 3A). To

evaluate whether the genes identified from publicly available RNA-seq datasets on ovarian aging ("UP

with ovarian aging" and "DOWN with ovarian aging") were significantly regulated in response to FMT-EF,

we conducted GSEA using significant age-regulated genes as input gene sets. Surprisingly (and contrary

to expectations based on previously reported observation at the somatic level for heterochronic FMT), our

findings revealed a significant negative enrichment score for genes that were up-regulated in the context

of ovarian aging (p-value ~0.0046; Fig. 3B), suggesting that the ovarian transcriptome of the

heterochronic FMT recipients (*i.e.* FMT-EF group) displays transcriptional profiles resembling those of

younger females.

As an orthogonal line of evidence, we next asked whether indicators of cellular senescence (as

measured by the highly curated senescent marker gene set, SenMayo[70]) were significantly impacted by

FMT-EF. Importantly, our results showed a significant negative enrichment score for the SenMayo gene

set in the FMT-EF group, consistent with an overall "rejuvenation" trend for the ovarian transcriptome of

FMT-EF mice (p-value ~0.0382; Fig. 3C).

A recently published study highlighted the role of Nicotinamide Adenine Dinucleotide (NAD+)

metabolism in ovarian aging, showing that increased NADase CD38 expression accelerated ovarian

aging [69]. CD38 deletion or pharmacological inhibition preserved fertility and follicle reserves in aged mice

[69]. Building on these findings, we examined the expression of *Cd38* as well as inflammation markers

discussed in the study, including *Cdkn1a* and *Il1a* (Fig. 3D). In our analysis, we observed reduced *Cd38*

expression in the FMT-EF group compared to the FMT-YF group, along with decreased expression of the

inflammation markers (Fig. 3D). These findings suggest the potential involvement of the CD38-NAD

pathway in the observed phenotypes of the FMT-EF group.

FMT-EF mice show improved ovarian health

Since we observed large ovarian transcriptome remodeling in the FMT-EF group consistent with

a less inflammatory, more youthful ovarian state, we next assessed the ovarian health state of the FMT-

YF *vs.* FMT-EF groups to determine whether FMT-EF led to enhanced ovarian function (Fig. 4). We

evaluated ovarian health function using a 2-pronged approach (Fig. 4A): (i) calculating the ovarian health

index, and (ii) evaluating effective fertility using litter size and time to first pregnancy after mating[71] (Fig. 4).

Consistent with our prediction from the ovarian transcriptomes, the FMT-EF mice exhibited

significantly higher scores on the composite ovarian health index than FMT-YF mice (p-value ~0.0255;

Fig. 4B, Extended Data Fig. 10A-D), supporting an enhanced ovarian health phenotype. The ovarian

health index incorporates multiple parameters, including serum levels of AMH, FSH, and Inhibin A, as well

as ovarian follicle counts (Extended Data Fig. 1A). Notably, for the FMT cohorts, FSH levels were

measured using two different assay kits during the study due to an update implemented by the core

facility conducting the assays. To ensure consistency and comparability of data across cohorts, we

applied a correction (see Methods). This adjustment ensured that FSH levels accurately reflected ovarian

health and were comparable between cohorts (Extended Data Fig. 10D). Next, we evaluated the result of

ovarian function – fertility – through measures of litter size and reproductive success rate. Specifically,

there was a suggestive trend for increased litter size in FMT-EF mice (p-value ~0.1726; Fig. 4C). In

addition, FMT-EF mice remained nulliparous for a shorter amount of time after pairing with a young health

male (p-value ~0.002; Fig. 4D), also suggestive of improved fertility. To determine whether the inclusion

of 0 pup count data influenced our findings, we re-analyzed the latency data after excluding animals with

0 pup counts. Importantly, the latency data remained statistically significant (p-value ~0.0159 without 0

pup counts vs. p-value ~0.002 with all data). Together, our findings are consistent with

improved/rejuvenated ovarian function after FMT-EF.

Importantly, no significant differences were observed in body weight between FMT-YF and FMT-

EF groups after Ab/Am treatment (Extended Data Fig. 10E and Extended Data Table 1). Additionally,

staging of the estrous cycle on the day of sample collection revealed no significant differences between

the groups (Extended Data Table 2).

FMT-YF and FMT-EF mice show distinct microbial profiles

Importantly, we next verified that fecal transplants from young or estropausal female mice led to

distinct microbial profiles in the recipient mice. For this purpose, we conducted and analyzed 16S rRNA

V3-V4 amplicon sequencing on the gut microbiota of FMT-YF and FMT-EF mice (Extended Data Fig. 11).

Principal component analysis of CLR-transformed ASV counts, complemented by Bray-Curtis

dissimilarity and Jaccard index, showed a clear segregation between the FMT-YF and FMT-EF groups

(Extended Data Fig. 11B). While no significant differences were noted across four alpha diversity indices

(Extended Data Fig. 11C), we identified several microbial genera with significant differential abundance

between the FMT-YF and FMT-EF groups (absolute effect size > 1, adjusted p-value < 0.05; Extended

Data Fig. 11D and 12). Notably, two genera that showed increased abundance in the FMT-EF recipient

group   were   also   elevated   in   the   donor   estropausal   females

(*Bacteria;Firmicutes;Clostridia;Clostridiales;Ruminococcaceae*   and   *Bacteria;Firmicutes;Clostridia;*

Clostridiales;Lachnospiraceae;Coprococcus 2;uncultured organism ; Extended Data Fig. 11E). These

results indicate that transplanting microbiota from young or estropausal female mice led to changes in the

microbial landscape of the recipient mice. Other studies have also reported the selective enrichment of

specific microbial species from donor microbiota in recipient subjects[22,72]. These observations may be

attributable to variations in the intestinal microenvironments, which allow the proliferation of certain

species.

Subsequently, we used PICRUSt2 to predict functional abundance differences between the FMT-

YF and FMT-EF groups (Extended Data Fig. 11F). We identified significant functional groups only with

up-regulated functional terms in the FMT-EF group, with no significant down-regulated terms. Notably,

there was a significant presence of terms related to the menaquinol biosynthesis pathway, including those

involved in 1,4-dihydroxy-6-naphthoate biosynthesis II, 1,4-dihydroxy-6-naphthoate biosynthesis I, and

the superpathway of menaquinol-8 biosynthesis II (Extended Data Fig. 11F). Menaquinols, the active

hydroquinone forms of vitamin K, also known as menaquinones (MKs), facilitate the carboxylation of

vitamin K-dependent proteins, which are crucial for biological processes such as blood clotting, bone

health, and heart health[73].

To pinpoint specific microbial species that may contribute to the observed changing in the ovarian

health state, we performed whole genome shotgun (WGS) metagenomics on the FMT cohort (Fig. 5A).

Consistent with the 16S amplicon sequencing results, PCA analysis of CLR-transformed data and beta

diversity analysis revealed distinct clustering between the FMT-YF and FMT-EF groups (Fig. 5B-C).

Similarly, no significant differences were observed in alpha diversity indices, reinforcing the similarity in

overall microbial diversity between the groups (Fig. 5D). WGS analysis identified 168 differentially

abundant microbial species (FDR < 5%; Fig. 5E-F and Extended Data Fig. 13), further confirming distinct

microbial compositions in the FMT-YF and FMT-EF groups. Intriguingly, microbial species previously

reported to be differentially enriched in the fecal microbiota of PCOS patients, such as *Bacteroides*

xylanisolvens, *Bacteroides thetaiotaomicron*, *Parabacteroides distasonis*, and *Bacteroides stercoris*[25],

were significantly depleted in the FMT-EF group (Extended Data Table 3).

Functional abundance analysis using HUMAnN3[74] revealed up-regulation of pathways related to

menaquinol biosynthesis, including PWY-7371: 1,4-dihydroxy-6-naphthoate biosynthesis II, consistent

with predictions from the 16S amplicon analysis (Fig. 5G and Extended Data Fig. 11F). Intriguingly, we

detected terms associated with the NAD biosynthesis pathway, such as PWY–7761: NAD salvage

pathway II (PNC IV cycle) and PYRIDNUCSYN–PWY: NAD de novo biosynthesis I (from

aspartate)|unclassified, which aligns with our ovarian RNA-seq data showing altered expression of *Cd38*

in the FMT-EF ovaries (Fig. 3D, 5G).

Given the stability of menaquinones and their primary synthesis by gut microbiota, we further

evaluated their levels in stool samples (MK6-MK13) to explore potential systemic effects post-absorption,

as they contribute significantly to the host's vitamin K status. However, no significant differences in MKs

concentrations in the stool samples of FMT-YF and FMT-EF animals were observed (Extended Data Fig.

14A,B), although this may be the result of high animal-to-animal variability and/or the high vitamin K

content of the standard mouse chow at the USC vivarium dietary (see Discussion).

FMT-YF and FMT-EF mice show distinct serum metabolomic profiles

It is well established that gut microbiota-derived molecules can enter the bloodstream and

influence health[75]. Additionally, metabolite levels in plasma have been shown to be affected by aging[76].

To assess whether shifts in the gut microbial profiles of the FMT animals induced changes in circulating

metabolites, we performed untargeted metabolomics on serum samples from two independent

experimental cohorts (Fig. 6A). First, we performed PCA of the identified metabolites to detect unbiased

separation between the FMT-YF and FMT-EF groups. Although non-linear batch effects were present and

could not be fully corrected, a clear separation was still evident within each batch (Fig. 6B). Subsequent

LDA analysis further showed the distinct metabolomic profiles of the two groups (Fig. 6C). Finally, we

extracted metabolites that changed consistently across both cohorts and selected the top 15 based on

the LDA loadings (Fig. 6D).

Among the metabolites identified, 3,4-dihydroxyphenylpropionic acid (3,4-DHPPA) and lactamide

were significantly decreased in FMT-EF animals at FDR < 0.1 (Fig. 6D, E). 3,4-DHPPA, a microbial

metabolite derived from polyphenol metabolism, has been linked to lipid metabolism and broader

metabolic regulation[77]. Given the critical role of lipid metabolism in hormonal balance and ovarian function,

its reduction in FMT-EF animals may reflect metabolic adaptations that contribute to improved ovarian

health. Additionally, while the direct role of lactamide in ovarian function is unclear, microbial-derived

lactic acid and its derivatives have been implicated in gut microbiota-driven regulation of reproductive

health[78]. For instance, lactic acid bacteria have been shown to influence sex hormone-related gut

microbiota composition, leading to beneficial effects on ovarian physiology in preclinical models[78]. The

observed decrease in lactamide levels may indicate broader microbiome-mediated metabolic changes

that support ovarian health.

In addition to these significant metabolic changes, palmitoylcarnitine, a key metabolite involved in

fatty acid β-oxidation and mitochondrial energy production[79], was elevated in FMT-EF animals according

to LDA loadings. This suggests enhanced mitochondrial efficiency and metabolic flexibility, which are

essential for oocyte and follicular health[80]. Furthermore, palmitoylcarnitine has been implicated in immune

modulation[81], and its increased levels may indicate a metabolic shift supporting both energy homeostasis

and an anti-inflammatory ovarian environment. This aligns with our bulk RNA-seq analysis, which

revealed a downregulation of inflammatory pathways in the ovaries of FMT-EF animals (Fig. 2C-E).

Conversely, LDA loadings for β-estradiol suggested that serum levels are decreased in FMT-EF

mice. Based on the improved ovarian function observed in FMT-EF mice, this may reflect a more

physiologically regulated hormonal environment, rather than a pathological decline. Estradiol plays a

critical role in ovarian function, lipid metabolism, and mitochondrial activity[82], and its levels may be

influenced by gut microbiota-driven metabolic pathways[27]. The gut microbiome is known to modulate

steroid hormone metabolism[27], and FMT from EF donors may have contributed to a systemic

reprogramming of lipid and hormonal pathways that promote ovarian resilience.

Together, these findings suggest that FMT from estropausal donors induces distinct metabolic

shifts that may contribute to the observed improvements in ovarian health. Further studies will be needed

to determine the functional impact of these metabolites on ovarian physiology and their potential role in

reproductive aging.

Causal mediation analyses reveal microbial species potentially influencing ovarian transcriptome

Lastly, we performed causal mediation analyses to characterize microbial species that may

directly influence the ovarian transcriptome remodeling that we observed between FMT-YF and FMT-EF

mice (Fig. 7A). From the features exhibiting significant abundance variations between the groups, we

identified four microbial species (*Bacteroides zhangwenhongii, Shigella flexneri, Bacteroides stercoris*,

and *Bacteroides graminisolvens*) with a significant mediation effect on ovarian transcriptome changes

(Fig. 7B). Interestingly, *Bacteroides stercoris* was previously highlighted as a key contributor to the distinct

fecal microbiota profiles between healthy individuals and those with PCOS[25]. Further investigation into the

functional roles of these identified species in modulating ovarian function will be essential in future studies.

Similarly,   we   identified   four   ASV   IDs   (850e0d58c1aa35238538564a8010e0ff,

e4f32cdc3407edb8423447421078fdae,   55ebdcf041fb6b8b4632993423011372,   and

d4840c45eb37593108d020fb6a63f8e2) with a significant mediation effect on ovarian transcriptome

changes (Extended Data Fig. 16).

To further explore the relationship between gut microbiota and ovarian health, we performed

Spearman correlation analysis using WGS metagenomics data and serum hormone quantification data

(Fig. 7C). This analysis revealed significant correlations between multiple microbial species and hormone

levels, including *Shigella flexneri, Parabacteroides merdae,* and *Porphyromonas sp. oral taxon 275,*

suggesting these species as potential candidates influencing ovarian function. These findings provide a

strong foundation for future studies aimed at experimentally validating the roles of these microbial species

in modulating ovarian function and hormone levels.

Discussion

In this study, we examined the fecal microbial profiles of three distinct groups: (1) young and

naturally aged estropausal female mice, (2) mice with chemically-induced premature ovarian failure and

their control counterparts, and (3) young and naturally aged male mice. Through our comparative analysis,

we identified and differentiated between functional pathways that may be unique to ovarian aging (*i.e.*

phospholipases pathway) and those specific to general aging (*i.e.* L-lysine biosynthesis I, pyridoxal 5-

phosphate biosynthesis I, and superpathway of pyridoxal 5-phosphate biosynthesis). Furthermore,

through fecal microbiota transplantation assays, we discovered that alterations in the gut microbiota can

significantly impact the ovarian landscape, affecting both fertility and the health state of the ovaries. We

also identified specific microbial species as potential mediators of the observed effects on ovarian

function. The vitamin K2 and CD38-NAD pathways emerged as potential mechanisms to explore for

investigating the underlying causes of the observed phenotypes in our FMT experiments (Fig. 7D). Future

investigations focusing on these species hold significant promise for crafting novel therapeutic strategies

aimed at mitigating or potentially reversing ovarian aging.

Ovarian RNA-seq data of the FMT recipients supports causal influence of the gut microbiota on

ovarian function

Our analysis of ovarian RNA-seq data from the FMT recipients (FMT-YF vs. FMT-EF) revealed

significant changes in the ovarian transcriptome following alterations in the gut microbiota. While existing

research underscores the broad effects of the gut microbiome on host health, evidence detailing its direct

effects on the transcriptome of various tissues remains sparse. For instance, earlier research

demonstrated that mice raised in a germ-free environment exhibit a markedly distinct liver transcriptome

compared to their conventionally raised counterparts[26]. To the best of our understanding, the direct

influence of gut microbiota on the ovarian transcriptome has not yet been documented. Our analysis not

only uncovered significant variations in the ovarian transcriptome among FMT recipient mice (*i.e.* reduced

inflammatory pathway in the FMT-EF group), but we also noted the transmission of phenotypic alterations

(*i.e.* improved fertility and ovarian health). Collectively, these results offer substantial evidence of gut

microbiota's direct role in influencing ovarian function and health. Furthermore, the microbial species and

genera, and functional pathways we identified serve as valuable assets for future research aimed at

investigating potential therapeutic strategies.

Unexpected advantageous effects of grafting estropausal female fecal microbiota may be due to

compensatory mechanisms

Based on previous research that has shown detrimental effects of gut microbiota from donors

with ovarian dysfunction, such as PCOS[25], we initially hypothesized that transplanting fecal microbiota

from estropausal mice might induce estropause-like phenotypes in the recipient mice. Contrary to our

expectations, however, we observed an ovarian transcriptome in the recipients that resembled that of

young ovaries, alongside improvements in ovarian health and fertility following the transplantation of

microbiota from estropausal females. The unexpected advantageous effects observed could be attributed

to a compensatory mechanism. For instance, FSH, required for follicle maturation and ovulation, is found

in elevated levels in post-menopausal women and estropausal mice[2,83,84]. After reproductive senescence,

the ovaries no longer produce estrogen or mature eggs, so the pituitary gland compensates by producing

more FSH in an attempt to stimulate the now non-responsive ovaries[30]. Similarly, it is conceivable that the

gut microbiota in reproductively senescent female mice could be enriched with beneficial microbes and/or

metabolites that support ovarian health, perhaps as a compensatory response. Cumulative studies have

highlighted the role of gut microbiota in compensatory processes. For instance, one study demonstrated

that gut microbiota can influence the levels of peptides that regulate energy balance in the hypothalamus,

serving as a mechanism to decrease body fat[85]. Furthermore, an increase in microbes that produce

metabolites critical for the nervous system's function—such as serotonin, acetylcholine, histamine, and

dopamine—was observed in patients with mild cognitive impairment[86]. Therefore, when young adult

female mice received fecal microbiota transplants from estropausal females, it is possible that the

enriched gut microbial environment contributed to the manifestation of younger or more reproductively

active phenotypes in recipient mice.

Potential involvement of MKs in ovarian health and aging

Although our analysis of functional abundance from both WGS and 16S rRNA amplicon

sequencing datasets predicted potential shifts in the biosynthesis pathway of MKs in the FMT mice, we

observed no significant differences in the stool concentrations of MKs between the FMT-YF and FMT-EF

groups. This lack of observable difference might be attributed to the vitamin K content of the standard

chow used by the USC vivarium during our experiments (LabDiet, PicoLab Rodent Diet 20). Specifically,

the high vitamin K dietary content could have saturated the system, thus masking subtle variations in MK

production by the gut microbiome. Such nuances are typically more pronounced under conditions of a low

vitamin K diet[87]. Published studies have shown that vitamin K status influences MK production because

gut bacteria require vitamin K substrates to synthesize MKs[87]. In our study, the presence of vitamin K3

(menadione) in the diet, which is converted into MK4 in the body, likely contributed to the high basal MK

levels observed in the fecal matter of the recipient mice. While there have been limited studies, MKs have

demonstrated potential benefits for ovarian health in specific contexts, such as certain ovarian cancers

and in patients with PCOS[88,89]. Additionally, some studies have underscored the anti-inflammatory

properties of MKs[90]. Given our observations of decreased inflammatory function and enhanced ovarian

function in the FMT-EF mice, coupled with the documented benefits of MKs, MKs may be therapeutic

candidates for improving ovarian health and function. However, further research into the direct effects of

MKs on ovarian function will be necessary.

Sex-differences in aging-associated gut microbial profiles

In our comparative study, while we detected certain similarities in the microbial profiles of both

females and males with aging, our analysis primarily revealed notable differences between the sexes.

Specifically, while aging was associated with significant changes in alpha diversity in female mice, such

changes were not observed in males. Moreover, our analyses of differential abundance of microbial

genera and predictions of functional abundance revealed substantial differences in the gut microbial

profiles and their functions between females and males as they age. Indeed, extensive research has

established that aging manifests differently in females and males, a phenomenon known as sex

dimorphism in aging[91]. Particularly, the susceptibility to diseases associated with aging, including

Alzheimer's disease and obesity, varies significantly between the sexes[92-94]. Furthermore, the disparity in

lifespan between females and males is consistently observed across various species[95,96]. In agreement

with our findings in this study, sex differences in the gut microbiome associated with aging have

previously been documented[97,98]. While the limited number of existing studies suggest a possible

influence of sex hormones on the observed sex differences in the gut microbial profiles[99-101], these

findings do not negate the potential effects of non-hormonal factors, such as those related to sex

chromosomes, on the differences noted between sexes. Given the significant distinctions between sexes

highlighted in this study and their varied impacts, it is imperative to consider sex as a contributing factor in

future gut microbiome research, not only to deepen our understanding but also to address the specific

differences effectively.

The potential role of the CD38-NAD pathway in mediating ovarian rejuvenation in the FMT-EF mice

CD38, a primary NADase in mammalian tissues, plays a pivotal role in regulating NAD+ levels[102].

Emerging evidence implicates the CD38-NAD axis in aging, with increased CD38 expression contributing

to age-related declines in NAD+ levels, impairing mitochondrial function and cellular metabolism[103,104]. In

the context of ovarian aging, studies have shown that CD38 overexpression accelerates follicular

depletion and diminishes ovarian function, whereas CD38 inhibition preserves follicle reserves and fertility

in aged mice[69].

In this study, we observed reduced *Cd38* expression in the ovaries of the FMT-EF group,

alongside downregulation of inflammatory markers such as *Cdkn1a* and *Il1a* (Fig. 3D). Functional

pathway predictions from the metagenomics data further supported these findings, revealing enrichment

of pathways related to NAD biosynthesis in the FMT-EF group (Fig. 5G). Specifically, pathways such as

NAD salvage pathway II (PNC IV cycle) and NAD de novo biosynthesis I (from aspartate) were

upregulated (Fig. 5G). These results suggest that the gut microbiota established through estropausal

fecal microbiota transplantation may suppress CD38 activity, influencing ovarian NAD metabolism and

creating a more favorable ovarian environment. This aligns with previous research linking CD38-mediated

NAD+ depletion to inflammation and cellular senescence, processes that are detrimental to ovarian

function[105,106].

Together, these findings indicate that modulation of the CD38-NAD pathway by gut microbiota

may alleviate ovarian inflammation and promote a more youthful transcriptomic profile, contributing to the

enhanced ovarian health observed in FMT-EF mice.

Limitations of the study

One potential caveat in our chemically induced ovarian failure model is that VCD, while

selectively depleting ovarian follicles, is a systemically administered agent and could potentially affect

other biological systems. However, studies have shown that VCD's effects are confined to ovarian tissue,

and its impact on other organ systems have not been reported[107,108]. To further assess off-target effects,

we conducted control experiments in male mice injected with VCD or vehicle, and no significant

differences in gut microbial beta diversity indices (Bray-Curtis dissimilarity and Jaccard index) were

observed. These findings suggest that the observed microbial changes in female mice are primarily due

to ovarian dysfunction rather than direct effects of VCD, though we acknowledge the potential for non-

specific effects. Additionally, it is possible that the improved fertility outcomes in the FMT-EF group,

compared to the FMT-YF group, may reflect differences in how aged and young donor microbiota

recolonize the gut following antibiotics treatment. For example, the aged microbiota might better adapt to

recolonization, reducing the likelihood of artificially poor fertility outcomes in the recipients. Nonetheless,

the changes in the microbiota induced by FMT resulted in significant changes in ovarian health, as

evidenced by differences in ovarian transcriptomic profiles, fertility outcomes, and overall ovarian health

indices in the FMT-YF and FMT-EF groups. Finally, another limitation of the study involves the use of

PICRUSt2, which predicts functional abundances based on marker gene sequences. While PICRUSt2

provides insights into potential functional pathways affected by changes in the microbiota, it is a predictive

tool and does not directly measure functional abundances. Future studies will require additional functional

assays to validate these predictions.

In summary, we performed a comparative analysis of microbial profiles among aging female,

aging male and a chemically-induced menopause model mice to characterize microbial genera that are

unique to general aging vs. ovarian aging. Our work highlights potential microbial species and genera,

and functional pathways that may play key roles in modulating ovarian function.

Acknowledgements

We thank Dr. Victor Ansere and Dr. Juan Bravo for feedback on our manuscript. We also thank Dr. Dario Valenzano for his insightful comments and guidance on microbiome data analysis.

This work was supported by the GCRLE-2020 post-doctoral fellowship from the Global Consortium for Reproductive Longevity and Equality at the Buck Institute, made possible by the Bia-Echo Foundation to M.K., USC Provost's Undergraduate Research Fellowship to J.W., National Institute on Aging (NIA) T32 AG052374 predoctoral fellowship and Diana Jacobs Kalman/AFAR Scholarships for Research in the Biology of Aging to R.J.L., and Pew Biomedical Scholar award #00034120 from the Pew Charitable Trust to B.A.B. Ovarian histological analysis was performed by the Translational Pathology Core at the USC Norris Comprehensive Cancer Center (supported by NCI P30 CA014089). Serum hormone quantification was performed by the University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core (supported by the Eunice Kennedy Shriver NICHD/NIH Grant R24HD102061). Computational analyses were performed using the Center for Advanced Research Computing (CARC) resources at the University of Southern California (USC). Stool vitamin K concentrations were supported by the USDA Agricultural Research Service under Cooperative Agreement No. 58-1950-7-707. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the USDA. We thank Whitaker Cohn, the Assistant Director of the USC Mann Multi-Omics Mass Spectrometry Core, for running the metabolomics samples on their instruments.

Author contributions

M.K. and B.A.B. designed the study. M.K. performed live animal experiments. M.K. and R.L.J. dissected mouse tissues. M.K., J.W. and Y.K analyzed ovarian histology data. A.X. generated R shiny application. M.L., X.F. and S.L.B quantified stool MK concentrations. S.E.P, M.K. and P.A.M performed and analyzed metabolomics data. M.K. and J.W. performed computational analyses. M.K. and B.A.B. wrote the manuscript with input from all authors. All authors edited and commented on the manuscript.

Declaration of interests

The authors declare no competing interests.

Methods

Mouse husbandry

All animals were treated and housed in accordance with the Guide for Care and Use of

Laboratory Animals. All experimental procedures were approved by the USC's Institutional Animal Care

and Use Committee (IACUC) and are in accordance with institutional and national guidelines. Samples

were derived from animals on approved IACUC protocol number 20770 and 21212.

For the "aging cohort," female and male C57BL/6JNia mice (4- and 20-month-old animals) were

obtained from the National Institute on Aging (NIA) colony at Charles River Laboratories. For the "VCD

cohort" and "fecal microbiota transplantation (FMT) cohort", female and male C57BL/6NTac mice (3.5-

month-old animals) were ordered from Taconic Biosciences. All animals were acclimated at the specific-

pathogen-free animal facility at USC for 2 weeks before any processing. All animals were fed PicoLab

Rodent Diet 20 (LabDiet, 5053). The facility is on a 12-h light/dark cycle and animal housing rooms are

maintained at 72 °F and 30 % humidity. All animals were euthanized between 8–11 am, by CO_2

asphyxiation followed by cervical dislocation.

For the "VCD cohort," animals were subjected to daily intraperitoneal injections of either vehicle

(safflower oil) or VCD (at a dosage of 160 mg/kg/day) for 15 consecutive days. All injections were

performed between 8:00 AM and 10:00 AM to minimize variability due to circadian effects. Lethality was

observed in male VCD-injected mice (4 out of 10 males in the experimental group within 16 days of the

start of injections). Intriguingly all lethality cases occurred in males that had to be single-housed prior to

injections due to fighting and wounds. Effects may have been aggravated due to lack of social

interactions and/or temperature regulation.

Ovarian follicle counts

Ovaries were fixed in Bouin's solution (Sigma, HT10132) for 24 hrs at room temperature and

transferred to 70 % ethanol for storage. Paraffin embedding, tissue sectioning and staining with

hematoxylin and eosin (H&E) were performed by the USC Norris Comprehensive Cancer Center

Translational Pathology Core Facility. H&E staining slides were imaged on the Keyence BZ-X All-in-One

Fluorescence Microscope platform. Three sections per ovary were used to count the number of primordial,

primary, secondary, and antral follicles and corpus luteum by three blinded observers. Median counts

across the 3 observers were used for data analysis. Statistical significance between groups was

assessed using the non-parametric Wilcoxon rank-sum test.

Quantification of serum AMH, FSH and Inhibin A concentrations

After euthanasia, blood was collected directly from the heart. Blood was allowed to clot at room

temperature for 1 hour, and serum was separated using MiniCollect® Serum Tube (Greiner, 450472),

then stored at -80 °C until further processing. Quantification of serum AMH (Rat and Mouse Anti-Müllerian

hormone (AMH) ELISA kit, Ansh Labs, AL-113), FSH (Millipore Pituitary Panel Multiplex kit, RPT86K and

Ultra-Sensitive Mouse & Rat FSH, UVA Ligand Core, in-house[109]), and Inhibin A (Inhibin A ELISA kit, Ansh

Labs, AL-161) levels were performed by the University of Virginia Center for Research in Reproduction

Ligand Assay and Analysis Core. Standard normalized values were provided by the core. Statistical

significance between groups was assessed using the non-parametric Wilcoxon rank-sum test.

For FSH quantification, two different kits were used during the course of the study, necessitating

correction of the dataset generated by the ultra-sensitive kit. To ensure consistency, we generated data

using both kits with the same serum samples and devised a polynomial model to fit the relationship

between the two datasets (R^2 = 0.90). The trained polynomial model was then applied to correct the data

generated by the Ultra-Sensitive Mouse & Rat FSH assay. This correction process was specifically

applied to the data for the FMT cohort. The raw data used to train the model has been uploaded to Github

(https://github.com/BenayounLaboratory/Ovarian_Aging_Microbiome).

Ovarian health index calculation

The ovarian health index was calculated by integrating two key components: (1) ovarian hormone

levels (AMH, FSH, and Inhibin A) and (2) follicle counts (combined counts of primordial, primary,

secondary, antral follicles, and corpus luteum). A 3-tier scoring system was used for each parameter.

Values beyond the young medians were assigned a score of 3, values between the young and

estropausal medians were assigned a score of 2, and values beyond the estropausal medians were

assigned a score of 1. The average hormone score (calculated from the AMH, FSH, and Inhibin A scores)

was then combined with the follicle score in a 1:1 ratio to generate the overall ovarian health index. This

combined score was scaled to a 0-100 range for ease of interpretation.

Statistical significance between groups was assessed using the Wilcoxon rank-sum test. Scripts

used to calculate the ovarian health index are available on the Benayoun lab GitHub at

https://github.com/BenayounLaboratory/Ovarian_Aging_Microbiome.

DNA extraction for 16S rRNA amplicon and shotgun metagenomics analyses

Fecal matter for 16S rRNA V3-V4 amplicon and shotgun metagenomics analyses were collected

aseptically post-euthanasia. Samples were immediately snap frozen in sterile tubes and stored at -80 °C.

DNA extraction was performed using QIAamp Fast DNA Stool Mini Kit (QIAGEN, 51604), following the

manufacturer's protocol.

V3-V4 region targeted amplification and sequencing were performed by Novogene Inc.

Amplification primers used were CCTAYGGGRBGCASCAG (FWD_341F) and

GGACTACNVGGGTWTCTAAT (REV_806R). Amplicon libraries were sequenced on the Illumina

NovaSeq 6000 machine. Shotgun metagenomics library preparation and sequencing were performed by

Novogene Inc. on the Illumina NovaSeq X machine. De-multiplexed FASTQ files were provided by

Novogene Inc. for data analysis.

16S rRNA V3-V4 amplicon sequencing data analysis

Data pre-processing, quality control and denoising

Data pre-processing, quality control and denoising were performed within the QIIME2[110] (v.

2023.7) platform. Each cohort was analyzed independently for memory and computing efficiency. FASTQ

files were imported to QIIME2 and denoised using DADA2[111]. Forward and reverse reads were truncated

at 226 bp and 224 bp, respectively, to obtain reads with quality scores above 25 and to ensure coverage

of the V3-V4 region (–p-trunc-len-f 226 –p-trunc-len-r 224).

Batch correction and principal component analysis of amplicon sequence variants

Previous studies have shown that effectiveness of batch correction tools are dataset-dependent[41].

For batch correction, a previously described benchmarking pipeline[41] was used to select the most

effective batch correction tool for our dataset. removeBatchEffect function from limma[112] (v. 3.50.1),

Combat from sva[113] (v. 3.42.0) and FAbatch from bapred[114] (v. 1.1) were benchmarked. Denoised feature

tables were imported to R using qiime2R[115] package (v. 0.99.6). Then, features with low counts (less than

0.01% of all the counts) were removed and filtered counts were centered log ratio (CLR)-transformed.

Based on principal component analysis results (method described below) and variance calculations,

FAbatch-corrected values were chosen for downstream analyses.

<u>Principal component analysis of amplicon sequence variants</u>

For principal component analysis (PCA), pca function from mixOmics[116] (v. 6.18.1) was used in R.

For female and male aging cohorts, FAbatch-corrected values were used for PCA. For VCD and FMT

cohorts, CLR-transformed values were used for PCA.

<u>Beta diversity and alpha diversity analyses</u>

For beta and alpha diversity analyses, phylogenetic trees were generated using the 'qiime

phylogeny align-to-tree-mafft-fasttree' command. Rooted phylogenetic trees were used to run the 'qiime

diversity core-metrics-phylogenetic' command for beta and alpha diversity analyses. Diversity analysis

results were imported to R using qiime2R[115] package (v. 0.99.6) for further plot generation and

significance tests. Bray-Curtis dissimilarity and Jaccard index principal coordinate analysis results were

used to generate beta diversity plots using ggplot2 (v. 3.4.2). To ensure comparability across different

cohorts, we normalized alpha diversity metrics within each cohort by dividing the values of each sample

by the median value of the corresponding metric (e.g., observed features, Shannon entropy) within that

cohort (e.g., CTL vs. VCD, FMT-YF vs. FMT-EF). This normalization allowed for combined cohort analysis.

The statistical significance of differences in alpha diversity between groups was determined using the

Wilcoxon rank-sum test.

<u>Differential abundance analysis of microbial genera</u>

Differential abundance analysis of microbial genera was performed using the ALDEx2[117] plugin

within the QIIME2[110] platform. Taxonomic classifier was generated using the SILVA[118] (132 release)

reference. Reference reads were extracted for CCTAYGGGRBGCASCAG (FWD_341F) and

GGACTACNVGGGTWTCTAAT (REV_806R) primer set and a classifier was trained using the 'qiime

feature-classifier extract-reads' and 'qiime feature-classifier fit-classifier-naive-bayes' commands,

respectively. Differential abundance analysis results and taxonomic classifier were imported to R using

qiime2R[115] for comparative analysis and plot generation. For female and male aging cohorts, differential

abundance analysis results for each cohort for each sex was combined based on the directionality of the

differential abundance measures, identifying features that consistently exhibited either an increase or

decrease across all cohorts for each sex. Using Fisher's method and Benjamini-Hochberg (BH) correction

method, p-values for each feature's differential abundance were combined and corrected, respectively.

Additionally, the mean of effect size across cohorts for each feature was used for analysis. For combined

female and aging cohorts, as well as VCD and FMT cohorts, features with an adjusted p-value < 0.05 and

an absolute effect size > 1 were identified as significantly different in abundance.

<u>Functional abundance prediction analysis</u>

Functional abundance prediction analysis was performed via the PICRUSt2[48] plugin within the

QIIME2[110] platform using default parameters. PICRUSt2 results were imported to R using qiime2R. For

differential abundance and statistical significance analyses, ALDEx2[117] was used (500 Monte Carlo

samples, Wilcoxon test for statistical testing, effect size calculation enabled, and the interquartile log ratio

(iqlr) for denominator calculation). To combine independent cohorts for female and male aging cohorts,

Fisher's method and Benjamini-Hochberg multiple testing correction were used to combine and correct

the p-values. Also, the mean values of effect sizes were calculated for each feature. Adjusted p-value less

than 0.05 and absolute effect size greater than 1 were used as criteria to filter functional pathways that

were significantly different in abundance.

Fecal microbiota transplantation

For fecal microbiota transplantation (FMT), animals underwent a 10-day course of

antibiotics/antimycotic (Ab/Am) treatment before transplantation. The regimen included ampicillin in

drinking water at 1 g/L (Sandoz, 00781), and oral gavage of amphotericin-B (Sigma, A9528) at 1

mg/kg/day, metronidazole (MFR VIONA PHARMACEUTICALS, 096988) at 100 mg/kg/day, vancomycin

(Alvogen, Vet-Rx-MW 090160) at 50 mg/kg/day, and neomycin (MFR VETONE, 510570) at 100

mg/kg/day. Starting on the third day after the end of the Ab/Am treatment, FMT was performed biweekly

for three weeks via oral gavage. Donor stool samples from more than 30 young or estropausal female

mice were pooled and prepared by suspending in saline (20 mg/ml), followed by vortexing and

centrifugation to obtain the supernatant. Each recipient animal received 100 µl of the fecal suspension.

Two independent cohorts were enrolled in the study, each using independently prepared pools of FMT

materials. Sample collection for subsequent processing and analysis occurred three days after the final

FMT session.

Ovarian RNA-seq sample preparation

Ovaries were harvested and immediately snap-frozen in sterile tubes, and then stored at -80 °C.

For RNA extraction, 600 µl of Trizol reagent (Ambion, 15596018) was added to the ovaries in Lysing

Matrix D tubes (MP, 6913500) and tissues were homogenized with a BeadBug 6 microtube homogenizer

(Benchmark Scientific, D1036) at 3,500 rpm for 30 seconds for 3 cycles in the cold room. Total RNA was

purified using the Direct-Zol RNA Miniprep (Zymo Research, R2052), following the manufacturer's

instructions. Integrity and quality of the purified total RNA were assessed on the 4200 TapeStation system

(Agilent, G2991A) using High Sensitivity RNA ScreenTapes (Agilent, 5067–5579) to determine the RNA

Integrity Number (RIN). Subsequently, mRNA-seq libraries were prepared and sequenced by Novogene

Inc. to obtain PE150 sequencing reads on the NovaSeq X Plus platform.

FMT cohort ovarian RNA-seq data analysis

Data quality control and pre-processing

Paired-end 150-bp reads were hard-trimmed to remove the first 6 bases and keep up to 100bp

using Fastx Trimmer[119] (v. 0.0.14) and processed using Trim Galore[120] using default parameters (v. 0.6.6).

Trimmed reads were mapped to the mm39 genome reference using STAR[121] (v. 2.7.9a). Read counts

were assigned to genes from the mm39 genome reference using subread[122] (v. 2.0.3). After assigning

read counts to genes, downstream processing was performed in R (v. 4.1.2). Low count genes (genes

with less than 1 count in at least 6 out of 14 RNA-seq libraries) were removed to improve memory and

speed efficiency of DESeq2[123], as recommended by the developers. We used surrogate variable analysis

to estimate and correct for unwanted experimental noise. R package sva[113] (v. 3.42.0) was used to

estimate surrogate variables and the removeBatchEffect function from limma[112] (v. 3.50.1) was used to

regress out the effects of surrogate variables and RNA integrity differences (RNA integrity number scores)

from raw read counts. To remove extreme values within the read counts, Winsorize function from

DescTools[124] (v. 0.99.50) was used to adjust the values above the 99.9th percentile to the values at the

99.9th percentile. DESeq2[123] package (v. 1.34.0) was used for downstream analyses of the RNA-seq data.

Genes with FDR < 5% were considered statistically significant. For the multidimensional scaling (MDS)

analysis, we calculated the distance between samples using the inverse of Spearman's rank correlation

coefficient (1-Rho) as the distance metric. These distances were then input into the cmdscale function in

R to compute the MDS coordinates.

Gene ontology analysis of RNA-seq dataset

We used the GSEA[55] approach through the phenoTest package. GO term annotations

(ENS.GO.BP, ENS.GO.MF, ENS.GO.CC and GTRD gene sets) were sourced from ENSEMBL (Ensembl

108) and Molecular Signature Database[125]. The t-statistic from DESeq2[123] was used to rank genes for

functional enrichment analysis. For readability, only the top ten most-significant pathways with negative

NES and top ten most-significant pathways with positive NES are shown in figures if more than that

passed the FDR < 5% significance threshold for GO terms and FDR < 10% for GTRD transcription

factor targets (see main text for explanation).

Deconvolution analysis of immune cell type proportion

To estimate any differences in the immune cell proportions between FMT-YF and FMT-EF groups

of the FMT cohort, single-cell RNA-seq reference-based deconvolution analyses were performed, using

CSCDRNA[126] and Granulator[127]. We downloaded the cell type annotated Seurat object of a publicly

available murine ovarian single-cell RNA-seq dataset (Open Science Framework dataset ID 924fz)[56]. In

the CSCDRNA analysis, the min.p parameter was set to 0.3 following the authors' recommendation[126], to

efficiently exclude the 30% least significant portion of marker genes for each cell type. This threshold was

chosen to increase efficiency and reduce noise. For Granulator, the qprogwc method[128] was used with

default parameters to estimate the immune cell proportions within the ovarian bulk RNA-seq dataset. Cell

proportions were tested for significance using a Wilcoxon test rank-sum test.

<u>Pre-processing and filtering candidate transcription factor-bound genes</u>

For potential transcription factors identified from GSEA[55] analysis against GTRD[61] transcription

factor targets, peak information was obtained for Cistrome.org. Peak annotation was performed using

HOMER[129] (v. 4.11.1). Genes with peaks annotated in TSS-promoter of protein coding genes were

considered direct target genes. Obtained gene lists were used to perform GSEA through the phenoTest (v.

1.42.0) package.

<u>Pre-processing of public ovarian aging datasets</u>

To evaluate the correlation between ovarian aging and FMT gene expression profiles of the

ovaries, we pre-processed and analyzed two publicly available ovarian aging datasets (2-3-month-old vs.

12-month-old female mice; BioProject dataset PRJNA100222 and GSA dataset CRA003645)[68,69]. For

each dataset, reads were pre-processed and aligned to the mm39 genome reference using the same

methods as for the FMT cohort ovarian RNA-seq dataset. Low count genes (genes with less than 1 count

in at least 4 out of 7 RNA-seq libraries) were removed. sva[113] (v. 3.42.0) was used to estimate surrogate

variables and the removeBatchEffect function from limma[112] (v. 3.50.1) was used to regress out the

effects of surrogate variables from raw read counts. DESeq2[123] package (v. 1.34.0) was used for

downstream analyses of the RNA-seq data. p-values for common genes between the two datasets were

combined and corrected using Fisher's method and BH correction method, respectively. FDR < 1e-5 was

used to catalog gene sets that are robustly up- or down-regulated with aging. Obtained gene lists were

used to perform GSEA through the phenoTest package.

Fertility assay

Fertility assay was performed as previously described[71]. FMT female mice were paired with 4-

month-old males at a 1:1 ratio 3 days after the last day of FMT administration until the birth of the first

litter. The number of pups in the first litter and the latency, defined as the duration between the pairing

date and the birth date of the first litter, were recorded and analyzed. Statistical difference in latency in the

1st litter after pairing was evaluated using a log-rank test. For pairs that did not produce pups within the

60-day timeframe, the latency was recorded as 60 days.

Whole genome shotgun metagenomics analysis

Data quality control and pre-processing

Paired-end 150-bp reads were pre-processed to remove sequencing adapters and low-quality

reads using Trimmomatic (v. 0.39)[130]. Trimmed reads were then mapped to the mouse reference genome

(mm10) using Bowtie2 (v. 2.4.4)[131] to remove host DNA contamination. Unmapped reads from paired-end

files were extracted using the --un-conc-gz option in Bowtie2 and used for subsequent analyses. Quality-

controlled and host-depleted reads were then processed for taxonomic classification and functional

profiling.

Taxonomic classification, abundance estimation and functional profiling

Kraken2 (v. 2.1.2)[132] was used for taxonomic classification of unmapped reads. Reads were

classified against a pre-built standard Kraken2 database containing bacterial, archaeal, and viral

sequences. Classification outputs were summarized using Bracken (v. 2.7)[133] to refine species- and

genus-level abundance estimates. Bracken was configured with a read length of 150 bp and default

settings for abundance profiling. Functional profiling was conducted using HUMAnN3 (v. 3.0)[74]. Reads

were processed in paired-end mode. Pathway and gene family tables were further merged and

normalized using human_join_tables and human_renorm_table functions.

Alpha and beta diversity and differential abundance analysis

Raw taxonomic abundance tables from Bracken (v. 2.7)[133] were loaded in R and filtered to

remove low-abundance and low-prevalence features. Features with total counts <20, relative abundance

<0.01%, or prevalence in fewer than 10% of samples were excluded. Samples with high host DNA

alignment rates (i.e., MHK271) were also removed. Centered log-ratio (CLR) transformation of filtered

relative abundances was performed. Shannon entropy was calculated using the mia[134] (v. 1.12.0) R

package. CLR-transformed relative abundance data were analyzed using mixOmics[116] (v. 6.28.0) and

Bray-Curtis and Jaccard distance matrices were calculated using vegan (v. 2.6-8). Differential abundance

of species was assessed using ALDEx2[117] (v. 1.36.0). Features with adjusted p-values of <0.05 were

deemed significant.

<u>Functional pathway analysis</u>

Relative abundance data for microbial pathways from HUMAnN3[74] (v. 3.9) was loaded in R and

low-abundance pathways with total abundance below 0.001 across all samples were removed.

Differential abundance of microbial pathways between groups was analyzed using ANCOMBC2[135] (v.

2.0.0). Significant pathways with Q-values below 0.1 were identified, and log2-fold change values were

used to assess the direction and magnitude of differential abundance.

Stool menaquinone concentration quantification

Frozen stool samples were shipped to the Vitamin K Laboratory at the Tufts University Human

Nutrition Research Center on Aging (HNRCA), where stool menaquinones concentrations were measured

using the LC/MS, as detailed elsewhere[136].

Serum untargeted metabolomics

<u>Sample processing</u>

Serum samples were stored at -80°C until they were thawed on ice for analysis. Metabolites were

extracted by adding 80 µL of ice-cold methanol to 20 µL of serum, followed by incubation on ice with

vortexing every 5 minutes. The samples were then centrifuged at 21,300 × g for 10 minutes at 4°C.

<u>Liquid Chromatography-Tandem Mass Spectrometry (LC-MS/MS)</u>

Untargeted metabolomics was performed at the USC Alfred E. Mann Multi-Omics Mass

Spectrometry Core. A volume of 10 µL from each sample was injected onto a SeQuant ZIC-pHILIC

Column (5 µm, 2.1 mm × 150 mm; Millapore Sigma) using a Vanquish Flex UHPLC System (Thermo

Fisher Scientific). Chromatography was conducted at a flow rate of 100 µL/min with an optimized gradient

of solvent A (water/20 mM ammonium carbonate, pH 9.8) and solvent B (acetonitrile) as follows: 0 min/97%

B, 1 min/97% B, 28 min/75% B, 39 min/50% B, 44 min/20% B, 45 min/20% B, 45 min/97% B, 60 min/97%

B. The chromatography system was coupled to a heated electrospray ionization (HESI) source and a

hybrid quadrupole-Orbitrap mass spectrometer (Q Exactive, Thermo Fisher Scientific). Mass spectra were

acquired in Full MS/data-dependent MS2 mode, alternating between positive and negative ion polarities.

Source parameters were set as follows: a source voltage of 3.5 kV, source temperature of 275°C,

sheath gas of 45, auxiliary gas of 10, sweep gas of 2, and capillary temperature of 320°C. Full scans

were performed across an m/z range of 70 - 1050 with the following settings: an automated gain control

(AGC) target of $5 \times 10^{\square}$, a maximum injection time of 100 ms, and a resolution of 70,000 at m/z 200. Up

to five consecutive MS/MS scans were acquired per full scan with the following parameters: quadrupole

isolation of charge states 1–7, isolation window of 0.6 m/z, normalized collision energies of 20, 30, and 40,

dynamic exclusion of 30 s, AGC target of $1 \times 10^{\square}$, maximum injection time of 100 ms, and resolution of

17,500 at m/z 200.

<u>Data analysis</u>

Raw data were processed using Compound Discoverer (Version 3.3 SP3, Thermo Fisher

Scientific) for metabolite identification, retention time alignment, and unknown compound detection across

all samples. Metabolites were identified by matching exact masses (delta ppm of >10 or <-10) and

fragmentation spectra to the mzCloud database (ddMS2; 'mzCloud Best Match' > 50). For quality control,

only ions with a peak rating > 4 in at least six samples within a group were retained. VSN normalized

filtered metabolic matrix via normalizeVSN function of limma[112] (v. 3.50.1) was used for PCA analysis. The

filtered metabolic matrix was then scaled, and Linear Discriminant Analysis (LDA) was performed via the

MASS (v. 7.3-60.2) R package. Metabolites with a consistent direction of $\log_2$ fold change ($\log_2$FC)

between two batches were subsequently retained for abundance analysis.

Causal mediation analysis

Causal mediation analysis to identify specific features and amplicon sequence variants (ASVs)

from WGS and 16S rRNA amplicon sequencing datasets from the FMT cohort acting as mediators

between the treatment (FMT-YF vs. FMT-EF) and outcome (changes in the ovarian transcriptome profiles)

was performed using mediation package[137] (v. 4.5.0) in R. Feature table, taxonomic reference and

differential abundance analysis results generated from QIIME2[110] were imported to R using qiime2R[115].

ASVs identified as significant in differential abundance analyses were tested. CLR-transformed ASV

counts were used for the mediation analyses. For the outcome variable, the first dimension from the

multidimensional scaling (MDS) analysis of the FMT ovarian mRNAseq dataset was used. Linear models

were constructed for both the mediator and the outcome, incorporating the treatment as an independent

variable in each. The mediate function from the mediation package was then employed to estimate the

Average Causal Mediation Effects (ACME), Average Direct Effects (ADE), and Total Effects.

Quantification and statistical analysis

944 All statistical analysis was performed using the R software, version 4.2.4. For all boxplots, the

945 data is shown with the median, the 25th and 75th percentile of the data, and the whiskers represent 1.5 *

946 the inter-quartile range (IQR). Individual datapoints are always overlayed for transparency and rigor.

947 Specific statistical tests used, the number of biological replicates and how many animals they are

948 indicated in the corresponding figure legends and associated method section.

Data availability

The raw FASTQ files have been deposited to the Sequence Read Archive under accession

PRJNA1076509. Raw microscopy pictures of ovarian hematoxylin and eosin staining are available on

Figshare (Figshare DOIs: 10.6084/m9.figshare.25749588, 10.6084/m9.figshare.25749594,

10.6084/m9.figshare.25749591, 10.6084/m9.figshare.28462274). Processed serum metabolomics data

can be found in Extended Data Table 4. Raw data can be found on Metabolomics Workbench, project

ST003750. All R code was run using R 4.1.2. Any additional information required to reanalyze the data

reported in this work paper is available from the corresponding author upon request.

Code availability

All scripts used to analyze the datasets are available on the Benayoun lab GitHub at

https://github.com/BenayounLaboratory/Ovarian_Aging_Microbiome.

Figure legends

Fig 1. Characterization of ovarian health state and fecal microbial profiles of young and

estropausal female mice. (A) Schematic diagram of the experimental setup of female aging cohorts.

Resources from Flaticon.com were used. (B) Boxplot of combined follicle count of young and estropausal

mice. (C) Boxplot of serum concentrations of AMH. (D) Boxplot of ovarian health index of young and

estropausal mice. (E) Principal component analysis result of CLR-transformed and batch-corrected ASV

counts from young and estropausal female mice. Animals from four independent cohorts (C1-C4), n=19-

20 per group (variation due to animal death prior to experiment; n=20 for young females; n=19 for

estropausal females). (F) Boxplots of Observed features and Shannon entropy indices of young and

estropausal female mice. Medians from young females for each cohort were used to normalize indices

per cohort. (G) Differential abundance analysis results of microbial genera of young vs. estropausal

female mice using ALDEx2. (H) Functional abundance prediction analysis of young vs. estropausal

female mice using PICRUSt2. Significance in nonparametric two-sided Wilcoxon rank-sum tests are

reported for (B) and (D). YF: Young female; EF: Estropausal female. C: Cohort.

Fig 2. Bulk RNA-seq analysis of ovaries from young and estropausal female mice fecal microbiota

transplantation recipient mice. (A) Schematic diagram of the experimental setup of fecal microbiota

transplantation (FMT) cohort. Fecal microbiota from either young (4 months old) or estropausal (20

months old) female mice were transplanted into young (4 months old) female recipient mice. FMT

recipient groups are referred to as FMT-YF (receiving young donor microbiota) and FMT-EF (receiving

estropausal donor microbiota). (B) Multidimensional scaling (MDS) analysis result of RNA expression

between young and estropausal female mice FMT recipient mice (FMT-YF and FMT-EF, respectively).

(C-F) Top GSEA enriched terms from (C) Gene Ontology Biological Process (GO BP), (D) Gene Ontology

Molecular Function (GO MF), (E) Gene Ontology Cellular Component (GO CC) and (F) Gene

Transcription Regulation Database (GTRD – TF targets). Only up to top ten most-upregulated and top ten

most-downregulated gene sets are plotted for readability.

Fig 3. Analysis of public ovarian aging bulk RNA-seq datasets. (A) Schematic diagram of the analysis setup of public ovarian aging bulk RNA-seq datasets, CRA003645 and PRJNA1002222, from young (2-3-months) and estropausal (12-months) female mice. (B-C) GSEA enrichment score plots of (B) genes that are up-regulated with ovarian aging ("UP with ovarian aging", left panel) and down-regulated with ovarian aging ("DOWN with ovarian aging, right panel), and (C) SenMayo gene set. In both (B) and (C), the black vertical lines indicate the positions of the genes from the gene set within the ranked list of genes. The curve represents the running enrichment score (ES), with the peak indicating the point of maximum enrichment for the gene set. The color scale at the bottom of the plot reflects the ranking metric, where red represents upregulated genes and blue represents downregulated genes. The p-value shown is calculated based on GSEA and adjusted for multiple testing using the Benjamini-Hochberg method. DGE: Differential gene expression. (D) Boxplots of DESeq2 normalized $\log_2$ counts of *Cd38*, *Cdkn1a* and *Il1a*.

Fig 4. Evaluation of fertility and ovarian health state of fecal microbiota transplantation recipient mice. (A) Schematic diagram of the experimental setup. (B) Boxplot of ovarian health index of FMT-YF and FMT-EF mice. Ovarian health index was calculated for two independent cohorts (n=5 or 7 per group and 10 or 14 per cohort). (C) Boxplot of pup counts of first litters of young and estropausal female mice fecal microbiota transplantation recipient mice (FMT-YF and FMT-EF, respectively). Pup counts were measured from two independent cohorts (n=8 or 10 per group and 16 or 20 per cohort). (D) Reproductive success rate analysis result shown as percentages of nulliparous between FMT-YF and FMT-EF. Significance in nonparametric two-sided Wilcoxon rank-sum tests are reported for (B) and (C), and logrank test for (D).

Fig 5. Characterization of fecal microbial profiles of fecal microbiota transplantation recipient mice. (A) Schematic diagram of the experimental setup. (B) Principal component analysis result of CLR-transformed and batch-corrected counts from young and estropausal female mice fecal microbiota transplantation recipient mice (FMT-YF and FMT-EF, respectively). (C) Principal coordinate analysis results of Bray-Curtis dissimilarity (left panel) and Jaccard (right panel) indices. (D) Boxplots of Observed features and Shannon entropy indices of FMT-YF and FMT-EF mice. (E) Volcano plot of differential

1018 abundance analysis results of microbial species of FMT-YF vs. FMT-EF mice using ALDEx2. (F) Bubble

1019 plot of top 10 up- and down-regulated microbial species in FMT-EF. (G) Functional abundance analysis of

1020 FMT-YF vs. FMT-EF mice using HUMAnN3 (Q-value < 0.1). Significance in nonparametric two-sided

1021 Wilcoxon rank-sum tests are reported for (D).

1022

Fig 6. Characterization of serum metabolic profiles of fecal microbiota transplantation recipient

mice. (A) Schematic diagram of the experimental setup. (B) PCA plots of identified serum metabolites

from two independent FMT cohorts. (C) LDA2 projection scatter plot displaying LDA1 scores for the FMT-

YF and FMT-EF groups. (D) Top 15 metabolites contributing to LDA-based separation, ranked by their

positive and negative contributions to the LDA axis. (E) Boxplots of 3,4-dihydroxyphenylpropionic acid

(3,4-DHPPA) and lactamide serum levels between FMT-YF and FMT-EF groups. Significance in

nonparametric two-sided Wilcoxon rank-sum tests are reported for (E).

Fig 7. Causal mediation analysis of microbial species and ovarian transcriptome changes in fecal

microbiota transplantation recipient mice. (A) Graphical representation of causal mediation analysis.

(B) Causal mediation analysis results of whole genome shotgun metagenomics counts that showed

significant mediation effects on changes in the ovarian transcriptome in fecal microbiota transplantation

recipient mice. Nonparametric bootstrap confidence intervals with the percentile method are reported. (C)

Heatmap of Spearman correlation analysis between CLR-transformed WGS count data and serum

hormone quantification data. #: p-value < 0.1; *: p-value < 0.05. (D) Summary diagram. The estropausal

female fecal microbiota exhibits increased levels of commensal microbes and metabolites through

compensatory mechanisms. As a result, fecal transplantation of estropausal microbiota enhances ovarian

health measures. Potential mechanisms, including NAD and vitamin K synthesis pathways, should be

further examined. Generic Diagramming Platform (GDP) was used to generate the summary diagram[138].

ACME: Average Causal Mediated Effect; ADE: Average Direct Effect.

Extended Data Fig. legend

1043

Extended Data Fig. 1. Characterization of ovarian health state of young and estropausal female

mice. (A) Schematic diagram of ovarian health index calculation. (B) Representative images of ovarian

hematoxylin-eosin stained sections. Scale bar: 50 μm. (C) Boxplots of ovarian follicle counts per section.

The schematic diagrams of the follicles were created using the Generic Diagramming Platform (GDP)[138].

(D) Boxplots of serum concentrations of FSH and Inhibin A. Significance in nonparametric two-sided

Wilcoxon rank-sum tests are reported for (C) and (D). YF: Young female; EF: Estropausal female.

1050

1051   **Extended Data Fig. 2. Characterization of fecal microbial profiles of young and estropausal female**

1052   **mice.** (A) Principal coordinate analysis results of Bray-Curtis dissimilarity (top panel) and Jaccard (bottom

1053   panel) indices. (B) Boxplots of base abundance of differentially abundant microbial genera. YF: Young

1054   female; EF: Estropausal female.

1055

1056   **Extended Data Fig. 3. Characterization of fecal microbial profiles of young and estropausal female**

1057   **mice.** Boxplots of base abundance of differentially abundant microbial genera. YF: Young female; EF:

1058   Estropausal female.

1059

Extended Data Fig. 4. Characterization of ovarian health state of VCD cohort mice. (A) Schematic

diagram of the experimental setup of VCD cohort. (B) Representative images of ovarian hematoxylin-

eosin stained sections. Scale bar: 50 μm. (C-F) Boxplots of (C) ovarian follicle counts per section, (D)

combined follicle counts, (E) ovarian health index, and (E) serum concentrations of AMH, FSH and Inhibin

A of control and VCD-injected mice. The schematic diagrams of the follicles were created using the

Generic Diagramming Platform (GDP)[138]. Significance in nonparametric two-sided Wilcoxon rank-sum

tests are reported for (C-F).

1067

1068   **Extended Data Fig. 5. Characterization of fecal microbial profiles of VCD cohort mice.** (A) Principal

1069   component analysis result of CLR-transformed and batch-corrected ASV counts from control and VCD-

injected mice. (B) Principal coordinate analysis results of Bray-Curtis dissimilarity (top panel) and Jaccard (bottom panel) indices. (C, D) Boxplots of normalized observed features and Shannon entropy indices of (C) females, and (D) males. Median from control-injected mice was used to normalize indices. (E) Differential abundance analysis results of microbial genera of control vs. VCD-injected female mice using ALDEx2. (F) Functional abundance prediction analysis of control vs. VCD-injected female mice using PICRUSt2. Significance in nonparametric two-sided Wilcoxon rank-sum tests are reported for (C,D).

Extended Data Fig. 6. Differential abundance analysis of 16S amplicon sequencing data of VCD cohort female mice. Boxplots of base abundance of differentially abundant microbial genera between control and VCD-injected female mice.

Extended Data Fig. 7. Characterization of fecal microbial profiles of young and old male mice. (A) Schematic diagram of the experimental setup of male aging cohorts. (B) Principal component analysis result of CLR-transformed and batch-corrected ASV counts from young and old male mice. Animals from three independent cohorts, n=14 per group (variation due to animal death or abnormal health issues prior to experiment). (C) Principal coordinate analysis results of Bray-Curtis dissimilarity (top panel) and Jaccard (bottom panel) indices. (D) Boxplots of observed features and Shannon entropy indices of young and old male mice. Medians from young males for each cohort were used to normalize indices per cohort. (E) Differential abundance analysis results of microbial genera of young and old male mice using ALDEx2. (F) Functional abundance prediction analysis of young vs. old male mice using PICRUSt2. Significance in nonparametric two-sided Wilcoxon rank-sum tests are reported for (D). YM: Young male; OM: Old male.

Extended Data Fig. 8. Differential abundance analysis of 16S amplicon sequencing data of young and old male mice. Boxplots of base abundance of differentially abundant microbial genera between young and old male mice.

Extended Data Fig. 9. Bulk RNA-seq analysis of ovaries from young and estropausal female mice fecal microbiota transplantation recipient mice. (A) Heat map of significant (FDR < 5%) genes

1098 differentially expressed in FMT-YF vs. FMT-EF mice. (B) Dot plots of immune cell proportions in

1099 percentages between young and estropausal fecal microbiota transplantation recipient mice (FMT-YF and

1100 FMT-EF, respectively) predicted by deconvolution analyses using CSCDRNA (left panel) and Granulator

1101 (right panel). (C) GSEA enrichment score plots Ncoa1-peak (left panel) and Usp7-peak (right panel)

1102 genes. (D) Bar plots of normalized $\log_2$ counts of *Ncoa1* and *Usp7* expression.

1103

Extended Data Fig. 10. Characterization of ovarian health state of FMT cohort mice. (A)

Representative images of ovarian hematoxylin-eosin stained sections. Scale bar: 50 μm. (B-E) Boxplots

of (B) combined ovarian follicle counts, (C) ovarian follicle counts per section, (D) serum concentrations of

AMH, FSH (Multiplex assay), FSH (Ultra-sensitive assay), FSH combined and INHBA, and (E) weight

percentage difference between day 0 and 16 of FMT regimen. The schematic diagrams of the follicles

were created using the Generic Diagramming Platform (GDP)[138].

1110

Extended Data Fig. 11. Characterization of fecal microbial profiles of FMT cohort mice. (A) Principal

component analysis result of CLR-transformed and batch-corrected counts from young and estropausal

female mice fecal microbiota transplantation recipient mice (FMT-YF and FMT-EF, respectively). (B)

Principal coordinate analysis results of Bray-Curtis dissimilarity (left panel) and Jaccard (right panel)

indices. (C) Boxplots of observed features and Shannon entropy indices of FMT-YF and FMT-EF mice. (D)

Differential abundance analysis results of microbial species of FMT-YF vs. FMT-EF mice using ALDEx2.

(E) Venn diagram of overlapping microbial genera between genera that increase in abundance with

female aging ("UP in EF") and increase in abundance in FMT-EF recipient mice ("UP in FMT-EF"). (F)

Functional abundance prediction analysis of FMT-YF vs. FMT-EF mice using PICRUSt2. Significance in

nonparametric two-sided Wilcoxon rank-sum tests are reported for (C). YF: Young female; EF:

Estropausal female.

1122

1123 **Extended Data Fig. 12. Differential abundance analysis of 16S amplicon sequencing data of FMT**

1124 **cohort mice.** Boxplots of base abundance of differentially abundant microbial genera between FMT-YF

1125 and FMT-EF mice.

1126

1127 **Extended Data Fig. 13. Differential abundance analysis of WGS data of FMT cohort mice.** Boxplots

1128 of base abundance of differentially abundant microbial species between FMT-YF and FMT-EF mice.

1129

1130 **Extended Data Fig. 14. Stool vitamin K quantification of FMT cohort mice.** (A) Stool concentrations

1131 of combined vitamin K (MK6-MK13) of FMT-YF and FMT-EF. (B) Fold change in stool concentrations of

1132 vitamin K (MK6-MK13) of FMT-YF and FMT-EF.

1133

1134 **Extended Data Fig. 15. Serum untargeted metabolomics analysis of FMT cohort mice.** Boxplots of

1135 metabolite abundances of top 15 metabolites contributing to LDA separation between FMT-YF and FMT-

1136 EF.

1137

Extended Data Fig. 16. Causal mediation analysis of microbial species and ovarian transcriptome

changes in fecal microbiota transplantation recipient mice. (A) Graphical representation of causal

mediation analysis. (B) Causal mediation analysis results of 16S amplicon ASVs that showed significant

mediation effects on changes in the ovarian transcriptome in fecal microbiota transplantation recipient

mice. Nonparametric bootstrap confidence intervals with the percentile method are reported. ACME:

Average Causal Mediated Effect; ADE: Average Direct Effect.

Extended Data Table

1145

1146 Extended Data Table 1. Weight measurements of FMT cohort.

1147 Extended Data Table 2. Estrous cycle staging data of FMT cohort.

1148 Extended Data Table 3. ALDEx2 differential abundance analysis result of metagenomics data of FMT

1149 cohort.

1150 Extended Data Table 4. MS2-filtered metabolite abundance matrix.

1151

References

References

1 Davis, S. R. *et al.* Menopause. *Nat Rev Dis Primers* **1**, 15004, doi:10.1038/nrdp.2015.4
(2015).

2 Broekmans, F. J., Soules, M. R. & Fauser, B. C. Ovarian aging: mechanisms and clinical
consequences. *Endocr Rev* **30**, 465-493, doi:10.1210/er.2009-0006 (2009).

3 Faddy, M. J. & Gosden, R. G. A model conforming the decline in follicle numbers to the
age of menopause in women. *Hum Reprod* **11**, 1484-1486,
doi:10.1093/oxfordjournals.humrep.a019422 (1996).

4 Fu, C. *et al.* Association of reproductive factors with dementia: A systematic review and
dose-response meta-analyses of observational studies. *EClinicalMedicine* **43**, 101236,
doi:10.1016/j.eclinm.2021.101236 (2022).

5 Shuster, L. T., Rhodes, D. J., Gostout, B. S., Grossardt, B. R. & Rocca, W. A. Premature
menopause or early menopause: long-term health consequences. *Maturitas* **65**, 161-166,
doi:10.1016/j.maturitas.2009.08.003 (2010).

6 Muka, T. *et al.* Association of Age at Onset of Menopause and Time Since Onset of
Menopause With Cardiovascular Outcomes, Intermediate Vascular Traits, and All-Cause
Mortality: A Systematic Review and Meta-analysis. *JAMA Cardiol* **1**, 767-776,
doi:10.1001/jamacardio.2016.2415 (2016).

7 Ossewaarde, M. E. *et al.* Age at menopause, cause-specific mortality and total life
expectancy. *Epidemiology* **16**, 556-562, doi:10.1097/01.ede.0000165392.35273.d4
(2005).

8 Hong, J. S. *et al.* Age at menopause and cause-specific mortality in South Korean women:
Kangwha Cohort Study. *Maturitas* **56**, 411-419, doi:10.1016/j.maturitas.2006.11.004
(2007).

9 Knobil, E. The neuroendocrine control of the menstrual cycle. *Recent Prog Horm Res* **36**,
53-88, doi:10.1016/b978-0-12-571136-4.50008-5 (1980).

10 Santoro, N., Roeca, C., Peters, B. A. & Neal-Perry, G. The Menopause Transition: Signs,
Symptoms, and Management Options. *J Clin Endocrinol Metab* **106**, 1-15,
doi:10.1210/clinem/dgaa764 (2021).

11 Randolph, J. F., Jr. *et al.* Change in follicle-stimulating hormone and estradiol across the
menopausal transition: effect of age at the final menstrual period. *J Clin Endocrinol*
Metab **96**, 746-754, doi:10.1210/jc.2010-1746 (2011).

12 Kelsey, T. W., Wright, P., Nelson, S. M., Anderson, R. A. & Wallace, W. H. A validated
model of serum anti-mullerian hormone from conception to menopause. *PLoS One* **6**,
e22024, doi:10.1371/journal.pone.0022024 (2011).

13 Weiss, G., Skurnick, J. H., Goldsmith, L. T., Santoro, N. F. & Park, S. J. Menopause and
hypothalamic-pituitary sensitivity to estrogen. *JAMA* **292**, 2991-2996,
doi:10.1001/jama.292.24.2991 (2004).

14 Xiong, J. *et al.* FSH blockade improves cognition in mice with Alzheimer's disease. *Nature*
603, 470-476, doi:10.1038/s41586-022-04463-0 (2022).

15 Mao, L., Wang, L., Bennett, S., Xu, J. & Zou, J. Effects of follicle-stimulating hormone on
fat metabolism and cognitive impairment in women during menopause. *Front Physiol* **13**,
1043237, doi:10.3389/fphys.2022.1043237 (2022).

16 Xiong, J. *et al.* FSH and ApoE4 contribute to Alzheimer's disease-like pathogenesis via
C/EBPbeta/delta-secretase in female mice. *Nat Commun* **14**, 6577, doi:10.1038/s41467-
023-42282-7 (2023).
17 Luckey, T. D. Introduction to intestinal microecology. *Am J Clin Nutr* **25**, 1292-1294,
doi:10.1093/ajcn/25.12.1292 (1972).
18 Savage, D. C. Microbial ecology of the gastrointestinal tract. *Annu Rev Microbiol* **31**, 107-
133, doi:10.1146/annurev.mi.31.100177.000543 (1977).
19 Sender, R., Fuchs, S. & Milo, R. Are We Really Vastly Outnumbered? Revisiting the Ratio
of Bacterial to Host Cells in Humans. *Cell* **164**, 337-340, doi:10.1016/j.cell.2016.01.013
(2016).
20 Biagi, E. *et al.* Ageing and gut microbes: perspectives for health maintenance and
longevity. *Pharmacol Res* **69**, 11-20, doi:10.1016/j.phrs.2012.10.005 (2013).
21 Zapata, H. J. & Quagliarello, V. J. The microbiota and microbiome in aging: potential
implications in health and age-related diseases. *J Am Geriatr Soc* **63**, 776-781,
doi:10.1111/jgs.13310 (2015).
22 Parker, A. *et al.* Fecal microbiota transfer between young and aged mice reverses
hallmarks of the aging gut, eye, and brain. *Microbiome* **10**, 68, doi:10.1186/s40168-022-
01243-w (2022).
23 Kundu, P. *et al.* Neurogenesis and prolongevity signaling in young germ-free mice
transplanted with the gut microbiota of old mice. *Sci Transl Med* **11**,
doi:10.1126/scitranslmed.aau4760 (2019).
24 Wu, J. *et al.* Association between premature ovarian insufficiency and gut microbiota.
BMC Pregnancy Childbirth **21**, 418, doi:10.1186/s12884-021-03855-w (2021).
25 Qi, X. *et al.* Gut microbiota-bile acid-interleukin-22 axis orchestrates polycystic ovary
syndrome. *Nat Med* **25**, 1225-1233, doi:10.1038/s41591-019-0509-0 (2019).
26 Weger, B. D. *et al.* The Mouse Microbiome Is Required for Sex-Specific Diurnal Rhythms
of Gene Expression and Metabolism. *Cell Metab* **29**, 362-382 e368,
doi:10.1016/j.cmet.2018.09.023 (2019).
27 Lombardi, P., Goldin, B., Boutin, E. & Gorbach, S. L. Metabolism of androgens and
estrogens by human fecal microorganisms. *J Steroid Biochem* **9**, 795-801,
doi:10.1016/0022-4731(78)90203-0 (1978).
28 Adlercreutz, H., Martin, F., Jarvenpaa, P. & Fotsis, T. Steroid absorption and
enterohepatic recycling. *Contraception* **20**, 201-223, doi:10.1016/0010-7824(79)90094-5
(1979).
29 Eriksson, H. & Gustafsson, J. A. Steroids in germfree and conventional rats. Distribution
and excretion of labelled pregnenolone and corticosterone in male and female rats. *Eur J
Biochem* **15**, 132-139, doi:10.1111/j.1432-1033.1970.tb00987.x (1970).
30 Finch, C. E. The menopause and aging, a comparative perspective. *J Steroid Biochem Mol
Biol* **142**, 132-141, doi:10.1016/j.jsbmb.2013.03.010 (2014).
31 Whitehead, J. C. *et al.* A clinical frailty index in aging mice: comparisons with frailty index
data in humans. *J Gerontol A Biol Sci Med Sci* **69**, 621-632, doi:10.1093/gerona/glt136
(2014).
32 Wallace, W. H. & Kelsey, T. W. Human ovarian reserve from conception to the
menopause. *PLoS One* **5**, e8772, doi:10.1371/journal.pone.0008772 (2010).

33   Hirshfield, A. N. Development of follicles in the mammalian ovary. *Int Rev Cytol* **124**, 43-
101, doi:10.1016/s0074-7696(08)61524-7 (1991).

34   Tehrani, F. R., Mansournia, M. A., Solaymani-Dodaran, M. & Azizi, F. Age-specific serum
anti-Mullerian hormone levels: estimates from a large population-based sample.
Climacteric **17**, 591-597, doi:10.3109/13697137.2014.912262 (2014).

35   Broer, S. L., Broekmans, F. J., Laven, J. S. & Fauser, B. C. Anti-Mullerian hormone: ovarian
reserve testing and its potential clinical implications. *Hum Reprod Update* **20**, 688-701,
doi:10.1093/humupd/dmu020 (2014).

36   Felicio, L. S., Nelson, J. F. & Finch, C. E. Prolongation and cessation of estrous cycles in
aging C57BL/6J mice are differentially regulated events. *Biol Reprod* **34**, 849-858,
doi:10.1095/biolreprod34.5.849 (1986).

37   Johnson, J. S. *et al.* Evaluation of 16S rRNA gene sequencing for species and strain-level
microbiome analysis. *Nat Commun* **10**, 5029, doi:10.1038/s41467-019-13036-1 (2019).

38   Gupta, S. *et al.* Amplicon sequencing provides more accurate microbiome information in
healthy children compared to culturing. *Commun Biol* **2**, 291, doi:10.1038/s42003-019-
0540-1 (2019).

39   Knight, R. *et al.* Best practices for analysing microbiomes. *Nat Rev Microbiol* **16**, 410-422,
doi:10.1038/s41579-018-0029-9 (2018).

40   Willis, A. D. Rarefaction, Alpha Diversity, and Statistics. *Front Microbiol* **10**, 2407,
doi:10.3389/fmicb.2019.02407 (2019).

41   Wang, Y. & LeCao, K. A. Managing batch effects in microbiome data. *Brief Bioinform* **21**,
1954-1970, doi:10.1093/bib/bbz105 (2020).

42   Gamez-Macias, P. E. *et al.* Intestinal Permeability, Gut Inflammation, and Gut Immune
System Response Are Linked to Aging-Related Changes in Gut Microbiota Composition: A
Study in Female Mice. *J Gerontol A Biol Sci Med Sci* **79**, doi:10.1093/gerona/glae045
(2024).

43   de la Cuesta-Zuluaga, J. *et al.* Age- and Sex-Dependent Patterns of Gut Microbial
Diversity in Human Adults. *mSystems* **4**, doi:10.1128/mSystems.00261-19 (2019).

44   Ghosh, T. S., Shanahan, F. & O'Toole, P. W. Toward an improved definition of a healthy
microbiome for healthy aging. *Nat Aging* **2**, 1054-1069, doi:10.1038/s43587-022-00306-
9 (2022).

45   Luo, M. *et al.* E. coli Nissle 1917 ameliorates mitochondrial injury of granulosa cells in
polycystic ovary syndrome through promoting gut immune factor IL-22 via gut
microbiota and microbial metabolism. *Front Immunol* **14**, 1137089,
doi:10.3389/fimmu.2023.1137089 (2023).

46   Biagi, E. *et al.* Gut Microbiota and Extreme Longevity. *Curr Biol* **26**, 1480-1485,
doi:10.1016/j.cub.2016.04.016 (2016).

47   Wang, N. *et al.* Enriched taxa were found among the gut microbiota of centenarians in
East China. *PLoS One* **14**, e0222763, doi:10.1371/journal.pone.0222763 (2019).

48   Douglas, G. M. *et al.* PICRUSt2 for prediction of metagenome functions. *Nat Biotechnol*
38, 685-688, doi:10.1038/s41587-020-0548-6 (2020).

49   Murali, R. *et al.* Deregulated Metabolic Pathways in Ovarian Cancer: Cause and
Consequence. *Metabolites* **13**, doi:10.3390/metabo13040560 (2023).

50   Ji, Z. *et al.* Deregulation of Lipid Metabolism: The Critical Factors in Ovarian Cancer. *Front*
Oncol **10**, 593017, doi:10.3389/fonc.2020.593017 (2020).

51   Borman, S. M., Christian, P. J., Sipes, I. G. & Hoyer, P. B. Ovotoxicity in female Fischer rats
and B6 mice induced by low-dose exposure to three polycyclic aromatic hydrocarbons:
comparison through calculation of an ovotoxic index. *Toxicol Appl Pharmacol* **167**, 191-
198, doi:10.1006/taap.2000.9006 (2000).

52   Van Kempen, T. A., Milner, T. A. & Waters, E. M. Accelerated ovarian failure: a novel,
chemically induced animal model of menopause. *Brain Res* **1379**, 176-187,
doi:10.1016/j.brainres.2010.12.064 (2011).

53   Mayer, L. P., Devine, P. J., Dyer, C. A. & Hoyer, P. B. The follicle-deplete mouse ovary
produces androgen. *Biol Reprod* **71**, 130-138, doi:10.1095/biolreprod.103.016113 (2004).

54   Boehme, M. *et al.* Microbiota from young mice counteracts selective age-associated
behavioral deficits. *Nat Aging* **1**, 666-676, doi:10.1038/s43587-021-00093-9 (2021).

55   Subramanian, A. *et al.* Gene set enrichment analysis: a knowledge-based approach for
interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-
15550, doi:10.1073/pnas.0506580102 (2005).

56   Morris, M. E. *et al.* A single-cell atlas of the cycling murine ovary. *Elife* **11**,
doi:10.7554/eLife.77239 (2022).

57   Jew, B. *et al.* Accurate estimation of cell composition in bulk expression through robust
integration of single-cell information. *Nat Commun* **11**, 1971, doi:10.1038/s41467-020-
15816-6 (2020).

58   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.
Nat Methods **12**, 453-457, doi:10.1038/nmeth.3337 (2015).

59   Lovasco, L. A. *et al.* Accelerated ovarian aging in the absence of the transcription
regulator TAF4B in mice. *Biol Reprod* **82**, 23-34, doi:10.1095/biolreprod.109.077495
(2010).

60   Baumann, C. *et al.* Acute irradiation induces a senescence-like chromatin structure in
mammalian oocytes. *Commun Biol* **6**, 1258, doi:10.1038/s42003-023-05641-0 (2023).

61   Yevshin, I., Sharipov, R., Valeev, T., Kel, A. & Kolpakov, F. GTRD: a database of
transcription factor binding sites identified by ChIP-seq experiments. *Nucleic Acids Res*
45, D61-D67, doi:10.1093/nar/gkw951 (2017).

62   Zheng, R. *et al.* Cistrome Data Browser: expanded datasets and new tools for gene
regulatory analysis. *Nucleic Acids Res* **47**, D729-D735, doi:10.1093/nar/gky1094 (2019).

63   Mei, S. *et al.* Cistrome Data Browser: a data portal for ChIP-Seq and chromatin
accessibility data in human and mouse. *Nucleic Acids Res* **45**, D658-D662,
doi:10.1093/nar/gkw983 (2017).

64   Huang, H. Y. *et al.* Association of polymorphisms for nuclear receptor coactivator 1 gene
with egg production traits in the maternal line of Shaobo hens. *Br Poult Sci* **52**, 328-332,
doi:10.1080/00071668.2011.577057 (2011).

65   Wang, M. *et al.* The USP7 Inhibitor P5091 Induces Cell Death in Ovarian Cancers with
Different P53 Status. *Cell Physiol Biochem* **43**, 1755-1766, doi:10.1159/000484062 (2017).

66   Briley, S. M. *et al.* Reproductive age-associated fibrosis in the stroma of the mammalian
ovary. *Reproduction* **152**, 245-260, doi:10.1530/REP-16-0129 (2016).

67 Zhang, Z., Schlamp, F., Huang, L., Clark, H. & Brayboy, L. Inflammaging is associated with
shifted macrophage ontogeny and polarization in the aging mouse ovary. *Reproduction*
159, 325-337, doi:10.1530/REP-19-0330 (2020).
68 Zhou, Z. *et al.* Temporal transcriptomic landscape of postnatal mouse ovaries reveals
dynamic gene signatures associated with ovarian aging. *Hum Mol Genet* **30**, 1941-1954,
doi:10.1093/hmg/ddab163 (2021).
69 Yang, Q. *et al.* NADase CD38 is a key determinant of ovarian aging. *Nat Aging* **4**, 110-128,
doi:10.1038/s43587-023-00532-9 (2024).
70 Saul, D. *et al.* A new gene set identifies senescent cells and predicts senescence-
associated pathways across tissues. *Nat Commun* **13**, 4827, doi:10.1038/s41467-022-
32552-1 (2022).
71 Mehalko, K. *et al.* Lack of accelerated ovarian aging in a follicle-stimulating hormone
receptor haploinsufficiency model. *Transl Med Aging* **7**, 1-8,
doi:10.1016/j.tma.2023.01.001 (2023).
72 Wrzosek, L. *et al.* Transplantation of human microbiota into conventional mice durably
reshapes the gut microbiota. *Sci Rep* **8**, 6854, doi:10.1038/s41598-018-25300-3 (2018).
73 Gong, X., Gutala, R. & Jaiswal, A. K. Quinone oxidoreductases and vitamin K metabolism.
Vitam Horm **78**, 85-101, doi:10.1016/S0083-6729(07)00005-2 (2008).
74 Beghini, F. *et al.* Integrating taxonomic, functional, and strain-level profiling of diverse
microbial communities with bioBakery 3. *Elife* **10**, doi:10.7554/eLife.65088 (2021).
75 Dekkers, K. F. *et al.* An online atlas of human plasma metabolite signatures of gut
microbiome composition. *Nat Commun* **13**, 5370, doi:10.1038/s41467-022-33050-0
(2022).
76 Pilley, S. E. *et al.* A metabolic atlas of mouse aging. *bioRxiv*, 2024.2005.2004.592445,
doi:10.1101/2024.05.04.592445 (2024).
77 Marquez Campos, E., Stehle, P. & Simon, M. C. Microbial Metabolites of Flavan-3-Ols and
Their Biological Activity. *Nutrients* **11**, doi:10.3390/nu11102260 (2019).
78 He, Y. *et al.* Lactic acid bacteria alleviate polycystic ovarian syndrome by regulating sex
hormone related gut microbiota. *Food Funct* **11**, 5192-5204, doi:10.1039/c9fo02554e
(2020).
79 Dikalov, S., Panov, A. & Dikalova, A. Critical Role of Mitochondrial Fatty Acid Metabolism
in Normal Cell Function and Pathological Conditions. *Int J Mol Sci* **25**,
doi:10.3390/ijms25126498 (2024).
80 Wang, Z. H. *et al.* Targeting mitochondria for ovarian aging: new insights into
mechanisms and therapeutic potential. *Front Endocrinol (Lausanne)* **15**, 1417007,
doi:10.3389/fendo.2024.1417007 (2024).
81 Zhang, I. W. *et al.* Palmitoylcarnitine impairs immunity in decompensated cirrhosis. *JHEP*
Rep **6**, 101187, doi:10.1016/j.jhepr.2024.101187 (2024).
82 Xu, X. L. *et al.* Estrogen Biosynthesis and Signal Transduction in Ovarian Disease. *Front*
Endocrinol (Lausanne) **13**, 827032, doi:10.3389/fendo.2022.827032 (2022).
83 Stachenfeld, N. S. Hormonal changes during menopause and the impact on fluid
regulation. *Reprod Sci* **21**, 555-561, doi:10.1177/1933719113518992 (2014).

84   Liu, T., Bowen, R. L., Wilson, A. C. & Atwood, C. S. Estropause, Sex Hormones and Metal
1368   Homeostasis in the Mouse Brain. *Front Neurol* **13**, 841822,
1369   doi:10.3389/fneur.2022.841822 (2022).
1370 85   Schele, E. *et al.* The gut microbiota reduces leptin sensitivity and the expression of the
1371   obesity-suppressing neuropeptides proglucagon (Gcg) and brain-derived neurotrophic
1372   factor (Bdnf) in the central nervous system. *Endocrinology* **154**, 3643-3651,
1373   doi:10.1210/en.2012-2151 (2013).
1374 86   Fan, K. C. *et al.* Altered gut microbiota in older adults with mild cognitive impairment: a
1375   case-control study. *Front Aging Neurosci* **15**, 1162057, doi:10.3389/fnagi.2023.1162057
1376   (2023).
1377 87   Ellis, J. L. *et al.* Dietary vitamin K is remodeled by gut microbiota and influences
1378   community composition. *Gut Microbes* **13**, 1-16, doi:10.1080/19490976.2021.1887721
1379   (2021).
1380 88   Tarkesh, F., Namavar Jahromi, B., Hejazi, N. & Tabatabaee, H. Beneficial health effects of
1381   Menaquinone-7 on body composition, glycemic indices, lipid profile, and endocrine
1382   markers in polycystic ovary syndrome patients. *Food Sci Nutr* **8**, 5612-5621,
1383   doi:10.1002/fsn3.1837 (2020).
1384 89   Sibayama-Imazu, T. *et al.* Induction of apoptosis in PA-1 ovarian cancer cells by vitamin
1385   K2 is associated with an increase in the level of TR3/Nur77 and its accumulation in
1386   mitochondria and nuclei. *J Cancer Res Clin Oncol* **134**, 803-812, doi:10.1007/s00432-007-
1387   0349-z (2008).
1388 90   Harshman, S. G. & Shea, M. K. The Role of Vitamin K in Chronic Aging Diseases:
1389   Inflammation, Cardiovascular Disease, and Osteoarthritis. *Curr Nutr Rep* **5**, 90-98,
1390   doi:10.1007/s13668-016-0162-x (2016).
1391 91   Sampathkumar, N. K. *et al.* Widespread sex dimorphism in aging and age-related
1392   diseases. *Hum Genet* **139**, 333-356, doi:10.1007/s00439-019-02082-w (2020).
1393 92   Podcasy, J. L. & Epperson, C. N. Considering sex and gender in Alzheimer disease and
1394   other dementias. *Dialogues Clin Neurosci* **18**, 437-446,
1395   doi:10.31887/DCNS.2016.18.4/cepperson (2016).
1396 93   Vina, J. & Lloret, A. Why women have more Alzheimer's disease than men: gender and
1397   mitochondrial toxicity of amyloid-beta peptide. *J Alzheimers Dis* **20 Suppl 2**, S527-533,
1398   doi:10.3233/JAD-2010-100501 (2010).
1399 94   Varghese, M. *et al.* Female adipose tissue has improved adaptability and metabolic
1400   health compared to males in aged obesity. *Aging (Albany NY)* **12**, 1725-1746,
1401   doi:10.18632/aging.102709 (2020).
1402 95   Austad, S. N. & Fischer, K. E. Sex Differences in Lifespan. *Cell Metab* **23**, 1022-1033,
1403   doi:10.1016/j.cmet.2016.05.019 (2016).
1404 96   Lemaitre, J. F. *et al.* Sex differences in adult lifespan and aging rates of mortality across
1405   wild mammals. *Proc Natl Acad Sci U S A* **117**, 8546-8553, doi:10.1073/pnas.1911999117
1406   (2020).
1407 97   Markle, J. G. *et al.* Sex differences in the gut microbiome drive hormone-dependent
1408   regulation of autoimmunity. *Science* **339**, 1084-1088, doi:10.1126/science.1233521
1409   (2013).

98   Steegenga, W. T. *et al.* Sexually dimorphic characteristics of the small intestine and colon
of prepubescent C57BL/6 mice. *Biol Sex Differ* **5**, 11, doi:10.1186/s13293-014-0011-9
(2014).
99   Org, E. *et al.* Sex differences and hormonal effects on gut microbiota composition in
mice. *Gut Microbes* **7**, 313-322, doi:10.1080/19490976.2016.1203502 (2016).
100   Vemuri, R. *et al.* The microgenderome revealed: sex differences in bidirectional
interactions between the microbiota, hormones, immunity and disease susceptibility.
Semin Immunopathol **41**, 265-275, doi:10.1007/s00281-018-0716-7 (2019).
101   Kaliannan, K. *et al.* Estrogen-mediated gut microbiome alterations influence sexual
dimorphism in metabolic syndrome in mice. *Microbiome* **6**, 205, doi:10.1186/s40168-
018-0587-0 (2018).
102   Hogan, K. A., Chini, C. C. S. & Chini, E. N. The Multi-faceted Ecto-enzyme CD38: Roles in
Immunomodulation, Cancer, Aging, and Metabolic Diseases. *Front Immunol* **10**, 1187,
doi:10.3389/fimmu.2019.01187 (2019).
103   Camacho-Pereira, J. *et al.* CD38 Dictates Age-Related NAD Decline and Mitochondrial
Dysfunction through an SIRT3-Dependent Mechanism. *Cell Metab* **23**, 1127-1139,
doi:10.1016/j.cmet.2016.05.006 (2016).
104   Gomes, A. P. *et al.* Declining NAD(+) induces a pseudohypoxic state disrupting nuclear-
mitochondrial communication during aging. *Cell* **155**, 1624-1638,
doi:10.1016/j.cell.2013.11.037 (2013).
105   Chini, C. C. S. *et al.* CD38 ecto-enzyme in immune cells is induced during aging and
regulates NAD(+) and NMN levels. *Nat Metab* **2**, 1284-1304, doi:10.1038/s42255-020-
00298-z (2020).
106   Covarrubias, A. J. *et al.* Senescent cells promote tissue NAD(+) decline during ageing via
the activation of CD38(+) macrophages. *Nat Metab* **2**, 1265-1283, doi:10.1038/s42255-
020-00305-3 (2020).
107   Wright, L. E. *et al.* Comparison of skeletal effects of ovariectomy versus chemically
induced ovarian failure in mice. *J Bone Miner Res* **23**, 1296-1303,
doi:10.1359/jbmr.080309 (2008).
108   National Toxicology, P. Toxicology and Carcinogenesis Studies of 4-Vinyl-1-cyclohexene
Diepoxide (CAS No. 106-87-6) in F344/N Rats and B6C3F1 Mice (Dermal Studies). *Natl
Toxicol Program Tech Rep Ser* **362**, 1-249 (1989).
109   Ongaro, L. *et al.* Development of a Highly Sensitive ELISA for Measurement of FSH in
Serum, Plasma, and Whole Blood in Mice. *Endocrinology* **162**,
doi:10.1210/endocr/bqab014 (2021).
110   Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat Biotechnol* **37**, 852-857, doi:10.1038/s41587-019-0209-9
(2019).
111   Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina amplicon
data. *Nat Methods* **13**, 581-583, doi:10.1038/nmeth.3869 (2016).
112   Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Res* **43**, e47, doi:10.1093/nar/gkv007 (2015).
113   Leek, J. T. & Storey, J. D. Capturing heterogeneity in gene expression studies by surrogate
variable analysis. *PLoS Genet* **3**, 1724-1735, doi:10.1371/journal.pgen.0030161 (2007).

114 Hornung, R., Boulesteix, A. L. & Causeur, D. Combining location-and-scale batch effect
adjustment with data cleaning by latent factor adjustment. *BMC Bioinformatics* **17**, 27,
doi:10.1186/s12859-015-0870-z (2016).
115 Bisanz, J. <https://github.com/jbisanz/qiime2R> (
116 Rohart, F., Gautier, B., Singh, A. & Le Cao, K. A. mixOmics: An R package for 'omics
feature selection and multiple data integration. *PLoS Comput Biol* **13**, e1005752,
doi:10.1371/journal.pcbi.1005752 (2017).
117 Fernandes, A. D. *et al.* Unifying the analysis of high-throughput sequencing datasets:
characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by
compositional data analysis. *Microbiome* **2**, 15, doi:10.1186/2049-2618-2-15 (2014).
118 Yilmaz, P. *et al.* The SILVA and "All-species Living Tree Project (LTP)" taxonomic
frameworks. *Nucleic Acids Res* **42**, D643-648, doi:10.1093/nar/gkt1209 (2014).
119 Gordon, A. <http://hannonlab.cshl.edu/fastx_toolkit/> (
120 Krueger, F. <https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/> (
121 Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
doi:10.1093/bioinformatics/bts635 (2013).
122 Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930,
doi:10.1093/bioinformatics/btt656 (2014).
123 Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biol* **15**, 550, doi:10.1186/s13059-014-0550-8
(2014).
124 Signorell, A. <https://andrisignorell.github.io/DescTools/> (
125 Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425, doi:10.1016/j.cels.2015.12.004 (2015).
126 Karimnezhad, A. More accurate estimation of cell composition in bulk expression
through robust integration of single-cell information. *Bioinform Adv* **2**, vbac049,
doi:10.1093/bioadv/vbac049 (2022).
127 Pfister, S., Kuettel, V., Ferrero, E. <https://github.com/Novartis/granulator> (
128 Gong, T. & Szustakowski, J. D. DeconRNASeq: a statistical framework for deconvolution
of heterogeneous tissue samples based on mRNA-Seq data. *Bioinformatics* **29**, 1083-
1085, doi:10.1093/bioinformatics/btt090 (2013).
129 Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime
cis-regulatory elements required for macrophage and B cell identities. *Mol Cell* **38**, 576-
589, doi:10.1016/j.molcel.2010.05.004 (2010).
130 Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114-2120, doi:10.1093/bioinformatics/btu170
(2014).
131 Langmead, B., Wilks, C., Antonescu, V. & Charles, R. Scaling read aligners to hundreds of
threads on general-purpose processors. *Bioinformatics* **35**, 421-432,
doi:10.1093/bioinformatics/bty648 (2019).
132 Wood, D. E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2.
Genome Biol **20**, 257, doi:10.1186/s13059-019-1891-0 (2019).

133   Lu, J. *et al.* Metagenome analysis using the Kraken software suite. *Nat Protoc* **17**, 2815-
1498   2839, doi:10.1038/s41596-022-00738-y (2022).
1499   134   mia: Microbiome analysis (2025).
1500   135   Lin, H. & Peddada, S. D. Multigroup analysis of compositions of microbiomes with
1501   covariate adjustments and repeated measures. *Nat Methods* **21**, 83-91,
1502   doi:10.1038/s41592-023-02092-7 (2024).
1503   136   Karl, J. P., Fu, X., Dolnikowski, G. G., Saltzman, E. & Booth, S. L. Quantification of
1504   phylloquinone and menaquinones in feces, serum, and food by high-performance liquid
1505   chromatography-mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci* **963**,
1506   128-133, doi:10.1016/j.jchromb.2014.05.056 (2014).
1507   137   Tingley, D., Yamamoto, T., Hirose, K., Keele, L. & Imai, K. mediation: R Package for Causal
1508   Mediation Analysis. *Journal of Statistical Software* **59**, 1 - 38, doi:10.18637/jss.v059.i05
1509   (2014).
1510   138   Jiang, S. *et al.* Generic Diagramming Platform (GDP): a comprehensive database of high-
1511   quality biomedical graphics. *Nucleic Acids Res* **53**, D1670-D1676,
1512   doi:10.1093/nar/gkae973 (2025).
1513

Figure 1

Figure 2

Figure 3

A

Public ovarian aging bulk RNA-seq datasets

Figure 4

A

FMT-YF

FMT-EF

Ovarian health state
- Ovarian health index ┬ ▪ Ovarian follicle count
                       └ ▪ Serum AMH, FSH
                                 & Inhibin A concentration

Fertility
- Litter size
- Reproductive success rate

B — Ovarian health index — pval ~0.0255

C — Litter size — pval ~0.1726

D — Reproductive success rate — pval ~0.002
FMT-YF (n=18)
FMT-EF (n=18)

Figure 5

Figure 6

A FMT-YF / FMT-EF → Serum untargeted metabolomics (LC-MS/MS)

B PCA - Cohort 1 / PCA - Cohort 2
FMT-YF (n=11)
FMT-EF (n=12)

C LDA1 projection
LDA1 Score
FMT-YF (n=11)   FMT-EF (n=12)

D Top 15 metabolites contributing to LDA separation
DOWN in FMT-EF / UP in FMT-EF

E 3,4-Dihydroxyphenylpropionic acid (0.07923) / Lactamide (0.09084)

Figure 7