SLC13A2-transported citrate is a metabolic signal for PKM2 acetylation and degradation to suppress tumor growth

Mengyao Qin[a,1], Li Shi[a,1], Hao Chen[a,1], Chan Liu[a], Zhiquan Tang[a], Donghao An[a], Wanting Yu[a], Dandan He[b], Chang Shao[b], Shengtao Yuan[c], Hong Yu[d], Haiping Hao[a,b]*, Jing Xiong[a]*

[a]*Department of Pharmacology, School of Pharmacy, China Pharmaceutical University, Nanjing, Jiangsu 210009, China*

[b]*Jiangsu Provincial Key Laboratory of Drug Metabolism and Pharmacokinetics, State Key Laboratory of Natural Medicines, State Key Laboratory of Natural Medicines, China Pharmaceutical University, Nanjing, Jiangsu 210009, China*

[c]*Jiangsu Key Laboratory of Drug Screening, China Pharmaceutical University, Nanjing, Jiangsu 210009, China*

[d] *Department of Pathology, Taizhou People's Hospital, Taizhou, Jiangsu 225300, China*

[1] These authors made equal contributions to this work.

Running title: SLC13A2 inhibits tumor growth via PKM2 acetylation

* Corresponding authors

Jing Xiong, Ph.D.

Department of Pharmacology, School of Pharmacy, China Pharmaceutical University, 24 Tongjiaxiang, Nanjing, Jiangsu 210009, China.

Tel: 86-25-86185328

E-mail address: jxiong@cpu.edu.cn

Haiping Hao, Ph.D.

28 Jiangsu Provincial Key Laboratory of Drug Metabolism and Pharmacokinetics, State Key

29 Laboratory of Natural Medicines, China Pharmaceutical University, 24 Tongjiaxiang, Nanjing,

30 Jiangsu 210009, China.

31 Tel: 86-25-86185566

32 E-mail address: haipinghao@cpu.edu.cn

Abstract

Metabolic reprogramming represents as a pivotal hallmark for cancer, but TCA cycle in tumorigenesis and progression has long been neglected. Solute carrier (SLC) transporters mediate the transport of TCA cycle intermediates across membrane, but their functions in cancer pathogenesis remains unclear. Using integrated analysis of solute carrier (SLC) transporters for TCA cycle intermediates, we found that SLC13A2 was consistently downregulated in hepatocellular carcinoma (HCC) cells and liver tissues from human patients and heterogeneous mouse models. Adeno-associated virus (AAV)-transduced liver-specific knockout or overexpression of SLC13A2 promoted or ameliorated HCC progression in the primary mouse model, demonstrating that SLC13A2 served as a protective factor during HCC pathogenesis. SLC13A2 inhibited HCC cell proliferation by decreasing mitochondrial function via suppressed oxygen consumption and ATP production. Combined with metabolic flux analysis, we found that SLC13A2 imported citrate, which secreted acetyl-CoA as a precursor for the acetylation of pyruvate kinase muscle isozyme M2 (PKM2), which led to its protein degradation. Decreased activity of pyruvate kinase depleted pyruvate for the TCA cycle, thus inhibiting amino acid synthesis and nucleotide metabolism. Additionally, a decrease in nuclear PKM2 protein transduced to reprogrammed gene transcription for cell proliferation and metabolism which is required for tumor growth. This study revealed that citrate transported by SLC13A2 acts as a signal to disrupt metabolic homeostasis for tumor growth and suggests potential drug targets for HCC therapy.

Keywords: TCA cycle, citric acid, pyruvate kinase muscle isozyme M2, glycolysis, mitochondrial function, hepatocellular carcinoma

Introduction

Cancer is characterized by metabolic reprogramming to fulfill its increased demands for uncontrolled cell proliferation. Malignant cells, including hepatocellular carcinoma (HCC), rapidly take up and breakdown nutrients, which contributes to cancer progression (1). Accumulating evidence indicates that elucidating the metabolic dysregulation of tumor cells provides an exciting new direction for targeted HCC therapy (2).

The tricarboxylic acid (TCA) cycle serves as a convergence point for nutrient metabolism and macromolecule synthesis (3). It completely catabolizes many types of fuel sources from the diet, including glucose, amino acids and fatty acids, and then provides precursors for building blocks in macromolecule synthesis, such as nonessential amino acids, purines and pyrimidines (4). In normal cells, glucose produces pyruvate, which enters the TCA cycle and acts as a major source of adenosine triphosphate (ATP) through oxidative phosphorylation. In contrast, cancer cells often truncate glucose metabolism away from the TCA cycle through aerobic glycolysis and therefore are more addicted to glutamine or rely on fatty acids to replenish TCA cycle intermediates (5). Aberrant TCA function due to somatic/germline enzyme mutations or altered activity, such as mutation of isocitrate dehydrogenase 2 (mIDH2) and increased activity of citrate synthase (CS), has been implicated in cancer pathogenesis (4). Moreover, changes in the expression of intermediate transporters have been shown to rewire the metabolism of cancer cells (6). Both oncogenes and tumor suppressors manipulate TCA cycle flux by regulating the expression of intermediate transporters and the activity of enzymes (4). In light of the well-accepted concept that cancer cells mainly obtain energy through aerobic glycolysis, the role of the TCA cycle in cancer metabolism and tumorigenesis has long been neglected and deserves further elucidation.

To ensure the function of the TCA cycle, the concentrations of intermediates are

precisely regulated by the expression of transporters across the plasma membrane

and mitochondrial inner membrane (7). Solute carrier (SLC) transporters, which are

highly expressed in the liver, represent the second largest group of membrane

proteins after G protein-coupled receptors (8). SLCs mediate the transmembrane

transport of various types of nutrients and metabolites, such as glucose, amino acids,

lipids and nucleotides, and thus serve as metabolic switches in cells (9). As most

SLCs transport small molecules, they are considered promising targets for drug

development (10). Some SLC transporters, such as SLC1A5 inhibitor, have been

successfully targeted for the treatment of cancer (11, 12). There are six main SLC

families with members identified as carriers that regulate TCA cycle flux by

transporting TCA cycle intermediates such as citrate, succinate and malate through

the plasma membrane or mitochondrial matrix according to an updated version of the

guide to transporters (13). The role of these SLC transporters in the regulation of

TCA cycle function and tumor growth is largely elusive. Based on previous findings, it

is speculated that these SLC family members control anaplerotic or cateplerotic flux

in response to metabolic adaptation of cancer cells.

Here, we identified SLC13A2 (also known as Na^+-dicarboxylate cotransporter 1,

NaDC1) as a downregulated SLC transporter for TCA cycle intermediates in HCC

liver tissues and cells. By examining a liver-specific SLC13A2

knockout/overexpression mouse HCC model, we further identified SLC13A2 as a

tumor suppressor for HCC progression. Combined with metabolomics and

transcriptomics analysis, SLC13A2, a carrier for citrate, depletes TCA cycle

intermediates, suppresses the biosynthesis of macromolecules, and remodels gene

transcription for tumor growth. Our findings imply that citrate acts as a metabolic

signal transported by SLC13A2, which has the potential to provide an attractive

strategy for treating HCC.

Materials and methods

Animal studies

All animal experiments were approved by the IACUC (Institutional Animal Care and Use Committee) of China Pharmaceutical University (Approval number: 2020-08-006). Male C57BL/6J mice aged 6-8 weeks were purchased from the University of Yangzhou and randomly grouped. Mice were injected with oncogene plasmids through the tail vein for 4-7 s via a technique called hydrodynamic transfection with oncogene delivery and Sleeping Beauty (SB)-mediated somatic integration for stable and long-term gene expression in hepatocytes, as previously described (14). The oncogenic plasmids used were pT3-EF1aH-c-MET (20 μg) and pT3-EF1aH-ΔN90-β-catenin (20 μg), along with SB transposase (5 μg). The oncogene plasmids were diluted with saline to 10% body weight (2 ml/20 g). For liver-specific SLC13A2 knockout experiments, single-guide RNA (sgRNA) sequences were designed based on a CRISPR design tool targeting the first coding exon (Table S1) and constructed together with thyroid hormone-binding globulin (TBG)-Cre into adeno-associated virus 8 (AAV8, 5×10^{11} genome copies/mouse), which was injected into the tail vein of *Rosa26-flox-STOP-flox-Cas9* knockin mice (*Rosa26-LSL-Cas9*, Jackson, #024857) as previously described (15). For liver-specific SLC13A2 overexpression experiments, adult male mice (6-8 weeks of age, ~20 g body weight) were injected with AAV8 (1×10^{11} genome copies/mouse) under the control of the liver-specific TBG promoter via the tail vein (AAV-TBG-GFP versus AAV-TBG-SLC13A2). The plasmids were constructed using the primers shown in Table S1 and used for the AAV8 package. All sufferings of the animal are minimized during the experiments.

Human studies

Human HCC liver samples were obtained during partial liver resection surgery at Jiangsu Taizhou People's Hospital (Taizhou, Jiangsu, China) under a protocol

approved by the Medical Ethics Committee of Taizhou People's Hospital (Approval

number: 2022-03-017). The study was conducted in accordance with the ethical

requirements of the Declaration of Helsinki. Fresh liver tissues from 8 HCC patients

(including cancerous tissues and adjacent paracancerous tissues) were obtained

during surgical resection at the Hospital of Jiangsu Taizhou People (Taizhou,

Jiangsu, China). Both male and female individuals were included. The average age

of the patients was 68.125 years. Written informed consent was provided by all

participants.

Cell culture

All cell lines were characterized by short tandem repeat profiling. The human HCC

cell line HepG2 was obtained from the Institute of Biochemistry and Cell Biology of

the Chinese Academy of Sciences (Shanghai, China). The mouse HCC cell line

Hepa1-6 was obtained from the American Type Culture Collection (CRL-1830) as

described previously (16). Mouse primary hepatocytes were isolated from wild-type

male C57BL/6J mice and cultured as described previously (16).

Quantitative reverse transcriptase–polymerase chain reaction (qRT–PCR)

Total RNA was extracted from cells or liver tissues using TRIzol reagent (Vazyme,

Nanjing, China). cDNA was synthesized using the PrimeScript™ RT Reagent Kit

(Vazyme, Nanjing, China). qRT–PCR was performed with the qPCR SYBR Green

Master Mix (High ROX Premixed) kit (AG Bio, Hunan, China). The primers designed

with NCBI primer BLAST are shown in Table S1.

Immunoblotting and immunoprecipitation

Protein samples were mixed with 3 × SDS loading buffer, separated using SDS–

PAGE, and then transferred onto a polyvinylidene fluoride membrane. The

membranes were incubated with primary antibodies at 4°C for 12-16 h and then with

secondary antibodies for 1 h at room temperature. Enhanced chemiluminescence

reagents were used to visualize the target proteins. For the immunoprecipitation

assay, cell extracts were incubated with primary antibodies or control IgG in a

rotating incubator overnight at 4°C, followed by incubation with protein A agarose

beads (Beyotime, Shanghai, China) for another 4 h. The immunoprecipitates were

washed five times with lysis buffer and analyzed by immunoblotting. The primary and

secondary antibodies used are shown in Table S2.

Histological analysis and immunohistological chemistry staining

Formalin-fixed and paraffin-embedded mouse liver sections were stained with

hematoxylin and eosin (H&E). For immunohistological chemistry staining, the

samples were dewaxed and hydrated. The antigen was retrieved with citric acid

buffer (pH 6.0), blocked with horse serum, and incubated with Ki67 (Abcam,

ab279653) antibody at 4°C for 12-16 h and then with a secondary antibody

conjugated with horseradish peroxidase for 1 h at room temperature. Images were

taken with a BX53 light microscope (Olympus, Japan).

Plasmid construction

186   The plasmids PLVX-GFP-N1 and PLVX-GFP-SLC13A2 were constructed using the

187   primers listed in Table S1. An empty control vector for pReceiver-M02 (EX-NEG-

188   M02) and a mouse PKM2 expression plasmid (EX-Mm04490-M02) were purchased

189   from Guangzhou FulenGen Co., Ltd. (Guangzhou, China).

siRNA and plasmid transfection

For siRNA or plasmid transfection, Hepa1-6 cells were seeded at 40% or 80%

confluence the next day and transfected with siRNAs or plasmids using Lipo8000

(Beyotime, Shanghai, China) according to the manufacturer's instructions (siRNA

sequences are shown in Table S1). HepG2 cells were infected with adenovirus

(pAdeno-MCMV-3XFlag versus pAdeno-MCMV-SLC13A2-3XFlag, OBiO, Shanghai,

China) at an MOI of 25. After 12-18 h of incubation, the medium was changed to

complete medium, and the cells were then cultured for the indicated period before

the experiments.

Assays for cell proliferation *in vitro*

For the knockdown experiment, Hepa1-6 cells were transfected with siRNAs for 24 h,

digested for seeding in 6-well plates at a density of 2.5×10^3 cells/well, and then cultured

for 7 days. For the overexpression experiment, Hepa1-6 and HepG2 cells were

transfected with the overexpression plasmid or adenovirus, respectively, for 24 h,

digested for seeding onto a 6-well plate at a density of 1×10^3 cells/well, and cultured

for 14 days. Cell colonies were fixed with 4% formalin for 15 min and stained with

0.4% crystal violet for 10 min. For the 5-ethynyl-2'-deoxyuridine (EdU) incorporation

assay, Hepa1-6 and HepG2 cells were incubated with EdU for 4 h, fixed for 30 min

and analyzed using a commercial kit (Beyotime, Shanghai, China). Similarly, cell

cycle arrest and cell viability were analyzed using a cell cycle analysis kit (Beyotime,

Shanghai, China) and a CCK8 assay kit (Vazyme, Nanjing, China), respectively,

according to the manufacturer's instructions.

Measurement of the oxygen consumption rate (OCR)

The cells were seeded into the wells of a Seahorse XF 96-well culture plate (Agilent,

Japan) at a density of 8×10^4 cells/well and cultured overnight at 37°C and 5% CO2.

Six wells of cells from each group were used for the analysis. The OCR was

measured with the assurance that the cells were even and approximately 90%

confluent in a monolayer. The sensor cartridge was incubated in a non-CO_2 incubator

at 37°C overnight. Cells in growth media were replaced with fresh and prewarmed XF

media (10 mM glucose and 1 mM sodium pyruvate, 1 mM glutamine) and then

incubated in a non-CO_2 incubator at 37°C for 45 min. Compounds were diluted with

fresh XF media and injected for measurement as follows (oligomycin 1.5 µM, FCCP

1 µM and rotenone/antimycin 0.5 µM). The data were analyzed directly by Seahorse

software. The basal and maximal OCRs were obtained for statistical analysis.

Mitochondrial membrane potential assay

Hepa1-6 cells were seeded at a density of 5×10^4 cells/well in a 24-well plate. After

transfection with the plasmid for 48 h, the cells were stained with JC-1 according to

the manufacturer's instructions (Beyotime, Shanghai, China). Images were taken

using an inverted fluorescence microscope (IX71, Olympus, Japan) with the imaging

software Olympus cellSens Standard software.

Isolation of subcellular fractions

The procedures were conducted on ice with prechilled buffers and equipment.

Hepa1-6 cells in 10 cm dishes were transfected with adenovirus and harvested after

72 h. After being washed with PBS three times, the cell pellets were suspended in

homogenization buffer (225 mM D-mannitol, 75 mM sucrose, 0.1 mM EDTA, 30 mM

Tris-HCl, pH 7.5) and then transferred to a 2-mL Dounce homogenizer to disrupt the

cells with several strokes of the pestle. For each stroke, the pestle was pressed

straight down the tube to maintain a firm and steady pressure. When checking the

degree of homogenization with a phase-contrast microscope, eight to nine naked

nuclei for every whole cell indicated a good result. The homogenate was transferred

to a centrifuge tube and spun down at 500 × g for 5 min. While the supernatant was

used for preparing the cytoplasm at 7000 × g for 10 min, the pellet was gently

suspended in homogenization buffer followed by centrifugation at 700 × g for 10 min

to collect the nuclei. The extracted subcellular fractions were confirmed by

immunoblotting for cytosolic and nuclear β-tubulin and PCNA, respectively.

ATP measurement

Transfected cells were seeded into 96-well culture plates at a density of 2×10^3

cells/well. After 72 h, ATP production was detected with a CellCounting-Lite® 2.0

Luminescent Cell Viability Assay kit (Vazyme Biotech, Nanjing, China) by a plate

reader (Beckman Coulter, Krefeld, Germany) according to the manufacturer's

instructions.

Metabolomic analysis

Hepa1-6 cells were collected, and metabolites were extracted after adenovirus

transfection for 72 h. Cell metabolites were dissolved in 0.5 mL of ice-cold 80%

methanol solution containing 1 μg/mL 4-chloro-DL-phenylalanine (Sigma, C6506).

The samples were then incubated for 30 min in a -80°C freezer, followed by

centrifugation at 13,000 × g for 15 min at 4°C, after which the supernatant was dried

using a vacuum centrifugal concentrator (Thermo, Massachusetts, USA). Before

analysis, the samples were reconstituted with methanol, followed by centrifugation at

18,000 × g for 10 min at 4°C twice, and untargeted metabolomics was conducted

using a 6546 LC/Q-TOF (Agilent, USA) instrument equipped with an electrospray

ionization (ESI) source. Separation was achieved on an XBridge Premier BEH Amide

VanGuard FIT column (2.5 μm, 4.6 mm × 150 mm, Waters, USA). MS data were

acquired in negative ESI mode over a range of m/z 50 to 850. The mobile phase

consisting of water (containing 0.3% ammonium acetate and 0.3% ammonia in phase

A, v/v) and acetonitrile (phase B) was delivered at a flow rate of 0.4 ml min^{-1}. The

total elution time was 36 min for the gradient program. The data were analyzed by

Progenesis QI with conditions of P<0.05, fold change>2, and VIP>1 to screen for

differentially abundant metabolites. Metabolic pathway analysis was performed

based on the MetaboAnalyst 5.0 platform.

For the quantitative determination of metabolites, cell samples and mixed standards

of gradient concentrations were measured on a TripleTOF 5600 system (Sciex, USA)

by direct infusion. The instrument was set to acquire data over a m/z range of 100-

2,000 for the TOF-MS/MS scan. Aliquots of reconstituted supernatants (5 µL) were

analyzed with an XBridge BEH Amide column (3.5 µm, 4.6 mm × 100 mm, Waters,

USA) on an LC-30 HPLC system (Shimadzu). The mobile phase consisting of

water/acetonitrile (95:5, v/v) (containing 0.1% ammonium acetate and 0.3% ammonia

in phase A, v/v) and acetonitrile (phase B) was delivered at a flow rate of

0.4 ml min^{-1}. The total elution time was 36 min for the gradient program. The

concentrations of the tested substances were calculated by comparing the parent ion

and daughter ion as well as the retention time with the standard curve formed by the

gradient standards.

282 Stable isotope resolved metabolomics (SIRM) analysis

The culture medium was replaced by U-13C^{6}-glucose (Cambridge) in glucose-free

DMEM (Sigma) after the transfection of Hepa1-6 cells for 60 h. The cells were

labeled at the indicated time points. An unlabeled culture was prepared in parallel by

adding equal concentrations of unlabeled glucose to the media to identify unlabeled

metabolites. Cell metabolites were extracted with 80% methanol and completely

concentrated to dryness using a Speedvac (Thermo Fisher Scientific, USA). The

samples were redissolved in pure methanol and used for LC–MS/MS analysis.

Standard metabolites were observed, and the MS/MS spectra of these compounds

were manually confirmed according to Metlin (www.metlin.scripps.edu) or HMDB

(www.HMDB.ca).

293 RNA-seq and bioinformatic analysis

Liver RNA sequencing was performed on DNBSEQ-T7 cells from GFP- or SLC13A2-

OE mice in the HTVi model. The fastq files were aligned to the mouse reference

genome (mm10) using the aligner STAR. RNA-seq fastq data have been deposited

into the Sequence Read Archive (PRJNA1083485). The published RNA-seq raw data

for heterogeneous liver cancer induced by diverse cancer driver genes were

downloaded from the NCBI Sequence Read Archive with accession code

PRJNA674008 (17). The published RNA-Seq raw data of human HCC cancerous

and paracancerous specimens were downloaded from the Gene Expression

Omnibus database (GSE193567) (18). TCGA HCC RNA-seq data were downloaded

from the NIH-GDC repository (https://portal.gdc.cancer.gov/), which included 371

primary HCC patients in level 3. The HCC samples were aligned according to the

expression level of PKM2, where top quarter considered as PKM2-high group was

compared with bottom quarter as PKM2-low group to screen out PKM2-regulated

gene cluster in clinical datasets. Differential gene expression analysis was performed

using the Deseq2 package in the R program. The ENSEMBL numbers in the analysis

results were converted to gene names using probes in the human gene annotation

package, and genes with a log2FoldChange greater than 1 or -1 and a P value less

than 0.05 were selected for plotting using the ggplot2 package. Heatmaps and

volcano plots were generated by R. Gene enrichment analysis was performed using

MetaScape (www.metascape.org). The overall differentially expressed genes and the

upregulated and downregulated genes in SLC13A2-OE mice were independently

subjected to motif enrichment analysis (HOMER, findMotif.pl).

Enzymatic activity assay

Approximately 100 mg of liver tissue was ground with a 1:9 weight/volume ratio of

PBS and centrifuged at 2500 × g at 4°C for 10 min. Hepa1-6 cells were transfected in

6-well plates with plasmids for 72 h. The collected cells were mixed with 500 μL of

PBS and homogenized manually. The enzymatic activity of the homogenate was

detected by alanine transaminase (ALT) and aspartate transaminase (AST) kits

(Jiancheng, Nanjing, China). Protein concentrations were quantified using the

Bradford protein assay and used for the normalization of enzymatic activity.

Cycloheximide (CHX) chase assay

To assess the half-life of the proteins, Hepa1-6 cells overexpressing SLC13A2 were

treated with 30 µg/mL CHX, a protein synthesis inhibitor, and harvested at 0 h, 3 h, 6

h or 9 h for immunoblot analysis. The densitometry of the bands was quantified using

ImageJ.

Statistical analysis

The data are expressed as the mean $\pm$ SEM. The differences between two groups

were statistically analyzed using a two-tailed Student's *t* test. Differences from the

growth curve study were analyzed by two-way ANOVA, followed by Bonferroni

multiple comparisons. P<0.05 was considered to indicate statistical significance.

Results

SLC13A2 is a downregulated SLC transporter for TCA cycle intermediates in

HCC

To explore the formerly overlooked role of the TCA cycle, we focused on transporters

of TCA cycle intermediates in HCC. To date, SLC families, including SLC5, SLC13,

SLC16, SLC25, SLC33 and SLC54, have been identified to transport TCA

intermediates such as citrate, succinate, a-KG, malate, L-lactate and pyruvate,

according to an updated version of the guide to transporters (13) (Figure 1A). We

then analyzed RNA-seq data from heterogeneous liver tumors induced by diverse

cancer driver genes selected by mutational frequency from human HCC cohorts with

different etiological factors and ethnic origins (17). SLC13A2 was potently and

consistently downregulated in all liver tumors with obviously heterogeneous

phenotypes, showing an extraordinary pattern compared with that of other detectable

SLC transporters (Figure 1B).

Based on the above data, we generated a primary HCC mouse model (HTVi model)

to evaluate the changes in the above SLC family members via hydrodynamic

transfection of oncogene plasmids in combination with SB transposase (Figure 1C).

Compared with those in the saline group, obvious tumor nodules were observed in

the livers of the mice that were transfected with the oncogene plasmids, as shown by

the presence of densely packed and multinucleated hepatocytes, as well as an

increased nucleoplasmic ratio according to histological staining (Figure 1D). The

liver/body weight ratio significantly increased in the oncogene-transduced group

(Figure 1E, left). The mRNA expression of genes involved in glucose metabolism and

fatty acid metabolism enzymes in the oncogene plasmid group was consistent with

the metabolic adaptation of tumor cells (Figure 1E, right). These findings suggested

that the primary HCC mouse model was generated successfully.

Next, we detected the expression of SLC transporters of interest in liver tissues and

found that the expression of *Slc16a3* and *Slc16a8* significantly increased, while that

of *Slc13a2, Slc25a1, Slc25a10, Slc25a11, Slc25a21, Slc54a1,* and *Slc54a2*

significantly decreased in liver tissues from mice with HCC (Figure 1F top left).

Among these genes, *Slc13a2* decreased 10-100 times and attracted our attention

because it exhibited the most abundant variation. Moreover, the expression of SLC

transporters was evaluated in mouse HCC Hepa1-6 cells and mouse primary

hepatocytes, and the results showed that the expression of *Slc25a21* and *Slc54a3*

significantly increased, while that of other SLC transporters of interest decreased,

except for *Slc16a1* (Figure 1E bottom left). Similarly, we detected SLC transporter

expression in mouse HCC Hepa1-6 cells compared with that in immortalized mouse

hepatocyte AML12 cells. The gene expression of *Slc25a1, Slc25a10 and Slc54a1*

increased, while that of *Slc13a2*, *Slc13a3*, *Slc13a5*, and *Slc54a3* decreased in the

HCC cell line (Figure 1E, upper right). Among the above SLC transporters, *Slc5a12*

was undetectable in both HCC liver tissues and cell lines. Using integrated analysis

of SLC expression in primary HCC liver tissues and cell lines, we found that *Slc13a2*

was the only gene consistently deregulated both *in vivo* and *in vitro* (Figure 1F

bottom right). In addition, reduced protein expression of SLC13A2 was further

confirmed in HCC liver tissues, while the levels of proteins and cell signaling

pathways that promote cell proliferation were significantly increased (Figure 2A).

These results indicate that SLC13A2 is downregulated in HCC and stands out from

the other SLC transporters for TCA cycle intermediates.

By analyzing a published Gene Expression Omnibus dataset (GES193567), we

identified 4247 differentially expressed genes (DEGs) between cancerous and

paracancerous tissues of human HCC patients. Gene enrichment analysis revealed

that the DEGs were enriched in macromolecular biosynthesis, membrane trafficking

and transmembrane transporters (Figure 2B, left). In this dataset, SLC13A2 was

identified as one of the three significantly altered SLC transporters for TCA cycle

intermediates between paratumor and tumor tissues (Figure 2B, right). As shown for

all DEGs, SLC13A2 was significantly reduced in tumor tissues (Figure 2C).

Moreover, Kaplan–Meier survival curves revealed that male patients with high

SLC13A2 expression had a better prognosis and a greater overall survival rate

(Figure 2D). Additionally, SLC13A2 mRNA was significantly lower in human HCC cell

lines than in the normal human liver cell line LO2 (Figure 2E). Furthermore, we

collected liver tissues from clinical patients with HCC and analyzed the expression of

SLC13A2. We confirmed that all HCC patients had lower SLC13A2 expression in

cancerous tissues than in paracancerous tissues (Figure 2F). The above data

suggest that reduced expression of SLC13A2 is conserved in mouse and human

HCC and is positively associated with the prognosis of this disease.

SLC13A2 inhibits HCC progression in a mouse model induced by

hydrodynamic transfection of oncogenes

The function of SLC13A2, a previously described transporter in renal tubules and

intestinal cells, in physiological homeostasis in the liver is still obscure. Initially, we

studied the role of SLC13A2 in the liver via hepatocyte-specific SLC13A2 knockout or

overexpression via AAV-mediated transduction. Recombinant AAV8 with a vector

expressing sgRNA constructed together with TBG-Cre was used to transduce Cas9$^{fl/fl}$

knock-in mice for specific ablation of SLC13A2 in the liver. When SLC13A2 was

specifically knocked out in hepatocytes for 3 months, no consistent differences in

blood glucose, body weight, liver weight, histological staining or the expression of

genes involved in glycolysis, the TCA cycle or lipogenesis were detected in the two

liver-specific SLC13A2-KO (LKO) groups (Figure S2A-E). Furthermore, we detected

the activation of signaling pathways regulating cell proliferation and found that p-

mTOR was slightly decreased, while p-GSK3β was significantly increased (Figure

S2F). Similarly, when we overexpressed SLC13A2 in the liver, no differences in body

weight, liver weight, blood glucose, histological morphology, gene expression or

signaling pathway activation were detected (Figure S3A-F). These data imply that

hepatocyte-specific knockout or overexpression of SLC13A2 primes the cell

metabolic status for HCC tumorigenesis but is not sufficient to directly induce the

initiation of HCC.

To explore the effect of SLC13A2 on the tumorigenesis and progression of HCC, we

first used CRISPR/Cas9-mediated LKO mice. The mice were injected with oncogene

plasmids using the HTVi model after 2 weeks of AAV8 injection and sacrificed after 8

more weeks (Figure 3A). As expected, the knockout efficiency of SLC13A2 was

verified at the mRNA and protein levels (Figure 3B). Although there were no

significant differences in body weight or blood glucose between the two groups, the

liver weight/body weight ratio significantly increased in the LKO group (Figure 3D).

The number of surface tumor nodules was significantly greater in LKO mice than in

control mice (Figure 3C). While the percentage of tumor incidence was similar

between the two groups, their tumor sizes were significantly different. The SLC13A2

LKO group had remarkably more mice with larger maximal tumor sizes than did the

control group (Figure 3E). As indicators of liver tumor burden (19), the enzymatic

activity of serum and liver ALT/AST also increased with SLC13A2 ablation (Figure

3F). Histological staining revealed larger nodules with more irregularly shaped nuclei

in the liver H&E-stained slides from the SLC13A2 LKO group (Figure 3G).

Immunohistochemical staining of Ki67 also revealed 16.6% nuclear positivity in the

control group and approximately 22.9% nuclear positivity in the SLC13A2 LKO group,

demonstrating a significant increase in cell proliferation induced by the knockout of

SLC13A2 *in vivo* (Figure 3H)*.* Consistently, the phosphorylation of mTOR, AKT and

GSK3β, which are involved in cellular signaling related to cell proliferation and

metabolism, was significantly elevated, consistent with the promotion of tumor growth

in liver-specific knockout mice (Figure 3I).

Likewise, a primary mouse model was established with liver-specific overexpression

of SLC13A2 using AAV8 under the control of the TBG promoter. After 2 weeks of

AAV injection, the mice were injected with oncogene plasmids via the HTVi method

and sacrificed after 9 more weeks, which is a longer period than that of the LKO

experiments, to better demonstrate tumor suppression by SLC13A2 (Figure 4A). The

overexpression efficiency of SLC13A2 was verified at the mRNA and protein levels

(Figure 4B). The number of surface tumor nodules and liver/body weight ratio were

significantly lower in the SLC13A2-overexpressing group than in the control group,

while no significant differences were observed in body weight or blood glucose

between the two groups (Figure 4C-D). The tumor burden severity was significantly

different, as was the percentage of tumor incidence. The maximal tumor size in the

SLC13A2 overexpression group was significantly reduced compared to that in the

GFP group (Figure 4E). The enzymatic activity of ALT and AST was consistently

inhibited in the blood and liver tissues harvested from mice with liver-specific

overexpression of SLC13A2 (Figure 4F). Histological staining revealed that

hepatocytes were smaller and more separated in the livers of the SLC13A2-

overexpressing group than in those of the control group, as indicated by a decreased

nucleo-plasmic ratio (Figure 4G). Consistent with the decrease in tumor growth,

immunohistochemical staining of Ki67 also revealed 8.0% nuclear positivity in the

control group and approximately 2.8% nuclear positivity in the SLC13A2

overexpression group, demonstrating a significant decrease in cell proliferation

induced by SLC13A2 *in vivo* (Figure 4H). Similarly, the RNA expression of cell

proliferation-associated genes, such as *Ccnb1*, *Foxm1, Fgf21* and *Mcm2,* decreased

in liver tissues overexpressing SLC13A2 *in vivo* (Figure 4I). Moreover, the expression

of genes involved in metabolism, such as *Hk2* and *Mdh2,* also significantly

decreased with the overexpression of SLC13A2 (Figure 4I). These findings indicated

a favorable metabolic status for HCC progression induced by SLC13A2.

SLC13A2 inhibits HCC cell proliferation by depleting TCA cycle intermediates

for macromolecular biosynthesis and suppressing oxidative respiration

Next, a mouse HCC cell line, Hepa1-6, and a human HCC cell line, HepG2, were

chosen to confirm the capacity of SLC13A2 to prevent the proliferation of HCC cells.

To further investigate the effect of SLC13A2 on the proliferation of HCC cells *in vitro*,

we used RNA interference (KD) to knock down SLC13A2 and plasmid/adenovirus

transfection to induce SLC13A2 expression (OE) in HCC cells. Notably, RNA

interference transduction significantly reduced the protein expression of SLC13A2 in

Hepa1-6 cells (Figure 5A). Strikingly, growth curve and colony formation experiments

revealed that the colony formation ability and viability of Hepa1-6 cells were

significantly increased with SLC13A2 knockdown (Figure 5B-C). Transfection of

Hepa1-6 or HepG2 cells with plasmids or adenoviruses increased the protein

expression of SLC13A2 (Figure 5A). Growth curve and colony formation assays

revealed that SLC13A2 overexpression significantly decreased the viability and

clonogenicity of Hepa1-6 and HepG2 cells (Figure 5B-C). To further confirm the

effect of SLC13A2 on cell proliferation, an EdU incorporation assay was applied. The

ratio of EdU-positive cells was significantly greater in SLC13A2-KD cells than in

SLC13A2-OE cells (Figure 5D). Consistently, a cell cycle distribution assay showed

that SLC13A2 knockdown decreased the proportion of cells in S phase, while

SLC13A2 overexpression increased S phase arrest (Figure 5E).

Since classical substrates of SLC13A2 are TCA cycle intermediates that control

cellular function and cell fate (3), we conducted UPLC-QTOF-MS-based untargeted

metabolomics to investigate the effect of SLC13A2 overexpression on metabolism in

HCC Hepa1-6 cells. The levels of intermediates of the TCA cycle, such as citrate, a-

KG, malate and aconitate; nonessential amino acids, such as glutamine and

aspartate; and metabolites involved in nucleotide metabolism, such as cytidine and

orotidine, decreased after SLC13A2 overexpression (Figure 6A). The differentially

abundant metabolites were largely enriched in alanine-aspartate-glutamate

metabolism, D-glutamine and D-glutamate metabolism and the aminoacyl-tRNA

biosynthesis pathway (Figure 6B). Then, the effects of SLC13A2 on oxidative

respiration were detected using Seahorse analysis. SLC13A2 overexpression

impaired basal and maximal respiratory capacity (Figure 6C). Consistently, ATP

production decreased with SLC13A2 overexpression and increased with SLC13A2

knockdown (Figure 6D). To detect the mitochondrial transmembrane potential, the

cells were stained with the sensitive fluorescent probe JC-1. Mitochondrial

depolarization was induced by SLC13A2, as the ratio of aggregates to monomers

significantly decreased with SLC13A2 overexpression and increased with SLC13A2

knockdown (Figure 6E). This finding suggested that SLC13A2 decreases cellular

mitochondrial capability by inhibiting oxidative phosphorylation and ATP synthesis

and increasing mitochondrial permeability.

Citrate-imported SLC13A2 acetylates and thus degrades PKM2, resulting in

TCA cycle restriction to suppress HCC cell proliferation

The mechanisms by which SLC13A2, a well-known citrate inward transporter,

restricts the TCA cycle to suppress HCC tumor growth are intriguing. Therefore, we

used metabolic flux analysis to better understand the effects of SLC13A2 on

glycolytic flux via the application of U-$^{13}C_6$-glucose at the indicated time points.

Surprisingly, after SLC13A2 was overexpressed, the levels of the upstream

glycolysis intermediates G-6-P, GAP/DHAP, 3-PG, and PEP significantly increased,

while the levels of the downstream metabolite pyruvate significantly decreased with

SLC13A2 overexpression (Figure 7A). Furthermore, targeted metabolomic analysis

confirmed that the intracellular pyruvate content decreased significantly with

SLC13A2 overexpression, which was consistent with the restoration of cell

proliferation by an additional supply of pyruvate (500 μM) in Hapa1-6 cells. (Figure

7B). These findings suggested that SLC13A2 controls the entrance of the TCA cycle,

which was consistent with its ability to suppress tumor growth and attenuate cell

proliferation.

Since citrate is one of the major substrates of SLC13A2 with the highest affinity (20),

we hypothesized that the citrate imported by SLC13A2 was catalyzed by ACLY to

produce oxaloacetate and acetyl-CoA. We used targeted metabolomic analysis to

detect intracellular oxaloacetate and found that it was significantly increased by

SLC13A2 overexpression, while extracellular citrate was significantly decreased

(Figure 7C, left). These results indicated that SLC13A2 imported extracellular citrate,

which produced oxaloacetate. Cell viability assays also revealed that treatment with

an ACLY inhibitor (50 μM) significantly reversed the SLC13A2-mediated inhibition of

cell growth (Figure 7C, right). As a major enzyme that catalyzes the transition of PEP

to pyruvate in HCC cells (21), PKM2 was therefore suggested to be regulated by

SLC13A2. We thus hypothesized that acetyl-CoA released by citrate catalysis may

act as a precursor for the acetylation of PKM2, which leads to its protein degradation

(22). To evaluate the protein distribution of PKM2 after SLC13A2 overexpression, we

prepared isolated subcellular fractions from the nucleus and cytosol and found that

SLC13A2 significantly decreased PKM2 protein levels in the cytosol and nucleus

(Figure 7D below). While the PKM2 protein was decreased by SLC13A2, the RNA

level was unchanged (Figure 7D up), which also indicated that posttranslational

modification resulted in a decrease in the PKM2 protein. Therefore, we performed

coimmunoprecipitation (co-IP) experiments in SLC13A2-overexpressing Hepa1-6

cells. The acetylation of PKM2 increased significantly with SLC13A2 overexpression

(Figure 7E). We further used a CHX-chase assay to examine whether SLC13A2

accelerated the turnover of the PKM2 protein. The data revealed that protein

degradation was the primary pathway involved in the SLC13A2-mediated decrease in

the PKM2 protein level (Figure 7F). Consistently, pyruvate kinase activity was

strongly attenuated by SLC13A2 overexpression (Figure 7G). Since K305 has been

reported to be acetylated for PKM2 protein degradation(22), we generated a PKM2-

K305Q acetylation-mimic mutant and transfected it into HCC cells with SLC13A2

knockdown. The results confirmed that the level of the PKM2 305Q mutant protein

significantly decreased compared with that of the WT protein. The knockdown of

SLC13A2 increased the levels of both the endogenous protein and the transfected

WT PKM2 protein but did not increase the levels of the K305Q mutant protein (Figure

7H). Furthermore, PKM2 overexpression reversed the inhibitory effect of SLC13A2

on HCC cell proliferation, as confirmed by growth curve and colony formation

analyses (Figure 7I). SLC13A2 suppresses HCC tumor growth in a PKM2-dependent

manner.

SLC13A2 remodels gene transcription for proliferation and metabolism, which

restrains tumor growth

To delineate the mechanisms of SLC13A2 in tumor suppression, we evaluated the

transcriptional regulation induced by SLC13A2 using transcriptomic RNA-seq

analysis of total liver RNA from the GFP and SLC13A2-OE groups in the HTVi model

(Figure 4A). We identified a total of 2205 hepatic DEGs in SLC13A2-OE mice

(FDR<0.05, fold change>2, adj P>0.05) (Figure 8A). Gene Ontology analysis

revealed that the downregulated genes were enriched in cell proliferation and nuclear

division, and the upregulated genes were enriched mainly in small molecule catabolic

pathways and nutrient metabolism pathways (Figure 8B). Gene set enrichment

analysis also revealed that 102 gene sets were significantly differentially regulated by

SLC13A2. The top two gene sets of organic acid catabolic process and amino acid

catabolic process were upregulated in the SLC13A2-OE group (Figure 8C).

It has been reported that PKM2 not only acts as a PK enzyme but is also is located in

the nucleus to promote gene transcription as a transcription factor or cotransporter by

physically interacting with other transcription factors, including HIF-1α, the NF-κB

subunit p65/RELA, STAT3 and ATF2 (23-25).   In addition, nuclear PKM2 also

induces the expression or regulates the protein stability of transcription factors such

as EGR1 and RUNX1 (26). By analyzing transcription factor binding motif

enrichment, relevant transcription factors, including RELA, EGR1, HNF4a, STAT3

and RUNX1, which are highly associated with PKM2 nuclear functions, were

predicted to be involved in the regulation of gene transcription by SLC13A2 during

HCC tumorigenesis (Figure 8D).

Furthermore, we analyzed the RNA-seq data of 371 primary HCC patients in level 3

from the TCGA clinical dataset and evaluated the differential genes in the PKM2-high

group compared with PKM2-low expression group (27). These differential genes

reflected the regulatory effects of PKM2 on gene expression. We identified 3975

differentially expressed genes that were conserved between human and mouse, with

a coverage of approximately 50% SLC13A2-regulated genes (Figure 8E top). Gene

ontology enrichment showed similar gene expression patterns between the

regulation of PKM2 and the induction by the overexpression of SLC13A2 (Figure 8E

bottom and 8F). By performing transcription factor motif enrichment analysis, we

observed that transcription factor-binding sites predicted for the genes whose

transcription was differentially regulated by SLC13A2 were also highly enriched for

the genes whose expression was differentially regulated by PKM2 in human HCC

patient datasets. The shared motifs enriched for the expression of both SLC13A2-

and PKM2-regulated genes corresponded to predicted binding sites for transcription

factors, including RELA, EGR1, HNF4a, RXR and PPARa (Figure 8G). The PKM2-

regulated genes covered about 90% of the predicted transcription factors by

SLC13A2, indicating the dependence of SLC13A2-regulated gene expression on

nuclear PKM2. The data establish a novel model of PKM2 protein degradation

through the SLC13A2-mediated inward transport of citrate, which acts as a signaling

metabolite that suppresses HCC progression (Graphic abstract).

Discussion

Since the 'Warburg effect' was proposed by Otto Warburg during the 1920s, an

increasing number of scholars have focused on the role of aerobic glycolysis in

tumors, but the role of the TCA cycle, not to mention the TCA cycle substrates, is still

largely obscure (28). In the present study, we found that SLC13A2, an SLC

transporter that imports citrate, is downregulated in human and mouse HCC cells.

Citrate produces acetyl-CoA for the acetylation and degradation of PKM2, thus

depleting TCA cycle intermediates and decreasing macromolecule biosynthesis and

gene transcription for HCC tumor growth (schematic diagram shown in the graphic

abstract). As deregulated metabolism is considered a key hallmark of HCC

progression, our findings suggest a novel transporter with potential for the design of

drugs or strategies for HCC treatment.

Citrate is a substance involved in the TCA cycle and de novo fatty acid/cholesterol

synthesis and plays a key role in cell survival. Deciphering the consequences of

citrate metabolism is important for dissecting the etiology of HCC. The classical

plasma membrane citrate transporters SLC13A2, SLC13A3, SLC13A5 and

mitochondrial SLC25A1 have nonnegligible effects on the metabolism of citrate.

SLC13A family transporters, which can be divided into two functionally distinct

groups, mediate the movement of Na+-coupled anion substrates across the

membrane (29). The group of Na^+-di- and tri-carboxylate cotransporters carrying TCA

cycle intermediates includes SLC13A2, SLC13A3 and SLC13A5, among which the

first is exclusively downregulated in HCC liver tissues and cells (Figure 1B, 1F and

2A-C, 2F). As a previously defined transporter expressed in the apical membrane of

renal proximal tubular cells and small intestine cells, SLC13A2 is considered a

primary protein involved in citrate reabsorption and is thus important for acid–base

homeostasis (30). In renal tubules, SLC13A2 integrates the movement of three

sodium ions with dicarboxylate and is triggered by a low intracellular sodium

concentration as well as intracellular electronegativity. SLC13A2 has high specificity

for dicarboxylates with different affinities for substrates (0.35 mM km for succinate

and 0.6 mM km for citrate) (31). Although SLC13A2 has a greater affinity for

succinate than citrate, as previously described, the latter is a much more abundant

metabolite in HCC cells than the former (the content ratio is more than 2800-fold)

(Figure S1C). There is also less intracellular a-KG than citrate (approximately 14-

fold). To date, few studies have focused on the role of SLC13A2 in addition to the

kidney, especially the liver. In contrast to the general belief that SLC13A2 is

expressed at very low levels in the liver, we found that SLC13A2 mRNA is normally

expressed in healthy liver tissue (Table 1), indicating its role in the physiological

homeostasis of liver function. Although we did not observe any obvious phenotype

with hepatocyte-specific overexpression or knockout of SLC13A2 (Figure S2, S3)

under physiological conditions, it possibly contributes to tuning the metabolic status

of hepatocytes for the maintenance of liver homeostasis. We speculate that a robust

phenotype might be shown with metabolic stresses such as HFD feeding. The role of

SLC13A2 in the regulation of liver metabolism still deserves further elucidation.

As another member of the SLC13 family, SLC13A5 has gained increasing attention

because of its high expression in the liver. SLC13A5 reportedly regulates hepatic

glucose and fatty acid metabolism by transporting citrate from the extracellular to the

intracellular compartment. Knockout of SLC13A5 prevents high-fat diet-induced

insulin resistance by attenuating hepatic gluconeogenesis and lipogenesis (32).

RNAi-mediated silencing of SLC13A5 expression in two human HCC cell lines,

HepG2 and Huh7, profoundly suppressed cell proliferation and colony formation and

induced cell cycle arrest through increased expression of the cyclin-dependent

kinase inhibitor p21 and decreased expression of cyclin B1 (33). However, the mRNA

expression of both *Slc13a3* and *Slc13a5,* two other members with similar functions in

the family, did not significantly or consistently change in HCC liver tissues (Figure 1B,

1F and 2B). The extracellular import of citrate induced by SLC13A2 represents a new

mechanism for regulating HCC metabolism that is distinct from that of SLC13A5.

SLC25 family members are considered important transporters of the inner

mitochondrial membrane and are involved in numerous metabolic pathways (34).

SLC25A1 (also known as mitochondrial citrate transport protein) is a mitochondrial

carrier that transports citrate or isocitrate into the cytosol in exchange for malate.

Increased transcription and increased activity of SLC25A1 have been revealed as

hallmarks of several metabolic disorders and several cancer types (35). It has been

reported that SLC25A1 is highly expressed in the metastatic lymph nodes of non-

small cell lung cancer patients. Lung cancer stem cells rely on SLC25A1 to maintain

the mitochondrial citrate pool and redox balance, resulting in the stimulation of

oxidative phosphorylation for energy production and cell survival (36). Studies have

shown that SLC25A1 promotes the entry of cytosolic citrate into mitochondria

through reverse import activity, revealing the obscure role of SLC25A1 in

tumorigenesis (36). Anchorage-independent tumor growth requires changes in the

metabolism of citrate, which is imported into mitochondria by SLC25A1, to inhibit

mitochondrial oxidative stress (37). In addition, SLC25A1 plays an important role in

lipid metabolism by exporting citrate from mitochondria, and pharmacological

inhibition of SLC25A1 prevents NASH pathogenesis (38). As another key

dicarboxylate carrier, SLC25A10 transports malate and succinate out of mitochondria

in exchange for phosphate, sulfate and thiosulfate. The expression of SLC25A10 has

been linked to cancer, but further understanding of its role in mitochondrial function

and TCA cycle intermediates is lacking (39, 40). A recent study revealed that

SLC25A10 is predominantly expressed in white adipose tissue and is responsible for

succinate transport from mitochondria to the cytosol. The exported succinate

therefore acts on its receptor SUCNR1 to inhibit adipocyte lipolysis and protect

against liver lipotoxicity upon HFD feeding (41). Both SLC25A1 and SLC25A10

transport substrates by electroneutral exchange (42), which is distinct from the

scenario induced by SLC13A2. Downregulation of SLC13A2 in HCC liver tissues and

cells maintains the mitochondrial membrane potential across the inner membrane

(Figure 6E), which ensures the functional operation of the TCA cycle for oxidative

phosphorylation and ATP production in HCC cells (Figure 6C-D).

Apart from playing a central role in mitochondrial metabolism and respiration, citrate

also plays fundamental roles in the cytosol by serving as a metabolic substrate, an

allosteric enzymatic modulator and a source of acetyl-CoA as an epigenetic modifier

and precursor for lipogenesis (43). The activities of enzymes involved in glycolysis,

lipogenesis and gluconeogenesis are regulated by binding to citrate, for example, the

activation of 1,6-bisphosphatase (FBP1) and acetyl-CoA carboxylase alpha (ACACA)

and the inhibition of phosphofructokinase (PFK) (44, 45). Our results reveal a novel

mechanism for subtly tuning the intracellular concentration of citrate for HCC cell

growth. SLC13A2 imports citrate, leading to the inhibition of glycolysis through the

acetylation and degradation of PKM2 (Figure 8D-F). This finding echoes previous

findings that the intracellular concentration of citrate remains low in cancer cells to

sustain tumor growth (46). Intriguingly, SLC13A2 also gates the entrance to the TCA

cycle and decreases its intermediates, including citrate (Figure 6A left), for cell

growth. This indicates that the depletion of TCA cycle intermediates is the eventual

consequence of SLC13A2, which ultimately suppresses cell growth. The present

findings provide further evidence that citrate serves as a metabolic signal to regulate

cell growth in addition to serving as a metabolite in the TCA cycle to produce energy.

Aerobic glycolysis is found in HCC and is a hallmark of liver cancer and is

responsible for the regulation of HCC proliferation, immune evasion, invasion,

metastasis, angiogenesis and drug resistance. Pyruvate kinase (PK) is an enzyme

that catalyzes the conversion of phosphoenolpyruvate and ADP to pyruvate and ATP

in glycolysis and plays a role in regulating cellular metabolism. Unlike other PKs,

PKM2 is highly upregulated in cancer cells and is associated with a poor prognosis

(47). The activity of PKM2 can also be regulated by posttranslational modifications,

including acetylation, oxidation, phosphorylation, prolyl hydroxylation, and

SUMOylation. These modifications have been shown to modulate PKM2 activity,

degradation, oligomerization, and subcellular localization. PKM2 is directly

phosphorylated on tyrosine residues, including Y105, allowing phosphorylated PKM2

to play a role in inactivating other PKM2 tetramers by stimulating FBP release (48).

Acetylation of PKM2 K305 reduces pyruvate kinase activity by reducing its affinity for

PEP and triggering degradation of the enzyme itself, and acetylation of PKM2 K433

disrupts FBP binding and reduces enzyme activity (22, 49). A decrease in PK activity

suppresses glycolysis and gates the access of nutrients to the TCA cycle (Figure

7G). These effects restrain the TCA cycle for the endogenous production of TCA

intermediates, including citrate (Figure 6A, left), and lead to an insufficient supply of

building blocks for the synthesis of macromolecules, including amino acids and

nucleotides (Figure 6A, right), which are essential for cancer cell growth (4). Our

study revealed that SLC13A2 inhibits HCC cell proliferation by inducing the

acetylation and degradation of PKM2 (Figure 7D-F).

Our data indicate that SLC13A2 is a potential drug target for HCC treatment. Recent

RNA-seq data analysis revealed that several anticancer drugs, including nilotinib and

imatinib (GSE19567), elevated the transcription of SLC13A2. It has been reported

that the transcription factor HNF4a is involved in the downregulated expression of

SLC13A2, suggesting that saturated fatty acid and inflammatory signals are factors

that inhibit SLC13A2 gene transcription (GSE210842). Similarly, a HFD potently

decreases the expression of SLC13A2 (GSE68360). A deep understanding of the

regulatory mechanisms of SLC13A2 expression during HCC tumorigenesis will

provide insightful suggestions for drug discovery strategies targeting this transporter.

To date, few studies have linked SLC13A2 with cancer, and no investigations have

investigated its roles and mechanisms in cancer pathogenesis. According to the

database of the Human Protein Atlas (www.proteinatlas.org) (50), the expression of

SLC13A2 has low cancer specificity. A high expression level of SLC13A2 is positively

associated with a better overall survival rate in patients with renal cancer, suggesting

its role in improving the prognosis of cancer patients. Transcriptome analysis of

primary colorectal cancer tissues from patients revealed that SLC13A2 is a

representative differentially expressed gene that is responsible for liver metastasis

(51). Another study conducted a genome-wide association analysis with familial

HBV-related HCC in comparison with non-HCC controls with chronic HBV to identify

susceptibility loci (52). The results revealed a single-nucleotide polymorphism cluster

located at the 3' end of SLC13A2, which is among the genes most strongly

associated with familial HBV-associated HCC (52). This finding is in accordance with

our findings suggesting that SLC13A2 is a hereditary genetic factor related to the

pathogenesis of HCC.

Acknowledgments

The authors would like to express their appreciation to Dr. Tongyu Liu for technical

support with the bioinformatic analysis and Dr. Xin Chen (UCSF) for sharing the

oncogene plasmids.

Declarations

Ethical Approval

All animal experiments were approved by the IACUC (Institutional Animal Care and Use Committee) of China Pharmaceutical University (Approval number: 2020-08-006). Human HCC liver samples were obtained in the partial liver resection surgery from Jiangsu Taizhou People's Hospital (Taizhou, Jiangsu, China) under the protocol approved by Medical Ethics Committee of Taizhou people's Hospital (Approval number: 2022-03-017). The study was conducted in accordance with the ethical requirements in the Declaration of Helsinki. Written informed consent was provided by all participants.

Competing interests

The authors declare no competing interests.

Authors' contributions

Q.M., S.L., C.H., L.C., T.Z., H.D. and S.C. performed the research; X.J. conceived the study; X.J. and H.H. designed the research study; X.J. and H.H. contributed essential reagents or tools; X.J., C.H., A.D. and Y.W. analyzed the RNA-seq datasets; Y.H. collected the clinical samples; Q.M., L.C. Y.S. and X.J. analyzed the data; and X.J., Q.M., and L.C. wrote the paper. All the authors have read and approved the final manuscript.

Funding

This study was supported by the National Natural Science Foundation of China (No. 82070883 and 82273982 to J.X.), the National Natural Science Foundation of Jiangsu Province (No. BK20221525 to J.X.) and the Scientific Research Foundation for High-level Faculty, China Pharmaceutical University (to J.X.).

Availability of data and materials

RNA-seq dataset generated in this study has been deposited into the Sequence

Read Archive (PRJNA1083485) and will be accessible with the following link upon

publication (

https://dataview.ncbi.nlm.nih.gov/object/PRJNA1083485?reviewer=l0oikm8iveidksmc

2bjcir9lp2).

References

1. Faubert B, Solmonson A, and DeBerardinis RJ. Metabolic reprogramming and
cancer progression. *Science.* 2020;368(6487).
2. Du D, Liu C, Qin M, Zhang X, Xi T, Yuan S, et al. Metabolic dysregulation and
emerging therapeutical targets for hepatocellular carcinoma. *Acta Pharm Sin*
B. 2022;12(2):558-80.
3. Martinez-Reyes I, and Chandel NS. Mitochondrial TCA cycle metabolites
control physiology and disease. *Nat Commun.* 2020;11(1):102.
4. Anderson NM, Mucka P, Kern JG, and Feng H. The emerging role and
targetability of the TCA cycle in cancer metabolism. *Protein & cell.*
2018;9(2):216-37.
5. Zhu J, and Thompson CB. Metabolic regulation of cell growth and
proliferation. *Nat Rev Mol Cell Biol.* 2019;20(7):436-50.
6. Payen VL, Mina E, Van Hee VF, Porporato PE, and Sonveaux P.
Monocarboxylate transporters in cancer. *Mol Metab.* 2020;33:48-66.
7. Choi I, Son H, and Baek JH. Tricarboxylic Acid (TCA) Cycle Intermediates:
Regulators of Immune Responses. *Life (Basel).* 2021;11(1).
8. Chatzikyriakidou Y, Ahn DH, Nji E, and Drew D. The GFP thermal shift assay for
screening ligand and lipid interactions to solute carrier transporters. *Nature*
protocols. 2021;16(12):5357-76.
9. Zhang Y, Zhang Y, Sun K, Meng Z, and Chen L. The SLC transporter in nutrient
and metabolic sensing, regulation, and drug development. *J Mol Cell Biol.*
2019;11(1):1-13.
10. Lin L, Yee SW, Kim RB, and Giacomini KM. SLC transporters as therapeutic
targets: emerging opportunities. *Nat Rev Drug Discov.* 2015;14(8):543-60.
11. Schulte ML, Fu A, Zhao P, Li J, Geng L, Smith ST, et al. Pharmacological
blockade of ASCT2-dependent glutamine transport leads to antitumor
efficacy in preclinical models. *Nat Med.* 2018;24(2):194-202.
12. Du D, Qin M, Shi L, Liu C, Jiang J, Liao Z, et al. RNA binding motif protein
45-mediated phosphorylation enhances protein stability of ASCT2 to promote
hepatocellular carcinoma progression. *Oncogene.* 2023;42(42):3127-41.
13. Alexander SP, Kelly E, Mathie A, Peters JA, Veale EL, Armstrong JF, et al. THE
CONCISE GUIDE TO PHARMACOLOGY 2021/22: Transporters. *Br J Pharmacol.*
2021;178 Suppl 1:S412-S513.
14. Chen X, and Calvisi DF. Hydrodynamic transfection for generation of novel
mouse models for liver cancer research. *Am J Pathol.* 2014;184(4):912-23.
15. Chu VT, Weber T, Graf R, Sommermann T, Petsch K, Sack U, et al. Efficient
generation of Rosa26 knock-in mice using CRISPR/Cas9 in C57BL/6 zygotes.
BMC biotechnology. 2016;16:4.
16. Xiong J, Liu T, Mi L, Kuang H, Xiong X, Chen Z, et al. hnRNPU/TrkB Defines a
Chromatin Accessibility Checkpoint for Liver Injury and Nonalcoholic

Steatohepatitis Pathogenesis. *Hepatology (Baltimore, Md)*. 2020;71(4):1228-
46.

17. Tang M, Zhao Y, Zhao J, Wei S, Liu M, Zheng N, et al. Liver cancer
heterogeneity modeled by in situ genome editing of hepatocytes. *Science*
advances. 2022;8(25):eabn5683.

18. Reparaz D, Ruiz M, Llopiz D, Silva L, Vercher E, Aparicio B, et al. Neoantigens
as potential vaccines in hepatocellular carcinoma. *J Immunother Cancer*.
2022;10(2).

19. Zhang LX, Lv Y, Xu AM, and Wang HZ. The prognostic significance of serum
gamma-glutamyltransferase levels and AST/ALT in primary hepatic carcinoma.
BMC cancer. 2019;19(1):841.

20. Bergeron MJ, Clémençon B, Hediger MA, and Markovich D. SLC13 family of
Na+-coupled di- and tri-carboxylate/sulfate transporters. *Molecular Aspects*
of Medicine. 2013;34(2):299-312.

21. Feng J, Li J, Wu L, Yu Q, Ji J, Wu J, et al. Emerging roles and the regulation of
aerobic glycolysis in hepatocellular carcinoma. *Journal of Experimental &*
Clinical Cancer Research. 2020;39(1):126.

22. Lv L, Li D, Zhao D, Lin R, Chu Y, Zhang H, et al. Acetylation targets the M2
isoform of pyruvate kinase for degradation through chaperone-mediated
autophagy and promotes tumor growth. *Molecular cell*. 2011;42(6):719-30.

23. Li M, Lu H, Wang X, Duan C, Zhu X, Zhang Y, et al. Pyruvate kinase M2
(PKM2) interacts with activating transcription factor 2 (ATF2) to bridge
glycolysis and pyroptosis in microglia. *Mol Immunol*. 2021;140:250-66.

24. Gao X, Wang H, Yang JJ, Liu X, and Liu ZR. Pyruvate kinase M2 regulates
gene transcription by acting as a protein kinase. *Mol Cell*. 2012;45(5):598-609.

25. Xu Q, Liu LZ, Yin Y, He J, Li Q, Qian X, et al. Regulatory circuit of PKM2/NF-
kappaB/miR-148a/152-modulated tumor angiogenesis and cancer
progression. *Oncogene*. 2015;34(43):5482-93.

26. Xia L, Jiang Y, Zhang XH, Wang XR, Wei R, Qin K, et al. SUMOylation
disassembles the tetrameric pyruvate kinase M2 to block myeloid
differentiation of leukemia cells. *Cell Death Dis*. 2021;12(1):101.

27. Lv WW, Liu D, Liu XC, Feng TN, Li L, Qian BY, et al. Effects of PKM2 on global
metabolic changes and prognosis in hepatocellular carcinoma: from gene
expression to drug discovery. *BMC Cancer*. 2018;18(1):1150.

28. Liberti MV, and Locasale JW. The Warburg Effect: How Does it Benefit
Cancer Cells? *Trends Biochem Sci*. 2016;41(3):211-8.

29. Pajor AM. Molecular properties of the SLC13 family of dicarboxylate and
sulfate transporters. *Pflugers Arch*. 2006;451(5):597-605.

30. Bergeron MJ, Clemencon B, Hediger MA, and Markovich D. SLC13 family of
Na(+)-coupled di- and tri-carboxylate/sulfate transporters. *Mol Aspects Med*.
2013;34(2-3):299-312.

31. Colas C, Pajor AM, and Schlessinger A. Structure-Based Identification of

Inhibitors for the SLC13 Family of Na(+)/Dicarboxylate Cotransporters.
Biochemistry. 2015;54(31):4900-8.

32. Birkenfeld AL, Lee HY, Guebre-Egziabher F, Alves TC, Jurczak MJ, Jornayvaz
FR, et al. Deletion of the mammalian INDY homolog mimics aspects of dietary
restriction and protects against adiposity and insulin resistance in mice. *Cell*
metabolism. 2011;14(2):184-95.

33. Li Z, Li D, Choi EY, Lapidus R, Zhang L, Huang SM, et al. Silencing of solute
carrier family 13 member 5 disrupts energy homeostasis and inhibits
proliferation of human hepatocarcinoma cells. *The Journal of biological*
chemistry. 2017;292(33):13890-901.

34. Ruprecht JJ, and Kunji ERS. The SLC25 Mitochondrial Carrier Family:
Structure and Mechanism. *Trends Biochem Sci*. 2020;45(3):244-58.

35. Mosaoa R, Kasprzyk-Pawelec A, Fernandez HR, and Avantaggiati ML. The
Mitochondrial Citrate Carrier SLC25A1/CIC and the Fundamental Role of
Citrate in Cancer, Inflammation and Beyond. *Biomolecules*. 2021;11(2).

36. Fernandez HR, Gadre SM, Tan M, Graham GT, Mosaoa R, Ongkeko MS, et al.
The mitochondrial citrate carrier, SLC25A1, drives stemness and therapy
resistance in non-small cell lung cancer. *Cell Death Differ*. 2018;25(7):1239-58.

37. Jiang L, Shestov AA, Swain P, Yang C, Parker SJ, Wang QA, et al. Reductive
carboxylation supports redox homeostasis during anchorage-independent
growth. *Nature*. 2016;532(7598):255-8.

38. Tan M, Mosaoa R, Graham GT, Kasprzyk-Pawelec A, Gadre S, Parasido E, et
al. Inhibition of the mitochondrial citrate carrier, Slc25a1, reverts steatosis,
glucose intolerance, and inflammation in preclinical models of NAFLD/NASH.
Cell Death Differ. 2020;27(7):2143-57.

39. Wang G, Xia J, Chen C, Qiu J, Sun P, Peng Z, et al. SLC25A10 performs an
oncogenic role in human osteosarcoma. *Oncol Lett*. 2020;20(4):2.

40. Zhou X, Paredes JA, Krishnan S, Curbo S, and Karlsson A. The mitochondrial
carrier SLC25A10 regulates cancer cell growth. *Oncotarget*. 2015;6(11):9271-
83.

41. An YA, Chen S, Deng Y, Wang ZV, Funcke JB, Shah M, et al. The
mitochondrial dicarboxylate carrier prevents hepatic lipotoxicity by inhibiting
white adipocyte lipolysis. *J Hepatol*. 2021;75(2):387-99.

42. Kunji ERS, King MS, Ruprecht JJ, and Thangaratnarajah C. The SLC25 Carrier
Family: Important Transport Proteins in Mitochondrial Physiology and
Pathology. *Physiology (Bethesda)*. 2020;35(5):302-27.

43. Iacobazzi V, and Infantino V. Citrate--new functions for an old metabolite.
Biol Chem. 2014;395(4):387-99.

44. Icard P, Fournel L, Coquerel A, Gligorov J, Alifano M, and Lincet H. Citrate
targets FBPase and constitutes an emerging novel approach for cancer
therapy. *Cancer Cell Int*. 2018;18:175.

45. Munday MR. Regulation of mammalian acetyl-CoA carboxylase. *Biochem*

Soc Trans. 2002;30(Pt 6):1059-64.

46. Icard P, and Lincet H. The reduced concentration of citrate in cancer cells:
An indicator of cancer aggressiveness and a possible therapeutic target. *Drug*
resistance updates : reviews and commentaries in antimicrobial and anticancer
chemotherapy. 2016;29:47-53.

47. Feng J, Li J, Wu L, Yu Q, Ji J, Wu J, et al. Emerging roles and the regulation of
aerobic glycolysis in hepatocellular carcinoma. *Journal of experimental &*
clinical cancer research : CR. 2020;39(1):126.

48. Hitosugi T, Kang S, Vander Heiden MG, Chung TW, Elf S, Lythgoe K, et al.
Tyrosine phosphorylation inhibits PKM2 to promote the Warburg effect and
tumor growth. *Science signaling.* 2009;2(97):ra73.

49. Lv L, Xu YP, Zhao D, Li FL, Wang W, Sasaki N, et al. Mitogenic and oncogenic
stimulation of K433 acetylation promotes PKM2 protein kinase activity and
nuclear localization. *Molecular cell.* 2013;52(3):340-52.

50. Thul PJ, and Lindskog C. The human protein atlas: A spatial map of the
human proteome. *Protein Sci.* 2018;27(1):233-44.

51. Wang S, Zhang C, Zhang Z, Qian W, Sun Y, Ji B, et al. Transcriptome analysis
in primary colorectal cancer tissues from patients with and without liver
metastases using next-generation sequencing. *Cancer Med.* 2017;6(8):1976-
87.

52. Lin YY, Yu MW, Lin SM, Lee SD, Chen CL, Chen DS, et al. Genome-wide
association analysis identifies a GLUL haplotype for familial hepatitis B virus-
related hepatocellular carcinoma. *Cancer.* 2017;123(20):3966-76.

Figure 1

Figure 1. SLC13A2 is a transporter for TCA cycle intermediates deregulated in

HCC.

A. The SLC transporters for TCA cycle intermediates are summarized.

B. Heatmap of detectable SLC transporters for TCA cycle intermediates in

heterogeneous primary liver cancer (PLC) models established using hydrodynamic tail

vein injection (HTVi)-generated genome editing of cancer driver genes selected by

mutational frequency from human HCC cohorts (SRA accession code PRJNA674008).

C. Schematic showing the study design of the HTVi model by injecting sleeping beauty

(SB) transposase plasmids combined with c-Met and ΔN90-β-catenin oncogene

plasmids (c-MET: ΔN90-β-catenin:SB = 20 μg:20 μg:5 μg).

D. Representative liver morphology after dissection (left, Scale bar: 5 mm) and

histological staining (right, Scale bar: 200 μm). The arrows indicate areas with densely

packed and multinucleated hepatocytes with an increased nucleo-plasma ratio.

E. Liver/body weight ratio (left) and mRNA expression of metabolic genes in liver tissues

of the HTVi model (right) (n=4 for the control group and n=5 for the HCC model

group).

F. Methodology to screen candidate SLC transporters for HCC progression based on the

relative expression of SLC transporters in the HTVi model, mouse HCC cells, AML12

cells and mouse primary hepatocytes (MPHs). The SLC transporters that are

detectable for TCA intermediates in liver tissues of the HTVi model (n=4 for the control

group and n=5 for the HCC model group), AML12 cells, primary hepatocytes and

mouse HCC Hepa1-6 cells were validated by qPCR (n=3 independent experiments).

The data are presented as the means ± SEMs. *P < 0.05, **P < 0.01, ns = not significant;

two-tailed unpaired Student's *t* test.

Figure 2

Figure 2. SLC13A2 expression is decreased in human HCC specimens and cell

lines, which is consistent with the findings in animal models.

A. Immunoblots of total liver lysates from the HTVi model.

B-C. Analysis of published RNA-Seq data of human HCC cancerous and paracancerous

specimens from GEO datasets (GSE193567). Ontology enrichment analysis of

differentially expressed genes (B, left). Heatmap showing the detectable SLC

transporters for TCA cycle intermediates (B, right). Volcano plots showing DEGs with

emphasis on SLC13A2 (C, left). The expression level of SLC13A2 in tumor tissues

compared with that in para-tumor tissues (C right). The data are presented as the

mean ± SEM (n=9). **P < 0.01, tumor versus para-tumor tissues, two-tailed unpaired

Student's *t* test.

D. Kaplan–Meier (KM) analysis of survival in male patients with HCC stratified by high or

low SLC13A2 expression is shown.

E. The RNA expression of SLC13A2 in human HCC cell lines versus L02 cells after being

split and cultured for 48 h.

F. The RNA and protein expression of SLC13A2 was analyzed in eight clinical HCC

patients. PC, paracancerous; C, cancerous.

Figure 3

Figure 3. Liver-specific knockout of SLC13A2 promotes HCC tumor growth.

A. A diagram of Cas9-mediated liver-specific knockout (LKO) of SLC13A2 and study

design. AAV-TBG-Cre-U6-Lacz (Ctrl) and AAV-TBG-Cre-U6-sgRNA (LKO) viruses (5

× 10^{11} VP/mouse) were injected into *Rosa26-FSF-Cas9* knock-in mice via the tail vein.

The transduced mice were subjected to the HTVi model by injecting SB in

combination with oncogene plasmids (c-MET: ΔN90-β-catenin: SB = 20 μg:20 μg:5

μg).

B. qPCR analysis (left, n=6 per group) and immunoblots (right) showing the SLC13A2

knockout efficiency in mouse liver tissues.

C. Representative liver morphology after dissection. Scale bar: 5 mm.

D. Liver/body weight ratio. n=16 per group.

E. Tumor incidence showing the number of mice with or without nodules (left) and the

maximal tumor size (right, n=13 per group) in the two groups.

F. Liver ALT/AST and serum ALT/AST activity.

G. Representative images of H&E staining.

H. Immunohistochemical staining for Ki67. The Ki67-positive ratio was calculated (n=6

per group).

I. Immunoblots of total liver lysates showing cell signaling for tumor growth.

The data are presented as the means ± SEMs. *P < 0.05, LKO vs Ctrl group, two-tailed

unpaired Student's *t* test.

 Figure 4

1015

Figure 4. Liver-specific overexpression of SLC13A2 inhibits HCC tumor

progression.

A. Schematic diagram of the liver-specific overexpression of SLC13A2 and study design.

AAV-TBG-GFP (GFP) and AAV-TBG-SLC13A2 (SLC13A2) viruses (2 × 10^{11}

VP/mouse) were injected into wild-type mice via the tail vein. The transduced mice

were subjected to the HTVi model by injecting SB in combination with oncogene

plasmids (c-MET: ΔN90-β-catenin: SB = 20 μg:20 μg:5 μg).

B. qPCR analysis (left, n=6 per group) and immunoblots (right) showing SLC13A2

overexpression efficiency in mouse HCC tissues.

C. Representative images showing liver morphology upon dissection. Scale bar: 5 mm.

D. Liver/body weight ratio (n=14 for the GFP group and n=12 for the SLC13A2 OE

group).

E. Tumor incidence showing the number of mice with or without nodules (left) and the

maximal tumor size (right, n=12 for the GFP group and n=10 for the SLC13A2 OE

group) in the two groups.

F. Liver and serum ALT/AST activity.

G. Representative images of H&E staining.

H. Immunohistochemical staining for Ki67. Ratio of Ki67-positive cells in the two groups

(right, n=6 per group).

I. qPCR analysis of liver tissues showing gene transcription for cell proliferation and

metabolism.

The data are presented as the means ± SEMs. *P < 0.05, **P < 0.01, SLC13A2-OE vs

GFP group, two-tailed unpaired Student's *t* test.

1039

 Figure 5

1042 **Figure 5. SLC13A2 inhibits the proliferation of HCC cells *in vitro*.**

A. Knockdown and overexpression efficiency *in vitro.* The protein level of SLC13A2 in Hepa1-6 cells was evaluated after transfection with NC/vector or SLC13A2 siRNA/expression plasmids/adenovirus for 48 h.

B. Colony formation assay showing the clonogenic potential of HCC Hepa1-6 cells after treatment with SLC13A2 KD (siSLC13A2, top), Hepa1-6 cells and HepG2 cells with SLC13A2 OE (bottom) and culture for 7 or 14 days. The number of colonies was counted.

C. The growth of HCC cells with SLC13A2 knockdown (left) or overexpression (right) was analyzed by CCK8 assays at 0, 24, 48, 72, 96, and 120 h after transfection. **$P <$ 0.01, two-way ANOVA, followed by Bonferroni multiple comparisons.

D. EdU incorporation was detected in Hepa1-6 cells with SLC13A2 KD (left) or in Hepa1-6 and HepG2 cells with OE (right) for 72 h, followed by incubation with EdU (20 µM) for 3-4 h (scale bar: 200 µm or 50 µm). EdU-positive cells were visualized under a fluorescence microscope, and the percentages are shown below.

E. Propidium iodide (PI) staining and flow cytometry analysis showing cell cycle arrest in Hepa1-6 cells after transfection with SLC13A2 siRNAs or control siRNA (NC) for 48 h (top) or in Hepa1-6 and HepG2 cells after overexpression of SLC13A2 for 72 h (bottom).

Bars represent the means ± SEMs (n=3 or 4 independent experiments). *$P < 0.05$ and **$P < 0.01$, two-tailed unpaired Student's *t* test (B, D-E).

 Figure 6

Figure 6. SLC13A2 restricts the TCA cycle and inhibits mitochondrial respiration.

A-B. SLC13A2 was overexpressed in Hepa1-6 cells, which were subsequently cultured

for an additional 60 h, after which the metabolites were analyzed via untargeted

metabolomics. (**A**) Relative content analysis of TCA cycle substrates (left), amino acid

metabolites (middle) and nucleotide metabolites (right). (**B)** Metabolite set enrichment

analysis (MSEA) of the metabolic pathways (n = 3). adj. P < 0.05 was considered to

indicate statistical significance.

C. Hepa1-6 cells were cultured for 24 h after transfection with the vector or SLC13A2

overexpression plasmid. After the cells were digested, 8,000 cells/well were seeded

for the Seahorse XF assay. Dashed vertical lines indicate the subsequent addition of

the ATPase inhibitor oligomycin (OligO: 1.5 µM), the uncoupling reagent (FCCP: 1.0

µM) and the inhibitors of the electron transport chain rotenone/antimycin A (ROT/AA:

0.5 µM). The oxygen consumption rate (OCR) under basal conditions and after

treatment with the uncoupler FCCP (maximal respiration) were analyzed.

D. HepG2 and Hepa1-6 cells were cultured for 72 h after transfection to detect ATP

production.

E. Confocal images of JC-1 fluorescence. The graph shows the ratio of aggregates (red)

to monomers (green), indicating changes in the mitochondrial membrane potential.

The data are presented as the means ± SEMs (n=3 independent experiments). *$P < 0.05$

and **$P < 0.01$, two-tailed unpaired Student's t test (A, C-E).

1085

Figure 7

Figure 7. SLC13A2 suppresses tumor growth through the acetylation and

degradation of PKM2 by imported citrate.

A. Hepa1-6 cells were transfected with vector or SLC13A2-OE adenovirus and cultured

for an additional 60 h. U-^{13}C$_6$-glucose was then added, and the cells were incubated

for 1 h (A-D) or the indicated period (E). The metabolites were analyzed using an

untargeted metabolomics method.

B. Hepa1-6 cells were transfected with vector or SLC13A2-OE adenovirus and cultured

for 60 h. The relative content of pyruvate was analyzed using targeted metabolomics

(left). The viability of Hepa1-6 cells with or without SLC13A2 overexpression was

measured after treatment with pyruvate for 72 hours (middle). Cell colony formation

assay showing the clonogenic potential of HCC cells after treatment with pyruvate for

7 days in the presence or absence of SLC13A2 overexpression (right). *P < 0.05 vs.

vector without pyruvate, #P < 0.05 vs. SLC13A2 without pyruvate.

C. Hepa1-6 cells were transfected with vector or SLC13A2-OE adenovirus and cultured

for 60 h. The relative levels of intracellular oxaloacetate (left) and extracellular citrate

(middle) were analyzed via targeted metabolomics. Hepa1-6 cell viability was

measured after treatment with an ACLY inhibitor for 72 hours in the presence or

absence of SLC13A2 overexpression (right). *P < 0.05 vs. vector without the ACLY

inhibitor, #P < 0.05 vs. SLC13A2 without the ACLY inhibitor.

D. PKM2 RNA or protein levels in whole-cell lysates, nuclei and cytosol of Hepa1-6 cells

after transfection with the vector or SLC13A2 and culture for 60 h.

E. Co-IP assay showing the acetylation of PKM2 in Hepa1-6 cells with or without

SLC13A2 overexpression. IgG served as a negative control.

F. Cycloheximide (CHX) chase assay showing the effect of SLC13A2 on the half-life of

the PKM2 protein in Hepa1-6 cells by immunoblotting (left) and densitometry analysis

(right).

G. Pyruvate kinase activity in Hepa1-6 cells after transfection for 60 h.

H. Hepa1-6 cells were transfected with vector or SLC13A2-OE adenovirus in the

presence or absence of WT or mutant PKM2. Immunoblots showing the PKM2 protein

level after culture for 60 h.

I. Immunoblot analysis showing the SLC13A2 and/or PKM2 overexpression efficiency in

Hepa1-6 cells after transfection. The growth curves of Hepa1-6 cells overexpressing

SLC13A2 and/or PKM2 were analyzed by CCK8 assays at 0, 24, 72 and 120 h after

transfection. Cell colony formation assay showing the clonogenic potential of HCC

cells after transfection with SLC13A2 and/or PKM2 and culture for 7 days. *P < 0.05

vs. Vector, #P<0.05 vs. SLC13A2.

The data are presented as the means ± SEMs (n=3 or 4 independent experiments). *P <

0.05, two-tailed unpaired Student's *t* test (A, B left, C left and middle, G); *P < 0.05,

two-way ANOVA, followed by Bonferroni multiple comparisons (B middle, C right, F, I).

Figure 8

1129

Figure 8. SLC13A2 suppresses tumor growth by reprograming PKM2-dependent

gene transcription for cell proliferation and metabolism.

A. Volcano plot of hepatic genes upregulated or downregulated in SLC13A2-OE mice

(left). Representative genes involved in cell proliferation and metabolism are shown

(right).

B. Enrichment of biological processes in the two clusters is indicated for SLC13A2-OE

mice.

C. Gene set enrichment analysis of all DEGs in SLC13A2-OE mice, showing the top two

DEGs.

D. Motif enrichment analysis of predicted transcription factors in SLC13A2-OE mice.

E. The Venn diagrams illustrates the conservative DEGs shared between PKM2-

regulated genes in the TCGA HCC level 3 clinical patient cohort and the SLC13A2-

regulated genes in HCC mouse model (top), and the enrichment of biological

processes shared by the regulatory gene clusters regulated by PKM2 and SLC13A2

(bottom).

F. Gene ontology enrichment by the PKM2-regulated gene cluster (left). The

representative enriched biological processes shared by the regulatory gene clusters of

PKM2 and those of SLC13A2 (right).

G. Motif enrichment showing representative predicted transcription factors shared

among the clusters of PKM2-regulated, SLC13A2-upregulated and SLC13A2-

downregulated genes.

1151

1152

1153

1154

1155 **Graphic abstract**

1156