Control of 3' splice site selection by the yeast splicing factor Fyv6

Katherine A. Senn[1]*, Karli A. Lipinski[2]*, Natalie J. Zeps[1], Amory F. Griffin[1],

Max E. Wilkinson[3, 4], Aaron A. Hoskins[1,2]

*indicates co-first authors

[1] Department of Biochemistry, University of Wisconsin-Madison, Madison, WI 53706 USA

[2] Department of Chemistry, University of Wisconsin-Madison, Madison, WI 53706 USA

[3] MRC Laboratory of Molecular Biology, Cambridge CB2 0QH UK

[4] Present Addresses: Broad Institute of MIT and Harvard, Cambridge MA 02142 USA and McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA 02139 USA

CORRESPONDING AUTHORS:

Aaron A. Hoskins, ahoskins@wisc.edu

Max E. Wilkinson, mwilkin@mit.edu

RUNNING TITLE: Structure and Function of the Splicing Factor Fyv6

KEYWORDS: yeast, spliceosome, pre-mRNA splicing, cryo-EM, snRNA, snRNP, structure, RNA-Seq

ABSTRACT

Pre-mRNA splicing is catalyzed in two steps: 5' splice site (SS) cleavage and exon ligation. A number of proteins transiently associate with spliceosomes to specifically impact these steps (1st and 2nd step factors). We recently identified Fyv6 (FAM192A in humans) as a 2nd step factor in *S. cerevisiae*; however, we did not determine how widespread Fyv6's impact is on the transcriptome. To answer this question, we have used RNA-seq to analyze changes in splicing. These results show that loss of Fyv6 results in activation of non-consensus, branch point (BP) proximal 3' SS transcriptome-wide. To identify the molecular basis of these observations, we determined a high-resolution cryo-EM structure of a yeast product complex spliceosome containing Fyv6 at 2.3 Å. The structure reveals that Fyv6 is the only 2nd step factor that contacts the Prp22 ATPase and that Fyv6 binding is mutually exclusive with that of the 1st step factor Yju2. We then use this structure to dissect Fyv6 functional domains and interpret results of a genetic screen for *fyv6Δ* suppressor mutations. The combined transcriptomic, structural, and genetic studies allow us to propose a model in which Yju2/Fyv6 exchange facilitates exon ligation and Fyv6 promotes usage of consensus, BP distal 3' SS.

INTRODUCTION

Precursor messenger RNA (pre-mRNA) splicing is catalyzed by a large macromolecular complex called the spliceosome. Spliceosomes are composed of 5 small nuclear ribonucleoproteins (snRNPs), each composed of a small nuclear RNA (snRNA; U1, U2, U4, U5, and U6) and several different protein splicing factors. The snRNPs and dozens of other proteins assemble *de novo* at each intron to form spliceosomes. Spliceosomes are highly dynamic and form different complexes as proteins and snRNAs join and leave. Many of these complexes have been characterized biochemically, genetically, and by cryo-electron microscopy (cryo-EM) (Plaschka et al., 2019). The splicing reaction itself is carried out in two sequential transesterification reactions (**Figure 1A**). First, the 5' splice site (5' SS) is cleaved during formation of an intron lariat (1st step). Second, the intron lariat is removed simultaneously with exon ligation at the 3' SS (2nd step).

The integrity of the genetic information contained within a pre-mRNA depends on correct identification of the 5' and 3' SS by the splicing machinery since a single nucleotide shift in either site could destroy a protein reading frame. In yeast, several DExD/H-box ATPases function to limit usage of suboptimal SS by enhancing splicing fidelity (Chung et al., 2023; Semlow and Staley, 2012). Two of these ATPases, Prp16 and Prp22, impact the fidelity of the 1st and 2nd catalytic steps, respectively. In addition to these roles, Prp16 is also required for remodeling of the spliceosome to permit the 1st-to-2nd step transition (Schwer and Guthrie, 1992), and Prp22 is essential for releasing the mRNA product (Company et al., 1991). While Prp16 does not need to be present during the 1st step (Chung et al., 2023), Prp22 is required to be present for exon ligation if the branch point (BP) to 3' SS distance is $\geq$ 21 nucleotides (nt) (Schwer and Gross, 1998). Despite many biochemical, single molecule, and structural studies, how spliceosomes promote usage of BP distal, Prp22-dependent 3' SS has remained elusive.

Identification of the correct 3' SS is an especially challenging problem given that the consensus sequence in *Saccharomyces cerevisiae* (yeast) and humans is just three nucleotides

(YAG, Y=U or C). Cryo-EM structures of spliceosome product (P) complexes have revealed how

these nucleotides can be recognized within the spliceosome active site (Bai et al., 2017; Fica et

al., 2019; S. Liu et al., 2017; Wilkinson et al., 2017; Zhang et al., 2019). The AG 3' SS dinucleotide

is recognized by formation of non-Watson Crick base-pairing interactions with the 5' SS +1G and

BP adenosine, while the pyrimidine at the -3 position of the 3' SS is recognized by the Prp8

protein. Given this short consensus sequence, multiple different splicing factors also help to

ensure that the correct 3' SS is utilized, including Cwc21 and the 2^{nd} step factors Slu7, Prp18, and

Prp22 (Crotti et al., 2007; Frank and Guthrie, 1992; Gautam et al., 2015; Kawashima et al., 2014;

Roy et al., 2023; Semlow et al., 2016). It is believed that Slu7 and Prp18 help to stabilize 3' SS

docking to the active site and that this contributes to 2^{nd} step efficiency. It has been proposed that

Prp22 antagonizes 3' SS docking to permit sampling of different 3' SS to ensure that optimal

sequences are used as a proofreading mechanism (Mayas et al., 2006; Semlow et al., 2016).

Importantly, identification of the correct 3' SS depends not just on identification of the YAG

sequence but also involves choosing which YAG sequence to use. Prp18 appears to aid in

selection of BP distal 3' SS and avoidance of BP proximal (and often non-consensus) 3' SS. This

observation could be due to Prp18 imposing a BP to 3' SS distance constraint, enforcing use of

canonical YAG 3' SS sequences, or a combination of the two activities.

Recently, Fyv6 was identified as a novel 2^{nd} step factor in yeast (Lipinski et al., 2023).

Fyv6 is a homolog of the human protein FAM192A, which was identified in pre-C* structures of

the human spliceosome (Zhan et al., 2022). While the impact of FAM192A on human spliceosome

activity has not yet been well-characterized, loss of Fyv6 in yeast decreases 2^{nd} step splicing

efficiency *in vitro* and results in use of an alternative 3' SS in the *SUS1* pre-mRNA *in vivo*. It is

likely that unassigned cryo-EM densities in yeast C* (just prior to exon ligation) and P complexes

correspond to Fyv6 since these densities are in analogous locations to that for FAM192A in the

human pre-C* structure (Zhan et al., 2022). However, the resolutions of the yeast C* and P

complex structures make unambiguous assignment of Fyv6 amino acid side chains and

interactions difficult.

Here, we use a combination of transcriptomic, structural, biochemical, and genetic assays

to elucidate how Fyv6 controls 3' SS usage in yeast. RNA-seq analysis reveals widespread

activation of alternative, BP proximal nonconsensus 3' SS in the absence of Fyv6 consistent with

biochemical assays that show Fyv6 facilitates splicing of 3' SS located $\geq$ 21 nt from the BP. To

further elucidate Fyv6 function, we determined a 2.3 Å cryo-EM structure of the yeast P complex

spliceosome in which Fyv6 can be modeled. The structure reveals interactions between Fyv6 and

Prp22 as well as a domain in Syf1 that interacts with either Fyv6 or the 1st step factor Yju2. 3D

classification allowed us to identify two additional conformational states of P complex and

provides insights into the coupling of 3' SS active site docking and the presence of 2nd step factors.

We then used this structural data to dissect the functional domains of Fyv6, interpret the results

of a genetic screen for suppressors of *fyv6Δ*, and to probe Fyv6 and Prp22 interactions.

Combined, we propose a model in which Fyv6 is recruited to the spliceosome via interactions with

the NTC component Syf1 in order to promote usage of BP distal, and Prp22-dependent, 3' SS.

RESULTS

Deletion of *FYV6* Results in Widespread Use of Alternative 3′ Splice Sites

Our previous work showed that loss of *FYV6* causes changes in 3' SS usage in the *SUS1*

pre-mRNA. To probe for changes transcriptome-wide, we used nonsense mediated decay (NMD)-

deficient (*upf1Δ*, WT) yeast cells expressing or lacking Fyv6. By suppressing NMD, we hoped to

limit degradation of alternatively spliced mRNA isoforms generated in the absence of Fyv6

(Sayani et al., 2008). Consistent with results from *fyv6Δ* strains, the *fyv6Δ upf1Δ* double mutant

strain also showed reduced growth at 30°C and cold-sensitive (*cs*) and temperature-sensitive (*ts*)

phenotypes relative to a *upf1Δ* control strain (**Figure 1B**). We then used RNA-Seq to analyze the

isolated RNAs from these strains. As expected, many more changes in splicing were observed in

the *fyv6Δ upf1Δ* strain relative to the *fyv6Δ* strain (**Figure 1-figure supplement 1A, 1B; supplementary tables 1, 2**). This is consistent with many mRNA isoforms generated due to loss of Fyv6 being substrates for NMD. Loss of Fyv6 resulted in a number of changes in gene expression with many non-intron containing genes also being up- or down-regulated (**Figure 1-figure supplement 1C**).

117   We compared and classified changes in splicing between strains containing or lacking *FYV6* under permissive growth conditions (30°C) or after the yeast had been shifted to a non-permissive temperature (16 or 37°C) for 1 h (**Figure 1C**). In this case, we only considered events resulting in at least a 10% change in the percent spliced in (PSI) value when *FYV6* is deleted. At all temperatures, we see changes in alternative 3' SS usage in the *fyv6Δ* strain relative to the *FYV6* background. For example, we detected use of the alternative 3' SS in the first intron of the *SUS1* transcript as we previously reported (**Figure 1-figure supplement 1D**; Lipinski et al., 2023). We also saw increased use of a cryptic, nonconsensus GAG 3' SS in the first, but not second, intron of the *YOS1* transcript in the *fyv6Δ* data sets and confirmed this result by RT-PCR (**Figure 1-figure supplement 1E, F**). We observed several cases of alternative 5' SS. However, the majority of these alternative 5' SS were used with equal efficiencies in the WT and *fyv6Δ* strains (**Figure 1-figure supplement 1G**).

129   The strains temperature shifted to 16°C had the largest number of changes due to *FYV6* deletion and the strains shifted to 37°C the least. The temperature shift to 16° caused an increase in the numbers of both alternative 3' SS and retained intron events detected (**Figure 1C**). The changes in alternative 3' SS used at each temperature were mostly unique with few events detected under all three conditions (**Figure 1D**). Collectively, we observed that 61 different RNAs (~20% of introns) changed splicing patterns due to loss of Fyv6. For many of these mRNAs, we detected multiple different 3' SS being used.

To investigate these splicing events more closely, we analyzed changes in usage of

alterative splice junctions by calculating the Fraction of Annotated Splicing (FAnS) in the presence

and absence of Fyv6 (Roy et al., 2023). FAnS ratios report the relative abundance of an alternative

splicing event in relation to the main spliced isoform. We calculated FAnS values for all exon

junction reads corresponding to usage of a canonical 5' SS and an alternative 3' SS for WT and

fyv6Δ strains at each temperature. We then calculated the ratios of the *fyv6Δ*/WT FAnS values

for each transcript. These results confirm changes in alternative 3' SS usage for both ribosomal

protein gene (RPG) and non-RPG transcripts at all conditions (**Figure 1E**). We could not detect

any significant differences between RPG and non-RPG splicing with this analysis, suggesting that

differences in gene expression and splicing efficiencies between these two classes are not

correlated with alternative 3' SS usage.   Combined these results show that Fyv6 plays a critical

role in defining mRNA isoform production in yeast, particularly under non-optimal growth

conditions such as temperature stress.

Fyv6 Facilitates Usage of Branch Point Distal 3′ Splice Sites

We next analyzed the features of the 3' SS impacted by Fyv6 loss. We used RNAs

collected from yeast grown at 30°C and sequenced the libraries at higher depth than those used

for the temperature-shift experiment described in **Figure 1** (~400 million vs. 200 million reads;

supplementary table 2). We first identified alternative 3' SS used in each strain, annotated each

as being upstream (5') or downstream (3') of the canonical 3' SS (Grate and Ares, 2002), and

then calculated the FAnS ratio for each site. This analysis shows that the majority of the alternative

3' SS arising from Fyv6 deletion are upstream (**Figure 2A**, pink points above the diagonal). When

these alternative 3' SS are mapped according to their nucleotide distance from the canonical site,

most of these are found within ~40 nt upstream (**Figure 2B**, purple violin). Plotting these

alternative 3' SS sites relative to their distance from the annotated BP reveals that the highest

density of sites is $\leq$ 20 nt from the BP for transcripts with the highest FAnS ratios (**Figure 2C**,

purple violin). It is important to note that we did not map BP in Fyv6 deletion strains, and it is

possible that in some cases a shift in 3' SS usage could also coincide with a shift in the BP. Indeed,

a small number of alternative 3' SS detected in the Fyv6 data sets are located upstream of the

annotated BP, suggesting activation of an alternative BP and 3' SS. However, we believe that the

changes are predominantly due to usage of alternative 3' SS located between the annotated BP

and canonical 3' SS.

The above results suggest that Fyv6 is important for splicing of 3' SS located distant from

the BP. To test this directly and systematically, we created a series of ACT1-CUP1 reporters with

various BP to 3' SS distances (9-50 nt, **Figure 2D**, **supplementary table 3**) based on sequences

used in previous studies of 3' SS selection (Brys and Schwer, 1996; Frank and Guthrie, 1992;

Schwer and Gross, 1998). We transformed strains containing or lacking Fyv6 and without

debranchase (Dbr1) to limit degradation of lariat intermediates. We then detected RNA products

by primer extension and calculated the 2^{nd} step (exon ligation) efficiency for each reporter (**Figure**

2E, F). While loss of Fyv6 has minimal impact on exon ligation when BP to 3' SS distances are

short (9-15 nt), exon ligation efficiency decreases significantly at distances of 21 nt or greater. In

addition, we do not see evidence of cryptic BP usage in these assays. This result is consistent

with the RNA-seq analysis and supports a function for Fyv6 in facilitating splicing at BP distal 3'

SS.

Finally, we analyzed the sequence features of the alternative 3' SS with increased use

after loss of Fyv6 (**Figure 2G, H**). For sites located upstream of the canonical 3' SS, we observed

much greater sequence variability relative to the canonical site. We detected increased use of

highly variable 3' SS including those with atypical HAU (H=A,C,U) and BG (B=C,U,G) motifs and

confirmed generation of new RNA isoforms for several of these by RT-PCR (**Figure 2H, figure 2-**

figure supplement 1). Increased use of variable 3' SS was also observed when the 2^{nd} step

factor *PRP18* was deleted (Roy et al., 2023). However, the subset of RNAs most impacted by

splicing factor deletion appear to be different. For example, we were not able to detect changes

in 3' SS usage for the *UBC12*, *MAF1*, *MUD1*, *PHO85*, *SPT14*, or *YCL002C* transcripts in the *fyv6Δ*

data sets as was reported for *prp18∆* (not shown). Prp18 and Fyv6 likely have transcript-specific

effects on splicing outcomes. Combined our results strongly support Fyv6 as a 2nd step splicing

factor that facilitates usage of consensus sites located distal to the BP *in vivo*.

A High-Resolution Spliceosome Structure Reveals Fyv6 Interactions

Previous structures of the yeast P-complex spliceosome were solved with resolutions at

the core ranging from 3.3 Å to 3.7 Å (Bai et al., 2017; S. Liu et al., 2017; Wilkinson et al., 2017),

and with peripheral components such as the Prp22 helicase, U2 snRNP, and Prp19 Complex

(NTC) at much lower resolutions (5-10 Å) that precluded detailed investigation. To improve the

resolution, we purified P complex as previously described by stalling spliceosome disassembly

with a dominant negative mutant of Prp22 (S635A) defective in mRNA release (Schwer and

Meszaros, 2000; Wilkinson et al., 2017). However, by collecting a much larger cryo-EM dataset

on a more modern electron detector, we were able to resolve the structure to 2.3 Å within the

catalytic core, as well as visualize peripheral components at resolutions from 3.0 Å – 3.7 Å (**Figure**

3A, B; figure 3-figure supplements 1-4; supplementary table 4). This is the highest resolution

spliceosome structure to date. The high-quality density at the active site shows the positions of

bases unambiguously, confirming the manner of the non-Watson-Crick base pairing that mediates

3' SS recognition (S. Liu et al., 2017; Wilkinson et al., 2017). Monovalent ions bound at the active

site were previously shown to be important during the 1st step of splicing (Wilkinson et al., 2021),

and our data indicate that the positions of these ions are preserved after exon ligation. Additionally,

the density also shows a metal ion, likely potassium, that bridges the base pair between +1G of

the 5' SS and -1G of the 3' SS (**Figure 3C**).

The 2nd-step splicing factor Prp22 is resolved at 3.0 Å resolution, allowing atomic

interpretation for the first time (**Figure 3B** and later figures). Bases 13 – 21 of the 3' exon are

visible between the two RecA domains of Prp22 consistent with biochemical footprinting data

(Schwer, 2008) (**figure 3-figure supplement 4**). The S635A mutation used to stall P complex is

also visible; however, the origin of defective mRNA release due to this mutation is not apparent

(Schwer and Meszaros, 2000). The loop containing S635A is in a similar conformation in this

structure and in the structure of WT human Prp22 (Felisberto-Rodrigues et al., 2019) (**figure 3-**

figure supplement 4). Part of the extensive N-terminal domain of Prp22 is resolved and

constitutes a long helix that bridges between the Prp22 RecA2 domain and the C-terminal domain

of Cwc22; this helix was unassigned in previous P-complex structures (**Figure 4B**).

To improve the resolution of the peripheral, flexible regions, we used a new data-driven

regularization strategy (Kimanius et al., 2024) that allows focused refinements of much smaller

domains than previously possible (**figure 3-figure supplement 2, 4**). Combined with AlphaFold2-

assisted modelling, we were able to obtain more accurate models for the U2 snRNP, NTC, U5

Sm ring, and Cwc22 N-terminal domains. Our model for the NTC within the P complex is similar

to our previous model for the NTC within C and C^i complexes (Wilkinson et al., 2021) (**Figure**

4C). Notably however, we observed a star-shaped density coordinated by several conserved

lysines from Clf1 and Ntc20, which we were able to model as inositol hexakisphosphate (IP_6)

(**Figure 4D**). The same density, at lower resolution, is visible in the C complex (Wilkinson et al.,

2021). A distinct, separate IP_6 molecule had previously been observed within all catalytic

spliceosomes between the B^{act} and P-complex stages coordinated by Prp8 and (for C* and P-

complexes) Slu7 (**figure 3-figure supplement 4**). We do not believe that the NTC-bound IP_6 is

essential for splicing since deletion of the Clf1 tetratricopeptide repeat that coordinates IP_6 does

not result in growth or splicing phenotypes in yeast and Ntc20 is a non-essential splicing factor

(Chen et al., 2001; Chung et al., 1999). Nonetheless, it is possible that IP_6 may regulate splicing

in some manner via interactions with both Prp8 and the NTC.

We were able to unambiguously assign three, connected long α-helices to Fyv6 (**Figure**

4A). These helices were visible in previous cryo-EM reconstructions of yeast C* and P complexes

but were either unassigned or misassigned. In the cases of misassigned densities, Fyv6 was

previously attributed to the C-terminal domain of the 1[st] step factor Yju2 (Wilkinson et al., 2017).

Yju2 also has an elongated helical architecture and adopts a similar position on C complex

(Wilkinson et al., 2021) and the post-P complex Intron Lariat Spliceosome (ILS) (Wan et al., 2017).

We closely inspected the density from previously determined spliceosome structures and

concluded that the Yju2 C-terminus is still a better fit than Fyv6 for densities in B*, C, and ILS

complexes, whereas Fyv6 is a better fit than Yju2 for densities in C* and P complexes (**figure 4-**

figure supplement 1). Therefore, this position of Fyv6 seems to be characteristic of a 2^{nd} step

conformation of the spliceosome and consistent with the position of its distant human homolog

FAM192A in the human pre-C* complex (Zhan et al., 2022).

Fyv6 makes a multitude of interactions with essential splicing factors (**Figures 3B; 4E, F,**

G). One conserved region of Fyv6 is the N-terminus with a FVSE motif (**figure 4-figure**

supplement 2). This motif forms a 'hook' that interacts with a hydrophobic patch on the surface

of the Prp22 RecA2 domain, where it also forms hydrogen bonds by β-sheet augmentation

(**Figure 4E**). The hook is followed by three α-helices. The start of helix 2 sits on top of the intron

between the BP and the active site docked 3' SS, where it may act as a steric block to prevent 3'

SS undocking and promote exon ligation (**Figure 4F**) (S. Liu et al., 2017). Helix 2 also contains a

patch of conserved arginines that form salt bridges with Asp317 and Glu318 of Slu7 (**Figure 4F**).

The linker between helices 2 and 3 interacts with the NTC component Cef1, and helix 3 interacts

with NTC component Syf1 (**Figure 4G**). The binding interface with Syf1 overlaps with the

Syf1/Yju2 interaction in the C complex (Wilkinson et al., 2021), suggesting these interactions are

mutually exclusive (**Figure 4G**). Fyv6 thus provides a direct link between the Prp22 ATPase, NTC

components, and other essential splicing factors over a distance of ~100 Å.

By 3D classification, we were able to resolve a total of three states of the P-complex

spliceosome (**Figure 3C-E**). State I, represented by 51% of particles, is the most complete and

has all the above-described factors visible, including a stably-docked 3' SS. State II, represented

by 20% of particles, is similar to State I but lacks density for the N-terminal half of the 2^{nd} step

factor Slu7. The endonuclease-like domain of Prp8, to which this domain of Slu7 binds, is

correspondingly shifted. Interestingly, this state retains strong density for the 3' SS, suggesting

the N-terminal regions of Slu7 are not involved in maintaining 3' SS docking after exon ligation.

Finally, State III, containing 29% of particles, entirely lacks density for the 2nd step factors Slu7,

Prp18, and Fyv6. Prp17 and the RNase H domain of Prp8 are still present but are weaker than in

States I and II, suggesting they are more flexible in State III. Most of Prp22 can still be observed

with weaker density except for the C-terminal domain that interacts with Prp8, which lacks density

entirely. Importantly, State III also lacks density for the 3' SS but has density for the 3' exon,

suggesting it is indeed post-catalytic and represents a state after loss of 2nd step factors and

undocking of the 3' SS from the catalytic core. The coincidence of loss of Fyv6 with loss of Prp18

and Slu7 reinforces the notion that these three factors act together and are important for 3' SS

docking, maintaining the 3' SS in the active site after exon ligation, and for stabilization of Prp17,

Prp22, and the RNase H domain of Prp8.

The Fyv6/Syf1 Interaction is Critical for Suppressing *fyv6Δ* Phenotypes

To determine the functionally critical regions and interactions of Fyv6, we used the cryo-

EM structure to design five truncation mutants (**Figure 5A**). The first three truncations remove

amino acids from the N-terminus of Fyv6: deletion of the first 16 amino acids, which are not

resolved in the cryo-EM structure (Δ1-16); deletion of the first 23 amino acids, which removes the

conserved hook region of Fyv6 that interacts with the Prp22 RecA2 domain (Δ1-23); and deletion

of amino acids 1 to 51, which additionally removes the first α-helix and interactions with Prp22,

Prp8, and Slu7 (Δ1-51). The remaining two mutants are C-terminal truncations. In the first, amino

acids 134 to 173, which do not have clear density in the cryo-EM structure and are not modeled

but which do contain a predicted nuclear localization signal (NLS), are deleted (Δ134-173)

(Nguyen Ba et al., 2009). In the second, amino acids 103 to 173 are deleted (Δ103-173), which

additionally removes the region of the protein that interacts with Syf1.

We tested each truncation mutant for its ability to suppress the *cs* and *ts* phenotypes of

the *fyv6Δ* strain when expressed from a plasmid under the native promoter (**Figure 5B**). All of the

N-terminal truncation mutations could suppress the *cs* or *ts* phenotypes to some degree.

Surprisingly, a strain with the complete deletion of the conserved hook domain that interacts with

Prp22 (Δ1-23) grew similarly to the strain expressing WT Fyv6. We wondered if splicing changes

could still be present even if the growth phenotype was suppressed; so, we assayed *SUS1*

isoforms by RT-PCR. We previously showed that loss of Fyv6 causes use of an alternative 3' SS

in the first intron of this pre-mRNA (Lipinski et al., 2023). In addition to suppression of the *fyv6Δ*

growth phenotype, the N-terminal truncation mutants also showed no evidence for changes in

SUS1 splicing (**Figure 5C**). The N-terminal region of Fyv6 composed of the hook domain and the

first α-helix is not required for suppression of either the temperature sensitivity or splicing

phenotypes observed when Fyv6 is lost despite its conservation and interactions with critical

splicing factors (Prp8, Prp22, and Slu7).

For the C-terminal truncations, the Δ134-173 truncation was also able to suppress the *cs*

and *ts* phenotypes, indicating that the predicted NLS is also not essential (**Figure 5B**). However,

this mutant grew more poorly at 16°C than did the Δ1-16 and Δ1-23 truncations and showed some

evidence of alternative 3' SS usage in *SUS1* (**Figure 5C**). The largest effects were seen when the

C-terminus was truncated further. The Δ103-173 truncation strain phenocopied the *fyv6Δ* strain

and grew poorly at 37°C and was dead at 23° or 16°C (**Figure 5B**). Additionally, RT-PCR showed

increased use of the alternative 3' SS in *SUS1* with this truncation (**Figure 5C**). To determine if

this phenotype could be due to loss of protein expression, we assayed expression of the epitope-

tagged proteins by western blot (**figure 5-figure supplement 1A**). In all cases, we could detect

protein; however, the abundance of the N-terminally epitope-tagged Δ103-173 variant was much

lower than the others. Therefore, we also constructed a C-terminally epitope-tagged version of

Δ103-173 (**figure 5-figure supplement 1B**). The C-terminally-tagged protein expressed at much

higher levels but also failed to suppress the *cs* and *ts* phenotypes. These data suggest that the

C-terminal region of Fyv6 that includes a portion of α-helix 3 and the Syf1 interaction domain is

critical for function.

To obtain additional evidence for the importance of the Fyv6/Syf1 interaction, we created

a *syf1Δ* shuffle strain containing WT Fyv6 and expressed C-terminal Syf1 truncations (**Figure**

5D). We hypothesized that loss of the Fyv6-interacting domain on Syf1 would phenocopy *fyv6Δ*.

We observed just this effect. Loss of the Fyv6-interacting region in Syf1 Δ778-859 and Δ634-859

mutants resulted in either modest (Δ778-859) or severe (Δ634-859) *cs* and *ts* phenotypes. In

addition, these Syf1 truncation mutants also triggered use of the alternative 3' SS in *SUS1* (**Figure**

5F). Interestingly, if the Δ778-859 and Δ817-859 Syf1 mutants are expressed in a *fyv6Δ*

background, it results in suppression of the *fyv6Δ ts* phenotype (**Figure 5G; *c.f.* 5B**). However,

the *SUS1* alternative 3' SS is still used (**Figure 5H**). This indicates that *ts* phenotype suppression

and triggering usage of the alternative 3' SS in *SUS1* are separable processes.

When considered together, these results likely originate from mutually exclusive Fyv6/Syf1

and Yju2/Syf1 interactions. It is possible that in the absence of Fyv6, Syf1 inappropriately binds

Yju2 during exon ligation. The presence of Yju2 during the 2^{nd} step could then lead to growth

defects that are suppressed when the Yju2-binding site on Syf1 is deleted. Alternatively or in

addition, the absence of Fyv6 could perturb the equilibrium between the 1^{st}- and 2^{nd}-step

conformations of the spliceosome and result in favoring of the 1^{st} step (L. Liu et al., 2007). Loss

of this equilibrium could lead to growth defects that are suppressed by destabilization of the 1^{st}-

step conformation by removing a binding site for Yju2 on Syf1 and restoring the equilibrium. In

both cases, truncation of Syf1 suppresses growth defects but not *SUS1* splicing defects since

Fyv6 function is still needed during the 2^{nd} step.

Mutations in Many Different Splicing Factors Can Suppress *fyv6Δ* Phenotypes

To gain additional insights into Fyv6 function, we carried out a genetic screen to detect

spontaneously arising suppressors that can correct temperature-dependent growth defects

(**Figure 6A**). This method has previously been used to illuminate many aspects of spliceosome

biochemistry and non-silent mutations within splicing factors are likely causative for suppression

(Montemayor et al. 2014; Kuhn and Brow, 2000; Brow, 2019).

We plated saturated *fyv6Δ* yeast cultures onto 20 different YPD plates and incubated them

at 37°C until colonies arose. Within several days, colonies of various sizes appeared and were

isolated. A subset of picked colonies were then tested for suppression of the *ts* phenotype. All of

these colonies were able to suppress growth defects at 37°C, and many also suppressed the *cs*

phenotype at 23°C (**figure 6-figure supplement 1**). We then performed whole genome

sequencing on the largest of the picked colonies as these were expected to be the strongest

suppressors. In 19 of 20 sequenced strains, we were able to identify at least one non-silent

mutation in a splicing factor (**Figure 6B**, **supplementary table 5**). When we carried out a similar

experiment aimed at identifying *cs* suppressors, we were unable to identify any likely causative

mutations in splicing factors. Therefore, we focused our analysis on the *ts* suppressor mutations.

Collectively, the identified mutations tended to fall into two categories: gain of positive

charge or substitution with a bulkier side chain. Gain of positive charge mutations were found in

Prp8 (G1868R) as well as the second step factors Slu7 (A24R, and N28K), and Prp22 (I1133R).

Structurally, amino acids within Prp8 and Prp22 are proximal to RNA within the spliceosome and

addition of charged amino acid side chains may stabilize interactions with the phosphodiester

RNA backbone. The N-terminus of Slu7 containing A24 and N28 is not resolved in any

spliceosome structure to our knowledge and normally contains many negatively charged

residues. The majority of identified suppressors involve a mutation to a bulkier side chain including

several in Prp8 (S1594F/Y, V1862L), Prp17 (A251V, G386W), Cef1 (A37V), and Clf1 (T33I). The

increase in hydrophobic surface area provided by these mutations may help to stabilize

interactions with nearby proteins, compensating for structural destabilization due to the absence

of Fyv6. Alternatively, since Prp8, Prp17, Cef1, and Clf1 are all present during 5'SS cleavage, it

is possible that these bulky mutations could destabilize a 1st step conformation of the spliceosome

and suppress phenotypes due to the loss of Fyv6 by facilitating formation of the 2nd step

conformation.

We isolated one suppressor mutation within the U2 snRNA (A25G) near where the C-

terminal domain of Prp22 inserts towards the active site and located beneath the Fyv6 binding

site. It has previously been observed that mutation of this specific U2 snRNA nucleotide is not

deleterious to growth (McPheeters and Abelson, 1992) and is therefore well-tolerated by yeast.

The mechanism of suppression by this mutation is not clear; however, it is interesting to note that

we also observed suppressor mutations in the U2 snRNP component Rse1 (**supplementary**

table 5) which is released from the spliceosome prior to catalysis. The Rse1 mutations always

occurred coincident with mutations in other splicing factors. It is possible that suppressors in

factors present prior to or after exon ligation are able to suppress (or enhance suppression of)

fyv6Δ by changing how pre-mRNAs compete for a limited set of splicing factors or by reducing

blocks to pre-spliceosome assembly caused by accumulation of stalled complexes (Mendoza-

Ochoa et al., 2019; Munding et al., 2013). Together these data indicate that mutations in many

different splicing factors including second-step factors and the U2 snRNA can be isolated from

fyv6Δ suppressor strains.

Multiple Second-step Splicing Defects can be Rescued by the Same Suppressor

We decided to study several of these mutations in greater detail and to confirm that they

were sufficient for *fyv6Δ ts* phenotype suppression. In the case of Cef1, we obtained several

mutants located near the interface between Cef1 and the 2nd step factor Prp17: A37V, M175I, and

Q193P (**Figure 6C**). This interface has only been observed in structures of C* and P-complex

spliceosomes, suggesting its importance for exon ligation. In addition, it has previously been

reported that another mutant of Cef1 A37—A37P—is able to suppress splicing defects due to a

BS adenine to guanosine substitution (BS-G) which causes stalling after 5' SS cleavage (Query

and Konarska, 2012). Therefore, we wondered if other Cef1 mutants capable of suppressing

defects due to BS-G could also suppress *fyv6Δ* (Cef1 A37P, V36R, and S48R). We created *fyv6Δ*

shuffle strains for Cef1 and tested each mutant's ability to suppress *cs* and *ts* phenotypes (**Fig.**

6D). All of these Cef1 mutants, both those isolated from our suppressor screen and those

identified as BS-G suppressors, could at least weakly suppress the *cs* phenotype at 23°C with

our isolated M175I mutant being a particularly strong suppressor. In addition, all of the mutants

except Cef1 A37V could suppress the *ts* phenotype at 37°C to varying degrees. These results

confirm that single point mutants in Cef1 can suppress *fyv6Δ* growth phenotypes and that at least

some BS-G suppressors in Cef1 also suppress *fyv6Δ*. A common mechanism could be that these

mutations stabilize the exon ligation conformation of the spliceosome to favor the 2nd step either

when a BS-G is present or Fyv6 is absent.

In the case of Prp8, we tested five different suppressor mutations in *prp8* shuffle strains

(**Figure 6E, F**). Two of these are located within the Prp8 α-finger domain (S1584F,Y; **Figure 6E**)

while the remainder are located within the RNase H-like domain β-finger (V1862L, G1868R,

T1982S; **Figure 6C, 6E**). We confirmed that four of the identified Prp8 suppressor mutations could

partially suppress either or both of the *cs* and *ts fyv6Δ* phenotypes (**Figure 6F**; S1574Y, V1862L,

G1868R, and T1982S). Even though we isolated the Prp8 S1584F mutant in our screen and this

substitution is very similar to S1584Y, we were not able observe suppression due to this mutant

in these strains. Identification of mutants in Prp8 that suppress *fyv6Δ* is not surprising since many

different alleles of Prp8 are known to modulate the conformational equilibrium between the 1st

and 2nd steps (Fica and Nagai, 2017; L. Liu et al., 2007; Query and Konarska, 2004) or other steps

in splicing. Indeed, suppressors in Prp8 have previously been identified at some of these same

positions: V1862A/Y/D and T1982A were identified as suppressors of, respectively, the *U4-cs*1

allele (Kuhn and Brow, 2000; Kuhn et al., 1999) or the 5' SS U2A mutant (Siatecka et al., 1999).

Finally, given our observations with the Cef1 and Prp8 mutants, we wondered about the

generality of 2nd step splicing factor suppression. Several dominant mutants in the 2nd step factor

Prp18 were previously shown to suppress *ts* phenotypes due to various mutations in Slu7

(Aronova et al., 2007). We wondered if these Prp18 mutants would also suppress the *ts* (or *cs*)

phenotype due to Fyv6 loss. Indeed, the Prp18 V191A suppressor of *slu7* alleles was able to

improve growth at 37°C of *fyv6Δ* yeast (**figure 6-figure supplement 2**). Together our results from

the suppressor analysis support a model in which a given 2nd step suppressor mutation may be

able to restore splicing activity lost by a variety of mechanisms.

A Prp22-Dependent Splicing Stall is Relieved by Fyv6 Deletion

The final *fyv6Δ* suppressor mutation that we studied was the Prp22 I1133R mutant. This

mutation is located within the C-terminal domain of Prp22 that enters the spliceosome active site

and is located proximal to U2 snRNA nucleotide A31 (**Figure 7A**). Fyv6 bridges over this Prp22-

U2 snRNA interaction. The I1133R mutation might stabilize Prp22/U2 snRNA contacts that can

compensate for Fyv6 loss and promote exon ligation. The Prp22 I1133R mutant is able to weakly

suppress both the *cs* and *ts* phenotypes in *fyv6Δ* yeast (**Figure 7B, C**). Given its location in the

C-terminal domain, we wondered if it or *fyv6Δ* would show synthetic genetic interactions or

epistasis with mutations in the helicase domain. We tested two helicase domain mutations, R805A

and G810A, both of which are known to impact Prp22-dependent proofreading (Mayas et al.,

2006). The Prp22 R805A mutant is by itself *cs* (Schwer and Meszaros, 2000) and showed a

synthetic lethal interaction with *fyv6Δ* (**Figure 7B, 7C**). This was only slightly rescued by

simultaneously introducing I1133R into Prp22. In contrast, Prp22 G810A mutant is viable in the

presence and absence of Fyv6 as well as in combination with the I1133R suppressor mutation

(**Figure 7B, 7C**). Prp22 G810A by itself does not suppress the *ts fyv6Δ* phenotype; however, the

Prp22 G810A/I1133R double mutant is a stronger suppressor of the *ts* phenotype than either

mutant alone. These additive genetic interactions suggest that the Prp22 *fyv6Δ* suppressor

mutation (I1133R) and the helicase domain mutations (R805A and G810A) impact distinct Prp22

functions. These functions could correspond to Prp22's role in promoting exon ligation during the

2nd step (the C-terminal domain mutation) and its role in promoting ATP-dependent mRNA release

and proofreading (the helicase domain mutations).

Recently, it was reported that the Prp22 G810A mutant could activate usage of some

alternative 3' SS that are also activated by *PRP18* deletion (Roy et al., 2023). We therefore

wondered if any of the Prp22 mutants could activate alternative 3' SS also activated by *FYV6*

deletion. We isolated RNA from these strains and analyzed *SUS1* splicing by RT-PCR (**Figure**

7D, E). None of the Prp22 mutants activated usage of the *SUS1* alternative 3' SS or enhanced

usage of this site beyond what is observed upon Fyv6 deletion. However, we noticed that the

Prp22 helicase mutants, R805A and G810A, both caused an accumulation of unspliced pre-

mRNA. This is consistent with these mutations causing a block during pre-mRNA splicing and

having a different effect on splicing than Prp22 I1133R. Surprisingly, the block due to the G810A

mutation was relieved in the *fyv6Δ* strains and little pre-mRNA accumulation was observed in the

absence of Fyv6. This result suggests that Fyv6 could act as a negative regulator of splicing of

the *SUS1* transcript when a Prp22 helicase domain mutant is present. Removing Fyv6 restores

splicing by activation of the *SUS1* BP proximal 3' SS.

To further explore the connections between Fyv6 and Prp22 fidelity, we used the ACT1-

CUP1 assay in which yeast growth in the presence of increasing concentrations of Cu^{2+} is

proportional to the extent of splicing of a reporter RNA (**Figure 7F, G**; **figure 7-figure supplement**

1, 2). Since the Prp22 R805A mutant grew poorly, we only assayed the Prp22 G810A, I1133R,

and G810A/I1133R (double) mutants in the presence and absence of Fyv6. Consistent with

previous results, we found that *fyv6Δ* is deleterious for splicing of the WT ACT1-CUP1 reporter

containing consensus splice sites. Copper tolerance was improved when *fyv6Δ* was combined

with Prp22 I1133R. Unexpectedly, Prp22 I1133R was deleterious when Fyv6 was present for this

reporter. If this Prp22 mutant and Fyv6 both promote the 2^{nd} step, then their combination may

have an additive, negative effect on splicing due to over-stabilization of this spliceosome

conformation.

Finally, we assayed strains containing a 3' SS UAG to gAG reporter since it was previously

shown that Prp22 helicase domain mutants could increase usage of nonconsensus 3' SS due to

faulty proofreading activity (**Figure 7F, G**). In agreement with this, we observed a dramatic

increase in copper tolerance when Prp22 contained both the C-terminal domain and helicase

mutants (double). This suggests that splicing of nonconsensus 3' SS splice sites is enhanced both

by stabilization of the 2^{nd} step conformation (I1133R) and perturbing mRNA release (G810A). This

effect on Prp22 is Fyv6-dependent since loss of Fyv6 results in very poor copper tolerance of

strains containing the gAG reporter, likely since this 3' SS is located far from the BP (38 nt). These

results show that *fyv6Δ* can be suppressed with a Prp22 C-terminal domain mutant and that

mutations in the Prp22 C-terminal domain can have distinct functional consequences from those

located in the helicase domain. In the case of *SUS1*, stalling of splicing due to the Prp22 G810A

mutant can be relieved by removing Fyv6 and activation of a BP proximal 3' SS.

DISCUSSION

Here, we establish Fyv6 as a 2^{nd} step splicing factor in yeast that promotes usage of BP-

distal 3' SS. Structural, genetic, and biochemical studies reveal critical contacts and interactions

between Fyv6 and the splicing machinery. From these data, we propose that Yju2/Syf1

interactions need to be disrupted for proper Fyv6 recruitment and that Fyv6 can enforce the

Prp22-dependence of at least some splicing events. Combined, our data allow us to propose a

model for Fyv6's role during the 2^{nd} step (**Figure 8**). In this model, Fyv6 is recruited to the C*

complex after Prp16-dependent release of Yju2, which exposes the binding site for Fyv6 on Syf1.

Once bound, Fyv6 (possibly in concert with Slu7 and Prp18) facilitates docking of BP distal 3' SS

into the active site. In the absence of Fyv6, the use of BP distal 3' SS becomes more difficult and

BP proximal 3' SS are favored. Under these conditions, the requirement for Prp22 during exon

ligation is reduced and Yju2 may also potentially rebind Syf1. One or both of these factors could

contribute to formation of a lower fidelity spliceosome with relaxed sequence constraints for exon

ligation.

Structural Insights into the Recruitment and Release of 2nd step Factors

493    A key principle of spliceosome structural biochemistry is the presence of mutually

494  exclusive interactions that coordinate the stepwise and directional progression of splicing

495  (Townsend et al., 2020). Our cryo-EM structure of the P complex spliceosome suggests another

496  such interaction is involved in the 1st to 2nd-step transition during which Fyv6 displaces Yju2. Yju2

498  and Fyv6 occupy overlapping binding sites on the splicing factor Syf1 and, in fact, unassigned

499  density in spliceosome P complexes due to Fyv6 was previously mis-interpreted as Yju2 due to

500  their structural similarity (Wilkinson et al., 2017; Wilkinson et al., 2021). We propose that Prp16-

501  promoted spliceosome remodeling results in release of Yju2 from the spliceosome C complex and

502  permits mutually exclusive binding of Fyv6. Fyv6 remains bound in the P complex but then may

503  dissociate to permit re-binding of Yju2, which was observed by cryo-EM to be a component of the

504  yeast ILS complex (Wan et al., 2017). It should be noted, however, that intron-lariats only weakly

505  co-immunoprecipitate with Yju2 (Y.-C. Liu et al., 2007) and that Yju2's presence in the cryo-EM

506  data for ILS complexes could originate from the complexes being purified with a TAP-tag on Yju2

507  itself (Wan et al., 2017). Since Yju2 has also not been observed in metazoan ILS complexes, its

508  role in spliceosome disassembly and the need for Fyv6-Yju2 exchange during the P to ILS

509  transition should be studied further.

510    Previously, Chiang and Cheng noted that while the N-terminal half of Yju2 was able to

511  restore 1st step activity to Yju2-depleted extracts, it is much more stably associated if a second

512  polypeptide containing the C-terminal, Syf1-interacting half was included (Chiang and Cheng,

513  2013). It is possible that competition between Yju2 and Fyv6 for Syf1 binding can explain these

514  results. If only the N-terminal half of Yju2 is present, Fyv6 could associate prematurely with Syf1

515  and result in destabilization of the 1st step conformation in favor of 2nd step interactions. This may

516  then result in weaker binding for the N-terminal half of Yju2. Failure to properly recruit and

517  exchange the Yju2/Fyv6 interaction on Syf1 may also be the origin of some of the yeast growth

518  and splicing phenotypes we observe (**Figure 5**).

The need for Yju2/Fyv6 exchange hints at two distinct functions for Fyv6: a role in Yju2

displacement to prevent its re-binding during exon ligation and a role in stabilization of the 2^{nd}

step spliceosome conformation. These two functions can potentially result in the distinct *cs* and

ts phenotypes in yeast. While more work is needed to dissect these phenotypes, it is possible that

the *cs* phenotype results from increased stability of the Yju2/Syf1 interaction at lower

temperatures and the need for Fyv6 to act as a competitor of this interaction. The *ts* phenotype

on the other hand may result from destabilization of the 2^{nd} step conformation of the spliceosome

active site due to loss of Fyv6. It is possible that loss of Fyv6 results in destabilization due to

removal of a buttress that holds the RNase H-like domain of Prp8 in place (along with Prp18),

changes in how the C-terminal domain of Prp22 interacts with the spliceosome active site,

removal of an impediment to reversal of the U2 snRNA/BS duplex conformational change that

occurs during the C to C* complex transition, or a combination of these factors.

The high quality of our cryo-EM data combined with 3D classification also provide new

insights into the requirements for 3' SS docking into the active site and potentially for how the P

complex spliceosome is disassembled into the ILS complex. It was previously speculated that the

N-terminal domain of Slu7 may be critical for 3' SS docking by reducing the entropic cost of this

process when the BP to 3' SS distance is long (Wilkinson et al., 2017). In our state II P complex

structure, the 3' SS remains docked but this domain of Slu7 is not observed. This suggests that

either Slu7 N-terminal domain interactions with Cwc22 and Prp8 are not essential for 3' SS

docking into the active site for exon ligation and/or that these interactions are not essential for

maintaining 3' SS docking within P complex once they form in C*.

In the state III P complex structure, densities for the 2^{nd} step factors Fyv6, Prp18, and Slu7

are entirely missing and the 3' SS is no longer docked into the active site. While it is possible that

this may reflect decomposition of the P complex (or another intermediate) by a pathway unrelated

to cellular spliceosome disassembly, the structure is suggestive that 3' SS undocking is coupled

to 2^{nd} step factor release. These steps may be necessary to help prepare the spliceosome for ILS

complex formation since the 3' SS is undocked in all ILS structures obtained to date (Vorländer et al., 2024; Wan et al., 2017; Yan et al., 2015; Zhang et al., 2019). Prp22 is still present in this structure, and it is possible that state III represents a conformation occurring after exon ligation but prior to Prp22-dependent mRNA release. In this case, the state III structure implies that Prp22 ATPase activity is not needed for dissociation of Fyv6, Prp18, and Slu7. It is also interesting to note that the C-terminal domain of Prp22 can no longer be resolved in the state III structure. What exactly activates the Prp22 ATPase to promote proofreading or mRNA release is not known. It is conceivable that undocking of the 3' SS from the active site can be sensed by the Prp22 C-terminal domain and this contributes to ATPase activation. This model differs from that previously proposed for Prp22-enabled 3' SS sampling in which that ATPase activity of Prp22 is needed to undock the 3' SS (Semlow et al., 2016). However, sampling could involve multiple steps: ATP-independent 3' SS undocking and ATP-dependent entry into the sampling state. Whether or not the 2nd step factors must release or alter conformation to enable sampling is unknown. Regardless of the exact mechanism, it is important to recognize that the P complex structures determined here were all obtained using a Prp22 mutant defective in mRNA release. The chemical and kinetic competencies of these states will require future investigations.

Multiple, Non-Redundant 2nd Step Factors Promote Use of BP Distal 3' SS

The challenge of 3' SS selection involves not just selection of the correct nucleotide sequence (YAG) but also identifying which YAG sequence to use. Here, we define a function for Fyv6 in facilitating use of BP distal 3' SS, especially those located $\geq$ 21 nt away from the branch. This function is similar to those of other 2nd step factors important for efficient splicing when the BP-3' SS distance is $\geq$12 nt (Slu7, Prp18) or $\geq$21 nt (Prp22) (Brys and Schwer, 1996; Schwer and Gross, 1998; Zhang and Schwer, 1997). Human Slu7 has additionally been reported to be important for preventing suppression of the correct 3' SS in favor of up- or downstream alternative 3' SS (Chua and Reed, 1999). Our previous study using competing 3' SS in the ACT1-CUP1 intron did not provide evidence for suppression of the correct 3' SS in the absence of Fyv6,

suggesting that this function is not shared between human Slu7 and yeast Fyv6. Nonetheless, it

is important to recognize that the splicing machinery employs multiple factors that enforce usage

of BP distal 3' SS. These functions are not redundant as changes in 3' SS are still observed in the

absence of Fyv6 but presence of Prp18 and vice versa (Roy et al., 2023).

Intron architecture can also contribute to 3' SS selection since canonical "AG" 3' SS are

depleted in humans in the region between the BP and annotated 3' SS (the AG exclusion zone,

or AGEZ) (Corvelo et al., 2010; Gooding et al., 2006). AG sites are also typically depleted in the

corresponding region in yeast introns (data not shown). This would suggest that annotated 3' SS

are also the most optimal sites located downstream of the BP. However, an important assumption

of this model is that non-AG 3' SS can also be effectively proofread and not used. Deletion of

either Prp18 or Fyv6 leads to activation of not just BP proximal 3' SS but also those with highly

divergent sequences from the YAG consensus (Roy et al., 2023). This suggests that an AGEZ

alone is insufficient for correct 3' SS usage and that mechanisms to ensure preferential use of

YAG sites are critical.

It is likely that Fyv6 and Prp18 enforce usage of YAG sites by common as well as factor-

specific mechanisms. Both proteins stabilize the conformation of the Prp8 RNase H-like domain

in P complex. Numerous splice site suppressor mutations map to this domain including those that

enhance splicing of non-YAG 3' SS (Galej et al., 2013). Fyv6 and Prp18 may be essential for

establishing a "high fidelity" conformation of the RNase H-like domain that enforces use of

canonical YAG 3' SS and preventing activation of cryptic sites within the AGEZ. In support of this,

we note that in our state III P complex structure that lacks both Fyv6 and Prp18 and a stably

docked 3' SS, the RNase H-like domain is also more flexible.

In addition to this possibility, Fyv6 and Prp18 may also contribute uniquely to 3' SS fidelity

and selection. The conserved loop region of Prp18 inserts into the spliceosome active site and

contacts the 3' SS. Partial deletion of this region of Prp18 phenocopies at least some of the

changes in 3' SS selection when Prp18 is deleted entirely (Crotti et al., 2007; Roy et al., 2023).

This indicates that even when the Prp8 RNaseH-like domain is still stabilized by Prp18 (and Fyv6)

disruptions to 3' SS selection can still occur. Unlike Prp18, Fyv6 does not contact the spliceosome

active site. However, it is the only 2^{nd} step factor that contacts exon ligation fidelity factor Prp22.

While these contacts are not essential for suppressing use of an alternative 3' in *SUS1*, a mutation

in Prp22 can nonetheless suppress a *fyv6Δ* phenotype and loss of Fyv6 relieves a Prp22 helicase

mutant-dependent stall in *SUS1* splicing. These observations suggest that Fyv6 and Prp22 are

functionally connected to one another.

Some 2^{nd} Step Factors May Enforce Usage of Prp22 during Exon Ligation

We were particularly struck by the similarities in the BP-3' SS distances for the requirement

of Prp22 during exon ligation and for promotion of BP distal 3' SS usage by Fyv6, both $\geq$21 nt

(Schwer and Gross, 1998). In addition, the alternative, BP proximal 3' SS that are activated when

Fyv6 is deleted show much less sequence conservation relative to the annotated 3' SS and

include sequences (GAG, BG) known to be proofread and rejected by Prp22. Together, this

suggests that one function of Fyv6 is to enforce Prp22-dependence of exon ligation by stabilizing

a spliceosome conformation that more easily enables usage of BP distal 3' SS (**Figure 8**). As

these sites require Prp22 for exon ligation, it is also likely that they are subject to Prp22-dependent

proofreading, and this may enforce accurate exon ligation at the YAG 3' SS consensus. When

Fyv6 is absent, many alternative, BP proximal 3' SS are activated. These sites could have a

relaxed requirement for Prp22 and proofreading. While the Fyv6-dependence of Prp22 occupancy

is beyond the scope of the current manuscript, loss of Prp18 (which also activates usage of non-

canonical, BP-proximal 3' SS) does result in lower occupancy of Prp22 on the *ACT1* intron

(Strittmatter et al., 2021). This suggests that Prp22 occupancy, and possible involvement in the

exon ligation step, can be influenced by 2^{nd} step factors.

There is precedent for nonessential splicing factors enforcing use of a proofreading

ATPase. Cus2 is not essential for pre-spliceosome formation (Yan et al., 1998). However, when

Cus2 is present then ATPase activity of Prp5 is essential for disruption of the Hsh155/Cus2

interaction (Perriman et al., 2003; Talkish et al., 2019). In this case, Cus2 is not essential for Prp5

proofreading of BP selection, and the connection between the ATP hydrolysis activity of Prp5,

removal of Cus2, and proofreading is unclear (Liang and Cheng, 2015; Talkish et al., 2019; Xu

and Query, 2007). It is possible that enforcement of Prp5 ATPase activity by Cus2 is important for

other aspects of pre-spliceosome formation *in vivo* and the impact of Prp5 and Cus2 on BP

selection has not yet been studied transcriptome-wide using high-resolution, sequencing-based

approaches. The C-terminal domain of the 1st step splicing factor Yju2 (the same domain that

competes with Fyv6 for Syf1 binding) is also not essential for 5' SS cleavage. In the absence of

this domain, the N-terminal domain of Yju2 is sufficient for promoting the 1st step reaction, and

the spliceosome can transition to the 2nd step and carry out exon ligation without Prp16, albeit

inefficiently (Chiang and Cheng, 2013). Importantly, addition of the C-terminal domain on a

separate polypeptide reduces Prp16-independent activity. This suggests that the Yju2 C-terminal

domain enforces the use of the Prp16 ATPase. As we propose with Fyv6 and Prp22, the C-

terminal domain of Yju2 may not just help to promote the 1st to 2nd step transition by enforcing

Prp16-dependence but could also facilitate Prp16-dependent proofreading during 5' SS cleavage

(Koodathingal et al., 2010; Semlow et al., 2016). General operating principles of the yeast splicing

machinery may not just include proofreading but also use of splicing factors that enforce usage

of the proofreading ATPases.

MATERIALS AND METHODS

Key Resources Table				
Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information

strain, strain backgr ound (*Sacch aromyc es cerevisi ae*)	Various S. cerevisia e strains			See Supplementary Table 6 for full list
recomb inant DNA reagent	Various plasmids			See Supplementary Table 7 for full list
Sequen ce-based reagent	Various oligonucl eotides			See Supplementary Table 8 for full list
antibod y	Anti-FLAG (Mouse monoclo nal)	Sigma-Aldrich	Cat #: F1804; RRID:A B_26204 4	WB (1:1000)
antibod y	Anti-actin (mouse monoclo nal)	Sigma-Aldrich	Cat #: MAB150 1; RRID:A B_22230 41	WB (1:5000)
antibod y	Anti-mouse IgG – HRP (Goat polyclon al)	Bio Rad	Cat #: 1706516 ; RRID:A B_11125 547	WB (1:8000)
peptide , recomb inant protein	Bovine Serum Albumi n	Sigma	Cat#: A7 906	

commercial assay or kit	Clarity Western ECL substrate	Bio Rad	Cat #: 1705060	
commercial assay or kit	Monarch RNA Cleanup Kit (50 ug)	New England Biolabs	Cat #: T2040	
commercial assay or kit	TURBO DNA-free kit	Thermo Fisher Scientific	Cat #: AM1907	
commercial assay or kit	TruSeq Stranded mRNA kit	Illumina	Cat #: 20020594	
commercial assay or kit	Master Pure Yeast RNA Purification kit	LGC Biosearch Technologies	Cat #: MPY03100	
commercial assay or kit	Access RT-PCR System	Promega	Cat #: A1250	
chemical compound, drug	Ni-NTA agarose	Qiagen	Cat #: 30210	
chemical compound, drug	Amylose resin	New England Biolabs	Cat #: E8022S	
chemical compo	Fluorescein-5-thiosem	Sigma-Aldrich	Cat #: 46985-100MG-F	

und, drug	icarbazide			
software, algorithm	RELION-5.0	Kimanius et al., 2024; doi: 10.1038/s41592-024-02304-8	RRID:SCR_016274	https://relion.readthedocs.io/en/latest/index.html
software, algorithm	Topaz	Bepler et al., 2019; doi: 10.1038/s41592-019-0575-8		https://cb.csail.mit.edu/topaz/
software, algorithm	Coot	Casañal et al., 2020; doi: 10.1002/pro.3791	RRID:SCR_014222	https://www2.mrc-lmb.cam.ac.uk/personal/pemsley/coot/
software, algorithm	ISOLDE	Croll, 2018; doi: 10.1107/S2059798318002425	RRID:SCR_025577	https://tristanic.github.io/isolde/
software, algorithm	PHENIX	Liebschner et al., 2019; doi: 10.1107/S2059798319011471	RRID:SCR_014224	https://phenix-online.org/
software, algorithm	UCSF ChimeraX	Petterson et al., 2021; doi: 10.1002/pro.3943	RRID:SCR_015872	http://www.cgl.ucsf.edu/chimerax/
software, algorithm	R Project for Statistical Computing	R Foundation for Statistical Computing	RRID:SCR_001905	https://www.r-project.org/
software, algorithm	FASTQC	Andrews, 2010	RRID:SCR_014583	http://www.bioinformatics.babraham.ac.uk/projects/fastqc/
software,	Fastp	Chen et al., 2018; doi:10.1093/bioi	RRID:SCR_016962	https://github.com/OpenGene/fastp

algorithm		nformatics/bty560		
software, algorithm	STAR	Dobin et al., 2013; doi: 10.1093/bioinformatics/bts635	RRID:SCR_004463	https://github.com/alexdobin/STAR/releases
software, algorithm	SAMtools	Li et al., 2009; doi: 10.1093/bioinformatics/btp352	RRID:SCR_002105	https://www.htslib.org/
software, algorithm	Salmon	Patro et al., 2017; doi: 10.1038/nmeth.4197	RRID:SCR_017036	https://combine-lab.github.io/salmon/
software, algorithm	DeSeq2	Love et al., 2014; doi: 10.1186/s13059-014-0550-8	RRID:SCR_015687	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
software, algorithm	featureCounts	Liao et al., 2014; doi: 10.1093/bioinformatics/btt656	RRID:SCR_012919	https://bioconductor.org/packages/release/bioc/html/Rsubread.html
software, algorithm	SpliceWiz	Wong et al., 2024; doi: https://doi.org/10.1093/bib/bbad468		https://github.com/alexchwong/SpliceWiz
software, algorithm	ImageJ 1.53v	Schneider et al., 2012; doi:10.1038/nmeth.2089	RRID:SCR_003070	https://imagej.net/ij/
other	Cryo-EM grid	Quantifoil	Cat #: Q350CR2-2NM	
other	NuPAGE Bis-Tris mini protein gel	Invitrogen	Cat #: NP0322	

other	NuPAGE™ MES SDS Running Buffer	Invitrogen	Cat #: NP0002	
other	Standard Vitrobot Filter Paper, Ø55/20 mm, Grade 595	Ted Pella	Cat #: 47000-100	
other	Mixed cellulose ester filter paper	Whatman	WHA7140104	
other	TRIzol	Thermo Fisher Scientific	Cat #: 15596018	
other	SuperScript III reverse transcriptase	Thermo Fisher Scientific	Cat #: 18080044	
other	Zymolyase	Zymo Research	Cat #: E1005	Used at 50 U/mL

Yeast growth and culture

Yeast transformation, plasmid shuffling/5-FOA selection, and growth were carried out using standard techniques and media (Treco and Lundblad 1993; Sikorski and Boeke 1991).

Cryo-electron microscopy

Purification of dominant-negative Prp22

Yeast Prp22 with a dominant-negative S635A mutation was cloned into pRS424 and

pRS426 vectors after an N-terminal CBP-His-TEV tag (Galej et al., 2013). The protein was

expressed in 24 L of BCY123 yeast cells essentially as described (Galej et al., 2013).

Recombinant Prp22 was then purified first by Calmodulin-Sepharose in a buffer containing 50

mM Tris-HCl pH 8.5, 350 mM NaCl, 2 mM $CaCl_2$, 1 mM Mg acetate, 1 mM imidazole, 0.1% NP-

40 substitute, 5 mM β-mercaptoethanol and eluted with 3 mM EGTA. The protein was dialyzed

into 20 mM Tris-HCl pH 7.4, 350 mM NaCl, 10 mM imidazole, 0.1% NP-40 substitute and further

purified over Ni-NTA resin at 500 mM NaCl and eluted with 200 mM imidazole. The eluted protein

was treated with His-tagged TEV protease during dialysis against 20 mM HEPES-KOH pH 7.9,

300 mM KCl, 0.2 mM EDTA, 20% glycerol, 0.5 mM DTT, before removal of the protease with Ni-

NTA resin. The final protein was frozen in liquid nitrogen and stored at -80 °C.

Preparation and purification of P complex for cryo-EM

P complex was purified as described in (Wilkinson et al., 2017). Brr2-TAPS yeast were

grown in a 120 L fermenter and splicing extract was prepared by the liquid nitrogen method

essentially as previously described (Lin et al., 1985). A pre-mRNA substrate consisting of 20 nt

5'-exon, 95 nt intron, and 32 nt 3'-exon from the UBC4 pre-mRNA followed by 3xMS2 stem loops

was generated by run-off in vitro transcription from an EcoRI-linearized plasmid template. The

RNA product was labelled at the 3' end with fluorescein-5-thiosemicarbazide (Wu et al., 1996). A

180 mL in vitro splicing reaction was performed containing 60 mM potassium phosphate pH 7, 2

mM ATP, 2.5 mM $MgCl_2$, 3% w/v PEG8000, 3 nM pre-mRNA, 37.5 nM MS2-MBP protein, and

40% (v/v) splicing extract treated with dominant negative Prp22 S635A. Reactions were incubated

for 30 min at 23°C, then incubated for 20 min with 5 mM of a DNA oligonucleotide complementary

to the 3'-exon (oligo sequence 5'-ATGAAGTAGGTGGAT-3') to induce cleavage of the 3'-MS2 tag

by the endogenous RNase H activity of the splicing extract. The reaction mixture was centrifuged

through a 40% glycerol cushion in buffer A (20 mM HEPES, pH 7.9, 75 mM KCl, 0.25 mM EDTA).

The cushion was collected and applied to amylose resin in the presence of 0.025% NP-40

substitute. After overnight incubation at 4°C the resin was washed and eluted with buffer A

containing 5% glycerol, 0.01% NP-40 substitute and 12 mM maltose. Fractions containing

spliceosomes were concentrated to 0.1 mg/mL then dialyzed against buffer A for 3 h.

For cryo-EM grid preparation, a freshly glow-discharged Cu300 R2/2 holey carbon grid

with a 2-nm layer of amorphous carbon (Quantifoil) was mounted in the chamber of a Vitrobot

Mark IV (Thermo Fisher Scientific) maintained at 4°C and 100% humidity. Spliceosomes (3 μL)

were applied and after 60 s was manually blotted using Ø55 grade 595 filter paper (Ted Pella),

followed by application of another 3 μL of spliceosomes, waiting 60 s, before blotting again and

plunging into liquid ethane.

Cryo-EM data collection

Cryo-EM data were collected as two datasets from two grids made from the same sample.

Dataset 1 was acquired using a Thermo Scientific Titan Krios cryo-TEM (LMB Krios 1) and dataset

2 was acquired using a Thermo Scientific Titan Krios G3i cryo-TEM (LMB Krios 3). Both used a

K3 direct detector (Gatan) operated in super-resolution mode with 2-fold binning, and an energy

filter with slit width of 20 eV. Micrographs were collected automatically using EPU in AFIS mode,

yielding a total of 51,113 movies at 81,000x or 130,000x magnification with a real nominal pixel

size of 0.93 Å or 0.669 Å, with defocus ranging from -1.3 μm to -3.1 μm and a total fluence per

micrograph of 40 e^{-}/Å^{2}, fractionated into 40 frames.

Cryo-EM data processing

All cryo-EM data were processed using RELION-5.0 (Kimanius et al., 2024). Both datasets

were initially processed separately. Movies were corrected for motion using the RELION

implementation of MotionCor2, with 4x4 patches and dose-weighting. CTF parameters were

estimated using CTFFIND-4.1. Particle picking was done using Topaz with the general model

(Bepler et al., 2019), yielding 1,820,457 particles. 3D classification was performed on 4-fold

(dataset 1) or 5-fold (dataset 2) binned particles using a P-complex reference map (EMD-10140),

and 299,741 (dataset 1) or 599,123 (dataset 2) were selected and refined to 2.99 Å or 2.78 Å

resolution respectively. After Bayesian Polishing and CTF refinement (per-particle defocus,

anisotropic magnification, beam tilt, trefoil, and 4^{th} order aberrations) the resolution improved to

2.67 Å or 2.32 Å respectively. To improve the general density quality, 3D classification without

alignment was performed with a soft mask around the whole complex with T=15. Particles with

good density, without selecting for Fyv6 occupancy, were selected (151,443 and 257,501 for

datasets 1 and 2) and refined to 2.66 Å or 2.27 Å respectively. The particles were then merged,

refined, and then refined for anisotropic magnification. The overall scaling factors from the

"anisotropic" magnification matrix was then used to calculate the corrected pixel size for dataset

1 as 0.94057 Å, keeping dataset 2 constant at 0.669 Å per pixel. Defocus values for dataset 1

were then scaled by the square of 0.94057 / 0.93, with an empirical constant correction of -10.06

Å (Wilkinson et al., 2019). Finally, another round of per-particle defocus refinement and

anisotropic magnification refinement was performed to reduce errors in defocus correction and

dataset scaling. The combined particles refined to 2.24 Å resolution. Resolution is reported using

the gold-standard Fourier Shell Correlation with 0.143 cutoff.

The refined particles after merging were subjected to 3D classification without alignment,

with a soft mask around the entire spliceosome, with T=15. The resulting four classes were used

to define States I, II and III. To improve the densities for the periphery of the spliceosome that

displayed local flexibility, focused refinements were used with Blush regularization in RELION-5

to reduce overfitting (Kimanius et al., 2024). The Prp22 density was improved using a soft-mask

just around the Prp22 helicase domain for refinement of particles from States I and II, yielding a

2.96 Å map. Densities for the U2 snRNP, NTC, U5 Sm ring, and Cwc22 N-terminal domain were

each improved first by 3D classification without alignment of particles from all three states,

selecting particles with strong density (72,869 for the U2 snRNP, 260,409 for the NTC, 113,561

for the U5 Sm ring, 201,984 for Cwc22-NTD), performing signal subtraction to remove density for

the rest of the spliceosome (recentering on the mask center of mass and re-boxing to 300 pixels,

or 192 pixels for Cwc22-NTD), and finally 3D refinement starting with 1.8 degree local angular

searches, giving 3.42 Å for the U2 snRNP, 3.72 Å for the NTC, 3.06 Å for the U5 Sm, and 3.57 Å

for Cwc22-NTD. In all cases, control refinements without Blush regularization did not produce

interpretable maps (**figure 3-figure supplement 4).**

<u>Model building</u>

The model was built using Coot (Casañal et al., 2020) and ISOLDE (Croll, 2018) starting

with PDB 6EXN (Wilkinson et al., 2017) as an initial model for the core and PDB 7B9V (Wilkinson

et al., 2021) as an initial model for the periphery. The core was improved by manual fixing in Coot,

while the peripheral U2 snRNP and NTC were improved by applying torsion and distance

restraints to reference models produced by AlphaFold2. The model was refined using PHENIX

real_space_refine (Liebschner et al., 2019), just performing one macro-cycle. Figures were

generated using UCSF ChimeraX.

Plasmid cloning

Fyv6 plasmids were made by PCR amplification from gDNA of the *FYV6* coding sequence

plus ~250 nt upstream and downstream, restriction digest with NotI and SalI, and ligation into

pRS414 or pRS416. An N-terminal 1x FLAG tag was added onto Fyv6 in the plasmids by PCR

mutagenesis. Fyv6 truncations were cloned by Genewiz (Azenta Life Sciences) from pAAH1572.

Restriction digest and ligation were used to place Fyv6 constructs into the pRS413 backbone.

Syf1 plasmids were made by PCR amplification from gDNA of the SYF1 coding sequence

plus ~275 nt upstream and downstream, restriction digest, and ligation into pRS414 or pRS416.

Cef1 mutant plasmids, Prp22 I1133R mutant plasmid, Prp8 mutant plasmids, ACT1-CUP1

BP-3' SS distance reporters, and Syf1 truncation plasmids were prepared by PCR-based site

directed mutagenesis of pAAH1611, pAAH1042, pAAH1440, pAAH0470, and pAAH1625

respectively. All plasmids were fully sequenced to confirm their nucleotide sequence.

Yeast strain creation

The *DBR1* and *FYV6* genes were deleted by replacement with a nourseothricin resistance cassette (NatMX4) or hygromycin resistance cassette (hphMX4), respectively, through homologous recombination (see **supplementary table 7**; Goldstein and McCusker, 1999). The *SYF1* and *CEF1* genes were deleted by replacement with a kanamycin resistance cassette (KanMX) through homologous recombination in strains transformed with pRS416-Syf1 or pRS316-Cef1 respectively. Gene deletion was confirmed by colony PCR and/or genomic DNA extraction from the strains and PCR amplification of the corresponding genomic locus.

RNA isolation

Yeast were grown in YPD media until OD_{600} 0.5-0.8. For RNA-sequencing of temperature-shifted samples, cultures were shifted to either 37°C or 16°C or remained at 30°C and grown for 1 hr. A 15 mL volume of culture was harvested via vacuum filtration with mixed cellulose ester filter paper (Whatman, 47 mm diameter) in a Buchner funnel. Filter papers were separated into 5 mL tubes and flash frozen. Filters were stored at –80°C until RNA isolation. Filters were washed with 1 mL TRIzol to collect cells. For all other experiments involving RNA isolation, 10 OD_{600} units of OD_{600} 0.5-0.8 yeast were harvested via centrifugation.

Cell walls were disrupted by incubation with Zymolyase (50 U/mL, Zymo Research) in Y1 buffer (1 M sorbitol, 100 mM EDTA pH 8.0, 13 mM β-mercaptoethanol) (primer extension, RNA-seq of non-temperature-shifted samples) or vortexing with silica disruption beads in TRIzol (RT-PCR, RNA-seq of temperature-shifted samples). Total RNA was then isolated using TRIzol reagent (Thermo Fisher Scientific) according to manufacturer's instructions. Samples were cleaned using a 50 µg Monarch RNA Cleanup Kit (New England Biolabs) and treated with TURBO DNase (Thermo Fisher Scientific) to remove residual contaminating DNA. For RNA-sequencing, quality of RNA samples was assessed prior to library preparation by NanoDrop (concentration, A_{260}/A_{280}, and A_{260}/A_{230}), Qubit (RNA High Sensitivity and RNA IQ), and TapeStation (RNA

775 ScreenTape Analysis). High quality samples showing minimal rRNA degradation with RIN >7 and

776 A260/A280 ratios of ~2.0 were selected for library preparation.

Library preparation and RNA sequencing

778 Library preparation was conducted by the University of Wisconsin-Madison Biotechnology

779 Center Gene Expression Center. mRNA was selected from total RNA samples through polyA

780 enrichment with the TruSeq Stranded mRNA kit (Illumina). Sequencing was performed with an

781 NovaSeq6000 sequencer (Illumina).

Bioinformatic analysis of RNA-seq datasets

FASTQC was used for quality control of reads pre- and post- trimming (Andrews, 2010).

Trimming of reads was accomplished using fastp (Chen et al., 2018) prior to mapping to the

SacCer3 genome (Ensembl, R64-1-1) with STAR (Dobin et al., 2013). All samples were aligned

with STAR in a first pass. Novel junctions from all samples (WT and *fyv6Δ*) were combined and

filtered for likely false-positive junctions (see Supplemental Methods). STAR was run in a second

pass with the additional input of the filtered set of novel junctions. Indexing of bam files was

accomplished with SAMtools (Li et al., 2009). Quantitation of read counts aligned to the SacCer3

transcriptome was accomplished with Salmon (Patro et al., 2017). Differential gene expression

analysis was conducted with DeSeq2 (Love et al., 2014). Junction reads within genes were

counted with featureCounts (Liao et al., 2014) and used to calculate fraction of Annotation Splicing

(FAnS) based on methods in Roy et al., 2023 (see Supplemental Methods). Canonical branch

points were used as defined in the Ares Intron Database (Grate and Ares, 2002). Differential

splicing analysis was conducted with SpliceWiz (Wong et al., 2024).

Primer extension

RNA was isolated using the protocol above. IR700 dye conjugated probes (Integrated DNA

Technologies) were used for primer extension with SuperScript III (Thermo Fisher) of the ACT1-

CUP1 reporters (2 pmol yAC6: /5IRD700/GGCACTCATGACCTTC) and U6 snRNA (0.4 pmol

yU6: /5IRD700/GAACTGCTGATCATGTCTG) (Carrocci et al., 2017; van der Feltz et al., 2021).

Primer extension products were visualized on a 7% (w/v) denaturing polyacrylamide gel run at 35

W for 80 min at RT. Gels were imaged with an Amersham Typhoon NIR laser scanner (Cytiva),

and band intensities were quantified with Image J (version 1.53v, 2022).

RT-PCR

Yeast cultures were inoculated from cultures grown to stationary phase overnight and

grown until OD_{600} = 0.7-0.9. RNA was isolated and depleted of contaminating DNA using the

MasterPure Yeast RNA Purification Kit (LGC Biosearch Technologies) protocol as previously

described (Carrocci et al., 2017) (RT-PCR shown in Fig. 4C) or using the RNA isolation protocol

above (all others).

RT-PCR reactions were set up using the Access RT-PCR system (Promega Corporation)

following the kit procedure with 100 ng of input RNA per 25 µl reaction. Primers to amplify *SUS1*

were SUS1-exon1 5'-TGGATACTGCGCAATTAAAGAGTC-3' and SUS1-exon3 5'-

TCATTGTGTATCTACAATCTCTTCAAG-3'. Reaction products were separated on 2% (w/v)

agarose-TBE gels and imaged using ethidium bromide fluorescence.

Selection of *fyv6Δ* suppressors

Strain yAAH3353 was previously described (Lipinski et al. 2023). Twenty single colonies

of yAAH3353 were selected from a YPD plate and used to inoculate 20 x 5mL YPD liquid culture.

Cultures were grown for 48 h at 30°C. A 1 mL volume of culture was spun down, resuspended in

200 µL YPD liquid medium, and spread on a YPD plate containing 40 mg/L adenine hemisulfate

(to prevent ade2 reversion). Duplicate plates for each overnight were placed in incubators at either

18°C or 37°C. Many colonies of various sizes appeared on each plate by 9 days for 37°C and 12

days for 18°C. Colonies were picked after that time and annotated first by growth temperature,

then by overnight number, and finally roughly by colony size with 01 generally being the largest

colony on a plate (ex: 370101 from a plate grown at 37°C inoculated from overnight 01 and the

largest colony selected from the plate). Picked colonies were grown overnight in 5mL YPD at 30°C

and frozen stocks were made. Anywhere from 0 to ~15 colonies were picked per plate, covering

the variety of colony sizes that arose.

Sequencing of yeast genomic DNA

829   Yeast were grown to saturated stationary phase in 5mL YPD and ~10 OD units were sent

830   as a frozen yeast pellet to Azenta or SeqCoast for genomic DNA extraction, library, preparation,

831   and sequencing with 30x/400 mbp coverage.

Bioinformatic analysis of WGS data

Azenta and SeqCoast utilized the SacCer3 genome (Ensembl, R64-1-1) for variant calling

and provided a list of >6,000 SNPs. Since the strain is not a perfect match for this reference, non-

unique SNPs were filtered from each dataset by taking the union of two sets with each set

obtained from colonies isolated on different plates (to reduce the chance of computationally

eliminating the same variant SNP). Filtering reduced the dataset to <100 unique variants per

colony. Few unique SNPs were within protein- or snRNA-coding genes. Of SNPs within protein-

coding regions, single amino acid changes were frequently found within splicing factors (see

Supplemental Table S4). Datasets were reexamined upon identification of suppressor mutations

for suppressor mutation duplication that were potentially filtered out during comparison of SNP

datasets.

Yeast temperature growth assays

844   Yeast were grown to stationary phase overnight in the appropriate liquid medium indicated

845   in the figure legends. The cultures were first diluted to OD_{600} 0.5, serially diluted 1:10, 1:100, and

846   1:1000, and stamped onto agar plates containing the appropriate growth medium. The plates

847   were incubated at 16°, 23°, 30°, and 37°C, and plates were imaged after the number of days

848   indicated in each figure.

ACT1-CUP1 copper tolerance assays

Yeast strains expressing ACT1-CUP1 reporters were grown to stationary phase in -Leu

DO media, diluted to OD_{600} = 0.5 in 10% (v/v) glycerol, and spotted onto -Leu DO plates containing

0 to 2.5 mM CuSO₄ (Lesser and Guthrie 1993; Carrocci et al., 2018). Plates were scored and

imaged after 48 h of growth at 30°C for WT strains and after 72 h of growth at 30°C for *fyv6Δ*

strains due to differential growth between strains.

Western blotting

Total soluble protein was isolated by TCA precipitation from 10 ODs of an overnight yeast

culture. Proteins were separated with a NuPAGE 4 to 12% Bis-Tris pre-cast gel (Invitrogen) using

1X NuPAGE MES running buffer (Invitrogen) and transferred to a PVDF membrane (0.2 μm pore

size) at 30V for 120 min at 4°C using 1x NuPAGE Transfer Buffer (Invitrogen) with 20% (v/v)

methanol in a XCell II Blot Module (Invitrogen). The membrane was blocked with 3% (w/v) BSA

(Sigma, A7906) in 1x TBST for 1h at room temperature with gentle shaking. All antibodies were

diluted in 1x TBST with 3% (w/v) BSA. The primary antibodies anti-FLAG (Millipore Sigma, F1804;

diluted 1:1000) and anti-actin (Sigma Aldrich, MAB1501; diluted 1:5000) were incubated overnight

at 4°C. The membrane was then washed with 1x TBST before incubation with goat anti-mouse

IgG HRP conjugate secondary antibody (Biorad, #1706516; diluted 1:8000) for 2h at room

temperature. The membrane was then washed with 1x TBST before developing with Clarity

Western ECL substrate (Biorad, #1705060) with a 20s exposure for anti-actin and a 100s

exposure for anti-FLAG.

869
ACKNOWLEDGEMENTS

We thank David Brow for his advice on the suppressor screen. We thank Justin Mabin for

his advice on RNA-sequencing data analysis. MEW is grateful to Feng Zhang for funding and

support, and to the labs of Kiyoshi Nagai and Kelly Nguyen for reagents and support. The authors

utilized the University of Wisconsin – Madison Biotechnology Center Gene Expression Center

(Research Resource Identifier – RRID:SCR_017757) for preparation of NGS libraries for RNA-

sequencing and the Biochemistry Computational Research Facility (BCRF) for analysis of RNA-

sequencing datasets. This work was supported by grants from the National Institutes of Health

(R35 GM136261 to AAH), NIH Biotechnology Training Program (T32 GM135066) and National

Science Foundation Graduate Research Fellowship Program (Grant No. DGE-1747503)

fellowships to KAS, a Helen Hay Whitney Postdoctoral Fellowship (to MEW). This study was

supported by the MRC Laboratory of Molecular Biology electron microscopy facility. We thank

Guillaume Chanfreau for advice and Florian Heyd and Tucker Carrocci for critical reading of the

manuscript.

CONTRIBUTIONS

AAH, KAS, KAL, and MEW conceptualized the project; MEW collected and analyzed the cryo-EM

data; KAL and NJZ prepared RNA for sequencing; KAL, NJZ, and AFG analyzed the RNA-

sequencing data; AAH, KAS, NJZ, and KAL created yeast strains; KAS, KAL, NJZ, and AFG

performed genetic and biochemical assays and analyzed data; KAS, KAL, MEW, and AAH wrote

the manuscript.

DATA AND MATERIALS AVAILABILITY

RNA-seq data has been uploaded to NCBI SRA with BioProject ID PRJNA1113593. R code for

calculation of FAnS values can be found at the following DOI: 10.5281/zenodo.13798743. The

cryo-EM maps and atomic model of the P complex (state I) have been deposited in the Electron

Microscopy Data Bank (https://www.ebi.ac.uk/pdbe/emdb/) and in the Protein Data Bank

(https://www.rcsb.org). The data are available under the following accession codes: P complex

state I (EMD-47157, EMD-47160, EMD-47161, EMD-47162, EMD47163, EMD-47164; PDB

9DTR), state II (EMD-47158), and state III (EMD-47159). Raw cryo-EM data is available from

EMPIAR (EMPIAR-12330).

COMPETING INTEREST STATEMENT

907
REFERENCES

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data* [Computer
software]. http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Aronova, A., Bacíková, D., Crotti, L. B., Horowitz, D. S., & Schwer, B. (2007). Functional
interactions between Prp8, Prp18, Slu7, and U5 snRNA during the second step of pre-
mRNA splicing. *RNA*, *13*(9), 1437–1444. https://doi.org/10.1261/rna.572807

Bai, R., Yan, C., Wan, R., Lei, J., & Shi, Y. (2017). Structure of the Post-catalytic Spliceosome
from Saccharomyces cerevisiae. *Cell*, *171*(7), 1589-1598.e8.
https://doi.org/10.1016/j.cell.2017.10.038

Bepler, T., Morin, A., Rapp, M., Brasch, J., Shapiro, L., Noble, A. J., & Berger, B. (2019).
Positive-unlabeled convolutional neural networks for particle picking in cryo-electron
micrographs. *Nature Methods*, *16*(11), Article 11. https://doi.org/10.1038/s41592-019-
0575-8

Brow, D. A. (2019). An Allosteric Network for Spliceosome Activation Revealed by High-
Throughput Suppressor Analysis in Saccharomyces cerevisiae. *Genetics*, *212*(1), 111–
124. https://doi.org/10.1534/genetics.119.301922

Brys, A., & Schwer, B. (1996). Requirement for SLU7 in yeast pre-mRNA splicing is dictated by
the distance between the branchpoint and the 3' splice site. *RNA*, *2*(7), 707–717.

Carrocci, T. J., Paulson, J. C., & Hoskins, A. A. (2018). Functional analysis of Hsh155/SF3b1
interactions with the U2 snRNA/branch site duplex. *RNA*, *24*(8), 1028–1040.
https://doi.org/10.1261/rna.065664.118

Carrocci, T. J., Zoerner, D. M., Paulson, J. C., & Hoskins, A. A. (2017). SF3b1 mutations
associated with myelodysplastic syndromes alter the fidelity of branchsite selection in
yeast. *Nucleic Acids Research*, *45*(8), 4837–4852. https://doi.org/10.1093/nar/gkw1349

Casañal, A., Lohkamp, B., & Emsley, P. (2020). Current developments in Coot for
macromolecular model building of Electron Cryo-microscopy and Crystallographic Data.
Protein Science, *29*(4), 1055–1064. https://doi.org/10.1002/pro.3791

Chen, C.-H., Tsai, W.-Y., Chen, H.-R., Wang, C.-H., & Cheng, S.-C. (2001). Identification and
Characterization of Two Novel Components of The Prp19p-associated Complex, Ntc30p
and Ntc20p *. *Journal of Biological Chemistry*, *276*(1), 488–494.
https://doi.org/10.1074/jbc.M006958200

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ
preprocessor. *Bioinformatics*, *34*(17), i884–i890.
https://doi.org/10.1093/bioinformatics/bty560

Chiang, T.-W., & Cheng, S.-C. (2013). A Weak Spliceosome-Binding Domain of Yju2 Functions
in the First Step and Bypasses Prp16 in the Second Step of Splicing. *Molecular and*
Cellular Biology, *33*(9), 1746–1755. https://doi.org/10.1128/MCB.00035-13

Chua, K., & Reed, R. (1999). The RNA splicing factor hSlu7 is required for correct 3′ splice-site
choice. *Nature*, *402*(6758), 207–210. https://doi.org/10.1038/46086

Chung, C.-S., Wai, H. L., Kao, C.-Y., & Cheng, S.-C. (2023). An ATP-independent role for Prp16
in promoting aberrant splicing. *Nucleic Acids Research*, *51*(20), 10815–10828.
https://doi.org/10.1093/nar/gkad861

Chung, S., McLean, M. R., & Rymond, B. C. (1999). Yeast ortholog of the Drosophila crooked
neck protein promotes spliceosome assembly through stable U4/U6.U5 snRNP addition.
RNA, *5*(8), 1042–1054.

Company, M., Arenas, J., & Abelson, J. (1991). Requirement of the RNA helicase-like protein
PRP22 for release of messenger RNA from spliceosomes. *Nature*, *349*(6309), 487–493.
https://doi.org/10.1038/349487a0

Corvelo, A., Hallegger, M., Smith, C. W. J., & Eyras, E. (2010). Genome-Wide Association
between Branch Point Properties and Alternative Splicing. *PLOS Computational Biology*,
6(11), e1001016. https://doi.org/10.1371/journal.pcbi.1001016

Croll, T. I. (2018). ISOLDE: A physically realistic environment for model building into low-
resolution electron-density maps. *Acta Crystallographica Section D: Structural Biology*,
74(6), 519–530. https://doi.org/10.1107/S2059798318002425

Crotti, L. B., Bačíková, D., & Horowitz, D. S. (2007). The Prp18 protein stabilizes the interaction
of both exons with the U5 snRNA during the second step of pre-mRNA splicing. *Genes &*
Development, *21*(10), 1204–1216. https://doi.org/10.1101/gad.1538207

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson,
M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*,
29(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Felisberto-Rodrigues, C., Thomas, J. C., McAndrew, C., Le Bihan, Y.-V., Burke, R., Workman,
P., & van Montfort, R. L. M. (2019). Structural and functional characterisation of human
RNA helicase DHX8 provides insights into the mechanism of RNA-stimulated ADP
release. *Biochemical Journal*, *476*(18), 2521–2543.
https://doi.org/10.1042/BCJ20190383

Feltz, C. van der, Nikolai, B., Schneider, C., Paulson, J. C., Fu, X., & Hoskins, A. A. (2021).
Saccharomyces cerevisiae Ecm2 modulates the catalytic steps of pre-mRNA splicing.
RNA, *27*(5), 591–603. https://doi.org/10.1261/rna.077727.120

Fica, S. M., & Nagai, K. (2017). Cryo-electron microscopy snapshots of the spliceosome:
Structural insights into a dynamic ribonucleoprotein machine. *Nature Structural &*
Molecular Biology, *24*(10), Article 10. https://doi.org/10.1038/nsmb.3463

Fica, S. M., Oubridge, C., Wilkinson, M. E., Newman, A. J., & Nagai, K. (2019). A human
postcatalytic spliceosome structure reveals essential roles of metazoan factors for exon
ligation. *Science*, *363*(6428), 710–714. https://doi.org/10.1126/science.aaw5569

Frank, D., & Guthrie, C. (1992). An essential splicing factor, SLU7, mediates 3' splice site choice
in yeast. *Genes & Development*, *6*(11), 2112–2124.
https://doi.org/10.1101/gad.6.11.2112

Galej, W. P., Oubridge, C., Newman, A. J., & Nagai, K. (2013). Crystal structure of Prp8 reveals active site cavity of the spliceosome. *Nature*, *493*(7434), 638–643. https://doi.org/10.1038/nature11843

Gautam, A., Grainger, R. J., Vilardell, J., Barrass, J. D., & Beggs, J. D. (2015). Cwc21p promotes the second step conformation of the spliceosome and modulates 3′ splice site selection. *Nucleic Acids Research*, *43*(6), 3309–3317. https://doi.org/10.1093/nar/gkv159

Goldstein, A. L., & McCusker, J. H. (1999). Three new dominant drug resistance cassettes for gene disruption in Saccharomyces cerevisiae. *Yeast*, *15*(14), 1541–1553. https://doi.org/10.1002/(SICI)1097-0061(199910)15:14<1541::AID-YEA476>3.0.CO;2-K

Gooding, C., Clark, F., Wollerton, M. C., Grellscheid, S.-N., Groom, H., & Smith, C. W. (2006). A class of human exons with predicted distant branch points revealed by analysis of AG dinucleotide exclusion zones. *Genome Biology*, *7*(1), R1. https://doi.org/10.1186/gb-2006-7-1-r1

Grate, L., & Ares, M. (2002). Searching yeast intron data at ares lab web site. In C. Guthrie & G. R. Fink (Eds.), *Methods in Enzymology* (Vol. 350, pp. 380–392). Academic Press. https://doi.org/10.1016/S0076-6879(02)50975-7

Hossain, M. A., Claggett, J. M., Nguyen, T., & Johnson, T. L. (2009). The cap binding complex influences H2B ubiquitination by facilitating splicing of the SUS1 pre-mRNA. *RNA*, *15*(8), 1515–1527. https://doi.org/10.1261/rna.1540409

Kawashima, T., Douglass, S., Gabunilas, J., Pellegrini, M., & Chanfreau, G. F. (2014). Widespread Use of Non-productive Alternative Splice Sites in Saccharomyces cerevisiae. *PLOS Genetics*, *10*(4), e1004249. https://doi.org/10.1371/journal.pgen.1004249

Kimanius, D., Jamali, K., Wilkinson, M. E., Lövestam, S., Velazhahan, V., Nakane, T., & Scheres, S. H. W. (2024). Data-driven regularization lowers the size barrier of cryo-EM structure determination. *Nature Methods*, *21*(7), 1216–1221. https://doi.org/10.1038/s41592-024-02304-8

Koodathingal, P., Novak, T., Piccirilli, J. A., & Staley, J. P. (2010). The DEAH Box ATPases Prp16 and Prp43 Cooperate to Proofread 5′ Splice Site Cleavage during Pre-mRNA Splicing. *Molecular Cell*, *39*(3), 385–395. https://doi.org/10.1016/j.molcel.2010.07.014

Kuhn, A. N., & Brow, D. A. (2000). Suppressors of a Cold-Sensitive Mutation in Yeast U4 RNA Define Five Domains in the Splicing Factor Prp8 That Influence Spliceosome Activation. *Genetics*, *155*(4), 1667–1682. https://doi.org/10.1093/genetics/155.4.1667

Kuhn, A. N., Li, Z., & Brow, D. A. (1999). Splicing Factor Prp8 Governs U4/U6 RNA Unwinding during Activation of the Spliceosome. *Molecular Cell*, *3*(1), 65–75. https://doi.org/10.1016/S1097-2765(00)80175-6

Lesser, C. F., & Guthrie, C. (1993). Mutational analysis of pre-mRNA splicing in Saccharomyces cerevisiae using a sensitive new reporter gene, CUP1. *Genetics*, *133*(4), 851–863. https://doi.org/10.1093/genetics/133.4.851

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Liang, W.-W., & Cheng, S.-C. (2015). A novel mechanism for Prp5 function in prespliceosome formation and proofreading the branch site sequence. *Genes & Development*, *29*(1), 81–93. https://doi.org/10.1101/gad.253708.114

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: An efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–930. https://doi.org/10.1093/bioinformatics/btt656

Liebschner, D., Afonine, P. V., Baker, M. L., Bunkóczi, G., Chen, V. B., Croll, T. I., Hintze, B., Hung, L. W., Jain, S., McCoy, A. J., Moriarty, N. W., Oeffner, R. D., Poon, B. K., Prisant, M. G., Read, R. J., Richardson, J. S., Richardson, D. C., Sammito, M. D., Sobolev, O. V., … Adams, P. D. (2019). Macromolecular structure determination using X-rays, neutrons and electrons: Recent developments in Phenix. *Acta Crystallographica. Section D, Structural Biology*, *75*(Pt 10), 861–877. https://doi.org/10.1107/S2059798319011471

Lin, R. J., Newman, A. J., Cheng, S. C., & Abelson, J. (1985). Yeast mRNA splicing in vitro. *Journal of Biological Chemistry*, *260*(27), 14780–14792. https://doi.org/10.1016/S0021-9258(17)38641-6

Lipinski, K. A., Senn, K. A., Zeps, N. J., & Hoskins, A. A. (2023). Biochemical and Genetic Evidence Supports Fyv6 as a Second-Step Splicing Factor in Saccharomyces cerevisiae. *RNA*, rna.079607.123. https://doi.org/10.1261/rna.079607.123

Liu, L., Query, C. C., & Konarska, M. M. (2007). Opposing classes of prp8 alleles modulate the transition between the catalytic steps of pre-mRNA splicing. *Nature Structural & Molecular Biology*, *14*(6), Article 6. https://doi.org/10.1038/nsmb1240

Liu, S., Li, X., Zhang, L., Jiang, J., Hill, R. C., Cui, Y., Hansen, K. C., Zhou, Z. H., & Zhao, R. (2017). Structure of the yeast spliceosomal postcatalytic P complex. *Science, 358*(6368), Article 6368. https://doi.org/10.1126/science.aar3462

Liu, Y.-C., Chen, H.-C., Wu, N.-Y., & Cheng, S.-C. (2007). A Novel Splicing Factor, Yju2, Is Associated with NTC and Acts after Prp2 in Promoting the First Catalytic Reaction of Pre-mRNA Splicing. *Molecular and Cellular Biology*, *27*(15), 5403–5413. https://doi.org/10.1128/MCB.00346-07

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Mayas, R. M., Maita, H., & Staley, J. P. (2006). Exon ligation is proofread by the DExD/H-box ATPase Prp22p. *Nature Structural & Molecular Biology*, *13*(6), Article 6. https://doi.org/10.1038/nsmb1093

McPheeters, D. S., & Abelson, J. (1992). Mutational analysis of the yeast U2 snRNA suggests a structural similarity to the catalytic core of group I introns. *Cell*, *71*(5), 819–831. https://doi.org/10.1016/0092-8674(92)90557-s

Mendoza-Ochoa, G. I., Barrass, J. D., Maudlin, I. E., & Beggs, J. D. (2019). Blocking late stages of splicing quickly limits pre-spliceosome assembly in vivo. *RNA Biology*, *16*(12), 1775–1784. https://doi.org/10.1080/15476286.2019.1657788

Montemayor, E. J., Curran, E. C., Liao, H. H., Andrews, K. L., Treba, C. N., Butcher, S. E., & Brow, D. A. (2014). Core structure of the U6 small nuclear ribonucleoprotein at 1.7-Å resolution. *Nature Structural & Molecular Biology*, *21*(6), Article 6. https://doi.org/10.1038/nsmb.2832

Mumberg, D., Müller, R., & Funk, M. (1995). Yeast vectors for the controlled expression of heterologous proteins in different genetic backgrounds. *Gene*, *156*(1), 119–122. https://doi.org/10.1016/0378-1119(95)00037-7

Munding, E. M., Shiue, L., Katzman, S., Donohue, J. P., & Ares, M. (2013). Competition between Pre-mRNAs for the Splicing Machinery Drives Global Regulation of Splicing. *Molecular Cell*, *51*(3), 338–348. https://doi.org/10.1016/j.molcel.2013.06.012

Nguyen Ba, A. N., Pogoutse, A., Provart, N., & Moses, A. M. (2009). NLStradamus: A simple Hidden Markov Model for nuclear localization signal prediction. *BMC Bioinformatics*, *10*(1), 202. https://doi.org/10.1186/1471-2105-10-202

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419. https://doi.org/10.1038/nmeth.4197

Perriman, R., Barta, I., Voeltz, G. K., Abelson, J., & Ares, M. (2003). ATP requirement for Prp5p function is determined by Cus2p and the structure of U2 small nuclear RNA. *Proceedings of the National Academy of Sciences*, *100*(24), 13857–13862. https://doi.org/10.1073/pnas.2036312100

Pettersen, E. F., Goddard, T. D., Huang, C. C., Meng, E. C., Couch, G. S., Croll, T. I., Morris, J. H., & Ferrin, T. E. (2021). UCSF ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Science*, *30*(1), 70–82. https://doi.org/10.1002/pro.3943

Plaschka, C., Newman, A. J., & Nagai, K. (2019). Structural Basis of Nuclear pre-mRNA Splicing: Lessons from Yeast. *Cold Spring Harbor Perspectives in Biology*, *11*(5), a032391. https://doi.org/10.1101/cshperspect.a032391

Query, C. C., & Konarska, M. M. (2004). Suppression of Multiple Substrate Mutations by Spliceosomal prp8 Alleles Suggests Functional Correlations with Ribosomal Ambiguity Mutants. *Molecular Cell*, *14*(3), 343–354. https://doi.org/10.1016/S1097-2765(04)00217-5

Query, C. C., & Konarska, M. M. (2012). CEF1/CDC5 alleles modulate transitions between catalytic conformations of the spliceosome. *RNA*, *18*(5), 1001–1013. https://doi.org/10.1261/rna.029421.111

Roy, K. R., Gabunilas, J., Neutel, D., Ai, M., Yeh, Z., Samson, J., Lyu, G., & Chanfreau, G. F. (2023). Splicing factor Prp18p promotes genome-wide fidelity of consensus 3′-splice sites. *Nucleic Acids Research*, gkad968. https://doi.org/10.1093/nar/gkad968

Sayani, S., Janis, M., Lee, C. Y., Toesca, I., & Chanfreau, G. F. (2008). Widespread Impact of Nonsense-Mediated mRNA Decay on the Yeast Intronome. *Molecular Cell*, *31*(3), 360–370. https://doi.org/10.1016/j.molcel.2008.07.005

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of image analysis. *Nature Methods*, *9*(7), 671–675. https://doi.org/10.1038/nmeth.2089

Schwer, B. (2008). A Conformational Rearrangement in the Spliceosome Sets the Stage for Prp22-Dependent mRNA Release. *Molecular Cell*, *30*(6), Article 6. https://doi.org/10.1016/j.molcel.2008.05.003

Schwer, B., & Gross, C. H. (1998). Prp22, a DExH-box RNA helicase, plays two distinct roles in yeast pre-mRNA splicing. *The EMBO Journal*, *17*(7), 2086–2094. https://doi.org/10.1093/emboj/17.7.2086

Schwer, B., & Guthrie, C. (1992). A conformational rearrangement in the spliceosome is dependent on PRP16 and ATP hydrolysis. *The EMBO Journal*, *11*(13), 5033–5039. https://doi.org/10.1002/j.1460-2075.1992.tb05610.x

Schwer, B., & Meszaros, T. (2000). RNA helicase dynamics in pre-mRNA splicing. *The EMBO Journal*, *19*(23), 6582–6591. https://doi.org/10.1093/emboj/19.23.6582

Semlow, D. R., Blanco, M. R., Walter, N. G., & Staley, J. P. (2016). Spliceosomal DEAH-Box ATPases Remodel Pre-mRNA to Activate Alternative Splice Sites. *Cell*, *164*(5), Article 5. https://doi.org/10.1016/j.cell.2016.01.025

Semlow, D. R., & Staley, J. P. (2012). Staying on message: Ensuring fidelity in pre-mRNA splicing. *Trends in Biochemical Sciences*, *37*(7), 263–273. https://doi.org/10.1016/j.tibs.2012.04.001

Siatecka, M., Reyes, J. L., & Konarska, M. M. (1999). Functional interactions of Prp8 with both splice sites at the spliceosomal catalytic center. *Genes & Development*, *13*(15), 1983–1993.

Sikorski, R. S., & Boeke, J. D. (1991). In Vitro Mutagenesis and Plasmid Shuffling: From Cloned Gene to Mutant Yeast. *Methods in Enzymology*, *194*(C), 302–318. https://doi.org/10.1016/0076-6879(91)94023-6

Strittmatter, L. M., Capitanchik, C., Newman, A. J., Hallegger, M., Norman, C. M., Fica, S. M., Oubridge, C., Luscombe, N. M., Ule, J., & Nagai, K. (2021). psiCLIP reveals dynamic RNA binding by DEAH-box helicases before and after exon ligation. *Nature Communications*, *12*(1), 1488. https://doi.org/10.1038/s41467-021-21745-9

Talkish, J., Igel, H., Hunter, O., Horner, S. W., Jeffery, N. N., Leach, J. R., Jenkins, J. L., Kielkopf, C. L., & Ares, M. (2019). Cus2 enforces the first ATP-dependent step of splicing by binding to yeast SF3b1 through a UHM–ULM interaction. *RNA*, *25*(8), 1020–1037. https://doi.org/10.1261/rna.070649.119

Townsend, C., Leelaram, M. N., Agafonov, D. E., Dybkov, O., Will, C. L., Bertram, K., Urlaub, H., Kastner, B., Stark, H., & Lührmann, R. (2020). Mechanism of protein-guided folding of the active site U2/U6 RNA during spliceosome activation. *Science*, *370*(6523), eabc3753. https://doi.org/10.1126/science.abc3753

Treco, D. A., & Lundblad, V. (1993). Preparation of Yeast Media. *Current Protocols in Molecular Biology*, *23*(1), 13.1.1-13.1.7. https://doi.org/10.1002/0471142727.mb1301s23

Vorländer, M. K., Rothe, P., Kleifeld, J., Cormack, E. D., Veleti, L., Riabov-Bassat, D., Fin, L., Phillips, A. W., Cochella, L., & Plaschka, C. (2024). Mechanism for the initiation of spliceosome disassembly. *Nature*, *632*(8024), 443–450. https://doi.org/10.1038/s41586-024-07741-1

Wan, R., Yan, C., Bai, R., Lei, J., & Shi, Y. (2017). Structure of an Intron Lariat Spliceosome from Saccharomyces cerevisiae. *Cell*, *171*(1), 120-132.e12. https://doi.org/10.1016/j.cell.2017.08.029

Wilkinson, M. E., Fica, S. M., Galej, W. P., & Nagai, K. (2021). Structural basis for conformational equilibrium of the catalytic spliceosome. *Molecular Cell*. https://doi.org/10.1016/j.molcel.2021.02.021

Wilkinson, M. E., Fica, S. M., Galej, W. P., Norman, C. M., Newman, A. J., & Nagai, K. (2017). Postcatalytic spliceosome structure reveals mechanism of 3′–splice site selection. *Science*, *358*(6368), 1283–1288. https://doi.org/10.1126/science.aar3729

Wilkinson, M. E., Kumar, A., & Casañal, A. (2019). Methods for merging data sets in electron
cryo-microscopy. *Acta Crystallographica Section D: Structural Biology*, *75*(9), 782–791.
https://doi.org/10.1107/S2059798319010519

Wong, A. C. H., Wong, J. J.-L., Rasko, J. E. J., & Schmitz, U. (2024). SpliceWiz: Interactive
analysis and visualization of alternative splicing in R. *Briefings in Bioinformatics*, *25*(1),
bbad468. https://doi.org/10.1093/bib/bbad468

Wu, T., Ruan, K., & Liu, W. (1996). A Fluorescence-Labeling Method for Sequencing Small RNA
on Polyacrylamide Gel. *Nucleic Acids Research*, *24*(17), 3472–3473.
https://doi.org/10.1093/nar/24.17.3472

Xu, Y.-Z., & Query, C. C. (2007). Competition between the ATPase Prp5 and Branch Region-U2
snRNA Pairing Modulates the Fidelity of Spliceosome Assembly. *Molecular Cell*, *28*(5),
838–849. https://doi.org/10.1016/j.molcel.2007.09.022

Yan, C., Hang, J., Wan, R., Huang, M., Wong, C. C. L., & Shi, Y. (2015). Structure of a yeast
spliceosome at 3.6-angstrom resolution. *Science*, *349*(6253), 1182–1191.
https://doi.org/10.1126/science.aac7629

Yan, D., Perriman, R., Igel, H., Howe, K. J., Neville, M., & Ares Jr., M. (1998). CUS2, a Yeast
Homolog of Human Tat-SF1, Rescues Function of Misfolded U2 through an Unusual
RNA Recognition Motif. *Molecular and Cellular Biology*, *18*(9), 5000–5009.
https://doi.org/10.1128/MCB.18.9.5000

Zhan, X., Lu, Y., Zhang, X., Yan, C., & Shi, Y. (2022). Mechanism of exon ligation by human
spliceosome. *Molecular Cell*, *82*(15), Article 15.
https://doi.org/10.1016/j.molcel.2022.05.021

Zhang, X., & Schwer, B. (1997). Functional and physical interaction between the yeast splicing
factors Slu7 and Prp18. *Nucleic Acids Research*, *25*(11), 2146–2152.
https://doi.org/10.1093/nar/25.11.2146

Zhang, X., Zhan, X., Yan, C., Zhang, W., Liu, D., Lei, J., & Shi, Y. (2019). Structures of the
human spliceosomes before and after release of the ligated exon. *Cell Research*, *29*(4),
Article 4. https://doi.org/10.1038/s41422-019-0143-x

1190

1191

1192

Figure 1. Widespread splicing changes due to loss of Fyv6. A) Pre-mRNA splicing reaction steps and products. **B)** Growth of WT (*cup1Δ*), *fyv6Δ*, *upf1Δ* and *fyv6Δupf1Δ* yeast at 16, 23, 30, and 37°C. Images were taken on the indicated days. White lines indicate where images were cropped, as all dilutions for all strains were on the same plate, but not necessarily directly adjacent to one another. **C)** Number of alternative splicing events observed in a *fyv6Δupf1Δ* strain relative to a strain containing FYV6 after a temperature shift to 16, 30, and 37°C. **D)** Venn diagram of shared alternative 3′ SS splicing events at 16 (blue), 30 (green), and 37°C (red). **E)** Box plots of the log$_{10}$ of the ratios of *fyv6Δ upf1Δ* to WT (*upf1Δ*) Fraction of Annotated Splicing (FAnS) values for RPGs (red) and non-RPGs (black) for 3′ SS splicing events. FAnS values represent the ratio of the alternative splicing event (here, selection of an alternate 3′ SS) to the annotated, canonical splice isoform.

Figure 2: Loss of Fyv6 activates BP proximal, non-consensus 3' SS. A) Plot of the $\log_{10}$ of Fraction of Annotated Splicing (FAnS) values for differences between alternative 3' SS usage in *fyv6Δ upf1Δ* and *upf1Δ* strains. High FAnS values indicate increased usage of the alternate 3' SS relative to the annotated site. Points are colored based on whether they are upstream (pink) or downstream (blue) to the canonical 3' SS. **B)** Violin plots of the ratios of *fyv6Δupf1Δ* to the *upf1Δ* FAnS values based on the distances between the alternative and canonical 3' SS (Δ3' SS = 3' $SS_{Alt.}$ - 3' $SS_{canonical}$). FAnS ratios >0 are indicative of the site being upregulated in *fyv6Δ*. **C)** Violin plots of the ratios of *fyv6Δupf1Δ* to the *upf1Δ* FAnS values based on the distances between the alternative 3' SS and the annotated BP. FAnS ratios >0 are indicative of the site being upregulated in *fyv6Δ*. **D)** Diagram of ACT1-CUP1 reporter showing BP-3' SS distances. **E)** Representative primer extension analysis of RNA products generated from splicing of the ACT1-CUP1 reporter in the presence (WT) or absence of Fyv6 (*fyv6Δ*). Bands for fully spliced mRNA and lariat intermediate are indicated. U6 snRNA was detected as a loading control. The * indicates an unknown product present in every lane. **F)** Quantification of the primer extension results from *N*=3 replicates represented by the ratio of band intensities for mRNA/(mRNA+lariat intermediate). Bars represent the average ratio of the replicates ±SD. Means between WT and *fyv6Δ* groups for each reporter were compared with an unpaired Welch's two-tailed t-test. Significance is indicated: n.s. no significance; ** p<0.01; *** p<0.001. **G)** Sequence logos of

alternative 3' SS with FAnS > 0 sorted by either upstream or downstream of the canonical 3' SS compared against the canonical 3' SS for genes with FAnS > 0. **H)** The $\log_{10}$ of the FAnS ratio for alternative 3' SS sorted by the sequences of the 3' SS.

Figure 3. Cryo-EM structure of the yeast P complex spliceosome at 2.3 Å resolution. A) A composite density map for P complex showing focused refinements of the Prp22, NTC, U2 snRNP, U5 Sm ring, and Cwc22 N-terminal domain regions. **B)** Overall model for the P complex spliceosome. **C)** Cryo-EM density for state I of P complex (above, low-pass filtered) and for the active site and 3' SS (below, sharpened). **D,E)** As for **C)** but for states II (**D**) and III (**E**).

Figure 4. Fyv6 and its interactors in the P complex cryo-EM structure. A) Cryo-EM density segmented around Fyv6. **B)** Structure of Prp22. **C)** Structure of the NTC within P complex. **D)** IP$_6$ site and cryo-EM density in NTC. **E)** Interaction of the hook domain of Fyv6 with the Prp22 RecA2 domain. **F)** Interaction of Fyv6 with Slu7 and the region of the intron between the BP and 3' SS. **G)** Interaction of Fyv6 with Syf1 and an analogous view of the interaction of Syf1 with Yju2 in C complex (Wilkinson et al., 2021).

Figure 5. Structure-based analysis of Fyv6 domain function. A) Diagram of Fyv6 protein structure, protein interactors, and protein truncations. NLS= nuclear localization signal **B)** Spot dilution assay of strains with Fyv6 truncations on –Trp DO plates at different temperatures. Plates were imaged after the number of days indicated. **C)** Representative gel image of *SUS1* RT-PCR in strains with an empty vector or expressing the indicated Fyv6 variant. (+RT reactions contain reverse transcriptase; -RT control reactions do not contain reverse transcriptase.) **D)** Diagram of Syf1 protein structure and truncations. **E)** Spot dilution assay of strains with Syf1 truncations on –Trp DO plates at different temperatures. Plates were imaged after the number of days indicated. **F)** Representative gel image of *SUS1* RT-PCR in strains with an empty vector or expressing the indicated Syf1 variant. **G)** Temperature growth assay of strains with Syf1 truncations in a *fyv6Δ* background on YPD plates. Plates were imaged after the number of days indicated. **H)** Representative gel image of RT-PCR of *SUS1* in strains with plasmids containing WT Syf1 or a Syf1 truncation in a *fyv6Δ* background.

Figure 6. Identification of novel suppressors of *fyv6Δ* temperature sensitivity. A) Workflow for the suppressor screen. **B)** P complex spliceosome structure with *fyv6Δ* suppressor mutations labelled. (See **Supplemental Table S5**) **C)** Close-up view of Cef1 and Prp8 suppressor mutations in P complex. **D)** Spot dilution temperature growth assay with Cef1 mutants on YPD plates. Plates were imaged after the number of days indicated. **E)** Close-up view of Prp8 mutations in P complex. **F)** Spot dilution temperature growth assay with Prp8 mutants on YPD plates. Plates were imaged after the number of days indicated.

Figure 7. Genetic interactions between Prp22 and Fyv6. A) Location of the Prp22 I1133R *fyv6Δ* suppressor in the P complex structure. **B)** Spot dilution growth assay with Prp22 mutants in WT or *fyv6Δ* backgrounds grown on –Trp DO or –Trp+5-FOA plates. Plates were imaged after 3 days at 30°C. **C)** Spot dilution growth assay with strains containing Prp22 mutants on YPD plates at different temperatures. Plates were imaged after the number of days indicated. **D)** Representative gel of *SUS1* RT-PCR products in strains with Prp22 mutants in the presence or

absence of Fyv6. **E)** Quantitation of band intensities for each *SUS1* splice isoform as a fraction of total product in the lane. Bars indicate average ± SD of *N*=3 replicates. **F)** ACT1-CUP1 reporter construct with the location of the 3' SS substitution noted. **G)** Copper tolerance for each ACT1-CUP1 reporter with each Prp22 allele in strains with (■) or without (○) Fyv6 present. Shown is a representative, single replicate of *N*=3.

Figure 8. Model for the role of Fyv6 during the 2nd step. Fyv6 interacts with Syf1 and Prp22 and promotes usage of BP distal 3' SS. When Fyv6 is absent, BP proximal 3' SS are favored. It is possible that lack of Fyv6 also results in the presence of Yju2 during exon ligation and relaxing of the requirement for Prp22 at this step.