Mitochondria-derived Vesicles mediate Mito-nuclear Trafficking of Cargo

Thejaswitha Rajeev[1][2], Dipak Shil[1], Swagato Bhattacharjee[1], Lariza Ramesh[1], Victor Samuel I[1], Nidhi Nair[1], Areeba Marib[1], Mrithulabhashini Jayakumar[1], Sai Gayathri[1], Arun Surendran[1], John B Johnson[1], Ananthalakshmy Sundararaman[1]*

[1] Rajiv Gandhi Centre for Biotechnology, Poojappura, Thycaud PO, Thiruvananthapuram, Kerala, India

[2] Manipal Academy of Higher Education, Tiger Circle Road, Madhav Nagar, Manipal, Karnataka, India

*Correspondence: ananthalakshmys@rgcb.res.in

Abstract

Mito-nuclear communication is key to cellular homeostasis. Metabolites from TCA cycle, calcium, ROS and other small molecules are known to mediate retrograde signalling of mitochondria to the nucleus. Recently, the direct transfer of mitochondrial proteins to the nucleus has been described. However, the mechanism for direct transfer of mitochondrial protein complexes to the nucleus is not understood. In this study, we demonstrate the transit of mitochondrial proteins to the nucleus through mitochondria-derived vesicles (MDVs) which represents a novel intracellular trafficking route. Following an unbiased proteomic approach to detecting cargo proteins in cardiac MDVs, we screen for the presence of MDV cargo in the nucleus. Pyruvate dehydrogenase complex (PDH) and Cytochrome c oxidase subunit IV isoform 1 (COX4I1) are packaged into MDVs in cardiomyocytes but only PDH is targeted to the nucleus in this cell type demonstrating that a subset of cargo-selective MDVs fuse with the nucleus. Further, several mitochondrial stresses that are known to increase MDV generation triage these MDVs to other cellular destinations thereby reducing the nuclear pool of PDH. The transit of PDH through MDVs to the nucleus is a basal physiological process in cardiomyocytes. Mito-nuclear trafficking thus represents a physiological pathway in cardiac cells to enable cargo shuttle from mitochondria to the nucleus.

Keywords: Mitochondria-derived vesicles (MDVs), mito-nuclear communication, Pyruvate dehydrogenase (PDH)

Introduction

Cardiomyocytes rely heavily on mitochondria for energy through oxidative phosphorylation which contributes to 90% of cellular ATP levels [1]. Mitochondrial dysfunction contributes to various cardiovascular diseases. Hence there exists various mitochondrial quality control pathways to ensure a healthy mitochondrial network [2]. One such quality control mechanism discovered recently is through mitochondria-derived vesicles (MDVs) that transport oxidatively damaged proteins to lysosomes for degradation [3]. While MDV biology is not yet explored in depth in cardiac cells, they are present in a basal physiological state in this system [4]. Interestingly, fission-fusion cycle is extremely rare in cardiomyocytes [5] which is an important quality control mechanism in other cell types. Therefore, we hypothesise that cardiac cells would be highly reliant on MDVs to maintain mitochondrial homeostasis and

for inter-organellar communication. Our study therefore aims to understand MDV biogenesis,
its trafficking, particularly to the nucleus and its relevance to the cardiac system.

Mitochondria-derived vesicles are cargo selective vesicles of 70-150 nm in size that originate
from mitochondria. They can be single or double membrane structures that can target cargo
to cell destinations like the peroxisome and lysosome [3, 6]. MDV generation and targeting is
aided by various proteins including Vps35, PINK1/PARKIN, syntaxin 17-SNARE complex,
Sorting nexin 9-Rab 9, Rab 7, and Tollip [7-13]. Previous studies have shown that MDVs are
formed independent of the fission protein, Drp1 [6, 10]. But, in the case of the widely studied
TOM20[+] MDVs, MIRO1/2 and Drp1 are required for MDV formation [14]. MDVs thus
generated play a variety of roles including peroxisome biogenesis, mitochondrial quality
control, mitochondrial antigen presentation, bacterial killing within phagosomes and in the
regeneration of ATP deficient mitochondria [3, 6, 10, 15, 16]. However, MDVs have not
been implicated in mito-nuclear cargo transit which we explore here.

Mitochondria in eukaryotes evolved from an α-proteobacterial ancestor capable of secreting
outer membrane vesicles (OMV) for inter-colony communication and interaction with host
cells [17-19]. As MDVs are speculated to be evolutionarily descended or closely related to
bacterial OMVs we fathom that inter-organellar communication, particularly between the
endosymbiont and host nuclei might be a key physiological role for MDVs. We therefore
hypothesised that MDVs mediate mito-nuclear protein transit. Several mitochondrial proteins
are known to have moonlighting functions in the nucleus. For example, Nuclear factor
erythroid 2-related factor 2 (NRF2), CLOCK-1 (CLK-1), fumarase, Lonp1 and TCA cycle
enzymes are mitochondrial proteins also found to function in the nucleus in different systems
[20-24]. In mouse embryo, pyruvate carboxylase (PCB), citrate synthase (CS), aconitase 2
(ACO2), isocitrate dehydrogenase 3 (IDH3A) and pyruvate dehydrogenase (PDHE1) are
required in the nucleus for early zygotic genome activation (ZGA) [25]. The same enzymes
have been found to play a role in somatic cell reprogramming as well [26]. In the only study
exploring moonlighting functions of TCA enzymes in the cardiac system, malate
dehydrogenase 2 (MDH2), pyruvate dehydrogenase subunit 1 (PDH-E1) and isocitrate
dehydrogenase 2 (IDH-2) were found in the nucleus of hiPSC-derived cardiomyocytes upon
doxorubicin-mediated cardiotoxicity [27]. A well studied nuclear translocated mitochondrial
protein is pyruvate dehydrogenase (PDH) [28, 29] which is also independently characterised
as an MDV cargo [3, 30].

Pyruvate dehydrogenase complex (PDC) is a ~10 MDa, multi-subunit complex of diameter
45 nm, linking glycolysis and TCA cycle through acetyl CoA generation. The subunits are
PDHE1 (pyruvate dehydrogenase), PDHE2/DLAT (dihydrolipoamide acetyltransferase),
PDHE3 (dihydrolipoamide dehydrogenase), and E3BP (E3 binding protein) [29, 31, 32]. It
has been shown that the nucleus accumulates only the mitochondria localization signal
(MLS) cleaved mature form of PDC subunits and since the peptidase to process this is
available only in the mitochondria it stands to reason that PDH reaches the nucleus from the
mitochondria. Nuclear translocation of PDC occurs when cells are exposed to proliferative
stimuli or mitochondrial stresses like hypoxia [28, 29]. But prolonged hypoxia can lead to
decreased nuclear PDH [33]. Within the nucleus, PDH is believed to regulate gene expression
through histone acetylation [26, 28, 29] as seen in the regulation of the expression of Sterol
regulatory element-binding transcription factor (SREBF) by nuclear PDC in prostate cancer
[34]. So, nuclear PDC has several cell type-specific roles like cardioprotective response to
cardiotoxicity, tumour progression, epigenetic reprogramming after T cell activation [27, 34,
35].

Here, we hope to address the question of the entry of pyruvate dehydrogenase complex which cannot freely diffuse nor pass through the nuclear pore complex (NPC) [29]. We propose that packaging of either the pyruvate dehydrogenase complex or their individual subunits into MDVs as a means of mito-nuclear trafficking. MDVs labelled with PDHE2/E3bp are reported [36] but these are believed to exclude PDHE1 [8]. On the other hand, some papers show PDHE1[+] MDVs but their cellular destinations are not explored [10, 37]. MDV transit and corresponding changes to nuclear pool of cargo proteins have not been studied in tandem. It is reported that PDH[+] MDV number increases in response to xanthine oxidase as well as antimycin-A treatment [4, 30]. PDH[+] MDV biogenesis also increases in cardiotoxicity after chemotherapy [4] but in these cases the cellular destination of PDH[+] MDVs remain unexplored. Interestingly, PDH[+] MDVs can triage to different cell destinations based on cellular context, for example Parkin dependent MDVs to lysosomes and Parkin independent MDVs to endosomes and extracellular vesicles [8, 9, 11]. In this study we unravel the cellular context under which PDH in MDVs could be directed to the nucleus thereby describing an unknown trafficking route for MDVs.

Results

Cardiac MDV proteome is enriched for mitochondrial proteins involved in TCA cycle and electron transport chain.

We initially took an unbiased proteomic approach to generate MDV cargo list from rat hearts through *in vitro* budding assay. This approach has been used previously [30, 36, 38] but our budding reactions additionally incorporate N-ethyl maleimide (NEM) [39]and we further analyse cargos based on their NEM sensitivity. Figure 1A shows the steps in our cell free MDV budding assay. A detailed protocol is available from our group in Nair et al., 2024 [40] .

To confirm that the control and NEM budding reactions contained vesicles, the reactions were visualised using TEM. We observed vesicular structures ranging from 70 to 150 nm in size which is within the size range of MDVs [6]. There were less vesicular structures in the NEM budding reaction as NEM is expected to reduce the budding of the vesicles (Figure 1B).

To further confirm our protocol, we immunoblotted the budding reaction with some of the previously reported MDV cargo like VDAC, Tom20, Opa1 and PDH [3, 11, 30, 38]. We confirmed that all these cargo proteins are incorporated into MDVs in an NEM sensitive manner (Figure 1C). For further validation of the packaging of these proteins in vesicles, we floated the control budding reaction in a sucrose gradient by ultracentrifugation. We observed that the matrix protein, PDH, was also present in the denser sucrose fractions compared to all other proteins from other compartments (Figure 1D). Together our TEM, immunoblotting of the budding reactions and floatation gradients validate our in vitro approach to generation of MDVs from rat heart.

Having validated the cell free MDV budding, we proceeded to identify the cargo packaged within these MDVs by mass spectrometry (Supplementary Data S1). We used multiple sources including MitoCarta 3.0 and Integrated Mitochondrial Protein Index (IMPI) database to confirm the mitochondrial localization and sublocalization of the detected proteins. Of the total 453 proteins detected in the three independent MS analyses of rat cardiac MDV proteome, 211 proteins were mitochondrial proteins. Out of the 211 mitochondrial proteins only those proteins that appeared twice in the three MS runs were analysed further as MDV cargo. There were 110 mitochondrial proteins primarily of matrix and inner mitochondrial

origin in our MDV proteome (Figure S1A). This preponderance of IMM and matrix proteins
in the in vitro generated MDVs is also reported previously [36]. Pathway analysis indicated
that these proteins belong mostly to the group of TCA cycle enzymes and electron transport
chain proteins (Figures S1C and S1D).

We next considered NEM-sensitive proteins (present in the control reaction but absent in the
NEM containing reaction) as a more stringent list of MDV cargo. This eliminates any
artifactually present proteins in the supernatant post-budding. When NEM sensitivity was
used as cut-off 61 proteins were identified as MDV cargo in at least two out of the three MS
runs (Figure 1G). These were also predominantly matrix and inner mitochondrial membrane
proteins playing roles in TCA cycle and electron transport chain (Figure S1B).

In our NEM sensitive list of MDV cargo, we noted multiple subunits of the PDC complex
like PDHA1 (PDHE1α), PDHB (PDHE1β), DLAT (PDHE2) and PDHX (PDHE3 binding
protein) (Figure 1G) which have previously been used to mark matrix positive MDVs in the
field [8, 10]. Additionally, we noticed both subunits of SDH and components of the complex
IV of the respiratory electron transport chain. We validated the new hits like COX4I and
SDH as being incorporated into MDVs in an NEM sensitive manner *in vitro* (Figure S1E)
and further demonstrated the presence of COX4I1 and SDH in MDVs floated on sucrose
gradients (Figure S1F).

Next, we validated the formation of MDVs in our cells of interest, H9C2 cardiomyoblasts.
PDH$^+$ MDVs were already known to exist basally in H9C2 cardiomyoblasts and were
upregulated in the presence of an oxidative stress induced by antimycin A [4]. We observed
COX4$^+$PDH$^-$, PDH$^+$COX4$^-$ and COX4$^+$PDH$^+$ dual positive MDVs basally and found that
these MDVs were antimycin sensitive (Figures S1 G-K).

Together our NEM sensitive MDV cargo part list, while likely incomplete, adds to the set of
high confidence MDV cargo that can be used to understand MDV biology in the cardiac
system.

PDH, but not COX4, is found in the nucleus of rat cardiac cells

To examine if any of the mitochondrial proteins that we identified as MDV cargo were
localized to the nucleus, we performed an initial screening in the nuclei of HEK293T cells.

We detected mitochondrial proteins from various sublocalizations in the nuclear isolates of
HEK293T (Figures 2A and S2A). We observed the outer mitochondrial membrane proteins
(VDAC and TOM20) in the nucleus. We also observed COX4, the newly identified MDV
cargo in the nucleus. COX4 is a subunit of the Complex IV of the ETC. We did not detect the
other subunits of Complex IV, COX1 and COX10 in the nucleus suggesting that not the
entire Complex IV but subassemblies may be transiting to the nucleus. Prohibitin and
NDUFS1 are inner membrane proteins detected in the nucleus in our system. While
Prohibitin is already known to be present in the nucleus [41], the other proteins are not
known to be present in the nucleus. Several of the other inner mitochondrial proteins that
were part of or closely associated with the ETC, including RISP, ACP, SDH and cytochrome
c were absent in the nucleus. Also, inner membrane protein involved in membrane fission
Opa1, was absent in the nucleus.

We observed trace amounts of PDH in the nuclei of HEK293T. PDH was previously reported
in the nucleus of other cell types [25, 28, 42, 43]. Other matrix proteins like SOD1 and

HSP60 were not detected in the nucleus. Of the screened proteins detected in the nucleus,
PDH, TOM20, VDAC and COX4 were MDV cargo validated by us and other groups.

Next, we investigated the presence of mitochondrial proteins in the nucleus of H9C2
cardiomyoblasts and rat heart nuclei. We observed VDAC in the nuclei from both H9C2 and
rat heart but detected TOM20 only in H9C2 cardiomyoblasts. We did not detect COX4 or any
of the other Complex IV subunits (COX1 and COX10) in the nuclei of either H9C2
cardiomyoblasts or rat heart. But PDH was detected in the nucleus in cardiac cells (Figures
2B-C, S2B-C). In H9C2 cardiomyoblasts overexpressing PDH mCherry, we observed PDH[+]
MitoTracker [negative] vesicular structures near the mitochondria before their fusion with the
nucleus to release PDH in the nucleoplasm(Figure S2E, Supplementary movie Data S2). We
also observed that in endothelial cells, PDH but not COX4 is present in the nucleus (Figure
S2D). Together, we confirm that nuclear transit of proteins including PDH and COX4 are cell
type specific and PDH but not COX4 is present in the nucleus of cardiac and endothelial
cells.

It is noteworthy that not all of the screened MDV cargo gets delivered to the nucleus as
observed in the cases of COX4, OPA1 and SDH. This highlights the specificity of mito-
nuclear protein transit. While the screen does not provide evidence for the transit being
MDV-mediated, lack of transit of some proteins and the cell type specificity in others ensure
that this is not an artifactual co-purification of some mitochondria with the nuclear pool. We
further explore PDH transit in detail as it is present in the nuclei of cardiomyoblast cells and
the rat heart tissue.

Actin and tubulin depolymerization leads to decreased PDH[+] MDVs.

It is known that TOM20[+] MDVs are trafficked to endolysosomes in a microtubule dependent
manner with the aid of MIRO1/2 and DRP1 [14]. However, other cytoskeletal elements like
actin which is known to control vesicle formation has not been studied in the context of MDV
biogenesis and trafficking [44, 45]. We wanted to explore the role of actin and tubulin in
PDH[+]MDV biogenesis in cardiomyocytes.

We had previously observed that antimycin A upregulates the number of PDH[+] MDVs
formed. We sought to investigate the potential impact of actin or tubulin depolymerization on
antimycin-driven mitochondrial-derived vesicles (MDVs). First, we had confirmed that the
actin and tubulin depolymerized at the concentrations (1µM and 10 µM respectively) used in
our study (Figures S3A and S3B). We observed a significant drop in the number of PDH[+]
MDVs formed in the presence of latrunculin B (Figures 3A and 3B) or nocodazole (Figures
3C and 3D). This is evidence that cytoskeletal elements like actin and tubulin play a role in
PDH[+] MDV biogenesis. To uncouple MDV trafficking and nuclear fusion events from
budding we undertook an *in vitro* reconstitution approach.

***In vitro* reconstitution of mito-nuclear trafficking shows that PDH delivery to the**
nucleus is NEM sensitive.

To determine the mechanism of transit of mitochondrial proteins to the nuclei, we incubated
rat cardiac MDVs with rat heart nuclei as shown in the graphical representation (Figure 4A).
GTP (0.5 µM) was added to all the reconstitution reactions. The nuclear purity of our
isolation could be noted by the absence of OPA1 and COX4 in the nuclear isolates as
observed previously. We observed increased nuclear PDH in rat nuclei incubated with
MDVs (Figures 4B and 4C) providing direct evidence for the delivery of PDH by MDVs to
the nucleus. By addition of NEM to the cell free reconstitution reaction, we inhibited
trafficking proteins required for the fusion of membranes. We noted that PDH entry into the

nucleus was NEM sensitive. This proved that the fusion of MDVs with nucleus and PDH delivery required NEM sensitive proteins. We also observed that some of the other MDV cargo like NDUFS1 and VDAC did not appear to increase upon MDV addition or further reduce with NEM (Figure S4A). Therefore, these two proteins might enter the nucleus without MDV mediated shuttling.

Mitochondrial stress represses PDH transit to the nucleus.

As our previous results indicate a transit of PDH to the nucleus in cardiac cells, we wanted to understand the physiological and pathophysiological conditions under which this transit is modulated. When cells are grown in galactose containing medium, they mostly rely on mitochondrial function through oxidative phosphorylation to generate ATP [46, 47]. When we probed for PDH in the nuclear isolates of cells grown in the presence of glucose and compared it with 20 mM galactose containing media (with a low-5 mM glucose) we found that the nuclear PDH levels increase almost 1.5 folds in galactose containing medium (Figures 5A and 5B). Concomitantly we noted that the total cellular PDH levels remain unchanged in H9C2 cells when cultured for 24 hours in galactose containing media (Figures S5A and S5B). This suggests that PDH is transiting from the mitochondria to the nucleus under these conditions. Previously, it was shown that PDH and other TCA cycle enzymes transit to the nucleus in a pyruvate dependent manner during ZGA [25]. We assessed the role of pyruvate in nuclear PDH transit in H9C2 cardiomyoblasts. However, in our system no significant change was observed in nuclear PDH levels in the absence or presence of pyruvate in cells cultured in either glucose or galactose medium (Figures 5A and 5B). Again, we confirmed that pyruvate did not affect the total levels of PDH in the cells (Figures S5A and S5B).

Further we were interested to see the status of PDH transit to the nucleus under differentiation of H9C2 cardiomyoblast to cardiomyocytes-like cells. Firstly, we confirmed that indeed differentiation is happening upon retinoic acid and 1% serum growth. Cells change in morphology and appear more flattened after differentiation as can be seen in phase contrast and phalloidin stained images (Figures S5C and S5D). Differentiation was also confirmed by an increase in the percentage of multinucleate cells (data not shown). Next, we checked the number of PDH$^+$ MDVs in control and differentiated cells and observed an increase PDH$^+$ MDVs in our system (Figures S5M and S5N) as has been observed previously [4]. Next, we queried whether nuclear PDH pool responds to this change in cell state. Indeed, we observed a twofold increase in nuclear PDH upon differentiation compared to control cells (Figures 5C and 5D). Previous reports showed that mitochondrial mass increases in the process of H9C2 differentiation [48]. Following the same trend, we also observed an increase in total cellular PDH levels (Figures S5E and S5F). Together our results suggest that nuclear PDH levels increase upon growth in galactose media and the differentiation process but not on pyruvate supplementation.

Several stress conditions that lead to mitochondrial depolarization increase MDV numbers. We next investigated the nuclear PDH levels under various stress conditions relevant to cardiac cells like doxorubicin treatment, ischemia-reperfusion injury and hypoxia. Doxorubicin, a cytotoxic drug, induces cardiotoxicity by causing mitochondrial stress and

damaging the mitochondria [49]. The mitochondrial stress is initiated by production of
superoxide, hydroxyl radicals, peroxynitrite etc [50]. We treated H9C2 cells with 25 µM
doxorubicin [4] for 1 hour and 3 hours as significant cell death was observed only by 6 hours
(Figure S5K). Nuclear PDH decreased drastically at 1 hour and 3 hours treatment of
doxorubicin (Figures 5E and 5F). We confirmed that the total cellular PDH level remains
unchanged (Figures S5G and S5H). Interestingly, PDH[+] MDVs biogenesis also increased in
doxorubicin treatment in both 1 and 3 hours (Figures S5O and S5P). This result shows that
PDH[+] MDVs is an early response to doxorubicin-induced stress but these MDVs are not
targeting the nucleus to enable mito-nuclear PDH transit. Ischemia-reperfusion injury is one
of the major causes of heart failure. Ischemia is caused by hypoxia and nutrient deficiency,
followed by reperfusion through reoxygenation process. The reperfusion process produces
free radicals, which damages the mitochondria [51]. We observed a reduction in nuclear PDH
during ischemia and reperfusion compared to control conditions (Figure 5G). Our confocal
data showed that the number of PDH[+] MDVs has increased during reperfusion, but no
significant increment observed in ischemia condition (Figures S5Q and S5R). Here again the
generated MDVs do not target the nucleus but perhaps other cellular destinations with a
reduction in PDH levels in the nucleus. Probably the PDH already present in the nucleus is
turned over with no fresh entry through MDVs under these conditions.

Cobalt chloride is a chemical agent which helps to stabilize and accumulate HIF-1α to mimic
hypoxia conditions [52].   Decreased nuclear PDH was observed in HeLa cells upon
prolonged hypoxia [33]. To understand the effect of hypoxia on nuclear PDH levels in H9C2,
we treated the cells with 100 µM cobalt chloride ($CoCl_2$) to establish hypoxic conditions for
different time intervals. We verified HIF1α stabilization under these conditions (data not
shown). No cell death was observed even after 24 hours of $CoCl_2$ treatment (Figure S5L).
Nuclear transit of PDH decreased at 24 hours of $CoCl_2$ treatment (Figures 5H and 5I),
although total cellular PDH pool remained unchanged (Figures S5I and S5J).

Our findings suggest that PDH[+] MDVs increase under most of the mitochondrial stress
conditions as expected, but it is not targeted to the nucleus and the PDH levels in the nucleus
remain unchanged as with experiments in Figure 3 for Antimycin A or reduce as shown here.
However, the PDH[+] MDVs generated under basal physiological conditions that necessitate a
higher mitochondrial functionality like growth in galactose media or differentiation are
targeted to the nucleus.

Discussion

Our study investigates the retrograde communication pathway between mitochondria and
nucleus through mitochondria-derived vesicles (MDVs). We show that PDHE1 is transported
to the nucleus through MDVs. PDH is a large protein complex and we are currently
investigating if other components of the complex like PDHE2 and E3 are also trafficked to
the nucleus using MDVs to form a functional active complex. Our proteomics data identified
multiple subunits of PDC complex as NEM sensitive MDV cargo proteins. Further we
identified and validated COX4 as a novel MDV cargo protein.

Several mitochondrial proteins have moonlighting functions in the nucleus [24]. Multiple
TCA cycle and ETC enzymes were found in the nucleus [25]. It is reported that PDH is
present in the nucleus of different cell lines like A549, BaF3 and 3T3-L1 adipocyte cells [28,

42, 43]. We found PDH in the nucleus of rat heart and H9C2 cardiomyoblast lines whereas
trace amount of PDH is also present in HEK293T cell nucleus as reported previously [29].
Moreover, we identified COX4 as present in the nucleus of HEK293T cells but not the
cardiac cells suggesting cell type specificity in the process of mito-nuclear protein transit. The
roles of trafficked proteins in helping mito-nuclear communication and homeostasis are also
therefore likely to be cell type specific.

With our live cell imaging data, we show that PDH[+] MDV-like structures that are tracker
negative fuse with the nucleus but the fusion mechanism is still unknown. A previous study
has shown that the entire PDH complex enters the nucleus through the lamina following
MFN2 mediated direct mito-nuclear tethering [29]. However, in mature cardiac cells with
spheroid, relatively less motile mitochondria embedded within the myofibrils [53], direct
tethering might be difficult to achieve. We believe that in this context MDVs may mediate
the transit of proteins like PDH. Using a reconstitution approach, we observe that nuclear
PDH levels increase upon incubation with MDVs and this is sensitive to the presence of
NEM, the inhibitor of NSF and vesicle fusion process. Interestingly the endogenous nuclear
pools of NDUFS1 and VDAC do not change suggesting that their nuclear pools aren't
dependent on MDV mediated protein transit and some of these could be shuttled directly
from the cytosol through the nuclear pore complex. Our reconstitution results in tandem with
literature [28] suggests that while PDH requires mitochondrial entry to fold and form a
functional complex following the cleavage of its mitochondrial localization signal; this
enzyme further transits into the nucleus through MDVs to enable other moonlighting
functions to take place. While we envisage trafficking to be the likely mode of PDH transit in
our system our data does not rule out the possibility of protein transits across mito-nuclear
tethers reported recently [29, 54].

Our results indicate that PDH[+]MDV biogenesis is actin dependent (Figure 3). We also show
that like Tom20[+] MDVs which were reported to be microtubule dependent [14], PDH+ MDV
biogenesis also appears to be tubulin dependent. Actin mediates all endocytic budding events
[55] and it would be interesting to further study how actin mediated force generation helps in
MDV budding from the mitochondrial surface. Role of actin in MDV budding has not been
explored. We need to decouple budding versus transit to understand the role of cytoskeletal
elements in carrying MDVs to the nucleus.

Our preliminary observations suggest that global acetyl histone (H3K9/14Ac) levels in
cardiac cells do not parallel the increase or decrease in nuclear PDH (data not shown). Unlike
previous reports [27], this nuclear transit of PDH is also not stress driven (Figure 5) and
happens more predominantly in basal states with higher mitochondrial oxidative
phosphorylation. This is contrary to the observation on several mitochondrial dehydrogenases
including PDHE1 translocating to the nucleus post chemotherapy-induced cardiotoxicity
[27]. This study uses hiPSC-cardiomyocytes and further they are glucose deprived to get a
pure cardiomyocytes line eliminating other cell types. In contrast, our studies are on shorter
time points of mitochondrial stress in rat cardiomyoblasts.

We found that PDH[+] MDV biogenesis increases under most of the mitochondrial stress
conditions as reported previously [4], but this does not result in an increased nuclear pool of
PDH. Surprisingly in most cases there is a reduction in nuclear pools commiserate with
nuclear protein turnover once delivery of new cargo is stopped.   Previous literature

demonstrates syntaxin-17, a mitochondrial SNARE protein, associates with mature MDVs to
mediate MDV-endolysosome fusion [9]. This indicates that appropriate v-SNARES that
mediate MDV to nuclear membrane fusion have to be incorporated in the outer membrane of
the subpopulation of PDH MDVs that deliver PDH to the nucleus. PDH itself, depending on
cellular context, in Parkin dependent and independent manner localizes to different cellular
compartments like the lysosome or multivesicular bodies (MVB) respectively [11]. This
observation indicates that various destinations are plausible for PDH⁺ MDVs. We
hypothesized that stress induced PDH packaging into MDVs is probably destined to the
endolysosomal pathway and may be a quality control arm whereas basal trafficking to the
nucleus could be to enable other moonlighting functions which require further probing.

Further, we establish a novel mito-nuclear trafficking route which could be employed by
other mitochondrial proteins that have dual roles in both compartments but are functional
only after mitochondrial import. Since MDVs form an ancient vesicle compartment [19] we
envisage MDV mediated nuclear trafficking of proteins as a general means of the
endosymbiont communicating and controlling the host nuclear program.

Key Resource Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
ACP/NDUFAB1	Cell Signaling Technology	Cat#71814, RRID: AB_3086728
COX1	Cell Signaling Technology (CST)	Cat# 55159T
COX10	Cell Signaling Technology (CST)	Cat# 24744T
COX4 (3E11) Rabbit mAb	Cell Signaling Technology (CST)	Cat# 4850, RRID: AB_2085424
COX4 Polyclonal Antibody	Thermo Fisher Scientific Invitrogen	Cat# PA5-19471, RRID: AB_11004909
Cytochrome c (136F3) Rabbit mAb	Cell Signaling Technology (CST)	Cat# 4280, RRID: AB_10695410
Histone 3	Cell Signaling Technology (CST)	Cat# 9715
Histone 3	Santa Cruz	Cat# 517576
HSP60 (D6F1) XP® Rabbit mAb	Cell Signaling Technology (CST)	Cat# 12165, RRID: AB_2636980
OPA1 Mouse mAb	BD Biosciences	Cat# 612606, RRID: AB_399888

PHB1 Polyclonal Antibody	Cell Signaling Technology (CST)	Cat# 2426, RRID: AB_823689
Pyruvate Dehydrogenase (C54G1) Rabbit mAb	Cell Signaling Technology (CST)	Cat# 3205, RRID: AB_2162926
SDHA (D6J9M) Rabbit mAb	Cell Signaling Technology (CST)	Cat# 11998, RRID: AB_2750900
SOD1 (71G8) Mouse mAb	Cell Signaling Technology (CST)	Cat# 4266, RRID: AB_2193898
TOM20 (D8T4N) Rabbit mAb	Cell Signaling Technology (CST)	Cat#42406, RRID: AB_2799825
Tubulin	Merck	Cat#AB9354
UQCRFS1/RISP	Cell Signaling Technology (CST)	Cat# 95231, RRID: AB_2922784
VDAC (D73D12) Rabbit mAb	Cell Signaling Technology (CST)	Cat#4661, RRID: AB_10557420
Chemicals, peptides, and recombinant proteins		
Acetone	Himedia	MB179
Acrylamide	BIO RAD	161-0101
2 mercaptoethanol	Himedia	MB041
ADP disodium salt	Himedia	MB191
Ammonium persulphate	Himedia	MB003
Antibiotic Antimycotic solution	Sigma Aldrich	A5955
Antimycin A	Sigma	A8674
ATP disodium salt	Himedia	RM439
Bis-acrylamide	Himedia	MB005
Bromophenol blue	Himedia	MB123
Cobalt chloride	Merck	S4321140
D- Glucose	Sigma Aldrich	G8270
DAPI	Himedia	MB097
D-Galactose	Sigma Aldrich	G0750
Dipotassium hydrogen phosphate	Himedia	TC596
di-Sodium hydrogen phosphate dodecahydrate	Himedia	GRM1418
DMEM low glucose Medium	Himedia	AL006A
DMEM Medium	Sigma Aldrich	D0547
DMSO	Sigma Aldrich	D2438
Dodecyl sulphate sodium salt	Himedia	TC698
Doxorubicin	Sigma Aldrich	44583
EGTA	Himedia	GRM1530
EM grade glutaraldehyde	Merck	G5882
Endothelial cell growth medium	Sigma Aldrich	211-500
Fetal Bovine Serum	Gibco	10270106
Fluoromount G	Invitrogen	4958-02
Formvar/Carbon 200 Mesh, Gold grid	Electron Microscopy	FCF200-AU25

	Sciences	
Glycerol	Himedia	TC503
Glycine	Himedia	MB013
HEPES	Himedia	MB016
Hydrocortisone	Sigma Aldrich	H0888
Insulin	Sigma Aldrich	I9278
Latrunculin B	Sigma Aldrich	428020
Lipofectamine 2000	Invitrogen	11668030
M-199 Medium	Sigma Aldrich	M0393
Magnesium acetate	Himedia	MB039
Mannitol	Himedia	MB198
Mdivi	Sigma Aldrich	M1099
Methanol	Thermo Fisher Qualigens	Q43637G
MitoTracker	Cell Signaling Technology (CST)	8778S
N-Ethylmaleimide	Sigma	E3876
Nocodazole	Sigma Aldrich	M1404
NP-40 solution	Himedia	MB143
Opti-MEM Medium	Gibco	31985070
Phosphotungstic acid hydrate	Sigma Aldrich	79690
Potassium chloride	Himedia	GRM3934
Potassium dihydrogen orthophosphate monobasic	Himedia	GRM3943
Retinoic acid	Sigma Aldrich	R2625
Sodium chloride	Himedia	GRM3954
Sodium pyruvate	Sigma Aldrich	S8636
Sodium succinate	Himedia	GRM418
Sucrose	Himedia, Sigma	PCT0616, S0389
TEMED	BIO RAD	161-0800
Tris-Base	BIO RAD, Himedia	1610719, TC072
Trypsin-EDTA solution	Sigma Aldrich	T4174
Tween-20	Himedia	TC287
Recombinant DNA		
pLV[Exp]-Puro-CMV>hPDHA1[NM_001173454.2]/mCherry	Vector Builder	VB221211-1356azz
Experimental models: Organisms/strains		
Rat (Wistar)		
Software and algorithms		
Fiji	NIH	https://fiji.sc/

| GraphPad Prism | GraphPad Software Inc | http://www.graphpad.com/ |

Materials and Methods

Cell culture and treatments

Rat embryonic cardiomyoblasts (H9C2), Human kidney cells (HEK293T) (ATCC) and Human umbilical vein endothelial cells (HUVEC) (Promocell) were used in this study. H9C2 cardiomyocytes and HEK293T cells were maintained in Dulbecco's modified eagle medium (DMEM) (Sigma-Aldrich) supplemented with 10% FBS (Gibco) and 1% antibiotic antimycotic solution (Sigma-Aldrich). The H9C2 cardiomyoblasts were additionally supplemented with 1 mM sodium pyruvate except in experiments where the role of pyruvate is tested (Sigma-Aldrich). HUVEC was maintained in endothelial cell growth media (Sigma-Aldrich) supplemented with 2% FBS and 2% antibiotic antimycotic solution (Sigma-Aldrich).

All cell culture chemicals are from Sigma Aldrich unless otherwise indicated. Please find details in the key resource table. Cell treatments are indicated in the respective figures, as appropriate, and were: antimycin A (20 µM, Sigma-Aldrich, 1h), latrunculin B (1 µM, 15 min), nocodazole (10 µM, 30 min), doxorubicin (25 µM, 1h, 3h), cobalt chloride (100 µM, Merck, 1h-24h). Appropriate time matched controls were set for every treatment condition. To investigate the effect of galactose, H9C2 cardiomyoblasts were cultured in DMEM low glucose (5 mM glucose, Himedia) with galactose (20 mM) for 24 hours.

Lentiviral transduction

To prepare stable cell line overexpressing PDH mCherry, initially viral particles were produced in HEK293T. HEK293T was transfected with pLVX PDH mCherry and packaging vectors, pMDG2 and pAX2 using lipofectamine 2000. After 48 hours, viral supernatant was added to H9C2 for 4 hours. Cells were selected for stable expression using puromycin (1.5 µg/ml).

H9C2 cardiomyoblast differentiation

H9C2 cardiomyoblasts were differentiated using retinoic acid in low serum containing media [56]. All-trans retinoic acid (Sigma-Aldrich) was prepared in DMSO and stored in dark at -20°C. A total of 81,000 cells for non-differentiation and 3,00,000 cells for differentiation were plated in 60 mm tissue culture dish and cultured one day in 10% FBS containing media in order to adhere cells. Non-differentiated cell plate was maintained in 10% FBS containing media. For differentiation, cells were cultured with 1µM all-trans retinoic acid in low serum containing media (1% FBS) for 5 days. On the 6th day, cells were harvested for nuclei isolation or whole cell lysate preparation following described procedures.

Ischemia-Reperfusion induction

H9C2 cardiomyoblasts were seeded in 100 mm tissue culture dishes. For ischemia conditions, cells were incubated with complete media in a 5% CO2 incubator for 2 hours, followed by incubation in hypoxia chamber (New Brunswick Galaxy 48R) with glucose free and serum free media for 2 hours. To establish reperfusion conditions, cells were incubated in hypoxia conditions as mentioned above, followed by incubation in a regular 5% CO_2 incubator with complete media for 2 hours. Post-treatment, cells were subjected to nucleus isolation following the described procedures.

Isolation of MDVs

The protocol followed as well as appropriate controls is detailed in Nair et al., 2024 [40]. Cardiac MDVs were budded from mitochondria isolated from rat hearts.

The budding reaction solution was constituted of 1 mM ATP, 5 mM succinate, 80 μM ADP and 2 mM K_2HPO_4, in 1X MIB. A parallel reaction containing NEM (20 mM) was also set up. Budding reactions were incubated at 37°C for 2 hours in the incubator with shaking. The budding reactions were centrifuged at 10,000g for 10 minutes and the supernatant containing the MDVs was collected.

To further purify the vesicles, MDVs were floated by sucrose gradient centrifugation (Beckman Coulter - Sw41Ti rotor). Samples were loaded at the bottom of the discontinuous sucrose gradient, by loading 0.5 ml of sample to 1.5 ml of 80% sucrose, resulting in 60% sucrose. The gradient from the bottom was as follows: 60%, 50%, 40%, 30%, 20%, and 10 % (2ml each). Samples were centrifuged at 32,500 rpm (100,000g) for 16 hours at 4°C and 1 ml of the fractions were collected from the top.

Proteomics analysis

All budding reactions were concentrated using 3kDa Amicon filter column to 50 μL and the concentrated samples were loaded on SDS PAGE. The samples were cut from gel upon reaching the interface between the separating gel and the resolving gel. The gel piece was destained and 100% acetonitrile solution was used to dry the gel. This was followed by overnight digestion with 13 ng/μL trypsin in 50 mM ammonium bicarbonate. 5% formic acid/acetonitrile (1:2, v/v) extraction buffer was used to extract peptide from the gel. Finally, the samples were reconstituted in 3% acetonitrile.

A nano ACQUITY UPLC System (Waters) chromatography system was used for the LC analysis for the first two sets of samples. The Mass Lynx 4.1 SCN781 software was used for the acquisition. The peptides were separated by reversed-phase chromatography. MS analysis of eluting peptides was carried out on a SYNAPT® G2 High-Definition MS™ system (HDMS[E] System) (Waters). The analyses were carried out in positive mode electrospray ionization (ESI). The third set of samples were analysed using Orbitrap Eclipse. Xcalibur software was used for data acquisition. Protein identification was done by analysing LC-MS data using Progenesis QI for Proteomics V4.2 (Waters) for the initial two MS runs and Proteome Discoverer v2.5 was used for the last MS run. All three runs were separated on a

binary phase system: the mobile phase-A was water, and the mobile phase-B was acetonitrile; both contain 0.1% formic acid. The proteins were identified from the *Rattus norvegicus* database on UniProt.

For further analysis, only proteins with at least one unique peptide were considered. Mitochondrial localization of the protein was established with the help of MitoCarta 3.0 and The Integrated Mitochondrial Protein Index (IMPI) database. The sublocalization of these proteins within the mitochondria was established with the help of these databases. In addition to that a literature survey done by us gave evidence for the mitochondrial sublocalization. To identify the high confidence NEM sensitive MDV cargo, we compared the control proteins with the NEM proteins, and the proteins detected only in the control were considered. A Reactome pathway analysis of the confirmed MDV cargo was performed. Our MDV proteome was compared to the Reactome database including the interaction data from IntAct.

TEM imaging

The budding reactions were loaded onto gold Formvar carbon coated TEM grid (FCF200-Au; Electron Microscopy Sciences) and incubated overnight at 4°C. To fix, 2.5% glutaraldehyde was added for 30s and negative staining was done using uranyl acetate for 1 min. Images were acquired using JEM-1011 electron microscope (JEOL) using Digital Micrograph software.

Isolation of Nucleus from Cells

Cells were grown on a 100 mm tissue culture dish for nucleus isolation. Cells were washed with ice-cold 1X PBS and scrapped in 0.1% NP-40 solution by the help of cell scrapper lysing only the plasma membrane. The sample was washed 2 times with 0.1% NP-40 solution by centrifugation at 700g at 4°C [57]. Then samples were loaded on top of the 60% sucrose cushion and processed for ultracentrifugation (Beckman Coulter - Sw41Ti rotor) at 30,000g for 1 hour at 4°C. After ultracentrifugation, the nuclear pellet was resuspended in Laemmli sample buffer and ultrasonicated to shear the genomic DNA before loading for western blots.

Isolation of nucleus from rat heart

Nucleus isolation from rat heart was adapted from previously established protocols [57, 58]. Rat hearts were collected in ice-cold PBS. The hearts were chopped and followed by centrifugation at 3000g for 10 min at 4°C. Supernatant was discarded and the pellet was washed in PBS by centrifugation. The pellet was resuspended in isolation buffer (Mannitol 220 mM sucrose 68 mM; KCl 80 mM ; EGTA 0.5 mM; Mg(C□H□O□)□ 2 mM; HEPES, pH 7.4 20 mM) and homogenized using a 50 ml Dounce tissue homogenizer, followed by centrifugation at 3000g for 10 min at 4°C. The pellet was resuspended in isolation buffer containing 0.1% NP-40 and passed through a 40 µm cell strainer (Himedia). The filtered suspension was washed by centrifugation at 500g for 10 min at 4°C. The resuspended pellet was loaded on a 60% sucrose cushion. This was followed by ultracentrifugation using swinging bucket rotor (Beckman Coulter - Sw41Ti rotor) at 30,000g for 1 hour at 4°C. The

nuclear pellet was resuspended in 200 µL of 1X Laemmli buffer for western blotting or 200
µL of MIB for reconstitution experiment. Whole tissue lysate was prepared from rat heart
tissue using 1X Laemmli buffer followed by sonication.

Immunoblotting

All the samples were prepared in 1X Laemmli buffer, sonicated and boiled at 95°C for 5 min.
Following separation of proteins by SDS-PAGE, the proteins were transferred to
nitrocellulose membrane. Ponceau S staining was done to determine efficient transfer and
equal load. The blots were immunoblotted with respective primary and secondary antibodies.
The primary antibodies used were anti-VDAC (1:1000) (CST 4661T), anti-TOM20 (1:1000)
(CST 42406S), anti-OPA1 (1:2000) (BD Biosciences cat 612606), anti-PDH (1:1000) (CST
3205S), anti-SDHA (1:1000) (CST 11998T), anti-COX4 (1:1000) (CST 3E11 4850S), anti-
COX4 (1:2000) (Invitrogen PA5-19471), anti-PHB1 (1:1000) (CST 2426T), anti-cytochrome
c (1:1000) (CST 4280t), anti HSP60 (1:1000) (CST), anti SOD1 (1:1000) (CST 4266T), anti-
histone 3 (1: 5000) (Santa Cruz sc517576), anti-histone 3 (1:5000) (CST 9715), anti-β-
tubulin (1:2000) (Merck), anti-NDUFS1 (1:1000) (CST 70264T), anti-COX10 (1:1000) (CST
24744T), anti-COX1 (1:1000) (CST 55159T), anti-ACP/NDUFAB1 (1:1000) (CST
71814T), anti-UQCRFS1/RISP (1:1000) (CST 95231T). The secondary antibodies are Goat
anti-rabbit (1:10,000) (CST) and Goat anti-mouse (1:10,000) (CST). The protein bands were
detected using ECL solution and developed either X ray or on a Chemidoc machine
(ImageQuant LAS500)

Antimycin A washout assay

70,000 H9C2 cardiomyoblasts were seeded on coverslips in 35 mm dishes. All the cells were
pre-treated with Drp1 inhibitor, Mdivi (Sigma-Aldrich) (1µM) for 1 hour and Mdivi is
present throughout the experiment. The control cells were treated with DMSO for 1 h while
the treatment cells were treated with antimycin A (20 µM) for 1 h. After treatment the
antimycin containing media was removed and Mdivi containing media was added. The
treated cells were fixed at 0 min, 30 min, 60 min, 90 min and 120 min after antimycin
washout. The cells were fixed with ice-cold methanol-acetone (1:1) for 5 min at -20°C.

Immunofluorescence

Cells were grown on coverslip and treated according to each experiment. Post-treatment
500nM MitoTracker (CST) was added to stain mitochondria and incubated at 37°C for 30
minutes. Cells were washed 3 times with 1X PBS. In order to fix the cells, ice-cold 100%
methanol was added and incubated in -20°C for 15 minutes. Then cells were washed 3 times
with 1X PBS and blocked with 1% BSA solution at room temperature for 1 hour. Cells were
incubated with primary antibodies for 1 hour at room temperature and subsequent 3 washes
with 1X PBS. The primary antibodies used were anti-PDH E1-α (1:25) (Santacruz, sc-
377092), anti-COX4 (1:100) (CST 3E11 4850S) and anti-tubulin (1:200) (). Then, cells were
incubated with respective Alexa Fluor secondary antibodies (Jackson Immunoresearch) for 1

hour at room temperature in dark, followed by 3 washes with 1X PBS. Actin was stained with
phalloidin (1:400) (CST). Cells were counterstained with DAPI (Himedia) (1:10, 000
dilution) for 5 minutes at room temperature, followed by 5 times washing with 1X PBS.
Finally, coverslips were mounted with Fluoromount G (Invitrogen).

Imaging

Confocal images (~0.3 μm step size) were acquired with Olympus Fluoview FV3000 60X,
Oil. The excitation wavelengths used were 405, 488, 561 and 640 nm. Image files were
analysed using ImageJ.

For live imaging, H9C2 cells overexpressing PDH-mCherry were seeded on a 35mm glass
bottom dish. Cells were tracked with MitoTracker Far Red (500nM) for 30 min prior to
imaging followed by three PBS washes. The nucleus was stained with Hoechst followed by
five PBS washes. Cells were imaged at 37°C in a chamber with 5% CO_2. Images were
acquired with Olympus Fluoview FV3000.

In vitro reconstitution

To study cargo delivery into nucleus by MDVs, we established cell free reconstitution
reactions between nucleus and MDVs. Nucleus and cardiac MDVs were isolated in parallel
from rat heart as described above. We reconstituted the nucleus and MDVs in the presence
and absence of NEM (20 mM). A nucleus only reaction was set up as a control. All the
reactions occurred in the presence of 0.5 μM GTP (Sigma-Aldrich). The reactions were
incubated at 37°C for 1h in a shaking water bath. The nucleus was re-isolated by
ultracentrifugation and the samples were prepared in 1X Laemmli buffer.

Quantification and Statistical Analysis

For western blots, band intensities were quantified using ImageJ software. All statistical
analyses were performed using GraphPad Prism software (version 8.4.3). Student t-test was
performed to compare fold changes against the normalized control to compare two samples.
Ordinary one-way ANOVA or repeated measure one-way ANOVA followed by Bonferroni
post-hoc analysis was performed to analyze a single independent variable with three or more
samples. Error bar is reported as mean ± SEM. ****p-value< 0.0001, ***p-value< 0.001,
**p-value< 0.01, *p-value< 0.0 5.

References

1. Thapa, D., et al., *Adropin regulates pyruvate dehydrogenase in cardiac cells via a novel GPCR-MAPK-PDK4 signaling pathway.* Redox Biol, 2018. **18**: p. 25-32.
2. Cadete, V.J.J., et al., *Mitochondrial quality control in the cardiac system: An integrative view.* Biochim Biophys Acta Mol Basis Dis, 2019. **1865**(4): p. 782-796.
3. Soubannier, V., et al., *A vesicular transport pathway shuttles cargo from mitochondria to lysosomes.* Curr Biol, 2012. **22**(2): p. 135-41.

4.   Cadete, V.J., et al., *Formation of mitochondrial-derived vesicles is an active and physiologically relevant mitochondrial quality control process in the cardiac system.* J Physiol, 2016. **594**(18): p. 5343-62.

5.   Dorn, G.W., 2nd, *Mitochondrial dynamism and heart disease: changing shape and shaping change.* EMBO Mol Med, 2015. **7**(7): p. 865-77.

6.   Neuspiel, M., et al., *Cargo-selected transport from the mitochondria to peroxisomes is mediated by vesicular carriers.* Curr Biol, 2008. **18**(2): p. 102-8.

7.   Braschi, E., et al., *Vps35 mediates vesicle transport between the mitochondria and peroxisomes.* Curr Biol, 2010. **20**(14): p. 1310-5.

8.   McLelland, G.L., et al., *Parkin and PINK1 function in a vesicular trafficking pathway regulating mitochondrial quality control.* Embo j, 2014. **33**(4): p. 282-95.

9.   McLelland, G.L., et al., *Syntaxin-17 delivers PINK1/parkin-dependent mitochondrial vesicles to the endolysosomal system.* J Cell Biol, 2016. **214**(3): p. 275-91.

10.   Matheoud, D., et al., *Parkinson's Disease-Related Proteins PINK1 and Parkin Repress Mitochondrial Antigen Presentation.* Cell, 2016. **166**(2): p. 314-327.

11.   Todkar, K., et al., *Selective packaging of mitochondrial proteins into extracellular vesicles prevents the release of mitochondrial DAMPs.* Nat Commun, 2021. **12**(1): p. 1971.

12.   Ryan, T.A., et al., *Tollip coordinates Parkin-dependent trafficking of mitochondrial-derived vesicles.* Embo j, 2020. **39**(11): p. e102539.

13.   Juhász, G., *A mitochondrial-derived vesicle HOPS to endolysosomes using Syntaxin-17.* J Cell Biol, 2016. **214**(3): p. 241-3.

14.   König, T., et al., *MIROs and DRP1 drive mitochondrial-derived vesicle biogenesis and promote quality control.* Nat Cell Biol, 2021. **23**(12): p. 1271-1286.

15.   Abuaita, B.H., T.L. Schultz, and M.X. O'Riordan, *Mitochondria-Derived Vesicles Deliver Antimicrobial Reactive Oxygen Species to Control Phagosome-Localized Staphylococcus aureus.* Cell Host Microbe, 2018. **24**(5): p. 625-636.e5.

16.   Hazan Ben-Menachem, R., et al., *Mitochondrial-derived vesicles retain membrane potential and contain a functional ATP synthase.* EMBO Rep, 2023. **24**(5): p. 17.

17.   Andrade-Navarro, M.A., L. Sanchez-Pulido, and H.M. McBride, *Mitochondrial vesicles: an ancient process providing new links to peroxisomes.* Curr Opin Cell Biol, 2009. **21**(4): p. 560-7.

18.   Gould, S.B., S.G. Garg, and W.F. Martin, *Bacterial Vesicle Secretion and the Evolutionary Origin of the Eukaryotic Endomembrane System.* Trends Microbiol, 2016. **24**(7): p. 525-534.

19.   König, T. and H.M. McBride, *Mitochondrial-derived vesicles in metabolism, disease, and aging.* Cell Metab, 2024. **36**(1): p. 21-35.

20.   Lo, S.C. and M. Hannink, *PGAM5 tethers a ternary complex containing Keap1 and Nrf2 to mitochondria.* Exp Cell Res, 2008. **314**(8): p. 1789-803.

21.   Monaghan, R.M., et al., *A nuclear role for the respiratory enzyme CLK-1 in regulating mitochondrial stress responses and longevity.* Nat Cell Biol, 2015. **17**(6): p. 782-92.

22.   Yogev, O., et al., *Fumarase: a mitochondrial metabolic enzyme and a cytosolic/nuclear component of the DNA damage response.* PLoS Biol, 2010. **8**(3): p. e1000328.

23.   Gibellini, L., et al., *Evidence for mitochondrial Lonp1 expression in the nucleus.* Sci Rep, 2022. **12**(1): p. 10877.

24.   Monaghan, R.M. and A.J. Whitmarsh, *Mitochondrial Proteins Moonlighting in the Nucleus.* Trends Biochem Sci, 2015. **40**(12): p. 728-735.

25. Nagaraj, R., et al., *Nuclear Localization of Mitochondrial TCA Cycle Enzymes as a Critical Step in Mammalian Zygotic Genome Activation.* Cell, 2017. **168**(1-2): p. 210-223.e11.

26. Li, W., et al., *Nuclear localization of mitochondrial TCA cycle enzymes modulates pluripotency via histone acetylation.* Nat Commun, 2022. **13**(1): p. 7414.

27. Srivastava, S., et al., *Nuclear translocation of mitochondrial dehydrogenases as an adaptive cardioprotective mechanism.* Nat Commun, 2023. **14**(1): p. 023-40084.

28. Sutendra, G., et al., *A nuclear pyruvate dehydrogenase complex is important for the generation of acetyl-CoA and histone acetylation.* Cell, 2014. **158**(1): p. 84-97.

29. Zervopoulos, S.D., et al., *MFN2-driven mitochondria-to-nucleus tethering allows a non-canonical nuclear entry pathway of the mitochondrial pyruvate dehydrogenase complex.* Mol Cell, 2022. **82**(5): p. 1066-1077.e7.

30. Soubannier, V., et al., *Reconstitution of mitochondria derived vesicle formation demonstrates selective enrichment of oxidized cargo.* PLoS One, 2012. **7**(12): p. e52830.

31. Lee, J., et al., *The plasticity of the pyruvate dehydrogenase complex confers a labile structure that is associated with its catalytic activity.* PLoS One, 2020. **15**(12): p. e0243489.

32. Patel, M.S., et al., *The pyruvate dehydrogenase complexes: structure-based function and regulation.* J Biol Chem, 2014. **289**(24): p. 16615-23.

33. Eguchi, K. and K. Nakayama, *Prolonged hypoxia decreases nuclear pyruvate dehydrogenase complex and regulates the gene expression.* Biochem Biophys Res Commun, 2019. **520**(1): p. 128-135.

34. Chen, J., et al., *Compartmentalized activities of the pyruvate dehydrogenase complex sustain lipogenesis in prostate cancer.* Nat Genet, 2018. **50**(2): p. 219-228.

35. Mocholi, E., et al., *Pyruvate metabolism controls chromatin remodeling during CD4(+) T cell activation.* Cell Rep, 2023. **42**(6): p. 112583.

36. Vasam, G., et al., *Proteomics characterization of mitochondrial-derived vesicles under oxidative stress.* Faseb j, 2021. **35**(4): p. e21278.

37. Towers, C.G., et al., *Mitochondrial-derived vesicles compensate for loss of LC3-mediated mitophagy.* Dev Cell, 2021. **56**(14): p. 2029-2042.e5.

38. Roberts, R.F., et al., *Proteomic Profiling of Mitochondrial-Derived Vesicles in Brain Reveals Enrichment of Respiratory Complex Sub-assemblies and Small TIM Chaperones.* J Proteome Res, 2021. **20**(1): p. 506-517.

39. Weidman, P.J., et al., *Binding of an N-ethylmaleimide-sensitive fusion protein to Golgi membranes requires both a soluble protein(s) and an integral membrane receptor.* J Cell Biol, 1989. **108**(5): p. 1589-96.

40. Nidhi, N., et al., *In Vitro Budding and Floatation based Enrichment of Mitochondria-derived Vesicles for Proteomics from Rat Heart.* bioRxiv, 2024: p. 2024.03.04.583262.

41. Wang, S., et al., *Prohibitin co-localizes with Rb in the nucleus and recruits N-CoR and HDAC1 for transcriptional repression.* Oncogene, 2002. **21**(55): p. 8388-96.

42. Chueh, F.Y., et al., *Nuclear localization of pyruvate dehydrogenase complex-E2 (PDC-E2), a mitochondrial enzyme, and its role in signal transducer and activator of transcription 5 (STAT5)-dependent gene transcription.* Cell Signal, 2011. **23**(7): p. 1170-8.

43. Richard, A.J., H. Hang, and J.M. Stephens, *Pyruvate dehydrogenase complex (PDC) subunits moonlight as interaction partners of phosphorylated STAT5 in adipocytes and adipose tissue.* J Biol Chem, 2017. **292**(48): p. 19733-19742.

44. Wu, X.S., et al., *Actin Is Crucial for All Kinetically Distinguishable Forms of Endocytosis at Synapses.* Neuron, 2016. **92**(5): p. 1020-1035.

45. Schuh, M., *An actin-dependent mechanism for long-range vesicle transport.* Nat Cell Biol, 2011. **13**(12): p. 1431-6.

46. Aguer, C., et al., *Galactose enhances oxidative metabolism and reveals mitochondrial dysfunction in human primary muscle cells.* PLoS One, 2011. **6**(12): p. e28536.

47. Kase, E.T., et al., *Remodeling of oxidative energy metabolism by galactose improves glucose handling and metabolic switching in human skeletal muscle cells.* PLoS One, 2013. **8**(4): p. e59972.

48. Comelli, M., et al., *Cardiac differentiation promotes mitochondria development and ameliorates oxidative capacity in H9c2 cardiomyoblasts.* Mitochondrion, 2011. **11**(2): p. 315-26.

49. Osataphan, N., et al., *Effects of doxorubicin-induced cardiotoxicity on cardiac mitochondrial dynamics and mitochondrial function: Insights for future interventions.* J Cell Mol Med, 2020. **24**(12): p. 6534-6557.

50. Wu, L., et al., *Mitochondrial quality control mechanisms as therapeutic targets in doxorubicin-induced cardiotoxicity.* Trends Pharmacol Sci, 2023. **44**(1): p. 34-49.

51. Pedriali, G., et al., *Perspectives on mitochondrial relevance in cardiac ischemia/reperfusion injury.* Front Cell Dev Biol, 2022. **10**: p. 1082095.

52. Prajapat, S.K., K.C. Maharana, and S. Singh, *Mitochondrial dysfunction in the pathogenesis of endothelial dysfunction.* Mol Cell Biochem, 2023.

53. Tepp, K., et al., *High efficiency of energy flux controls within mitochondrial interactosome in cardiac intracellular energetic units.* Biochim Biophys Acta, 2011. **1807**(12): p. 1549-61.

54. Desai, R., et al., *Mitochondria form contact sites with the nucleus to couple prosurvival retrograde response.* Sci Adv, 2020. **6**(51).

55. Galletta, B.J. and J.A. Cooper, *Actin and endocytosis: mechanisms and phylogeny.* Curr Opin Cell Biol, 2009. **21**(1): p. 20-7.

56. Branco, A.F., et al., *Gene Expression Profiling of H9c2 Myoblast Differentiation towards a Cardiac-Like Phenotype.* PLoS One, 2015. **10**(6): p. e0129303.

57. Nabbi, A. and K. Riabowol, *Rapid Isolation of Nuclei from Cells In Vitro.* Cold Spring Harb Protoc, 2015. **2015**(8): p. 769-72.

58. Bergmann, O. and S. Jovinge, *Isolation of cardiomyocyte nuclei from post-mortem tissue.* J Vis Exp, 2012(65).

Ethics statement

All animal experiments were performed after valid ethical clearance from the Institutional (RGCB) Animal Bioethics Committee (IAEC/867/ANANTHA/2021 and IAEC/940/ANANTHA/2023) which is registered under Committee for the Purpose of Control and Supervision of Experiments on Animals (CPCSEA), Govt. of India [326/GO/ReBiBt/S/2001/CPCSEA].

Data Availability

All relevant data supporting the key findings of this study are availa

ble within the article and the supplementary data file S1 and S2. All data reported in this paper will be shared by lead contact/ corresponding author (AS) upon request.

Acknowledgements

This work is supported by the Ramalingaswami Re-entry Fellowship (BT/RLF/Re-entry/51/2019) and SERB POWER grant (SPG/2021/003267) awarded to AS by the Department of Biotechnology (DBT) and Science and Engineering Research Board, Department of Science and Technology (DST), Government of India respectively, and JRF fellowship from University Grants Commission to TR. Authors are grateful to the confocal, proteomics and animal research facilities as well as the Instrumentation Department at Rajiv Gandhi Centre for Biotechnology (RGCB) for their help with experiments.

Author contributions

Conceptualization, A.S. and T.R.

Methodology T.R., D.S., S.B., L.R., V.S.I, N.N, A.M., M.J., S.G., J.B.J., A.S.

Investigation T.R., D.S., S.B., L.R., V.S.I, N.N, A.M., M.J., S.G., J.B.J., A.S.

Writing – Original Draft, A.S, T.R, D.S.,

Writing – Review & Editing, all authors

Funding Acquisition A.S.

Supervision A.S.

Conflict of interest

The authors declare no competing interests.

Main Figure legends

Figure 1: Cardiac MDV proteome was enriched for mitochondrial proteins involved in TCA cycle and electron transport chain.

(A) Schematic of the MDV isolation protocol, created with BioRender.com. Mitochondria isolated from rat heart were budded in a budding reaction buffer (composition is as described in methods) for 2h at 37°C. In parallel reactions, the budding was performed in the presence or absence of vesicular transport inhibitor, N-ethylmaleimide (NEM: 20mM). The mitochondria was pelleted and the supernatant containing the MDVs was used for downstream experiments.

(B) TEM imaging of the budding reactions performed in the presence or absence of NEM showed vesicular structures. Scale bar shows 100 nm.

(C) Immunoblot of the budding reactions generated in the presence or absence of NEM. Equal volumes of the budding reactions was loaded and were probed for outer mitochondrial membrane proteins, Voltage-dependent anion channel (VDAC) and Translocase of outer mitochondrial membrane 20 (TOM20); inner mitochondrial membrane and intermembrane space protein, Optic atrophy 1 (OPA1) and mitochondrial matrix protein, Pyruvate dehydrogenase E1 (PDHE1). Mt. pellet: mitochondria pellet.

(D) Immunoblot of fractions collected after sucrose density gradient (10-60%) ultracentrifugation of budding reaction. Blot was probed for VDAC, TOM20, OPA1 and PDHE1. Mt. pellet: mitochondria pellet.

(E) Venn diagram showing the overlap of the mitochondrial proteins detected in the three MS runs.

(F) Venn diagram showing the overlap of the NEM sensitive proteins.

(G) List of the NEM sensitive proteins. These proteins appeared at least twice in three independent MS runs.

Figure 2: PDH, but not COX4, is found in the nucleus of rat cardiac cells.

(A) Immunoblot of the nuclei isolated from HEK293T. The blot was probed for mitochondrial proteins(VDAC, TOM20, COX4, Prohibitin 1, OPA1, Cytochrome, SDH, PDHE1, HSP60, SOD1); cytoplasmic marker, Tubulin and nuclear marker, Histone 3. WCL: whole cell lysate.

(B) Immunoblot of the nuclei isolated from H9C2 cardiomyoblasts. The blot was probed for mitochondrial proteins(VDAC, TOM20, OPA1, SDH, PDHE1, HSP60) and Histone 3. WCL: whole cell lysate.

(C) Immunoblot of the nuclei isolated from rat heart. The blot was probed for mitochondrial proteins(VDAC, NDUFS1, OPA1, PDHE1) and Histone 3. WTL: whole tissue lysate

Figure 3: Actin and tubulin depolymerization leads to decreased PDH⁺ MDVs.

(A) H9C2 rat cardiomyoblasts were pre-treated with MitoTracker (500nM) for 30 min. Subsequently the cells were treated with vehicle (NT), AA (20µM) and Lat B (1µM) with AA. The cells were fixed after 15 min with 100% methanol at -20°C. The cells were immunolabeled for PDH E1-α (green) and the nucleus was stained with DAPI (blue). MitoTracker appears red. The arrowhead indicates MDVs. Scale bar:10 µm (cell), 1 µm (inset). (B) Quantification of PDH⁺ MDVs (N=3, 5 cells/condition). One-way ANOVA followed by Bonferroni post-hoc analysis, ****p-value< 0.0001, **p-value< 0.01. Error bars, SEM.

(C) H9C2 rat cardiomyoblasts were pre-treated with MitoTracker (500nM) for 30 min. Subsequently the cells were treated with vehicle (NT), AA (20µM) and Noc (10µM) with AA. The cells were fixed after 30 min with 100% methanol at -20°C. The cells were immunolabeled for PDH E1-α (green) and the nucleus was stained with DAPI (blue). MitoTracker appears red. The arrowhead indicates MDVs. Scale bar:10 µm (cell), 1 µm (inset). (D) Quantification of PDH⁺ MDVs (N=3, 5 cells/condition). One-way ANOVA followed by Bonferroni post-hoc analysis, ****p-value< 0.0001, ***p-value< 0.001. Error bars, SEM.

Figure 4: PDH delivery to the nucleus is NEM sensitive.

(A) Schematic of the *in vitro* reconstitution between rat cardiac MDVs and rat nuclei, created with BioRender.com. Cardiac MDVs and nuclei were incubated together *in vitro* for 1 hour at 37°C. Three parallel reconstitution conditions were set up: nucleus only; nucleus and cardiac MDVs; and nucleus and MDVs in the presence of NEM (20mM). The nuclei were subsequently isolated by ultracentrifugation.

(B) Immunoblot of the nuclei isolated after reconstitution as described in (A). The whole tissue lysate (WTL) was prepared from rat heart tissue using Laemmli buffer followed by sonication. The blot was probed OPA1, PDHE1, COX4 and Histone 3 (C) Quantification of nuclear PDH after reconstitution. Nuclear PDH was normalized to Histone 3. One-way ANOVA followed by Bonferroni post-hoc analysis, ***p-value< 0.001, **p-value< 0.01. Error bars, SEM.

Figure 5: Regulation of nuclear PDH in basal and pathophysiological conditions.

(A,B) H9C2 cardiomyoblasts were cultured in 25 mM glucose and 20 mM galactose-containing media (with a low 5 mM glucose) in the presence or absence of pyruvate (1 mM) for 24 hours; (A) Immunoblot of nuclear PDHE1, (B) Quantification of nuclear PDH (n=3). Nuclear PDH was normalized to Histone 3. WCL: Whole cell lysate, Glu: Glucose, Gal: Galactose. RM one-way ANOVA followed by Bonferroni post-hoc analysis, *p-value< 0.05, ns: non-significant. Error bars, SEM.

(C,D) H9C2 cardiomyoblasts were differentiated into cardiomyocytes using retinoic acid (1 μM) for 5 days;(C) Immunoblot of nuclear PDHE1, (D) Quantification of nuclear PDH (n=5). Nuclear PDH was normalized to Histone 3, WCL: Whole cell lysate. Paired t-test, **p-value< 0.01. Error bars, SEM.

(E,F) H9C2 cardiomyoblasts were treated with Doxorubicin (25 μM) for 1 hour and 3 hours. (E) Immunoblot of nuclear PDHE1, (F) Quantification of nuclear PDH (n=3), Nuclear PDH was normalized to Histone 3, WCL: Whole cell lysate. One-way ANOVA followed by Bonferroni post-hoc analysis, **p-value< 0.01. Error bars, SEM.

(G) Immunoblot of nuclear PDHE1 in ischemia condition (2 hours normoxia followed by 2 hours hypoxia) and reperfusion condition (2 hours hypoxia followed by 2 hours normoxia). WCL: Whole cell lysate.

(H,I) H9C2 cardiomyoblasts were treated with $CoCl_2$ (100 μM) to mimic hypoxia conditions. (H) Immunoblot of nuclear PDH E1-α. (I) Quantification of nuclear PDH (n=3). Nuclear PDH was normalized to Histone 3. Paired t-test, **p-value< 0.01, ns: non-significant. Error bars, SEM.

Figure 1

A Budding reaction: Control / NEM → TEM Imaging, Floatation Ultracentrifugation, Mass spectroscopy

B Control budding reaction; NEM budding reaction

C Mt. pellet; Control; NEM — VDAC, TOM20, OPA1, PDH

D Mt. pellet; Sucrose gradient 10% 20% 30% 40% 50% 60% (1 2 3 4 5 6 7 8 9 10 11) — VDAC, TOM20, OPA1, PDH

E MDV proteome — MS1 (28), MS2 (34), MS3 (39), 42, 11, 49, 8

F NEM sensitive proteins — MS1 (22), MS2 (32), MS3 (54), 24, 7, 15, 15

G NEM SENSITIVE MDV CARGO

MATRIX
ACADS PDHX IARS2 PCCA
SUCLG1 DBT ACSS1 ACADM
PCCB DLAT IDH2 HIBCH
MCCC2 ALDH2 ETFA ECHS1 ETFB
ALDH5A1 BPHL ALDH7A1 MLYCD
ALDH9A1 KYAT3
PGK1 ECI2 CLPP SUCLG2 OXCT1
PPA2 IDH3A TST HIBADH BCAT2
PDHA1 PDHB ACAA2 MCEE
HSPA1A ACTB

IMM
COX2 ATP5PB ATP5PD
MPST NDUFA9 ETFDH SDHA
BCKDHA SLC25A4 ATP5ME
ATP5MF NDUFA12 ATP5PO
IMMT COX4I1 SLC25A11
SDHB CYC1

IMM/MATRIX
PPIF

IMS
AIFM1 ES1

OMM
ACSL1 VDAC1

Figure 2

Figure 3

Figure 4

A

Rat heart nuclei

In vitro reconstitution for 1h

Nucleus isolation

Rat cardiac MDVs

B

OPA1

PDH

COX4

Histone 3

C

Figure 5