Spatial risk of disease transmission between wild bovids and livestock in Thailand

Wantida Horpiencharoen[1*], Jonathan C. Marshall[1], Renata L. Muylaert[1], Reju Sam John[2], David T. S.
Hayman[1]

[1]Molecular Epidemiology and Public Health Laboratory, Hopkirk Research Institute, Massey University,
Palmerston North 4472 New Zealand
[2]Department of Physics, The University of Auckland, 38 Princes St, Auckland 1010, New Zealand

* Corresponding author Email: wantidah@gmail.com

Abstract

The livestock-wildlife interface is one of the most essential issues threatening wildlife
conservation and public health. Identifying interface areas can help to prioritise disease
surveillance and implement mitigation measures and control programs for targeting
threatened wildlife. We predicted interface areas which were assumed to be areas at risk of
infectious disease transmission based on the spatial overlap between three Thai wild bovids
(including gaur, banteng and wild water buffalo) habitat suitability and domestic cattle. We
assumed that domestic cattle are the reservoir of bovine infectious disease, and that high
cattle density is a proxy for a higher risk of disease transmission. Our study indicated that the
highest risk areas for the native species are at the forest edges where overlap exists between
high habitat suitability and high cattle density. Wild water buffalo showed the largest
proportion of high-risk areas (8%), while gaur and banteng showed similar risk areas (4%) in
Thailand. The largest proportion of risk areas overlapping with protected areas was Namtok
Sam Lan PAs at 89% for gaur, 84% for banteng and 65% for wild water buffalo. Kuiburi NP has
the largest risk area around 274 km^2 (around 28% of the total protected area) for gaur and
banteng, whereas wild water buffalo has the largest risk area overlapping with Huai
Thabthan-Had Samran around 126 km^2 (10% of the PA). Kaengkrachan Forest Complex
showed the second largest risk area from 249 km^2 for gaur and 273 km^2 for banteng (8-9% of
the PA). Our results address how habitat suitability might be helpful for infectious disease
prevention and control strategies focused on native fauna and One Health. Furthermore, this
work may also support the wild bovid habitat conservation initiatives and land use planning
by informing decision-making about balancing wildlife habitats and livestock farming.

Keywords: Cattle, infectious disease, interface area, mapping, wildlife

Introduction

Wild Bovidae (Mammalia: Artiodactyla) are distributed worldwide and play crucial
ecosystem roles, because they determine the forest and ecosystem structure, transport
micronutrients, and disperse plant seeds (1, 2) and are also important prey species of
predators (3). In Asia, wild bovid populations are threatened by multiple factors, including
habitat loss and hunting, especially in South to Southeast Asia (4). Natural habitat loss often

comes with increased free-grazing livestock interaction, which can lead to problems as varied
as resource competition, reducing wildlife population abundance (5), interbreeding between
domestic and wild water buffalo (6), and infectious disease transmission (7).

Among the twenty-seven recognized wild bovid species as of 2020 (IUCN), five species
remain in Thailand, including gaur (*Bos gaurus*), banteng (*Bos javanicus*), wild water buffalo
(*Bubalus arnee*), mainland serow (*Capricornis sumatraensis*) and Chinese goral (*Naemorhedus*
griseus). Their habitat and populations have been threatened by human activities such as
deforestation and hunting (8-10). Wild bovids, especially the large herbivores (e.g. gaur,
banteng and wild water buffalo), gradually adapt their distribution and behaviour to land use
change. For example, gaur has been found close to agricultural areas and forest edges where
they forage on crop plantations (e.g. grass, cassava) (11). Banteng is also well-adapted to
secondary forests near villages and logging sites (12). These wild bovids are, therefore, able
to share natural resources with free-grazing domestic bovids, which can potentially cause
disease transmission via direct and indirect contact with the sources of infections (e.g.
infectious cattle, host, environmental reservoirs (13).

The livestock-wildlife interface is one important issue threatening wildlife
conservation and global public health because 72% of reported emerging diseases originate
from wildlife to humans and/or livestock (14). Bovine infectious diseases, such as bovine
tuberculosis, brucellosis and foot and mouth disease, can be transmitted and circulate in
domestic and wild bovid populations (15). These diseases and their impact on wildlife and
livestock population health have been studied in Europe (16), North America (17) and Africa
(18), but less so in Asia, including Thailand.

Several factors can drive disease transmission between livestock and wildlife
populations, such as the expanding livestock production (19), the shrinking of wildlife habitat
(20) and changes in wildlife distribution, demography and behaviour (21). Among these
factors, high host density is potentially a determinant risk factor that can lead to successful
disease transmission as it may translate to a higher probability of between- and among-host
species interactions, contact and pathogen exposure (22, 23). The movement and spatial
overlap of wildlife and livestock can lead to increased infectious disease transmission risk.
Areas where there is potential for interaction between a new susceptible host and a reservoir
can increase the chance of disease transmission through increasing contact rates and time
(24). For wild and domesticated species, these areas are usually the transition areas between
two or more land use types, such as the edges of forest and agricultural areas, which are likely
to have more species activities leading to a greater chance of interaction and so disease
transmission among wildlife and livestock (25). Previous studies indicated the presence of
some infectious diseases, such as babesiosis and leptospirosis (26, 27) in domestic cattle at
the edge of forest, making these domestic animals a potential reservoir of disease
transmission to the wild bovids.

Bovine infectious diseases can circulate between livestock and wild ungulates with
varying levels of virulence (15). Certain pathogens can circulate within either livestock or
wildlife populations without causing visible clinical signs but have a significant impact on other
species. For example, foot and mouth disease (FMD) might not affect African buffalo, but can
lead to mortality in gaur (28). Similarly, haemorrhagic septicaemia (*Pasteurella multocida*)
might be identified within the farm environment as non-pathogenic with limited mortality
except under certain circumstances but can cause mass mortality in saiga (29). Chronic
diseases like bovine tuberculosis and brucellosis with long incubation and relatively low
fatality rates could potentially have long-term consequences by reducing populations due to
disease, while FMD, which has a higher transmission rate with short incubation periods or
even highly fatal infections such as anthrax, may have lower impacts on populations (30)

Moreover, in the past twenty years, there have been numerous transboundary
emerging disease outbreaks among domestic animals in Thailand. For example, there have
been outbreaks of lumpy skin disease among cattle (31), African horse sickness among horses
(32), and African swine fever among pigs (33). Hence, it is crucial to investigate where there
are high risk areas to prevent disease transmission to wild populations, considering their
susceptibility to similar pathogens shared by livestock. Targeting the potential risks of disease
transmission in wildlife and livestock interface areas can support the implementation of
surveillance and control measures that may help prevent cross-species transmission (34).

In this study, we aim to 1) identify the potential risk of disease transmission of wild
bovids and livestock in Thailand and 2) provide suggestions for disease surveillance and
conflict mitigation measures in the wildlife-livestock interface areas of Thailand. The outputs
could be used to prioritise local surveillance and mitigation measures for optimising resource
allocation.

Methods

Study area

Thailand is located on the Indochina Peninsula, part of mainland Southeast Asia.
Thailand borders four countries, Myanmar, Laos, Cambodia and Malaysia, with the Gulf of
Thailand on the southeast connected to the Pacific Ocean and the southwest connected to
the Andaman Sea. The total country area is around 514,000 km², with agricultural land
covering 41% and forested areas comprising around 32% of the country area. Most
agricultural areas consist of rice fields (51%) and crop plantations (34%), while livestock
farming covers only around 0.6% of the total agricultural area or around 0.3% of the total
country (land use data source: https://agri-map-online.moac.go.th/). There is high cattle
production in the central west, which includes free-range cattle and buffalo in some rural
areas. These free-ranging livestock have encroached into wildlife habitats and share the same
resources. Moreover, there is shared land use, for example, domestic buffalo may use rice
fields.

Populations

This study focuses on wildlife and livestock populations, and we calculate the largest potential for wildlife-livestock interface areas. We selected the remaining wild bovid species in Thailand because they are widely distributed and likely to share the same resources and pathogens as livestock, especially the large bovids (gaur, banteng and wild water buffalo) distribution, which tends to overlap with free-ranging cattle and agricultural areas. For the livestock population, we used the cattle population estimates as cattle production is all over the country, with varied production scales and systems from intensive farming to free grazing. We assume that domestic cattle can be a pathogen reservoir and transmit diseases to the wild bovid population, and our focus is on livestock transmitting infection to wild species, though the alternative is possible. Therefore, for our analysis, a high cattle density is assumed to have a higher risk transmission risk and a lower cattle density have a lower risk, as reported in previous studies (25, 35).

Identifying the potential risk

Briefly, we identified the risk area using two types of datasets: 1) wild bovid distribution and 2) cattle density. Then, we overlaid these together and calculated the overlapping areas in 1-km^2 cells as a sampling unit.

Wild bovid potential distribution

We assumed wild bovid distributions correlate with their suitable habitat we previously predicted by ecological niche models (36). Ecological niche models used 28 as predictor environmental variables using 8 algorithms. We conducted the ensemble models using the weighted mean method and used True Skill Statistics as a threshold to convert the ensemble models to binary values (1 = suitable areas and 0 = unsuitable). We selected three wild bovid species for further analysis: gaur (*B. gaurus*), banteng (*B. javanicus*), and wild water buffalo (*B. arnee*), and excluded Mainland serow (*C. sumatraensis*) or Chinese goral (*N. griseus*) from the analyses because the ecological niche models did not perform well. Full methods and model results can be found in Horpiencharoen et al. (2023) (36).

Cattle density

We downloaded cattle density data from Global Livestock of the World 2015, GLW 4 (link). This data gives values of cattle density at an original spatial resolution of 10 km^2. We cropped the raster layer to Thailand limits and disaggregated the raster to 1 km^2 per cell to make it compatible with the habitat suitability raster using the raster package (37). Then, we rescaled the density values to 0 - 1 using this equation:

$$X_2 = (X_1 - X_{minimum})/(X_{maximum} - X_{minimum})$$

Where X is the value in the cattle density cell. Then, we used the mean of cattle density (0.14 cattle/km^2) in Thailand calculated from the GLW 4 raster as a cut-off value for converting the cattle density raster into the binary values of high and low. The raster cells containing

values greater than the mean were converted to 1 (high density), and the values lower than
the mean were converted to 0 (low density).

We assumed that higher cattle density correlates with a greater risk of infectious
diseases. Therefore, we counted the number of outbreaks in low and high cattle density areas
to test this assumption. We divided the total number of outbreak events by the total area of
cattle density for each category to check whether the higher number of outbreaks in high-
density areas was not simply due to larger areas, as the following calculation:

157
$$\frac{Number\ of\ outbreak\ in\ high\ (or\ low)\ cattle\ density}{Total\ area\ of\ high\ (or\ low)\ cattle\ density}$$

The results found a higher incidence of outbreak events in high cattle density areas
compared to low cattle density areas. Thus, we used cattle density as the major risk factor to
identify the potential disease transmission areas (more details in the resultsTable 2**Error!**
Reference source not found.).

The potential risk areas
In this study, the potential high risk areas refer to the interface areas between wildlife
and livestock that potentially share the resources (e.g. water bodies, grassland, mineral lick)
and have a higher risk of disease transmission due to the increased opportunity of direct and
indirect contact with disease reservoirs and environment, compared to other areas (25, 35,
38).

To define the risk areas, we overlapped the cattle density raster with the species'
binary maps and calculated the percentages of the potential risk areas in Thailand. Then, we
intersected the risk areas with the national protected areas (PA) (39) to calculate the risk
areas inside and outside PAs and also classified the risk areas by land use types to prioritise
where to implement the disease surveillance. Lastly, we counted the occurrence of disease
outbreaks reported by the Department of Livestock Development, Thailand, within the
interface areas to explore the distribution of highly frequent diseases in the high risk areas
(see below). All spatial analyses were programmed in R 4.3.1 (40). The code is available at a
public repository (https://github.com/Wantidah/BovidRiskMaps). Data is available upon
reasonable request.

Disease occurrence data
We used the national database of livestock disease outbreak reports from 2013 to 2021
generated by the Department of Livestock Development, Thailand. The data collection starts
when an outbreak in livestock is reported by local authorities or farm owners. Then
epidemiological data are recorded, including the date of the index case, animal type, clinical
signs, and number of animals infected, followed by collecting the samples for laboratory
diagnosis. If a diagnosis is made and the causative pathogen and disease known, the
authorities will record this in the database. If the authorities cannot find the causing
pathogen, they will add the tentative diagnosis from the clinical signs. Each outbreak will be

reported as confirmed if the causative pathogen is identified by laboratory diagnostics.
However, if there is no laboratory result, the authority will fill in the tentative diagnosis
according to animal clinical signs. The GPS location of the outbreak refers to the centroid of
the sub-district (average area of districts of Thailand = 87 km^2, range: 0.88 - 2,387 km^2) where
the outbreak occurred.

Here we selected five globally or regionally common bovine infectious diseases
considered important for livestock health: 1) foot and mouth disease (FMD), 2) haemorrhagic
septicaemia (HS - *Pasteurella multocida*), 3) bovine Tuberculosis (*Mycobacterium bovis* - bTB),
4) lumpy skin disease (LSD) and 5) brucellosis (*Brucella abortus*) from the national database.
We selected outbreaks from these five diseases in cattle, then cleaned the outbreak events
by excluding incorrect coordinates falling outside Thailand using R. Lastly, we counted the
number of outbreaks within overlapping areas for each species and cattle population
densities using the 'extract' function in the raster R package (37).

Results

The high-risk areas with high wild bovid habitat suitability and high cattle density are
mostly found in the central-western through the southern part of Thailand for the three
species (Figure 1). The districts that showed the highest percentages of the risk areas are
Nakhon Si Thammarat (south), Ratchaburi and Prachuap Khiri Khan) for all three species. Wild
water buffalo showed the largest of the total interface areas, covering ~44,000 km^2 (8% of
Thailand), due to their potential habitat suitability predicted across the country. However, in
the actual species distribution, only one population remains in the Huai Kha Khaeng Wildlife
Sanctuary. Banteng and Guar showed similar potential habitat suitability species, which also
resulted in the closest number of interface areas, ~22,000 km^2 (4% of Thailand) (Table 1).

The highest percentage of risk areas inside the PA were identified in Namtok Sam Lan
(also known as Phra Budda Chai) National Park (NP) (45 km^2) in Saraburi Province, covering
approximately 89% for gaur, 83% for banteng and 65% for buffalo. The second highest
percentage for gaur and banteng is Namtok Huai Yang (160 km^2) NP in Prachuap Khiri Khan,
covering 60% (~100 km^2) of the total PA, and for wild water buffalo is Huai Thabthan-Had
Samran (498 km^2) representing 25% (125 km^2) of the total PA. However, for gaur and banteng,
the largest risk area is located in the same PA—Kuiburi NP (970 km^2), representing 273 km^2
(28% of the PA). This is followed by the Kaengkrachan forest complex, representing 249 km^2
for gaur (8% of the PA) and 261 km^2 for banteng (9% of the PA). These two protected areas
are in close proximity, with high-risk areas situated along the western forest edge, connected
to agricultural areas with high cattle density, while the western side is connected to the
Myanmar forest (Figure 1). Moreover, the large intact forests like the Western, Eastern and
Dong Payayen - Khoa Yai forest complex illustrated high habitat suitability with low cattle
density within the PA, but showed high risk at the border of the forests, while the fragmented
forests in the north illustrated the potential high risk of disease transmission with high cattle
density and low habitat suitability.

A

Bivariate maps of the potential risk area

B

High cattle density overlapped with high habitat suitability

Figure 1 Bivariate maps of the potential risk area (A) between the habitat suitability (blue) and cattle density (yellow) reveal the interface areas between three wild bovid species and domestic cattle populations in Thailand. High-risk areas are represented in dark red, while low-risk areas are represented in grey.The potential high risk areas for disease transmission between three wild bovid species and cattle populations in Thailand (B). High-risk areas are represented in dark red, extracted from the interface areas in (A).

235 Table 1 The percentage of interface areas overlapped with protected areas by three wild
236 bovid species.

Species	Interface area with livestock (km^2)			Percentage (%) of overlap area in Thailand
	inside PA	outside PA	total	
Gaur	2,018	18,367	20,385	3.9
Banteng	2,089	20,477	22,566	4.4
Wild water buffalo	747	43,642	44,389	8.7

According to the national disease surveillance, the total number of outbreak events is
7,522 events for the five selected bovine infectious diseases from 2013 to 2021. LSD (6,913)
has the most recorded outbreak events among the others, followed by FMD (563) and
brucellosis (39), while HS and bTB have only 5 and 2 events, respectively (Table 2 and Figure
2). This is because the first LSD outbreak occurred in cattle herds in Thailand in 2021, leading
to a large number of events reported across the country in a short period, while the other
infections, which have lower records, are endemic in this area.

The cattle density demonstrated correlations with the number of infectious disease
outbreaks, and this correlation is proportional to the area size. We observed that in high cattle
density areas (190,076 km^2), there were higher outbreak events, totalling around 6,894
events (0.036 events per km^2), 18 times more than low cattle density areas (324,335 km^2),
which had 640 events (0.002 events per km^2). The results of outbreak events by cattle density
areas are presented in Table 2

Table 2 The occurrence of outbreak events classified by infectious disease and cattle density
from 2013 to 2021.

Infectious disease	Cattle density		Total
	Low (<=mean)	High (>mean)	
bTB	2	0	2
HS	3	2	5
LSD	445	6,473	6,918
FMD	175	395	563
Brucellosis	15	24	39
Total	640	6,894	7,534
Area of Thailand (km^2)	324,335	190,076	514,411
Total outbreak event per area (km^2)	0.0020	0.0363	0.0146

Figure 2 Occurrence of recorded bovine infectious disease outbreaks in Thailand from 2013
to 2021 and protected area distribution (PA; green). The maps show that the outbreaks
occurred across Thailand, particularly in proximity to protected and forest areas that overlap
with suitable habitats for wild bovids.

Table 3 The reported outbreak events of bovine infectious disease occurrences in the high
risk areas.

Bovine Infectious Diseases	All occurrences	Within high cattle density - high habitat suitability		
		Gaur	Banteng	Wild water buffalo
Tuberculosis	2	0	0	0
HS	5	0	0	1
LSD	6,918	166	108	442
FMD	563	32	51	190
Brucellosis	39	1	3	14
Total	7,534	199	162	647
High risk areas (km^2)		20,385	22,566	44,389
Thailand	514,411			
Total outbreak event per risk area (km^2)	0.015	0.010	0.008	0.015

We found that the density of outbreak events in cattle within the potential high risk
area of gaur (0.01) and especially for wild water buffalo (0.015) were similar to the average
density calculated for the country (0.0146) (Table 3). Wild water buffalo showed the highest
events (647) within the risk areas as they have the largest potential habitat areas across the
country while gaur (199) and banteng (162) show close results to each other. Similar to Table

2, the greatest numbers of disease events within high risk areas was for LSD and FMD in all
species and the other diseases presented only small numbers.

Moreover, according to the land use types, the most extensive interface areas were
found in close unknown forests (meaning they did not match any of the other forest
definitions), followed by cropland for three species. Closed evergreen forests also contain risk
areas for gaur and banteng. The open deciduous forest had no interface areas detected
(Figure 3).

Figure 3 Interface areas of three wild bovid species are categorised based on land use types,
from the land cover layers dataset ((41), https://zenodo.org/records/3243509). The term
closed unknown forest denotes a type of forest that does not match any of the other
definitions.

Discussion

We examined potential risk areas of disease transmission between wild bovids and
livestock and provided the preliminary focus area that should be considered for disease
surveillance and mitigation in Thailand. Total risk areas in Thailand are between 4% (gaur and
banteng) to 9% (wild water buffalo) of the country, with the most high risk areas being from
the central west (Ratchaburi) to the south (Nakhon Si Thammarat). The highest risk
proportion inside PAs was at Namtok Sam Lan National Park (NP) in the central, and the
largest risk areas were Kuiburi NP and Kaengkrachan NP in the western forest, related to the
highest cattle density in Thailand. Gaur and banteng have similar risk areas mostly around the

edge of forests, while wild water buffalo have risk widely across the country because models
predicted extensive amounts of suitable areas in the central down to the southern part that
coincide with the high cattle density areas.

Although the wild water buffalo showed the largest areas of predicted suitable habitat
among all three bovid species, it is also the most endangered. This is due to its current
distribution being confined solely to the Huai Kha Khaeng Wildlife Sanctuary, with the
population remaining stable and not exceeding 69 individuals for decades (42, 43). Thus,
many areas identified as high risk are unlikely to be current high risk areas but are important
for future planning should wild buffalo ranges expand to these or they be relocated. This
species is highly susceptible to endemic infectious diseases that could rapidly lead to serious
decline or even local extinction. For instance, diseases like HS can cause a high fatality rate of
up to 80% in domestic buffalo (44). The highly contagious and fatal nature of diseases like
rinderpest and FMD may be contributing factors to population disappearances in Nepal and
India (45). An outbreak could also lead to local extinction in a single fragmented population,
as the recovery process is prolonged and potentially results in a lack of gene flow (46),
especially with the independent mother origin of Thai wild water buffalo (47).

In contrast to wild water buffalo, gaur and banteng have more opportunities for contact
with domestic cattle and humans, while wild water buffalo may encounter livestock and
humans encroaching into the protected areas and suitable habitat. Gaur and banteng can
share habitats, making the interface areas similar to each other. However, gaur uses a wider
range of habitat types (e.g., evergreen, deciduous dipterocarp, mixed deciduous forests) than
banteng, which is restricted to dry and open forests (e.g., dry dipterocarp, mixed deciduous
forest) (48, 49). These two species show evidence of contact with livestock and humans due
to their ability to adapt and tolerate human activities, resulting in conflicts in overlapping
areas (11, 12, 50). However, their habitat suitability decreases when the distance is closer to
human settlement and the presence of domestic cattle grazing (51).

Our study identified the potential high-risk areas in the northeastern and southern
parts, which have the highest cattle density but low or even an absence of the actual species
distribution in some areas. This caveat is observed in the ecological niche modelling of wild
water buffalo, where high habitat suitability represents potential distribution and may not
necessarily correspond to the actual species distribution. Nevertheless, this caveat could be
mitigated by collecting and regularly updating occurrences of these bovid species, as well as
data on livestock density and distribution, or by restricting the analyses to areas with
sufficient data.

Urbanisation and expansion of agricultural areas increase the opportunity for contact
between domestic livestock and wildlife. Contact rate, the probability of transmission and the
location shifts of animals at each time step, are the major factors that need to be considered
for the spatial disease transmission model (52). The direct contact between wildlife and
livestock is unlikely but indirect contact in the same space at different times via shared

resources (e.g. water, grassland, supplement) with domestic animals potentially causes a
chance of wildlife exposure to pathogens and disease transmission (13, 53). However, to
succeed in cross-species infection, several factors should converge to drive pathogens
through the natural barriers before having a pathogenic infection into a new host (54). We
found that the highest interfaced areas were identified in cropland and unclassified forests,
which potentially can be shared by free-ranging livestock and wild bovids. Thailand also
experienced significant land cover changes (55) primarily driven by the increase of crop
plantations and urbanisation with concurrent population growth, which leads to changing
wildlife and livestock interactions and risk of disease transmission as per the previous studies
(56).

Host density is one of the main risk factors in wildlife and livestock disease transmission
(57, 58). We used host density and distribution as the main risk factors to identify and
prioritise the potential risk areas of wildlife and livestock disease transmission, as has been
used in the other studies (53, 59, 60). The advantage is knowing the target place for
implementing the disease surveillance system, but the disadvantages can arise from the
complexity of disease transmission dynamics, which depend on factors such as host species
movement (60), contact pattern (61), high adaptability of wildlife behaviour, transmission
modes (e.g. density or frequency-dependent) (58, 62) and population size (63).

Understanding the underlying factors that contribute to disease outbreaks in a specific
potential risk area is essential for planning effective disease mitigation and control strategies.
Multidisciplinary approaches incorporating key elements like pathogens, hosts, and
environmental factors (supplementary materials Table S1-S2, describe the generic risk factors
for disease transmission and mitigation methods between wildlife and livestock) also help
policymakers develop disease control and mitigation measures. Pathogen spillover events are
complicated, with a convergence of risk factors, which are difficult to approach. Integrating a
complex system of human, animal and environmental will benefit prevention efforts or at
least mitigate the impact of the next spill-over event (64).

Mitigation strategies will likely vary according to local socioeconomic conditions, but
among the preventive actions are using vaccination of livestock or even wild species, targeted
reduction of infected individual animals, herds or farms (ideally with compensation), along
with reducing livestock herd sizes and densities, transport of livestock among farms, and
contacts between farmed animals and wild species (65, 66). Contact reduction might be
through measures such as altering land use at the local level, or with "natural" (e.g. plant-
based) or artificial (e.g. metal) fencing or barriers and zoning of forests, livestock and human
settlements to minimise the contact (17, 67), which may lead to reduce pathogen spillover
(68). Longer term strategies might include societal transitions to lower meat-based, more
plant-based proteins to reduce demand for meat and dairy products. Conserving intact forests
with effective surveillance can mitigate the risk of disease transmission at the interface,
especially in edge or transition areas. In contrast, fragmented forests increase the likelihood

of wildlife being exposed to livestock and humans, leading to an elevated risk of disease
transmission (20, 69, 70).

Livestock vaccination is crucial for reducing outbreak incidences of endemic diseases,
requiring approximately 80% coverage to effectively prevent disease transmission for many
pathogens based on pathogen specific, particularly in high-risk areas and populations, as part
of routine practice (71, 72). However, capturing and delivering parenteral-route vaccinations
to free-ranging wild bovids pose significant challenges, especially in tropical forests where
animals might be hidden. Consequently, various aspects must be carefully considered in the
vaccination plan, including the target population, coverage, safety, and efficiency, to
effectively stimulate herd immunity (73). Non-invasive vaccination methods, like tuberculosis
oral vaccination, have been tested in domestic cattle and some wildlife and are planned for
use in wild cattle (74). Research and development for other endemic diseases like FMD, HS,
and brucellosis is still ongoing (75-77). Culling livestock infected with zoonotic diseases (e.g.,
bTB, brucellosis) is commonly implemented in Thailand (78, 79). However, infected animals
often undergo illegal translocation, potentially spreading the disease to other locations. To
manage this issue, the government should rigorously regulate animal movement, regulating
the guidelines for isolation of infectious animals during outbreaks, and providing appropriate
compensation for culling cases. The effectiveness of these mitigation actions is influenced by
the presence and use of effective infection and disease surveillance, as discussed below.

Livestock and, likely, wildlife disease surveillance in Thailand is based on the DLD, DNP
(Thailand) and WOAH guidelines (80), which cover significant transboundary disease
outbreaks in the country. Even though the passive surveillance system is useful for recording
the obvious clinical signs and emerging infectious diseases (like FMD and LSD), there is a gap
in collecting non-clinical to subclinical signs of disease due to these being challenging to
detect. Moreover, passive surveillance leads to underreporting by the farmers for some
zoonoses like bovine tuberculosis and brucellosis, for which animals must be condemned due
to the slow and partial (not less than 75% of the market price, but often not 100%)
compensation from the government. Another drawback is the clinical signs reported from
passive surveillance may not refer to the place where the animal got infected if those are
moved from the original area. Therefore, active surveillance such as risk-based (81), disease
surveys (Arjkumpa et al., 2020) screening or detecting seroprevalence are necessary in
hotspots or endemic areas to effectively allocate resources for disease mitigation and control
strategies. Furthermore, even when reported, further work must be undertaken to
understand the disease risk in depth. Reported data might refer to one event being a single
case or multiple cases, and infectious diseases are, by default, dynamic in their nature.
Without significant further work, passive surveillance data may offer a limited and biased
understanding of the true disease risk in a location at a particular time. The impact of the
detection of some infections on trade must be addressed, as this might be a barrier to
effective surveillance and reporting (82, 83). The involvement of field authorities is another
crucial aspect of data collection, indispensable for obtaining real-time information. Modern

technologies or data sharing can help identify risk areas and plan preparedness implementations (62, 84, 85). Therefore, one should consider investing more in field data collection and incorporating field practitioners or epidemiologists into the team before formulating policies (86).

Using a One Health approach, a framework for disease surveillance has been developed, incorporating essential considerations of spillover events into the processes. To sustainably manage data collection and surveillance systems, collaboration among government organisations and stakeholders is a key step in the process, involving considerations of political, ethical, administrative, regulatory, and legal (PEARL) aspects through all approaches (87). An effective surveillance system, characterised by rapid detection and accurate results, not only monitors emerging diseases but also reduces the risk of disease transmission and minimizes the impact on lives, economies, and biodiversity during disease outbreaks (88). Also, the use of non-invasive data collection for willdife disease surveillance and surveys, such as faces, urine, saliva and environmental samples (e.g. soil, water) should be considered to avoid direct contact and reduce disturbing wildlife during capturing and data collection (89). Further studies may consider including other risk factors such as multi-species host distribution, the distance of risk factors and contact pattern (90, 91), as well as improving the model by using updated disease surveillance data and wild bovid species occurrences, especially for areas where the uncertainty of model predictions is high.

Conclusion

Our study predicted the potential risk areas by using the interface areas between wildlife and domestic cattle, where livestock disease is frequently reported. We overlaid suitable habitats of three large wild bovids in Thailand with cattle density to create potential risk maps. High-risk areas were identified in locations with both high cattle density and high habitat suitability, particularly at the edges of forest-protected areas. Notably, small, fragmented forest areas with high cattle density presented the highest proportion of the high-risk areas. Among various land-use types, cropped land and some closed forests exhibited the largest interface areas. Our findings highlight the importance of wildlife habitat and intact forest conservation to mitigate contact and reduce vulnerability to extinction, reduce shared areas and address the potential risk areas for disease transmission between wild bovids and livestock. This methodology not only supports disease surveillance but also facilitates the implementation of effective mitigation and control measures.

Acknowledgements

WH was supported by Manaaki New Zealand Scholarship. D.T.S.H., RSJ & RLM were supported by Bryce Carmine and Anne Carmine (née Percival), through the Massey University Foundation (grant no. RM22688), and D.T.S.H. was supported by the Percival Carmine Chair in Epidemiology and Public Health and Royal Society Te Apūrangi (grant no. RDF-MAU1701). Thank you to Dr Weerapong Thanapngtham from DLD, Thailand, for sharing the disease surveillance data.

Supplementary Materials

Table S1: Generic risk factors for disease transmission in wildlife and livestock interface areas, including studies and perspectives in Thailand and internationally. Table S2: Mitigation methods for infectious disease transmission.

References

1. Ripple WJ, Newsome TM, Wolf C, Dirzo R, Everatt KT, Galetti M, et al. Collapse of the world's largest herbivores. Science advances. 2015;1(4):e1400103-e.
2. Romero GQ, Gonçalves-Souza T, Vieira C, Koricheva J. Ecosystem engineering effects on species diversity across ecosystems: a meta-analysis. Biological Reviews. 2015;90(3):877-90.
3. Simcharoen A, Simcharoen S, Duangchantrasiri S, Bump J, Smith JLD. Tiger and leopard diets in western Thailand: Evidence for overlap and potential consequences. Food Webs. 2018;15:e00085.
4. Tilker A, Abrams JF, Mohamed A, Nguyen A, Wong ST, Sollmann R, et al. Habitat degradation and indiscriminate hunting differentially impact faunal communities in the Southeast Asian tropical biodiversity hotspot. Commun Biol. 2019;2(1):1-11.
5. Bhandari S, Crego RD, Stabach JA. Spatial segregation between wild ungulates and livestock outside protected areas in the lowlands of Nepal. PLOS ONE. 2022;17(1):e0263122.
6. Bhattarai RK, Joshi R, Gautam J, Kandel B, Singh B. Reproductive Behavior of *Bubalus arnee* (Kerr, 1792) (Mammalia: Artiodactyla: Bovidae) in Koshi Tappu Wildlife Reserve and Chitwan National Park, Nepal. International Journal of Zoology. 2023;2023:e6305614.
7. Hassell JM, Begon M, Ward MJ, Fèvre EM. Urbanization and Disease Emergence: Dynamics at the Wildlife–Livestock–Human Interface. Trends in Ecology & Evolution. 2017;32(1):55-67.
8. Carr E, Amir Z, Mendes C, Moore J, Nursamsi I, Luskin M. The highs and lows of serow (*Capricornis sumatraensis*): multi-scale habitat associations inform large mammal conservation strategies in the face of synergistic threats of deforestation, hunting, and climate change. 2023.
9. Harrison RD, Sreekar R, Brodie JF, Brook S, Luskin M, O'Kelly H, et al. Impacts of hunting on tropical forests in Southeast Asia. Conservation Biology. 2016;30(5):972-81.
10. Nguyen MH. The status of Vulnerable gaur Bos gaurus and Endangered banteng *Bos javanicus* in Ea So Nature Reserve and Yok Don and Cat Tien National Parks, Vietnam. Oryx. 2009;43(1):129-35.
11. Chaiyarat R, Prasopsin S, Bhumpakphan N. Food and nutrition of Gaur (*Bos gaurus* C.H. Smith, 1827) at the edge of Khao Yai National Park, Thailand. Scientific Reports. 2021;11(1):3281.
12. Journeaux K, Gardner P, Lim H, Wern J, Goossens B. Herd demography, sexual segregation and the effects of forest management on Bornean banteng *Bos javanicus lowi* in Sabah, Malaysian Borneo. Endang Species Res. 2018;35:141-57.
13. Bacigalupo SA, Dixon LK, Gubbins S, Kucharski AJ, Drewe JA. Towards a unified generic framework to define and observe contacts between livestock and wildlife: a systematic review. PeerJ. 2020;8:e10221.

14.   Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in
emerging infectious diseases. Nature. 2008;451(7181):990-3.
15.   Caron A, Miguel E, Gomo C, Makaya P, Pfukenyi DM, Foggin C, et al. Relationship
between burden of infection in ungulate populations and wildlife/livestock interfaces.
Epidemiology & Infection. 2013;141(7):1522-35.
16.   Martin C, Pastoret P-P, Brochier B, Humblet M-F, Saegerman C. A survey of the
transmission of infectious diseases/infections between wild and domestic ungulates in
Europe. Veterinary Research. 2011;42(1):70.
17.   Miller RS, Farnsworth ML, Malmberg JL. Diseases at the livestock–wildlife interface:
Status, challenges, and opportunities in the United States. Preventive Veterinary
Medicine. 2013;110(2):119-32.
18.   Garine-Wichatitsky MD, Caron A, Kock R, Tschopp R, Munyeme M, Hofmeyr M, et al. A
review of bovine tuberculosis at the wildlife–livestock–human interface in sub-Saharan
Africa. Epidemiology & Infection. 2013;141(7):1342-56.
19.   Morand S. Emerging diseases, livestock expansion and biodiversity loss are positively
related at global scale. Biological Conservation. 2020;248:108707.
20.   Wilkinson DA, Marshall JC, French NP, Hayman DTS. Habitat fragmentation, biodiversity
loss and the risk of novel infectious disease emergence. Journal of The Royal Society
Interface. 2018;15(149):20180403.
21.   Eby P, Peel AJ, Hoegh A, Madden W, Giles JR, Hudson PJ, et al. Pathogen spillover driven
by rapid changes in bat ecology. Nature. 2023;613(7943):340-4.
22.   Habib TJ, Merrill EH, Pybus MJ, Coltman DW. Modelling landscape effects on density–
contact rate relationships of deer in eastern Alberta: Implications for chronic wasting
disease. Ecological Modelling. 2011;222(15):2722-32.
23.   Sirdar MM, Fosgate GT, Blignaut B, Mampane LR, Rikhotso OB, Du Plessis B, et al. Spatial
distribution of foot-and-mouth disease (FMD) outbreaks in South Africa (2005–2016).
Tropical Animal Health and Production. 2021;53(3):376.
24.   Clifford DL, Schumaker BA, Stephenson TR, Bleich VC, Cahn ML, Gonzales BJ, et al.
Assessing disease risk at the wildlife–livestock interface: A study of Sierra Nevada
bighorn sheep. Biological Conservation. 2009;142(11):2559-68.
25.   Miguel E, Grosbois V, Caron A, Boulinier T, Fritz H, Cornélis D, et al. Contacts and foot
and mouth disease transmission from wild to domestic bovines in Africa. Ecosphere.
2013;4(4):art51.
26.   Simking P, Yatbantoong N, Saetiew N, Saengow S, Yodsri W, Chaiyarat R, et al.
Prevalence and risk factors of Babesia infections in cattle trespassing natural forest
areas in Salakpra Wildlife Sanctuary, Kanchanaburi Province. Journal of Tropical
Medicine and Parasitology. 2014;37(1):10-9.
27.   Yatbantoong N, Chaiyarat R. Factors Associated with Leptospirosis in Domestic Cattle in
Salakphra Wildlife Sanctuary, Thailand. International Journal of Environmental Research
and Public Health. 2019;16(6):1042.
28.   Chandranaik BM, Hegde R, Shivashankar BP, Giridhar P, Muniyellappa HK, Kalge R, et al.
Serotyping of foot and mouth disease virus and *Pasteurella multocida* from Indian gaurs
(*Bos gaurus*), concurrently infected with foot and mouth disease and haemorrhagic
septicaemia. Tropical Animal Health and Production. 2015;47(5):933-7.
29.   Robinson S, Milner-Gulland EJ, Grachev Y, Salemgareyev A, Orynbayev M, Lushchekina
A, et al. Opportunistic bacteria and mass mortality in ungulates: lessons from an
extreme event. Ecosphere. 2019;10(6):e02671.

30. Horpiencharoen W, Marshall JC, Muylaert RL, John RS, Hayman DTS. Impact of
Infectious Diseases on Wild Bovidae Populations in Thailand: Insights from Population
Modelling and Disease Dynamics. bioRxiv. 2023:2023.08.29.554960.
31. Arjkumpa O, Suwannaboon M, Boonrod M, Punyawan I, Liangchaisiri S, Laobannue P,
et al. The First Lumpy Skin Disease Outbreak in Thailand (2021): Epidemiological
Features and Spatio-Temporal Analysis. Frontiers in Veterinary Science. 2022;8.
32. King S, Rajko-Nenow P, Ashby M, Frost L, Carpenter S, Batten C. Outbreak of African
horse sickness in Thailand, 2020. Transboundary and Emerging Diseases.
2020;67(5):1764-7.
33. Thanapongtharm W, Wongphruksasoong V, Sangrat W, Thongsrimoung K,
Ratanavanichrojn N, Kasemsuwan S, et al. Application of Spatial Risk Assessment
Integrated With a Mobile App in Fighting Against the Introduction of African Swine
Fever in Pig Farms in Thailand: Development Study. JMIR Formative Research.
2022;6(5):e34279.
34. Kelly TR, Karesh WB, Johnson CK, Gilardi KVK, Anthony SJ, Goldstein T, et al. One Health
proof of concept: Bringing a transdisciplinary approach to surveillance for zoonotic
viruses at the human-wild animal interface. Preventive Veterinary Medicine.
2017;137:112-8.
35. Khanyari M, Robinson S, Morgan ER, Brown T, Singh NJ, Salemgareyev A, et al. Building
an ecologically founded disease risk prioritization framework for migratory wildlife
species based on contact with livestock. Journal of Applied Ecology. 2021;58(9):1838-
53.
36. Horpiencharoen W, Muylaert RL, Reju Sam J, Marshall JC, Lynam AJ, Riggio A, et al.
Mapping threatened Thai bovids provides opportunities for improved conservation
outcomes in Asia. bioRxiv. 2023:2023.08.25.554763.
37. Hijmans RJ. raster: Geographic Data Analysis and Modeling. 2023.
38. Barasona JA, Latham MC, Acevedo P, Armenteros JA, Latham ADM, Gortazar C, et al.
Spatiotemporal interactions between wild boar and cattle: implications for cross-
species disease transmission. Veterinary Research. 2014;45(1):122.
39. Protected Planet: The World Database on Protected Areas (WDPA) [Online] [Internet].
UNEP-WCMC and IUCN. 2021 [cited March 2021]. Available from:
www.protectedplanet.net.
40. R Core Team. R: A language and environment for statistical computing. Vienna, Austria:
R Foundation for Statistical Computing; 2023.
41. Buchhorn M, Smets B, Bertels L, Lesiv M, Tsendbazar N-E, Herold M, et al., editors.
Copernicus Global Land Service: Land Cover 100m: collection 2: epoch 2015: Globe
(Version V2.0.2) [Data set]. ESA Living Planet Symposium 2019 (LPS2019); 2019; Milan,
Italy: Zenodo.
42. Chaiyarat R, Lauhachinda V, Kutintara U, Bhumpakphan N, Prayurasiddhi T. Population
of wild water buffalo (*Bubalus bubalis*) in Huai Kha Khaeng Wildlife Sanctuary, Thailand.
Natural history bulletin of the Siam Society. 2004;52(2):151-62.
43. Khiowsree N, Sukmasuang R, Kutintara U. Population monitoring and distribution range
of wild buffalo (*Bubalus bubalis* LINNAEUS, 1758) in Huai Kha Khaeng wildlife sanctuary,
Uthai Thani province. Journal of Wildlife in Thailand. 2015;22(1):112 - 8.
44. Deepak, Gulia D, Aly SS. Modeling Vaccination Programs in Outbreaks of Hemorrhagic
Septicemia in India. JAR. 2020;10(1).

45.   Choudhury A. Wild water buffalo *Bubalus arnee* (Kerr, 1792). In: Burton J, Melletti M,
581   editors. Ecology, Evolution and Behaviour of Wild Cattle: Implications for Conservation.
582   Cambridge: Cambridge University Press; 2014. p. 255-301.
583   46.   McKnight DT, Schwarzkopf L, Alford RA, Bower DS, Zenger KR. Effects of emerging
584   infectious diseases on host population genetics: a review. Conservation genetics.
585   2017;18:1235-45.
586   47.   Sarataphan N, Narongwanichgarn W, Maneerat S. Phylogenetic Analysis of a Thai Wild
587   Water Buffalo (*Bubalus arnee*) Through Mitochondrial Control Region. International
588   Journal of Conservation Science. 2017;8(1):105-12.
589   48.   Prayurasiddhi T. The ecological separation of gaur (*Bos gaurus*) and banteng (*Bos
590   javanicus*) in Huai Kha Khaeng Wildlife Sanctuary, Thailand: University of Minnesota;
591   1997.
592   49.   Steinmetz R. Gaur (*Bos gaurus*) and Banteng (*B. javanicus*) in the lowland forest mosaic
593   of Xe Pian Protected Area, Lao PDR: abundance, habitat use, and conservation.
594   Mammalia. 2004;68(2-3):141-57.
595   50.   Chaiyarat R, Ingudomnukul P, Yimphrai N, Nakbun S, Youngpoy N. The Preferred Habitat
596   of Reintroduced Banteng (*Bos javanicus*) at the Core and the Edge of Salakphra Wildlife
597   Sanctuary, Thailand. Animals. 2023;13(14):2293.
598   51.   Jornburom P, Duangchantrasiri S, Jinamoy S, Pattanavibool A, Hines JE, Arnold TW, et
599   al. Habitat use by tiger prey in Thailand's Western Forest Complex: What will it take to
600   fill a half-full tiger landscape? Journal for Nature Conservation. 2020;58:125896.
601   52.   Böhm M, Hutchings MR, White PCL. Contact Networks in a Wildlife-Livestock Host
602   Community: Identifying High-Risk Individuals in the Transmission of Bovine TB among
603   Badgers and Cattle. PLOS ONE. 2009;4(4):e5016.
604   53.   Yang A, Boughton RK, Miller RS, Wight B, Anderson WM, Beasley JC, et al. Spatial
605   variation in direct and indirect contact rates at the wildlife-livestock interface for
606   informing disease management. Preventive Veterinary Medicine. 2021;194:105423.
607   54.   Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to
608   zoonotic spillover. Nat Rev Microbiol. 2017;15(8):502-10.
609   55.   Wang Y, Hu Y, Niu X, Yan H, Zhen L. Land Use/Cover Change and Its Driving Mechanism
610   in Thailand from 2000 to 2020. Land. 2022;11(12):2253.
611   56.   Shah HA, Huxley P, Elmes J, Murray KA. Agricultural land-uses consistently exacerbate
612   infectious disease risks in Southeast Asia. Nature Communications. 2019;10(1):4299.
613   57.   Daszak P, Cunningham AA, Hyatt AD. Emerging Infectious Diseases of Wildlife-- Threats
614   to Biodiversity and Human Health. Science. 2000;287(5452):443-9.
615   58.   Smith KF, Acevedo-Whitehouse K, Pedersen AB. The role of infectious diseases in
616   biological conservation. Animal Conservation. 2009;12(1):1-12.
617   59.   Miller RS, Sweeney SJ, Slootmaker C, Grear DA, Di Salvo PA, Kiser D, et al. Cross-species
618   transmission potential between wild pigs, livestock, poultry, wildlife, and humans:
619   implications for disease risk management in North America. Scientific Reports.
620   2017;7(1):7821.
621   60.   Morgan ER, Lundervold M, Medley GF, Shaikenov BS, Torgerson PR, Milner-Gulland EJ.
622   Assessing risks of disease transmission between wildlife and livestock: The Saiga
623   antelope as a case study. Biological Conservation. 2006;131(2):244-54.
624   61.   Craft ME. Infectious disease transmission and contact networks in wildlife and livestock.
625   Philosophical Transactions of the Royal Society B: Biological Sciences.
626   2015;370(1669):20140107.

62. Pruvot M, Musiani M, Boyce MS, Kutz S, Orsel K. Integrating livestock management and
telemetry data to assess disease transmission risk between wildlife and livestock.
Preventive Veterinary Medicine. 2020;174:104846.
63. Lambert S, Ezanno P, Garel M, Gilot-Fromont E. Demographic stochasticity drives
epidemiological patterns in wildlife with implications for diseases and population
management. Scientific Reports. 2018;8(1):16846.
64. Vora NM, Hannah L, Walzer C, Vale MM, Lieberman S, Emerson A, et al. Interventions
to Reduce Risk for Pathogen Spillover and Early Disease Spread to Prevent Outbreaks,
Epidemics, and Pandemics. Emerging Infectious Diseases. 2023;29(3):e221079.
65. Carrasco-Garcia R, Barasona JA, Gortazar C, Montoro V, Sanchez-Vizcaino JM, Vicente J.
Wildlife and livestock use of extensive farm resources in South Central Spain:
implications for disease transmission. European Journal of Wildlife Research.
2016;62(1):65-78.
66. Kuiken T, Cromie R. Protect wildlife from livestock diseases. Science. 2022;378(6615):5-
5.
67. Jori F, Etter E. Transmission of foot and mouth disease at the wildlife/livestock interface
of the Kruger National Park, South Africa: Can the risk be mitigated? Prev Vet Med.
2016;126:19-29.
68. Plowright RK, Reaser JK, Locke H, Woodley SJ, Patz JA, Becker DJ, et al. Land use-induced
spillover: a call to action to safeguard environmental, animal, and human health. The
Lancet Planetary Health. 2021;5(4):e237-e45.
69. Muylaert RL, Wilkinson DA, Kingston T, D'Odorico P, Rulli MC, Galli N, et al. Using drivers
and transmission pathways to identify SARS-like coronavirus spillover risk hotspots.
Nature Communications. 2023;14(1):6854.
70. Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, et al. Impacts of
biodiversity on the emergence and transmission of infectious diseases. Nature.
2010;468(7324):647-52.
71. Hopker A, Pandey N, Bartholomew R, Blanton A, Hopker S, Dhamorikar A, et al. Livestock
vaccination programme participation among smallholder farmers on the outskirts of
National Parks and Tiger Reserves in the Indian states of Madhya Pradesh and Assam.
PLoS ONE. 2021;16(8):e0256684.
72. Porco A, Chea S, Sours S, Nou V, Groenenberg M, Agger C, et al. Case report: Lumpy skin
disease in an endangered wild banteng (*Bos javanicus*) and initiation of a vaccination
campaign in domestic livestock in Cambodia. Frontiers in Veterinary Science. 2023;10.
73. Treanor JJ, Johnson JS, Wallen RL, Cilles S, Crowley PH, Cox JJ, et al. Vaccination
strategies for managing brucellosis in Yellowstone bison. Vaccine. 2010;28:F64-F72.
74. Buddle BM, Vordermeier HM, Chambers MA, de Klerk-Lorist LM. Efficacy and Safety of
BCG Vaccine for Control of Tuberculosis in Domestic Livestock and Wildlife. Front Vet
Sci. 2018;5:259.
75. Gortazar C, Diez-Delgado I, Barasona JA, Vicente J, De La Fuente J, Boadella M. The Wild
Side of Disease Control at the Wildlife-Livestock-Human Interface: A Review. Frontiers
in Veterinary Science. 2015;1.
76. Olsen SC, Christie RJ, Grainger DW, Stoffregen WS. Immunologic responses of bison to
vaccination with Brucella abortus strain RB51: Comparison of parenteral to ballistic
delivery via compressed pellets or photopolymerized hydrogels. Vaccine.
2006;24(9):1346-53.

77. Lee B-R, Lee H-J, Kim N-H, Kim Y-S, Park KI. Increased Effect of Foot-and-Mouth Disease
Virus Vaccine Structural Protein Antibody Positivity Rates in Piglets Orally Treated with
Amino–Zinc Complex. Animals (Basel). 2023;13(12):2027.

78. Peck ME, Chanachai K, Jenpanich C, Amonsin A, Alexander B, Bender JB. Seroprevalence
of Brucellosis in Goats and Sheep in Thailand: Results from the Thai National Brucellosis
Surveillance System from 2013 – 2015. Transboundary and emerging diseases.
2018;65(3):799-805.

79. Singhla T, Boonyayatra S, Punyapornwithaya V, VanderWaal KL, Alvarez J, Sreevatsan S,
et al. Factors Affecting Herd Status for Bovine Tuberculosis in Dairy Cattle in Northern
Thailand. Veterinary Medicine International. 2017;2017:2964389.

80. WOAH. Animal Diseases - WOAH - World Organisation for Animal Health 2024 [Available
from: https://www.woah.org/en/what-we-do/animal-health-and-welfare/animal-
diseases/?_tax_animal=terrestrials.

81. Sansamur C, Arjkumpa O, Charoenpanyanet A, Punyapornwithaya V. Determination of
Risk Factors Associated with Foot and Mouth Disease Outbreaks in Dairy Farms in
Chiang Mai Province, Northern Thailand. Animals. 2020;10(3).

82. Khengwa C, Jongchansittoe P, Sedwisai P, Wiratsudakul A. A traditional cattle trade
network in Tak province, Thailand and its potential in the spread of infectious diseases.
Anim Prod Sci. 2015;57(1):152-60.

83. Wiratsudakul A, Sekiguchi S. The implementation of cattle market closure strategies to
mitigate the foot-and-mouth disease epidemics: A contact modeling approach.
Research in Veterinary Science. 2018;121:76-84.

84. Goutard FL, Binot A, Duboz R, Rasamoelina-Andriamanivo H, Pedrono M, Holl D, et al.
How to reach the poor? Surveillance in low-income countries, lessons from experiences
in Cambodia and Madagascar. Preventive Veterinary Medicine. 2015;120(1):12-26.

85. Innes GK, Lambrou AS, Thumrin P, Thukngamdee Y, Tangwangvivat R, Doungngern P, et
al. Enhancing global health security in Thailand: Strengths and challenges of initiating a
One Health approach to avian influenza surveillance. One Health. 2022;14:100397.

86. Iamsirithaworn S, Chanachai K, Castellan D. Field Epidemiology and One Health:
Thailand's Experience. In: Yamada A, Kahn LH, Kaplan B, Monath TP, Woodall J, Conti L,
editors. Confronting Emerging Zoonoses: The One Health Paradigm. Tokyo: Springer
Japan; 2014. p. 191-212.

87. Hayman DTS, Adisasmito WB, Almuhairi S, Behravesh CB, Bilivogui P, Bukachi SA, et al.
Developing One Health surveillance systems. One Health. 2023;17:100617.

88. Ryser-Degiorgis M-P. Wildlife health investigations: needs, challenges and
recommendations. BMC Veterinary Research. 2013;9(1):223.

89. Schilling A-K, Mazzamuto MV, Romeo C. A Review of Non-Invasive Sampling in Wildlife
Disease and Health Research: What's New? Animals. 2022;12(13):1719.

90. Pepin KM, Brown VR, Yang A, Beasley JC, Boughton R, VerCauteren KC, et al. Optimising
response to an introduction of African swine fever in wild pigs. Transbound Emerg Dis.
2022;69(5):e3111-e27.

91. Sangrat W, Thanapongtharm W, Poolkhet C. Identification of risk areas for foot and
mouth disease in Thailand using a geographic information system-based multi-criteria
decision analysis. Preventive Veterinary Medicine. 2020;185:105183.