Host-encoded DNA methyltransferases modify the epigenome and host tropism of invading phages

Michiko Takahashi[1,¶,*], Satoshi Hiraoka[2,¶], Yuki Matsumoto[3], Rikako Shibagaki[1], Takako Ujihara[4], Hiromichi Maeda[1], Satoru Seo[1], Keizo Nagasaki[3], Hiroaki Takeuchi[5], Shigenobu Matsuzaki[6]

[1]Department of Surgery, Kochi Medical School, Kochi University, Nankoku, Kochi, Japan
[2]Research Center for Bioscience and Nanoscience (CeBN), Research Institute for Marine Resources Utilization, Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka, Kanagawa, Japan
[3]Faculty of Agriculture and Marine Science, Kochi University, Nankoku, Kochi, Japan
[4]Science Research Center, Kochi University, Nankoku, Kochi, Japan
[5]Department of Medical Laboratory Sciences, Health and Science, International University of Health and Welfare Graduate School, Narita, Chiba, Japan
[6]Faculty of Health Sciences, Kochi Gakuen University, Kochi, Kochi, Japan
*Correspondence: michikot@kochi-u.ac.jp
¶These authors contributed equally to this work.

SUMMARY

Restriction modification (RM) systems are ubiquitous bacterial defense systems; however, some phages evade RM system and adapt to their bacterial hosts. In such cases, phages are thought to stochastically acquire DNA methylation from host-encoded DNA methyltransferases (MTases), facilitating host adaptation. However, no studies have directly compared the methylomes of host bacteria and their infecting phages. Here, we demonstrate the epigenetic landscape of adapted phages with diverse infection histories, focusing on the broad host-range phage KHP30T as its adapts to three *Helicobacter pylori* strains. Using a multistage infection system, we observed that the adapted phages displayed significantly high titers against the last infected *H. pylori* strain, suggesting an attendant change in host tropism. Single-molecule real-time sequencing revealed that methylated motifs were predominantly shared between the adapted phages and their most recent host. Our findings enhance our understanding of epigenetic phage–host interactions, which have significant implications for microbial ecology.

KEYWORDS

Epigenome, methyltransferases, phage adaptation, phage-host interaction, phase variation, restriction-modification system, SMRT sequencing, *Helicobacter pylori*

INTRODUCTION

Epigenetic modifications regulate the activity of specific genes and cellular functions without altering DNA sequences. In eukaryotes, epigenetic modifications play a major role in various biological processes through modulating chromatin structure and organization.[1] In particular, DNA methylation, catalyzed by methyltransferases (MTases), is a key epigenetic signal discovered early in studies of DNA as genetic material.[2,3] In mammals, DNA methylation controls gene expression and chromatin structure, contributing to various cellular processes, including cell differentiation, genome imprinting, and X chromosome inactivation.[4]

As bacteria lack essential elements such as histones to structurally modify DNA, DNA methylation serves as a primary epigenetic regulator. [2] In prokaryotes, DNA methylation generally occurs in three forms: N6-methyladenine (m6A) and C5-methylcytosine (m5C), both also found in eukaryotes, and N4-methylcytosine (m4C), specific to bacteria and archaea.[2,5,6] Prokaryotic DNA methylation regulates multiple biological processes, including cell cycle, gene expression, virulence, and host defense.[6]

Historically, one of the most well-studied prokaryotic systems involved in DNA methylation is the restriction modification (RM) system that defends against foreign DNA such as conjugative plasmids and phages.[7–9] Most of these systems comprise an MTase and a restriction endonuclease (REase) that recognizes a specific target sequence for cleavage. These RM systems are classified into Types I, II, and III based on their subunit composition, cofactor requirements, recognition sequence structure, and mode of action.[10] REases interact with their target DNA sequences to cleave the DNA, whereas MTases methylate these recognition sites to protect them from cleavage.[10] Consequently, host genomic DNA is methylated by endogenous MTases and protected against REase digestion, whereas unmethylated foreign DNA is cleaved by REases. In contrast to these three major groups, Type IV systems consist solely of REase(s)

that recognize and cleave methylated DNA.[10,11]

Methods for detecting DNA methylation have been particularly well developed for

eukaryotic m5C, because the biological importance of m5C in mammalian cells over recent

decades.[12] However, these techniques are not directly applicable to bacterial methylomes, where

m6A and m4C are more prevalent. Several approaches, such as restriction enzyme assays, and

bisulfite sequencing, have been used to detect bacterial methylomes with higher prevalences of

m6A and m4C; however, these methods have some limitations: restriction enzyme assays cannot

detect unknown motifs, Sanger sequencing has low throughput, and bisulfite sequencing cannot

detect m6A and m4C.[13] Therefore, further development of DNA methylation detection methods

applicable to bacterial methylomes is needed.

Recent advances in sequencing technologies, including single-molecule real-time

(SMRT) sequencing have addressed this problem. SMRT sequencing was the first to enable

simultaneous detection of all three major forms of prokaryotic DNA methylation, leading to

extensive analysis of bacterial and archaeal methylomes and significant expansion of REBASE,

a database of information about components of RM systems.[13–18] In addition, methylome analysis

has been extended to viruses and phages.[19–22] Recent studies revealed that some DNA viruses

encode MTases in their genomes and use them as part of their replication strategy.[21,23,24] In

contrast, viruses that do not encode MTases can stochastically acquire DNA methylation from

host-encoded MTases during their adaptation to the host strain. Experiments with microbes

possessing well-characterized RM systems such as *Escherichia coli, Bacillus subtilis,*

Haemophilus spp., and *Salmonella* spp. suggest that phage genomes are modified by host-

encoded MTases during infection, allowing adapted phages to develop a "protective suit" against

the host RM systems.[25] However, no study has directly compared the methylomes of hosts and

phages to determine whether adapted phages acquire DNA methylation from host-encoded

MTases.

Here, we demonstrate the epigenomic landscape of phages with distrinct histories of

infection. To model bacteria-phage interactions mediated by host-encoded MTases, we

established a multistage infection system and generated nine adapted phages with different

infection histories using three strains of *Helicobacter pylori* and the *H. pylori* phage KHP30T. The

adapted phages showed high infectious titers against the strains of *H. pylori* infected in the last

generation but low titers against other strains. SMRT sequencing revealed that methylated motifs

in the KHP30T genome were shared with those in the last-infected *H. pylori* strains. These findings

suggest that the phage genome is modified by host-encoded MTases, resulting in the phage DNA

evading cleavage by cognate REases. Our results provide direct evidence that adapted phages

stochastically acquire DNA methylation from host bacteria and offer new insights relevant to

phage therapy design.

RESULTS

Titration of adapted phages

We designed multistage infection systems using the KHP30T phage and three *H. pylori* strains

(26695, 3401T, and HPK5) to obtain adapted phages with different infection histories (Figures 1

and S1). Using this strategy, nine adapted phages (K2, K23, K232, K23H, K23H3, K2H, K2H2,

K2H3, and K2H3H) were retrieved. For example, K23 represents a <u>K</u>HP30T phage adapted to *H.*

pylori <u>2</u>6695 in the first stage and then to *H. pylori* <u>3</u>401T in the second stage. Infectious titers

were measured using a plaque assay. Adapted phages showed high titers against the *H. pylori*

strain in the last generation of their adaptation but lower titers against non-adapted or previously

adapted hosts (Figures 2 and S2). For example, K2, which was adapted to *H. pylori* 26695 in the

first stage (Figure 1), showed a significantly higher titer (9.2 $\log_{10}$ plaque forming units [PFU]/mL)

against 26695, but a markedly lower titer against 3401T and HPK5 (3.1 and 3.9 $\log_{10}$ PFU/mL,

respectively) ($p < 0.01$, Tukey–Kramer test). Focusing on second-stage phages, K23 showed a

significantly higher titer (8.9 $\log_{10}$ PFU/mL) against *H. pylori* 3401T than against 26695 and HPK5

(0.8 and 1.8 $\log_{10}$ PFU/mL, respectively) ($p < 0.01$), even though K23 had adapted to 26695 in

the first stage. K2H also showed the same trend: a titer against *H. pylori* HPK5 (8.2 $\log_{10}$ PFU/mL)

that was significantly higher than that against 26695 and 3401T (1.9 and 2.4 $\log_{10}$ PFU/mL,

respectively, $p < 0.01$) (Figure 2).

Figure 1. Overview of multistage infection system. Nine KHP30T phages were prepared by
adapting to three strains of *H. pylori* (26695, 3401T, HPK5) through four infection stages and four
lines. Details of the adaptation procedure are provided in Transparent Methods and Figure S1.

Figure 2. Infectious titer of adapted KHP30T. Orange, blue, and green bars represent KHP30T

titers against *H. pylori* 26695, 3401T, and HPK5, respectively. Error bars represent standard

deviation (*n* = 3).

Similarly, third-stage adapted phages showed titers against their most recent host strain

that were higher than those against other strains. For example, K232 and K2H2 (lines 1 and 3 in

Figure 1, respectively) showed significantly higher titers against *H. pylori* 26695 (9.6, and 8.2 $\log_{10}$

PFU/mL, respectively) than those against the other two strains (*p* < 0.01). A similar trend was

observed in the fourth stage: K23H3 (line 2) showed a higher titer against 3401T (8.6 $\log_{10}$

PFU/mL) than that against HPK5 (1.8 $\log_{10}$ PFU/mL) (*p* < 0.01). K2H3H (line 4) showed a higher

titer against *H. pylori* HPK5 (9.2 $\log_{10}$ PFU/mL) than that against 3401T (3.1 $\log_{10}$ PFU/mL, *p* <

0.01) (Figure 2).

When comparing between adapted phages, the infectious titers against each of the last

host strains were generally similar. For example, K2, K232, and K2H2, which all shared *H. pylori*

26695 as their last host, showed titers >8.0 $\log_{10}$ PFU/mL against 26695 (range: 8.2–9.6 $\log_{10}$

PFU/mL) but significantly lower titers against other strains (range: 3.1–3.9 $\log_{10}$ PFU/mL).

Likewise, K23 and K23H3 showed significantly higher titers against 3401T (range: 8.5–8.6 $\log_{10}$

PFU/mL) than those against any other strain. Phages K2H, K23H, and K2H3H also showed

significantly higher titers against HPK5 (8.2–9.2 $\log_{10}$ PFU/mL) than those against the others

(Figure 2).

Additionally, we measured the infectious titers of the three adapted phages (K2, K23,

and K2H) after large-scale cultivation to evaluate whether host tropism changed with passaging.

The reproliferated phages exhibited the same trends as the original phages across all host

combinations; titers were highest against the most recent host but were lower against the other

hosts, including those previously adapted strains (Figure S3). Taken together, these results

suggest that the generation gap during reproliferation did not significantly affect the infectivity of

the adapted phages in this study.

Genomic analysis of host H. pylori strains and adapted phages

SMRT sequencing of two *H. pylori* strains (3401T and HPK5) and three adapted phages (K2, K23,

and K2H) was performed to determine the epigenomes of both host strains and adapted phages

during infection. Genomic and epigenomic data for *H. pylori* 26695 were retrieved from the

National Center for Biotechnology Information (NCBI) RefSeq database (accession no.

CP003904.1) and a previously published report.[26] The complete circular genomes of *H. pylori*

3401T and HPK5 were obtained by de novo assembly using high fidelity (HiFi) reads (Table S1).

The general characteristics of these genomes (e.g., genome size, coding sequence [CDS]

number, and GC content) are similar to those of other *H. pylori* genomes deposited in the NCBI

database. The reconstructed *H. pylori* 3401T genome showed >99.99% average nucleotide

identity (ANI) with the ancestral strain *H. pylori* 3401 (AP024599.1).[27] Minor differences in the

sequences most likely arose from genomic mutations during serial subculturing and/or from

different sequencing technologies and bioinformatic software tools. No extrachromosomal content

was observed in any of the strains.

Using functional annotation with REBASE, we predicted 32–36 MTase and 16–21

REase genes in the three *H. pylori* genomes (Tables S1 and S2). The most abundant MTase type

was Type II, whereas Types I and III were nearly equivalent. A systematic search for phage

defense systems in the genomes revealed 14, 9, and 14 RM systems in *H. pylori* 26695, 3401T,

and HPK5, respectively (Table S3). More than half of the detected MTases were orphaned. The

most common modification was m6A, followed by m5C and m4C. Although it has been reported

that DNA methylation is also involved in mediating other defense systems (e.g., BREX[28] and

DISARM[29]), no other systems associated with DNA modification have been identified. In addition

to modification-mediated defense systems, six to nine other systems (e.g., AbiL[30] and

dynamins[31,32]) were identified in the three strains (Table S3).

Mapping KHP30T reads to the original KHP30 phage genome (AB647160.1)[33] identified

an approximately 2-kilobase (kb) insertion in all three phages. No single nucleotide

polymorphisms or short insertions/deletions (indels) were detected. The insertion matched a

segment of the *H. pylori* 3401 genome, indicating it may have been transferred from the host

during serial passaging. This insertion likely does not significantly affect the general

characteristics of the phage, such as infection efficiency or host tropism, consistent with the

titration results above (Figure 2) and previous reports.[33–35] The complete circular genome of

KHP30T was reconstructed using HiFi reads from the K2 phages (Table S1). Read mapping

analysis showed no obvious genomic variations fixed within the phage population. No MTase or

REase genes, as well as phage defense systems, were detected in the phage genome.

Epigenomic analysis of H. pylori strains and adapted phages

We successfully identified numerous base modifications across both the *H. pylori* and phage

genomes (Figures 3 and S4, Table S4). From motif predictions, we detected 15 and 20 motifs in

the *H. pylori* 3401T and HPK5 genomes, respectively, as well as 17 motifs in *H. pylori* 26695, the

latter retrieved from a previous study[26] (Table S4). Among these motifs, the most abundant

modification type was m6A, followed by m4C and m5C. There was a slight discrepancy in the

modification types of the encoded MTases, possibly due to the SMRT sequencing's low detection

sensitivity for m5C modifications. As expected, the detected m5C motifs showed low modification

ratios (<50.0%) across all three *H. pylori* strains. Only two motifs, C**A**TG and G**A**TC (with the

modified bases in bold and their reverse complements underlined), were shared among all three

strains. However, 28 of 39 (72%) motifs were present in either of the 3 strains, indicating high

diversity in methylation patterns across these strains. This trend aligns with previous findings that

H. pylori strains possess highly divergent combinations of modified motifs.[26,36]

Next, we compared the methylation patterns in *H. pylori* strains and their adapted

phages. Overall, the methylated motifs were shared between pairs of *H. pylori* strains and their

respective adapted phages (Figure 3). For example, among the motifs detected in *H. pylori*

HPK5, with the exception of m5C motifs, 17 of 19 were highly modified in phage K2H (motif

modification ratios >68.9% and >63.6% in the host and phage genomes, respectively). Similarly,

12 of 13 motifs were shared between *H. pylori* 3401T and phage K23 with ratios >73.9% and

>41.7%, respectively. In addition, 12 of 14 motifs were concordantly methylated between *H. pylori*

26695 and phage K2, with ratios >95.2% and >71.3%, respectively. In contrast, sets of methylated

motifs were not shared between non-corresponding host-phage pairs.

Figure 3. Methylomes of adapted phages and *H. pylori* strains. Methylated motifs were predicted from the three *H. pylori* strains. Motifs are shown separately according to methylation type. Phages and *H. pylori* strains are colored in red and black, respectively, in the bottom text. Text on the left side indicates the methylated motifs with methylated nucleotide position and methylation type. The top dendrogram was calculated using Euclidean distance and Ward method. For *H. pylori* 26695, methylated motifs and ratios were retrieved from the previous study reported by Krebes et al. (2014), and gray cells indicate motifs for which no data are available. Some motifs showed comparatively high modification ratios due to motif overlap. For example, the modification

ratio of CGR**A**T (R=A/G) in *H. pylori* 3401T is explained by motif overlapping with CGR**A**, which

was detected from epigenomic analysis of the strain. In addition, CGW**A**G (W=A/T) in *H. pylori*

3401T partly overlapped with CGR**A**, i.e., CGA**A**G. A motif G**A**GG in *H. pylori* HPK5 also

overlapped with the detected G**A**GGA from the strain, and in contrast, G**A**GGA in K2 overlap with

G**A**GG reported from *H. pylori* 26695 previously. Among all seven CCGC**A**T sites in K23, two and

one overlapped with other motifs C**A**TG (i.e., CCGC**A**TG) and **A**TCC (i.e., CCGC**A**TCC),

respectively. Hence, CCGC**A**T showed a comparatively higher modification ratio in K23.

Notably, several motifs were identified that were highly methylated in the host genomes

but showed minimal methylation in the adapted phage genomes (Figure 3). For instance, the

m6A-type motif G**A**TGG was methylated in the *H. pylori* 3401T genome with a high modification

ratio (73.9%), but was undetected in the K23 genome. Similarly, the motif CCTTG**A** was highly

modified in *H. pylori* HPK5 (81.1%) but not in K2H (7.1%). Additionally, motifs A<u>T</u>TA**A**T and

CGR**A**T have been reported to be methylated at high modification ratios (98.5% and 98.9%,

respectively) in *H. pylori* 26695[26] but showed much lower modification in phage K2 (0.0% and

22.6%, respectively).

Phase-variable RM systems and associated genes

In the *H. pylori* HPK5 genome, a G9-tract was identified within the HPYHPK5_00170−00190

region (Figure S5A), displaying high sequence similarity to the M.Hpy300X Type III m6A MTase

gene. This gene is known to induce methylome variation through simple sequence repeats

(SSRs).[37] The basal state of the poly-G tract results in a stop codon and a truncated protein

product. However, the addition of a single nucleotide to the tract results in a frameshift, generating

a fully functional fusion protein. Similar to the Hpy300X system, this tract likely modulates MTase

activity, suggesting that HPYHPK5_00150-00160 functions as a phase-variable Type III RM

system in conjunction with a Type III REase (HPYHPK5_00170). Similar structures were also

predicted in *H. pylori* 26695, specifically in Type III RM systems AFV42588−42590 (HpyAXVII)

(Figure S5B) and AFV42742−42744 (Figure S5C), with the former previously characterized.[38]

In the *H. pylori* 3401T genome, HPY3401T_09270 and HPY3401T_09280 harbor the

C19- and C18-tracts, respectively (Figure S5D). These sequences showed the highest similarity

to the sequences of genes associated with the HpyAXVI system (89.8% and 89.0% identity to

HpyAXVI-mut2 and HpyAXVI-mut1, respectively). The HpyAXVI system, composed of Type II RM

genes (REase and MTase fusion genes), was originally identified in *H. pylori* 26695[26] (Figure S5E).

This system is inactive when a stop codon separates the two CDSs fillowing the first tract but

becomes active when a frameshift near the tract fuses the two CDSs. When active, the tract

produces a truncated protein that regulates MTase sequence specificity. Therefore,

HPY3401T_09270 and HPY3401T_09280 are likely phase-variable Type II RM systems. In

addition, HPYHPK5_13900 and HPYHPK5_13910 contain C14- and G15-tracts, respectively

(Figure S5F). The former showed the highest sequence similarity (88.2%) to Hpy300XI found in

H. pylori BCM-300 strain. This system uses two tracts to facilitate phase variation, likely operating

similarly to HpyAXVI[26]. Thus, HPYHPK5_13900 and HPYHPK5_13910 may also comprise

phase-variable Type II RM systems.

Beyond MTases, certain SSRs were associated with REase and specificity (S) subunit

genes. In all *H. pylori* genomes, four tracts were identified within Type I REase CDSs (Figure S5G,

S5H, S5I, and S5J). In contrast, cognate MTases and S genes were not associated with SSRs.

As Type I MTases require an S subunit complex for functionality, these tracts likely impact only

the cleavage activity of Type I RM systems, leaving methylation functions unaffected. In addition,

a C17-tract was found in the HPY3401T_11810 REase, part of a Type III RM system with the

HPY3401T_11820 MTase (Figure S5K). While Type III MTases function independently, REases

require a cognate MTase for cleavage. Tandem short-sequence repeats, including SSRs, are

common in Type III MTase genes and act as ON/OFF switches regulating phase variation,[39] as

also observed in this study (Figure S5A, S5B, and S5C). Therefore, this C17-tract may specifically

regulate only REase activity, while the cognate MTase may be constituitively active. AFV41850

from *H. pylori* 26695 and HPY3401T_02460 from 3401T showed the highest sequence similarity

to Type IV REase EcoCTGmrSD and possessed T15- and T19-tracts located 47 and 46 base

paris (bp) upstream of their CDSs, respectively (Figure S5L and S5M). Type IV REases cleave

methylated DNA strands in contrast to other REase types, suggesting that these tracts likely

regulate the ON/OFF switch for Type IV RM systems. Finally, consistent with a previous study's

findings, we confirmed that G14-tracts within the S gene comprise a BcgI-like Hpy99XXII system

accompanied by an upward-adjusted RM gene, though its function is still unclear[26] (Figure S5N).

DISCUSSION

DNA methylation-mediated biological processes affect a wide range of microbial ecologies,

including phage-host interactions. There is growing interest in epigenomic systems of diverse

prokaryotes and viruses, given their importance in microbial physiology, genetics, evolution, and

ecology.[2,40,41] To investigate the role of methylation in phage-host interactions, *H. pylori* and phage

KHP30 were chosen as model organisms.

H. pylori is a major human pathogen that causes chronic inflammation in the stomach

and gastric cancer. It harbors multiple RM systems with highly variable composition, resulting in

diverse methylomes across strains.[26,36,42–44] However, epigenomic evaluation has not been

performed to determine whether the MTases encoded in the *H. pylori* genome play a role against

invading phages. We previously reported that KHP30, a double-stranded DNA phage belonging

to the *Schmidvirus* genus, infects *H. pylori*[33–35] and has potential as a therapeutic agent against

antibiotic-resistant *H. pylori*, owing to its broad host range.[34] However, a significant barrier to

phage therapy for *H. pylori* lies in the limited understanding of its defense mechanisms against

KHP30 infection.

To clarify the relationship between DNA methylation and phage-host interactions, we

designed a multistage infection system (Figure 1) and analyzed the methylomes of three *H. pylori*

strains (26695, 3401T, and HPK5) along with three adapted KHP30T phages (K2, K23, and K2H)

with varying infection histories. The adapted phages showed high titers against their most recent

host strain but significantly lower titers against other strains, even if they had previously adapted

to them (Figure 2). Considering the short generation time during cultivation, gene expression

disruptions or enzymatic inactivation from mutations were unlikely to affect tropism significantly.

As expected, the reproliferated phages consistently showed high titers only against the latest

infected host (Figure S3).

Our epigenomic analysis revealed that each *H. pylori* strain contained distinct sets of

MTase genes, many positioned near REase genes within the RM systems (Tables S1 and S2).

Adapted phages were found to share methylated motifs with their final adapted host strain,

indicating that phage epigenomes were modified by host MTases during infection. Taken together,

our work demonstrates that adapted phages acquire DNA methylation from their host to evade

from the host's RM systems, thereby increasing infection efficiency, as has been previously

hypothesized.[45] To the best of our knowledge, this is the first study to directly observe epigenomic

changes in the phage genome during infection (Figure S4). Although methylated motifs were

shared between adapted phages and their host, exceptions were noted. Specifically, several

motifs were methylated in the host genome but not in the adapted phage (Figure 3). These

exceptions may partially be due to phase variations in several RM systems of *H. pylori* (Figure

S5).

Phase variation is a random and reversible switching mechanism in gene expression

that generates transcriptional, translational, and/or phenotypic diversity within bacterial

populations.[46] Bacterial MTases within RM systems often undergo phase variation, resulting in

heterogeneous cells with varied RM activity within a clonal population.[47] This heterogeneity

enables phages to selectively infect cells with reduced RM activity, as their epigenomes are

methylated by the MTases in these hosts. A primary mechanism regulating phase variation is the

presence of SSRs within or upstream of RM genes, with phase-variable ON/OFF switching

occuring in the former, resulting in reversible changes in the number of SSRs.[47–51] Our analysis

found several phase-variable regulons or phasevarions associated with RM genes in all three *H.*

pylori genomes (Figure S5). Further single-cell evaluation such as combining

immunofluorescence imaging to track phase-variated proteins[52] with fluorescence *in situ*

hybridization for phage detection, are necessary to clarify the link between phase variation and

phage variation. Moreover, bioinformatic tools for detecting modifications needs to be improved,

as current modification detection tools do not account for phase-variable epigenomes and instead

treat modifications on a binary, population-wide manner. Another possible explanation for the

methylated motifs found exclusively in the host genome is the potential presence of anti-RM

systems in the phage that block host MTase activity. For example, DarA and DarB in P1 phages

protect recognition sites on their genome from host RM activity[53,54]. Similarly, Ocr in T7 phages

mimics DNA structure to block the DNA-binding groove of Type I REases/MTases.[55] However, no

genes associated with known anti-RM systems[56] were found in the KHP30T genome.

Defense systems are involved in the evolutionary arms race between phages and

hosts.[57] Epigenomic modifications significantly influence their co-evolution and promote microbial

diversification.[58,59] Our findings revealed that the phage epigenome was altered by host bacteria

during infection, enhancing the phage's ability to reinfect the same host strain (Figures 2 and 3).

To counter the epigenomic synchronization of invading phages and avoid extinction of the entire

population by phage persistence, some bacteria employ variable phage-defense systems. For

example, many species of gut bacteria, such as *Bacteroides fragilis* and *H. pylori*, show strain-

specific variations in gene sets associated with phage defense, including MTase genes.[60–62] While

gene acquisition or deletion rely on genomic events such as horizontal gene transfer and

recombination, such events occur at infrequently. Alternatively, a more dynamic strategy by

bacteria involves phase variation to prevent phage circulation. For example, *Vibrio cholerae*

modulates the expression of the lipopolysaccharide O1 antigen through phase variation to escape

predation by O1-specific phages.[63] In *H. pylori*, inter-strain variation in methylomes caused by

MTase phase variation has been reported.[36,37] Therefore, antagonistic coevolution between

prokaryotes and phages, along with associated epigenetic changes, is of great interest for

understanding the diversity within microbial ecosystems.

In summary, our study demonstrates that phages can overcome host RM systems by

acquiring methylation modifications from host-encoded MTases during infection. Adapted phages

showed significantly higher titers against their most recent host strain, likely due to neutralization

of the host RM system through acquired methylation. Our results enhance our understanding of

DNA methylation-mediated phage–host interactions and their impact on microbial ecology across

diverse environments.

Our approach provides new insights into the design of therapeutic phage cocktails.

Phage therapy holds promise across various fields, including human and plant diseases,[64–67]

owing to phages' advantages such as host specificity, self-amplification, biofilm degradation, and

low toxicity to humans.[68,69] One of the primary challengess in phage therapy is bacterial

resistance; however, phage cocktails, as shown in studies on *E. coli* O157:H7 and *Klebsiella*

pneumoniae, offer a strategy to combat resistance by incorporating diverse phage variants.[70,71]

We propose that a cocktail of adapted phages, each with distinct methylation motifs, could

improve the chances of overcoming host modification-mediated defense systems, such as RM

systems. Further research to understand the mechanism of phage adaptation and evaluate their

therapeutic potential will contribute to the development of tailored phage cocktails for medical and

industrial applications.

Limitation of the study

This study has few limitations. First, we did not experimentally verify the activity and specificity of

the MTases and REases identified, leaving uncertainty about which RM systems are actively

involved in phage defense and the specific MTases responsible for the modified motifs. Second,

although SMRT sequencing technology efficiently identifies m6A and m4C modifications,[26] it has

limited sensitivity for detecting m5C modifications, one of the most common methylation types in

nature after m6A and m4C. In total, five m5C motifs were detected across all three *H. pylori* strains

(Table S4), but they showed low modification ratios (Figure 3, Table S4), presumably due to the

low detection accuracy for this modification type using current SMRT technology. In addition to

methylation, other epigenetic modifications, such as phosphorothioate modification, have recently

been linked to phage defense.[72] Expanding sequencing technologies to detect a wider range of

DNA modifications is crucial for a comprehensive understanding of epigenomic functions. Third,

although HiFi reads are highly accurate (>99.9%), sequencing of homopolymer regions remains

challenging,[73–75] complicating estimatations of tract diversity. Nonetheless, given the high genome

coverage achieved in this study (Table S1), the tracts associated with RM genes (Figure S5) likely

represent the predominant phase variation state in the populations studied.

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the lead contact, Michiko Takahashi (michikot@kochi-u.ac.jp).

Data availability

The assembled genomes were deposited in DDBJ/ENA/GenBank (S1 Table). All data are

registered under BioProject ID PRJDB17787

[https://ddbj.nig.ac.jp/resource/bioproject/PRJDB17787]. Any additional information required to

reanalyze the data reported in this paper is available from the lead contact upon request.

METHODS

All methods can be found in the Transparent Methods.

ACKNOWLEDGMENTS

The authors are greatful to Waka Ishida, Yusaku Funaoka, and Takahiro Ishikawa for experimental

support, and to Daichi Morimoto for helpful comments on this work. This study was supported by

JSPS KAKENHI (grant numbers JP20K15582 and JP22K08875), ACT-X, Japan Science and

Technology Agency (grant numbers JPMJAX21BD and JPMJAX22BK), and the budget of Kochi

University Hospital (Japan). The experiments were conducted at the Division of Biological

Research, Science Research Center, Kochi University. Computations were partially performed on

the supercomputer system at the Institute for Chemical Research, Kyoto University, and the NIG

supercomputer system at ROIS National Institute of Genetics, and the Data Analysis and the

Earth Simulator systems at JAMSTEC.

AUTHOR CONTRIBUTIONS

MT designed the study, acquired funding, performed the experiments, and drafted the manuscript.

SH designed the study, performed the bioinformatics analyses, and wrote the manuscript. YM,

RS, and TU performed the experiments. HM acquired funding and provided resources. SS, KN,

and HT supervised the project. SM conceived the original idea, acquired funding, and supervised

the project. All authors have read and approved the final manuscript.

DECLARATION OF INTERESTS

The authors declare no conflicts of interest associated with this manuscript.

SUPPLEMENTAL INFORMATION

Figure S1. Overview of preparation procedure for adapted phages.

Figure S2. Tukey-Kramer pairwise comparisons of phage titers. Corresponding titers are

shown in Figure 2. Combination of phage and host strain are indicated as 'adapted phage': 'host

strain used for titration' format in the x-axis: for example, 'K2:2' means a titer of K2 against *H.*

pylori 26695. Solid black bar within each box represents median. Upper alphabet summarizes the

results of the Tukey-Kramer test; letters denote significant differences ($p < 0.01$) in mean titers

with groups a > b > c > d > e > f.

Figure S3. Infectious titer of reproliferated phages. (A) Orange, blue, and green bars

represent titer of reproliferated KHP30T phages to *H. pylori* 26695, 3401T, and HPK5, respectively.

Error bars indicate standard deviation (n = 3). (B) Tukey-Kramer pairwise comparisons of

reproliferated phage titers. Phage and host combinations are indicated in the same format as

Figure S2. Infectious titer of the 'Original phages' is derived from Figure 2. Solid black bar within

each box represents median. Upper alphabet summarizes the results of the Tukey-Kramer test;

letters denote significant differences ($p < 0.01$) in mean titers with groups a > b > c > d > e > f.

Figure S4. Methylomes of adapted phages. Circos plot showing the distribution of DNA

methylation. Orange, blue, and red bars represent m4C, m5C, and m6A modifications,

respectively. K2 and K23 were sequenced using PacBio Sequel II in CCS mode, whereas K2H

was sequenced using PacBio Sequel in CLR mode.

Figure S5. Genetic organization of RM system genes associated with SSRs. Homopolymeric

tracts inside or upstream of RM genes were specified as SSRs. RM gene types are indicated by

color and text inside to each CDS. CDSs not assigned to RM system genes are colored in gray.

Table S1. Descriptions and statistics of *H. pylori* strains and KHP30T phages.

Table S2. Detected RM system genes in *H. pylori* strains.

Table S3. Predicted phage defense systems in *H. pylori* strains.

Table S4. Methylated motifs predicted from *H. pylori* strains.

Transparent Methods

REFERENCES

1. Liyanage, V., Jarmasz, J., Murugeshan, N., Del Bigio, M., Rastegar, M., and Davie, J. (2014). DNA modifications: Function and applications in normal and disease states. Biology *3*, 670–723.

2. Sánchez-Romero, M.A., and Casadesús, J. (2020). The bacterial epigenome. Nat. Rev. Microbiol. *18*, 7–20.

3. Avery, O.T., Macleod, C.M., and McCarty, M. (1944). Studies on the chemical nature of the substance inducing transformation of pneumococcal types : Induction of transformation by a desoxyribonucleic acid fraction isolated from Pneumococcus Type III. J. Exp. Med. *79*, 137–158.

4. Razin, A., and Riggs, A.D. (1980). DNA methylation and gene function. Science *210*, 604–610.

5. Weigele, P., and Raleigh, E.A. (2016). Biosynthesis and function of modified bases in bacteria and their viruses. Chem. Rev. *116*, 12655–12687.

6. Seong, H.J., Han, S.-W., and Sul, W.J. (2021). Prokaryotic DNA methylation and its functional roles. J. Microbiol. *59*, 242–248.

7. Labrie, S.J., Samson, J.E., and Moineau, S. (2010). Bacteriophage resistance mechanisms. Nat. Rev. Microbiol. *8*, 317–327.

8. Ando, T., Ishiguro, K., Watanabe, O., Miyake, N., Kato, T., Hibi, S., Mimura, S., Nakamura, M., Miyahara, R., Ohmiya, N., et al. (2010). Restriction-modification systems may be associated with *Helicobacter pylori* virulence. J. Gastroenterol. Hepatol. *25*, S95-8.

9. Arber, W. (1974). DNA modification and restriction. Prog. Nucleic Acid Res. Mol. Biol. *14*, 1–37.

10. Ershova, A.S., Rusinov, I.S., Spirin, S.A., Karyagina, A.S., and Alexeevski, A.V. (2015). Role of restriction-modification systems in prokaryotic evolution and ecology. Biochemistry *80*, 1373–1386.

11. Mruk, I., and Kobayashi, I. (2014). To be or not to be: regulation of restriction-modification systems and other toxin-antitoxin systems. Nucleic Acids Res. *42*, 70–86.

12. Plongthongkum, N., Diep, D.H., and Zhang, K. (2014). Advances in the profiling of DNA modifications: cytosine methylation and beyond. Nat. Rev. Genet. *15*, 647–661.

13. Beaulaurier, J., Schadt, E.E., and Fang, G. (2019). Deciphering bacterial epigenomes using modern sequencing technologies. Nat. Rev. Genet. *20*, 157–172.

14. Roberts, R.J., Vincze, T., Posfai, J., and Macelis, D. (2015). REBASE--a database for DNA restriction and modification: enzymes, genes and genomes. Nucleic Acids Res. *43*, D298-9.

15. Forde, B.M., McAllister, L.J., Paton, J.C., Paton, A.W., and Beatson, S.A. (2019). SMRT sequencing reveals differential patterns of methylation in two O111:H- STEC isolates from a

479   hemolytic uremic syndrome outbreak in Australia. Sci. Rep. *9*, 9436.

480   16.   Murray, I.A., Clark, T.A., Morgan, R.D., Boitano, M., Anton, B.P., Luong, K., Fomenkov, A.,

481   Turner, S.W., Korlach, J., and Roberts, R.J. (2012). The methylomes of six bacteria. Nucleic

482   Acids Res. *40*, 11450–11462.

483   17.   Hiraoka, S., Okazaki, Y., Anda, M., Toyoda, A., Nakano, S.-I., and Iwasaki, W. (2019).

484   Metaepigenomic analysis reveals the unexplored diversity of DNA methylation in an

485   environmental prokaryotic community. Nat. Commun. *10*, 159.

486   18.   Roberts, R.J., Vincze, T., Posfai, J., and Macelis, D. (2023). REBASE: a database for DNA

487   restriction and modification: enzymes, genes and genomes. Nucleic Acids Res. *51*, D629–

488   D630.

489   19.   Coy, S.R., Gann, E.R., Papoulis, S.E., Holder, M.E., Ajami, N.J., Petrosino, J.F., Zinser, E.R.,

490   Van Etten, J.L., and Wilhelm, S.W. (2020). SMRT sequencing of Paramecium bursaria

491   chlorella virus-1 reveals diverse methylation stability in adenines targeted by restriction

492   modification systems. Front. Microbiol. *11*, 887.

493   20.   Sun, C., Chen, J., Jin, M., Zhao, X., Li, Y., Dong, Y., Gao, N., Liu, Z., Bork, P., Zhao, X.-M.,

494   et al. (2023). Long-read sequencing reveals extensive DNA methylations in human gut

495   phagenome contributed by prevalently phage-encoded methyltransferases. Adv. Sci. *10*,

496   e2302159.

497   21.   Jeudy, S., Rigou, S., Alempic, J.-M., Claverie, J.-M., Abergel, C., and Legendre, M. (2020).

498   The DNA methylation landscape of giant viruses. Nat. Commun. *11*, 2657.

499   22.   Hiraoka, S., Sumida, T., Hirai, M., Toyoda, A., Kawagucci, S., Yokokawa, T., and Nunoura, T.

500   (2022). Diverse DNA modification in marine prokaryotic and viral communities. Nucleic Acids

501   Res. *50*, 1531–1550.

502   23.   Hoelzer, K., Shackelton, L.A., and Parrish, C.R. (2008). Presence and role of cytosine

503   methylation in DNA viruses of animals. Nucleic Acids Res. *36*, 2825–2837.

504   24.   Wilson, W.H., Van Etten, J.L., and Allen, M.J. (2009). The Phycodnaviridae: the story of how

505   tiny giants rule the world. Curr. Top. Microbiol. Immunol. *328*, 1–42.

506   25.   Krüger, D.H., and Bickle, T.A. (1983). Bacteriophage survival: multiple mechanisms for

507   avoiding the deoxyribonucleic acid restriction systems of their hosts. Microbiol. Rev. *47*, 345–

508   360.

509   26.   Krebes, J., Morgan, R.D., Bunk, B., Spröer, C., Luong, K., Parusel, R., Anton, B.P., König,

510   C., Josenhans, C., Overmann, J., et al. (2014). The complex methylome of the human gastric

511   pathogen *Helicobacter pylori*. Nucleic Acids Res. *42*, 2415–2432.

512   27.   Takahashi, M., Matsumoto, Y., Ujihara, T., Maeda, H., Hanazaki, K., Nagasaki, K., Takeuchi,

513   H., and Matsuzaki, S. (2021). Complete genome sequence of *Helicobacter pylori* strain 3401,

514   a suitable host for bacteriophages KHP30 and KHP40. Microbiol Resour Announc *10*,

515   e0064721.

28. Goldfarb, T., Sberro, H., Weinstock, E., Cohen, O., Doron, S., Charpak-Amikam, Y., Afik, S., Ofir, G., and Sorek, R. (2015). BREX is a novel phage resistance system widespread in microbial genomes. EMBO J. *34*, 169–183.

29. Ofir, G., Melamed, S., Sberro, H., Mukamel, Z., Silverman, S., Yaakov, G., Doron, S., and Sorek, R. (2017). DISARM is a widespread bacterial defence system with broad anti-phage activities. Nat. Microbiol. *3*, 90–98.

30. Deng, Y.-M., Liu, C.-Q., and W. Dunn, N. (1999). Genetic organization and functional analysis of a novel phage abortive infection system, AbiL, from *Lactococcus lactis*. J. Biotechnol. *67*, 135–149.

31. Millman, A., Melamed, S., Leavitt, A., Doron, S., Bernheim, A., Hör, J., Garb, J., Bechon, N., Brandis, A., Lopatina, A., et al. (2022). An expanded arsenal of immune systems that protect bacteria from phages. Cell Host Microbe *30*, 1556-1569.e5.

32. Guo, L., Sattler, L., Shafqat, S., Graumann, P.L., and Bramkamp, M. (2022). A bacterial dynamin-like protein confers a novel phage resistance strategy on the population level in *Bacillus subtilis*. MBio *13*. https://doi.org/10.1128/mbio.03753-21.

33. Uchiyama, J., Takeuchi, H., Kato, S.-I., Takemura-Uchiyama, I., Ujihara, T., Daibata, M., and Matsuzaki, S. (2012). Complete genome sequences of two *Helicobacter pylori* bacteriophages isolated from Japanese patients. J. Virol. *86*, 11400–11401.

34. Uchiyama, J., Takeuchi, H., Kato, S.-I., Gamoh, K., Takemura-Uchiyama, I., Ujihara, T., Daibata, M., and Matsuzaki, S. (2013). Characterization of *Helicobacter pylori* bacteriophage KHP30. Appl. Environ. Microbiol. *79*, 3176–3184.

35. Kamiya, R., Uchiyama, J., Matsuzaki, S., Murata, K., Iwasaki, K., and Miyazaki, N. (2022). Acid-stable capsid structure of *Helicobacter pylori* bacteriophage KHP30 by single-particle cryoelectron microscopy. Structure *30*, 300-312.e3.

36. Furuta, Y., Namba-Fukuyo, H., Shibata, T.F., Nishiyama, T., Shigenobu, S., Suzuki, Y., Sugano, S., Hasebe, M., and Kobayashi, I. (2014). Methylome diversification through changes in DNA methyltransferase sequence specificity. PLoS Genet. *10*, e1004272.

37. Nell, S., Estibariz, I., Krebes, J., Bunk, B., Graham, D.Y., Overmann, J., Song, Y., Spröer, C., Yang, I., Wex, T., et al. (2018). Genome and methylome variation in *Helicobacter pylori* with a *cag* pathogenicity island during early stages of human infection. Gastroenterology *154*, 612-623.e7.

38. Prasad, Y., Kumar, R., Chaudhary, A.K., Dhanaraju, R., Majumdar, S., and Rao, D.N. (2019). Kinetic and catalytic properties of M.HpyAXVII, a phase-variable DNA methyltransferase from *Helicobacter pylori*. J. Biol. Chem. *294*, 1019–1034.

39. Rao, D.N., Dryden, D.T.F., and Bheemanaik, S. (2014). Type III restriction-modification

enzymes: a historical perspective. Nucleic Acids Res. *42*, 45–55.

40.   Srikhanta, Y.N., Fox, K.L., and Jennings, M.P. (2010). The phasevarion: phase variation of

type III DNA methyltransferases controls coordinated switching in multiple genes. Nat. Rev.

Microbiol. *8*, 196–206.

41.   Furuta, Y., and Kobayashi, I. (2012). Mobility of DNA sequence recognition domains in DNA

methyltransferases suggests epigenetics-driven adaptive evolution. Mob. Genet. Elements

2, 292–296.

42.   Ando, T., Xu, Q., Torres, M., Kusugami, K., Israel, D.A., and Blaser, M.J. (2000). Restriction-

modification system differences in *Helicobacter pylori* are a barrier to interstrain plasmid

transfer. Mol. Microbiol. *37*, 1052–1065.

43.   Alm, R.A., Ling, L.S., Moir, D.T., King, B.L., Brown, E.D., Doig, P.C., Smith, D.R., Noonan,

B., Guild, B.C., deJonge, B.L., et al. (1999). Genomic-sequence comparison of two unrelated

isolates of the human gastric pathogen *Helicobacter pylori*. Nature *397*, 176–180.

44.   Bubendorfer, S., Krebes, J., Yang, I., Hage, E., Schulz, T.F., Bahlawane, C., Didelot, X., and

Suerbaum, S. (2016). Genome-wide analysis of chromosomal import patterns after natural

transformation of *Helicobacter pylori*. Nat. Commun. *7*, 11995.

45.   Isaev, A.B., Musharova, O.S., and Severinov, K.V. (2021). Microbial arsenal of antiviral

defenses - Part I. Biochemistry *86*, 319–337.

46.   de Vries, N., Duinsbergen, D., Kuipers, E.J., Pot, R.G.J., Wiesenekker, P., Penn, C.W., van

Vliet, A.H.M., Vandenbroucke-Grauls, C.M.J.E., and Kusters, J.G. (2002). Transcriptional

phase variation of a type III restriction-modification system in *Helicobacter pylori*. J. Bacteriol.

184, 6615–6623.

47.   Atack, J.M., Tan, A., Bakaletz, L.O., Jennings, M.P., and Seib, K.L. (2018). Phasevarions of

bacterial pathogens: methylomics sheds new light on old enemies. Trends Microbiol. *26*,

715–726.

48.   Gor, V., Ohniwa, R.L., and Morikawa, K. (2021). No change, no life? What we know about

phase variation in *Staphylococcus aureus*. Microorganisms *9*, 244.

49.   Nahar, N., Tram, G., Jen, F.E.-C., Phillips, Z.N., Weinert, L.A., Bossé, J.T., Jabbari, J.S.,

Gouil, Q., Du, M.R.M., Ritchie, M.E., et al. (2023). *Actinobacillus pleuropneumoniae* encodes

multiple phase-variable DNA methyltransferases that control distinct phasevarions. Nucleic

Acids Res. *51*, 3240–3260.

50.   Moxon, R., Bayliss, C., and Hood, D. (2006). Bacterial contingency loci: the role of simple

sequence DNA repeats in bacterial adaptation. Annu. Rev. Genet. *40*, 307–333.

51.   Bayliss, C.D. (2009). Determinants of phase variation rate and the fitness implications of

differing rates for bacterial pathogens and commensals. FEMS Microbiol. Rev. *33*, 504–520.

52.   Ben-Assa, N., Coyne, M.J., Fomenkov, A., Livny, J., Robins, W.P., Muniesa, M., Carey, V.,

Carasso, S., Gefen, T., Jofre, J., et al. (2020). Analysis of a phase-variable restriction

modification system of the human gut symbiont *Bacteroides fragilis*. Nucleic Acids Res. *48*,

11040–11053.

53. Iida, S., Streiff, M.B., Bickle, T.A., and Arber, W. (1987). Two DNA antirestriction systems of

bacteriophage P1, *darA*, and *darB*: characterization of *darA*- phages. Virology *157*, 156–166.

54. Teklemariam, A.D., Al-Hindi, R.R., Qadri, I., Alharbi, M.G., Ramadan, W.S., Ayubu, J., Al-

Hejin, A.M., Hakim, R.F., Hakim, F.F., Hakim, R.F., et al. (2023). The battle between bacteria

and bacteriophages: a conundrum to their immune system. Antibiotics *12*, 381.

55. Roberts, G.A., Stephanou, A.S., Kanwar, N., Dawson, A., Cooper, L.P., Chen, K., Nutley, M.,

Cooper, A., Blakely, G.W., and Dryden, D.T.F. (2012). Exploring the DNA mimicry of the Ocr

protein of phage T7. Nucleic Acids Res. *40*, 8129–8143.

56. Samson, J.E., Magadán, A.H., Sabri, M., and Moineau, S. (2013). Revenge of the phages:

defeating bacterial defences. Nat. Rev. Microbiol. *11*, 675–687.

57. Hampton, H.G., Watson, B.N.J., and Fineran, P.C. (2020). The arms race between bacteria

and their phage foes. Nature *577*, 327–336.

58. Scanlan, P.D. (2017). Bacteria-bacteriophage coevolution in the human gut: implications for

microbial diversity and functionality. Trends Microbiol. *25*, 614–623.

59. Dimitriu, T., Szczelkun, M.D., and Westra, E.R. (2020). Evolutionary ecology and interplay of

prokaryotic innate and adaptive immune systems. Curr. Biol. *30*, R1189–R1202.

60. O'Connell Motherway, M., Zomer, A., Leahy, S.C., Reunanen, J., Bottacini, F., Claesson, M.J.,

O'Brien, F., Flynn, K., Casey, P.G., Munoz, J.A.M., et al. (2011). Functional genome analysis

of *Bifidobacterium breve* UCC2003 reveals type IVb tight adherence (Tad) pili as an essential

and conserved host-colonization factor. Proc. Natl. Acad. Sci. U. S. A. *108*, 11217–11222.

61. Zhu, A., Sunagawa, S., Mende, D.R., and Bork, P. (2015). Inter-individual differences in the

gene content of human gut bacterial species. Genome Biol. *16*, 82.

62. Yawata, Y., Kiyokawa, T., Kawamura, Y., Hirayama, T., Takabe, K., and Nomura, N. (2019).

Intra- and interspecies variability of single-cell innate fluorescence signature of microbial cell.

Appl. Environ. Microbiol. *85*, e00608-19.

63. Seed, K.D., Faruque, S.M., Mekalanos, J.J., Calderwood, S.B., Qadri, F., and Camilli, A.

(2012). Phase variable O antigen biosynthetic genes control expression of the major

protective antigen and bacteriophage receptor in *Vibrio cholerae* O1. PLoS Pathog. *8*,

e1002917.

64. Pires Diana P., Cleto Sara, Sillankorva Sanna, Azeredo Joana, and Lu Timothy K. (2016).

Genetically engineered phages: a review of advances over the last decade. Microbiol. Mol.

Biol. Rev. *80*, 523–543.

65. Abedon, S.T., Kuhl, S.J., Blasdel, B.G., and Kutter, E.M. (2011). Phage treatment of human

infections. Bacteriophage *1*, 66–85.

66. Balogh, B., Jones, J.B., Iriarte, F.B., and Momol, M.T. (2010). Phage therapy for plant disease
control. Curr. Pharm. Biotechnol. *11*, 48–57.

67. Loc Carrillo, C., Atterbury, R.J., el-Shibiny, A., Connerton, P.L., Dillon, E., Scott, A., and
Connerton, I.F. (2005). Bacteriophage therapy to reduce *Campylobacter jejuni* colonization
of broiler chickens. Appl. Environ. Microbiol. *71*, 6554–6563.

68. Lin, D.M., Koskella, B., and Lin, H.C. (2017). Phage therapy: an alternative to antibiotics in
the age of multi-drug resistance. World J. Gastrointest. Pharmacol. Ther. *8*, 162–173.

69. Loc-Carrillo, C., and Abedon, S.T. (2011). Pros and cons of phage therapy. Bacteriophage *1*,
111–114.

70. Tanji, Y., Shimada, T., Yoichi, M., Miyanaga, K., Hori, K., and Unno, H. (2004). Toward rational
control of *Escherichia coli* O157:H7 by a phage cocktail. Appl. Microbiol. Biotechnol. *64*, 270–
274.

71. Gu, J., Liu, X., Li, Y., Han, W., Lei, L., Yang, Y., Zhao, H., Gao, Y., Song, J., Lu, R., et al.
(2012). A method for generation phage cocktail with great therapeutic potential. PLoS One
7, e31698.

72. Jiang, S., Chen, K., Wang, Y., Zhang, Y., Tang, Y., Huang, W., Xiong, X., Chen, S., Chen, C.,
and Wang, L. (2023). A DNA phosphorothioation-based Dnd defense system provides
resistance against various phages and is compatible with the Ssp defense system. MBio *14*,
e0093323.

73. Wenger, A.M., Peluso, P., Rowell, W.J., Chang, P.-C., Hall, R.J., Concepcion, G.T., Ebler, J.,
Fungtammasan, A., Kolesnikov, A., Olson, N.D., et al. (2019). Accurate circular consensus
long-read sequencing improves variant detection and assembly of a human genome. Nat.
Biotechnol. *37*, 1155–1162.

74. Mc Cartney, A.M., Shafin, K., Alonge, M., Bzikadze, A.V., Formenti, G., Fungtammasan, A.,
Howe, K., Jain, C., Koren, S., Logsdon, G.A., et al. (2022). Chasing perfection: validation
and polishing strategies for telomere-to-telomere genome assemblies. Nat. Methods *19*,
687–695.

75. Hebert, P.D.N., Braukmann, T.W.A., Prosser, S.W.J., Ratnasingham, S., deWaard, J.R.,
Ivanova, N.V., Janzen, D.H., Hallwachs, W., Naik, S., Sones, J.E., et al. (2018). A Sequel to
Sanger: amplicon sequencing that scales. BMC Genomics *19*, 219.