Complex genetic architecture underlying the plasticity of tobacco leaf width provides insight into across-environment genomic prediction

Li Liu[1#], Yu Han[2#], Wei Liu[2#], Yanjun Zan[1], Dong Wang[3], Yan Ji[1], Yiwen Sun[1], Xin Liu[4], Ran Hao[5], Wenjia Zhang,[1] Linjie Guo[1], Jiarui Zhao[1], Zhimei Song[6], Dan Liu[1], Aiguo Yang[1], Caineng Zhao[7*], Haizhou Hu[1,6*], Lirui Cheng[1*], Huan Si[1*]

[1]Tobacco Research Institute, Chinese Academy of Agricultural Sciences, Qingdao, CN-266000, China
[2]Key Laboratory of Bio-resource and Eco-environment of Ministry of Education, College of Life Sciences, Sichuan University, Chengdu, CN-610065, China
[3]China Tobacco Hunan Industrial Co. Ltd., Changsha, China
[4]China Tobacco Yunnan Industrial Co. Ltd., Kunming, China
[5]College of Forestry, Northwest Agriculture and Forestry University, Yangling, CN-712100, China
[6]Qingdao Zhongyan Seed Co., LTD
[7]China Tobacco Jiangsu Industrial Co. Ltd., Nanjing, China

[#]These authors contributed equally

[*]**Correspondence:**

Huan Si

Email: sihuan@caas.cn

Lirui Cheng

Email: chenglirui@caas.cn

Haizhou Hu

Email: huhaizhou@caas.cn

Caineng Zhao

Email: 119139698@qq.com

30   **Keywords: Tobacco, Leaf width, Phenotypic plasticity, QTL-by-environment**
31   **interaction**

32
33

Abstract

Phenotypic plasticity refers to the ability of a given genotype to produce multiple

phenotypes in response to changing environmental conditions. Understanding the

genetic basis of phenotypic plasticity and developing predictive models for agronomic

traits are crucial for future agricultural adaptation to climate change. In this study, we

investigated the genetic basis of leaf width mean (LWm) and plasticity (LWp) in a

tobacco multiparent advanced generation inter-cross (MAGIC) population which

consisting of 594 individuals. We identified 14 quantitative trait loci (QTLs)

significantly associated with LWm, 43 with LWp. Our findings suggest that dynamic

changes in QTL effects across environments, along with polygenic effects, may

underlie the genetic basis of leaf width plasticity. Among them, *qLW14* was narrowed

down to a 3 Mb structural variation region. When this fragment was deleted in

tobacco, plants exhibited increased leaf width, but only under specific environmental

conditions. This finding suggests that the key gene within *qLW14* may act as a

negative regulator of leaf width through interactions with specific environmental

factors. By integrating genetic diversity, environments variation, and their interactions

into a GEAI model, we were able to build a framework for cross-environment

prediction, improving prediction accuracy by 8.7% when compared to traditional

model. Overall, this study highlights the complex genetic basis underlying LWp,

involving multiple alleles and genotype-environment interactions. These findings

provide valuable insights into the role of phenotypic plasticity in plant adaptation to

environmental changes.

56
57

1. Introduction

Phenotypic plasticity refers to the phenomenon where individuals of the same genotype exhibit different phenotypes under different environmental conditions (Bradshaw, 1965; Pigliucci, 2005), or the ability of organisms to produce different phenotypes in response to varying environmental factors (Kliman, 2016; Nicotra et al., 2010). In organisms, phenotypic plasticity is a trait and possibly an important strategy to quickly adjust phenotypes according to environments, which potentially have effects on evolution (Kusmec et al., 2018; Sommer, 2020). As sessile organisms, plants have evolved unique mechanisms to cope with rapid climate changes, including physiological, metabolic, growth, and developmental adjustments (Brooker et al., 2022; Des Marais et al., 2013). To understand the mechanism of plastic responses is important because agricultural systems will face immense pressure from rapid population growth and climate change in the future.

Previous studies on model organisms, such as yeast, and major crops like rice, maize, and cotton have demonstrated that phenotypic plasticity is widespread and under complex genetic regulation (Alvarez Prado et al., 2014; Kikuchi et al., 2017; Kusmec et al., 2017; Pigliucci, 2005; Zan and Carlborg, 2020). However, fully leveraging phenotypic plasticity requires a comprehensive understanding of its genetic architecture. Three genetic models have been proposed to explain the regulation of phenotypic plasticity, including overdominance model (Gillespie and Turelli, 1989), allelic sensitivity model (Via, 1993) and regulatory gene model (Scheiner and Lyman, 1989; Schneider et al., 2020). The overdominance model suggests that heterozygosity at genes causes plastic responses. However, both the allelic sensitivity model and the regulatory gene model describe genotype-environment interaction (G×E) as the variation in how different genotypes respond to environmental changes. The key distinction lies in their proposed mechanisms: the allelic sensitivity model attributes plastic responses to environmentally sensitive alleles of genes that influence mean phenotypes, whereas the regulatory gene model suggests that plasticity arises from the regulation of mean phenotype genes by other genes that integrate environmental

stimuli.

When comparing the genetic structures underlying phenotypic means and

plasticity, some studies have demonstrated that they are neither entirely unified nor

entirely independent, but rather exhibit partial overlap (Alvarez Prado et al., 2014;

Guo et al., 2020; Jin et al., 2023). For example, a study on maize grain size and

weight plasticity revealed that 30% of the QTLs associated with plasticity were co-

localized with those governing phenotypic means, indicating a shared genetic basis

(Li et al., 2019). However, in another research on 23 maize traits, only 6% of the

QTLs associated with plasticity also co-localized with phenotypic mean loci, which

showed that the candidate genes associated with mean phenotypes and their plasticity

are structurally and functionally distinct across all 23 phenotypes (Kusmec et al.,

2017). To explore the genetic mechanisms underlying phenotypic plasticity have

significant implications for plant breeding.

In recent years, the combination of genome-wide association study (GWAS) and

genome selection (GS) has been widely reported to play a significant role in

accelerating the breeding processes, by integrating the QTLs from GWAS into GS

programs to enhance the prediction accuracy of phenotypes (Desta and Ortiz, 2014;

Huang et al., 2016, 2023; Resende et al., 2012; Spindel et al., 2016; Tong et al., 2020;

Wang et al., 2015; Si et al., 2025). However, the prediction performance of GS model

often varies significantly across different environments, primarily due to the lack of

consideration for G×E. Therefore, uncovering the genetic basis of phenotypic

plasticity and integrating plasticity-associated QTLs into GS models is of great

significance for enhancing the accuracy of phenotypic prediction in heterogeneous

environments.

Tobacco (*Nicotiana tabacum* L.) originated from interspecific hybridization

between *Nicotiana sylvestris* and *Nicotiana tomentosiformis*. As an allotetraploid

species, tobacco exhibits high diversity and plasticity in leaf morphology, plant

architecture, and metabolic traits (Honarnejad and Shoai-Deylami, 2004; Legg and

Collins, 1975; White et al., 1979), making it an important model organism for

research in plant genetics and developmental biology. Over the past two decades,

advancements in high-throughput sequencing technologies and the decreasing cost of sequencing have significantly facilitated research on the genetic mechanisms underlying key traits in plants and animals, including tobacco (Edwards et al., 2017; Khan et al., 2019; Resende et al., 2012; Wang et al., 2022). Leaves are the primary harvestable organ of tobacco and play a crucial role in its growth, development, and agricultural production. Recently, GWAS have been widely applied to investigate the genetic basis of leaf traits in tobacco, leading to the identification of numerous QTLs associated with leaf number, width, and length (Ahmed et al., 2019; Liu et al., 2022b; Safdar et al., 2020; Zuo et al., 2022), which provided valuable resources for revealing the genetic basis of plant growth and development. However, the majority of these genetic studies have focused on mean values of phenotypes, while the genetic basis of plasticity of traits remains largely unexplored.

In this study, we conducted a GWAS in a tobacco MAGIC population consisting of 594 individuals and identified a total of 45 QTLs for LWm, LWp and. Our findings suggest that dynamic changes in major QTL effects across environments, along with polygenic effects, may underlie the genetic basis of leaf width plasticity. Further analysis identified the candidate region of *qLW14* associated with leaf width plasticity Further analysis identified the candidate region of *qLW14* associated with leaf width plasticity and refined it to a ~3 Mb structural variation located at the distal end of chromosome 14.. By integrating genetic diversity, environmental variation, and their interactions, we developed a framework for cross-environment prediction, improving prediction accuracy by 8.7% compared to traditional models. Overall, this study elucidates the complex genetic architecture underlying both leaf width mean and plasticity, involving multiple alleles and G×E. These findings offer valuable insights into the role of phenotypic plasticity in plant adaptation to environmental changes and provide a theoretical foundation for its application in crop breeding.

2. Materials and methods

2.1 Plant materials and growth conditions

A total of eight different tobacco varieties were utilized as parents to develop the

MAGIC population. These included *Beinhart1000-1* and *Florida301*, which is cigar

tobacco varieties; *Basma* and *Samsun*, belonging to oriental tobacco; *Xiaohuaqing*

(XHQ) and *Tangpeng* (TP) from sun-cured tobacco; *Vam*, a variety of Virginia

tobacco; and finally, the flue-cured tobacco variety *Honghuadajinyuan* (HD). Initially,

these eight parents were crossed in pairs to produce four two-way crosses. The

offspring from these crosses then underwent incomplete diallel crossing to generate

four-way crosses, which were further crossed to produce eight-way crosses. These

eight-way crosses were then subjected to random mating, yielding a total of 320 eight-

way crossbreds. From these 320 eight-way crossbreds, three individual plants were

selected per crossbred, and then propagated for five generations through the single

seed descent (SSD) method, resulting in the establishment of over 800 eight-way

tobacco MAGIC inbred lines.

In 2019 and 2020, the MAGIC population, along with the eight parents, was

planted separately in Zhucheng, Shandong Province, China, and Guiyang, Hunan

Province, China, respectively. A randomized complete block design was employed

with two replicates. Each replicate consisted of teo rows of 10 plants per line, with a

row length of 10 m, a row spacing of 1.2 m, and a plant spacing of 0.5 m.

2.2 Phenotype and Genotype

During each experiment, five randomly selected plants were selected for

phenotypic evaluation, and the average value was analyzed. Leaf width was defined

by measuring the width of the blade perpendicular to the main vein for the middle leaf.

In addition to LWm, we assessed LWp using SP and OP (Methods 2.5), which

quantify the variation in LW across different environments.

Genotyping was conducted using a tobacco SNP chip containing 432,362

markers from the Zhengzhou Tobacco Research Institute of the China Tobacco

Corporation. Detailed information regarding genotyping and SNP calling procedures

can be found in a previous study (Yuan et al., 2023).

2.3 Calculation of narrow-sense heritability and genetic correlation

We used a mixed linear model to estimate the narrow-sense heritability：

$$\overline{y} = \mu + Zb + e \quad (\text{Equation 1})$$

Among them, $\overline{y}$ represents LWm, with a total of 594 rows of data. μ is the intercept,

which represents the population mean. Z is the design matrix, satisfying $ZZ^T = G$,

where G is the genetic relationship matrix estimated by GCTA (Yang et al., 2011).

Therefore, b follows a normal distribution $N(0, I\sigma_g^2)$, and e is the residual, which

follows $e \sim N(0, I\sigma_e^2)$. The narrow-sense heritability (h^2) is calculated by the interclass

correlation, with the formula $h^2 = \frac{\sigma_g^2}{\sigma_g^2 + \sigma_e^2}$. All analysis were performed in GCTA.

Furthermore, we used multivariate mixed linear models in ASReml to estimate

pairwise genetic correlations between LWm measured in the four environments.

2.4 Phenotypic Variance Analysis and Estimation of BLUP

To quantify the contribution of each factor to phenotypic variation, we conducted

an analysis of variance (ANOVA) based on a linear model. The model included

genotype (id), environment (loc), and their interaction (id * loc) as explanatory

variables:

$$y_{ij} = u + id_i + loc_j + id_i * loc_j + e_{ij} \quad (\text{Equation 2})$$

y_{ij} ($i = 1 \ldots n$, n=560, number of individuals; $j=1 \ldots m$, m=4, number of sites)

represents the leaf width measured for each individual across four environments; u

represents the population mean; The id_i term denotes the genotype of each individual

and was treated as a categorical variable to capture genetic differences among lines;

The loc_j term represents the environmental factor, and the interaction term $id_i * loc_j$

captures genotype-by-environment interactions (G × E); The e term accounts for

residual error. The proportion of total phenotypic variance explained by each

component was estimated using the sum of squares (SS) obtained from the ANOVA.

We used the following model to estimate the BLUP of leaf width:

$$\overline{y}_{ij} = u + Z\mu + e_{ij} \quad \text{(Equation 3)}$$

where $\overline{y}_{ij}$ represents the leaf width measurements of 594 individuals across four

environments; u represents the population mean; Z is the design matrix linking the

BLUPs estimates with the observed phenotypic values; μ is a vector of random effects

representing the BLUP estimates of 594 individuals, assumed to follow a normal

distribution $N\left(0, I\sigma_g^2\right)$; and e_{ij} represents the residual error.

2.5 Quantification of plasticity measurements

Since all the MAGIC lines were phenotyped for LW across four environments,

we quantified and studied the genetics of LW plasticity in response to environmental

variation. Here, the phenotypic plasticity was classified into two categories. (1)

overall plasticity, which describes plasticity across all the studied environments, and

(2) the second category-specific plasticity is more unique to certain pairs of sites,

which only captures the plasticity across two environments. The motivation

underlying such classification is that some individuals are more robust across most of

the studied sites except only one or a few sites, while other individuals are plastic

among most of the sites. Site-specific plasticity was quantified using pairwise

differences in phenotypic values between two environments. In addition, two

additional approaches were used to quantify the overall plasticity. First, the across

environment variance (VarR) (Vanous et al., 2019) of the rank transformed phenotype

was used to account for the mean difference. The fourth score of the overall plasticity

(FWR) (Finlay and Wilkinson, 1963; Lian and de Los Campos, 2016) applies Finlay-

Wilkinson Regression to partition the phenotype into two components, one that

constant across environments and another that responds dynamically to environmental

changes. Using the linear mixed model, the phenotype of each line is partitioned into

these two components and the plasticity component is used as a measurement of

plasticity. In total, the described approaches resulted in two measurements of

phenotypic plasticity, abbreviated as SP, OP.

2.6 Genome-wide association analysis for LW mean/BLUP and plasticity

measurements

Genome-wide association analysis was conducted using a linear mixed model

implemented in GCTA. We applied a significance threshold of $p < 0.05/Me$ for SNPs,

where Me represents the number of independent SNPs estimated following methods

described by Li et al. (Li et al., 2012). Effect sizes of identified quantitative trait loci

(QTL) were extracted from the GCTA output. The effect sizes for each SNP were

estimated using the previously mentioned mixed linear model.

2.7 Predicting the site-specific performance of the LW

We fitted the following models to predict leaf width phenotypes:

$$Y = X_1\beta_1 + Zu + e \quad \text{(Equation 4)}$$

$$Y = X_2\beta_2 + Zu + e \quad \text{(Equation 5)}$$

Here, Y is a phenotypic value vector of size 560×4, representing the leaf width

measurements of 560 individuals across four environments. The term $X_1\beta_1$ represents

the fixed effects component, where X_1 is the design matrix for fixed effects, including

a column of 1 to represent the overall mean and three additional columns to represent

the environmental effects of the remaining three sites relative to a reference site. The

vector β_1 contains the corresponding fixed effect coefficients, indicating the

contribution of each fixed effect to the phenotypic values. The term Zu represents the

random effects component, where Z is the design matrix satisfying the condition

$ZZ' = G \otimes I$, with G being the genomic relationship matrix (GRM)(Yang et al., 2011)

that captures the genetic similarity among individuals and I being the identity matrix.

The vector u represents the random effects, specifically the breeding values of the

individuals, reflecting their genetic potential contribution to the phenotypic values.

The term e represents the randomly distributed residuals.

We define equation (4) as the GBLUP model, while equation (5) is defined as the

GEAI model. Compared to X_1 in equation (4), X_2 in equation (5) is an extended

design matrix that incorporates QTL genotypes and QTL-by-environment interaction

terms in addition to the columns in X_1. This extension allows the GEAI model to

simultaneously capture systematic effects, genetic effects, and genotype-by-

environment interactions, thereby providing more accurate predictions, particularly in

cross-environment and novel environment predictions. These models were fitted using

the rrBLUP (Endelman, 2011) package in R (https://www.R-project.org/).

We employed two methods to predict leaf width phenotypes. In the first method,

for each of the four environments, 80% of the individuals were randomly selected as

the training set, while the remaining 20% were used as the test set, with their

phenotypic values masked as *NA*. The coefficient of determination (r^2) between the

predicted and measured phenotypes was used to evaluate prediction accuracy within

each environment. In the second method, all individuals from any three of the four

environments were used as the training set, while all individuals from the remaining

environment were used as the test set, with their phenotypic values masked as *NA*.

The coefficient of determination (r^2) between the measured and predicted phenotypes

within each site was used to assess prediction accuracy.

3. Results

3.1 Tobacco leaf width mean variation within each of the four environments

To investigate the genetic basis of leaf width mean (LWm) and plasticity (LWp),

we analyzed 594 tobacco lines derived from a tobacco MAGIC population. Genotypes

were performed using the 430K tobacco SNP array, which includes 432,362 markers,

Leaf width was measured under four different environments in Guiyang (25°27′N, 112°

13′E) and Zhucheng (35°59′N, 119°24′E) during 2019 and 2020 (Figure 1A, Table S1).

The studied environments were abbreviated as follows: Guiyang in 2019 (GY19),

Guiyang in 2020 (GY20), Zhucheng in 2019 (ZC19) and Zhucheng in 2020 (ZC20).

Phenotypic analysis across different environments indicated that tobacco leaf

width exhibited continuous variation, ranging from 13.5 cm to 37.5 cm.

Environmental changes significantly altered the mean across the four environments,

with a widespread reaction norm observed for LWm. For example, when the

environment shifted from ZC20 to GY19, the leaf width in Guiyang (LWm = 27.8 cm ,

CV= 0.14, Fig. 1D) showed a marked increase compared to Zhucheng (LWm = 21.8

cm,CV = 0.15), resulting in variation in LWp (Fig. 1E).

When location was held constant, the leaf width in GY20 averaged 24.1 cm,

representing a 13% reduction compared to GY19. Similarly, in ZC20, the LWm

reached at 21.8 cm, reflecting a 4% decrease relative to ZC19. Within the same year,

the LWm in GY19 was 27.8 cm, 23% greater than that in ZC19, whereas in 2020, the

LWm in GY20 reached at 24.1 cm, 11% higher than in ZC20. Overall, the variation in

LWm across different locations (up to 23%) was substantially greater than the

interannual variation at the same location (4%–13%), indicating that geographic

location had a more pronounced effect on LWm (Fig. 1B, C). Nevertheless, year-to-

year fluctuations were also evident, suggesting that LWm is highly responsive to

environmental changes and exhibits significant phenotypic plasticity.These results

indicate that environmental changes affect both LWm and LWp.

To further dissect the phenotypic variance components, we performed analysis of

variance (ANOVA) considering genotype (G), environment (E), and genotype-by-

environment (G×E) interactions. The results revealed that genotype explained 40.27%

of the total variance in leaf width, while environment contributed 12.02%. Notably,

the G×E interaction accounted for 31.29%, indicating a significant role of

environmental responsiveness in shaping leaf width variation (Fig. S1). These

findings suggest that genetic effects predominantly determine leaf width, but

substantial G×E interactions underscore the importance of plastic responses to

environmental fluctuations.

Fig. 1 Variation in leaf width of MAGIC population across environments A)
Geographical illustration of two filed plantations, Zhucheng and Guiyang, where the
MAGIC population was phenotyped. **B)** Leaf width phenotype distribution histogram
of ZC19 and ZC20. **C)** Leaf width phenotype distribution histogram of GY19 and
GY20. **D)** Distribution histogram of CV of phenotype. **E)** Boxplot of LWm measured
at four environments.

3.2 Environmental perturbation buffers/releases additive variation for LWm by

altering the underlying genetic architecture

Moderate variation (0.13–0.61; Fig. 2A ,Fig.S2) in the estimated narrow-sense

heritability (h^2) across different environmental conditions was observed. Since

phenotypic variation (Vp) varied significantly among the four environments (Fig S3),

the amount of additive genetic variation (Va), estimated as Vp × h^2, ranged from 4.1 to

11.2 (Fig. 2B). These fluctuations in Va were likely driven by changes genetic effects

from one or more QTLs or by the activation and deactivation of a completely new set

of QTLs, indicating that the genetic architecture of LWm was dynamic across the four

environments.

This hypothesis was supported by the variation in pairwise genetic correlations

from the multivariate mixed model analysis (Fig. 2C). The estimated genetic

correlation between environments was as low as 0.4, suggesting considerable changes

in polygenic scores and breeding values across individuals. These changes may be

attributable to genotype-environment interactions, where certain alleles or QTLs exert

different effects in different environments. Collectively, these results illustrate that the

genetic architecture of LWm varied across these environments due to environmental

changes. In the following sections, we will quantify this variation and dissect the

genetic basis underlying LWm variation across environments.

Fig.2. Variation of genetic effects across four environments. (A) Estimated narrow-
sense heritability (h^2) for LWm at each environment. Each dot represents a point
estimate of h^2 indicated on the y-axis and the corresponding environment was labelled
in the x-axis. Standard errors were plotted as error bars. **(B)** Estimated additive
variance at each environment. **(C)** Upper triangular: Pairwise Spearman rank
correlation among LWm measured under four environments. **(C)** Lower triangular:
Pairwise genetic correlation among LWm measured under four environments.

3.3 Polygenic architecture underlying tobacco LWm variation and LWp variation

A widespread reaction norm was observed for LWm. At the population level, the

median LWm in Zhucheng was significantly smaller than that in Guiyang (Fig. 1B),

resulting in variation in LWp. Therefore, we quantified LWp using two types of

metrics. The first type included two measures: Finlay–Wilkinson Regression (FWR),

which evaluates genotype-by-environment interactions, and the across-environment

variance in rank-transformed LWm (VarR), which quantifies the overall plasticity (OP

of LW across all environments. The second metric involved assessing pairwise LWm

differences for all possible pairs of environments, quantifying specific changes across

paired environments, resulting in a total of six specific plasticity (SP) measures.

Almost all plasticity measurements showed a continuous distribution with low to

high narrow-sense heritability (0.015–0.61, Fig. S2, Fig. S4). These results indicate

that LWp is a heritable trait influenced by both genetic and environmental factors,

providing a solid foundation for further dissection of its genetic architectureA total of

45 QTL were identified for LWm and LWp by performing GWAS (Fig. 3A, Table S2,

Fig S5, Fig S6), including 14 QTL for LWm, 9 QTL for lead width BLUP (LW$_{BLUP}$)

and 43 QTL for LWp, LWm and SP share 12 overlapping loci, accounting for 27% of

all identified QTLs. Additionally, LWm and BLUP exhibit complete overlap at 9 loci.

Furthermore, LWm, SP, and BLUP share 8 common loci, representing 18% of all

QTLs. These results indicate a high degree of overlap among the three methods,

indicating a shared genetic basis between LWm, LWp and BLUP (Fig. 3C, Table S3).

The majority of loci show a high degree of overlap across different methods. This

may be due to the variation in the strength of genotype-environment interactions in

the environment, which leads to partial differences in the detected QTLs and

ultimately to the phenotypic plasticity of these traits. By contrasting genetic effects of

QTL across sites, two types of QTL, whose effects changed in magnitude, were

associated with LWp For example, *qLW1-3* showed a significant effect for LW$_{ZC19}$

(5.6 ± 0.7cm; $P = 4.8 \times 10^{-16}$; Fig. 3D), LW$_{ZC20}$ (3.7 ± 0.6cm; $P = 4.4 \times 10^{-9}$) as well as

on the specific plasticity measures LW$_{GY19-ZC20}$ ($P = 5.9 \times 10^{-8}$; Fig. 3D) and LW$_{ZC19-}$

$_{GY19}$ ($P = 2.2 \times 10^{-14}$; Fig. 3D), indicating changes in the magnitude of genetic effects

contributed to the variation of LW plasticity. In contrast, *qLW9-1* (Fig. 3E), was

exclusively detected for several LWp but was not associated with any LWm or

LW$_{BLUP}$ measurements. Altogether, these results suggested that changes in magnitude

of genetic effects across sites caused variation in plasticity. The changing genetic

effects highlighted the role of QTL by environment interaction in the variation of LWm and LWp.

Fig. 3. Summary of the QTLs associated with mean and plasticity measures for LW. (A) Manhattan plots overlaying genome-wide association analysis results for the mean, plasticity measurements and BLUP for LW. The black horizontal dashed line indicated the Bonferroni-corrected genome-wide significance threshold derived as 0.05/Me (Me is the effective number of independent markers; Methods) **(B)** QTL associated with the LWm and LWp measured at four environments and the LW_{BLUP} (y-axis). Each dot represents a SNP, and the size of the dot is proportional to its -Log10 p value, as indicated in the legend on the right. Loci with a p value above the genome-wide significance threshold are colored in blue. **(C)** Venn diagram illustrating the overlap of QTL detected for the 3 types of LW measurements. **(D) and (E)** Genotype-to-phenotype maps for LWm and LW_{SP} at Chr 1: 38450718bp, Chr 9: 55539734bp.

3.4 qLW14 associated with LWm and LWp measurements

We detected a QTL , *qLW14* located at chromosome 14: 116-123 Mb, associated

with LWm at Zhucheng (LW$_{ZC19}$, LW$_{ZC20}$) and multiple LWp measurements (SP-

LW$_{GY19-ZC20}$, SP-LW$_{ZC19-GY19}$, SP-LW$_{ZC19-GY20}$, SP-LW$_{ZC20-GY20}$, Fig. 4A, Fig S5).

Linkage disequilibrium (LD) analysis revealed a high level of LD in this region (Fig.

4B). Within the high level LD block, we identified three major haplotypes (hap1-3).

Phenotypic comparisons of these haplotypes under two environmental conditions

showed that in Zhucheng, lines carrying hap2 exhibited approximately 16% wider

leaves compared to hap1 and hap3 (Fig. 4C), whereas no significant differences were

observed in Guiyang (Fig. 4D). Further analysis revealed that the hap2 haplotype

originated from the parent *Vam* which harbored a ~3 Mb deletion between 121 and

124 Mb, overlapped with the *qLW14* region (Fig. 4E).

To explore the phenotypic effect of this deletion, we compared leaf width

between deletion and non-deletion lines in both environments. Phenotypic analysis

showed that , in Guiyang, lines without the deletion exhibited leaves approximately

18.5% wider than those grown in Zhucheng. In contrast, delete that fragment did not

significantly affect leaf width in either environment (Fig. 4F). Further analysis

revealed that 125 genes were annotated within this region, including 15 tandemly

duplicated genes encoding MADS-box proteins. Members of the MADS-box family

have been reported to regulate plant development in response to light and temperature.

Overall these results highlighted a general contribution from *qLW14* by environment

interaction to variation of LWp .

Fig. 4. GWAS of the LW$_{ZC19\text{-}GY20}$. (A) Manhattan plot for plasticity measurement LW$_{ZC19\text{-}GY20}$. The black horizontal dashed line represents the Bonferroni-corrected genome-wide significance threshold of 0.05/Me (Me is the effective number of independent markers; Methods). **(B)** LD heatmap showing the region surrounding a strong peak (chromosome 14: 116 Mb–124 Mb). One-way ANOVA was used to compare the leaf width of three haplotypes lines *qLW14* in the year of 2019 and 2020 at Zhucheng, Shandong **(C)** and Guiyang, Hunan **(D).(E)** Haplotype analysis from re-sequenced eight parents for MAGIC population (*Vam, Beinhart1000-1, Florida301, Samsun, Basma*, HD: *honghuadajianyuan*, TP: *tangpeng*, XHQ: *xiaohuaqing*) revealed a 3Mb deletion in *Vam*. Difference in leaf width of the deletion and non-deletion*qLW14* in the year of 2019 and 2020 at Zhucheng, Shandong and Guiyang, Hunan **(F).**

3.5 Accounting for dynamics in genetic architecture improved LWm prediction across environment

To improve the effectiveness of cross-environment prediction of leaf width traits, we evaluated the potential of integrating genetic diversity, environmental variation, and their interaction in complex trait prediction by jointly modelling genotype, environment, and their interaction (referred to as the GEAI model, Methods). First, we explored the ability to predict untested lines (Newly developed lines) at the four environments using five-fold cross-validation. Compared to GBLUP, which provides a universal prediction for all sites, our model not only offered site-specific predictions but also improved prediction accuracy. The average prediction accuracy for LW increased by 8.7%. (Fig. 5A, 5B).

In the second scenario, we performed prediction for novel sites without any tested lines by utilizing phenotypic data from three out of the four environments as the training set, while the phenotype data from the remaining environment served as the test set. Compared to the GBLUP model, our GEAI model exhibited an average improvement of 7.2% in prediction accuracy (Fig. 5C, 5D). This substantial enhancement underscores the effectiveness of our GEAI model in capturing and leveraging genotype-environment interactions to enhance prediction accuracy across different environmental contexts. Overall, our study highlighted the potential of integrating QTL by environment interaction in understanding complex traits variation and predictions.

Fig. 5. Prediction of leaf width phenotype using GEAI model. (A) Use 5-fold
cross-validation to predict the phenotypes of the remaining 20% lines in each
environment. **(B)** Use the phenotypes of all strains in three environments in the four
environments as training set for LW, and the remaining environments as a test set. **(C)**
Schematic illustration of the prediction design for **(A)**. **(D)** Schematic illustration of
the prediction design for **(B)**, with LW_{ZC20}, LW_{GY20}, and LW_{GY19} as training set and
LW_{ZC19} as the tested set.

4. Discussion

4.1 Genetic and Environmental Regulation of Leaf Width Plasticity

Consistent with studies in maize and rice (Guo et al., 2020; Ji et al., 2023), in this

study, we found that 33% of the QTLs associated with LWp were shared with those

associated with LWm. This overlap suggests a common genetic basis between LWm

and LWp, indicating that some genetic mechanisms controlling leaf width are

conserved across different environmental conditions. These shared QTLs may

regulate fundamental developmental processes that influence leaf morphology.

However, despite this genetic overlap, the observed phenotypic differences between

LWm and LWp emphasize the important role of phenotypic plasticity in allowing

plants to adapt to varying environmental conditions.

Due to the lack of detailed environmental factor data in our study, we were

unable to definitively determine which specific environmental factors, such as soil

moisture, temperature, light intensity, or nutrient availability contribute to the

plasticity of leaf width. Environmental factors often interact in complex ways to shape

phenotypic outcomes, and their effects may vary depending on the plant's

developmental stage or the intensity and duration of stress. For example, drought

stress may trigger specific signaling pathways that alter leaf development, while

temperature fluctuations could influence cell expansion and division rates, further

modulating leaf width. Additionally, epigenetic modifications, such as DNA

methylation or histone modifications, could mediate the plant's response to

environmental cues, potentially influencing phenotypic plasticity (Guo et al., 2025;

Liu et al., 2022a; Ma et al., 2024; Prasad et al., 2015; Yang et al., 2025). Further

research is needed to explore the specific roles of environmental factors in the

variation of LWm and LWp.

4.2 Potential role of MADS-box transcription factors at the qLW14 locus in leaf

development

Haplotype analysis of the *qLW14* locus revealed the presence of multiple

duplicated MADS-box transcription factors. The MADS-box gene family has been

shown to play a role in leaf development in both *Arabidopsis* and *tomato*. In

Arabidopsis, the MADS-box member FRUITFULL (FUL) regulates leaf morphology

by modulating vascular bundle development (Bemer et al., 2017), while in *tomato*,

SlMBP21 influences leaf development by suppressing genes associated with seed

development (Wang et al., 2021). These findings suggest that MADS-box genes may

have conserved functions in leaf development.

Although the candidate interval of *qLW14* remains relatively large,

advancements in multi-gene editing technologies, such as CRISPR/LbCpf1 (Li et al.,

2020), now make it feasible to perform fine mapping and functional validation of

potential candidate genes within this QTL. Future research could integrate gene

editing, expression analysis, and genetic mapping to narrow down the candidate

region and identify key genes, thereby elucidating the molecular mechanisms through

which MADS-box genes regulate leaf development. These studies will not only

enhance our understanding of the genetic regulatory networks underlying leaf

development but also provide valuable genetic resources for crop breeding.

4.3 Optimization of Genomic Prediction Model Incorporating Genotype-

Environment Interactions

We developed a genomic prediction model that incorporates genotype-

environment interactions. When using data from three environments as the training set

to predict phenotypes in the remaining environment, this model outperformed the

traditional GBLUP. This superiority is mainly due to the model's ability to capture

interaction effects between detectable QTL and environmental conditions. However,

due to the lack of diverse environmental factor data, we were unable to fully elucidate

the specific interaction mechanisms between QTLs and particular environmental

factors, which warrants further investigation.

Despite its limitations, we recommend prioritizing the GEAI model for cross-

environment trait prediction, as it significantly enhances prediction accuracy. By

establishing a prediction model that is responsive to environmental changes, we can

effectively predict phenotypes, such as leaf width, across different environments. This

approach lays the foundation for improving the accuracy of genome-wide predictions

of agronomic traits across multiple environments. Future research could integrate

more environmental factor data (e.g., temperature, humidity, light) to further optimize

model parameters and explore the specific interaction mechanisms between QTLs and

environmental factors. Such efforts will provide essential theoretical support and

technical tools for precise phenotype prediction and genotype selection in crop

breeding.

Funding

This research was funded by the State Key Research & Development Project-Youth
Scientist Program (2023YFD1202400), National Science Foundation of China
(32200503), China National Tobacco Corporation (110202101040), Taishan Young
Scholar Program and Distinguished Overseas Young Talents Program from Shandong
province (2024HWYQ-079), Agricultural Science and Technology Innovation
Program (ASTIP-TRIC01) from Chinese Academy Agriculture Sciences.

Author Contributions

Conceptualization, L.L., W.L., Y.H., R.H.; methodology, Y.H., R.H., S.Y., J.Y., G.J.,
Y.Z., H.S.; formal analysis, L.L., W.L., Y.H., W.D.; investigation, Y.S., L.X., Z.W.,
W.Z., L.C.; data collection and curation, L.C., S.Z., Y.Z., H.S., W.Z., Z.J., Y.S., L.G.;
writing original draft preparation, L.L., W.L.; writing reviewing and editing, L.L.,
W.L., Y.Z., Z.C., H.S.; visualization, L.L., W.L, Y.H; supervision, H.H., Y.Z., H.S.;
funding acquisition, Y.Z., H.S., L.C.

All authors have read and agreed to the published current version of the manuscript.
570

571 **Declaration of competing interest**
572

573 The authors declare that they have no known competing financial interests or personal
574 relationships that could have appeared to influence the work reported in this paper.
575

References

Ahmed, W., Xia, Y., Zhang, H., Li, R., Bai, G., Siddique, K.H., Guo, P., 2019. Identification of
conserved and novel miRNAs responsive to heat stress in flowering Chinese cabbage
using high-throughput sequencing. Sci. Rep. 9, 14922.
Alvarez Prado, S., Sadras, V.O., Borrás, L., 2014. Independent genetic control of maize (Zea mays
L.) kernel weight determination and its phenotypic plasticity. J. Exp. Bot. 65, 4479–4487.
Bemer, M., van Mourik, H., Muiño, J.M., Ferrándiz, C., Kaufmann, K., Angenent, G.C., 2017.
FRUITFULL controls SAUR10 expression and regulates Arabidopsis growth and

architecture. J. Exp. Bot. 68, 3391–3403.

Bradshaw, A.D., 1965. Evolutionary significance of phenotypic plasticity in plants. Adv. Genet. 13,
115–155.

Brooker, R., Brown, L.K., George, T.S., Pakeman, R.J., Palmer, S., Ramsay, L., Schöb, C.,
Schurch, N., Wilkinson, M.J., 2022. Active and adaptive plasticity in a changing climate.
Trends Plant Sci. 27, 717–728.

Des Marais, D.L., Hernandez, K.M., Juenger, T.E., 2013. Genotype-by-Environment Interaction
and Plasticity: Exploring Genomic Responses of Plants to the Abiotic Environment. Annu.
Rev. Ecol. Evol. Syst. 44, 5–29.

Desta, Z.A., Ortiz, R., 2014. Genomic selection: genome-wide prediction in plant improvement.
Trends Plant Sci. 19, 592–601.

Edwards, K.D., Fernandez-Pozo, N., Drake-Stowe, K., Humphry, M., Evans, A.D., Bombarely, A.,
Allen, F., Hurst, R., White, B., Kernodle, S.P., Bromley, J.R., Sanchez-Tamburrino, J.P.,
Lewis, R.S., Mueller, L.A., 2017. A reference genome for Nicotiana tabacum enables
map-based cloning of homeologous loci implicated in nitrogen utilization efficiency.
BMC Genomics 18, 448.

Endelman, J.B., 2011. Ridge Regression and Other Kernels for Genomic Selection with R Package
rrBLUP. Plant Genome 4, 250–255.

Finlay, K.W., Wilkinson, G.N., 1963. The analysis of adaptation in a plant-breeding programme.
Aust. J. Agric. Res. 14, 742–754.

Gillespie, J.H., Turelli, M., 1989. Genotype-environment interactions and the maintenance of
polygenic variation. Genetics 121, 129–138.

Guo, T., Mu, Q., Wang, J., Vanous, A.E., Onogi, A., Iwata, H., Li, X., Yu, J., 2020. Dynamic
effects of interacting genes underlying rice flowering-time phenotypic plasticity and
global adaptation. Genome Res. 30, 673–683.

Guo, Y.-W., Liu, Y., Huang, P.-C., Rong, M., Wei, W., Xu, Y.-H., Wei, J.-H., 2025. Adaptive
Changes and Genetic Mechanisms in Organisms Under Controlled Conditions: A Review.
Int. J. Mol. Sci. 26, 2130.

Honarnejad, R., Shoai-Deylami, M., 2004. Gene effects, combining ability and correlation of
characteristics in F2 populations of burley tobacco cultivars (Nicotiana tabacum L.).
JWSS-Isfahan Univ. Technol. 8, 135–148.

Huang, M., Cabrera, A., Hoffstetter, A., Griffey, C., Van Sanford, D., Costa, J., McKendry, A.,
Chao, S., Sneller, C., 2016. Genomic selection for wheat traits and trait stability. Theor.
Appl. Genet. 129, 1697–1710.

Huang, Y., Qi, Z., Li, J., You, J., Zhang, X., Wang, M., 2023. Genetic interrogation of phenotypic
plasticity informs genome-enabled breeding in cotton. J. Genet. Genomics 50, 971–982.
https://doi.org/10.1016/j.jgg.2023.05.004

Ji, Y., Han, Y., Dai, Y., Hao, F., Feng, X., Chen, Q., Hao, R., Chen, Z., Zhao, W., Zhang, W., 2023.
Polygenic basis of strong and rapid flowering time response to environment perturbations
in wild Arabidopsis thaliana population. bioRxiv 2023–07.

Jin, M., Liu, H., Liu, X., Guo, T., Guo, J., Yin, Y., Ji, Y., Li, Z., Zhang, J., Wang, X., Qiao, F., Xiao,
Y., Zan, Y., Yan, J., 2023. Complex genetic architecture underlying the plasticity of maize
agronomic traits. Plant Commun. 4, 100473.

Khan, S.U., Yangmiao, J., Liu, S., Zhang, K., Khan, M.H.U., Zhai, Y., Olalekan, A., Fan, C., Zhou,

Y., 2019. Genome-wide association studies in the genetic dissection of ovule number,
seed number, and seed weight in Brassica napus L. Ind. Crops Prod. 142, 111877.

Kikuchi, S., Bheemanahalli, R., Jagadish, K.S.V., Kumagai, E., Masuya, Y., Kuroda, E., Raghavan,
C., Dingkuhn, M., Abe, A., Shimono, H., 2017. Genome☐wide association mapping for
phenotypic plasticity in rice. Plant Cell Environ. 40, 1565–1575.

Kliman, R.M., 2016. Encyclopedia of evolutionary biology. Academic Press.

Kusmec, A., de Leon, N., Schnable, P.S., 2018. Harnessing phenotypic plasticity to improve maize
yields. Front. Plant Sci. 9, 1377.

Kusmec, A., Srinivasan, S., Nettleton, D., Schnable, P.S., 2017. Distinct genetic architectures for
phenotype means and plasticities in Zea mays. Nat. Plants 3, 715–723.

Legg, P.D., Collins, G.B., 1975. Genetic parameters in aKy 14 xKy Ex 42 burley population
ofNicotiana tabacum L. Theor. Appl. Genet. 45, 264–267.

Li, B., Liang, S., Alariqi, M., Wang, F., Wang, G., Wang, Q., Xu, Z., Yu, L., Zafar, M.N., Sun, L.,
2020. The application of temperature sensitivity CRISPR/LbCpf1 (LbCas12a) mediated
genome editing in allotetraploid cotton (G. hirsutum) and creation of nontransgenic,
gossypol-free cotton. Plant Biotechnol. J. 19, 221.

Li, C., Wu, X., Li, Yongxiang, Shi, Y., Song, Y., Zhang, D., Li, Yu, Wang, T., 2019. Genetic
architecture of phenotypic means and plasticities of kernel size and weight in maize.
Theor. Appl. Genet. 132, 3309–3320.

Li, M.-X., Yeung, J.M.Y., Cherny, S.S., Sham, P.C., 2012. Evaluating the effective numbers of
independent tests and significant p-value thresholds in commercial genotyping arrays and
public imputation reference datasets. Hum. Genet. 131, 747–756.

Lian, L., de Los Campos, G., 2016. FW: an R package for Finlay–Wilkinson regression that
incorporates genomic/pedigree information and covariance structures between
environments. G3 Genes Genomes Genet. 6, 589–597.

Liu, Y., Wang, J., Liu, B., Xu, Z., 2022a. Dynamic regulation of DNA methylation and histone
modifications in response to abiotic stresses in plants. J. Integr. Plant Biol. 64, 2252–2274.

Liu, Y., Yuan, G., Si, H., Sun, Y., Jiang, Z., Liu, D., Jiang, C., Pan, X., Yang, J., Luo, Z., 2022b.
Identification of QTLs associated with agronomic traits in tobacco via a biparental
population and an eight-way MAGIC population. Front. Plant Sci. 13, 878267.

Ma, L., Xing, L., Li, Z., Jiang, D., 2024. Epigenetic control of plant abiotic stress responses. J.
Genet. Genomics.

Nicotra, A.B., Atkin, O.K., Bonser, S.P., Davidson, A.M., Finnegan, E.J., Mathesius, U., Poot, P.,
Purugganan, M.D., Richards, C.L., Valladares, F., 2010. Plant phenotypic plasticity in a
changing climate. Trends Plant Sci. 15, 684–692.

Pigliucci, M., 2005. Evolution of phenotypic plasticity: where are we going now? Trends Ecol.
Evol. 20, 481–486.

Prasad, P.V.V., Staggenborg, S.A., Ristic, Z., 2015. Impacts of Drought and/or Heat Stress on
Physiological, Developmental, Growth, and Yield Processes of Crop Plants, in: Ahuja,
L.R., Reddy, V.R., Saseendran, S.A., Yu, Q. (Eds.), Advances in Agricultural Systems
Modeling. American Society of Agronomy and Soil Science Society of America, Madison,
WI, USA, pp. 301–355.

Resende, M.F.R., Muñoz, P., Acosta, J.J., Peter, G.F., Davis, J.M., Grattapaglia, D., Resende,
M.D.V., Kirst, M., 2012. Accelerating the domestication of trees using genomic selection:

accuracy of prediction models across ages and environments. New Phytol. 193, 617–624.

Safdar, L.B., Almas, F., Sarfraz, S., Ejaz, M., Ali, Z., Mahmood, Z., Yang, L., Tehseen, M.M., Ikram, M., Liu, S., Quraishi, U.M., 2020. Genome‐wide association study identifies five new cadmium uptake loci in wheat. Plant Genome 13, e20030.

Scheiner, S.M., Lyman, R.F., 1989. The genetics of phenotypic plasticity I. Heritability. J. Evol. Biol. 2, 95–107.

Schneider, H.M., Klein, S.P., Hanlon, M.T., Kaeppler, S., Brown, K.M., Lynch, J.P., 2020. Genetic control of root anatomical plasticity in maize. Plant Genome 13, e20003.

Si, H., Wang, D., Zan, Y., Liu, W., Pu, W., Li, X., Mao, H., Yang, X., Song, S., Wang, Y., 2025. Development of two sets of tobacco chromosome segment substitution lines and QTL mapping for agronomic and disease resistance traits. Ind. Crops Prod., 226, 120622,

Sommer, R.J., 2020. Phenotypic plasticity: from theory and genetics to current and future challenges. Genetics 215, 1–13.

Spindel, J.E., Begum, H., Akdemir, D., Collard, B., Redoña, E., Jannink, J.L., McCouch, S., 2016. Genome-wide prediction models that incorporate de novo GWAS are a powerful new tool for tropical rice improvement. Heredity 116, 395–408.

Tong, H., Küken, A., Nikoloski, Z., 2020. Integrating molecular markers into metabolic models improves genomic selection for Arabidopsis growth. Nat. Commun. 11, 2410.

Vanous, A., Gardner, C., Blanco, M., Martin‐Schwarze, A., Wang, J., Li, X., Lipka, A.E., Flint‐Garcia, S., Bohn, M., Edwards, J., Lübberstedt, T., 2019. Stability Analysis of Kernel Quality Traits in Exotic‐Derived Doubled Haploid Maize Lines. Plant Genome 12, 170114.

Via, S., 1993. Adaptive Phenotypic Plasticity: Target or By-Product of Selection in a Variable Environment? Am. Nat. 142, 352–365.

Wang, C., Ma, Q., Xie, X., Zhang, X., Yang, D., Su, J., Ma, X., Lin, H., 2022. Identification of favorable haplotypes/alleles and candidate genes for three plant architecture-related traits via a restricted two-stage multilocus genome-wide association study in upland cotton. Ind. Crops Prod. 177, 114458.

Wang, Y., Mette, M.F., Miedaner, T., Wilde, P., Reif, J.C., Zhao, Y., 2015. First insights into the genotype–phenotype map of phenotypic stability in rye. J. Exp. Bot. 66, 3275–3284.

Wang, Y.-S., Guo, P.-Y., Zhang, J.-L., Xie, Q.-L., Hui, S., Hu, Z.-L., Chen, G.-P., 2021. Overexpression of the MADS-box gene SIMBP21 alters leaf morphology and affects reproductive development in tomato. J. Integr. Agric. 20, 3170–3185.

White, F.H., Pandeya, R.S., Dirks, V.A., 1979. Correlation studies among and between agronomic, chemical, physical and smoke characteristics in flue-cured tobacco (*Nicotiana tabacum L.*). Can. J. Plant Sci. 59, 111–120.

Yang, J., Lee, S.H., Goddard, M.E., Visscher, P.M., 2011. GCTA: a tool for genome-wide complex trait analysis. Am. J. Hum. Genet. 88, 76–82.

Yang, W., Feng, M., Yu, K., Cao, J., Cui, G., Zhang, Y., Peng, H., Yao, Y., Hu, Z., Ni, Z., 2025. The TaCLE24b peptide signaling cascade modulates lateral root development and drought tolerance in wheat. Nat. Commun. 16, 1952.

Yuan, G., Sun, K., Yu, W., Jiang, Z., Jiang, C., Liu, D., Wen, L., Si, H., Wu, F., Meng, H., Cheng, L., Yang, A., Wang, Y., 2023. Development of a MAGIC population and high-resolution quantitative trait mapping for nicotine content in tobacco. Front. Plant Sci. 13.

716   Zan, Y., Carlborg, Ö., 2020. Dynamic genetic architecture of yeast response to environmental
717   perturbation shed light on origin of cryptic genetic variation. PLoS Genet. 16, e1008801.
718   Zuo, J.-F., Ikram, M., Liu, J.-Y., Han, C.-Y., Niu, Y., Dunwell, J.M., Zhang, Y.-M., 2022.
719   Domestication and improvement genes reveal the differences of seed size-and oil-related
720   traits in soybean domestication and improvement. Comput. Struct. Biotechnol. J. 20,
721   2951–2964.
722

Fig. 1 Variation in leaf width of MAGIC population across environments A) Geographical illustration of two filed plantations, Zhucheng and Guiyang, where the MAGIC population was phenotyped. **B)** Leaf width phenotype distribution histogram of ZC19 and ZC20. **C)** Leaf width phenotype distribution histogram of GY19 and GY20. **D)** Distribution histogram of CV of phenotype. **E)** Boxplot of LWm measured at four environments.

Fig.2. Variation of genetic effects across four environments. (A) Estimated narrow-sense heritability (h^2) for LWm at each environment. Each dot represents a point estimate of h^2 indicated on the y-axis and the corresponding environment was labelled in the x-axis. Standard errors were plotted as error bars. **(B)** Estimated additive variance at each environment. **(C)** Upper triangular: Pairwise Spearman rank correlation among LWm measured under four environments. **(C)** Lower triangular: Pairwise genetic correlation among LWm measured under four environments.

Fig. 3. Summary of the QTLs associated with mean and plasticity measures for LW. (A) Manhattan plots overlaying genome-wide association analysis results for the mean, plasticity measurements and BLUP for LW. The black horizontal dashed line indicated the Bonferroni-corrected genome-wide significance threshold derived as 0.05/Me (Me is the effective number of independent markers; Methods) **(B)** QTL associated with the LWm and LWp measured at four environments and the LW_{BLUP} (y-axis). Each dot represents a SNP, and the size of the dot is proportional to its -Log10 p value, as indicated in the legend on the right. Loci with a p value above the genome-wide significance threshold are colored in blue. **(C)** Venn diagram illustrating the overlap of QTL detected for the 3 types of LW measurements. **(D) and (E)** Genotype-to-phenotype maps for LWm and LW_{SP} at Chr 1: 38450718bp, Chr 9: 55539734bp.

Fig. 4. GWAS of the LW$_{ZC19-GY20}$. (A) Manhattan plot for plasticity measurement LW$_{ZC19-GY20}$. The black horizontal dashed line represents the Bonferroni-corrected genome-wide significance threshold of 0.05/Me (Me is the effective number of independent markers; Methods). **(B)** LD heatmap showing the region surrounding a strong peak (chromosome 14: 116 Mb–124 Mb). One-way ANOVA was used to compare the leaf width of three haplotypes lines *qLW14* in the year of 2019 and 2020 at Zhucheng, Shandong **(C)** and Guiyang, Hunan **(D).(E)** Haplotype analysis from re-sequenced eight parents for MAGIC population (*Vam*, *Beinhart1000-1*, *Florida301*, *Samsun*, *Basma*, HD: *honghuadajianyuan*, TP: *tangpeng*, XHQ: *xiaohuaqing*) revealed a 3Mb deletion in *Vam*. Difference in leaf width of the deletion and non-deletion*qLW14* in the year of 2019 and 2020 at Zhucheng, Shandong   and

Guiyang, Hunan **(F).**

Fig. 5. Prediction of leaf width phenotype using GEAI model. (A) Use 5-fold cross-validation to predict the phenotypes of the remaining 20% lines in each environment. **(B)** Use the phenotypes of all strains in three environments in the four environments as training set for LW, and the remaining environments as a test set. **(C)** Schematic illustration of the prediction design for **(A)**. **(D)** Schematic illustration of the prediction design for **(B)**, with LW_{ZC20}, LW_{GY20}, and LW_{GY19} as training set and LW_{ZC19} as the tested set.