Growth regulation bringing modularity to morphogenesis of complex three-dimensional exoskeletons

Hiroshi C. Ito[a] and Yu Uchiumi[b]

[a] Research Center for Integrative Evolutionary Science, The Graduate University for Advanced Studies, SOKENDAI, Hayama, Kanagawa 240-0193, Japan

[b] Department of Liberal Arts, Nihon University School of Medicine, Itabashi-ku, Tokyo 173–8610, Japan

Abstract

Diverse three-dimensional morphologies of arthropods' appendages, including beetle horns, are formed through the non-uniform growth of epidermis. Prior to moulting, an epidermal tissue peels off from the old cuticle and grows non-uniformly to shape protruding appendages, which are often branching, curving, or twisting, from the planar epidermis. Such non-uniform growth is possibly regulated by the distribution of morphogens on the epidermal cell sheet. Previous studies have identified molecules and signalling pathways related to such morphogenesis, yet leaving unclear how morphogen intensity regulates the cell growth rate locally on the epidermis and how the distribution corresponds to the shapes of three-dimensional appendages. In the present study, to reveal the relationship between the epidermal growth regulation and generated structures, we theoretically examined under a deductive growth model how various shapes can be generated from a planar epidermis via non-uniform growth regulated by the morphogen distributions. Our result shows that the heterochronic expression of some morphogens can fuse multiple simple shapes to generate various structures emulating complex appendages of beetles. This result implies that morphogenesis through such a mechanism may have developmental stability and modularity, providing insights into the evolution of the diverse morphology of arthropods.

Introduction

Arthropods have diverse three-dimensional morphologies. In particular, with beetle horns being a typical example, the morphology of their appendages varies among species, sexes, developmental stages, and environmental conditions [1]. Although horns often have extreme sizes and elaborate shapes, they can evolve rapidly and dramatically and are easy to gain and lose evolutionarily [2]. Moreover, those horns should have mechanical strength as a weapon of competition in males and developmental plasticity depending on the whole body size [3,4], implying that the morphogenesis has ambivalence between evolvability and developmental stability. Therefore, such appendages of arthropods can be regarded as a representative example of the classical paradox in evolutionary developmental biology [5].

The morphogenesis of exoskeletal appendages, such as beetle horns, depends on the non-uniform growth of epidermal tissues during metamorphosis or moulting [6,7]. Prior to moulting, an epidermal cell sheet lining the old cuticle detaches from the cuticle and grows non-uniformly to transform from the planar shape constituting the larval epidermis into the shape of an adult appendage, producing a new cuticle [8]. Because the synthesised cuticle does not have strong bending elasticity, the growing cell sheet and the produced cuticle are compacted inside the old cuticle in a wrinkled and folded state [9]. They extend to form an elastic or rigid three-dimensional shape during moulting, along with breaking out of the old cuticle [9]. The non-uniform growth of epithelial sheets involves non-uniform cell division, rearrangement, and growth, which are integratively controlled to achieve harmonious three-dimensional shapes [8]. Recent studies have identified genes and pathways involved in the integrative control to form the appendages [10–13]. For example, in the case of beetle horns [1], the genes related to leg patterning, e.g., *Distal-less* and *dachshund,* also contribute to the patterning of their proximo-distal axis; the Hox gene *Sex-combs reduced* affects their positioning; and several pathways, including insulin signalling, regulate their relative size. *Notch* and *CyclinE* affect the depth and pattern, respectively, of the primordial furrow of the epidermal tissue inside the prepupal cuticle, which is related to the proliferation of epidermal cells and determines the shape of the adult horn [14].

However, understanding how such non-uniform growth of epidermal sheets is controlled harmoniously to generate three-dimensional structures, including horns and other appendages, is still challenging. Previous studies have shown how local cell mechanics allow the construction of three-dimensional structures, e.g., the respiratory tubes of *Drosophila* eggs [15] and folding patterns similar to an insect primordium [16]. Another study revealed that the folding patterns on the primordium can expand into its horn structure [17]. Because these studies assume in advance certain distributions of mechanical forces or folding patterns, the processes to form

such distributions or patterns via non-uniform growth have not been clarified. For revealing morphogenesis of the appendages via non-uniform growth, Ito [18] proposed a model, called here the "MGS (morphogen-gradient-squared) growth model," where the direction of sheet growth is parallel to the morphogen intensity gradient at that position and its growth rate is determined by the gradient strength. This growth model can transform a flat epidermal sheet into a three-dimensional structure that materializes the shape of a morphogen distribution, if the shape is sufficiently simple such as a straight and unbranched horn or spine (Figure 1). In addition, the generated shape can be further curved and twisted by incorporating the expression of additional morphogens (Figure 2h, 2i). However, the exoskeletal appendages as beetle horns have more complex shapes involving branches, and the morphogenesis of such appendage remains unclear, particularly in the relationship between the growth regulation and the resulting shape of an extended sheet.

In the present study, we investigated how complex three-dimensional structures emulating beetle horns or mandibles can be generated through the non-uniform growth of epidermal sheets regulated by morphogen distributions. To facilitate the analysis, we adopted the MGS growth model proposed by Ito [18], because as above, this model provides simple correspondence of a morphogen distribution to the shape of an extended sheet. For efficient analysis on the relationship between the growth regulation and the extended sheet shape, our analysis did not incorporate the epidermal folding and remodelling process in the pupation of a beetles [19]. Our results show that heterochronic expression of several morphogens on the epidermal sheet can fuse different simple shapes to generate various complex shapes. The remainder of this paper is organised as follows. Section 2 explains how the nonuniform sheet growth in the MGS growth model is described geometrically, and how the three-dimensional shape of the extended sheet is analysed. Section 3.1 shows how multiple simple shapes can be fused in the extended sheet and how the overlap of expression periods among morphogens affects the shape of the extended sheet. Section 3.2 shows that the fusion of simple shapes can flexibly generate various three-dimensional shapes. Finally, section 4 discusses our results in relation to previous studies.

METHODS

(a) Description of sheet growth

In the MGS growth model, the epidermal sheet grows only in the direction tangential to the sheet surface (corresponding to the surface area increase) at each position; growth in the normal direction (corresponding to the thickness increase) is not considered. Therefore, for notational

simplicity, the sheet is assumed to be infinitesimally thin (although the sheet thickness was

explicitly considered in the finite element method analysis for the shape of the extended sheet).

Initially, the sheet is assumed to lie on the x-y plane ($z = 0$) in the three-dimensional spatial

coordinate system $\mathbf{r} = (x, y, z)^{\mathrm{T}}$, where the superscript T denotes the transposition of a matrix or

vector (Figure 1a). On this sheet, a two-dimensional coordinate system, $\mathbf{u} = (X, Y)^{\mathrm{T}}$, is set,

providing the material coordinates of the sheet, called the Lagrangian coordinate system in

continuum mechanics (Figure 1b). The material coordinate system stretches or contracts in the

same manner as the sheet stretches or contracts. Hence, the sheet maintains the same shape in the

material coordinate system as in its initial state, regardless of how it has deformed in the spatial

coordinate system (Figures 1b, 1d, and 1f). Accordingly, the position of each microelement in the

sheet is invariant in the material coordinate system.

For a microelement of the sheet at an arbitrary position $\mathbf{u}$ in the material coordinate system,

its spatial coordinates at time t are denoted as $\tilde{\mathbf{r}}(\mathbf{u}, t) = (\tilde{x}(\mathbf{u}, t), \tilde{y}(\mathbf{u}, t), \tilde{z}(\mathbf{u}, t))^{\mathrm{T}}$. In the initial

state, the (x, y) components of the spatial coordinates coincide with the material coordinates,

satisfying the conditions $\tilde{x}(\mathbf{u}, 0) = X$, $\tilde{y}(\mathbf{u}, 0) = Y$, and $\tilde{z}(\mathbf{u}, 0) = 0$.

For describing the non-uniform growth of the sheet, a short line segment from $\mathbf{u} = (X, Y)^{\mathrm{T}}$ to

$\mathbf{u} + \boldsymbol{\delta}\mathbf{u}$ with $\boldsymbol{\delta}\mathbf{u} = (\delta X, \delta Y)^{\mathrm{T}}$ is considered in the material coordinates. Then, its "unstrained

length" at time t, denoted by $l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, t)$, is defined as its length in the spatial coordinates,

provided that the microelement including the line segment undergoes no stress. On this basis, the

ratio of the unstrained length at time t to its initial length $|\boldsymbol{\delta}\mathbf{u}|$ describes the growth amount of

the sheet at $\mathbf{u}$. Specifically, the growth amount $L(\mathbf{u}, \mathbf{e}, t)$ at $\mathbf{u}$ along direction $\mathbf{e} = \boldsymbol{\delta}\mathbf{u} \, / \, |\boldsymbol{\delta}\mathbf{u}|$ at

time t is given by

115
$$L(\mathbf{u}, \mathbf{e}, t) = \lim_{|\boldsymbol{\delta}\mathbf{u}| \to 0} \frac{l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, t)}{l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, 0)} = \sqrt{\mathbf{e}^{\mathrm{T}} \mathbf{M}(\mathbf{u}, t) \mathbf{e}} \tag{1}$$

116 (see Appendix A.1 for the transformation), where $\mathbf{M}(\mathbf{u}, t)$ is a 2-by-2 symmetric matrix given by

117
$$\mathbf{M}(\mathbf{u}, t) = \begin{pmatrix} M_{XX}(\mathbf{u}, t) & M_{XY}(\mathbf{u}, t) \\ M_{XY}(\mathbf{u}, t) & M_{YY}(\mathbf{u}, t) \end{pmatrix} = \begin{pmatrix} |\partial_X \tilde{\mathbf{r}}|^2 & [\partial_X \tilde{\mathbf{r}}]^{\mathrm{T}} [\partial_Y \tilde{\mathbf{r}}] \\ [\partial_X \tilde{\mathbf{r}}]^{\mathrm{T}} [\partial_Y \tilde{\mathbf{r}}] & |\partial_Y \tilde{\mathbf{r}}|^2 \end{pmatrix} \tag{2}$$

118 with $\partial_X \tilde{\mathbf{r}} = \frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial X}$ and $\partial_Y \tilde{\mathbf{r}} = \frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial Y}$. Here $\mathbf{M}(\mathbf{u}, t)$ is referred to as the growth metric, containing

119 all the information on the amount of growth in all directions. For example, $\mathbf{e} = (1,0)^{\mathrm{T}}$ gives

120 $L(\mathbf{u}, \mathbf{e}, t) = M_{XX}(\mathbf{u}, t)$, whereas $\mathbf{e} = (0,1)^{\mathrm{T}}$ gives $L(\mathbf{u}, \mathbf{e}, t) = M_{YY}(\mathbf{u}, t)$. In the initial state, the

121 growth amount is assumed to be equal to 1 in any direction, satisfying $L(\mathbf{u}, \mathbf{e}, 0) = 1$ for any $\mathbf{e}$,

122 meaning that $\mathbf{M}(\mathbf{u}, 0) = \mathbf{I}$ holds with $\mathbf{I}$ describing the identity matrix.

(b) Sheet growth in MGS growth model

Mathematical description for non-uniform sheet growth often results in formulas that are not easy for analytical treatment. In contrast, as shown below, the MGS growth model describes the non-uniform sheet growth in a considerably simple form, allowing us to obtain deep insights.

In the MGS growth model, the sheet at each position in the spatial coordinates grows in parallel with the intensity gradient of a morphogen along the sheet at that position, and its growth rate is proportional to the square of the gradient strength. The morphogen intensity at each position $\mathbf{u}$ at time t, denoted by $\beta(\mathbf{u}, t)$, is assumed to be unaffected by the sheet growth. Sheet growth is assumed to be unstrained in the sense that the actual growth of each microelement of the sheet in spatial coordinates follows the growth amount given by equation (1). Note that the morphogen intensity gradient along the sheet in the spatial coordinates can still be weaken by the sheet growth. Under these assumptions, the time change of the growth metric is independent of the sheet shape in spatial coordinates and is derived in Appendix A as

$$\frac{\partial \mathbf{M}(\mathbf{u}, t)}{\partial t} = \nabla \beta(\mathbf{u}, t) \nabla \beta(\mathbf{u}, t)^{\mathrm{T}}, \tag{3}$$

where $\nabla \beta(\mathbf{u}, t) = \left(\frac{\partial \beta(\mathbf{u},t)}{\partial X}, \frac{\partial \beta(\mathbf{u},t)}{\partial Y}\right)^{\mathrm{T}}$ describes the gradient of $\beta(\mathbf{u}, t)$ in the material coordinates evaluated at $\mathbf{u}$, and where the $\nabla \beta(\mathbf{u}, t) \nabla \beta(\mathbf{u}, t)^{\mathrm{T}}$ is a 2-by-2 symmetric matrix given by

$$\nabla \beta(\mathbf{u}, t) \nabla \beta(\mathbf{u}, t)^{\mathrm{T}} = \begin{pmatrix} \partial_X \beta \\ \partial_Y \beta \end{pmatrix} \begin{pmatrix} \partial_X \beta & \partial_Y \beta \end{pmatrix} = \begin{pmatrix} [\partial_X \beta]^2 & [\partial_X \beta][\partial_Y \beta] \\ [\partial_X \beta][\partial_Y \beta] & [\partial_Y \beta]^2 \end{pmatrix} \tag{4}$$

with $\partial_X \beta = \frac{\partial \beta(\mathbf{u},t)}{\partial X}$ and $\partial_Y \beta = \frac{\partial \beta(\mathbf{u},t)}{\partial Y}$. Because $\mathbf{e}$ does not depend on time, we see from equations (1) and (3) that

$$\frac{\partial L(\mathbf{u}, \mathbf{e}, t)^2}{\partial t} = [\mathbf{e}^{\mathrm{T}} \nabla \beta(\mathbf{u}, t)]^2. \tag{5}$$

Hence, $\mathbf{e}$ being parallel to $\nabla \beta(\mathbf{u}, t)$ gives $\partial L(\mathbf{u}, \mathbf{e}, t)/\partial t = |\nabla \beta(\mathbf{u}, t)|^2 / L(\mathbf{u}, \mathbf{e}, t)$, while $\mathbf{e}$ being orthogonal to $\nabla \beta(\mathbf{u}, t)$ gives $\partial L(\mathbf{u}, \mathbf{e}, t)/\partial t = 0$.

When the morphogen distribution can be expressed as $\beta(\mathbf{u}, t) = \rho(t)\beta(\mathbf{u})$ with $\rho(t)$ describing the temporal expression pattern (satisfying $\rho(t) = 0$ for $t \geq \tau$) for a static morphogen distribution $\beta(\mathbf{u})$, the time integration of equation (3) gives the growth metric at time $t = \tau$:

$$\mathbf{M}(\mathbf{u}, \tau) = \mathbf{M}(\mathbf{u}, 0) + \int_0^{\tau} \frac{d\mathbf{M}(\mathbf{u}, t)}{dt} dt = \mathbf{I} + \hat{\rho}^2 \nabla \beta(\mathbf{u}) \nabla \beta(\mathbf{u})^{\mathrm{T}} \tag{6}$$

with $\hat{\rho}^2 = \int_0^{\tau} \rho(t)^2 \, dt$. As derived in Appendix B, this growth metric, $\mathbf{M}(\mathbf{u}, \tau)$, is identical to the growth metric $\mathbf{M}_\beta(\mathbf{u}, \tau)$ attained when the sheet shape is given by

$$\tilde{\mathbf{r}}_\beta(\mathbf{u}) = \begin{pmatrix} \tilde{x}_\beta(\mathbf{u}) \\ \tilde{y}_\beta(\mathbf{u}) \\ \tilde{z}_\beta(\mathbf{u}) \end{pmatrix} = \begin{pmatrix} X \\ Y \\ \hat{\rho}\beta(\mathbf{u}) \end{pmatrix}. \tag{7}$$

In other words, for each microelement of the sheet at its initial state $\left(\tilde{x}(\mathbf{u}), \tilde{y}(\mathbf{u}), \tilde{z}(\mathbf{u})\right) = (X, Y, 0)$, only the z-coordinate is raised from 0 to $\hat{\rho}\beta(\mathbf{u})$ (Appendix B). Therefore, the grown sheet with its growth metric $\mathbf{M}(\mathbf{u}, \tau)$ can be extended into a smooth shape $\tilde{\mathbf{r}}_\beta(\mathbf{u})$ keeping the sheet unstrained (provided that the sheet is infinitesimally thin), where $\tilde{\mathbf{r}}_\beta(\mathbf{u})$ is called an isometric embedding for metric $\mathbf{M}(\mathbf{u}, \tau)$ in differential geometry. Hence, the MGS growth can generate a three-dimensional structure from a flat epithelial sheet corresponding to the shape of the morphogen distribution.

For convenience, $\beta(\mathbf{u})$ is rescaled so that $\hat{\rho} = 1$ holds without loss of generality. We then refer to the three-dimensional shape $\tilde{\mathbf{r}}_\beta(\mathbf{u})$ and its corresponding growth metric $\mathbf{M}_\beta(\mathbf{u}, \tau)$ as the "basic shape" and the "basic growth metric," respectively, induced from the morphogen distribution $\beta(\mathbf{u})$.

(c) Analysis of the three-dimensional shape of the extended sheet

To investigate the extended shape of the grown sheet, we numerically analysed the sheet deformation in spatial coordinates by means of the finite element method with the assumption of neo-Hookean elasticity (using the modular finite element method [MFEM] library in C++ language [20]; see Appendix C.1 for details). The initial state of the sheet is a thin square object placed on the x-y plane (side length = 8.0, along the x- and y-axes, and thickness = 0.2 along the z-axis). Differently from the sheet growth process assuming unstrained growth, the extended sheet may have some strain so that the "extension metric" (Figure 1f) calculated from the shape of the extended sheet (Appendix C.1) may deviate from the growth metric (Figure 1d) given by equation (6).

In actual moulting in arthropods, the epidermal sheet detached from the old cuticle grows and produces a new cuticle with weak bending elasticity, forming wrinkled primordia [16]. After removing the old cuticle, these primordia are then unfolded by water pressure into a smooth shape with stronger elasticity [21]. However, such a wrinkled state causes difficulty in calculating the sheet deformation. Therefore, to avoid the wrinkled state in our analysis, we assumed that the sheet always had sufficiently strong elasticity to allow its steady extension into a smooth, three-dimensional shape. For the same purpose, we chose the initial shape as the basic shape of the morphogen distribution or a linear combination of morphogen distributions with some height shrinking (defined by $Z_{\text{init}}(X, Y)$ in Appendix C.1).

When the morphogen distribution $\beta(\mathbf{u})$ is defined as a normal distribution (Figure 2a), the
grown sheet (Figure 2d) is extended to have a three-dimensional shape (Figure 2g) similar to its
corresponding basic shape (Figure 2b). The extended sheet can then be curved (Figure 2h) or
twisted (Figure 2i) by assuming additional morphogens that modify the degree and direction of
local sheet growth (see Appendix D for details).

We also analysed the shapes of the extended sheets that mimic the head and thorax of beetles
with horns (see Appendix C.2 for details). Although the finite element method can calculate sheet
deformation under realistic physical assumptions, the calculation may not be stable when the sheet
is considerably thin as in the case of insect exoskeletons. Hence, in this analysis, we used the one-
layer spring-vertex method, which can be more stable for thin sheets as it does not require an
explicit description of the sheet thickness. In this analysis, water pressure was assumed for the
sheet extension, as in actual moulting [17].

Results

195   In this section, we first show that the MGS growth model can fuse multiple basic shapes through
196   the heterochronic expression of multiple morphogen distributions on the sheet. We then show that
197   various three-dimensional shapes of extended sheets can be generated by fusing basic shapes.

(a) Fusion of basic shapes

199   We can fuse multiple basic shapes by expressing multiple morphogen distributions on the same
200   sheet. Specifically, we can fuse the basic shapes of two static morphogen distributions, denoted
201   by $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$, by defining $\beta(\mathbf{u}, t)$ as

$$\beta(\mathbf{u}, t) = \rho_1(t)\beta_1(\mathbf{u}) + \rho_2(t)\beta_2(\mathbf{u}) \tag{8}$$

203   with $\rho_1(t)$ and $\rho_2(t)$ describing the temporal expression patterns for $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$,
204   respectively, and where both expressions end by time $t = \tau$ (i.e., $\rho_1(t) = 0$ and $\rho_2(t) = 0$ for
205   $t \geq \tau$) and satisfy $\int_0^\tau \rho_1(t)^2 dt = \int_0^\tau \rho_2(t)^2 dt = 1$. The final growth metric at $t = \tau$ is then given
206   by

$$\mathbf{M}(\mathbf{u}, \tau) = \mathbf{M}(\mathbf{u}, 0) + \int_0^\tau \nabla\beta(\mathbf{u}, t)\nabla\beta(\mathbf{u}, t)^{\mathrm{T}} dt$$
$$= \mathbf{I} + \mathbf{M}_{11}(\mathbf{u}) + \mathbf{M}_{22}(\mathbf{u}) + \psi_{12}\mathbf{M}_{12}(\mathbf{u}) \tag{9}$$

208   with

$$\mathbf{M}_{11}(\mathbf{u}) = \nabla\beta_1(\mathbf{u})\nabla\beta_1(\mathbf{u})^{\mathrm{T}},$$
$$\mathbf{M}_{22}(\mathbf{u}) = \nabla\beta_2(\mathbf{u})\nabla\beta_2(\mathbf{u})^{\mathrm{T}},$$
$$\mathbf{M}_{12}(\mathbf{u}) = \nabla\beta_1(\mathbf{u})\nabla\beta_2(\mathbf{u})^{\mathrm{T}} + \nabla\beta_2(\mathbf{u})\nabla\beta_1(\mathbf{u})^{\mathrm{T}}, \tag{10}$$
$$\psi_{12} = \int_0^\tau \rho_1(t)\rho_2(t)dt.$$

Hence, if we modify $\rho_1(t)$ and $\rho_2(t)$, keeping their temporal overlap ψ_{12} unchanged, then $\mathbf{M}(\mathbf{u},\tau)$ is also unchanged. In particular, when $\psi_{12} = 0$, switching of the expression order between $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$ does not affect $\mathbf{M}(\mathbf{u},\tau)$.

Figure 3 shows how ψ_{12} affects the shape of the extended sheet when $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$ are given respectively by a wide normal distribution (green gradient in panel (a)) and by a narrow normal distribution (blue contours in panel (a)). If $\rho_1(t) = \rho_2(t)$, then ψ_{12} attains its maximum value of 1. In this case, the growth metric (Figure 3e) induced from Eq. (8) is identical to the growth metric (Figure 3c) induced from a single morphogen distribution defined by $\beta(\mathbf{u}) = \beta_1(\mathbf{u}) + \beta_2(\mathbf{u})$. Therefore, the shape of the extended sheet (Figure 3h) resembles the basic shape for $\beta(\mathbf{u}) = \beta_1(\mathbf{u}) + \beta_2(\mathbf{u})$ (Figure 3d).

In contrast, if $\rho_1(t)$ and $\rho_2(t)$ have no temporal overlap, then ψ_{12} attains its minimum value of 0. In this case, the orientations of the substructure induced from $\beta_1(\mathbf{u})$ (left object in Figure 3b) and that induced from $\beta_2(\mathbf{u})$ (right object in Figure 3b) tend to be orthogonal to each other in the extended sheet (Figure 3j). Note that the sheet growth induced from $\beta_1(\mathbf{u})$ at the position of $\beta_2(\mathbf{u})$ broadens the shape of the substructure induced from $\beta_2(\mathbf{u})$ (Figure 3j) in comparison with its basic shape (right object in Figure 3b; see Appendix B.2 for its mathematical derivation). When ψ_{12} has an intermediate value between 0 and 1, the orientations of the substructures tend to have an angle between parallel and orthogonal (Figure 3i).

Analogously, we can fuse the basic shapes for an arbitrary number of morphogen distributions. For example, Figure 4 shows the fusion of three basic shapes specified by three morphogen distributions: $\beta_1(\mathbf{u})$ (green gradations in Figures 4a–4c), $\beta_2(\mathbf{u})$ (blue contour), and $\beta_3(\mathbf{u})$ (red contour), where only ψ_{23} is non-zero and affects the position of $\beta_3(\mathbf{u})$ as well as additional curving and twisting. The resulting shape variation depending on ψ_{23} may resemble the trimorphism of male mandibles of stag beetle species in genus *Dorcus*, depending on their body sizes [22,23].

(b) Morphogenesis of complex shapes

Fusion of multiple basic shapes with additional curving and twisting can generate various complex three-dimensional shapes for the extended sheet (Figure 5). Note that these complex shapes are generated from comparably much simpler settings for morphogen distributions given

by normal distributions or their linear combinations (Appendix C.1), and which may provide an

efficient genetic coding for various three-dimensional morphologies as discussed in Discussion

(b).

Figures 6–8 show the extended sheets obtained with the spring-vertex model, where the

morphogen distributions and their combinations are designed for mimicking heads and thoraxes

of beetles with horns: Figure 6 mimics genus *Dynastes* (e.g., *Dynastes hercules*) and genus

Xylotrupes, Figure 7 mimics genus *Chalcosoma* (e.g., *Chalcosoma chiron*), and Figure 8 mimics

genus *Sulcophanaeus* (e.g., *Sulcophanaeus faunus*) and genus *Diabroctis* (e.g., *Diabroctis*

mimas). Analogously to Figure 5, these three-dimensional shapes were generated by fusing

multiple basic shapes with additional curving and twisting, where all assumed morphogen

distributions are normal distributions or their linear combinations.

Discussion

(a) Growth regulation of epithelial sheet and developmental constraints

The present study showed that the non-uniform growth in the MGS growth model [18] can

transforms a planar epidermal sheet into various complex shapes emulating the horn shapes of

actual beetles. The MGS growth model is a deductively devised minimal model for achieving

smooth three-dimensional shapes, rather than representing the actual and concrete process of

arthropod morphogenesis. Nevertheless, the generated structures not only resemble the actual

horns of beetles as a whole but also share local shapes. For example, in the morphogenesis of

these horn-like shapes, the maximum sheet growth (corresponding to the maximum morphogen

gradient) is attained in the middle part of the horn and oriented toward the apex (Figure 2d),

following the empirically observed sheet growth for horn morphogenesis in beetles [9]. Moreover,

for the morphogenesis of such complex shape, the generated structures deviate spontaneously

from the shape of morphogen distributions, and the deviations bring local distortions that are in

common with the actual horns. In particular, the curving and/or twisting of the horns shown in

Figures 6–8 spontaneously transforms their cross-sections from round shapes into angular ones

that resemble actual beetle horns [4].

As the angular cross-sections, spontaneous deformation that deviates from the shape

corresponding to a morphogen distribution is caused by not being able to unfold a sheet as much

as it grew. The extended shape of a grown sheet corresponds to the minimization of the sum of

stretching energy (tangential to the sheet surface) and bending energy (orthogonal to the surface)

over the sheet. Provided that the sheet is sufficiently thin, the stretching energy has the dominant

contribution to this minimization. If the extended sheet has a shape such that the local sheet

growth measured in spatial coordinates exactly follows the growth metric deriving from morphogen distributions (equation (6)), then such a shape is called an isometric embedding for the growth metric, giving zero stretching energy over the sheet. For a sheet grown in the MGS growth model under a single morphogen distribution without additional curves and twists, the basic shape of the morphogen distribution gives the isometric embedding of the sheet, and hence the extended sheet can have a similar shape to the basic shape. In contrast, for more complex shapes generated from multiple morphogens with added curves and twists, neither basic shapes of the morphogen distributions nor their linear combinations do not give isometric embeddings, and hence the shape of the extended sheet can have significant local deviations from the basic shapes. Such deviations cause further deformation of the extended sheet as a byproduct of the global minimisation of local elastic energies. The angular cross-section of the central horn in Figure 7 and the concavity near the base of the left and right horns in Figure 8 can be interpreted as examples of such shapes. The byproduct shapes via deviating from the shape of morphogen distributions might be regarded as developmental constraints that bring about common morphological features shared by arthropods, and our model might represent some essential elements for such constraints.

Since the byproduct shapes are formed through the minimisation of elastic energies, they will contribute to the structure-mechanical stability of appendages. In particular, the horns of beetles as a weapon in male-male competition require high stability, and their angular cross-sections would be an adaptation to bring stability [4]. Notably, a rhinoceros beetle generates the angular shape of the horn by remodelling (i.e., adhesion and shrinkage) of a horn with round cross section developed through pupation [19]. Our result implies that the horn before the remodelling might already be oriented to have an angular shape.

(b) Modularity and evolution of exoskeletal morphogenesis

In a complex shape made by fusing multiple basic shapes under the MGS growth model, the substructure corresponding to each basic shape tends to maintain its shape similar to the basic shape. Notably, the size and shape of each basic shape can be changed separately by modifying the corresponding morphogen distribution. Moreover, the relative orientations among the substructures can also be changed by modifying the temporal overlaps among morphogen expressions. These properties of the MGS growth model imply that substructures of appendages, such as multiple horns, spines, or branches of horns, have developmental independency such that the morphogenesis of exoskeletal appendages is modular. This modularity explains why our simple growth model can easily generate complex structures.

Modularity results in developmental stability and evolutionary robustness against mutations. Even if an exoskeletal structure is structure-mechanically stable, mutations in the epidermal sheet growth may collapse the structure, because in principle, it is not warranted that an arbitrary grown sheet disturbed by mutations can be extended into a smooth three-dimensional shape with structure-mechanical stability. Such collapse of exoskeletal structure caused by mutation will be effectively reduced, if an exoskeletal appendage has modularity allowing its parts to develop independently, so that a mutation in one part may have little interference with other parts. In particular, under the MGS growth model, increasing/decreasing morphogen expression tends to induce only adding/removing appendages (such as spines and branches) or modifications (such as curving and twisting). Moreover, changes of the expression timings of morphogens tend to modify the aspect ratios and relative orientations of the substructures without breaking down the whole structure.

Since the modularity of the MGS growth model brings developmental stability and ease of adding modifications to the morphogenesis of exoskeletal appendages, it may contribute to explaining the rapid and dramatic evolution of beetle horns with extreme shapes and diversity. The horns of beetles have been evolving rapidly and dramatically despite their extreme sizes and elaborate shapes; they maintain structure-mechanical strength and developmental plasticity as weapons for males with good physiques [2]. This ambivalent feature between evolvability and developmental stability is a classical paradox in evolutionary developmental biology. One way to resolve this paradox is the modularity of developmental units at various scales [5]. Modularity via the MGS growth may allow the horns of beetles to evolve rapidly without losing their stability.

More fundamentally, developmental systems that enhance compatibility among mutations through modular morphogenesis, such as the MGS growth model, would be favoured by selection as they bring high developmental stability. Such systems might have been acquired by arthropods in their early evolution, paving the efficient way to evolve further novel and stable exoskeletons. Once developmental systems have evolved to generate highly stable exoskeletons, they can be hardened to that extent, thereby increasing performance in locomotion, attacks on prey, and defence against enemies. Thus, the emergence and rapid diversification of arthropods with hard exoskeletons during the Cambrian Explosion [24,25] may be related to the evolutionary emergence of developmental systems that efficiently produce novel and stable exoskeletons through modular morphogenesis.

(c) Limitations and future directions in connection with relevant studies

Because the MGS growth model is a deductively devised minimal model that relates shapes of extended sheets with hypothesized morphogen distributions, its relationship with actual molecules and biochemical processes must be examined carefully. For example, empirical studies on the morphogenesis of beetle horns have shown that the grown epithelial sheet has an intricate wrinkled structure [16,26], which unfolds and extends to form the horn through moulting to the pupa [9]. The wrinkle pattern is related to the direction and amount of local sheet growth; however, it remains unclear whether (i) the wrinkle formation *induces* sheet growth or (ii) it *is induced by* sheet growth. This study shows that it is possible to generate extended sheets of various shapes without wrinkle formation. The results support the possibility of (ii) but do not rule out the possibility of (i). Hence, it is necessary to analyse the the sheet deformation in spatial coordinates as a continuous dynamics that involves both the wrinkle structure formation through the sheet growth and its unfolding into a smooth three-dimensional shape.

Although the MGS growth model assumes an unstrained growth of the epidermal sheet, actual sheet growth and its folding can cause mechanical stress on the sheet. This mechanical stress would be a signal for each cell to determine the appropriate growth amounts. Further investigation is needed to clarify the distribution of the mechanical stress and how each microelement responds to it to grow in appropriate amounts (equations (6) or (9)) to generate a harmonious three-dimensional overall shape [8]. In our study, the head and thorax of each horned beetle were mimicked by a single sheet without separation (Figures 6–8). Such a global regulation of morphogenesis across multiple body segments is observed in some arthropods [27], and which might be essential for the harmonious morphogenesis of the whole body in terms of functional mechanics.

The MGS growth model is one of growth regulation models for cell sheet deformation. Since it assumes a single (type of) layered sheet, the growth metric corresponding to a protrusion (extending outward from the body surface) may be identical to that corresponding to a depression (sinking inward from the body surface). Hence, whether a protrusion or depression is formed depends on the initial growth direction and the surrounding structures. In contrast, some growth models proposed for plant leaves and petals assume multi-layered sheets such that the three-dimensional shape of the extended sheet can be uniquely determined under a given growth metric for each layer [28]. Even in morphogenesis of a hard exoskeleton like those of beetles and crabs, such a multi-layered sheet may work for orientating the sheet extension, as long as only one layer hardens afterwards (otherwise slight differences in the growth metric among hardened layers at the same position of the sheet can produce a large elastic energy that may cause structure-

372　mechanical instability).

Conclusion

We showed that the MGS growth model allows an epidermal sheet to grow into various complex structures emulating arthropods' appendages including beetle horns, by fusion of multiple basic shapes controlled separately by different morphogen distributions. This suggests that such a developmental system can bring modularity and developmental stability to the morphogenesis of exoskeletal appendages and provide insights into the diversity of arthropod morphology.

Literature Cited

1.　　Kijimoto T, Pespeni M, Beckers O, Moczek AP. 2013 Beetle horns and horned beetles: emerging models in developmental evolution and ecology. *WIREs Dev. Biol.* **2**, 405–418. (doi:10.1002/wdev.81)

2.　　Emlen DJ, Corley Lavine L, Ewen-Campen B. 2007 On the origin and evolutionary diversification of beetle horns. *Proc. Natl. Acad. Sci.* **104**, 8661–8668. (doi:10.1073/pnas.0701209104)

3.　　Emlen DJ. 1996 ARTIFICIAL SELECTION ON HORN LENGTH-BODY SIZE ALLOMETRY IN THE HORNED BEETLE ONTHOPHAGUS ACUMINATUS (COLEOPTERA: SCARABAEIDAE). *Evolution* **50**, 1219–1230. (doi:10.1111/j.1558-5646.1996.tb02362.x)

4.　　McCullough EL, Tobalske BW, Emlen DJ. 2014 Structural adaptations to diverse fighting styles in sexually selected weapons. *Proc. Natl. Acad. Sci.* **111**, 14484–14488. (doi:10.1073/pnas.1409585111)

5.　　Gilbert SF, Epel D. 2008 *Ecological Developmental Biology: Integrating Epigenetics, Medicine, and Evolution*. Sunderland, Mass: Sinauer Associates Inc.

6.　　Valentine JW, Collins AG. 2000 The significance of moulting in Ecdysozoan evolution. *Evol. Dev.* **2**, 152–156. (doi:10.1046/j.1525-142x.2000.00043.x)

7.　　Minelli A, Boxshall G, Fusco G. 2013 *Arthropod Biology and Evolution: Molecules, Development, Morphology*. 2013th edition. Berlin ; New York: Springer.

8.　　Zhou Z, Alégot H, Irvine KD. 2019 Oriented Cell Divisions Are Not Required for Drosophila Wing Shape. *Curr. Biol.* **29**, 856-864.e3. (doi:10.1016/j.cub.2019.01.044)

9.　　Matsuda K, Gotoh H, Adachi H, Inoue Y, Kondo S. 2021 Computational analyses decipher the primordial folding coding the 3D structure of the beetle horn. *Sci. Rep.* **11**, 1017. (doi:10.1038/s41598-020-79757-2)

10.   Hu Y, Linz DM, Moczek AP. 2019 Beetle horns evolved from wing serial
homologs. *Science* **366**, 1004–1007. (doi:10.1126/science.aaw2980)

11.   Miura T. 2019 Juvenile hormone as a physiological regulator mediating
phenotypic plasticity in pancrustaceans. *Dev. Growth Differ.* **61**, 85–96.
(doi:10.1111/dgd.12572)

12.   Nijhout HF. 2019 The multistep morphing of beetle horns. *Science* **366**, 946–
947. (doi:10.1126/science.aaz9010)

13.   Tripathi BK, Irvine KD. 2022 The wing imaginal disc. *Genetics* **220**, iyac020.
(doi:10.1093/genetics/iyac020)

14.   Adachi H, Matsuda K, Niimi T, Kondo S, Gotoh H. 2020 Genetical control of
2D pattern and depth of the primordial furrow that prefigures 3D shape of the
rhinoceros beetle horn. *Sci. Rep.* **10**, 18687. (doi:10.1038/s41598-020-75709-y)

15.   Osterfield M, Du X, Schüpbach T, Wieschaus E, Shvartsman SY. 2013 Three-
Dimensional Epithelial Morphogenesis in the Developing Drosophila Egg. *Dev. Cell* **24**,
400–410. (doi:10.1016/j.devcel.2013.01.017)

16.   Inoue Y, Tateo I, Adachi T. 2020 Epithelial tissue folding pattern in confined
geometry. *Biomech. Model. Mechanobiol.* **19**, 815–822. (doi:10.1007/s10237-019-
01249-8)

17.   Matsuda K, Gotoh H, Tajika Y, Sushida T, Aonuma H, Niimi T, Akiyama M,
Inoue Y, Kondo S. 2017 Complex furrows in a 2D epithelial sheet code the 3D structure
of a beetle horn. *Sci. Rep.* **7**, 13939. (doi:10.1038/s41598-017-14170-w)

18.   Ito HC. 2016 Metric dynamics for membrane transformation through regulated
cell proliferation. (doi:10.48550/arXiv.1605.01893)

19.   Matsuda K, Adachi H, Gotoh H, Inoue Y, Kondo S. 2024 Adhesion and
shrinkage transform the rounded pupal horn into an angular adult horn in Japanese
rhinoceros beetle. *Development* **151**, dev202082. (doi:10.1242/dev.202082)

20.   Anderson R *et al.* 2021 MFEM: A modular finite element methods library.
Comput. Math. Appl. **81**, 42–74. (doi:10.1016/j.camwa.2020.06.009)

21.   Matsuda K, Gotoh H, Tajika Y, Sushida T, Aonuma H, Niimi T, Akiyama M,
Inoue Y, Kondo S. 2017 Complex furrows in a 2D epithelial sheet code the 3D structure
of a beetle horn. *Sci. Rep.* **7**, 13939. (doi:10.1038/s41598-017-14170-w)

22.   Iguchi Y. 2013 Male mandible trimorphism in the stag beetle Dorcus rectus
(Coleoptera: Lucanidae). *EJE* **110**, 159–163. (doi:10.14411/eje.2013.022)

23.   Wang B-X, Wang Y-N, Zhang F, Xia W. In press. Male trimorphism in Dorcus
hopei: evaluating the superiority of different morphs in Mate competition. *Chin. J. Appl.*
Entomol. **60**, 178–185.

24.   Edgecombe GD, Legg DA. 2014 Origins and early evolution of arthropods. *Palaeontology* **57**, 457–468. (doi:10.1111/pala.12105)

25.   Zhang X, Shu D. 2021 Current understanding on the Cambrian Explosion: questions and answers. *PalZ* **95**, 641–660. (doi:10.1007/s12542-021-00568-5)

26.   Adachi H, Matsuda K, Niimi T, Inoue Y, Kondo S, Gotoh H. 2018 Anisotropy of cell division and epithelial sheet bending via apical constriction shape the complex folding pattern of beetle horn primordia. *Mech. Dev.* **152**, 32–37. (doi:10.1016/j.mod.2018.06.003)

27.   Kaji T, Møller OS, Tsukagoshi A. 2011 A bridge between original and novel states: ontogeny and function of "suction discs" in the Branchiura (Crustacea). *Evol. Dev.* **13**, 119–126. (doi:10.1111/j.1525-142X.2011.00462.x)

28.   Kennaway R, Coen E. 2019 Volumetric finite-element modelling of biological growth. *Open Biol.* **9**, 190057. (doi:10.1098/rsob.190057)

29.   Hobson MP, Efstathiou GP, Lasenby AN. 2006 *General Relativity: An Introduction for Physicists*. Cambridge: Cambridge University Press. (doi:10.1017/CBO9780511790904)

Figure legends

Figure 1

Initial state, grown state, and extended state of epidermal sheet in spatial coordinate system $\mathbf{r} = (x, y, z)^{\mathrm{T}}$ and material coordinate system $\mathbf{u} = (X, Y)^{\mathrm{T}}$. (a) Initial flat sheet in the spatial coordinates. Green gradient indicates morphogen intensity (light green: low, deep green: high). (b) Initial sheet in the material coordinates. Green gradient indicates morphogen intensity as in (a). The small triangle indicates the maximum morphogen intensity. (c) Grown sheet in the spatial coordinates (not analysed in this study and hence unknown). (d) Grown sheet in the material coordinates, represented with the growth metric $\mathbf{M}(\mathbf{u})$ given by equation (6). The cyan-red gradient indicates the square-root of the maximum eigenvalue of $\mathbf{M}(\mathbf{u})$ at each position (i.e., how many times each microelement of the sheet is elongated) with its eigenvector (i.e., growth direction) indicated with the short cyan line segment at that position. The small yellow triangle indicates a maxima of Riemann scalar curvature (expected to become a projection or follow in the extended sheet). (e) Extended sheet in the spatial coordinates. The cyan-red gradient indicates the maximum eigenvalue of $\mathbf{M}(\mathbf{u})$, and which is equivalent to (d). (f) Extended sheet in the material coordinates represented with the extension metric $\mathbf{M}_{\mathrm{ext}}(\mathbf{u})$ that was calculated from the extended sheet (Appendix B.1) and indicated in the same manner as in (d).

Figure 2

Initial state, grown state, and extended state for generation of simple horn-like shape and its modification. (a) Morphogen intensity distribution $\beta(\mathbf{u})$ defined as an isotropic Normal distribution. (b) Basic shape for $\beta(\mathbf{u})$, i.e., representation of morphogen intensities as heights. (c) Basic growth metric $\mathbf{M}_{\beta}(\mathbf{u})$, i.e., the growth metric calculated from the basic shape (see Appendix B). (d) Grown sheet produced from $\beta(\mathbf{u})$ under the MGS growth model, represented with the growth metric $\mathbf{M}(\mathbf{u})$ given by equation (6). (e) Grown sheet for curving the basic shape. (f) Grown sheet for curving and twisting the basic shape. (g), (h), and (i) show extended sheets from (d), (e), and (f), respectively, in the spatial coordinates (upper panels) and the material coordinates (lower panels, represented with the extension metric $\mathbf{M}_{\mathrm{ext}}(\mathbf{u})$). See Appendix D for how to add curve and twist to the basic shape.

  Figure 3

  Fusion of two basic shapes by heterochronic expression of two morphogen distributions. (a) Two

  morphogen distributions $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$ plotted respectively with green gradient and blue

  contours with small triangles indicating their maximums. (b) Basic shapes for $\beta_1(\mathbf{u})$ (left) and

  $\beta_2(\mathbf{u})$ (right). (c) Basic growth metric for $\beta_1(\mathbf{u}) + \beta_2(\mathbf{u})$. (d) Basic shape for $\beta_1(\mathbf{u}) + \beta_2(\mathbf{u})$.

  (e), (f), and (g) show grown sheets for $\psi_{12} = 1$, $\psi_{12} = 0.5$, and $\psi_{12} = 0$, respectively. (h), (i),

  and (j) show extended sheets for $\psi_{12} = 1$, $\psi_{12} = 0.5$, and $\psi_{12} = 0$, respectively. Plotting format:

  panel (a) is the same as figure 2a, panels (b), (d), and (h-j) are the same as figure 2g-i, and panels

  (e-g) are the same as figure 2d-f. See Appendix C.1 for morphogen distributions and parameters.

  Figure 4

  Extended sheets mimicking three mandible types of stag beetles depending on their body sizes

  (large-, middle-, and small-types). The extended sheets are generated by fusing the basic shapes

  of three morphogen distributions, $\beta_1(\mathbf{u})$, $\beta_2(\mathbf{u}, \psi_{23})$, and $\beta_3(\mathbf{u}, \psi_{23})$, where only β_2 and β_3 have

  temporal overlap in their expression (i.e., $\psi_{12} = \psi_{13} = 0$ and $0 \le \psi_{23} \le 1$). (a), (b), and (c)

  show morphogen distributions for the large-type ($\psi_{23} = 1$), middle-type ($\psi_{23} = 0.5$), and small-

  type ($\psi_{23} = 0$). Morphogen distributions β_1, β_2, and β_3 are plotted respectively as green

  gradient, blue contours, and red contours, with small triangles indicating their maximums. (d),

  (e), and (f) show grown sheets for the large-type ($\psi_{23} = 1$), middle-type ($\psi_{23} = 0.5$), and small-

  type ($\psi_{23} = 0$). (g), (h), and (i) show extended sheets (upper panels) and the corresponding

  extension metrics (lower panels) for the large-type ($\psi_{23} = 1$), middle-type ($\psi_{23} = 0.5$), and

  small-type ($\psi_{23} = 0$). Plotting formats are the same as in figure 3. See Appendix C.1 for

  morphogen distributions and parameters.

  Figure 5

  Extended sheets of complex three-dimensional shapes generated by fusing multiple basic shapes.

  (a-c) Morphogen distributions. (d-f) Grown sheets represented with growth metrics. (g-i)

  Extended sheets. Plotting formats are the same as in figures 3 and 4. See Appendix C.1 for

  morphogen distributions and parameters.

  Figure 6

  Extended sheet mimicking the head and thorax of beetles in genus *Dynastes* (e.g., *D. hercules*)

  and genus *Xylotrupes* (e.g., *X. florensis*). The upper left panel shows the grown sheet represented

  with growth metric, and the other panels show its extended shape viewed from different

522 perspectives. See Appendix C.2 for morphogen distributions and parameters.

Figure 7

524 Extended sheet mimicking the head and thorax of beetles in genus *Chalcosoma* (e.g., *C.*
525 *caucasus*). Plotting formats are the same as in figure 6. See Appendix C.2 for morphogen
526 distributions and parameters.

Figure 8

528 Extended sheet mimicking the head and thorax of beetles in genus *Sulcophanaeus* (e.g., *S.*
529 *faunus*). Plotting formats are the same as in figure 6. See Appendix C.2 for morphogen
530 distributions and parameters.

Appendix A: Derivation of equation (3)

Here we derive equation (3) in the main text in a more general form than [18].

A.1. Description of non-uniform sheet growth

Since the microelement including an arbitrary position $\mathbf{u}$ is assumed to be unstrained in the spatial coordinates, the square of the unstrained length $l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta u}, t)$ for a short line segment from $\mathbf{u} = (X, Y)^{\mathrm{T}}$ to $\mathbf{u} + \boldsymbol{\delta u} = (X + \delta X, Y + \delta Y)^{\mathrm{T}}$ is expressed as

$$l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta u}, t)^2 = |\tilde{\mathbf{r}}(\mathbf{u} + \boldsymbol{\delta u}, t) - \tilde{\mathbf{r}}(\mathbf{u}, t)|^2 + O(|\boldsymbol{\delta u}|^3), \tag{A.1}$$

where $O(|\boldsymbol{\delta u}|^3)$ contains the third- and higher-order terms of $\boldsymbol{\delta u}$, so that there exists a constant C such that $\lim\limits_{|\boldsymbol{\delta u}| \to 0} \dfrac{O(|\boldsymbol{\delta u}|^3)}{|\boldsymbol{\delta u}|^3} < C$ holds. To transform this equation, we expand $\tilde{\mathbf{r}}(\mathbf{u} + \boldsymbol{\delta u}, t)$ around $\mathbf{u}$ as

$$\tilde{\mathbf{r}}(\mathbf{u} + \boldsymbol{\delta u}, t) = \begin{pmatrix} \tilde{x}(\mathbf{u} + \boldsymbol{\delta u}, t) \\ \tilde{y}(\mathbf{u} + \boldsymbol{\delta u}, t) \\ \tilde{z}(\mathbf{u} + \boldsymbol{\delta u}, t) \end{pmatrix} = \begin{pmatrix} \tilde{x}(\mathbf{u}, t) + \dfrac{\partial \tilde{x}(\mathbf{u}, t)}{\partial X} \delta X + \dfrac{\partial \tilde{x}(\mathbf{u}, t)}{\partial Y} \delta Y + O(|\boldsymbol{\delta u}|^2) \\ \tilde{y}(\mathbf{u}, t) + \dfrac{\partial \tilde{y}(\mathbf{u}, t)}{\partial X} \delta X + \dfrac{\partial \tilde{y}(\mathbf{u}, t)}{\partial Y} \delta Y + O(|\boldsymbol{\delta u}|^2) \\ \tilde{z}(\mathbf{u}, t) + \dfrac{\partial \tilde{z}(\mathbf{u}, t)}{\partial X} \delta X + \dfrac{\partial \tilde{z}(\mathbf{u}, t)}{\partial Y} \delta Y + O(|\boldsymbol{\delta u}|^2) \end{pmatrix}$$

$$= \tilde{\mathbf{r}}(\mathbf{u}, t) + \mathbf{P}(\mathbf{u}, t)\boldsymbol{\delta u} + \begin{pmatrix} O(|\boldsymbol{\delta u}|^2) \\ O(|\boldsymbol{\delta u}|^2) \\ O(|\boldsymbol{\delta u}|^2) \end{pmatrix}, \tag{A.2}$$

$$\mathbf{P}(\mathbf{u}, t) = \begin{pmatrix} \partial_X \tilde{x} & \partial_Y \tilde{x} \\ \partial_X \tilde{y} & \partial_Y \tilde{y} \\ \partial_X \tilde{z} & \partial_Y \tilde{z} \end{pmatrix} := \begin{pmatrix} \partial \tilde{x}(\mathbf{u}, t)/\partial X & \partial \tilde{x}(\mathbf{u}, t)/\partial Y \\ \partial \tilde{y}(\mathbf{u}, t)/\partial X & \partial \tilde{y}(\mathbf{u}, t)/\partial Y \\ \partial \tilde{z}(\mathbf{u}, t)/\partial X & \partial \tilde{z}(\mathbf{u}, t)/\partial Y \end{pmatrix}.$$

Substitution of equation (A.2) into (A.1) gives

$$\begin{aligned} l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta u}, t)^2 &= [\tilde{\mathbf{r}}(\mathbf{u} + \boldsymbol{\delta u}, t) - \tilde{\mathbf{r}}(\mathbf{u}, t)]^{\mathrm{T}}[\tilde{\mathbf{r}}(\mathbf{u} + \boldsymbol{\delta u}, t) - \tilde{\mathbf{r}}(\mathbf{u}, t)] + O(|\boldsymbol{\delta u}|^3) \\ &= [\mathbf{P}(\mathbf{u}, t)\boldsymbol{\delta u}]^{\mathrm{T}}[\mathbf{P}(\mathbf{u}, t)\boldsymbol{\delta u}] + O(|\boldsymbol{\delta u}|^3) \\ &= \boldsymbol{\delta u}^{\mathrm{T}} \mathbf{P}(\mathbf{u}, t)^{\mathrm{T}} \mathbf{P}(\mathbf{u}, t)\boldsymbol{\delta u} + O(|\boldsymbol{\delta u}|^3) \\ &= \boldsymbol{\delta u}^{\mathrm{T}} \mathbf{M}(\mathbf{u}, t)\boldsymbol{\delta u} + O(|\boldsymbol{\delta u}|^3) \end{aligned} \tag{A.3}$$

with

$$\mathbf{M}(\mathbf{u}, t) = \begin{pmatrix} M_{XX}(\mathbf{u}, t) & M_{XY}(\mathbf{u}, t) \\ M_{XY}(\mathbf{u}, t) & M_{YY}(\mathbf{u}, t) \end{pmatrix} = \mathbf{P}(\mathbf{u}, t)^{\mathrm{T}} \mathbf{P}(\mathbf{u}, t),$$

$$M_{XX}(\mathbf{u}, t) = [\partial_X \tilde{x}]^2 + [\partial_X \tilde{y}]^2 + [\partial_X \tilde{z}]^2 = \left| \frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial X} \right|^2,$$

$$M_{YY}(\mathbf{u}, t) = [\partial_Y \tilde{x}]^2 + [\partial_Y \tilde{y}]^2 + [\partial_Y \tilde{z}]^2 = \left| \frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial Y} \right|^2, \tag{A.4}$$

$$M_{XY}(\mathbf{u}, t) = [\partial_X \tilde{x}][\partial_Y \tilde{x}] + [\partial_X \tilde{y}][\partial_Y \tilde{y}] + [\partial_X \tilde{z}][\partial_Y \tilde{z}] = \left[\frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial X} \right]^{\mathrm{T}} \left[\frac{\partial \tilde{\mathbf{r}}(\mathbf{u}, t)}{\partial Y} \right].$$

Then, the growth amount for the line segment from the initial state to time t, is defined by

548    $l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, t)/l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, 0)$, and which for an infinitesimally short line segment (i.e., $|\boldsymbol{\delta}\mathbf{u}| \to$

549   0) describes the growth amount for the sheet at position $\mathbf{u}$ along the direction $\mathbf{e} = \boldsymbol{\delta}\mathbf{u} \, / \, |\boldsymbol{\delta}\mathbf{u}|$ at

550   time t:

$$
\begin{aligned}
L(\mathbf{u}, \mathbf{e}, t) &= \lim_{|\boldsymbol{\delta}\mathbf{u}| \to 0} \frac{l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, t)}{l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, 0)} = \lim_{|\boldsymbol{\delta}\mathbf{u}| \to 0} \frac{\sqrt{\boldsymbol{\delta}\mathbf{u}^{\mathrm{T}}\mathbf{M}(\mathbf{u}, t)\boldsymbol{\delta}\mathbf{u} + O(|\boldsymbol{\delta}\mathbf{u}|^3)}}{|\boldsymbol{\delta}\mathbf{u}|} \\
&= \sqrt{\left[\frac{\boldsymbol{\delta}\mathbf{u}}{|\boldsymbol{\delta}\mathbf{u}|}\right]^{\mathrm{T}} \mathbf{M}(\mathbf{u}, t)\left[\frac{\boldsymbol{\delta}\mathbf{u}}{|\boldsymbol{\delta}\mathbf{u}|}\right]} = \sqrt{\mathbf{e}^{\mathrm{T}}\mathbf{M}(\mathbf{u}, t)\mathbf{e}}.
\end{aligned} \tag{A.5}
$$

A.2. Locally geodesic coordinate system

To describe the morphogen intensity gradient along the sheet in the spatial coordinates, we

introduce a local two-dimensional coordinate system $\mathbf{u}^\circ = (X^\circ, Y^\circ)^{\mathrm{T}}$ on the sheet around each

focal position $\tilde{\mathbf{r}}(\hat{\mathbf{u}}, t)$ corresponding to $\hat{\mathbf{u}} = (\hat{X}, \hat{Y})^{\mathrm{T}}$. For notational simplicity, we assume $\hat{\mathbf{u}} = \mathbf{0}$

without loss of generality and drop the arguments "$(\hat{\mathbf{u}}, t)$" from variables evaluated at $\hat{\mathbf{u}}$ and t.

Since the sheet growth in the spatial coordinates is assumed to be unstrained, the unstrained length

from $\mathbf{u}$ to $\mathbf{u} + \boldsymbol{\delta}\mathbf{u}$ can be expressed as $l(\mathbf{u}, \mathbf{u} + \boldsymbol{\delta}\mathbf{u}, t) \simeq |\boldsymbol{\delta}\mathbf{u}^\circ|^2$ for sufficiently small $|\mathbf{u} - \hat{\mathbf{u}}|$

and $|\boldsymbol{\delta}\mathbf{u}|$. Such a coordinate system is called the locally geodesic coordinate system [29], and in

which $\boldsymbol{\delta}\mathbf{u}$ is expressed as $\boldsymbol{\delta}\mathbf{u}^\circ$ satisfying

$$
\boldsymbol{\delta}\mathbf{u}^\circ = \mathbf{W}\boldsymbol{\delta}\mathbf{u} + \frac{1}{2}\mathbf{W}\begin{pmatrix} \boldsymbol{\delta}\mathbf{u}^{\mathrm{T}}\begin{pmatrix} \Gamma_{XX}^X & \Gamma_{XY}^X \\ \Gamma_{XY}^X & \Gamma_{YY}^X \end{pmatrix}\boldsymbol{\delta}\mathbf{u} \\ \boldsymbol{\delta}\mathbf{u}^{\mathrm{T}}\begin{pmatrix} \Gamma_{XX}^Y & \Gamma_{XY}^Y \\ \Gamma_{XY}^Y & \Gamma_{YY}^Y \end{pmatrix}\boldsymbol{\delta}\mathbf{u} \end{pmatrix}. \tag{A.6}
$$

562   Here, $\mathbf{W}$ is a two-by-two matrix obtained by the diagonalisation of $\mathbf{M}$ as

$$
\begin{aligned}
\mathbf{M} &= \mathbf{E}\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}\mathbf{E}^{\mathrm{T}} = \left[\begin{pmatrix} \sqrt{\lambda_1} & 0 \\ 0 & \sqrt{\lambda_2} \end{pmatrix}\mathbf{E}^{\mathrm{T}}\right]^{\mathrm{T}}\left[\begin{pmatrix} \sqrt{\lambda_1} & 0 \\ 0 & \sqrt{\lambda_2} \end{pmatrix}\mathbf{E}^{\mathrm{T}}\right] = \mathbf{W}^{\mathrm{T}}\mathbf{W}, \\
\mathbf{W} &= \begin{pmatrix} \sqrt{\lambda_1} & 0 \\ 0 & \sqrt{\lambda_2} \end{pmatrix}\mathbf{E}^{\mathrm{T}},
\end{aligned} \tag{A.7}
$$

where λ_1 and λ_2 are the two eigenvalues of $\mathbf{M}$ with their eigenvectors $\mathbf{v}_1$ and $\mathbf{v}_2$ being combined

into the matrix $\mathbf{E} = (\mathbf{v}_1, \mathbf{v}_2)$. Note that all Γs are negligible because all terms that can belong to

$O(|\boldsymbol{\delta}\mathbf{u}|^2)$ in equation (A.6) vanish when the limit $|\boldsymbol{\delta}\mathbf{u}| \to 0$ is taken afterwards. (Γs are called the

Christoffel symbols of the second kind in differential geometry of surfaces, and which are

adjusted first derivatives of the growth metric, defined by $\Gamma_{ji}^k = \sum_m \frac{1}{2} M_{km}^{-1}\left[\frac{\partial M_{im}(\mathbf{u}, t)}{\partial j} + \right.$

$\left. \frac{\partial M_{jm}(\mathbf{u}, t)}{\partial i} - \frac{\partial M_{ji}(\mathbf{u}, t)}{\partial j}\right]$ with $i, j, k, m \in \{X, Y\}$ and M_{km}^{-1} describing the (k, m) component of $\mathbf{M}^{-1}$

[29]). Therefore, we transform equation (A.6) into

$$\delta\mathbf{u}^\circ = \mathbf{W}\delta\mathbf{u} + \begin{pmatrix} O(|\delta\mathbf{u}|^2) \\ O(|\delta\mathbf{u}|^2) \end{pmatrix},$$

(A.8)

$$\delta\mathbf{u} = \mathbf{W}^{-1}\delta\mathbf{u}^\circ + \begin{pmatrix} O(|\delta\mathbf{u}^\circ|^2) \\ O(|\delta\mathbf{u}^\circ|^2) \end{pmatrix}.$$

A.3. Unidirectional and isotropic sheet growth

In the locally geodesic coordinates, $\mathbf{u}^\circ = (X^\circ, Y^\circ)^\mathrm{T}$, we consider both of unidirectional growth and isotropic growth for the sheet with their rates denoted by η and ζ, respectively. We describe the direction of the unidirectional growth at time t by a unit vector $\mathbf{e}^{*\circ} = (e_X^{*\circ}, e_Y^{*\circ})^\mathrm{T}$. For convenience, we introduce the orthogonal unit vector to $\mathbf{e}^{*\circ}$, given by $\mathbf{n}^{*\circ} = (n_X^{*\circ}, n_Y^{*\circ})^\mathrm{T} = (e_Y^{*\circ}, -e_X^{*\circ})^\mathrm{T}$. Then, we can decompose $\delta\mathbf{u}^\circ$ into the element along with $\mathbf{e}^{*\circ}$ and that along with $\mathbf{n}^{*\circ}$:

$$\begin{aligned} \delta\mathbf{u}^\circ &= q_e\mathbf{e}^{*\circ} + q_n\mathbf{n}^{*\circ}, \\ q_e &= [\mathbf{e}^{*\circ\mathrm{T}}\delta\mathbf{u}^\circ], \\ q_n &= [\mathbf{n}^{*\circ\mathrm{T}}\delta\mathbf{u}^\circ]. \end{aligned}$$

(A.9)

From time t to time $t' = t + \Delta t$, the q_e and q_n become $q_e' = [1 + \eta\Delta t + \zeta\Delta t + O(\Delta t^2)]q_e \simeq [1 + \eta\Delta t + \zeta\Delta t]q_e$ and $q_n' = [1 + \zeta\Delta t + O(\Delta t^2)]q_n \simeq [1 + \zeta\Delta t]q_n$, and which approximately gives

$$\begin{aligned} \delta\mathbf{u}^{\circ\prime} &= q_e'\mathbf{e}^{*\circ} + q_n'\mathbf{n}^{*\circ} \\ &= [1 + \eta\Delta t + \zeta\Delta t]q_e\mathbf{e}^{*\circ} + [1 + \zeta\Delta t]q_n\mathbf{n}^{*\circ} \\ &= [1 + \zeta\Delta t][q_e\mathbf{e}^{*\circ} + q_n\mathbf{n}^{*\circ}] + \eta\Delta t q_e\mathbf{e}^{*\circ} \\ &= [1 + \zeta\Delta t]\delta\mathbf{u}^\circ + \eta\Delta t[\mathbf{e}^{*\circ\mathrm{T}}\delta\mathbf{u}^\circ]\mathbf{e}^{*\circ} \\ &= (\mathbf{I} + [\zeta\mathbf{I} + \eta\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}]\Delta t)\delta\mathbf{u}^\circ \\ &= [\mathbf{I} + \mathbf{B}\Delta t]\delta\mathbf{u}^\circ = [\mathbf{I} + \mathbf{B}\Delta t]\mathbf{W}\delta\mathbf{u} + \begin{pmatrix} O(|\delta\mathbf{u}|^2) \\ O(|\delta\mathbf{u}|^2) \end{pmatrix}, \end{aligned}$$

(A.10)

where $\mathbf{B} = \zeta\mathbf{I} + \eta\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}$. Hence, the growth amount at $\hat{\mathbf{u}} = \mathbf{0}$ along with $\mathbf{e} = \delta\mathbf{u} / |\delta\mathbf{u}|$ at time t' is transformed as

$$\begin{aligned} L(\hat{\mathbf{u}}, \mathbf{e}, t') &= \lim_{|\delta\mathbf{u}|\to 0} \frac{l(\hat{\mathbf{u}}, \delta\mathbf{u}, t')}{l(\hat{\mathbf{u}}, \delta\mathbf{u}, 0)} = \lim_{|\delta\mathbf{u}|\to 0} \sqrt{\frac{\delta\mathbf{u}^{\circ\prime\mathrm{T}}\delta\mathbf{u}^{\circ\prime} + O\left(|\delta\mathbf{u}^{\circ\prime}|^3\right)}{|\delta\mathbf{u}|^2}} \\ &= \lim_{|\delta\mathbf{u}|\to 0} \sqrt{\frac{\delta\mathbf{u}^\mathrm{T}\mathbf{W}^\mathrm{T}[\mathbf{I} + \mathbf{B}\Delta t]^\mathrm{T}[\mathbf{I} + \mathbf{B}\Delta t]\mathbf{W}\delta\mathbf{u} + O(|\delta\mathbf{u}|^3)}{|\delta\mathbf{u}|^2}} \\ &= \sqrt{\mathbf{e}^\mathrm{T}\mathbf{W}^\mathrm{T}[\mathbf{I} + \mathbf{B}\Delta t]^\mathrm{T}[\mathbf{I} + \mathbf{B}\Delta t]\mathbf{W}\mathbf{e}} \\ &= \sqrt{\mathbf{e}^\mathrm{T}\mathbf{M}\mathbf{e} + 2\mathbf{e}^\mathrm{T}[\mathbf{W}^\mathrm{T}\mathbf{B}\mathbf{W}]\mathbf{e}\Delta t + \mathbf{e}^\mathrm{T}\mathbf{W}^\mathrm{T}\mathbf{B}^\mathrm{T}\mathbf{B}\mathbf{W}\mathbf{e}\Delta t^2} \\ &= \sqrt{L(\hat{\mathbf{u}}, \mathbf{e}, t)^2 + 2\mathbf{e}^\mathrm{T}[\mathbf{W}^\mathrm{T}\mathbf{B}\mathbf{W}]\mathbf{e}\Delta t + O(\Delta t^2)}, \end{aligned}$$

(A.11)

where $\mathbf{B}^\mathrm{T} = [\zeta\mathbf{I} + \eta\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}]^\mathrm{T} = \zeta\mathbf{I} + \eta\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}} = \mathbf{B}$ is used. From equations (A.11) and (A.7) we get

$$\begin{aligned}
\frac{dL(\hat{\mathbf{u}}, \mathbf{e}, t)^2}{dt} &= \lim_{\Delta t \to 0} \frac{L(\hat{\mathbf{u}}, \mathbf{e}, t')^2 - L(\hat{\mathbf{u}}, \mathbf{e}, t)^2}{\Delta t} \\
&= \lim_{\Delta t \to 0} \frac{1}{\Delta t}[2\mathbf{e}^{\mathrm{T}}[\mathbf{W}^{\mathrm{T}}\mathbf{BW}]\mathbf{e}\Delta t + O(\Delta t^2)] \\
&= 2\mathbf{e}^{\mathrm{T}}[\mathbf{W}^{\mathrm{T}}\mathbf{BW}]\mathbf{e} \\
&= 2\mathbf{e}^{\mathrm{T}}\mathbf{W}^{\mathrm{T}}[\zeta\mathbf{I} + \eta\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}]\mathbf{We} \\
&= \mathbf{e}^{\mathrm{T}}[2\zeta\mathbf{M} + 2\eta\mathbf{W}^{\mathrm{T}}\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}\mathbf{W}]\mathbf{e}.
\end{aligned} \tag{A.12}$$

Since $\mathbf{e}$ is invariant against time, the following relationship holds good:

$$\frac{dL(\hat{\mathbf{u}}, \mathbf{e}, t)^2}{dt} = \frac{d[\mathbf{e}^{\mathrm{T}}\mathbf{Me}]}{dt} = \mathbf{e}^{\mathrm{T}}\frac{d\mathbf{M}}{dt}\mathbf{e}. \tag{A.13}$$

Because equations (A.12) and (A.13) both hold for an arbitrary $\mathbf{e}$, we obtain

$$\frac{d\mathbf{M}}{dt} = 2\eta\mathbf{W}^{\mathrm{T}}\mathbf{e}^{*\circ}\mathbf{e}^{*\circ\mathrm{T}}\mathbf{W} + 2\zeta\mathbf{M}. \tag{A.14}$$

A.4. Sheet growth regulation by morphogen intensity gradient

Now we assume that the direction of the unidirectional growth, $\mathbf{e}^{*\circ}$, is given by the gradient $\nabla^\circ \beta$ of morphogen intensity $\beta(\mathbf{u})$ evaluated at $\hat{\mathbf{u}} = \mathbf{0}$ in the locally geodesic coordinates, $\mathbf{u}^\circ$:

$$\mathbf{e}^{*\circ} = \frac{\nabla^\circ\beta}{|\nabla^\circ\beta|},$$
$$\nabla^\circ\beta = \begin{pmatrix} \dfrac{\partial\beta}{\partial X^\circ} \\ \dfrac{\partial\beta}{\partial Y^\circ} \end{pmatrix}. \tag{A.15}$$

Since equation (A.8) gives $\boldsymbol{\delta}\mathbf{u}^{\mathrm{T}} = \boldsymbol{\delta}\mathbf{u}^{\circ\mathrm{T}}\mathbf{W}^{-1\mathrm{T}}$ for infinitesimally small $|\boldsymbol{\delta}\mathbf{u}^\circ|$, we see that $\mathbf{W}^{-1\mathrm{T}}$ can be expressed as

$$\mathbf{W}^{-1\mathrm{T}} = \begin{pmatrix} \dfrac{\partial X}{\partial X^\circ} & \dfrac{\partial Y}{\partial X^\circ} \\ \dfrac{\partial X}{\partial Y^\circ} & \dfrac{\partial Y}{\partial Y^\circ} \end{pmatrix}. \tag{A.16}$$

By using equation (A.16), we transform $\nabla^\circ\beta$ as

$$\nabla^\circ\beta = \begin{pmatrix} \dfrac{\partial\beta}{\partial X^\circ} \\ \dfrac{\partial\beta}{\partial Y^\circ} \end{pmatrix} = \begin{pmatrix} \dfrac{\partial X}{\partial X^\circ}\dfrac{\partial\beta(\mathbf{u})}{\partial X} + \dfrac{\partial Y}{\partial X^\circ}\dfrac{\partial\beta(\mathbf{u})}{\partial Y} \\ \dfrac{\partial X}{\partial Y^\circ}\dfrac{\partial\beta(\mathbf{u})}{\partial X} + \dfrac{\partial Y}{\partial Y^\circ}\dfrac{\partial\beta(\mathbf{u})}{\partial Y} \end{pmatrix} = \begin{pmatrix} \dfrac{\partial X}{\partial X^\circ} & \dfrac{\partial Y}{\partial X^\circ} \\ \dfrac{\partial X}{\partial Y^\circ} & \dfrac{\partial Y}{\partial Y^\circ} \end{pmatrix} \begin{pmatrix} \dfrac{\partial\beta(\mathbf{u})}{\partial X} \\ \dfrac{\partial\beta(\mathbf{u})}{\partial Y} \end{pmatrix}$$
$$= \mathbf{W}^{-1\mathrm{T}}\nabla\beta. \tag{A.17}$$

Substituting equation (A.15) with equation (A.17) into equation (A.14), we get

$$\frac{d\mathbf{M}}{dt} = 2\eta \mathbf{W}^{\mathrm{T}} \left[\frac{\mathbf{W}^{-1\mathrm{T}}\nabla\beta}{|\nabla^\circ\beta|}\right]\left[\frac{\mathbf{W}^{-1\mathrm{T}}\nabla\beta}{|\nabla^\circ\beta|}\right]^{\mathrm{T}} \mathbf{W} + 2\zeta\mathbf{M}$$

604
$$= 2\eta \mathbf{W}^{\mathrm{T}} \left[\frac{\mathbf{W}^{-1\mathrm{T}}\nabla\beta\nabla\beta^{\mathrm{T}}\mathbf{W}^{-1}}{|\nabla^\circ\beta|^2}\right]\mathbf{W} + 2\zeta\mathbf{M} \tag{A.18}$$

$$= 2\eta \frac{\nabla\beta\nabla\beta^{\mathrm{T}}}{|\nabla^\circ\beta|^2} + 2\zeta\mathbf{M}.$$

605 Now we assume that the unidirectional growth rate η is proportional to $|\nabla^\circ\beta|^k$ with k being a

606 positive integer:

607
$$\eta = \frac{1}{2}|\nabla^\circ\beta|^k = \frac{1}{2}|\mathbf{W}^{-1\mathrm{T}}\nabla\beta|^k = \frac{1}{2}[\nabla\beta^{\mathrm{T}}\mathbf{M}^{-1}\nabla\beta]^{k/2} \tag{A.19}$$

608 Substituting equation (A.19) into equation (A.18), we get

609
$$\frac{d\mathbf{M}(\mathbf{u},t)}{dt} = \nabla\beta(\mathbf{u},t)\nabla\beta(\mathbf{u},t)^{\mathrm{T}}[\nabla\beta(\mathbf{u},t)^{\mathrm{T}}\mathbf{M}(\mathbf{u},t)^{-1}\nabla\beta(\mathbf{u},t)]^{[k-2]/2} + 2\zeta(\mathbf{u},t)\mathbf{M}(\mathbf{u},t) \tag{A.20}$$

610 where the arguments "$(\mathbf{u},t)$" are recovered. Finally, assuming $k = 2$ and $\zeta(\mathbf{u},t) = 0$ gives

611 equation (3) in the main text. If $\zeta(\mathbf{u},t)$ and $\beta(\mathbf{u},t)$ are constant along time, the growth metric at

612 time τ is obtained as

613
$$\mathbf{M}(\mathbf{u},\tau) = \int_0^\tau [\nabla\beta\nabla\beta^{\mathrm{T}} + 2\zeta\mathbf{M}(\mathbf{u},t)]\, dt$$
$$= \begin{cases} \mathbf{M}(\mathbf{u},0) + \tau\nabla\beta\nabla\beta^{\mathrm{T}} & \text{for} \quad \zeta = 0 \\ \left[\mathbf{M}(\mathbf{u},0) + \dfrac{\nabla\beta\nabla\beta^{\mathrm{T}}}{2\mu}\right]\exp(2\mu\tau) - \dfrac{\nabla\beta\nabla\beta^{\mathrm{T}}}{2\mu} & \text{for} \quad \zeta \neq 0. \end{cases} \tag{A.21}$$

Appendix B: Isometric embeddability

B.1. Initial growth metric being identity

617 Here we show that the growth metric given by equation (6) in the main text,

618
$$\mathbf{M}(\mathbf{u},\tau) = \mathbf{M}(\mathbf{u},0) + \hat{\rho}^2 \nabla\beta(\mathbf{u})\nabla\beta(\mathbf{u})^{\mathrm{T}} \tag{B.1}$$

619 with $\mathbf{M}(\mathbf{u},0) = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, is identical to the growth metric calculated from the sheet shape given

620 by equation (7),

621
$$\tilde{\mathbf{r}}_\beta(\mathbf{u}) = \begin{pmatrix} X \\ Y \\ \hat{\rho}\beta(\mathbf{u}) \end{pmatrix}. \tag{B.2}$$

622 We confirm this relationship by deriving that the growth amount $L_\beta(\mathbf{u},\mathbf{e}) = \sqrt{\mathbf{e}^{\mathrm{T}}\mathbf{M}_\beta(\mathbf{u},\tau)\mathbf{e}}$

623 induced from equation (B.2) is identical to the growth amount $L(\mathbf{u},\mathbf{e},\tau) = \sqrt{\mathbf{e}^{\mathrm{T}}\mathbf{M}(\mathbf{u},\tau)\mathbf{e}}$ given

624 by equation (B.1) for any $\mathbf{e}$. Specifically, we see for $\boldsymbol{\delta u} = (\delta X, \delta Y)^{\mathrm{T}}$, $\mathbf{e} = (e_X, e_Y)^{\mathrm{T}} = \boldsymbol{\delta u}\,/$

625    $|\boldsymbol{\delta u}|$, and $\varepsilon = |\boldsymbol{\delta u}|$ that

$$
\begin{aligned}
L_\beta(\mathbf{u}, \mathbf{e})^2 &= \lim_{|\boldsymbol{\delta u}|\to 0} \frac{\left|\tilde{\mathbf{r}}_\beta(\mathbf{u}+\boldsymbol{\delta u}) - \tilde{\mathbf{r}}_\beta(\mathbf{u})\right|^2}{|\boldsymbol{\delta u}|^2} \\
&= \lim_{|\boldsymbol{\delta u}|\to 0} \frac{1}{|\boldsymbol{\delta u}|^2}\left|\begin{pmatrix} X+\delta X \\ Y+\delta Y \\ \hat{\rho}\beta(\mathbf{u}+\boldsymbol{\delta u}) \end{pmatrix} - \begin{pmatrix} X \\ Y \\ \hat{\rho}\beta(\mathbf{u}) \end{pmatrix}\right|^2 \\
&= \lim_{\varepsilon\to 0} \frac{1}{\varepsilon^2}\left|\begin{pmatrix} \varepsilon e_X \\ \varepsilon e_Y \\ \hat{\rho}\left[\dfrac{\partial\beta(\mathbf{u})}{\partial X}\varepsilon e_X + \dfrac{\partial\beta(\mathbf{u})}{\partial Y}\varepsilon e_Y + O(\varepsilon^2)\right] \end{pmatrix}\right|^2 \\
&= e_X^2 + e_Y^2 + \hat{\rho}^2\left[\frac{\partial\beta(\mathbf{u})}{\partial X}e_X + \frac{\partial\beta(\mathbf{u})}{\partial Y}e_Y\right]^2 \\
&= \begin{pmatrix} e_X \\ e_Y \end{pmatrix}^{\mathrm{T}}\left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \hat{\rho}^2\begin{pmatrix} \dfrac{\partial\beta(\mathbf{u})}{\partial X} \\ \dfrac{\partial\beta(\mathbf{u})}{\partial Y} \end{pmatrix}\begin{pmatrix} \dfrac{\partial\beta(\mathbf{u})}{\partial X} \\ \dfrac{\partial\beta(\mathbf{u})}{\partial Y} \end{pmatrix}^{\mathrm{T}}\right]\begin{pmatrix} e_X \\ e_Y \end{pmatrix} \\
&= \mathbf{e}^{\mathrm{T}}\left[\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \hat{\rho}^2\nabla\beta(\mathbf{u})\nabla\beta(\mathbf{u})^{\mathrm{T}}\right]\mathbf{e} = \mathbf{e}^{\mathrm{T}}\mathbf{M}_\beta(\mathbf{u})\mathbf{e},
\end{aligned}
\tag{B.3}
$$

and which is identical to $L(\mathbf{u}, \mathbf{e}, \tau)^2 = \mathbf{e}^{\mathrm{T}}\mathbf{M}(\mathbf{u}, \tau)\mathbf{e}$ given by equation (B.1).

Therefore, as long as the basic shape for $\beta(\mathbf{u})$ has a simple horn-like shape, we expect that
the extended sheet calculated numerically using the finite element or vertex-spring methods,
denoted by $\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u}) = \tilde{\mathbf{r}}(\mathbf{u}, t_{\mathrm{end}})$, will have a shape similar to $\tilde{\mathbf{r}}_\beta(\mathbf{u})$, so that the realised growth
amount $L_{\mathrm{real}}(\mathbf{u}, \mathbf{e}; \tilde{\mathbf{r}}_{\mathrm{ext}})$ calculated from $\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u})$ is similar to $L_\beta(\mathbf{u}, \mathbf{e})$. This means that the
realised growth, denoted by the extension metric $\mathbf{M}_{\mathrm{ext}}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}})$, is similar to $\mathbf{M}_\beta(\mathbf{u})$, where
$L_{\mathrm{ext}}(\mathbf{u}, \mathbf{e}; \tilde{\mathbf{r}}_{\mathrm{ext}})$ and $\mathbf{M}_{\mathrm{ext}}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}})$ are defined as

$$
\begin{aligned}
L_{\mathrm{ext}}(\mathbf{u}, \mathbf{e}; \tilde{\mathbf{r}}_{\mathrm{ext}})^2 &= \lim_{|\boldsymbol{\delta u}|\to 0} \frac{|\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u}+\boldsymbol{\delta u}) - \tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u})|}{|\boldsymbol{\delta u}|^2} = \mathbf{e}^{\mathrm{T}}\mathbf{M}_{\mathrm{ext}}(\mathbf{u})\mathbf{e}, \\
\mathbf{M}_{\mathrm{ext}}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) &= \begin{pmatrix} M_{\mathrm{ext},XX}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) & M_{\mathrm{ext},XY}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) \\ M_{\mathrm{ext},XY}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) & M_{\mathrm{ext},YY}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) \end{pmatrix}, \\
M_{\mathrm{ext},XX}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) &= \left|\frac{\partial\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u})}{\partial X}\right|^2, \\
M_{\mathrm{ext},YY}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) &= \left|\frac{\partial\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u})}{\partial Y}\right|^2, \\
M_{\mathrm{ext},XY}(\mathbf{u}; \tilde{\mathbf{r}}_{\mathrm{ext}}) &= \left[\frac{\partial\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u}, t)}{\partial X}\right]^{\mathrm{T}}\left[\frac{\partial\tilde{\mathbf{r}}_{\mathrm{ext}}(\mathbf{u}, t)}{\partial Y}\right].
\end{aligned}
\tag{B.4}
$$

B.2. Initial growth metric not being identity

Next, we consider the case that the initial growth metric $\mathbf{M}(\mathbf{u}, 0)$ is not identity but constant over
the sheet. Such a case arises when we fuse two basic shapes corresponding to $\beta_1(\mathbf{u})$ with a wide

basic shape and $\beta_2(\mathbf{u})$ with a very narrow basic shape located at the middle of the slope of the

basic shape of $\beta_1(\mathbf{u})$, as shown in figure 3. In this case, after the sheet growth induced by $\beta_1(\mathbf{u})$,

the sheet around the basic shape of $\beta_2(\mathbf{u})$, denoted here by $\beta(\mathbf{u})$, has approximately a constant

growth metric, denoted here by $\mathbf{M}(\mathbf{u}, 0)$.

An appropriate rotation of the material coordinates makes $\mathbf{M}(\mathbf{u}, 0)$ diagonal with positive

constant entries, which is expressed as

$$\mathbf{M}(\mathbf{u}, 0) = \begin{pmatrix} p^2 & 0 \\ 0 & q^2 \end{pmatrix}. \tag{B.5}$$

645 Then, as derived below, the growth metric $\mathbf{M}(\mathbf{u}, \tau) = \mathbf{M}(\mathbf{u}, 0) + \hat{\rho}^2 \nabla\beta(\mathbf{u})\nabla\beta(\mathbf{u})^{\mathrm{T}}$ of the grown

646 sheet has a smooth isometric embedding in the spatial coordinates given by

$$\tilde{\mathbf{r}}_\beta(\mathbf{u}) = \begin{pmatrix} pX \\ qY \\ \hat{\rho}\beta(\mathbf{u}) \end{pmatrix}. \tag{B.6}$$

648 We confirm this in a manner analogous to the case of the initial growth metric being identity

649 (Appendix B.1). Specifically, we see for $\boldsymbol{\delta u} = (\delta X, \delta Y)^{\mathrm{T}}$, $\mathbf{e} = (e_X, e_Y)^{\mathrm{T}} = \boldsymbol{\delta u}\,/\,|\boldsymbol{\delta u}|$, and $\varepsilon =$

650 $|\boldsymbol{\delta u}|$ that

$$
\begin{aligned}
L_\beta(\mathbf{u}, \mathbf{e})^2 &= \lim_{|\boldsymbol{\delta u}|\to 0} \frac{|\tilde{\mathbf{r}}_\beta(\mathbf{u} + \boldsymbol{\delta u}) - \tilde{\mathbf{r}}_\beta(\mathbf{u})|}{|\boldsymbol{\delta u}|^2} \\
&= \lim_{|\boldsymbol{\delta u}|\to 0} \frac{1}{|\boldsymbol{\delta u}|^2} \left| \begin{pmatrix} p[X + \delta X] \\ q[Y + \delta Y] \\ \hat{\rho}\beta(\mathbf{u} + \boldsymbol{\delta u}) \end{pmatrix} - \begin{pmatrix} pX \\ qY \\ \hat{\rho}\beta(\mathbf{u}) \end{pmatrix} \right|^2 \\
&= \lim_{\varepsilon\to 0} \frac{1}{\varepsilon^2} \left| \begin{pmatrix} p\varepsilon e_X \\ q\varepsilon e_Y \\ \hat{\rho}\left[\dfrac{\partial\beta(\mathbf{u})}{\partial X}\varepsilon e_X + \dfrac{\partial\beta(\mathbf{u})}{\partial Y}\varepsilon e_Y + O(\varepsilon^2)\right] \end{pmatrix} \right|^2 \\
&= p^2 e_X^2 + q^2 e_Y^2 + \hat{\rho}^2 \left[\frac{\partial\beta(\mathbf{u})}{\partial X} e_X + \frac{\partial\beta(\mathbf{u})}{\partial Y} e_Y\right]^2 \\
&= \begin{pmatrix} e_X \\ e_Y \end{pmatrix}^{\mathrm{T}} \left[\begin{pmatrix} p^2 & 0 \\ 0 & q^2 \end{pmatrix} + \hat{\rho}^2 \begin{pmatrix} \dfrac{\partial\beta(\mathbf{u})}{\partial X} \\ \dfrac{\partial\beta(\mathbf{u})}{\partial X} \end{pmatrix} \begin{pmatrix} \dfrac{\partial\beta(\mathbf{u})}{\partial X} \\ \dfrac{\partial\beta(\mathbf{u})}{\partial X} \end{pmatrix}^{\mathrm{T}} \right] \begin{pmatrix} e_X \\ e_Y \end{pmatrix} \\
&= \mathbf{e}^{\mathrm{T}}[\mathbf{M}(\mathbf{u}, 0) + \hat{\rho}^2 \nabla\beta(\mathbf{u})\nabla\beta(\mathbf{u})^{\mathrm{T}}]\mathbf{e} = \mathbf{e}^{\mathrm{T}}\mathbf{M}_\beta(\mathbf{u})\mathbf{e},
\end{aligned}
\tag{B.7}
$$

652 And which is identical to $L(\mathbf{u}, \mathbf{e}, \tau)^2 = \mathbf{e}^{\mathrm{T}}\mathbf{M}(\mathbf{u}, \tau)\mathbf{e}$.

Appendix C: Simulation method

C.1. Finite element method

In the finite element method used in the present study, the epidermal sheet is divided into first-order hexagonal finite elements, and the spatial coordinates of vertices of these elements are all collected into a single vector $\mathbf{p}$. The motion equation for $\mathbf{p}$ is written as

$$\mathbf{Q}\frac{d^2\mathbf{p}}{dt^2} = \mathbf{f}_{\text{NH}}(\mathbf{p}) + \mathbf{S}\frac{d\mathbf{p}}{dt}, \tag{C.1}$$

where $\mathbf{Q}$ and $\mathbf{S}$ respectively describe the mass and viscosity matrices. $\mathbf{f}_{\text{NH}}(\mathbf{p})$ is a vector describing the stress due to strain from the unstrained shape of each element, under the neo-Hookean hyper-elastic material model. This equation of motion is time-integrated with singly diagonally implicit Runge-Kutta (SDIRK) schemes, by using a C++ library MFEM-3.3.2.

In the initial state of the calculation, these finite elements were generated by equally dividing a square epidermal sheet along the x- and y-axes, where the sheet contains a single layer of elements along the z-axis. The time for the initial state is set at time $t = \tau$ (i.e., the time when the sheet growth ends), and the initial state for $\mathbf{p}$ corresponds to the $\tilde{\mathbf{r}}(\mathbf{u}, \tau) = \left(X, Y, Z_{\text{init}}(X, Y)\right)^{\text{T}}$ given by the basic shape for a morphogen distribution (or a linear combination of the basic shapes of morphogen distributions) with some scaling. A slight velocity in the positive z-direction was initially set to control the direction of sheet inflation in the positive z-direction. For each rectangular finite element, the unstrained lengths for the sides of the rectangles on the upper and lower surfaces are elongated according to the metric $\mathbf{M}_{\text{sim}}(\mathbf{u}, t)$. $\mathbf{M}_{\text{sim}}(\mathbf{u}, t)$ starts at time $t = \tau$ from its initial state $\mathbf{M}_{\text{init}}(\mathbf{u})$ calculated from $\tilde{\mathbf{r}}(\mathbf{u}, \tau) = \left(X, Y, Z_{\text{init}}(X, Y)\right)^{\text{T}}$ and then monotonically approaches the growth metric $\mathbf{M}(\mathbf{u})$ given by equations (6) or (9). After reaching $\mathbf{M}(\mathbf{u})$ at $t = \tau + t_{\text{ext}}$, $\mathbf{M}_{\text{sim}}(\mathbf{u}, t)$ is kept equal to $\mathbf{M}(\mathbf{u})$ until the calculation ends at $t = \tau + t_{\text{end}}$. Specifically, $\mathbf{M}_{\text{init}}(\mathbf{u})$ is defined as

$$\mathbf{M}_{\text{sim}}(\mathbf{u}, t) = \begin{cases} \mathbf{M}_{\text{init}}(\mathbf{u})\dfrac{t_{\text{ext}} - [t - \tau]}{t_{\text{ext}}} + \dfrac{t - \tau}{t_{\text{ext}}}\mathbf{M}(\mathbf{u}) & \text{for} \quad 0 \le t - \tau \le t_{\text{ext}} \\ \mathbf{M}(\mathbf{u}) & \text{for} \quad t_{\text{ext}} \le t - \tau \le t_{\text{end}}. \end{cases} \tag{C.2}$$

From $\mathbf{M}_{\text{sim}}(\mathbf{u}, t)$ the unstrained side length of an element connecting vertices i and j is given by

$$l(\mathbf{u}_i, \mathbf{u}_j, t) = \sqrt{[\mathbf{u}_i - \mathbf{u}_j]^{\text{T}}\mathbf{M}_{\text{sim}}\left(\frac{\mathbf{u}_i + \mathbf{u}_j}{2}, t\right)[\mathbf{u}_i - \mathbf{u}_j]} \tag{C.3}$$

[Parameters]

Initial sheet size: side length 8.0 along the x- and y-axes and thickness 0.2 along the z-axis

Time step: 0.5, Viscosity: 0.1

Bulk modulus: 5.0, Shear modulus: 0.05 (in Neo-Hookean hyperelastic model)

Initial velocity: $0.0001(x - 8)x(y - 8)y$ (only in the z-direction)

Fig. 2:

$\beta(X, Y) = 21N(X, Y; 1.6, 1.6)$, $n_{\text{element}} = 80$ (i.e., Number of elements along the X- or Y-axis,

giving $n_{\text{element}} \times n_{\text{element}} = 6400$ elements in total), $Z_{\text{init}}(X, Y) = 0.5\beta(X, Y)$.

For panels (d,g): $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0,0,0)$, $(t_{\text{ext}}, t_{\text{end}}) = (20, 50)$.

For panels (e,h): $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.24,0,0)$, $(t_{\text{ext}}, t_{\text{end}}) = (40, 100)$.

For panels (f,i): $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.24,0,0.05)$, $(t_{\text{ext}}, t_{\text{end}}) = (40, 100)$.

Fig. 3:

$\beta_1(X, Y) = 49N(X, Y; 1.6, 1.6)$, $\beta_2(X, Y) = 49N(X - 1.8, Y; 0.5, 0.5)$, $n_{\text{element}} = 80$,

$Z_{\text{init}}(X, Y) = 0.5\beta(X, Y)$, $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0,0,0)$, $(t_{\text{ext}}, t_{\text{end}}) = (20, 50)$.

For panels (e,h): $\psi_{12} = 1.0$.

For panels (f,i): $\psi_{12} = 0.5$.

For panels (g,j): $\psi_{12} = 0.0$

Fig. 4:

$\beta_1(X, Y) = 49N(X, Y; 1.7, 1.0)$, $\beta_2(X, Y) = 49 \cdot 2.8N(X - 0.45, Y; 1.7, 0.7)$, $\beta_3(X, Y) = 49 \cdot$

$0.98[\psi_{23} + 1]N(X - 2.6 + 0.6\psi_{23}, Y + 2.1 + 0.84\psi_{23}; 1.0, 0.5)$, $n_{\text{element}} = 80$, $Z_{\text{init}}(X, Y) =$

$\beta_1(X, Y) + 0.1\beta_2(X, Y) + 0.1\beta_3(X, Y)$, $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.16,0,0.01)$, $(t_{\text{ext}}, t_{\text{end}}) = (40, 100)$.

For panels (d,g): $\psi_{23} = 0.0$

For panels (e,h): $\psi_{23} = 0.5$

For panels (f,i): $\psi_{23} = 1.0$

Fig. 5a,d,g:

$\beta_1(X, Y) = 400N(X, Y; 1.7, 1.0)$, $\beta_2(X, Y) = 400 \cdot 1.8N(X - 1.6, Y - 0.36; 0.28, 0.32)$, $\psi_{12} =$

0.0 , $n_{\text{element}} = 128$, $Z_{\text{init}}(X, Y) = 0.6\beta_1(X, Y) + 0.4\beta_2(X, Y)$, $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.6,0,0.005)$,

$(t_{\text{ext}}, t_{\text{end}}) = (60, 200)$.

Fig. 5b,e,h:

$\beta_1(X, Y) = 400N(X, Y; 1.7, 1.1)$, $\beta_2(X, Y) = 400 \cdot 1.5\sum_{i=1}^{3} N\left(X + \frac{1.9}{3}i - 2.9, Y; 0.24 + \right.$

$0.02i, 0.4 + 0.025i\Big)$, $\psi_{12} = 0.0$, $n_{\text{element}} = 128$, $Z_{\text{init}}(X,Y) = \beta_1(X,Y) + 0.1\beta_2(X,Y)$,

$(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.2, 0, 0.01)$, $(t_{\text{ext}}, t_{\text{end}}) = (40, 100)$.

Fig. 5c,f,i:

$\beta_1(X,Y) = 196N(X,Y; 2.4, 2.4)$, $\beta_2(X,Y) = 196[0.6N(X, Y - 1.1; 0.7, 0.4) + 0.6N(X, Y +$

$1.1; 0.7, 0.4) + 0.18N(X + 2.2, Y - 1.1; 0.49, 0.28) + 0.18N(X + 2.2, Y + 1.1; 0.49, 0.28)]$,

$\beta_3(X,Y) = 196[0.24N(X + 0.3, Y - 1.54; 0.42, 0.24) + 0.24N(X + 0.3, Y +$

$1.54; 0.42, 0.24)]$, $\psi_{12} = 0.0$, $n_{\text{element}} = 80$, $Z_{\text{init}}(X,Y) = 0.4\beta_1(X,Y) + 0.01\beta_2(X,Y) +$

$0.01\beta_3(X,Y)$, $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.16, 0, 0.01)$, $(t_{\text{ext}}, t_{\text{end}}) = (40, 100)$.

C.2. Vertices-spring method

[Main equation]

We assume that the epithelial sheet has no thickness and initially has a square shape with its sides are parallel with the X- or Y-axis in the material coordinates. We discretize the sheet into equally spaced vertices, denoted by (i, j) with $i, j = 1, \dots, N$, connected by springs. The spatial coordinates of vertex (i, j) are given by $\tilde{\mathbf{r}}_{(i,j)} = \tilde{\mathbf{r}}(\mathbf{u}_{(i,j)}, t)$. We describe the force acting on vertex (i, j) as

$$\mathbf{f}_{(i,j)} = \mathbf{k}_{(i,j)} + \mathbf{b}_{(i,j)} + \mathbf{w}_{(i,j)} + \mathbf{o}_{(i,j)}, \tag{C.4}$$

where $\mathbf{k}_{(i,j)}$, $\mathbf{b}_{(i,j)}$, $\mathbf{w}_{(i,j)}$, and $\mathbf{o}_{(i,j)}$ are forces due to springs (i.e., stretching elasticity), bending elasticity, the water pressure (from inside of the sheet), and outer force, respectively. We assume that the system is kept almost valanced so that $|\mathbf{f}_{(i,j)}|$ is kept small at each time step, so that we can calculate the time evolution of the coordinates of vertex (i, j) by

$$\frac{d\tilde{\mathbf{r}}_{(i,j)}}{dt} = \mathbf{f}_{(i,j)}. \tag{C.5}$$

[Spring force, water pressure, and bending elasticity]

The spring force on vertex (i, j) is described as

$$\mathbf{k}_{(i,j)} = \sum_{(i',j') \in \text{con}(i,j)} c_k \left[\frac{\left| \tilde{\mathbf{r}}_{(i',j')} - \tilde{\mathbf{r}}_{(i,j)} \right| - l_{(i',j'):(i,j)}}{l_{(i',j'):(i,j)}} \right] \frac{\tilde{\mathbf{r}}_{(i',j')} - \tilde{\mathbf{r}}_{(i,j)}}{\left| \tilde{\mathbf{r}}_{(i',j')} - \tilde{\mathbf{r}}_{(i,j)} \right|} \tag{C.6}$$

with a positive constant parameter c_k describing the spring constant per unit length, and where $l_{(i',j'):(i,j)}$ describes the spring length connecting vertices (i, j) and (i', j') chosen from the eight connected vertices,

$$\begin{aligned} \text{con}(i, j) = &\{(i - 1, j), (i + 1, j), (i, j - 1), (i, j + 1)\} \\ &\cup \{(i - 1, j - 1), (i + 1, j - 1), (i - 1, j + 1), (i + 1, j + 1)\}. \end{aligned} \tag{C.7}$$

745    $l_{(i',j'):(i,j)}$ at time t is given by $l\left(\mathbf{u}_{(i,j)}, \mathbf{u}_{(i',j')}, t\right)$ in equation (D.3).

746   The bending-elastic force $\mathbf{b}_{(i,j)}$ on the vertex (i,j) is described as the sum of the bending-

747   elastic force $\mathbf{b}^X_{(i,j)}$ along the X-axis: $\{(i-1,j),(i,j),(i+1,j)\}$ and the bending-elastic force

748    $\mathbf{b}^Y_{(i,j)}$ along the Y-axis: $\{(i,j-1),(i,j),(i,j+1)\}$. Specifically, we describe $\mathbf{b}_{(i,j)}$ as:

749
$$\mathbf{b}_{(i,j)} = \mathbf{b}^X_{(i,j)} + \mathbf{b}^Y_{(i,j)}. \tag{C.8}$$

750   We denote by $\theta^X_{(i,j)}$ the bending angle of the sheet at vertex (i,j) along X, as illustrated in figure

751   9a. The $\theta^X_{(i,j)}$ is the angle between the segment $(i-1,j):(i,j)$ and the segment $(i,j):(i+1,j)$,

752   given by the inner product of the corresponding two unit vectors:

753
$$\begin{aligned}
\cos\theta_{(i,j)} &= \left(\mathbf{e}^{x-}_{(i,j)}, \mathbf{e}^{x+}_{(i,j)}\right), \\
\mathbf{e}^{x-}_{(i,j)} &= \frac{\tilde{\mathbf{r}}_{(i,j)} - \tilde{\mathbf{r}}_{(i-1,j)}}{\left|\tilde{\mathbf{r}}_{(i,j)} - \tilde{\mathbf{r}}_{(i-1,j)}\right|}, \\
\mathbf{e}^{x+}_{(i,j)} &= \frac{\tilde{\mathbf{r}}_{(i+1,j)} - \tilde{\mathbf{r}}_{(i,j)}}{\left|\tilde{\mathbf{r}}_{(i+1,j)} - \tilde{\mathbf{r}}_{(i,j)}\right|}.
\end{aligned} \tag{C.9}$$

The $\theta^X_{(i,j)}$ produces the forces $\mathbf{m}^{X-}_{(i,j)}$ and $\mathbf{m}^{X+}_{(i,j)}$ on vertices $(i-1,j)$ and $(i+1,j)$, respectively.

At the same time, due to the law of action and reaction, vertex (i,j) undergoes $-\left[\mathbf{m}^{X-}_{(i,j)} + \mathbf{m}^{X+}_{(i,j)}\right]$.

We assume that the bending elasticity caused by the bending angle $\theta_{(i,j)}$ is proportional to the

inverse of the curvature radius $R_{(i,j)} = a/\tan\frac{\theta_{(i,j)}}{2}$ (illustrated in figure 9b, where the

corresponding circle is tangent to line segments $(i-1,j):(i,j)$ and $(i,j):(i+1,j)$ respectively

at points $\mathbf{a}^-_{(i,j)}$ and $\mathbf{a}^+_{(i,j)}$ both of which have the constant distance a from $\tilde{\mathbf{r}}_{(i,j)}$). To keep $R^{-1}_{(i,j)} = $

$a^{-1}\tan\frac{\theta_{(i,j)}}{2} = a^{-1}\sqrt{\frac{1-\cos^2\theta_{(i,j)}}{1+\cos\theta_{(i,j)}}}$ from being infinite, we introduce a constant saturation parameter

$c_{bs} > 0$ and describe the torque as

762
$$\varsigma(\cos\theta_{(i,j)}) = c_b \frac{\sqrt{1-\cos^2\theta_{(i,j)}}}{1+\cos\theta_{(i,j)}+c_{bs}} \tag{C.10}$$

763   where a^{-1} is included in a positive constant parameter c_b. Then $\mathbf{m}^{X-}_{(i,j)}$ and $\mathbf{m}^{X+}_{(i,j)}$ are calculated

764   as

765
$$\begin{aligned}
\mathbf{m}^{X-}_{(i,j)} &= \frac{\varsigma\left(\cos\theta_{(i,j)}\right)}{l_{(i-1,j):(i,j)}}\mathbf{n}^{x-}, \\
\mathbf{m}^{X+}_{(i,j)} &= \frac{\varsigma\left(\cos\theta_{(i,j)}\right)}{l_{(i+1,j):(i,j)}}\mathbf{n}^{x+},
\end{aligned} \tag{C.11}$$

766   where $\mathbf{n}^{x-}_{(i,j)}$ and $\mathbf{n}^{x+}_{(i,j)}$ are unit vectors included in the plane that contains the three vertices

767   ($\tilde{\mathbf{r}}_{(i-1,j)}$, $\tilde{\mathbf{r}}_{(i,j)}$, and $\tilde{\mathbf{r}}_{(i+1,j)}$) and are orthogonal to $\mathbf{e}^{x-}_{(i,j)}$ and $\mathbf{e}^{x+}_{(i,j)}$, respectively. Specifically,

768   $\mathbf{n}_{(i,j)}^{x-}$ and $\mathbf{n}_{(i,j)}^{x+}$ are given by

$$\mathbf{n}_{(i,j)}^{x-} = \frac{\mathbf{v}_{\mathbf{n}(i,j)}^{x-}}{\left|\mathbf{v}_{\mathbf{n}(i,j)}^{x-}\right|},$$

$$\mathbf{n}_{(i,j)}^{x+} = \frac{\mathbf{v}_{\mathbf{n}(i,j)}^{x+}}{\left|\mathbf{v}_{\mathbf{n}(i,j)}^{x+}\right|}, \tag{C.12}$$

$$\mathbf{v}_{\mathbf{n}(i,j)}^{x-} = -\mathbf{e}_{(i,j)}^{x+} - \left(-\mathbf{e}_{(i,j)}^{x+}, \mathbf{e}_{(i,j)}^{x-}\right)\mathbf{e}_{(i,j)}^{x-},$$

$$\mathbf{v}_{\mathbf{n}(i,j)}^{x+} = \mathbf{e}_{(i,j)}^{x-} - \left(\mathbf{e}_{(i,j)}^{x-}, \mathbf{e}_{(i,j)}^{x+}\right)\mathbf{e}_{(i,j)}^{x+}.$$

770   On this basis, we obtain the bending-elastic force $\mathbf{b}_{(i,j)}^{X}$ on vertex (i,j) along the X-axis as the

771   sum of $-\left[\mathbf{m}_{(i,j)}^{X-} + \mathbf{m}_{(i,j)}^{X+}\right]$ from the bending at vertex (i,j), $\mathbf{m}_{(i-1,j)}^{X+}$ from $(i-1,j)$, and $\mathbf{m}_{(i+1,j)}^{X-}$

772   from $(i+1,j)$:

773
$$\mathbf{b}_{(i,j)}^{X} = -\left[\mathbf{m}_{(i,j)}^{X-} + \mathbf{m}_{(i,j)}^{X+}\right] + \mathbf{m}_{(i-1,j)}^{X+} + \mathbf{m}_{(i+1,j)}^{X-}. \tag{C.13}$$

774   Analogously, we obtain the bending-elastic force $\mathbf{b}_{(i,j)}^{Y}$ on vertex (i,j) along the Y-axis from

775   equations (C.9-C.13) with replacement of the superscript "X" with "Y." Then, the total bending

776   elastic force $\mathbf{b}_{(i,j)}$ on vertex (i,j) is calculated by equation (C. 8).

777   The force exerted by water pressure on the vertex (i,j) is described as

778
$$\mathbf{W}_{(i,j)} = c_w \mathbf{n}_{(i,j)} \tag{C.14}$$

with a positive constant parameter c_w, where $\mathbf{n}_{(i,j)}$ is the normal vector of the sheet at $\tilde{\mathbf{r}}_{(i,j)}$, given

by the average of the four normal vectors of the four triangles sharing vertex (i,j) (i.e.,

$\{(i,j), (i+1,j), (i,j+1)\}$, $\{(i,j), (i,j+1), (i-1,j)\}$, $\{(i,j), (i-1,j), (i-1,j-1)\}$, and

$\{(i,j), (i-1,j), (i+1,j-1)\}$.

Finally, the upward outer force on the vertices (i,j) (acting only on the sheet edges) could be

described as follows:

785
$$\mathbf{o}_{(i,j)} = \begin{cases} c_{\text{outer}} \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} & \text{for} \quad i = 1 \text{ or } i = N \text{ or } j = 1 \text{ or } j = N \\ \mathbf{0} & \text{for} \qquad\qquad \text{otherwise} \end{cases} \tag{C.15}$$

with a positive constant parameter c_{outer}.

[Parameters]

Sheet size: 4.0 (along X-axis), 4.0 (Y-axis), 0.0 (Z-axis)

Number of vertices: 129×129 (along X- and Y-axes)

Time step: 8.0×10^{-4} (Explicit Euler method)

Elasticity parameters: $c_k = 10$, $c_b = 0.23$, $c_{bs} = 0.0001$, $c_{\text{outer}} = 40$

Morphogen distribution: $\beta(\mathbf{u}, t) = \sum_{i=1}^{5} \rho_i(t)\beta_i(\mathbf{u})$ with $\psi_{ij} = \int \rho_i(t)\rho_j(t)dt = 0$ for all $i \neq j$

and $Z_{\text{init}}(\mathbf{u}) = \sum_{i=1}^{5} \beta_i(\mathbf{u})$

Function for linear combination of similar Normal distributions:

$$F\big(X,Y;a_0,a_1;X_0,X_1,Y_0,Y_1;\sigma_{x0},\sigma_{x1},\sigma_{y0},\sigma_{y1},K\big) = \sum_{k=1}^{K} a_k N\big(X-X_k,Y-Y_k,\sigma_{x\,k},\sigma_{y\,k}\big)$$

with $\sigma_{x0} = \sigma_{y0} = 0.5, X_0 = 0.2, Y_0 = -1.0,$

$a_k = \frac{k}{K}[a_1 - a_0] + a_0, X_k = \frac{k}{K}[X_1 - X_0] + X_0, Y_k = \frac{k}{K}[Y_1 - Y_0] + Y_0,$

$\sigma_{xk} = \frac{k}{K}[\sigma_{x1} - \sigma_{x0}] + \sigma_{x0}, \sigma_{yk} = \frac{k}{K}[\sigma_{y1} - \sigma_{y0}] + \sigma_{y0},$

Fig. 6:

$\beta_1(\mathbf{u}) = 0.48N(X,Y-1.0;1.3,0.5) + 0.48N(X,Y-0.3;0.7,1.0) + 4.5N(X,Y+1.7;1.5,1.0),$

$\beta_2(\mathbf{u}) = 0.8N(X,Y-1.0;0.4,0.4) + 2N(X,Y+1.7;1.0,1.0)$

$\qquad +0.8N(X-1.5,Y+1.7;1,1) + 0.8N(X+1.5,Y+1.7;1,1)$

$\qquad +2.8F(X,Y;0.3,0.2;1.5X_0,1.5X_0,0.9Y_0,0.9Y_0;1.4\sigma_{x0},0.07,1.4\sigma_{y0},0.07,20),$

$\beta_3(\mathbf{u}) = 0,$

$\beta_4(\mathbf{u}) = 0,$

$\beta_5(\mathbf{u}) = 2.52F(X,Y;0.3,0.2;0,0,0.6Y\_0,0.6Y\_0;0.6\sigma_{x0},0.07,0.8\sigma_{y0},0.07,20),$

$(\gamma_{X1},\gamma_{Y1},\omega_1) = (0,-0.06,0), (\gamma_{X2},\gamma_{Y2},\omega_2) = (\gamma_{X5},\gamma_{X5},\omega_5) = (0,0.15,0),$

$\big(c_{w,\text{ext}},c_{w,\text{end}}\big) = (2,0.2), (t_{\text{ext}},t_{\text{end}}) = (60000,480000).$

Fig. 7:

$\beta_1(\mathbf{u}) = 0.48N(X,Y-1.0;1.3,0.5) + 0.48N(X,Y-0.3;0.7,1.0) + 4.5N(X,Y+1.7;1.0,1.0)$

$\qquad +0.5N(X-1.4,Y-1.2;0.7,0.7) + 1.5N(X-1.4,Y-1.2;0.4,0.4)$

$\qquad +0.5N(X+1.4,Y-1.2;0.7,0.7) + 1.5N(X+1.4,Y-1.2;0.4,0.4)$

$\qquad +0.3F(X,Y;0.3,0.2;0,0,-0.5,-0.5;0.5\,\sigma_{x0},0.07,0.5\,\sigma_{y0},0.07,20),$

$\beta_2(\mathbf{u}) = 0.8N(X,Y-1.0;0.4,0.4) + 2N(X,Y+1.7;1.0,1.0) + b_2(X,Y;1) + b_2(X,Y;-1)$,

$\quad b_2(X,Y;s) = 0.8N(X+1.5s,Y+1.7;1,1)$

$\qquad +0.3F\big(X,Y;0.3,0.2;0.7sX_0,0.7sX_0,Y_0,Y_0;\sigma_{x0},\sigma_{x0},\sigma_{y0},\sigma_{y0},20\big)$

$\qquad +0.3F\big(X,Y;0.3,0.2;3sX_0,3sX_0,0.9Y_0,0.9Y_0;\sigma_{x0},0.1,\sigma_{y0},0.1,20\big)$

$\qquad +0.7F\big(X,Y;0.3,0.2;6sX_0,6sX_0,0.8Y_0,0.8Y_0;\sigma_{x0},0.07,\sigma_{y0},0.07,20\big)$

$\qquad +0.4F\big(X,Y;0.3,0.2;6sX_0,8sX_0,0.7Y_0,0.7Y_0;\sigma_{x0},0.07,\sigma_{y0},0.07,20\big),$

$\beta_3(\mathbf{u}) = 3.0F(X,Y;0.3,0.2;-6X_0,-6X_0,0.8Y_0,0.8Y_0;0.3,0.07,0.3,0.07,20),$

$\beta_4(\mathbf{u}) = 3.0F(X,Y;0.3,0.2;6X_0,6X_0,0.8Y_0,0.8Y_0;0.3,0.07,0.3,0.07,20),$

$\beta_5(\mathbf{u}) = F\big(X,Y;0.3,0.2;0,0,0.6Y_0,0.6Y_0;0.7\sigma_{x0},0.07,0.8\sigma_{y0},0.07,20\big),$

$(\gamma_{X1},\gamma_{Y1},\omega_1) = (0,-0.06,0), (\gamma_{X2},\gamma_{Y2},\omega_2) = (0,0.05,0), (\gamma_{X3},\gamma_{Y3},\omega_3) = (0.15,0,-0.015),$

$(\gamma_{X4},\gamma_{Y4},\omega_4) = (-0.15,0,0.015), (\gamma_{X5},\gamma_{Y5},\omega_5) = (0,0.09,0),$

$\left(c_{w,\text{ext}}, c_{w,\text{end}}\right) = (3.5, 0.03), (t_{\text{ext}}, t_{\text{end}}) = (120000, 480000).$

Fig. 8:

$\beta_1(\mathbf{u}) = 0.48N(X, Y - 1.0; 1.3, 0.5) + 0.48N(X, Y - 0.3; 0.7, 1.0) + 4.5N(X, Y + 1.7; 1.0, 1.0)$

$+0.5N(X - 1.4, Y - 1.2; 0.7, 0.7) + 1.5N(X - 1.4, Y - 1.2; 0.4, 0.4)$

$+0.5N(X + 1.4, Y - 1.2; 0.7, 0.7) + 1.5N(X + 1.4, Y - 1.2; 0.4, 0.4)$

$\beta_2(\mathbf{u}) = 0.8N(X, Y - 1.0; 0.4, 0.4) + 2N(X, Y + 1.7; 1.0, 1.0) + b_2(X, Y; 1) + b_2(X, Y; -1),$

$b_2(X, Y; s) = 0.8N(X + 1.5, Y + 1.7; 1, 1)$

835             $+0.4F\left(X, Y; 0.3, 0.2; X_0, X_0, Y_0, Y_0; \sigma_{x0}, 0.07, \sigma_{y0}, 0.07, 20\right)$

836             $+0.7F\left(X, Y; 0.3, 0.2; 3X_0, 3X_0, 0.9Y_0, 0.9Y_0; \sigma_{x0}, 0.07, \sigma_{y0}, 0.07, 20\right)$

837             $+0.4F\left(X, Y; 0.3, 0.2; 6X_0, 6X_0, 0.8Y_0, 0.8Y_0; \sigma_{x0}, 0.07, \sigma_{y0}, 0.07, 20\right)$

838             $+0.4F\left(X, Y; 0.3, 0.2; 6X_0, 8X_0, 0.7Y_0, 0.7Y_0; \sigma_{x0}, 0.07, \sigma_{y0}, 0.07, 20\right)$

$\beta_3(\mathbf{u}) = F(X, Y; 0.3, 0.2; -6X_0, -6X_0, 0.5Y_0, 0.5Y_0; 0.3, 0.07, 0.3, 0.07, 20)$

$+F(X, Y; 0.3, 0.2; -6X_0, -6X_0, 0.8Y_0, 0.8Y_0; 0.3, 0.07, 0.3, 0.07, 20),$

$\beta_4(\mathbf{u}) = F(X, Y; 0.3, 0.2; 6X_0, 6X_0, 0.5Y_0, 0.5Y_0; 0.3, 0.07, 0.3, 0.07, 20)$

$+F(X, Y; 0.3, 0.2; 6X_0, 6X_0, 0.8Y_0, 0.8Y_0; 0.3, 0.07, 0.3, 0.07, 20),$

$\beta_5(\mathbf{u}) = F\left(X, Y; 0.3, 0.2; 0, 0, 0.6Y_0, 0.6Y_0; 0.7\sigma_{x0}, 0.07, 0.8\sigma_{y0}, 0.07, 20\right),$

$(\gamma_{X1}, \gamma_{Y1}, \omega_1) = (0, -0.06, 0), \ (\gamma_{X2}, \gamma_{Y2}, \omega_2) = (0, 0.05, 0), \ (\gamma_{X3}, \gamma_{Y3}, \omega_3) = (0.15, 0, -0.06),$

$(\gamma_{X4}, \gamma_{Y4}, \omega_4) = (-0.15, 0, 0.06), (\gamma_{X5}, \gamma_{Y5}, \omega_5) = (0, 0.12, 0),$

$\left(c_{w,\text{ext}}, c_{w,\text{end}}\right) = (3, 0.2), (t_{\text{ext}}, t_{\text{end}}) = (60000, 240000).$

Appendix D: Modification of extended shapes

We can curve and twist the shape of the extended sheet corresponding to the basic shape for the

morphogen distribution $\beta(\mathbf{u})$, by introducing two additional kinds of morphogens, denoted by γ

and ω. These additional morphogens can modify the sheet growth regulated mainly by $\beta(\mathbf{u})$, as

explained below.

For simplicity of explanation, we assume that $\beta(\mathbf{u})$ has a simple horn-like basic shape and

has a peak at $\mathbf{u} = \mathbf{0}$ in the material coordinates, so that it is symmetric with respect to both the X-

and Y-axes (figure 2a). If the sheet elements in the region of $X < 0$ grow more than those in $X >$

0 (Fig. 2e), this horn-like shape can be curved in the positive X-direction. As the morphogen

gradient in the X-direction, $\partial\beta(\mathbf{u})/\partial X$, is positive for $X < 0$ and negative for $X > 0$, the curved

horn can also be attained if the region of $\partial\beta(\mathbf{u})/\partial X > 0$ grows more than that of $\partial\beta(\mathbf{u})/\partial X < 0$.

Specifically, we get the curved horn by modifying equation (3) into

$$\frac{\partial \mathbf{M}(\mathbf{u},t)}{\partial t} = \mathbf{b}(\mathbf{u})\mathbf{b}(\mathbf{u})^{\mathrm{T}},$$

861
$$\mathbf{b}(\mathbf{u}) = \exp\left(\hat{\gamma}_X \frac{\partial\beta(\mathbf{u})}{\partial X}\right)\nabla\beta(\mathbf{u}) \tag{D.1}$$

862 with a constant $\hat{\gamma}_X$ describing the direction and strength of the curve. Note that equation (D.1)

863 applies even when the peak of the basic shape is not located at the origin. We can curve the horn-

864 like shape along a direction other than the X-axis, by introducing the morphogen distribution $\gamma(\mathbf{u})$

865 with its gradient being constant, given by $\nabla\gamma(\mathbf{u}) = (\hat{\gamma}_X, \hat{\gamma}_Y)^{\mathrm{T}}$, and by extending $\mathbf{b}(\mathbf{u})$ into

$$\mathbf{b}(\mathbf{u}) = Q\left(\nabla\gamma(\mathbf{u})^{\mathrm{T}}\nabla\beta(\mathbf{u})\right)\nabla\beta(\mathbf{u}),$$

866
$$Q(w) = \exp(w). \tag{D.2}$$

Regarding the function $Q(w)$, a function other than $\exp(w)$ (e.g., $Q(w) = 1 + w$) may also be

used, as long as it is no less than 1 and monotonically increases with respect to w.

The curved horn-like shape attained by equation (D.1) or (D.2) can be further twisted by slight

rotation of the local growth direction at each position of the sheet. We can achieve this by

introducing a uniform morphogen distribution $\omega(\mathbf{u}) = \hat{\omega}$ with a constant $\hat{\omega}$, and by further

modifying equation (D.2) into

$$\mathbf{b}(\mathbf{u}) = Q\left(\nabla\gamma(\mathbf{u})^{\mathrm{T}}\nabla\beta(\mathbf{u})\right)\mathbf{R}(\omega(\mathbf{u}))\nabla\beta(\mathbf{u}),$$

873
$$\mathbf{R}(\omega(\mathbf{u})) = \begin{pmatrix} \cos\omega(\mathbf{u}) & \sin\omega(\mathbf{u}) \\ -\sin\omega(\mathbf{u}) & \cos\omega(\mathbf{u}) \end{pmatrix} \tag{D.3}$$

874 (Figure 2f,i).

875 For fusion of the two basic shapes corresponding to $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$ by defining $\beta(\mathbf{u},t) =$

876 $\rho_1(t)\beta_1(\mathbf{u}) + \rho_2(t)\beta_2(\mathbf{u})$ (i.e., equation (8) in the main text), we can choose different degrees of

877 curves and twists for these basic shapes by allowing $\omega(\mathbf{u})$ and $\gamma(\mathbf{u})$ to change along time:

$$\gamma(\mathbf{u},t) = \frac{\rho_1(t)\gamma_1(\mathbf{u}) + \rho_2(t)\gamma_2(\mathbf{u})}{\rho_1(t) + \rho_2(t)},$$

878
$$\omega(\mathbf{u},t) = \frac{\rho_1(t)\omega_1(\mathbf{u}) + \rho_2(t)\omega_2(\mathbf{u})}{\rho_1(t) + \rho_2(t)}. \tag{D.4}$$

879 Collecting equations (B.1), (B.3), and (B.4), we get

$$\frac{\partial\mathbf{M}(\mathbf{u},t)}{\partial t} = \mathbf{b}(\mathbf{u},t)\mathbf{b}(\mathbf{u},t)^{\mathrm{T}},$$

880
$$\mathbf{b}(\mathbf{u},t) = Q\left(\nabla\gamma(\mathbf{u},t)^{\mathrm{T}}\nabla\beta(\mathbf{u},t)\right)\mathbf{R}(\omega(\mathbf{u},t))\nabla\beta(\mathbf{u},t). \tag{D.5}$$

881 If $\beta_1(\mathbf{u})$ and $\beta_2(\mathbf{u})$ have no temporal overlap, i.e., $\psi_{12} = \int \rho_1(t)\rho_2(t)dt = 0$, then we see that

$$\begin{aligned}
\mathbf{M}(\mathbf{u}, \tau) &= \mathbf{M}(\mathbf{u}, 0) + \int_0^\tau \mathbf{b}(\mathbf{u}, t)\mathbf{b}(\mathbf{u}, t)^{\mathrm{T}} \, dt \\
&= \mathbf{M}(\mathbf{u}, 0) + \mathbf{M}_{11}(\mathbf{u}) + \mathbf{M}_{22}(\mathbf{u})
\end{aligned} \tag{D.6}$$

with

$$\begin{aligned}
\mathbf{M}_{11}(\mathbf{u}) &= \mathbf{b}_1(\mathbf{u})\mathbf{b}_1(\mathbf{u})^{\mathrm{T}} \\
\mathbf{M}_{22}(\mathbf{u}) &= \mathbf{b}_2(\mathbf{u})\mathbf{b}_2(\mathbf{u})^{\mathrm{T}}, \\
\mathbf{b}_1(\mathbf{u}) &= Q\left(\nabla\gamma_1(\mathbf{u})^{\mathrm{T}}\nabla\beta_1(\mathbf{u})\right)\mathbf{R}\big(\omega_1(\mathbf{u})\big)\nabla\beta_1(\mathbf{u}), \\
\mathbf{b}_2(\mathbf{u}) &= Q\left(\nabla\gamma_2(\mathbf{u})^{\mathrm{T}}\nabla\beta_2(\mathbf{u})\right)\mathbf{R}\big(\omega_2(\mathbf{u})\big)\nabla\beta_2(\mathbf{u}).
\end{aligned} \tag{D.7}$$

In Figure 2 in the main text, $\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}$ are chosen as $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.15, 0, 0)$ for (e) and (h), and $(\hat{\gamma}_X, \hat{\gamma}_Y, \hat{\omega}) = (0.15, 0, 0.05)$ for (f) and (i).

Figure 1   Ito & Uchiumi

Initial state

Grown state

Extended state

(a) Initial sheet

(c) Grown sheet

(e) Extended sheet

?

(b) Morphogen density

(d) Growth metric

Growth mean: 1.85   (max: 3.17)

(f) Extension metric

Deviation mean: 0.0112 (max: 0.1163)
Growth mean: 1.87   (max: 3.20)

Spatial coordinates

Material coordinates

Figure 2

Ito & Uchiumi

Initial state

Grown state

Extended state

Figure 3   Ito & Uchiumi

Initial state

(a)

(b)

(c)

Growth mean: 3.05   (max: 13.01)

(d)

Grown state

(e) $\psi_{12} = 1$

Growth mean: 3.05   (max: 13.01)

(f) $\psi_{12} = 0.5$

Growth mean: 3.01   (max: 11.45)

(g) $\psi_{12} = 0$

Growth mean: 2.98   (max: 9.96)

Extended state

(h) $\psi_{12} = 1$

(i) $\psi_{12} = 0.5$

(j) $\psi_{12} = 0$

Initial state

Grown state

Extended state

Figure 5 Ito & Uchiumi

Initial state

Grown state

Extended state

Figure 6   **Ito & Uchiumi**

Figure 7   Ito & Uchiumi

Figure 8   Ito & Uchiumi

(a)

(b)