The carnivoran adaptive landscape reveals functional trade-offs among the skull,

appendicular, and axial skeleton

Chris J. Law[1,2], Leslea J. Hlusko[3], Z. Jack Tseng[2]

[1]Burke Museum and Department of Biology, University of Washington

[2]Department of Integrative Biology, University of California Berkeley

[3]National Research Center on Human Evolution (CENIEH), Burgos, Spain

Abstract

Analyses of form-function relationships are widely used to understand links between

morphology, ecology, and adaptation across macroevolutionary scales. However, few have

investigated functional trade-offs and covariance within and between the skull, limbs, and

vertebral column simultaneously. In this study, we investigated the adaptive landscape of skeletal

form and function in carnivorans to test how functional trade-offs between these skeletal regions

contribute to ecological adaptations and the topology of the landscape. We found that

morphological proxies of function derived from carnivoran skeletal regions exhibit trade-offs

and covariation across their performance surfaces, particularly in the appendicular and axial

skeletons. These functional trade-offs and covariation correspond as adaptations to different

adaptive landscapes when optimized by various factors including phylogeny, dietary ecology,

and, in particular, locomotor mode. Lastly, we found that the topologies of the optimized

adaptive landscapes and underlying performance surfaces are largely characterized as a single

gradual gradient rather than as rugged, multipeak landscapes with distinct zones. Our results

suggest that carnivorans may already occupy a broad adaptive zone as part of a larger

mammalian adaptive landscape that masks the form and function relationships of skeletal traits.

26
27   **Key words**: Adaptive peak, Carnivora, ecomorphology, functional morphology, phylogenetic

28   comparative methods

29
30   **Introduction**

31   The diversity of animal forms is one of the most salient patterns across the tree of life,

and how morphological variation relates to the ecological diversity and survival of species across

macroevolutionary time remains a core question in evolutionary biology. The varying strength of

form–function relationships provide biologists with insight into the specificity of morphological

structure in determining species' abilities to carry out ecological tasks (i.e., performances),

especially when behavioral observations are scarce. Thus, performance is considered the link

between morphology, ecology, and fitness (Arnold 1983; Wainwright 1994; Higham et al. 2021).

Functional traits are morphological, phenological, and physiological traits that affect fitness and

are often used to estimate performance (Higham et al. 2021). Many researchers have examined

the form-function relationship of the skull (Santana et al. 2010; Collar et al. 2014; Law et al.

2018; Tseng et al. 2023), appendicular skeleton (Sustaita et al. 2013; Dickson and Pierce 2019;

Sansalone et al. 2020; Amson and Bibi 2021), and axial skeleton (Polly et al. 2016; Jones et al.

2018, 2021; Law et al. 2019; Stayton 2019), often finding trade-offs that are hypothesized to

facilitate distinct ecological adaptations. For example, the gradient from short, broad jaws to

long, narrow jaws is associated with a functional trade-off between generating stronger bites and

quicker bites or wider gapes (Herring and Herring 1974; Dumont and Herrel 2003; Slater and

Van Valkenburgh 2009; Slater et al. 2009; Forsythe and Ford 2011; Santana 2015), and similarly,

the gradient from gracility to robustness in limb bones is associated with a functional trade-off

between increasing cost of transport associated with cursoriality and resisting stresses associated

with locomoting through resistant media (Martín-Serra et al. 2014a, 2014b; Kilbourne 2017;

Hedrick et al. 2020; Muñoz 2020; Marshall et al. 2021; Rickman et al. 2023). However, most of

these studies investigate trade-offs within individual bones (e.g., the mandible, humerus, or

femur) and few have investigated functional trade-offs and covariation within and among the

three major skeletal systems.

The rise of adaptive landscape analyses enables researchers to investigate the adaptive

evolution of performance by elucidating the underlying links among morphology, ecology, and

fitness benefits (i.e., adaptiveness) at the macroevolutionary level (Arnold et al. 2001). Although

Ornstein–Uhlenbeck (OU) models (Hansen 1997; Butler and King 2004; Beaulieu et al. 2012;

Uyeda and Harmon 2014; Bastide et al. 2018) are widely used to test for the presence of adaptive

zones or peaks (e.g., Collar et al. 2014; Price and Hopkins 2015; Friedman et al. 2016; Zelditch

et al. 2017; Arbour et al. 2019; Law 2022; Slater 2022), it remains difficult to characterize the

full topology (i.e., peaks, valleys, and slope) of the adaptive landscape as well as assess the

relative importance of multiple performance traits and their contributions to overall adaptive

landscape using these models. Adaptive landscape analyses (Polly et al. 2016; Dickson and

Pierce 2019; Dickson et al. 2021) can overcome these limitations by examining the distribution

of species in morphospace and its relationship to the relative importance of various functional

traits on the topology of the adaptive landscape. While an increasing number of studies have

used functional adaptive landscapes to examine links between morphological diversity and

functional performance (Polly et al. 2016; Dickson and Pierce 2019; Stayton 2019; Dickson et al.

2021; Jones et al. 2021; Tseng et al. 2023), no study has yet to investigate these relationships

among the skull, limbs, and vertebral column.

Here, we examined the trade-offs and covariation among individual performance surfaces

derived from functional traits of the skull, appendicular skeleton, and axial skeleton as well as

assessed their relative contributions to ecological adaptations and the overall landscape. To

explore these patterns, we used terrestrial carnivorans (e.g., bears, cats, dogs, weasels, and their

relatives) as our model because of their high species richness and well-studied broad

morphological and ecological diversity. Numerous researchers have investigated the

morphological diversity of the carnivoran skull (Radinsky 1981; Van Valkenburgh 2007;

Figueirido et al. 2011; Law et al. 2018, 2022; Tseng and Flynn 2018; Slater and Friscia 2019),

appendicular skeleton (Van Valkenburgh 1985, 1987; Iwaniuk et al. 1999; Samuels et al. 2013;

Martín-Serra et al. 2014a, 2014b), vertebral column (Randau et al. 2017; Figueirido et al. 2021;

Martín-Serra et al. 2021), and overall body plan (Law 2021a, 2021b; Slater 2022). This diversity

is attributed to mosaic evolution, in which different skeletal components exhibit distinct modes

of evolution either from phylogenetic natural history (Uyeda et al. 2018) or from selection for

ecological adaptations (Law et al. 2024). The ability of individual skeletal components to adapt

to specific ecological factors independently from each other may have contributed to the clade's

hierarchical evolution. The hierarchical evolution is primarily framed by dental adaptations along

an axis of dietary resource use, which are hypothesized to facilitate the early radiation of

carnivorans across a rugged, multi-peak adaptive landscape (Slater and Friscia 2019).

Subsequent evolution led to the continual partitioning between clades, resulting in the origination

of extant carnivoran families that occupy different adaptive zones (Humphreys and Barraclough

2014) with distinct morphologies in the skull, appendicular, and axial skeletons (Law 2021a;

Law et al. 2022, 2024). Skeletal variation in the mandible, hindlimb, and post-diaphragmatic

region of the vertebral column then arose along shared ecological axes among taxa, theoretically

leading to distinct ecological zones across the adaptive landscape (Law et al. 2022, 2024).

Despite this large body of knowledge in carnivoran morphology, the functional implications of

these skeletal traits remain to be tested across the adaptive landscape; that is, how do

morphological traits in the skull, appendicular skeleton, and vertebral column dictate the

ecological performance of carnivoran species?

Our goals of this study were three-fold. First, we described functional trade-offs and

covariation among individual performance surfaces derived from functional traits from the skull,

appendicular skeleton, and axial skeleton. Second, because morphological traits are often

associated with locomotor and dietary adaptive peaks (Slater and Friscia 2019; Law et al. 2022,

2024; Slater 2022) and their functional trade-offs are often hypothesized to facilitate distinct

ecological adaptations (Slater and Van Valkenburgh 2009; Slater et al. 2009; Martín-Serra et al.

2014b, 2014a), we tested how these performance surfaces contribute to unique adaptive

landscapes and the formation of adaptive zones along locomotor, dietary, and phylogenetic axes.

Third, we explored the topology of the adaptive landscape of carnivorans. Previous work using

OU modeling provided evidence that morphological proxies for appendicular function exhibit

relatively low ruggedness across the adaptive landscape despite also exhibiting distinct adaptive

zones (Slater 2022). Adaptive landscape analyses will further clarify whether if these adaptive

zones are steep peaks or broad plateaus as well as how functional traits from the skull and axial

skeleton contributes to the adaptive landscape. Overall, this work provides a baseline

understanding of the relative contributions of the skull, appendicular skeleton, and axial skeleton

to the adaptive landscape, setting a foundation for future hypothesis testing on the processes that

influence the evolution of animal form and function.

Methods

Morphospace and functional proxies

We created a morphospace of 109 terrestrial carnivoran species based on 136 linear and

angular measurements that capture morphological variation across the entire skeleton (Fig. S1).

This dataset includes seven cranial traits, seven mandibular traits, 13 forelimb traits, 13 hindlimb

traits, and 11 traits each in the third cervical, fifth cervical, first thoracic, middle thoracic,

diaphragmatic thoracic, last thoracic, first lumbar, middle lumbar and last lumbar vertebrae. We

were able to incorporate representatives from 12 of the 13 extant terrestrial carnivoran families;

specimens from Prionodontidae were unavailable. We removed size effects on linear

measurements by calculating the log shape ratio (i.e., ln[trait/size]) of each skeletal trait, where

size is the geometric mean of all linear trait measurements (see Sensitivity Analyses 1 in the

Supplementary Materials for analyses based on non-size corrected data). We used only adult

male specimens because carnivorans exhibit differing degrees of sexual dimorphism (Law 2019).

We then conducted a principal component analysis (PCA) using the covariance matrix on all

size-corrected measurements and used the first two PC axes (46.7% of the total variance) to

create the morphospace (Fig. 1; see Table S1 for PC loadings). Carnivoran species are widely

distributed across the morphospace except for the bottom-left region in which no species occupy

(-PC1, -PC2). We chose not to run a phylogenetic PCA because we are interested in the primary

dimensions of morphological variation regardless of phylogenetic structuring. In addition, pPCA

is more difficult to interpret because it is a mixture of major axes that describe non-phylogenetic

variation and scores that contain phylogenetic components of variation, and pPC axes are not

orthogonal to each other, meaning that the first two axes (which we use for the adaptive

landscape analyses) may include less variance explained than PCA by containing correlated

variance components rather than independent ones (Polly et al. 2013).

From the 136 morphological traits, we then calculated 27 morphological proxies of

function as proxies for functional traits (hereinafter called "functional proxies"; Table 1; see

Supplementary Materials for full biomechanical and ecomorphological justification). These

functional proxies are often used to capture the functional diversity of the skull (Greaves 2012),

limbs (Davis 1964; Samuels et al. 2013), and vertebral column (Boszczyk et al. 2001; Jones et al.

2020, 2021).

Ecological traits

We classified the 109 carnivoran species into one of seven locomotor regimes: arboreal

(species that primarily live and forage in trees and rarely come down to the ground), cursorial

(species that displays rapid bounding locomotion, particularly during hunting), scansorial

(species that spend equal time in trees and on the ground), semi-aquatic (species that regularly

swim for dispersal and/or foraging), semi-fossorial (species that regularly dig for shelter and/or

foraging), and terrestrial (species that primarily live on the ground and rarely run, climb, dig, or

swim during foraging). Terrestrial species were further categorized as terrestrial hunters (species

that exhibit ambush and/or pouncing behaviors to kill prey) and terrestrial non-hunters (species

that rarely hunt for prey). We also classified each species into one of seven dietary regimes: large

prey hypercarnivory (consist of >70% terrestrial vertebrate prey that exceeds the predator's own

body mass), medium prey hypercarnivory (consist of >70% terrestrial vertebrate prey that are up

to the predator's own body mass), small prey hypercarnivory (consist of >70% terrestrial

vertebrate prey that are up to 20% of the predator's own body mass), omnivory (consist of >50%

terrestrial vertebrates), insectivory (consisting of >70% invertebrates), aquatic carnivory (consist

of >90% aquatic prey), and herbivory (consist of >90% plant material). These locomotor and

dietary regimes are widely used to describe carnivoran ecology and have demonstrated

significant associations with various traits of the cranial, appendicular, and axial skeletons of

carnivorans (Van Valkenburgh 1987, 2007; Friscia et al. 2007; Samuels et al. 2013).

Categorization for locomotor and dietary regimes were obtained from previous work

(Van Valkenburgh 1985; Samuels et al. 2013; Law 2021a) with minor edits based on literature

review.

To check the relationship between the morphospace (i.e., PCs 1 and 2) with ecological

regimes, we conducted two multivariate ANOVA models (i.e., morphospace ~ locomotor mode

and morphospace ~ dietary ecology) in the R package RRPP v1.4.0 (Adams and Collyer 2018).

We found that the morphospace exhibited a significant relationship with locomotor mode (R^2 =

0.29, F = 6.78, P = 0.001) and with dietary ecology (R^2 = 0.16, F = 3.12, P = 0.001). Post-hoc

pairwise comparison tests also indicated significance differences among most ecological regimes

(Table S2). Testing using multivariate PGLS models (Clavel et al. 2019; Clavel and Morlon

2020) also indicated that the exhibited a significant relationship with locomotor mode (Pagel's λ

= 0.97, Pillai's trace = 0.202, P = 0.041) and with dietary ecology (Pagel's λ = 0.97, Pillai's trace

= 0.213, P = 0.017).

Performance surfaces and adaptive landscapes

We investigated the functional optimality of the skeleton using adaptive landscape

analyses (Polly et al. 2016; Dickson and Pierce 2019; Dickson et al. 2021; Jones et al. 2021) in

the R package Morphoscape (Dickson et al. 2021). We first created 27 performance surfaces by

interpolating each of the 27 functional proxies across the morphospace surface using ordinary

kriging. Initial inspection of these surfaces revealed that the performance surfaces of four

functional proxies (masseter mechanical advantage [masMA], scapula index [SI], gluteal index

[GI], and tibial robustness [TRI]) exhibited topological peaks and valleys that outlined clusters of

species and even single species (Fig. 2; see section below about the downsides to using empirical

data instead of theoretical data). Therefore, we removed these four functional proxies from

subsequent analyses.

We computed a combined adaptive landscape (W) as the summation of all 23

performance surfaces (F_n), each weighted by their relative importance or contribution to overall

fitness (w_n) (Polly et al. 2016):

$$W = w_1F_1 + w_2F_2 + \bullet \bullet \bullet + w_nF_n$$

where W is optimized as the likelihood of combinations of performance surface and relative

fitness, under the definition that the total fitness sums to 1 and the variance of all surfaces is

equal (Polly et al. 2016). We tested all possible combinations of weights, ranging from 0 to 1 in

increments of 0.25, across a total of 27,405 possible adaptive landscapes (see Sensitivity

Analyses 2 in the Supplementary Materials for further investigation of the concern with number

of partition weight increments).

We identified the optimally weighted landscape that maximizes the fitness of each

locomotor regime using the function calcWprimeBy and tested whether these optimal landscapes

are significantly different among locomotor ecological groups using the function

multi.lands.grp.test. Significance testing for differences among landscapes was performed by

comparing the number of landscapes shared by the top 5% of each group with the total number

of landscapes in the top 5% of models (Jones et al. 2021). The top percentile of each group was

determined using a chi-squared test. We also investigated differences in adaptive landscapes

among dietary groups and carnivoran families that contained more than one species.

Creating adaptive landscapes using theoretical traits

A great concern in using empirical data in creating performance surfaces and adaptive

landscapes is that empirical-specific values are always less evenly distributed in the

morphospace. The unevenness contributes to heterogeneous resolution of the interpolation

applied to the space by the ordinary kriging method, and thus unevenness in the landscape itself.

Denser sampled regions will be more likely to have topological relief (i.e., peaks and valleys)

than sparsely sampled regions when using actual specimen values, not necessarily because of a

real underlying peak there. To mitigate these issues, many researchers have used theoretical data

across an evenly spaced grid to create performance surfaces and adaptive landscapes (Polly et al.

2016; Dickson and Pierce 2019; Smith et al. 2021; Tseng et al. 2023; Sansalone et al. 2024).

Therefore, we investigated whether there is consistency in our adaptive landscape

analyses when using functional proxies derived from actual species versus using functional

proxies derived from theoretical species. To fully sample skeletal variation throughout the

morphospace, we generated 63 theoretical species evenly across the morphospace in a 9 x 7 grid

along the first two PCs. We then generated theoretical morphological traits from each of the 63

theoretical species and calculated the 27 functional proxies (Table 1). We then performed the

same procedures as described above to generate the 27 performance landscapes and adaptive

landscapes optimized for locomotor ecology, diet, and family. We found that the patterns found

in adaptive landscapes are similar with only slight differences. To avoid confusion between the

two approaches, we report the results using the theoretical morphologies in the Supplementary

Results of the Supplementary Materials.

Sensitivity Analyses

Lastly, we conducted sensitivity analyses to investigate whether (1) removing the effects

of size on the morphological traits in creating the morphospace and (2) the coarse increments of

partition weights may potentially influence our interpretation of the results. We found that these

factors do not affect our findings (see Sensitivity Analyses section in the Supplementary

Materials).

Results

Performance surfaces reveal trade-offs and covariation within skeletal systems

Each functional proxy mapped onto the morphospace revealed both unique and similar

performance surfaces that characterize trait groups, suggesting that functional trade-offs and

covariation are present within the skull, appendicular skeleton, and axial skeleton. In the skull,

mechanical advantage of the temporalis (temMA) highest towards the left and bottom-left (-PC1,

-PC2) of morphospace and declining towards the top-right (+PC1, +PC2). In contrast, there is no

distinct pattern in masseter mechanical advantage (masMA) (Fig. 2).

In the forelimb, there is a trade-off between limb elongation and elbow robustness:

functional proxies of radius (BI) and metacarpal (MAN) elongation are highest in the top-right

(+PC1, +PC2) of morphospace. BI transitions towards increased robustness to the bottom-right

(+PC1, -PC2), whereas MAN transitions towards increased robustness to the left (-PC1). Proxies

associated with increased mechanical advantage of elbow extension (OLI) and attachment sites

for forearm flexor, pronator, and supinator muscles on the humeral epicondyles (HEI) and ulna

(URI) are highest in the bottom-left and lowest in the top-right. Overall robustness of humerus

(HRI) is highest on the left side of the morphospace (-PC1) and transitions towards increased

elongation to the right (+PC1), following a similar distribution as the latter indices. There is no

distinct pattern in scapula index (SI). The hindlimb also exhibits a trade-off between elongation

and robustness: indices of tibial (CI) and metatarsal (PES) elongation tend to be highest in the

right side (+PC1, with CI also trending towards -PC2) and transitions to increased robustness

towards to the left side (-PC2), whereas indices for femoral (FEI, FRI) robustness tend to be

highest on the bottom (-PC2) but transitions towards less robustness in the top-right (+PC1,

+PC2). There are no distinct patterns in indices for gluteal muscles (GI) and tibial robustness

(TRI).

In the vertebral joints, the performance surfaces show a trade-off between joint torsional

angle (JTA) as a proxy for range of rotational motion and joint verticality (JV) as a proxy for

sagittal mobility. For the cervical, diaphragmatic, and lumbar joints, JTA is highest in the

bottom-left of the morphospace (-PC1, -PC2) and declines diagonally to the top-right (+PC1,

+PC2), whereas JV exhibits the opposite pattern (i.e., highest in the top-right and lowest in the

bottom-left). The thoracic vertebra exhibits similar JTA and JV distributions but in the horizontal

plane (i.e., highest JTA in the left side of morphospace and highest JV in the right). In all

vertebral joints, second moment of area (SMA) as a proxy for stiffness tends to be greatest

towards the top-left of morphospace (-PC1, +PC2) and declines towards the right side of

morphospace (+PC1).

275
Optimized adaptive landscapes reveal trade-offs and covariation among skeletal systems

After summation of all performance surfaces based on optimized weights, we found that

the combined adaptive landscape is heavily weighted for sagittal mobility of the pre-

diaphragmatic thoracic ($w_{JV} = 0.41$) and lumbar ($w_{JV} = 0.38$) regions (Fig. 3A; Table S3). When

adaptive landscapes are optimized by locomotor ecologies, we found different degrees to which

the 23 functional proxies are incorporated among the different adaptive landscapes. The cursorial

landscape is characterized by lengthening of the forelimb, particularly in the metacarpal ($w_{MAN} =$

0.56) and, to a lesser extent, the radius ($w_{BI} = 0.04$) (Fig. 3C; Table S3). The cursorial landscape

is also strongly weighted with functional proxy associated with increased sagittal mobility of the

pre-diaphragmatic thoracic joints ($w_{JV} = 0.38$). The semi-aquatic and semi-fossorial landscapes

do not significantly differ from one another ($P = 0.184$; Table 2), and both are similarly weighted

for larger humeral epicondyles (semi-aquatic $w_{HEI} = 0.28$; semi-fossorial $w_{HEI} = 0.40$) and

increased joint torsion in the cervical joints (semi-aquatic $w_{JTA} = 0.39$; semi-fossorial $w_{JTA} =$

0.11) (Fig. 3E, F; Table S3). The semi-aquatic landscape is further strongly weighted for more

robust ulna ($w_{URI} = 0.16$), whereas the semi-fossorial landscape is further strongly weighted for

increased joint torsion in the diaphragmatic joint ($w_{JTA} = 0.37$) (Fig. 3E, F; Table S3). The

terrestrial non-hunter landscape is not significantly different from the cursorial and semi-

fossorial landscapes (Table 2). This landscape is strongly weighted for increased joint torsion in

the diaphragmatic joint ($w_{JTA} = 0.60$) and lengthening of the metacarpal ($w_{MAN} = 0.21$) (Fig. 3H;

Table S3). The remaining locomotor landscapes (i.e., arboreal, scansorial, and terrestrial hunter)

do not significantly differ from one another (Table 2) and are heavily weighted for sagittal

mobility of the pre-diaphragmatic thoracic and lumbar joints (Fig. 3B, D, G; Table S3).

When adaptive landscapes are optimized by dietary ecologies, we found that significant

differences in landscapes appear associated with piscivory (Fig. S2; Table S4). The piscivorous

adaptive landscape is similarly weighted for larger humeral epicondyles ($w_{HEI} = 0.32$), more

robust ulna ($w_{URI} = 0.22$), and increased joint torsion in the cervical joints ($w_{JTA} = 0.22$) (Fig. S2;

Table S5). In contrast, adaptive landscapes based on other diets are not significantly different

from each other and are largely characterized by increased sagittal mobility of the pre-

diaphragmatic thoracic and/or lumbar joints (Fig. S2; Table S4; Table S5).

Lastly, we found different adaptive landscapes when they are optimized by family.

Adaptive landscapes for felids, viverrids, euplerids, herpestids, canids, and procyonids are not

significantly different from each other and all remain heavily weighted for sagittal mobility of

the pre-diaphragmatic thoracic and/or lumbar joints (Fig. 4; Table S6; Table S7). The canid

landscape is also weighted for lengthening of the metacarpal (Fig. 4G; Table S6). The mephitid

and mustelid landscapes resemble semi-fossorial and semi-aquatic landscapes, respectively. The

mephitid landscape is equally weighted by larger humeral epicondyles ($w_{HEI} = 0.38$) and

increased joint torsion in the diaphragmatic joint ($w_{JTA} = 0.38$) (Fig. 4I; Table S6), whereas the

mustelid landscape is weighted by increased sagittal mobility of the pre-diaphragmatic thoracic

joints ($w_{JV} = 0.22$), more robust ulna ($w_{URI} = 0.17$), larger humeral epicondyles ($w_{HEI} = 0.13$),

and increased joint torsion in the cervical ($w_{JTA} = 0.18$) and diaphragmatic ($w_{JTA} = 0.13$) joints

(Fig. 4K; Table S6). Only the mephitid landscape significantly differs from all other families

(Table S7). The hyaenid landscape weighted heavily for elongation of the radius ($w_{BI} = 0.50$) and

metacarpal ($w_{MAN} = 0.50$) (Fig. 4D). Lastly, the ursid landscape weighted heavily for larger

humeral epicondyles ($w_{HEI} = 0.38$), increased joint torsion in the thoracic joints ($\sum w_{JTA} = 0.41$),

and increased robustness of the ulna ($w_{URI} = 0.14$) and humerus ($w_{HRI} = 0.08$) (Fig. 4H). Both

hyaenid and ursid landscapes significantly differ with most other family-specific landscapes

(Table S7).

Discussion

The diversity found in the carnivoran skeletal system is attributed to mosaic evolution, in

which the mandible, hindlimb, and post-diaphragmatic vertebrae showed evidence of adaptation

towards ecological regimes whereas the cranium, forelimb, and pre-diaphragmatic vertebrae

reflect clade-specific evolutionary shifts (Law et al. 2024). Using adaptive landscape analyses,

we further found that functional proxies derived from this morphological diversity exhibit trade-

offs and covariation, particularly within and between the appendicular and axial skeletons. These

functional trade-offs and covariation corresponded as adaptations to different adaptive

landscapes when optimized by various factors including phylogeny, dietary ecology, and, in

particular, locomotor mode. Lastly, these adaptive landscapes and underlying performance

surfaces were characterized by rather broad slopes, hinting that carnivorans occupy a broad

adaptive zone as part of a larger mammalian adaptive landscape that masks the form and

function relationships of skeletal traits.

337
Performance surfaces reveal trade-offs and covariation among skeletal systems

In the appendicular skeleton, we found support that the gradient from gracility to

robustness often found as the primary source of variation across limb bone morphospace

(Martín-Serra et al. 2014a, 2014b; Kilbourne 2017; Hedrick et al. 2020; Muñoz 2020; Marshall

et al. 2021; Rickman et al. 2023) signifies a functional trade-off between increasing cost of

transport associated with cursoriality and resisting stresses associated with locomoting through

resistant media (Fig. 2B). Specifically, long, gracile limb bones particularly on the distal ends of

the limbs facilitate increased stride length and decreased moment of inertia of limbs, which in

turn decreases the energetic cost of transport and increases running speeds (Kram and Taylor

1990; Strang and Steudel 1990; Garland and Janis 1993; Polly 2007; Pontzer 2007a, 2007b;

Kilbourne and Hoffman 2013). In contrast, short, robust limb bones facilitate resistance to

bending and shearing stresses and increased mechanical advantage for forceful movements by

reducing the out-lever of the limb and increasing the in-lever of muscle forces (Hildebrand

1985a; Nakai and Fujiwara 2023). Robustness also permits increased surface area of the bone for

more muscles to attach. Enlargement of the humeral and femoral epicondyles increases the

attachment sites of several muscles (i.e., flexors, pronators, and supinators in the forelimb and

gastrocnemius and soleus muscles in hindlimb) responsible for generating power, force, and

stability (Davis 1964; Hildebrand 1985a; Lessa and Stein 1992; Lagaria and Youlatos 2006).

Additionally, an enlarged olecranon process facilitates stronger extension and flexion of the

elbow and wrist by increasing mechanical advantage of the triceps brachii and

dorsoepitrochlearis muscles and providing greater attachment sites for the ulnar head of the

flexor carpi ulnaris (Davis 1964; Hildebrand 1985a; Lessa and Stein 1992; Lagaria and Youlatos

2006). These adaptations facilitate the ability to generate large forces during certain locomotor

behaviors such as digging (Hildebrand 1985a; Lessa and Stein 1992; Lagaria and Youlatos 2006;

Samuels et al. 2013; Rose et al. 2014; Rickman et al. 2023) or swimming (Fish 2000; Samuels

and Van Valkenburgh 2008; Samuels et al. 2013; Kilbourne 2017).

In the axial skeleton, our investigation using functional proxies suggested that

carnivorans exhibit trade-offs between joint mobility and range of axial rotation. That is, high

sagittal mobility covaries with low range of axial rotation whereas low sagittal mobility covaries

with high range of axial rotation. This pattern was surprising because, compared to other

tetrapods, mammals exhibit intervertebral joints that are characterized by high sagittal mobility

and high axial rotation (Jones et al. 2021). A possible explanation was that the covariation

between high sagittal mobility and low axial rotation may serve as a further adaptation to

increasing forward locomotion by prioritizing flexibility in the sagittal plane through the

reduction of torsional twisting. High mobility of the backbone in the sagittal plane has long been

recognized as a key adaptation facilitating the diversity of different locomotor habits in

mammals, particularly asymmetrical gaits (e.g., gallop, half-bound, bound) by enabling extensive

dorsoventral flexion of the body axis (Hildebrand 1959, 1985b; Gambaryan 1974; Schilling and

Hackert 2006). The reduction of torsional twisting in carnivorans may, therefore, prioritize force

generation in the sagittal plane (rather than parasagittal or transverse plane) needed for these

locomotor behaviors. Evidence for this hypothesis was found in comparisons with ungulates,

where carnivorans exhibit up to ~38° more sagittal mobility in the lumbar region and up to 200%

less axial rotational mobility in the thoracic region compared to ungulates (Belyaev et al. 2021,

2022, 2023). Increased rotational mobility of the backbone in ungulates is hypothesized to

enhance agile maneuvering such as sharp cornering and quick directional changes when escaping

from predators (Belyaev et al. 2023).

Functional covariation between appendicular and axial skeletons optimizes the adaptive

landscapes of some locomotor ecologies

The optimized adaptive landscape is heavily weighted for sagittal mobility of the pre-

diaphragmatic thoracic and lumbar joints (Fig. 3A), indicating that flexibility in the sagittal plane

serves an important functional role for all carnivorans. When the adaptive landscapes are

optimized based on locomotor mode, diet, or family, we found that locomotor behavior could

provide an explanation for most landscape patterns. In our analyses of locomotor landscapes, we

found that semi-aquatic and semi-fossorial landscapes were not significantly different to each

other but are distinct to other locomotor landscapes (Fig. 3; Table 2). The peaks of both semi-

aquatic and semi-fossorial landscapes occur in the bottom-left regions of morphospace, and

species with either locomotor modes occur in overlapping regions of the landscapes. In contrast,

the peaks of most of the remaining locomotor landscapes, particularly the cursorial and terrestrial

hunter landscapes, occur near the top-right region of morphospace. The opposing locations of

these landscape peaks corresponds to the functional trade-offs identified by the performance

surfaces, suggesting that covariation of the appendicular and axial skeletons facilitate adaptations

to each of these locomotor behaviors at these extreme ends. These functional trade-offs are

largely independent of size effects because body size variation scales from the top-left to bottom-

right of morphospace whereas the trade-offs scale from the top-right to the bottom-left (Fig. 3;

Fig. 4; Fig. S2). Sensitivity analyses examining the performance surfaces and adaptive

landscapes using non-size corrected morphological traits confirm this pattern (see Supplementary

Materials).

Covariation of the appendicular and axial skeletons and its role in facilitating adaptations

to locomotor behaviors is apparent in semi-aquatic and semi-fossorial species. Both behaviors

require large force generation for swimming and digging, and the appendicular and axial

skeletons of semi-aquatic and semi-fossorial carnivorans are functionally adapted for increased

elbow extension through enlarged elements of the limbs and increased axial rotation in the

vertebral column (Fig. 3E, F; Table S3). It is well documented that adaptations in the elbows and

knees facilitate the ability to generate large power strokes for turning and stabilizing the body

while swimming (Fish 2000; Samuels and Van Valkenburgh 2008; Samuels et al. 2013;

Kilbourne 2017). These similar adaptations also enable semi-fossorial species to generate large

forces to dig (Hildebrand 1985a; Lessa and Stein 1992; Lagaria and Youlatos 2006; Samuels et

al. 2013; Rose et al. 2014; Rickman et al. 2023) and improve stability and load transfer during

clearing (Hildebrand 1985a; Casinos et al. 1993; Samuels and Van Valkenburgh 2008; Rickman

et al. 2023). What remains largely undetermined is the importance of increased axial joint

rotation during swimming or digging. Presumably, for swimmers, torsional rotation of the

intervertebral joints increases the maneuverability and ability to perform rapid turns in water

(Fish 1994; Fish et al. 2003). Fully aquatic seals exhibit more flexible and compliant

intervertebral joints compared to terrestrial mammals (Gál 1993); whether their joints are also

more capable for torsional rotation remains to be studied. For semi-fossorial carnivorans,

increased axial rotation of the vertebral column may provide additional leverage when digging

through sediment. Evidence for increased axial rotation has been observed in the semi-fossorial

nine-banded armadillo; experimentation on intervertebral joint flexion in this species revealed

rotational motion in the joints despite not being explicitly tested (Oliver et al. 2016).

Nonetheless, the benefits of increased joint rotation for digging remains puzzling. Interestingly,

no carnivorans, including semi-aquatic and semi-fossorial species, occupied the highest regions

(bottom-left) of semi-aquatic or semi-fossorial landscapes. A likely explanation was that further

axial rotation is biologically impossible for these carnivorans given their vertebral morphology

that may be under evolutionary constraints having originated from terrestrial carnivorans. Thus,

semi-aquatic and semi-fossorial carnivorans may already be at the highest region of the adaptive

landscape that is biologically feasible.

Cursorial species tend to be most concentrated in the top-right region of morphospace,

and thus appears to serve as the opposing extreme to the semi-aquatic and semi-fossorial

landscapes. The majority of cursorial carnivorans occupy regions of morphospace that

corresponded to the highest regions of the cursorial landscape. This landscape indicates that the

appendicular and axial skeletons of cursorial carnivorans are functionally adapted for increased

stride length through elongation of forelimb and increased sagittal flexibility of the full vertebral

column (Fig. 3C; Table S3). As described above, these adaptations increase running speeds and

reduce the energetic cost of transport by prioritizing dorsoventral flexion and extension in the

sagittal plane (Hildebrand 1959, 1985b; Gambaryan 1974; Kram and Taylor 1990; Strang and

Steudel 1990; Garland and Janis 1993; Schilling and Hackert 2006; Pontzer 2007a, 2007b;

Kilbourne and Hoffman 2013; Belyaev et al. 2023).

The remaining locomotor landscapes were heavily weighted for sagittal mobility of the

pre-diaphragmatic thoracic and/or lumbar joints (Fig. 3; Table S3). That the arboreal landscape

was not heavily weighted by additional functional proxies is surprising considering that

arboreality is often described as a specialized form of locomotion (Young 2023). A possible

explanation was that carnivorans do not display the full diversity of arboreal behaviors (e.g.,

brachiation, leaping, suspensory climbing) performed by other mammals. Another possibility

was that we did not include all possible functional proxies in our analyses. For example, the ratio

between proximal manual phalanx length and metacarpal length has been shown to accurately

predict climbing frequency in rodents (Nations et al. 2019), and thus may have altered the

optimized weights of the arboreal adaptive landscape if this proxy or others were included in this

current study. These unaccounted sources may also explain why arboreal species lie away from

the highest regions of the adaptive landscape (Fig. 3B).

Adaptive landscapes optimized based on family also demonstrated similar patterns as

locomotor-specific landscapes (Fig. 4; Table S6). Most family-specific landscapes were heavily

weighted for sagittal mobility of the pre-diaphragmatic thoracic and/or lumbar joints. Canids,

which primarily exhibit cursorial or terrestrial hunting behaviors, exhibited similar patterns with

the cursorial landscape with increased sagittal mobility in other regions of the vertebral column

and elongation of the metacarpal (Fig. 4G; Table S6). Likewise, mephitids and mustelids

comprise of many semi-aquatic and semi-fossorial species and thus exhibited similar patterns

with the semi-aquatic and semi-fossorial landscapes of increased intervertebral joint rotation and

enlarged limb joints (Fig. 4I, K). Adaptive landscapes for hyaenids and ursids were both unique

compared to other family-specific landscapes (Table S6). The hyaenid landscape was heavily

weighted for relative elongation of the forelimb (Fig. 4D); however, a caveat was that our low

sample size of just two species reflected a biased representation of taxa with elongate forelimb

and sloped back found in their extant diversity. Lastly, the ursid landscape was heavily weighted

for increased robustness of the stylopodia of the limbs (Fig. 4H). These results were unsurprising

as ursids are the biggest terrestrial carnivorans and these traits support their heavy bodies against

the effects of gravity (Polly 2007; Jones et al. 2021).

Lastly, we found that only the piscivorous landscape was significantly different from all

other landscapes optimized based on dietary ecology (Table S4). Unsurprisingly, the piscivorous

landscape resembles the semi-aquatic landscape and was heavily weighted for larger humeral

femoral epicondyles and more robust ulna as well as increased joint torsion in the cervical joints

(Fig. S2; Table S5). Similarly, the insectivorous landscape resembles the semi-fossorial

landscape with heavy weights towards increased joint torsion and more robust forelimbs. These

results demonstrate the adaptations facilitate not only locomotor behaviors such as swimming

and digging but also dietary ecologies in concert; that is, they need to swim or dig for their prey.

Nevertheless, our functional proxies for feeding consisted of just the mechanical advantage of

jaw closure and thus may not capture the full functional diversity found in carnivorans. The

mammalian skull contains many functional trade-offs such as among bite strength, bite velocity,

and gape size (Herring and Herring 1974; Dumont and Herrel 2003; Slater and Van Valkenburgh

2009; Slater et al. 2009; Forsythe and Ford 2011; Santana 2015). Thus, inclusion of additional

functional proxies from the cranium, mandible, and dentition may uncover further important

contributions of the skull in the evolution of carnivorans (Tseng and Flynn 2018; Slater and

Friscia 2019; Law et al. 2022; Sansalone et al. 2024).

Although biomechanical and ecomorphological studies have linked many of our selected

functional proxies with performance traits (Davis 1964; Boszczyk et al. 2001; Greaves 2012;

Samuels et al. 2013; Jones et al. 2020, 2021), we acknowledge that our analyses were based on

morphological proxies of function rather than empirical performance traits. These may affect the

findings presented in this study. Future work incorporating empirical functional traits in adaptive

landscape analyses requires the continual collection of performance, behavioral, and natural

history data across the entire clade.

Is the carnivoran adaptive landscape relatively flat?

Many carnivoran skeletal components (e.g., mandible, dentition, hindlimb, and post-

diaphragmatic region of the vertebral column) exhibit a short phylogenetic half-life relative to

the age of Carnivora, suggesting that skeletal traits are strongly pulled towards distinct ecological

peaks or clade-based adaptive zones across the adaptive landscape (Slater and Friscia 2019; Law

et al. 2022, 2024; Slater 2022). We found that the overall carnivoran landscape based on

functional proxies from the skull, appendicular, and axial skeletons can be characterized as a

single gradual gradient rather than distinct zones (Fig. 3A). Although some of the performance

surfaces and adaptive landscapes show slight ruggedness and multiple peaks and valleys (Fig. 2–

4), we remain cautious in interpreting these as distinct adaptive zones. The use of empirical data

in creating performance surfaces and adaptive landscapes can lead to heterogeneous resolution of

the interpolation resulting in artificial unevenness in the landscape itself. Our analyses based on

theoretical data are more aligned with our views that the topologies of the carnivoran

performance surfaces and optimized adaptive landscapes are largely characterized as smooth,

gradual gradients with small topographical changes rather than as rugged, multipeak landscapes.

Many-to-one mapping (Wainwright et al. 2005) may explain this decoupling between form and

function. Multiple combinations of morphological traits may lead to the same functional

outcome, resulting in a flat landscape that does not capture the rugged morphological landscape

that was previously hypothesized.

The presence of a relatively flat topology may also indicate that carnivorans already

occupy a broad adaptive zone relative to the overall mammalian adaptive landscape. Although

carnivorans exhibit diverse locomotor modes and correspondingly diverse morphological

adaptations, this diversity does not match the extreme locomotor and morphological

specialization found in other mammalian clades, especially in the appendicular skeleton such as

cranially facing forelimbs in subterranean moles (Lin et al. 2019), digit reduction in cursorial

perissodactyls (Economou et al. 2021), and bipedalism in many saltatorial mammals (McGowan

and Collins 2018). Many of these specialized mammals may be constrained by their highly

derived morphology and thus are adapted to optimize performance for just a single specialized

locomotor behavior. That is, most locomotor modes cannot be maximized simultaneously and

must trade off with other locomotor modes. In contrast, most carnivorans can perform multiple

locomotor behaviors well including running, climbing, digging, and swimming. For example,

even the most cursorial carnivoran, the cheetah, can climb trees whereas no cursorial ungulate

can. Therefore, the relatively flat landscape in carnivorans signals that functional trade-offs

among locomotor performances cannot lead to highly derived specializations, which, in turn,

may lead to rugged, multi-peak landscapes in other mammals. Instead, this carnivoran topology

highlights that the even slight functional trade-offs across smooth, gradual gradients among

appendicular and axial functional proxies can facilitate diverse locomotor modes as well as be

flexible enough to enable additional locomotor behaviors. Future work "zooming out" of the

carnivoran landscape will further elucidate how the relationships among functional trade-offs,

relative degrees of morphological and ecological specializations, and landscape topologies differ

among the various clades across Mammalia.

Acknowledgements

We are grateful to the staff and collections at the American Museum of Natural History,

California Academy of Sciences, Field Museum of Natural History, Natural History Museum of

Los Angeles County, Museum of Vertebrate Zoology, Natural History Museum London, San

Diego Natural History Museum, Texas Vertebrate Paleontology Collection, National Museum of

Natural History, and Burke Museum of Natural History and Culture. We thank Blake Dickson

and Stephanie Pierce for helpful discussions about adaptive landscapes, Miriam Zelditch and

Dave Polly for suggestions on generating theoretical morphological traits from PCAs, and the

Santana lab at UW for helpful discussions.

Funding

Funding was supported by the National Science Foundation (grant no. DBI-2128146) to C.J.L.,

L.J.H., and Z.J.T.; a University of Texas Early Career Provost Fellowship and Stengl-Wyer

Endowment Grant (grant no. SWG-22-02) to C.J.L.; and the European Research Council

(Tied2Teeth, grant no. 101054659) to L.J.H.

Authors' contributions

C.J.L.: conceptualization, data curation, formal analysis, funding acquisition, investigation,

methodology, resources, validation, visualization, writing—original draft, writing—review and

editing; L.J.H.: funding acquisition, investigation, writing— review and editing; Z.J.T.: funding

acquisition, investigation, project administration, writing—review and editing.

Figures

Fig. 1. Phylomorphospace of the carnivoran skeletal system defined by principal components

(PCs) 1 and 2. PCA was conducted using 136 linear and angular measurements that capture

morphological variation across the skull, appendicular, and axial skeletons (Table S1 shows trait

loadings). Linear measurements were size-corrected using log shape ratios. Size of points are

scaled to estimated body size based on the geometric mean of all measurements.

Fig. 2. Performance surfaces for each functional proxy. Color represents height on the

performance surface. See Table 1 for definitions of functional proxies. We removed masMA, SI,

GI, and TRI in subsequent adaptive landscape analyses.

Fig. 3. Adaptive landscapes optimized for all carnivorans and each locomotor group. Landscapes were produced by combining the performance surfaces and optimizing their weightings to maximize the height of the landscape at the group mean. Pie charts show the relative weights of each performance surface on each landscape (Table S3 shows breakdown of weights). Functional proxies with weights > 0.07 were labeled. Table 2 shows statistical tests comparing adaptive landscapes among locomotor groups. Size of points are scaled to estimated body size based on the geometric mean of all measurements.

Fig. 4. Adaptive landscapes optimized for all carnivorans and each family. Landscapes were

produced by combining the performance surfaces and optimizing their weightings to maximize

the height of the landscape at the group mean. Pie charts show the relative weights of each

performance surface on each adaptive landscape (Table S6 shows breakdown of weights).

Functional proxies with weights > 0.09 were labeled. Table S7 shows statistical tests comparing

adaptive landscapes among families. Size of points are scaled to estimated body size based on

the geometric mean of all measurements.

613 **Tables**

614 **Table 1.** Functional proxies capturing the functional diversity of the skull (Greaves 2012), limbs

615 (Davis 1964; Samuels et al. 2013), and vertebral column (Boszczyk et al. 2001; Jones et al. 2020,

616 2021). The full performance, biomechanical, and/or ecomorphological justification for each

617 functional proxy is expanded upon in the Supplementary Materials.

Functional proxy	Description
Skull	
temporalis mechanical advantage (temMA)	Estimates how much force is produced at the bite point from force being input by the temporalis muscle
masseter mechanical advantage (masMA)	Estimates how much force is produced at the bite point from force being input by the masseter muscle
Forelimb	
Scapula index (SI)	Describes the expansion of shoulder musculature versus contribution of scapula to limb elongation.
Brachial index (BI)	Estimates the relative proportions of the proximal and distal elements of the forelimb and serves as an index of the relative distal out-lever length.
Humeral robustness index (HRI)	Estimates the robustness of the humerus and its ability to resist bending and shearing stresses.
Humeral epicondylar index (HEI)	Estimates the relative area of the distal end of the humerus available for the origins of the forearm flexors, pronators, and supinators.
Olecranon length index (OLI)	Estimates the relative mechanical advantage of the triceps brachii and dorsoepitrochlearis muscles used in elbow extension.
Ulnar robustness index (URI)	Estimates the robustness of the ulna and its ability to resist bending and shearing stresses, and relative area available for the origin and insertion of forearm and manus flexors, pronators, and supinators.
Manus proportions index (MAN)	Estimates the relative proportions of proximal and distal elements of the forelimb, and relative size of the hand.
Hindlimb	
Crural index (CI)	Estimates relative proportions of proximal and distal elements of the hind limb.

Femoral robustness index (FRI)	Estimates robustness of the femur and its ability to resist bending and shearing stresses.
Gluteal index (GI)	Estimates the relative mechanical advantage of the gluteal muscles used in retraction of the femur.
Femoral epicondylar index (FEI)	Estimates relative area available for the origins of the gastrocnemius and soleus muscles used in extension of the knee and plantar-flexion of the pes.
Tibial robustness index (TRI)	Estimates robustness of the tibia and its ability to resist bending and shearing stresses.
Pes length index (PES)	Estimates relative proportions of proximal and distal elements of the hind limb, and relative size of the hind foot.
Vertebrae	
Sagittal second moment of area (sSMA)	Estimates stiffness in the of the vertebral joint in the sagittal plane.
Lateral second moment of area (lSMA)	Estimates stiffness in the of the vertebral joint in the lateral plane.
Joint torsional angle (JTA)	Estimates the degree of axial torsion of the vertebrae.
Joint verticality (JV)	Estimates the relative importance of sagittal bending versus lateral bending of vertebral joints.

623 **Table 2.** Pairwise significance tests among locomotor adaptive landscapes. Top triangle: number

624 of landscape models shared in the top 5% between the paired groups. Bottom triangle: P-values

625 for difference between groups. Bolded P-values indicate significance.

	arboreal	cursorial	scansorial	semi-aquatic	semi-fossorial	terrestrial hunter	terrestrial non-hunter
arboreal	-	4	7	0	0	6	6
cursorial	0.100	-	10	0	0	7	34
scansorial	0.800	0.077	-	0	0	14	15
semi-aquatic	**0.001**	**0.001**	**0.001**	-	88	0	0
semi-fossorial	**0.001**	**0.001**	**0.001**	0.184	-	0	0
terrestrial hunter	0.500	0.231	0.417	**0.001**	**0.001**	-	12
terrestrial nonhunter	**0.001**	0.077	**0.001**	**0.001**	0.174	**0.001**	-

References

Adams DC, Collyer ML. 2018. Phylogenetic ANOVA: Group-clade aggregation, biological
challenges, and a refined permutation procedure. Evolution 72:1204–15.

Amson E, Bibi F. 2021. Differing effects of size and lifestyle on bone structure in mammals.
BMC Biol 19:87.

Arbour JH, Curtis AA, Santana SE. 2019. Signatures of echolocation and dietary ecology in the
adaptive evolution of skull shape in bats. Nat Commun 10:2036.

Arnold SJ. 1983. Morphology, performance and fitness. Amer Zool 23:347–61.

Arnold SJ, Pfrender ME, Jones AG. 2001. The adaptive landscape as a conceptual bridge
between micro- and macroevolution. Genetica 112–113:9–32.

Bastide P, Ané C, Robin S, Mariadassou M. 2018. Inference of Adaptive Shifts for Multivariate
Correlated Traits. Syst Biol 67:662–80.

Beaulieu JM, Jhwueng D-C, Boettiger C, O'Meara BC. 2012. Modeling stabilizing selection:
Expanding the Ornstein-Uhlenbeck model of adaptive evolution. Evolution 66:2369–83.

Belyaev RI, Kuznetsov AN, Prilepskaya NE. 2021. How the even□toed ungulate vertebral
column works: Comparison of intervertebral mobility in 33 genera. J Anat 239:1370–99.

Belyaev RI, Kuznetsov AN, Prilepskaya NE. 2022. From dorsomobility to dorsostability: A
study of lumbosacral joint range of motion in artiodactyls. J Anat 241:420–36.

Belyaev RI, Nikolskaia P, Bushuev AV, Panyutina AA, Kozhanova DA, Prilepskaya NE. 2023.
Running, jumping, hunting, and scavenging: Functional analysis of vertebral mobility and
backbone properties in carnivorans. J Anat.

Boszczyk BM, Boszczyk AA, Putz R. 2001. Comparative and functional anatomy of the
mammalian lumbar spine. Anatomical Rec 264:157–68.

Butler MA, King AA. 2004. Phylogenetic comparative analysis: a modeling approach for
adaptive evolution. Am Nat 164:683–95.

Casinos A, Quintana C, Viladiu C. 1993. Allometry and adaptation in the long bones of a
digging group of rodents (Ctenomyinae). Zoo□l J Linn Soc 107:107–15.

Clavel J, Aristide L, Morlon H. 2019. A Penalized Likelihood Framework for High-Dimensional Phylogenetic Comparative Methods and an Application to New-World Monkeys Brain Evolution. Systematic Biol 68:93–116.

Clavel J, Morlon H. 2020. Reliable Phylogenetic Regressions for Multivariate Comparative Data: Illustration with the MANOVA and Application to the Effect of Diet on Mandible Morphology in Phyllostomid Bats. Syst Biol 69:927–43.

Collar DC, Reece JS, Alfaro ME, Wainwright PC. 2014. Imperfect morphological convergence: variable changes in cranial structures underlie transitions to durophagy in moray eels. Am Nat 183:E168–84.

Davis DD. 1964. The giant panda: a morphological study of evolutionary mechanisms Fieldiana (Zoology Memoirs).

Dickson BV, Clack JA, Smithson TR, Pierce SE. 2021. Functional adaptive landscapes predict terrestrial capacity at the origin of limbs. Nature 589:242–45.

Dickson BV, Pierce SE. 2019. Functional performance of turtle humerus shape across an ecological adaptive landscape. Evolution 73:1265–77.

Dumont ER, Herrel A. 2003. The effects of gape angle and bite point on bite force in bats. J Exp Biol 206:2117–23.

Economou G, McGrath M, Wajsberg J, Granatosky MC. 2021. Perissodactyla Locomotion. In: Vonk J, Shackelford TK, editors. Encyclopedia of Animal Cognition and Behavior Switzerland: Springer Nature. p. 1–8.

Figueirido B, MacLeod N, Krieger J, Renzi MD, Pérez-Claros JA, Palmqvist P. 2011. Constraint and adaptation in the evolution of carnivoran skull shape. Paleobiology 37:490–518.

Figueirido B, Martín-Serra A, Pérez-Ramos A, Velasco D, Pastor FJ, Benson RJ. 2021. Serial disparity in the carnivoran backbone unveils a complex adaptive role in metameric evolution. Communications Biology 1–15.

Fish FE. 1994. Association of propulsive swimming mode with behavior in river otters (Lutra canadensis). J Mammal 75:989–97.

Fish FE. 2000. Biomechanics and Energetics in Aquatic and Semiaquatic Mammals: Platypus to Whale*. Physiol Biochem Zool 73:683–98.

Fish FE, Hurley J, Costa DP. 2003. Maneuverability by the sea lion Zalophus californianus: turning performance of an unstable body design. J Exp Biol 206:667–74.

Forsythe EC, Ford SM. 2011. Craniofacial Adaptations to Tree☐Gouging Among Marmosets. Anat Rec: Adv Integr Anat Evol Biol 294:2131–39.

Friedman ST, Price SA, Hoey AS, Wainwright PC. 2016. Ecomorphological convergence in
planktivorous surgeonfishes. J Evol Biol 29:965–78.

Friscia AR, Van Valkenburgh B, Biknevicius AR. 2007. An ecomorphological analysis of extant
small carnivorans. J Zool 272:82–100.

Gál JM. 1993. Mammalian Spinal Biomechanics: I. Static and Dynamic Mechanical Properties
of Intact Intervertebral Joints. J Exp Biol 174:247–80.

Gambaryan PP. 1974. How Mammals Run: Anatomical Adaptations J. Wiley.

Garland T, Janis CM. 1993. Does metatarsal/femur ratio predict maximal running speed in
cursorial mammals? J Zool 229:133–51.

Greaves WS. 2012. The Mammalian Jaw. 3–25.

Hansen TF. 1997. Stabilizing selection and the comparative analysis of adaptation. Evolution
51:1341–51.

Hedrick BP, Dickson BV, Dumont ER, Pierce SE. 2020. The evolutionary diversity of locomotor
innovation in rodents is not linked to proximal limb morphology. Scientific Reports 1–11.

Herring SW, Herring SE. 1974. The Superficial Masseter and Gape in Mammals. Am Nat
108:561–76.

Higham TE, Ferry LA, Schmitz L, Irschick DJ, Starko S, Anderson PSL, Bergmann PJ,
Jamniczky HA, Monteiro LR, Navon D, Messier J, Carrington E, Farina SC, Feilich KL,
Hernandez LP, Johnson MA, Kawano SM, Law CJ, Longo SJ, Martin CH, Martone PT, Rico-
Guevara A, Santana SE, Niklas KJ. 2021. Linking ecomechanical models and functional traits
to understand phenotypic diversity. Trends in Ecology & Evolution 1–14.

Hildebrand M. 1959. Motions of the Running Cheetah and Horse. J Mammal 40:481–95.

Hildebrand M. 1985a. Digging of Quadrupeds. In: Hildebrand M, Bramble DM, Liem KF, Wake
DB, editors. Functional Vertebrate Morphology Cambridge, MA: Harvard University Press.
p. 89–109.

Hildebrand M. 1985b. Walking and Running. In: Hildebrand M, Bramble DM, Liem KF, Wake
DB, editors. Functional Vertebrate Morphology Cambridge, MA: Harvard University Press.
p. 38–57.

Humphreys AM, Barraclough TG. 2014. The evolutionary reality of higher taxa in mammals.
Proc Royal Soc B Biological Sci 281:20132750.

Iwaniuk AN, Pellis SM, Whishaw IQ. 1999. The relationship between forelimb morphology and
behaviour in North American carnivores (Carnivora). Can J Zool 77:1064–74.

Jones KE, Benitez L, Angielczyk KD, Pierce SE. 2018. Adaptation and constraint in the evolution of the mammalian backbone. Bmc Evol Biol 18:172.

Jones KE, Dickson BV, Angielczyk KD, Pierce SE. 2021. Adaptive landscapes challenge the "lateral-to-sagittal" paradigm for mammalian vertebral evolution. Curr Biol 31:1883-1892.e7.

Jones KE, Gonzalez S, Angielczyk KD, Pierce SE. 2020. Regionalization of the axial skeleton predates functional adaptation in the forerunners of mammals. Nat Ecol Evol 4:470–78.

Kilbourne BM. 2017. Selective regimes and functional anatomy in the mustelid forelimb: Diversification toward specializations for climbing, digging, and swimming. Ecol Evol 66:2369–12.

Kilbourne BM, Hoffman LC. 2013. Scale Effects between Body Size and Limb Design in Quadrupedal Mammals. PLoS ONE 8:e78392.

Kram R, Taylor CR. 1990. Energetics of running: a new perspective. Nature 346:265–67.

Lagaria A, Youlatos D. 2006. ANATOMICAL CORRELATES TO SCRATCH DIGGING IN THE FORELIMB OF EUROPEAN GROUND SQUIRRELS (SPERMOPHILUS CITELLUS). J Mammal 87:563–70.

Law CJ. 2019. Solitary meat-eaters: solitary, carnivorous carnivorans exhibit the highest degree of sexual size dimorphism. Sci Rep-uk 9:15344.

Law CJ. 2021a. Ecological drivers of carnivoran body shape evolution. Am Nat 198:406–20.

Law CJ. 2021b. Evolutionary and morphological patterns underlying carnivoran body shape diversity. Evolution 75:365–75.

Law CJ. 2022. Different Evolutionary Pathways Lead to Incomplete Convergence of Elongate Body Shapes in Carnivoran Mammals. Systematic Biol 71:788–96.

Law CJ, Blackwell EA, Curtis AA, Dickinson E, Hartstone Rose A, Santana SE. 2022. Decoupled evolution of the cranium and mandible in carnivoran mammals. Evolution 76:2959–74.

Law CJ, Duran E, Hung N, Richards E, Santillan I, Mehta RS. 2018. Effects of diet on cranial morphology and biting ability in musteloid mammals. J Evol Biol 31:1918–31.

Law CJ, Hlusko LJ, Tseng ZJ. 2024. Uncovering the mosaic evolution of the carnivoran skeletal system. Biol Lett 20:20230526.

Law CJ, Slater GJ, Mehta RS. 2019. Shared extremes by ectotherms and endotherms: Body elongation in mustelids is associated with small size and reduced limbs. Evolution 73:735–49.

Lessa EP, Stein BR. 1992. Morphological constraints in the digging apparatus of pocket gophers (Mammalia: Geomyidae). Biol J Linn Soc 47:439–53.

Lin Y-F, Konow N, Dumont ER. 2019. How moles destroy your lawn: the forelimb kinematics of eastern moles in loose and compact substrates. J Exp Biol 222:jeb182436.

Marshall SK, Spainhower KB, Sinn BT, Diggins TP, Butcher MT. 2021. Hind Limb Bone Proportions Reveal Unexpected Morphofunctional Diversification in Xenarthrans. J Mamm Evol 28:599–619.

Martín-Serra A, Figueirido B, Palmqvist P. 2014a. A three-dimensional analysis of the morphological evolution and locomotor behaviour of the carnivoran hind limb. BMC Evol Biol 14:1–13.

Martín-Serra A, Figueirido B, Palmqvist P. 2014b. A Three-Dimensional Analysis of Morphological Evolution and Locomotor Performance of the Carnivoran Forelimb. Plos One 9:e85574.

Martín-Serra A, Pérez-Ramos A, Pastor FJ, Velasco D, Figueirido B. 2021. Phenotypic integration in the carnivoran backbone and the evolution of functional differentiation in metameric structures. Evolution Letters 1–14.

McGowan CP, Collins CE. 2018. Why do mammals hop? Understanding the ecology, biomechanics and evolution of bipedal hopping. J Exp Biol 221:jeb161661.

Muñoz NA. 2020. Locomotion in Rodents and Small Carnivorans: Are they So Different? Journal of Mammalian Evolution 1–12.

Nakai D, Fujiwara S. 2023. Fossorial mammals emphasise the forelimb muscle moment arms used for digging: New indices for reconstruction of the digging ability and behaviours in extinct taxa. J Anat 242:846–61.

Nations JA, Heaney LR, Demos TC, Achmadi AS, Rowe KC, Esselstyn JA. 2019. A simple skeletal measurement effectively predicts climbing behaviour in a diverse clade of small mammals. Biol J Linn Soc 128:323–36.

Oliver JD, Jones KE, Hautier L, Loughry WJ, Pierce SE. 2016. Vertebral bending mechanics and xenarthrous morphology in the nine-banded armadillo (Dasypus novemcinctus). J Exp Biol 219:2991–3002.

Polly PD. 2007. Limbs in mammalian evolution. In: Hall BK, editor. Fins into Limbs Evolution, Development, and Transformation. Fins into Limbs Evolution, Development, and Transformation Fins into Limbs Evolution, Development, and Transformation. p. 245–68.

Polly PD, Lawing AM, Fabre A-C, Goswami A. 2013. Phylogenetic principal components analysis and geometric morphometrics. Hystrix, the Italian Journal of Mammalogy 24:33–41.

Polly PD, Stayton CT, Dumont ER, Pierce SE, Rayfield EJ, Angielczyk KD. 2016. Combining Geometric Morphometrics and Finite Element Analysis with Evolutionary Modeling: Towards a Synthesis. J Vertebr Paleontol 36:e1111225.

Pontzer H. 2007a. Effective limb length and the scaling of locomotor cost in terrestrial animals. J Exp Biol 210:1752–61.

Pontzer H. 2007b. Predicting the energy cost of terrestrial locomotion: a test of the LiMb model in humans and quadrupeds. J Exp Biol 210:484–94.

Price SA, Hopkins SSB. 2015. The macroevolutionary relationship between diet and body mass across mammals. Biol J Linn Soc 115:173–84.

Radinsky LB. 1981. Evolution of skull shape in Carnivores .1. Representative modern Carnivores. Biol J Linn Soc 15:369–88.

Randau M, Cuff AR, Hutchinson JR, Pierce SE, Goswami A. 2017. Regional differentiation of felid vertebral column evolution: a study of 3D shape trajectories. Org Divers Evol 17:1–15.

Rickman J, Burtner AE, Linden TJ, Santana SE, Law CJ. 2023. Size And Locomotor Ecology Have Differing Effects on the External and Internal Morphologies of Squirrel (Rodentia: Sciuridae) Limb Bones. Integr Org Biol 5:obad017.

Rose J, Moore A, Russell A, Butcher M. 2014. Functional osteology of the forelimb digging apparatus of badgers. J Mammal 95:543–58.

Samuels JX, Meachen JA, Sakai SA. 2013. Postcranial morphology and the locomotor habits of living and extinct carnivorans. J Morpho 274:121–46.

Samuels JX, Van Valkenburgh B. 2008. Skeletal indicators of locomotor adaptations in living and extinct rodents. J Morpho 269:1387–1411.

Sansalone G, Castiglione S, Raia P, Archer M, Dickson B, Hand S, Piras P, Profico A, Wroe S. 2020. Decoupling Functional and Morphological Convergence, the Study Case of Fossorial Mammalia. Frontiers Earth Sci 8:112.

Sansalone G, Wroe S, Coates G, Attard MRG, Fruciano C. 2024. Unexpectedly uneven distribution of functional trade-offs explains cranial morphological diversity in carnivores. Nat Commun 15:3275.

Santana SE. 2015. Quantifying the effect of gape and morphology on bite force: biomechanical modelling and in vivomeasurements in bats. Funct Ecol 30:557–65.

Santana SE, Dumont ER, Dumont E, Davis JL. 2010. Mechanics of bite force production and its relationship to diet in bats. Funct Ecol 24:776–84.

Schilling N, Hackert R. 2006. Sagittal spine movements of small therian mammals during
asymmetrical gaits. J Exp Biol 209:3925–39.

Slater GJ. 2022. Topographically distinct adaptive landscapes for teeth, skeletons, and size
explain the adaptive radiation of Carnivora (Mammalia). Evolution.

Slater GJ, Dumont E, Van Valkenburgh B. 2009. Implications of predatory specialization for
cranial form and function in canids. J Zool 278:181–88.

Slater GJ, Friscia AR. 2019. Hierarchy in adaptive radiation: A case study using the Carnivora
(Mammalia). Evolution 73:524–39.

Slater GJ, Van Valkenburgh B. 2009. Allometry and performance: the evolution of skull form
and function in felids. Journal of Evolutionary Biology 22:2278–87.

Smith SM, Stayton CT, Angielczyk KD. 2021. How many trees to see the forest? Assessing the
effects of morphospace coverage and sample size in performance surface analysis. Methods
Ecol Evol 12:1411–24.

Stayton CT. 2019. Performance in three shell functions predicts the phenotypic distribution of
hard☐shelled turtles. Evolution 73:720–34.

Strang KT, Steudel K. 1990. Explaining the scaling of transport costs: the role of stride
frequency and stride length. J Zoo☐l 221:343–58.

Sustaita D, Pouydebat E, Manzano A, Abdala V, Hertel F, Herrel A. 2013. Getting a grip on
tetrapod grasping: form, function, and evolution. Biol Rev 88:380–405.

Tseng ZJ, Flynn JJ. 2018. Structure-function covariation with nonfeeding ecological variables
influences evolution of feeding specialization in Carnivora. Sci Adv 4:eaao5441.

Tseng ZJ, Garcia-Lara S, Flynn JJ, Holmes E, Rowe TB, Dickson BV. 2023. A switch in jaw
form–function coupling during the evolution of mammals. Philos Trans R Soc B
378:20220091.

Uyeda JC, Harmon LJ. 2014. A novel Bayesian method for inferring and interpreting the
dynamics of adaptive landscapes from phylogenetic comparative data. Syst Biol 63:902–18.

Uyeda JC, Zenil-Ferguson R, Pennell MW. 2018. Rethinking phylogenetic comparative
methods. Syst Biol 67:1091–1109.

Van Valkenburgh B. 1985. Locomotor diversity within past and present guilds of large predatory
mammals. Paleobiology 11:406–28.

Van Valkenburgh B. 1987. Skeletal indicators of locomotor behavior in living and extinct
carnivores. J Vertebr Paleontol 7:162–82.

Van Valkenburgh B. 2007. Deja vu: the evolution of feeding morphologies in the Carnivora.
Amer Zool 47:147–63.

Wainwright PC. 1994. Functional morphology as a tool in ecological research. In: Wainwright
PC, Reilly SM, editors. Ecological Morphology Integrative Organismal Biology. Ecological
Morphology Integrative Organismal Biology Ecological Morphology: Integrative ….

Wainwright PC, Alfaro ME, Bolnick DI, Hulsey CD. 2005. Many-to-one mapping of form to
function: a general principle in organismal design? Integr Comp Biol 45:256–62.

Young JW. 2023. Convergence of arboreal locomotor specialization: morphological and
behavioral solutions for movement on narrow and compliant supports. In: Convergent
evolution: Animal form and function. Fascinating Life Sciences Springer International
Publishing. p. 289–322.

Zelditch ML, Ye J, Mitchell JS, Swiderski DL. 2017. Rare ecomorphological convergence on a
complex adaptive landscape: Body size and diet mediate evolution of jaw shape in squirrels
(Sciuridae). Evolution 71:633–49.

868