1

TRPγ regulates lipid metabolism through Dh44 neuroendocrine cells

Dharmendra Kumar Nath[1,#], Subash Dhakal[1,#], and Youngseok Lee[1*]

[1]Department of Bio and Fermentation Convergence Technology, Kookmin University, Seoul, 02707, Republic of Korea

[#] These authors contributed equally to this work

*Correspondence and requests for materials should be addressed to:

Y.L. (email: ylee@kookmin.ac.kr; ORCID: 0000 0003 0459 1138)

ABSTRACT

Understanding how the brain controls nutrient storage is pivotal. Transient receptor potential (TRP) channels are conserved from insects to humans. They serve in detecting environmental shifts and in acting as internal sensors. Here, we found that a TRPγ mutant exhibited in *Drosophila melanogaster* are required for maintaining normal lipid and protein levels. In animals, lipogenesis and lipolysis control lipid levels in response to food availability. Lipids are mostly stored as triacylglycerol in the fat bodies (FBs) of *D. melanogaster*. Interestingly, *trpγ* deficient mutants exhibited elevated TAG levels and our genetic data indicated that *Dh44* neurons are indispensable for normal lipid storage but not protein storage. The *trpγ* mutants also exhibited reduced starvation resistance, which was attributed to insufficient lipolysis in the FBs. This could be mitigated by administering lipase or metformin orally, indicating a potential treatment pathway. Gene expression analysis indicated that *trpγ* knockout downregulated *brummer*, a key lipolytic gene, resulting in chronic lipolytic deficits in the gut and other fat tissues. The study also highlighted the role of specific proteins, including neuropeptide DH44 and its receptor DH44R2 in lipid regulation. Our findings provide insight into the broader question of how the brain regulates nutrient storage.

Keywords: TRP channel, Dh44, metabolism, lipid

INTRODUCTION

Nutrient storage is not a simple matter of accumulation but involves a sophisticated network of signals and receptors, feedback loops, and hormonal influences, all choreographed by the brain. The mechanisms involve various tissues and organs, including the liver, muscles, and adipose tissue, each playing a distinct role in the storage and release of vital nutrients like glucose, fats, and proteins in mammals [1]. Key to this regulation are specialized cells and molecular pathways that respond to dietary intake, energy expenditure, and changes in the internal and external environment [2]. For instance, the brain receives and interprets signals about the body's energy status through hormones such as insulin and leptin, and nutrients themselves, which inform decisions on whether to store or mobilize energy reserves in mammals [3,4]. Moreover, the role of Transient Receptor Potential (TRP) channels as molecular sensors that detect changes in the environment and internal metabolic status highlights the level of molecular sophistication involved in nutrient storage and energy balance [5-8]. These channels bridge external stimuli with internal responses, ensuring the body adapts efficiently to maintain homeostasis [9]. This concert of mechanisms, from the molecular to the systemic level, exemplifies the elegance with which the brain ensures our energy needs are met in a changing environment, highlighting the intricate interplay of biological process that sustains life [10]. Understanding these processes opens up new vistas in our approach to diet, disease management, and overall health optimization.

The use of a genetically tractable model organism such as *Drosophila melanogaster* could provide fundamental insights into the potential relationship between

TRP channels and metabolic syndrome. This study delves into the function of TRPγ

channels, specifically their role in managing lipid and protein levels. Through genetic

manipulation, we explored the consequences of TRPγ mutations on the metabolic

processes, revealing critical insights into the regulation of lipid storage and breakdown.

Our investigation sheds light on the balance between lipogenesis and lipolysis,

processes that respectively synthesize and degrade lipids based on the availability of

food, and how these processes are affected by TRPγ activity. By examining the

metabolic anomalies in *trpγ* deficient mutants, including alterations in lipid storage and

the consequent impact on starvation resistance, this study advances our understanding

of metabolic regulation. Furthermore, we uncover the potential therapeutic benefits of

lipase and metformin in counteracting the metabolic deficiencies caused by TRPγ

mutations. Through detailed gene expression analysis, we identify the downregulation

of crucial genes involved in lipid metabolism, offering new perspectives on the genetic

underpinnings of nutrient storage regulation. The implications of this research are far-

reaching, not only enhancing our comprehension of basic biological functions but also

providing avenues for therapeutic interventions in metabolic disorders.

79 RESULTS

TRPγ mutants exhibit reduced sugar level alongside elevated lipid and protein

levels

In order to understand how *D. melanogaster* regulates the maintenance of major

nutrient levels, including carbohydrates, lipids, and proteins, TRP channel mutants were

examined as potential candidates. A recent study revealed that among these mutants,

the *trpγ* mutant displayed only reduced carbohydrate levels [11], suggesting the

involvement of additional TRP channels in regulating major nutrients. *D. melanogaster*

possesses 13 members of the TRP channel family. Flies with mutations in *trpM*, *trpML*,

and *nompC* showed high mortality rates with minimal survival, whereas those with

mutations in the remaining 10 TRP superfamily genes were viable and healthy when

homozygous. Investigation into the lipid and protein levels of other available TRP

superfamily mutants revealed that, apart from the *trpγ* mutant, all tested mutants

exhibited normal levels (**Figure 1A and B**). Specifically, the *trpγ^1* mutant showed 1.3

times higher triacylglycerol (TAG) levels and 1.5 times higher protein levels compared to

the control. These metabolic changes were unique and specific to the *trpγ* mutants

among the evaluated TRP superfamily mutants. Previous observations indicated that

trpγ mutants had lower cellular sugar levels and stored glycogen. Consistently, *trpγ*

mutants exhibited reduced levels of cellular sugars (glucose and trehalose), stored

glycogen, and sugar levels in the hemolymph (**Figure 1C–E**). In conclusion, it was

found that *trpγ^1* mutants displayed lower carbohydrate levels but higher lipid and protein

levels compared to the control. Importantly, the increased TAG and protein levels

observed in *trpγ^1* mutants were confirmed with a second *trpγ^1* allele (*trpγ^{G4}*) (**Figure 1F**

and G). Furthermore, we restored the deficiency in lipid and protein levels using a

genomic *trpγ$^+$* transgene, *g(trpγ)*, and by introducing a *UAS-trpγ* cDNA, controlled by

GAL4, into the *trpγ* locus (*trpγ^{G4}* flies; +).

TRPγ regulates lipid metabolism through Dh44 neuroendocrine cells in the pars

intercerebralis

We recently proposed that TRPγ expression in the six DH44 neuroendocrine cells in the

pars intercerebralis (PI) located in the dorsal medial area of the brain is essential to

maintain normal carbohydrate levels in tissues (*Figure 2–figure supplement 1A1–A3*)

[11]. Therefore, we next sought to determine whether protein and lipid levels can be

regulated by the same neuroendocrine cells. To test this, we used a transgenic fly

encoding an inwardly rectifying K$^+$ channel (*UAS-Kir2.1*) to inactivate specific neurons

(*Figure 2A and B*) [12]. Interestingly, inactivating the *Dh44* neurons significantly

increased the lipid levels but not protein levels (*Figure 2A and B*). Approximately 16

neurons in the PI express dILP2, which are also overlapped by two DH44-positive

neurons in 5 to 10 days old adult male flies (*Figure 2–figure supplement 1B1–B3*) [13].

However, inactivation of *dILP2* neurons did not impair either lipid or protein levels

(*Figure 2A and B*). In conclusion, our findings indicated that DH44 neuroendocrine

cells contribute to lipid regulation but not protein regulation.

To assess whether *trpγ* expression in *Dh44* neurons is sufficient to restore

normal lipid and protein levels in *trpγ* mutant flies, we expressed the *UAS-trpγ* under the

control of *Dh44-GAL4* in the *trpγ^1* mutant background (*Figure 2C and D*). Our findings

indicated that the increased lipid levels in the *trpγ^1* mutant background decreased to

normal levels via the expression of the *trpγ* transgene in the DH44 neuroendocrine cells

but not in its parent strains (*Figure 2C*). Again, the expression of the *trpγ* transgene in

the dILP2 neurons had no appreciable effects (*Figure 2C*). In contrast, the expression

of the *trpγ* transgene did not decrease the protein levels in either the DH44 or the dILP2

neurons (***Figure 2D***). This indicates that TRPγ is sufficient for the regulation of lipid

levels in the DH44 cells. Next, RNAi-mediated knockdown experiments were conducted

to further examine the role of *trpγ* in *Dh44* neurons. Interestingly, *trpγ* knockdown in

Dh44 neurons significantly increased lipid levels, whereas flies harboring only the *Dh44-*

GAL4 or the *UAS-trpγ*[RNAi] transgenes displayed normal lipid levels (***Figure 2E***). This

indicated that TRPγ is needed for the regulation of lipid levels. Furthermore, the total

TAG level in the *trpγ*[1] flies was higher in both males and females (***Figure 2–figure***

supplement 1C), meaning that the functions of *trpγ* are not sex-specific.

In *D. melanogaster*, lipids are mainly stored in the form of TAG and cholesterol

ester in the adipose tissue [i.e., fat bodies (FBs)] as lipid droplets (LDs) [14]. Consistent

with the increased TAG in tissues, Nile red staining of the FBs of *trpγ*[1] and *trpγ*[G4] flies

exhibited larger lipid mass compared to control animals (***Figure 2F–I***). These LDs

returned to their normal size through the expression of *trpγ* in *Dh44* neurons but not

dILP2 neurons (***Figure 2F and 2J–L***). Additionally, inactivating the *Dh44* cells (*Dh44-*

GAL4/*UAS-Kir2.1*) recapitulated the enlarged LD phenotype of the *trpγ* mutant flies

(***Figure 2–figure supplement 1D–G***). Overall, our findings indicated that *trpγ*

expression regulates lipid and carbohydrate homeostasis but not protein levels in DH44

neuroendocrine cells in the PI.

TRPγ mutants exhibit starvation susceptibility and deficits in lipolysis

Higher lipid cellular levels may decrease lifespan [15,16]. Therefore, we measured the

lifespan of control and *trpγ[1]* flies fed with a standard cornmeal diet (**Figure 3A**).

However, the lifespans of control and *trpγ[1]* flies were not significantly different under

normal conditions. The LT_{50} of the control was 59.34 ± 0.92 days and that of *trpγ[1]* was

55.75 ± 2.35 days. In contrast, *trpγ* appeared to be required for proper metabolism

under starvation conditions, as demonstrated by the decreased starvation resistance of

the *trpγ[1]* and *trpγ[RNAi]* knockdown flies in *Dh44* cells (**Figure 3B**) [11]. This defect was fully

recovered by the expression of *UAS-trpγ* under the control of *Dh44-GAL4* (**Figure 3B**).

The starvation sensitive phenotype in the *trpγ* mutants may have been due to

decreased carbohydrate storage in tissues, including glucose, trehalose, and glycogen.

However, we previously demonstrated that *trpγ* mutants could utilize carbohydrates

under starved conditions [11]. Furthermore, the elevated whole-body TAG levels in the

trpγ mutant flies might suggest that they were unable to break down stored lipids even

under starving conditions. Metazoans must coordinate the metabolism of glycogen, lipid,

and protein to maintain metabolic homeostasis during fasting periods, thus providing an

appropriate energy supply across tissues. Therefore, the total TAG levels of the control

and *trpγ* mutant flies were assessed under sated and starved (starvation for 24 h)

conditions (**Figure 3C–F)**. When the control flies were deprived of food for 24 h, their

TAG levels decreased significantly (**Figure 3C**). In contrast, the *trpγ[1]* mutants exhibited

no changes in whole-body TAG levels before and after starvation (**Figure 3C**). To

further confirm that lipolysis was restricted in the *trpγ* mutants under starvation

conditions, LDs were stained, after which we measured the sizes of individual LDs

accumulated throughout the whole fat bodies of the flies (**Figure 3D–F**). Despite

considerable variations, the sizes of the LDs in the control flies were significantly

reduced after starvation. In contrast, no significant differences in LD sizes were

identified between the sated and starved conditions in the $trp\gamma^1$ flies. We examined how

overexpression and knockdown of $trp\gamma$ in $Dh44$ neurons affect the starvation phenotype.

Overexpressing $trp\gamma$ in $Dh44$ cells resulted in similarity to the wild-type in both sated and

starved conditions, as well as normal survival time under starvation conditions (**Figure**

3B and Figure 3–figure supplement 1A). Conversely, $trp\gamma^{RNAi}$ knockdown flies in $Dh44$

neurons reproduced phenotypic traits observed in TRPγ mutants, including decreased

lipid levels (**Figure 3–figure supplement 1B**) and reduced survival time under

starvation conditions (**Figure 3B**).

Adenosine monophosphate-activated protein kinase (AMPK) serves as the

master controller for maintaining energy balance in cells, coordinating metabolic

pathways [17]. It regulates the balance between building up and breaking down

substances to ensure cellular stability during metabolic stress. AMPK is a key target for

treating metabolic diseases like type 2 diabetes and obesity, as its activation increases

fatty acid oxidation [18]. We wonder if activating AMPK in $Dh44$ neurons serves as a

rescue mechanism for lipolysis. The expression of AMPK^{TD}, activated form of AMPK [19]

in $Dh44$ neurons indeed restored the elevated TAG levels observed in $trp\gamma^1$ (**Figure 3G**).

This suggests that AMPK functions as a downstream component of $Dh44$ neuronal

activation.

Recovery of the starvation resistance of the TRPγ mutant via metformin treatment

Metformin is widely used to treat many metabolic diseases such as type II diabetes [20].

We previously demonstrated that 1 to 5 mM metformin can induce hypoglycemia, in

addition to suppressing fat storage in flies [21]. Therefore, we next sought to test whether

the increased lipid levels of the *trpγ* mutant can be reduced to a normal level via oral

administration of metformin. First, we measured TAG levels at 0, 7, and 14 days after

dietary administration of 1 mM or 5 mM of metformin (the treatments were prepared by

mixing the appropriate metformin concentrations into standard cornmeal diets) (**Figure**

4A and B). The high TAG levels in the *trpγ* mutant (41.91 ±4.56) were decreased to

levels similar to those of the control at 7 and 14 days (30.03 ±4.34 and 33.79 ±3.20,

respectively) after treatment with 1 mM metformin (**Figure 4A**). Furthermore, although

oral administration of 1 mM metformin did not affect the TAG level in the control, the 5

mM metformin treatment significantly reduced the TAG level of the control flies at 14

days (**Figure 4B**). Moreover, the 5 mM metformin treatment was more effective in

reducing the TAG level in the *trpγ* mutant. These findings indicated that oral

administration of metformin can either suppress lipogenesis or enhance lipolysis in *D.*

melanogaster. To further confirm the effects of metformin treatment, LDs were analyzed

under the same condition (**Figure 4C and D**). The sizes of the LDs in the control and

trpγ mutant flies were significantly reduced after treatment with 1 mM metformin at 14

days or 5 mM metformin at 7 and 14 days (**Figure 4D**).

Next, we sought to assess whether metformin could increase starvation

resistance in the control and *trpγ* mutant flies. To test this hypothesis, the starvation

resistance of the flies was measured with or without metformin. Neither of the metformin

concentrations appeared to affect the starvation resistance of the control flies (**Figure**

4E and F). However, although metformin did not restore the starvation resistance of the

trpγ mutant flies to the level of the controls, it did significantly extend their survival

(**Figure 4E and F**; LT_{50} of the control at 1 mM and 5 mM: 56.06 ±1.40 h and 52.39

±2.09 h, respectively; LT_{50} of the $trpγ^1$ mutant at 1 mM and 5 mM: 50.35 ±3.34 h and

46.42 ±1.54 h, respectively). Our findings thus demonstrated that oral metformin

administration can rescue TRPγ-mediated metabolic syndrome.

Recovery of the TRPγ mutant through oral lipase administration or lipid

absorption

The intestine is a key organ for lipid absorption and metabolism [22] and disturbances in

intestinal lipid metabolism are associated with hyperlipidemia [23]. Therefore, we next

sought to determine whether *trpγ* mutation reduces lipase production in the abdomen

by comparing lipid accumulation in the intestine. The Drosophila midgut is divided into

R1–R5 regions, each with distinct functions in nutrient processing and absorption.

Notably, the R2 region serves as a critical interface between the fly's diet and metabolic

processes [24]. To assess lipid distribution, we conducted Nile red staining in control and

mutant animals (**Figure 5A–D**). Mutant flies displayed elevated lipid deposits on the

intestinal wall of the R2 region **(Figure 5B)**, a phenotype restored to normal levels by

expressing *trpγ* in its own cells or *Dh44* neurons (**Figure 5C–E**). Consequently, we

quantified lipid droplets specifically within the intestinal wall of the R2 region in *trpγ[1]* flies,

with no such accumulation observed in control or rescued flies (**Figure 5E**). These

findings indicate that TRPγ plays a crucial role in the brain-gut axis, specifically in

controlling lipid metabolism in the intestine, as evidenced by the lipid accumulation

observed in the gut region of *trpγ[1]*. To clarify if our observations were due to insufficient

activity of lipase in the gut, lipase enzyme mixed with 1% agar food was supplied to

both control and mutants flies. Given that the brief activity of lipase in the intestine,

lipase-containing agar food was provided to the flies every 12 h. Surprisingly,

supplementation with lipase, but not denatured lipase, under starvation conditions

increased the survival of *trpγ* mutant flies to levels comparable to those of the control

flies (**Figure 5F**; LT_{50} of lipase treated-control and *trpγ[1]*: 50.35 ±0.93 h and 44.94 ±1.33

h, respectively). To support the hypothesis that the extended lifespan of mutant flies

during starvation is associated with dietary lipase intake, we examined lipid

accumulation (**Figure 5–figure supplement 1A–E**). Lipase treatment resulted in the

breakdown of lipid deposits in the mutant R2 region of intestine. Importantly, only active

lipase showed an enhanced survival time under starvation conditions in mutant flies,

while denatured lipase did not exhibit this effect. Therefore, our findings suggest that

lipase may play a role in digesting accumulated lipid in the intestine, though the

mechanisms of lipase absorption remain unclear.

Next, we sought to assess the involvement of each of the components of TAG in

the recovery of starvation susceptibility. TAG is composed of glycerol and three fatty

acids. Therefore, we first fed the flies with 1% glycerol only. Glycerol extended the

survival of the control and $trp\gamma^1$ flies by 3–4-fold (**Figure 5G**). However, the LT_{50} of the

control and $trp\gamma^1$ were still significantly different (**Figure 5H**; LT_{50} of control and $trp\gamma^1$:

244.5 ± 20.10 h and 117.75 ± 3.32 h, respectively). Next, the flies were fed with 0.2%

and 0.5% hexanoic acid (HA) (**Figure 5H and Figure 5–figure supplement 1F**; 0.2%

HA, LT_{50} of control and $trp\gamma^1$: 65.44 ± 1.94 h and 52.32 ±2.05 h, respectively; 0.5% HA,

LT_{50} of control and $trp\gamma^1$: 56.27 ±1.30 h and 46.27 ±1.47 h, respectively). Additionally,

we also tested the effects of 0.2% and 0.5% concentrations of a TAG mixture

(combination of mono-, di-, and tri- acylglycerol) (**Figure 5I and Figure 5–figure**

supplement 1G; 0.2% TAG, LT_{50} of control and $trp\gamma^1$: 52.93 ±5.12 h and 50.1 ±3.74 h,

respectively; 0.5% TAG, LT_{50} of control and $trp\gamma^1$: 65.1 ±2.6 h and 57.75 ±3.32 h,

respectively). The HA and mixed TAG treatments significantly extended the survival of

the $trp\gamma^1$ mutants, albeit not to levels similar to those of the control. This indicated that

the TRPγ mutants could not fully absorb and burn the digested lipid. Furthermore, lipid-

only dietary supplementation markedly limited the LT_{50} of the control. Importantly,

supplementation with lipolytic drugs, lipase, TAG, and free fatty acids effectively

rescued the survival of $trp\gamma$ mutants under starvation conditions. This observation was

consistent with our finding that $trp\gamma$ mutants are unable to degrade lipid stores under

starvation conditions. The dietary rescue of $trp\gamma$ mutants suggests that although the $trp\gamma$

mutation increases the levels of stored lipids, the flies were not able to utilize them

when starved. Taken together, these observations indicate that TRPγ plays an important

role in maintaining systemic lipid levels through proper expression of triglyceride lipase.

Downregulation of the lipolytic gene *brummer* in *trpγ* mutants

The *trpγ[1]* mutants exhibited marked alterations in the levels of major nutrients. Therefore, we next sought to analyze the transcriptional levels of genes related to gluconeogenesis, lipogenesis, and lipolysis. Gluconeogenesis is a metabolic process that produces glucose from non-carbohydrate carbon substrates. The genome of *D. melanogaster* harbors two gluconeogenic genes: *fructose-1,6-bisphosphatase* (*fbp*) and *phosphoenolpyruvate carboxykinase 1* (*pepck1*) [25]. The results of our real-time quantitative reverse transcription PCR (qRT-PCR) analyses indicated that the transcription levels of *fbp* and *pepck1* were not significantly different between control flies under sated and starved conditions (Fig. 6A). Next, we analyzed two lipogenic genes: *acetyl–CoA carboxylase* (*acc*) and *desaturase 1* (*desat1*) (***Figure 6A***) [26,27]. No significant differences were identified between the control and *trpγ[1]* flies under both conditions. Finally, we investigated the expression of the lipolytic gene *brummer* (*bmm*) (***Figure 6A***) [28]. Under starved conditions, the transcriptional level of *bmm* was highly increased in the control. In contrast, *bmm* was downregulated in *trpγ[1]* under sated and starved conditions. Therefore, the transcriptional regulation of *w[1118]* and *trpγ[1]* during starvation was markedly and significantly different. Brummer, a homolog of human adipocyte triglyceride lipase (ATGL), is associated with lipid storage in the form of LDs [29,30]. These results suggest that the lipolytic process is not sufficient to deplete stored lipids in the *trpγ[1]* mutant, resulting in excessive lipid storage. Therefore, the survival time of the *trpγ[1]* mutant during starvation was significantly shorter than that of the control.

Transgenic flies expressing *bmm*::*GFP* (*bmm* promoter *GFP*) in their FB cells

were used to examine *bmm* expression in LDs [30]. The fluorescent images obtained from

the transgenic flies were then used to confirm our qRT-PCR results. Two images of

control flies under sated and starved conditions were first analyzed both qualitatively

and quantitatively (**Figure 6B and C**). Next, *trpγ¹* was analyzed under the two conditions

(**Figure 6B and C**). A reduction in fluorescent intensity was clearly detectable in the

trpγ¹ mutant using the transgenic reporter line. These results further confirmed that *trpγ¹*

has aberrant regulation in the lipolytic pathway.

To further substantiate the idea that the reduction in the expression level of *bmm*

is the primary cause of the enlarged LDs in *trpγ* mutant flies, we measured LDs in the

FB after expressing *UAS-bmm* using specific *GAL4* drivers: *trpγ^{G4}, Dh44-GLA4, r4-*

GAL4 (fat body specific)*, Myo1A-GAL4* (gut enterocyte specific), and *Dmef-GAL4*

(muscle specific). Only *trpγ^{G4}, Dh44-GAL4* and *r4-GAL4* were able to restore the

enlarged LDs defects in the FB after expressing *UAS-bmm* (**Figure 6D–I**). Overall,

these findings suggest that the downregulation of *bmm* is one of the factors contributing

to the elevated lipid defects in *trpγ¹*.

To conclusively explore whether metformin targets enzymes involved in lipid

metabolism, specifically lipolytic or lipogenic genes, we administered 5 mM metformin

supplemented in normal food for one day and measured the expression levels of *bmm*,

acc, and *desat1*. Oral supplementation of metformin resulted in an increase in the

expression level of the lipolytic gene, *bmm* lipase of the control as well as *trpγ¹* (**Figure**

6J). Importantly, there was no observed effect of metformin on the expression levels of

lipogenic genes. This data provides insights into the mode of action of metformin,

suggesting a specific impact on lipolysis.

Assessment of the role of DH44 and its receptors (DH44R1 and DH44R2) in

relation to the utilization of fat

In Drosophila, two receptors have been identified for DH44 [31]. In order to investigate the

potential roles of DH44 in fat utilization and identify the receptor responsible for nutrient

regulation, we examined lipid and protein levels in $Dh44^{Mi}$, $Dh44R1^{Mi}$, and $Dh44R2^{Mi}$.

$Dh44^{Mi}$ and $Dh44R1^{Mi}$ showed normal total TAG levels, while $Dh44R2^{Mi}$ exhibited higher

lipid levels like $trp\gamma^1$ (**Figure 7A**). Additionally, Nile red staining of LDs in the FBs

revealed that $Dh44^{Mi}$ and $Dh44R2^{Mi}$ mutants possessed larger LDs (**Figure 7B–F**),

although protein levels remained unaffected across all three mutants (**Figure 7G**).

Further investigations quantified the expression of the bmm in the FBs of $Dh44^{Mi}$,

$Dh44R1^{Mi}$, and $Dh44R2^{Mi}$ under both sated and starved conditions. In sated conditions,

bmm expression was found to be downregulated in $Dh44R2^{Mi}$ compared to the control

(**Figure 7H**). Interestingly, under starved conditions, while bmm levels significantly

increased in the control, indicating a normal response to starvation, such an increase

was not observed in $Dh44^{Mi}$ and $Dh44R2^{Mi}$. Instead, in these mutants, bmm levels

significantly decreased compared to their levels under sated conditions (**Figure 7H**).

This suggests a unique role of DH44 and its receptor Dh44R2 in regulating lipid

metabolism and response to nutritional status in Drosophila. To explore the connection

between TRPγ and DH44 signaling, we expressed UAS-$trp\gamma$ under the control of

$Dh44R1$-$GAL4$ and $Dh44R2$-$GAL4$. Only $Dh44R2$-$GAL4$ restored the lipid defect in $trp\gamma^1$,

unlike *Dh44R1-GAL4* (**Figure 7I**). Furthermore, *Dh44R2-GAL4*, but not *Dh44R1-GAL4*,

rescued the excessive lipid accumulation in the FB of *trpγ^1^* mutants (**Figure 7J–N**).

Because *Dh44R2* are mostly expressed in the intestine (**Figure 7O and P**), further

study regarding DH44R2 and fat usage are required. Overall, our findings suggest that

trpγ expression regulates lipid homeostasis specifically in *Dh44* and *Dh44R2* cells.

355   DISCUSSION

Sugars such as D-glucose, D-trehalose, and D-fructose are commonly detected in

DH44 neurons in the PI due to the inherent nature and functions of these cells [32]. Here,

we found that the function of TRPγ, one of the TRPC channels of *D. melanogaster*, in

DH44 neuroendocrine cells plays an essential role in lipid regulation, which is linked to

alterations in membrane lipids [33,34]. The *trpγ^1^* mutant exhibited clear signs of metabolic

syndrome, as demonstrated by reduced carbohydrate level coupled with much higher

protein and lipid levels in the body. More importantly, we found that the expression of

trpγ was necessary and sufficient to regulate the carbohydrate and lipid levels in the

DH44 neurons of *trpγ* mutants. However, we failed to identify the specific cells required

for regulating protein homeostasis because the increased protein levels in the *trpγ*

mutants were not recovered by pilot screening (data not shown) except in the *trpγ-GAL4*.

In conclusion, our findings suggested that the DH44 system has important roles in

regulating the metabolic homeostasis of carbohydrates and lipids. Recent studies have

suggested that corticotropin-releasing hormone (CRH) is the mammalian homolog of

DH44 [32]. Therefore, we propose that CRH neurons may also regulate lipid levels.

The metabolic dysregulation of carbohydrates and lipids observed in the *trpγ*

mutants was phenocopied by the cellular inactivation of DH44 cells. Compared to the

controls, the *Dh44-GAL4*/*UAS-Kir2.1* flies exhibited lower carbohydrate levels and

increased lipid levels. We previously proposed that TRPγ holds DH44 neurons in a state

of afterdepolarization, thus reducing firing rates by inactivating voltage-gated Na$^+$

channels [11]. At the physiological level, this induces the consistent release of DH44 and

depletion of DH44 stores, resulting in nutrient utilization and storage malfunctions.

Likewise, our findings revealed that *Dh44* mutant and its receptor mutant, *Dh44R2*,

displayed defects analogous to those observed in the *trpγ* mutant. Finally, it would also

be interesting to investigate the potential roles of TRPC members in humans, especially

the involvement of TRPC4 and TRPC5 in metabolic syndrome.

Multiple lines of evidence have suggested that DH44 neurons communicate with

the intestine. First, DH44 at the PI in the brain is an incretin-like hormone that acts as a

hemolymph nutrient sensor. The secretion of DH44 is highly dependent on the calorie

value of a given diet, as DH44 secretion is only mediated by nutritional sugars but not

non-nutritional sugars [32,35]. Second, the expression of TRPγ in the PI is sufficient to

recover any metabolic defects such as reduced carbohydrate and increased lipid levels,

as well as related phenomena such as starvation resistance. DH44 and kinin

neuropeptides influence desiccation and starvation tolerance in *D. melanogaster* [36].

Although *Dh44^{Mi}* exhibited normal TAG levels, the presence of larger LDs indicated that

DH44 plays a regulatory role in lipid metabolism. LDs consist of a hydrophobic core of

neutral storage fats, such as triglycerides or cholesterol esters, encased by a protein-

coated phospholipid monolayer. From our findings, it is suggested that DH44 and its

receptor DH44R2, but not DH44R1, are involved in the control of lipid storage,

indicating a specific pathway through which DH44 influences lipid metabolism distinct

from its role in carbohydrate metabolism and overall energy homeostasis. Third, *trpγ*

gene knockdown through RNAi in the DH44 neurons is enough to phenocopy all of the

metabolic defects observed in the *trpγ* mutants. Fourth, *trpγ* mutants exhibit low

expression of *brummer* lipase, which results in accumulation of LDs in the body, which

is a reciprocal process regulated by DH44 neurons. Fifth, the activation of AMPK in the

DH44 neurons can recover the defects of lipid level of *trpγ[1]*. Sixth, the reduced

starvation resistance can be significantly recovered through dietary lipase

supplementation. These findings support the assertion that the sugar-sensing DH44

neurons send downward signals to the intestine by absorbing the carbohydrates and

lipids derived from dietary intake. However, DH44 neurons do not regulate protein levels

in the body because the increased protein levels of *trpγ* mutants were not recovered by

the expression of *trpγ* in the DH44 neurons. Nevertheless, it is also known that DH44

neurons mediate the influence of dietary amino acids on promoting food intake in flies [37].

These findings confirm that *trpγ* regulates *bmm* expression in the adult bowel, enabling

dietary lipid digestion and maintaining systemic lipid homeostasis.

Treatment with metformin, a lipolytic agent, at two different concentrations (1 mM

and 5 mM) decreased the total TAG levels and reduced the size of the LDs in *D.*

melanogaster. Similarly, metformin increased the starvation resistance of the *trpγ[1]*

mutants but not the controls. These findings were consistent with the widely

acknowledged anti-obesity properties of metformin [21]. Furthermore, our results were in

agreement with the evidence that metformin helps to reduce lipid accumulation caused

by diets rich in saturated fatty acids [38]. Although the exact target of metformin is not

known, its lipolytic effect is significant and widely documented. Oral administration of 5

mM of metformin reduced normal TAG levels of wild-type flies, whereas the 1 mM dose

had no significant effect. Therefore, low metformin doses were effective at decreasing

stored lipid levels in flies. The accumulation of intracellular LDs is regulated by

autophagy. Metformin improves autophagic flux through AMPKα-mediated mechanisms

[38]. In the Drosophila context, we propose a mechanism involving AMPK activation within

DH44 neurons as a rescue mechanism for starvation resistance in $trp\gamma^1$, highlighting its

crucial role in this process. AMPK activation is known to enhance an organism's ability

to endure environmental stresses by facilitating autophagy-mediated cellular cleanup,

promoting mitochondrial biogenesis for increased energy production, and optimizing

metabolic pathways for enhanced energy efficiency [38]. While the precise nature of the

connection between TRPγ, DH44 neuronal activity, and AMPK activation remains

unspecified, it is conceivable that TRPγ influences AMPK activation or regulation within

DH44 neurons, contributing to stress responses, ATP supply, and metabolic

adaptations. Metformin can influence lipid metabolism through various mechanisms,

one of which is the promotion of lipolysis [39]. This effect results in a gradual reduction of

stored lipid levels over time. Orally administering metformin to flies led to a significant

reduction in lipid levels, accompanied by an increase in survival. Subsequent qPCR

analysis post-metformin feeding revealed a significant upregulation in the expression

level of the lipolytic gene (*bmm*). These results strongly support our hypothesis that

metformin induces lipolysis when orally administered.

The exact mechanism through which orally administered lipase leads to gastric

lipolysis remains unclear. However, our experimental data indicated that 0.1% lipase

increased the survival time of $trp\gamma^1$ under starvation. Lipase exhibits a high degree of

chemical selectivity and has the ability to catalyze triglyceride into glycerol ester,

monoglycerides, glycerol, and fatty acids [40]. Free fatty acids, HA, and glycerol at

concentrations of 0.2%–0.5% can be utilized as energy sources and increase survival

time under starvation conditions, suggesting that $trp\gamma^1$ participates in the utilization of

HA and glycerol under starvation conditions.

The therapeutic properties of metformin as a lipolytic and anti-diabetic drug have

been widely documented in humans. However, additional studies are needed to identify

the exact target of metformin. In this context, our findings demonstrate that *D.*

melanogaster can be an excellent model to study the mode of action and biosafety of

metformin.

MATERIALS AND METHODS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Mouse anti-GFP	Molecular probe	Cat # A11120
Rabbit anti-DsRed	Takara	Cat # 632496
Goat anti-mouse Alexa Fluor 488	Thermo Fisher/Invitrogen	Cat # A11029
Goat anti-mouse Alexa Fluor 568	Thermo fisher/Invitrogen	Cat # A11004
Rabbit anti-Dh44	Dr.J.A. Veenstra	NA
Goat anti-rabbit Alexa Fluor 488	Thermo fisher/Invitrogen	Cat # A11034
Goat anti-rabbit Alexa Fluor 568	Thermo fisher/Invitrogen	Cat # A11036
Chemicals, peptides, and recombinant proteins		
Trehalase from porcine kidney	Sigma-Aldrich Co.	Cat # 9025-52-9
Amyloglucosidase from Aspergillus	Sigma-Aldrich Co.	Cat # 9032-08-0
Glucose (HK) Assay Kit	Sigma-Aldrich Co.	Cat # GAHK-20

Glucose (HK) Assay reagent	Sigma-Aldrich Co.	Cat # G3293
The Pierce™ BCA protein assay kit	Thermo Fischer Scientific	Cat # 23225
Triglyceride mix	Sigma-Aldrich Co.	Cat #17810-1AMP-S
LiquiColor triglyceride test kit	Stanbio Laboratory	Cat # 2100-225
Lipase from *Candida rugosa*	Sigma-Aldrich Co.	Cat # L1754
Metformin	Sigma-Aldrich Co.	Cat # D150959
Hexanoic acid	Sigma-Aldrich Co.	Cat # W255912
Sucrose	Sigma-Aldrich Co.	Cat # 57-50-1
LiquiColor Triglyceride Test kit	Stanbio Laboratory	Cat # 2100-225
Glycerol	Sigma-Aldrich Co.	Cat # G5516
Nile red	Sigma-Aldrich Co.	Cat # N3013
Experimental models: Organisms/strains		
Genotype	Source	Identifier
$trp\gamma^1$	Dr. C. Montell	BDSC:64311
$trp\gamma^{G4}$	Dr. C. Montell	BDSC:64313
$trp\gamma^1,UAS\text{-}trp\gamma/CyO$	Dr. C. Montell	[41]
$trp\gamma^1;g(trp\gamma)$	Dr. C. Montell	[41]
$trpA1^1$	Dr. C. Montell	BDSC:26504
amo^1	Dr. C. Montell	[42]
$trpml^2$	Dr. C. Montell	BDSC:42230
trp^{343}	Dr. C. Montell	BDSC:25082
pyx^3	Dr. Y. Lee	[43]
$wtrw^{ex}$	Dr. C. Montell	BDSC:59038
$trpl^{29134}$	Bloomington Drosophila Stock center	BDSC:29134
iav^{3621}	Bloomington Drosophila Stock center	BDSC: 24768
$pain^2$	Dr. S. Benzer	[44]
nan^{36a}	Dr. C. Kim	[45]
dilp2-GAL4	KDRC	KDRC:200
Dh44-GAL4	Dr. Y. Kim	BDSC: 1453
UAS-mCD8::GFP	Bloomington Drosophila Stock Center	BDSC: 5130
UAS-Kir2.1	Bloomington Drosophila Stock Center	BDSC: 6596
$UAS\text{-}trp\gamma^{RNAi}$	Bloomington Drosophila Stock Center	Transformant ID107656
r4-GAL4	Korea Drosophila Resource Center	GIST: 2166
Myo1A-GAL4	Dr. S. Hyun	[46]
Dmef-GAL4	Dr. S. Hyun	[47]
$UAS\text{-}AMPK^{TD}$	Korea Drosophila Resource Center	KDRC:10099
UAS-bmm	Bloomington Drosophila Stock Center	BDSC: 76600
$Dh44^{MI}$	Dr. Y. Kim	
$Dh44R1^{MI}$	Dr. Y. kim	
$Dh44R2^{MI}$	Dr. Y. Kim	
Dh44R1-GAL4	Dr. Y. Kim	
Dh44R2-GAL4	Dr. Y. Kim	

yw;+;bmm::GFP	Dr. Kaeko Kamei	[30]
Software and algorithms		
Origin Pro Version	OriginLab corporation	RRID: SCR_002815 https://www.originlab.com
Graphpad Prism	GraphPad	RRID: SCR_002798 https://www.graphpd.com
Fiji(ImageJ-win64)	https://fiji.sc	

Chemical sources

Trehalase from porcine kidney (cat. # 9025-52-9), amyloglucosidase (cat. # 9032-08-0),

the glucose (HK) assay kit (cat. # GAHK-20-1KT), and triglyceride mix (cat. # 17810-

1AMP-S), lipase from *Candida rugosa* (cat. # OT705690000), metformin (cat. #

D150959), Nile Red (cat. # N3013), glycerol (cat. # G5516) were purchased from

Sigma-Aldrich Co. Hexanoic acid (cat. # W255912) were purchased from Wako Pure

Chemical Industry Ltd. The Pierce™ BCA protein assay kit (cat. # 23225) and the

LiquiColor triglyceride test kit (cat. # 2100-225) were purchased from Thermo Fisher

Scientific and Stanbio Laboratory, respectively.

TAG level measurement

TAG levels were quantified as described previously with some modifications using a

LiquiColor Triglyceride Test kit (cat. # 2100-225, Stanbio Laboratory, Germany) [21]. Ten

5–10 day-old male flies were weighed and crushed in 1 mL of PBST (1X PBS and 0.2

percent Triton X-100). The homogenate was incubated at 70 °C for 5 minutes and

centrifuged for 3 minutes at 9,500 g. Next, 100 μL of supernatant was transferred into a

1.5 mL Eppendorf tube and mixed with 1 mL of Stanbio LiquiColor Triglyceride Test kit

reagent or 1 mL of deionized water to provide a baseline. The reaction mixture was kept

at 37 °C for 15 minutes. Finally, the absorbance of the sample solution was measured

at 500 nm using a spectrophotometer and the TAG level was calculated based on a

standard calibration curve.

Protein level measurement

Protein assays were performed as previously described using the Pierce[TM] BCA Protein

Assay Kit with some modifications. Briefly, ten 5–10 day-old male flies were weighed

and crushed in 1 mL of PBST (1X PBS and 0.2 percent Triton X-10) and incubated at

70 °C for 5 minutes. The homogenate was then centrifuged for 3 minutes at 9,500 g,

after which 300 µL of supernatant was mixed with 600 µL of Pierce[TM] BCA Protein

Assay Kit (UF289330). After a 30-minute incubation period at 37 °C, the absorbance of

the samples was measured at 530 nm using a spectrophotometer and compared to a

standard calibration curve for quantification.

Trehalose and glucose measurements in whole adult flies or hemolymph

Whole-body glucose and trehalose levels were measured in adult flies as previously

described [48]. Briefly, 10 freshly hatched males were collected, weighed, and crushed in

250 µL of 0.25 M Na_2CO_3 buffer. The homogenates were then incubated in a water bath

(95°C) for 5 min to inactivate all enzymes. Next, 600 µL of 0.25 M sodium acetate and

150 µL of 1 M acetic acid (pH 5.2) were added to the samples, after which the mixtures

were centrifuged at 12,500 g for 10 min at 24 °C. Afterward, 200 µL of supernatant was

transferred a new microfuge tube and 2 μL porcine kidney trehalase (Sigma: T8778 UN)

was added to the sample mixture and incubated overnight at 37 °C to convert trehalose

into glucose. Next, 1 mL of glucose hexokinase solution (Sigma: GAHK-20) was added

to 100 μL of the sample and incubated for 20 min at 37 °C. Optical density (OD) values

were measured at 340 nm. Finally, total glucose and trehalose levels were calculated

using a standard glucose curve generated through similar reactions with standard

trehalose and glucose.

Glucose and trehalose level measurement in hemolymph were extracted from

the male flies. Flies were punctured in the thorax with the help of fine injection needle

and kept in 0.5 mL tubes having a punctured base with 21-gauge needle. The

punctured flies were adjusted by keeping the shoulder down to prevent leakage from

the genital tract. The tubes were set into the 1.5 mL microfuge tube and centrifuged at

4°C for 5 minutes at 2800 g force. 0.5 μL of hemolymph was added to 14.5 μL PBS. 2

μL of porcine kidney trehalase (Sigma: T8778 UN) was added to the sample mixture

and incubated overnight at 37°C to convert trehalose into glucose. 1 mL of glucose

hexokinase solution (Sigma: GAHK-20) was added to the 100 μL of the sample and

incubated at 37°C for 20 minutes. Optical density (OD) values were measured at 340

nm. Total glucose and trehalose levels were calculated using a standard glucose curve

generated through similar reactions with standard trehalose and glucose.

Glycogen measurements

Tissue glycogen levels from whole-body extracts of adult flies were quantified as

previously described [49]. Briefly, 5 males were homogenized in 100 μL of ice-chilled 1X

PBS after measuring the weight. The homogenates were kept at 70 °C for 5 min to

inactivate all metabolic enzymes. The homogenates were then centrifuged at 12,500 g

for 3 min at 4 °C, after which 20 µL of the supernatant was transferred to 1.5 mL

microfuge tubes and diluted with 1X PBS to a 1:3 ratio. An amyloglucosidase dilution

was prepared by mixing 1.5 µL of amyloglucosidase (Sigma A1602) suspension in

998.5 µL 1X PBS. Finally, a 20 µL aliquot of the diluted sample was added to 20 µL of

the diluted amyloglucosidase solution, as well as to the glycogen standard. Both the

glycogen standard and test samples were than incubated at 37 °C for 1 h. A commercial

glucose (HK) assay reagent (Sigma: G3293 VER) was used to measure total glucose at

340 nm. The glycogen level was quantified by comparing it with standard curve plotted

from standard glycogen samples.

Survival assay

Survival experiments were conducted as previously described [50]. Normal cornmeal diets

were used to conduct the normal survival assays. To measure survival under starvation

conditions, twenty 3–4-day-old male flies were fed with 1% agar food supplemented

with or without various concentrations of lipase (0.1%; active or denatured), glycerol

(1%), hexanoic acid (0.2%, 0.5%), glycerol (1%), TAG mix (0.2%, 0.5%), or metformin

(1 mM, 5 mM). Every 12 hours, the flies were monitored/counted and then transferred to

fresh vials with the same food supply. The experiments were conducted until the food

vials had been cleared.

Immunohistochemistry

We performed immunohistochemistry as previously described [11]. Briefly, to fix and block

the specimens, we placed freshly dissected tissues (brain and intestine) into a well of

24-well tissue culture plate (Costar Corp.) placed on ice, which contained 940 μL of

fixing buffer (1 mM EGTA, 0.1M Pipes pH 6.9, 2 mM $MgSO_4$, 1% TritonX-100, 150 mM

NaCl). 60 μL formaldehyde (37%) was add up to the wells and mixed instantly before

the tissue were added. As many as tissues that were dissected within 15 minutes were

fixed and then the samples were incubated for 30 min more. Samples were washed with

wash buffer (1X PBS, 0.1% saponin) for three times (15 min. for each washing), and

blocked with 1 mL blocking buffer (1X PBS, 0.1% saponin, and 5 mg/mL BSA) at 4 °C

for 4-8 hours.

To perform immunostaining, primary antibodies were inserted into the sample at 4°C

for 18 hours [4]. Samples were washed with wash buffer three times (15 min each) and

added secondary antibodies {(1:200) goat anti-mouse Alexa Fluor 488 (cat. # A11029),

goat anti-mouse Alexa Fluor 568 (cat. # A11004), goat anti-rabbit Alexa Fluor 488 (cat.

A11034), and goat anti-rabbit Alexa 568 (cat. # A11036) at 4 °C for 4 hours. Finally,

samples were washed three times and stored into 1.25X PDA (187.5 mM NaCl, 37.5%

glycerol, 62.5 mM Tris pH 8.8), and kept at 4°C for more than 1 hr. Samples were

mounted and examined using a Leica Stellaris 5 Confocal Microscope.

Nile red staining

Nile Red is a dark purplish-red powder (Sigma N-3013), the stock solution must be

prepared in acetone (1000 μg/mL) and kept in a tightly sealed, lightproof container at

4 °C. Briefly, 5–10 day-old male flies were fixed in a sagittal position on a glass slide

and submerged in a 1 X PBS solution. FBs were gently dissected from a dorsal

abdominal region along with thorax or the gut under a stereomicroscope. The dissected

tissues were fixed with 4% formaldehyde solution for 15 minutes at room temperature.

The fixed tissues were gently washed three times with 1X PBS (5 min for each wash).

Nile Red (1:1000 dilution) were added to the tissue samples for 5 minutes. Finally, the

stained tissues were washed with 1 mL 1X PBS and mounted in 50% glycerol on a

glass slide. LD deposition exhibits greater density in the upper abdominal region

compared to the lower abdominal area. This comparison was meticulously conducted

within the identical segments 2-3 of the abdomen to ensure accuracy.

For intestinal lipid staining, 5-10 days old male flies were fixed on a glass slide

and submerged in 1X PBS solution. Full intestine was dissected very carefully. The

dissected intestine was fixed with 4% formaldehyde solution for 15 minutes at room

temperature. The fixed intestine was washed 3 times with 1X PBS. Nile red was added

to the sample for 5 minutes. The stained samples were washed with 1mL 1X PBS and

mounted in 50% glycerol on a glass slide. The stained tissues were examined using a

Leica Stellaris 5 Confocal Microscope.

qRT-PCR

qRT-PCR assays were performed as described previously [51]. Briefly, ten 6–10 day-old

male flies (control and mutant) were selected for whole body and 15 flies were selected

for fat body samples in Supplement Figure 6K. The experiments were conducted under

sated (0 h starvation) and starved (24 h starvation) conditions. Total RNA was extracted

using the TRizol reagent (Invitrogen) followed by DNase (Promega) treatment. cDNA

was synthesized using the AMV reverse transcriptase system (Promega). RT-qPCR

experiments were carried out using a Bio-rad CFX system. The Takara TB Green

Premix was used to assess the mRNA expression level of each gene according to the

manufacturer's instructions. Relative gene expression was calculated using the $2^{-\Delta\Delta Ct}$

method. Three biological samples were used and transcript levels were normalized to

the *D. melanogaster* housekeeping gene *tubulin*. The experiments were conducted

using the following primer pairs: *acc*, 5'-ACG AGG GCG AGC AGC GTT AC-3′ (forward)

and 5′-TAG GGC GAC TTG GTG GGC AT-3′ (reverse); *bmm*, 5′-ATG ACT TCG GAC

TTC TTC AGG G-3′ (forward) and 5′-CCA ATT CAG ATG GAA GAG CTG-3′ (reverse);

fbp, 5′-CTC CAA CGA GCT GTT CAT CA-3′ (forward) and 5′-TGA ACC GAT CGA CAC

CAG GC-3′ (reverse); *pepck1*, 5′-AGG TGC ACA TCT GCG ATG GC-3′ (forward) and

5′-CCA CCA CGT AAG CAG AGT CC-3′ (reverse); *desat1*, 5'-AAG CCG GTG CCC

AGT CCA TC-3' (forward) and 5'-ATG GTC GCG AGC CCA ATG GT-3' (reverse); and

tubulin, 5′-TCC TTG TCG CGT GTG AAA CA-3' (forward) and 5′-CCG AAC GAG TGG

AAG ATG AG-3' (reverse).

Statistics and Reproducibility

D. melanogaster was selected as a model organism in this study. For the experiments,

male flies were mostly used unless we did not mention the sex. All of the experiments

were conducted under laboratory conditions. The appropriate number of replicates was

established based on previous research. A large enough sample size was used in all of

our assays to ensure that our results were representative and repeatable. No data

points were left out of the analysis. For each genotype, the data points indicate the

values of individual replicates. The error bars in all of the figures represent the standard

error of the mean (SEM). The analysis of the RT-qPCR data was conducted using the

C_T values. Comparisons between multiple experimental groups were conducted via

single-factor ANOVA and Scheffe's *post hoc* test. Pair-wise comparisons were

conducted via Student's t-test. Survival curves were estimated for each group, using a

Kaplan-Meier method and compared statistically using the log-rank tests. The asterisks

in the figures indicate statistical significance (*$P<0.05$, **$P<0.01$). All statistical analyses

were conducted using the Origin Pro 8 software for Windows (ver. 8.0932; Origin Lab

Corporation, USA).

Competing interest statement

624 The authors have no competing interests.

Acknowledgments

This work was supported by grants to Youngseok Lee from the National Research

Foundation of Korea (NRF) funded by the Korea government (MIST) (NRF-

2021R1A2C1007628) and the Biomaterials Specialized Graduate Program through the

Korea Environmental Industry & Technology Institute (KEITI) funded by the Ministry of

Environment (MOE). D.K.N. and S.D. were supported by the Global Scholarship

Program for Foreign Graduate Students at Kookmin University in Korea.

636 Author contributions

637 D.K.N. and S.D. performed the experiment, analyzed the data, generated the figures,

638 and wrote the original draft. Y.L. supervised, reviewed and edited the draft, wrote the

639 original manuscript, and obtained the funding.

References

1. Efeyan, A., Comb, W.C., and Sabatini, D.M. (2015) **Nutrient-sensing mechanisms and pathways** *Nature* **517**, 302-310 https://doi.org/10.1038/nature14190

2. Tran, L.T., Park, S., Kim, S.K., Lee, J.S., Kim, K.W., and Kwon, O. (2022) **Hypothalamic control of energy expenditure and thermogenesis** *Expe. Molecul. Med.* **54**, 358-369 https://doi.org/10.1038/s12276-022-00741-z

3. Roh, E., and Kim, M.-S. (2016) **Brain regulation of energy metabolism** *Endocrinol. Metab.* **31**, 519 https://doi.org/103803EnM2016.31.4.519

4. Jais, A., and Brüning, J.C. (2022) **Arcuate nucleus-dependent regulation of metabolism—pathways to obesity and diabetes mellitus** *Endocr. Rev.* **43**, 314-328 https://doi.org/10.1210/endrev/bnab025

5. Clapham, D.E. (2003) **TRP channels as cellular sensors** *Nature* **426**, 517-524.

6. Fowler, M.A., and Montell, C. (2013) *Life Sci.* **92**, 394-403 https://doi.org/10.1038/nature02196

7. Dhakal, S., and Lee, Y. (2019) **Transient receptor potential channels and metabolism** *Mol. Cell.* **42**, 569-578 https://doi.org/10.14348/molcells.2019.0007

8. Turner, H.N., Armengol, K., Patel, A.A., Himmel, N.J., Sullivan, L., Iyer, S.C., Bhattacharya, S., Iyer, E.P.R., Landry, C., and Galko, M.J. (2016) **The TRP channels Pkd2, NompC, and Trpm act in cold-sensing neurons to mediate unique aversive behaviors to noxious cold in Drosophila** *Curr. Biol.* **26**, 3116-3128 https://doi.org/10.1016/j.cub.2016.09.038

9.   Venkatachalam, K., and Montell, C. (2007) **TRP channels** *Annu. Rev. Biochem.*

76, 387-417 https://doi.org/ 10.1146/annurev.biochem.75.103004.142819

10.   Picard, M., McEwen, B.S., Epel, E.S., and Sandi, C. (2018) **An energetic view**

of stress: Focus on mitochondria *Front. Neuroendocrinol.* **49**, 72-85

https://doi.org/10.1016/j.yfrne.2018.01.00

11.   Dhakal, S., Ren, Q., Liu, J., Akitake, B., Tekin, I., Montell, C., and Lee, Y. (2022)

Drosophila TRPγ is required in neuroendocrine cells for post-ingestive

food selection *Elife* **11**, e56726-e56726 https://doi.org/10.7554/eLife.56726

12.   Hodge, J.J. (2009) **Ion channels to inactivate neurons in Drosophila** *Front.*

Mol. Neurosci. **13** https://doi.org/10.3389/neuro.02.013.2009

13.   Ohhara, Y., Kobayashi, S., Yamakawa-Kobayashi, K., and Yamanaka, N. (2018)

Adult-specific insulin-producing neurons in Drosophila melanogaster *J.*

Comp. Neurol. **526**, 1351-1367 https://doi.org/10.1002/cne.24410

14.   Liu, Z., and Huang, X. (2013) **Lipid metabolism in Drosophila: development**

and   disease.   *Acta   Biochim*   *Biophys.   Sin.*   **45,   44-50**

https://doi.org/10.1093/abbs/gms105

15.   Tatar, M., Post, S., and Yu, K. (2014) **Nutrient control of Drosophila longevity**

Trends Endocrinol. Metab. **25**, 509-517 https://doi.org/10.1016/j.tem.2014.02.006

16.   Johnson, A.A., and Stolzing, A. (2019) **The role of lipid metabolism in aging,**

lipespan   regulation,   and   age-related   disease *Aging   cell* **18**, e13048

https://doi.org/10.1111/acel.13048

685 17. Hardie, D.G., Scott, J.W., Pan, D.A., and Hudson, E.R. (2003) **Management of**

686   **cellular energy by the AMP-activated protein kinase system** *FEBS letters*

687   **546**, 113-120 https://doi.org/10.1016/S0014-5793(03)00560-X

688 18. Kim, J., Yang, G., Kim, Y., Kim, J., and Ha, J. (2016) **AMPK activators:**

689   **mechanisms of action and physiological activities** *Expe. Mol. Med.* **48**, e224-

690   e224 https://doi.org/10.1038/emm.2016.16

691 19. Lee, J.H., Koh, H., Kim, M., Kim, Y., Lee, S.Y., Karess, R.E., Lee, S.-H., Shong,

692   M., Kim, J.-M., and Kim, J. (2007) **Energy-dependent regulation of cell**

693   **structure by AMP-activated protein kinase** *Nature* **447**, 1017-1020

694   https://doi.org/10.1038/nature05828

695 20. Lv, Z., and Guo, Y. (2020) **Metformin and its benefits for various diseases**

696   *Front. Endocrinol.* **11**, 191 https://doi.org/10.3389/fendo.2020.00191

697 21. Sang, J., Dhakal, S., and Lee, Y. (2021) **Cucurbitacin B suppresses**

698   **hyperglycemia associated with a high sugar diet and promotes sleep in**

699   **Drosophila melanogaster** *Mol. and Cell.* **44**, 68

700   https://doi.org/10.14348/molcells.2021.2245

701 22. Han, J., Guo, X., Meng, X.-J., Zhang, J., Yamaguchi, R., Motoo, Y., and Yamada,

702   S. (2020) **Acupuncture improved lipid metabolism by regulating intestinal**

703   **absorption in mice** *World J. Gastroenterol.* **26**, 5118

704   1https://doi.org/10.3748/wjg.v26.i34.5118

705 23. Jia, X., Xu, W., Zhang, L., Li, X., Wang, R., and Wu, S. (2021) **Impact of gut**

706   **microbiota and microbiota-related metabolites on hyperlipidemia** *Front. Cell.*

707   *Infect. Microbiol.* **11**, 634780 https://doi.org/10.3389/fcimb.2021.634780

708   24.   Capo, F., Wilson, A., and Di Cara, F. (2019) **The intestine of Drosophila**
709   **melanogaster: An emerging versatile model system to study intestinal**
710   **epithelial homeostasis and host-microbial interactions in humans**
711   *Microorganisms* **7**, 336 https://doi.org/10.3390/microorganisms7090336
712   25.   Miyamoto, T., and Amrein, H. (2019) **Neuronal gluconeogenesis regulates**
713   **systemic glucose homeostasis in Drosophila melanogaster** *Curr. Biol.* **29**,
714   1263-1272. e1265 https://doi.org/10.1016/j.cub.2019.02.053
715   26.   Wang, Y., Yu, W., Li, S., Guo, D., He, J., and Wang, Y. (2022) **Acetyl-CoA**
716   **carboxylases and diseases** *Front. Oncol.* **12,** 836058
717   doi.org/10.3389/fonc.2022.836058 https://doi.org/10.1007/s11306-013-0511-3
718   27.   Yee, J.K., Wahjudi, P.N., Vega, J., Lim, S., Martin, A., Patterson, M.E., Cohen,
719   J.N., Mao, C.S., and Lee, W.-N.P. (2013) **Stearoyl-CoA desaturase enzyme 1**
720   **inhibition reduces glucose utilization for de novo fatty acid synthesis and**
721   **cell proliferation in 3T3-L1 adipocytes** *Metabolomics* **9**, 809-816
722   https://doi.org/10.1007/s11306-013-0511-3
723   28.   Blumrich, A., Vogler, G., Dresen, S., Diop, S.B., Jaeger, C., Leberer, S., Grune,
724   J., Wirth, E.K., Hoeft, B., and Renko, K. (2021) **Fat-body brummer lipase**
725   **determines survival and cardiac function during starvation in Drosophila**
726   **melanogaster** *iScience* **24**, 102288 https://doi.org/10.1016/j.isci.2021.102288
727   29.   Grönke, S., Mildner, A., Fellert, S., Tennagels, N., Petry, S., Müller, G., Jäckle, H.,
728   and Kühnlein, R.P. (2005) **Brummer lipase is an evolutionary conserved fat**
729   **storage regulator in Drosophila** *Cell Metab.* **1**, 323-330
730   https://doi.org/10.1016/j.cmet.2005.04.003

30. Men, T.T., Thanh, D.N.V., Yamaguchi, M., Suzuki, T., Hattori, G., Arii, M., Huy,

N.T., and Kamei, K. (2016) **A Drosophila model for screening antiobesity**

agents *BioMed Res. Internat.* 2016 https://doi.org/10.1155/2016/6293163

31. Hector, C.E., Bretz, C.A., Zhao, Y., and Johnson, E.C. (2009) **Functional**

differences between two CRF-related diuretic hormone receptors in

Drosophila *J. Exp. Biol.* **212**, 3142-3147 https://doi.org/10.1242/jeb.033175

32. Dus, M., Lai, J.S.-Y., Gunapala, K.M., Min, S., Tayler, T.D., Hergarden, A.C.,

Geraud, E., Joseph, C.M., and Suh, G.S. (2015) **Nutrient sensor in the brain**

directs the action of the brain-gut axis in Drosophila *Neuron* **87**, 139-151

https://doi.org/10.1016/j.neuron.2015.05.032

33. Overgaard, J., Sørensen, J.G., Petersen, S.O., Loeschcke, V., and Holmstrup, M.

(2005) **Changes in membrane lipid composition following rapid cold**

hardening in Drosophila melanogaster *J. Insect physiol.* **51**, 173-1182

https://doi.org/10.1016/j.jinsphys.2005.06.007

34. Cobb, T., Damschroder, D., and Wessells, R. (2021) **Sestrin regulates acute**

chill coma recovery in Drosophila melanogaster *Insect Biochem. Mol. Biol.*

133, 103548 https://doi.org/10.1016/j.ibmb.2021.103548

35. Oh, Y., Lai, J.S.-Y., Min, S., Huang, H.-W., Liberles, S.D., Ryoo, H.D., and Suh,

G.S. (2021) **Periphery signals generated by Piezo-mediated stomach stretch**

and Neuromedin-mediated glucose load regulate the Drosophila brain

nutrient sensor. *Neuron* **109**, 1979-1995

https://doi.org/10.1016/j.neuron.2021.04.028

36.   Cannell, E., Dornan, A.J., Halberg, K.A., Terhzaz, S., Dow, J.A., and Davies, S.-

A. (2016) **The corticotropin-releasing factor-like diuretic hormone 44 (DH44)**

and kinin neuropeptides modulate desiccation and starvation tolerance in

Drosophila melanogaster *Peptides* **80**, 96-107

https://doi.org/10.1016/j.peptides.2016.02.004

37.   Yang, Z., Huang, R., Fu, X., Wang, G., Qi, W., Mao, D., Shi, Z., Shen, W.L., and

Wang, L. (2018). **A post-ingestive amino acid sensor promotes food**

consumption in Drosophila. *Cell Res.* **28**, 1013-1025

https://doi.org/10.1038/s41422-018-0084-9

38.   Kim, H.-S., Ren, G., Kim, T., Bhatnagar, S., Yang, Q., Bahk, Y.Y., and Kim, J.-.

(2020) **Metformin reduces saturated fatty acid-induced lipid accumulation**

and inflammatory response by restoration of autophagic flux in endothelial

cells *Sci. Rep.* **10**, 1-14 https://doi.org/10.1038/s41598-020-70347-w

39.   Szkudelski, T., Konieczna, K., and Szkudelska, K. (2022) **Regulatory Effects of**

Metformin, an Antidiabetic Biguanide Drug, on the Metabolism of Primary

Rat Adipocytes *Molecules* **27**, 5250 https://doi.org/10.3390/molecules27165250

40.   Liu, T.-T., Liu, X.-T., Chen, Q.-X., and Shi, Y. (2020) **Lipase inhibitors for**

obesity: A review *Biomed. Pharmacother.* 128, 110314

https://doi.org/10.1016/j.biopha.2020.110314

41.   Akitake, B., Ren, Q., Boiko, N., Ni, J., Sokabe, T., Stockand, J.D., Eaton, B.A.,

and Montell, C. (2015) **Coordination and fine motor control depend on**

***Drosophila* TRPγ** *Nat. Commun.* **6**, 7288 Akitake, B., Ren, Q., Boiko, N., Ni, J.,

Sokabe, T., Stockand, J.D., Eaton, B.A., and Montell, C. (2015) Coordination and

fine motor control depend on Drosophila TRPγ Nat. Commun. 6, 7288

https://doi.org/10.1038/ncomms8288

42. Kwon, Y., Shim, H.-S., Wang, X., and Montell, C. (2008) **Control of**

thermotactic behavior via coupling of a TRP channel to a phospholipase C

signaling cascade *Nat. Neurosci.* **11**, 871 https://doi.org/10.1038/nn.2170

43. Watnick, T.J., Jin, Y., Matunis, E., Kernan, M.J., and Montell, C. (2003) **A**

flagellar polycystin-2 homolog required for male fertility in *Drosophila Curr.*

Biol. **13**, 2179-2184 https://doi.org/10.1016/j.cub.2003.12.002

44. Wang, T., Jiao, Y., and Montell, C. (2005). **Dissecting independent channel**

and scaffolding roles of the *Drosophila* **transient receptor potential**

channel *J. Cell Biol.* **171**, 685-694 https://doi.org/10.1083/jcb.200508030

45. Lee, Y., Lee, Y., Lee, J., Bang, S., Hyun, S., Kang, J., Hong, S.-T., Bae, E.,

Kaang, B.-K., and Kim, J. (2005) **Pyrexia is a new thermal transient receptor**

potential channel endowing tolerance to high temperatures in *Drosophila*

melanogaster Nat. Genet. **37**, 305 https://doi.org/10.1038/ng1513

46. Niemeyer, B.A., Suzuki, E., Scott, K., Jalink, K., and Zuker, C.S. (1996) **The**

Drosophila light-activated conductance is composed of the two channels

TRP and TRPL *Cell* **85**, 651-659 https://doi.org/10.1016/S0092-8674(00)81232-5

47. Tracey Jr, W.D., Wilson, R.I., Laurent, G., and Benzer, S. (2003) **painless, a**

Drosophila gene essential for nociception *Cell* **113**, 261-273

https://doi.org/10.1016/S0092-8674(03)00272-1

48. Kim, J., Chung, Y.D., Park, D.-y., Choi, S., Shin, D.W., Soh, H., Lee, H.W., Son,

W., Yim, J., and Park, C.-S. (2003) **A TRPV family ion channel required for**

hearing in Drosophila *Nature* **424**, 81-84 https://doi.org/10.1038/nature01733

49. Knapp, J.-M., Chung, P., and Simpson, J.H. (2015) **Generating customized**

transgene landing sites and multi-transgene arrays in Drosophila using

phiC31 integrase *Genetics* **199**, 919-934

https://doi.org/10.1534/genetics.114.173187

50. Meunier, N., Belgacem, Y.H., and Martin, J.-R. (2007) **Regulation of feeding**

behaviour and locomotor activity by takeout in *Drosophila* *J. Exp. Biol.* **210**,

1424-1434 https://doi.org/10.1242/jeb.02755

51. Dus, M., Min, S., Keene, A.C., Lee, G.Y., and Suh, G.S. (2011) **Taste-**

independent detection of the caloric content of sugar in *Drosophila* *Proc*

Natl Acad Sci U S A. **108**, 11644-11649

https://doi.org/10.1073/pnas.1017096108

52. Lee, Y., Poudel, S., Kim, Y., Thakur, D., and Montell, C. (2018) **Calcium taste**

avoidance in Drosophila *Neuron* **97**, 67-74. e64

https://doi.org/10.1016/j.neuron.2017.11.038

53. Yoshinari, Y., Kosakamoto, H., Kamiyama, T., Hoshino, R., Matsuoka, R., Kondo,

S., Tanimoto, H., Nakamura, A., Obata, F., and Niwa, R. (2021) **The sugar-**

responsive enteroendocrine neuropeptide F regulates lipid metabolism

through glucagon-like and insulin-like hormones in Drosophila

melanogaster *Nat. comm.* **12**, 4818 https://doi.org/10.1038/s41467-021-25146-

w

Figure Legends

Figure 1. TRPγ mutants exhibit altered carbohydrate, lipid, and protein levels. (**A**) TAG level measurement in the whole-body extract from control (w^{1118}) and TRP channel mutant lines (n=9). (**B**) Protein level measurement in the whole-body extract from control (w^{1118}) and TRP channel mutant lines (n=6). (**C**) Measurement of total glucose and trehalose levels (μg/mg) in the whole-body extracts of control (w^{1118}) and $trp\gamma^1$ adult males (n=6). (**D**) Measurement of tissue glycogen levels (μg/mg) in adult control (w^{1118}) and $trp\gamma^1$ males (n=6). (**E**) Measurement of Hemolymph glucose + trehalose level in the male flies of control (w^{1118}) and $trp\gamma^1$ (n=6). (**F**) Measurement of TAG level in adult males to test for rescue of the TAG defect in $trp\gamma$ flies with the *UAS-trpγ* and the $trp\gamma^{G4}$ or with the *trpγ* genomic transgene (n=6-10). (**G**) Rescue experiments showing the measurement of protein levels in adult flies with the indicated genotypes (n=6). Comparisons between multiple experimental groups were conducted via single-factor ANOVA coupled with Scheffe's *post hoc* test. The asterisks indicate significant differences from the controls (**P<0.01).

Figure 2. *Dh44* neurons are essential for regulating lipid tissue levels. (**A**) Tissue TAG level measurement in whole-body extracts of adult male flies after silencing of *Dh44-GAL4* and *dILP2-GAL4* with *UAS-Kir2.1* (n=6). (**B**) Measurement of tissue protein level in whole-body extracts of adult male flies. Brain-specific *Dh44-GAL4* and *dILP2-GAL4* neurons were ablated using *UAS-Kir2.1* (n=6). (**C**) Measurement of TAG from whole-body extract of adult male flies in the indicated genotypes (n=6). (**D**) Measurement of

tissue protein level from the whole-body extracts of adult males (n=6). (**E**) TAG levels in

whole-body extracts after RNAi knockdown of *trpγ* mutants under control of the *Dh44-*

GAL4 (n=6). (**F**) Measurement of area (μm^2) of LDs in adult fat body across the

indicated genotypes involved the selection of the 30 largest LDs, choosing the top 10

LDs from each sample for analysis (n=3). (**G**–**L**) Nile red stating of fat body from the

male of indicated genotypes. The scale bar represents 50 μm.

All values are reported as means ± SEM. Comparisons between multiple experimental

groups were conducted via single-factor ANOVA coupled with Scheffe's *post hoc* test.

The asterisks indicate significant differences from the controls (*$P<0.05$, **$P<0.01$).

Each dot indicates the distribution of individual sample values. (+) and (-) indicate the

presence or absence of the indicated transgenes, respectively.

Figure 3. *trpγ^1* have deficits in the lipolytic pathway under starvation conditions. (**A**)

Survival assay to measure the total survival time (days) of control (*w^{1118}*) and *trpγ^1* male

flies fed with a normal corn meal diet (n=8). (**B**) Survival assay to measure the survival

time (h) of the indicated genotypes with male flies under starvation conditions (n=4). (**C**)

TAG level measurement in control (*w^{1118}*) and *trpγ^1* adult male flies in both sated (0 h

starvation) and starved (24 h starvation) conditions (n=8). (**D**–**E**) Nile red staining of the

LDs extracted from FB of (D) *w^{1118}* and (E) *trpγ^1* flies under sated (**D1** and **E1**) and

starved (**D2** and **E2**) conditions, respectively. The scale bar represents 50 μm. (**F**)

Measurement of area (μm^2) of the LDs extracted from the FBs of *w^{1118}* and *trpγ^1* flies

under sated (0 h starvation) and starved (24 h starvation) conditions (n=3). (**G**)

Measurement of TAG level with controls and the flies after expressing *UAS-AMPK^{TD}*

under the control of *Dh44-GAL4* in the *trpγ^1* mutant background (n=4).

All values are reported as means ± SEM. Survival curves in A and B were estimated for

each group, using a Kaplan-Meier method and compared statistically using the log-rank

tests. Comparisons between multiple experimental groups in C, F, G, and H were

conducted via single-factor ANOVA coupled with Scheffe's *post hoc* test. The asterisks

indicate significant differences from the controls (**P<0.01).

Figure 4. Rescue of starvation susceptibility phenotype using the lipolytic drug

metformin**.** (**A**) TAG level measurement at 0, 7, and 14 days in the control (*w^{1118}*) and

trpγ^1 adult male flies after dietary exposure to 1 mM metformin (n=6). (**B**) TAG

measurement at 0, 7, and 14 days in *w^{1118}* and *trpγ^1* adult male flies after dietary

exposure to 5 mM metformin (n=6). (**C**) Pictures of Nile red staining of LDs after dietary

exposure to 1 mM and 5 mM metformin in standard fly food for 7 and 14 days in *w^{1118}*

and *trpγ^1* flies. The scale bars represent 50 μm. (**D**) Measurement of area (μm^2) of LDs

extracted from the FB of *w^{1118}* and *trpγ^1* flies after dietary exposure to 1mM and 5 mM

metformin in standard fly food for 7 and 14 days (n=3). (**E**) Survival assay to measure

the survival time (h) of control (*w^{1118}*) and *trpγ^1* males after dietary exposure to 1 mM

metformin in 1% agar food (n=6). (**F**) Starvation survival assay of control (*w^{1118}*) and

trpγ^1 males after dietary exposure to 5 mM metformin in 1% agar food (n=6).

All values are reported as means ± SEM. Comparisons between multiple experimental

groups were conducted via single-factor ANOVA coupled with Scheffe's *post hoc* test.

Each dot indicates the distribution of individual sample values. Survival curves in E and

F were estimated for each group, using a Kaplan-Meier method and compared

statistically using the log-rank tests. The asterisks indicate significant differences from

the controls (**P<0.01).

Figure 5. Dietary supply of lipase, glycerol, mixed TAG, and free FA to rescue the

starvation sensitivity phenotype. (**A**–**D**) Nile red staining of LDs in full gut and the

magnified R2 region of adult male flies. (**A**) w^{1118}, (**B**) $trp\gamma^1$, (**C**) $trp\gamma^1$,UAS-$trp\gamma$/$trp\gamma^{G4}$, (**D**)

$trp\gamma^1$,UAS-$trp\gamma$/$trp\gamma^1$;Dh44-GAL4/+. The scale bar represents 50 μm. The arrow indicates

the orientation of intestine from anterior to posterior. (**E**) Measurement of area of LDs in

the R2 region of $trp\gamma^1$ (n=10). Note that control and rescued flies have no LD. (**F**)

Survival assays of w^{1118} and $trp\gamma^1$ flies under starvation condition by feeding 0.1% lipase

or 0.1% denatured lipase mixed into 1% agar food (n=8). (**G**) Starvation survival assay

to measure the survival time (h) of w^{1118} and $trp\gamma^1$ male flies after feeding 1% glycerol

mixed into 1% agar food (n=4). (**H**) Survival assay to measure the survival time (h) of

w^{1118} and $trp\gamma^1$ flies after feeding 0.2% hexanoic acid (HA) supplemented into 1% agar

food (n=8). (**I**) Survival assay to measure the survival time (h) of w^{1118} and $trp\gamma^1$ flies

after feeding 0.2% mixed (mono-, di-, and tri-) glycerides (n=4).

All values are reported as means ± SEM. Survival curves were estimated for each

group, using a Kaplan-Meier method and compared statistically using the log-rank tests.

(**P<0.01).

Figure 6. Quantitative analysis of the expression of gluconeogenic, lipogenic, and

lipolytic genes and the effect of *bmm* expression and metformin feeding. (**A**) qRT-PCR

analysis to measure the expression of gluconeogenic genes (*fbp*, *pepck1*), lipogenic

genes (*acc*, *desat1*), and a lipolytic gene (*bmm*) under sated (0 h starvation) and

starved (24 h starvation) conditions in w^{1118} and $trp\gamma^1$ flies. The relative fold change in

the expression of starvation-induced genes (gluconeogenic genes: *fbp* and *pepck1*;

lipogenic genes: *acc* and *desat1;* and lipolytic gene: *bmm*) was determined in whole-

body samples of male adult w^{1118} and $trp\gamma^1$ flies by qRT-PCR. *Tubulin* was used as an

internal control to standardize the samples. Each graph shows the number of evaluated

samples (n=3). (**B**) Fluorescence microscopic imaging of *bmm*::*GFP* expression in the

FBs of w^{1118} and $trp\gamma^1$ under sated (0 h starvation) and starvation (24 h starvation)

conditions. The scale bar represent 50 μm. (**C**) Quantification of intensity level of

bmm::*GFP* in the FBs of w^{1118} and $trp\gamma^1$ under sated and starvation conditions. (**D–H**)

Nile red staining of lipids in the FB of flies with the indicated genotypes. The scale bar

represent 50 μm. (**I**) Measurement of area of LDs from samples D–H entailed selecting

a total of 30 LDs, with the 10 largest LDs chosen from each sample for analysis (n=3).

(**J**) qRT-PCR analysis to measure the expression of *acc*, *desat1*, and *bmm* from the

whole body samples of flies after feeding 5 mM metformin for 1 day (n=3).

All values are reported as means ± SEM. Comparisons between multiple experimental

groups were conducted via single-factor ANOVA coupled with Scheffe's *post hoc* test.

The asterisks indicate significant differences from the controls (*$P<0.05$, **$P<0.01$).

Figure 7. Functional analysis of *Dh44*, *Dh44R1*, and *Dh44R2* mutants and their roles in lipid accumulation. (**A**) Total TAG level (μg TAG/mg fly) measurement in whole body extracts from control (w^{1118}), $Dh44^{Mi}$, $Dh44R1^{Mi}$, and $Dh44R2^{Mi}$ mutants (n=5). (**B**–**E**) Nile red stains of the fat body from w^{1118}, $Dh44^{Mi}$, $Dh44R1^{Mi}$, and $Dh44R2^{Mi}$ respectively. Scale bars represent 50 μm. (**F**) Area of LDs in each indicated genotype (n=3). (**G**) The protein (μg protein/mg fly) measurement in the whole-body extracts from control (w^{1118}) and $Dh44^{Mi}$, $Dh44R1^{Mi}$, and $Dh44R2^{Mi}$ mutants (n=4). (**H**) Quantification (qRT-PCR) of lipolytic gene (bmm) expression level in the fat body of w^{1118}, $Dh44^{Mi}$, $Dh44R1^{Mi}$, and $D44R2^{Mi}$ flies under sated and starved conditions (n=3). (**I**) Total TAG level measurement in whole body extracts from the indicated genotypes (n=4). (**J**–**M**) Nile red stains of the fat body from (**J**) w^{1118}, (**K**) $trp\gamma^{1}$ (**L**) $trp\gamma^{1}$,UAS-$trp\gamma^{1}$/$trp\gamma^{1}$,Dh44R1-GAL4, and (**M**) $trp\gamma^{1}$,UAS-$trp\gamma^{1}$/$trp\gamma^{1}$,Dh44R2-GAL4. (**N**) Area of LDs in each indicated genotype (n=3). Scale bar represent 50 μm. (O-P) Expression of *Dh44R2* in the brain and intestine. (**O**) Expression of *Dh44R2* in the subesophageal zone (SEZ) of the brain. Scale bars represent 50 μm. (**P**) Expression of *Dh44R2* in the intestine. Scale bars represent 200 μm in full intestine and 500 μm in magnified form.

Means ±SEMs. Single factor ANOVA with Scheffe's analysis was used as a post hoc test to compare multiple sets of data. The asterisks indicate significance from control (*$P < 0.05$, **$P < 0.01$). Each dot indicates distribution of individual sample value.

Figure 1

Figure 2

Figure 3

A

survival (%) vs Time (days)

— w^{1118} — $trp\gamma^1$

B

survival % vs Time (h)

** n.s.
**

— w^{1118} — $UAS\text{-}dicer2;trp\gamma^{RNAi}$

— $trp\gamma^1$ — $UAS\text{-}dicer2;trp\gamma^{RNAi}/+;$ $Dh44\text{-}GAL4/+$

— $trp\gamma^1,UAS\text{-}trp\gamma/trp\gamma^1;Dh44\text{-}GAL4/+$

C

μg TAG/mg fly

** ** n.s.
**

w^{1118} $trp\gamma^1$

□ sated
■ starved

control $trp\gamma^1$

sated: D1 E1
starved: D2 E2

F

area of LDs (μm²)

n.s.
**

w^{1118} $trp\gamma^1$

□ sated
■ starved

G

μg TAG/mg fly

** ** **

$trp\gamma^1$	-	+	+	+	+
$UAS\text{-}AMPK^{TD}$	-	-	+	-	+
$Dh44\text{-}GAL4$	-	-	-	+	+

Figure 4

Figure 5

Figure 6

Figure 7