Semaglutide interferes with postnatal development of juvenile mice with

compromised growth

Amélie Joly[1]*, Lucas Rebiffé[1]*, Yves Dusabyinema[1], Julien Dellinger[1], Karine Gauthier[1],

François Leulier[1] & Filipe De Vadder[1] †

[1] Institut de Génomique Fonctionnelle de Lyon, École Normale Supérieure de Lyon, CNRS

UMR 5242, UCBL Lyon-1, F-69007 Lyon, France

* These authors contributed equally to this work

† To whom correspondence should be addressed: filipe.de_vadder@ens-lyon.fr

Keywords: semaglutide, stunting, somatotropic axis, energy metabolism

Grants and fellowships

This work was supported by funding from Fondation pour la Recherche Médicale (Équipe

FRM EQU202203014629), and the Agence Nationale de la Recherche (grant ANR PRC

2023 CE14 "GutStunting"). AJ was supported by a grant from Fondation pour la Recherche

Médicale (FDT202304016501) and LR is supported by a doctoral grant from the French

Ministry of Research.

Disclosure

The authors have nothing to disclose.

Abstract

GLP-1 receptor agonists, including semaglutide, are increasingly used to treat obesity—and

to a lesser extent, type 2 diabetes—in children and adolescents. Despite their therapeutic

promise, the impact of semaglutide on key endocrine and metabolic developmental

processes during this period is not yet fully understood. To address the limited understanding

of GLP-1 receptor agonists during development, we investigated the effects of chronic

semaglutide treatment on growth and metabolism of weanling male mice. In juvenile mice

displaying a normal growth pattern, semaglutide lowered blood glucose without affecting food

intake or weight gain. In contrast, chronic semaglutide treatment in growth-stunted juveniles

exacerbated linear growth impairment, despite preserving glucose-lowering effects.

Transcriptomic analysis of the liver, complemented by gene expression studies in other

metabolically active tissues, suggested altered energy expenditure, indicating a complex

interaction between semaglutide, energy metabolism, and growth during juvenile

development. These findings underscore the importance of developmental context when

evaluating the safety and efficacy of GLP-1 receptor agonists in pediatric populations.

<u>Introduction</u>

In recent years, GLP-1 receptor agonists (GLP-1RAs) have emerged as significant

therapeutic options for managing type 2 diabetes and obesity. As of 2025, seven GLP-1RAs

are approved by the US Food and Drug Administration for the treatment of type 2 diabetes[1].

Among them, semaglutide was optimized to have a longer half-life in humans and a better

resistance to Dipeptidyl Peptidase IV inhibitors than other GLP-1RAs[2]. Semaglutide improves

glycemic control through pleiotropic mechanisms, including increased insulin secretion in

response to glucose stimulation and improved beta cell survival and proliferation[3]. Besides its

effects on glucose homeostasis, semaglutide significantly lowers energy intake and induces

remarkable weight loss[1,4]. Hence, it is now widely used to treat obesity in adults[5,6], with

growing evidence supporting its efficacy in children as well[7,8]. Although semaglutide has

been widely studied in adult mouse models of diet-induced obesity, its effects during early

developmental windows, such as infancy and adolescence, remain poorly understood. Yet,

treatment timing may be critical, as semaglutide profoundly reshapes energy homeostasis.

The juvenile period, a phase characterized by intensive growth both in humans and mice, is

a highly energy-demanding process. In mammals, linear growth is regulated by the

somatotropic axis, governed by growth hormone (GH) and insulin-like growth factor 1 (IGF-

1). Interestingly, both *in vitro* and *in vivo* work in rodents demonstrated that GLP-1RAs can

promote growth-related phenotypes by enhancing osteoblast proliferation and activity[9,10].

Besides, several studies in mice have shown that GLP-1 and GLP-1RA may enhance IGF-1

receptor (IGF-1R) signaling pathways[11,12], favor bone metabolism homeostasis[9,13] and even

directly cross-activate the IGF-1R in human cardiac fibroblasts[14]. In addition, short- or long-

term administration of GLP-1RAs in healthy adult volunteers has been shown to increase

circulating GH levels[15]. Thus, it has been suggested that GLP-1RAs, such as semaglutide,

could interact with the somatotropic growth axis[16,17].

Despite these insights, how GLP-1RAs affect energy balance and growth at steady state

during the juvenile period remains largely unexplored. In this study, we examined the effects

of chronic semaglutide treatment on growth and energy metabolism in non-diabetic juvenile

mice under optimal and impaired growth conditions. We found that semaglutide improves

glucose clearance of non-obese, non-diabetic juvenile mice, irrespective of their growth
phenotype. However, while it does not affect growth in normally developing mice,
semaglutide further compromises linear growth in stunted juveniles. Notably, these growth-
inhibitory effects appear largely independent of alterations in the somatotropic axis.
Transcriptomic profiling of the liver, completed by gene expression analyses in other
metabolically active organs, suggests that semaglutide may impair growth in stunted
juveniles by downregulating pathways involved in basal energy expenditure and anabolic
metabolism.

Materials and methods

<u>Animals and treatments</u>

All animal procedures have been approved by a local animal care and use committee
(CECCAPP) and subsequently authorized by the French Ministry of Research (APAFIS
#40481-2022103017295302 v3).

21-day-old C57BL/6N male mice were purchased from Charles River and weaned onto either
the standard AIN-93G diet (Control Diet, CD) or on an isocaloric, protein-restricted
modification of the AIN-93G designed to induce stunting (Stunting-Inducing Diet, SID), as
previously described in our study[18]. The detailed composition of both diets is provided in
Table 1. Animals were weighted and body size was measured weekly under brief isoflurane
anesthesia. They received subcutaneous injections of semaglutide (Ozempic®, Novo
Nordisk 0.5 mg injectable solution) at a dose of 30 nmol/kg body weight, or vehicle (0.9%
NaCl), twice weekly until postnatal day 56 (P56). Injections were administered one day prior
to the oral glucose tolerance test (OGTT), and one day before sacrifice.

At P56, animals were fasted for 6 hours starting at 8 AM and killed by cervical dislocation at
2 PM. Blood and organs were sampled. The left femur and tibia were dissected and
measured precisely using a caliper.

For OGTT, mice were fasted for 6 hours starting at 8 AM and gavaged with a 20% D-(+)-
glucose solution (Sigma Aldrich, G7021) at 1 g/kg body weight. Blood glucose was measured
from the tail vein at 0 (baseline), 15, 30, 60, 90, and 120 min post-gavage using a handheld
glucometer (OneTouch Verio Reflect, LifeScan Europe). Blood samples were collected from
the tail vein at 0, 15 and 30 min, serum was extracted and stored at −20°C until further
processing.

Only male mice were included in this study, as we previously demonstrated that SID induces
greater growth impairment in juvenile males compared to females[18].

<u>Insulin measurements</u>

Serum insulin levels were measured using the Ultra Sensitive Insulin ELISA kit (Crystal
Chem, 90080), following the manufacturer's instructions.

<u>Western blotting</u>

Frozen liver samples were processed for Western blotting as described by Schwarzer et
al.[19]. Briefly, samples were lysed in RIPA Plus lysis buffer (50 mM Tris, 1% IGEPAL, 0.5%
sodium deoxycholate, 0.1% SDS, 150 mM NaCl, 2 mM EDTA, 50 mM NaF, protease
inhibitors). Protein concentration was measured using the BCA assay (Thermo Scientific,
23227). Equal amounts of protein (30 µg per well) were separated in Mini-PROTEAN TGX
Stain Free precast gels (Bio-Rad, 4868096), and transferred to nitrocellulose membranes
using the Trans-Blot Turbo system (Bio-Rad). Membranes were blocked and incubated for 1
hour at room temperature with either rabbit monoclonal anti-total Akt (C67E7) (1:1,000, Cell
Signaling, #4691), or rabbit monoclonal anti-phospho S473 Akt (D9E) (1:2,000, Cell
signaling, #4060). After washing, membranes were incubated for 1 hour at room temperature
with HRP-conjugated anti-Rabbit IgG secondary antibody (1:2,500, Promega, W4011).
Detection was performed using ECL Western Blotting Substrate (Promega, W1001), and
signals were captured with the Amersham IQ800 imager (Cytiva). Quantification was
performed using Image Lab software version 6.1 (Bio-Rad), using total protein
normalization[20].

<u>IGF-1 quantification</u>

Liver tissues were weighed and homogenized in lysis buffer containing PBS, 1% Triton X100
and protease inhibitors (Roche, cOmplete™, EDTA-free). Homogenates were centrifuged
(12,000 x g, 10 min, 4°C), supernatants were used to quantify IGF-1 levels using the
Mouse/Rat IGF-1 Quantikine ELISA (R&D systems, MG100). Total protein content was
measured using the BCA assay for normalization. IGF-1 concentrations are expressed as
pg/µg total protein. The same kit was used to quantify circulating IGF-1.

<u>Gene expression analysis</u>

Total RNA was extracted from brown adipose tissue (BAT) and subcutaneous white adipose
tissue (scWAT) using TRI Reagent (Invitrogen, M9738). Briefly, tissues were homogenized in
TRI reagent, centrifuged to remove debris, and RNA was isolated via chloroform phase
separation followed by isopropanol precipitation. For liver and skeletal muscle, total RNA was
extracted using the Nucleospin RNA kit (Macherey-Nagel, 740955.250).

Reverse transcription was performed on 1 µg of RNA using the SensiFAST cDNA Synthesis
Kit (Meridian Biosciences, BIO-65053). Quantitative PCR was performed using Takyon No
ROX SYBR Mastermix blue dTTP (Eurogentec, UF-NSMT-B0701) using a Bio-Rad CFX96
apparatus. Analysis followed MIQE guidelines[21]. Amplification efficiency was evaluated for
each primer pair using standard curve assays performed prior to experimental runs. All
reactions were carried out in technical duplicates to ensure reproducibility. At least two
validated reference genes were used for normalization, with gene stability confirmed using
the geNorm algorithm available within the CFX96 software. Primers sequences for all
reference and target genes are listed in Table 2. Data are expressed as geometric means of
relative expression normalized to the reference genes.

<u>Liver RNA-seq analysis</u>

The concentration and quality of 10 DNAse-treated RNA samples (from SID-fed juveniles; 5
vehicle-treated and 5 semaglutide-treated) were assessed using the Qubit 4.0 fluorometer
(Thermo Fisher Scientific) and the Tapestation 4150 system (Agilent Technologies),
respectively. For each sample, mRNA was purified from 1 µg of total RNA using oligo(dT)-
based magnetic bead isolation (Poly(A) mRNA Magnetic Isolation Module, New England
Biolabs), following the manufacturer's protocol.

Single-indexed libraries were built using the CORALL Total RNA-Seq Library Prep Kit
(Lexogen), according to the manufacturer's instructions. This fragmentation-free protocol
ensures full-length transcript coverage. Unique Molecular Identifiers (UMIs; 12bp tags) were
introduced during the linker Ligation step, enabling the identification and removal of PCR
duplicates during downstream analysis to reduce amplification bias.

After quality control (Qubit 4.0 and Tapestation) analysis, libraries were pooled together in
equimolar concentrations and sequenced on two high-output runs in a single-end mode
(1×86 bp) using a NextSeq500 Illumina sequencer (Illumina). A total of 1.082 billion reads
were generated, yielding an average of 36 million reads per sample (range: 31-44 million),
after demultiplexing.

Read preprocessing was performed using a custom pipeline derived from Lexogen's
recommendations. Briefly, adapter trimming was performed using the Cutadapt package

(version 3.5)[22] and read quality was assessed using FastQC. Read mapping was achieved
with STAR (Spliced Transcripts Alignment to a Reference, version 2.7.11a)[23] using the *Mus*
musculus GRCm39 genome reference (Ensembl, release 110). UMI-based deduplication
was carried out using UMI-tools dedup (version 1.1.2)[24], enabling accurate removal of PCR
duplicates. Gene-level quantification was performed with htseq-count (version 2.0.3)[25],
generating raw count tables for each sample.

Differential analysis was performed in R (v.4.3.1), using the DESeq2 package (v.1.42.0)[26].
Visualization of differential gene expression was done using the Enhanced Volcano package
(v.1.20.0)[27], applying cut-offs of adjusted p-value (FDR) ≤ 0.01 and $|\log_2$ Fold Change$| \geq 1$.
One vehicle-treated sample was excluded from analysis following unsupervised clustering
and quality control, as it appeared to be an outlier.

The RNA-seq dataset has been deposited in the NCBI Gene Expression Omnibus[28] (GEO)
under   accession   number   GSE266477
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE266477).

Statistical analysis

Statistical analysis was performed using GraphPad Prism v10.2.1 and R v. 4.4.3. For
comparisons between two groups, unpaired *t*-tests were used when data met assumptions of
normality and homogeneity of variance; otherwise, non-parametric Mann-Whitney *U*-tests
were applied.

Longitudinal data—including body weight, linear size, and glycemia during the OGTT—were
analyzed using linear mixed-effects models in R, employing the lme4 (v. 1.1-4), lmerTest (v.
1.1-36), and emmeans (v. 1.11.0) packages. Mouse ID was included as a random effect to
account for repeated measures. Two independent experimental batches were used, and
batch was included in all models as a fixed effect along with its interactions with treatment
and age. Although the fixed effects involving batch were not statistically significant (p > 0.05),
model comparison showed that including the full $\mathrm{Treatment} \times \mathrm{Age} \times \mathrm{Batch}$ interaction
significantly improved model fit. Additionally, the variance attributed to batch as a random
effect was consistently small. Therefore, to account for possible batch-related structure while
minimizing visual redundancy, representative data from one batch are shown in the figures.
Linear mixed-effects model results are reported in Table 3.

Results

Chronic semaglutide treatment improves glucose tolerance without altering growth of non-diabetic non-obese juvenile mice

To determine the effects of chronic semaglutide treatment during the juvenile developmental window, we characterized glucose metabolism of animals fed an experimental diet meeting the nutritional requirements for optimal growth conditions (AIN-93G[29], hereafter referred to as control diet, CD), and treated with either vehicle (Veh) or semaglutide (Sema). Consistent with findings in adult mouse models of obesity[30], chronic semaglutide treatment lowered fasting glycemia and insulinemia in juvenile mice (Fig. 1A-B). Semaglutide also enhanced glucose clearance during an OGTT (Fig. 1C), likely reflecting increased insulin sensitivity (Fig. 1D). In contrast to its known effects in mouse models of diet-induced obesity[31], chronic semaglutide treatment did not reduce body weight or food intake in juvenile animals maintained under optimal nutritional conditions (Fig. 1E-F). Linear growth was also unaffected, as shown by similar body size (Fig. 1G), and bone measurements (femur and tibia lengths; Fig. 1H-I) at the end of the treatment. Assessment of the somatotropic axis revealed no detectable changes in circulating GH/IGF-1 signaling pathway (Fig. S1), consistent with the absence of growth alterations. Furthermore, semaglutide treatment did not impact body composition, with similar weights of subcutaneous white adipose tissue and gastrocnemius muscle between groups (Fig. 1J-K). Altogether, these results show that chronic semaglutide treatment modulates glucose metabolism in healthy juvenile mice without affecting growth dynamics or body composition.

Chronic semaglutide treatment exacerbates growth impairment in stunted juvenile mice

Given the importance of linear growth as a hallmark of juvenile physiology and the observed metabolic modulation by semaglutide under optimal nutritional conditions, we next investigated its effects in a context of disrupted growth. Stunting, defined as low height-for-age, was induced in juvenile males by feeding a low-protein variant of the AIN-93G (referred to as the stunting-inducing diet, SID; Table 1)[18]. Unexpectedly, semaglutide treatment further impaired linear growth in SID-fed juveniles, resulting in a 7% reduction in body size compared to vehicle-treated controls, along with corresponding reductions in femur and tibia length (Fig. 2A-C). These findings indicate that semaglutide specifically disrupts body development under a growth-limiting nutritional environment, highlighting the context-dependency of its physiological effects.

Semaglutide is known to reduce body weight, largely via appetite suppression, in patients with obesity[5,6] and in adult mouse models[30,31]. In SID-fed juveniles, semaglutide reduced weight gain (32% lower than vehicle-treated counterparts), but notably did not reduce food

intake (Fig. 2D-E), indicating that the observed growth impairment is not attributable to
reduced caloric intake.

Given previous reports of GLP-1 receptor agonists interfering with IGF-1 signaling[16], we
examined the somatotropic axis in SID-fed mice treated with semaglutide. Circulating IGF-1
levels were reduced in semaglutide-treated animals (Fig. 2F), consistent with impaired
growth. However, liver expression of *Ghr* and *Igf1*, as well as hepatic IGF-1 protein levels,
were unchanged between semaglutide- and vehicle-treated animals (Fig. 2G–I), arguing
against a disruption of GH signaling at the hepatic level. Since IGF-1 promotes anabolic
processes in peripheral tissues such as muscle[32], we further assessed IGF-1 signaling in this
tissue. Semaglutide treatment did not affect *Igf1r* expression or downstream AKT
phosphorylation in skeletal muscle (Fig. 2J-K). Taken together, these results do not support
major alterations in somatotropic axis function as the primary mechanism underlying
semaglutide's growth-inhibitory effects in stunted juveniles.

Interestingly, semaglutide-treated SID-fed mice displayed altered body composition, with
lower relative weights of subcutaneous white adipose tissue and gastrocnemius muscle (Fig.
2L-M). In addition, semaglutide improved glycemic response to OGTT and lowered fasting
glycemia, similar to effects observed in CD-fed animals (Fig. S2A-B). However, fasting
insulinemia and HOMA-IR were not significantly altered (Fig. S2C-D). Taken together, these
findings demonstrate that chronic semaglutide treatment modulates energy metabolism and
body composition in growth-impaired juveniles, independently of food intake or major
changes in somatotropic axis activity.

<u>Semaglutide alters energy metabolism–related gene expression in metabolically active</u>
<u>tissues of stunted juvenile mice</u>

To better understand how semaglutide impacts growth-compromised juveniles, we
performed transcriptomic analysis of the liver, a central organ in both metabolic and growth
regulation. Among ~13,000 detected genes, only 126 were differentially expressed between
semaglutide- and vehicle-treated animals (47 upregulated, 79 downregulated) with no
signature of alterations in metabolic- or growth-related processes, indicating relatively mild
hepatic transcriptional reprogramming (Fig. 3A).

Given the limited hepatic response, we next examined semaglutide's impact on selected
gene's expression profiles in peripheral metabolically active tissues that are more direct
targets of GLP-1 signaling: brown adipose tissue (BAT), subcutaneous white adipose tissue
(scWAT), and gastrocnemius muscle. We focused on genes involved in key energy-
expending pathways, including fatty acid oxidation (*Cpt1b, Lpl, Acadm, Acadl*),

thermogenesis (*Ucp1, Ucp3, Pgc1a, Dio2*), basal energy metabolism (*Sdhb, Pfkm*), and
energy-wasting futile cycles (*Alpl, mCBK, Serca2a, Serca2b*).

In BAT (Fig. 3B), semaglutide-treated mice showed reduction of thermogenesis-related
genes (*Ucp1* and *Pgc1a*) and *Cpt1b*, suggesting decreased thermogenic and oxidative
activity. Interestingly, *Lpl* expression was slightly increased, possibly reflecting a
compensatory mechanism—despite reduced thermogenic drive. Expression of futile cycle
markers was mixed, with decreased *Alpl* and increased *mCBK*, suggesting altered energy-
wasting activity, though the direction of change remains unclear. These data point to altered
energy expenditure in BAT following semaglutide treatment in stunted juveniles.

In scWAT (Fig. 3C), *Ucp1* was similarly downregulated, indicating reduced beigeing capacity.
In gastrocnemius muscle (Fig. 3D), we observed decreased expression of *Acadl* and *Acadm*
(fatty acid oxidation enzymes), along with lower *Sdhb* levels, a marker of mitochondrial
oxidative capacity. Collectively, these transcriptional changes evoke that semaglutide may
suppress catabolic and energy-expending pathways along multiple peripheral tissues.

Catabolic activity is essential during growth tissues to sustain the high energy requirements
of tissue development. Therefore, reduced thermogenic and oxidative capacity in
semaglutide-treated stunted juveniles may contribute to an overall energy-conserving
phenotype that is incompatible with normal tissue growth.

Discussion

Understanding the potential impacts of the increasingly prescribed GLP-1 receptor agonist
semaglutide in the clinics is essential to guarantee its safety, particularly as growing
evidence highlights its broad physiological effects. However, the mechanisms underlying
some of these actions remain poorly defined. Given the expanding use of semaglutide in
pediatric populations, we aimed to investigate its interaction with a key developmental
process: linear growth.

In juvenile mice raised under nutritionally optimal conditions, semaglutide enhanced glucose
clearance, consistent with effects described in adult rodents and humans[3,33]. Notably, these
metabolic effects were uncoupled from changes in energy balance considerations, as food
intake and weight gain remained unchanged. To our knowledge, this is the first study to
report a dissociation between semaglutide's effects on glucose homeostasis and weight gain
in juveniles, suggesting age-specific mechanisms of action[17]. Chronic semaglutide treatment
did not affect linear growth in these animals, consistent with clinical studies in children with
obesity treated with semaglutide or liraglutide, which reported no evidence of altered growth
outcomes[7,34,35].

As linear growth is a key indicator of juvenile health, we explored the potential interaction
between GLP-1 receptor agonists and the somatotropic growth axis[16], in the context of
growth disruption. Using a model of diet-induced stunting, we found that semaglutide further
impaired linear growth of those juveniles, despite maintaining its beneficial effects on glucose
management. However, our investigation of the GH/IGF-1 somatotropic axis did not reveal
major alterations in hormone production and signaling, challenging our initial hypothesis that
semaglutide's effects on growth might be mediated through disruption of this pathway.

Semaglutide has been shown to influence many physiological processes regulating energy
expenditure[36–38]. Given that semaglutide- and vehicle-treated stunted juveniles had similar
food intake, we hypothesized that the observed growth impairment could be due to reduced
energy availability, rather than decreased intake. Transcriptomic and gene expression
analyses in metabolically active tissues revealed signs of decreased catabolic activity and
suppressed thermogenesis, particularly in BAT. These findings suggest that semaglutide
treatment in stunted juveniles leads to a shift in energy allocation—reducing energy-
expensive processes such as tissue anabolism and thermogenesis, which may impair
growth.

In conclusion, our study reveals unexpected consequences of chronic semaglutide treatment
during juvenile development in mice, particularly under growth-limiting nutritional conditions.
Our findings challenge the notion that GLP-1R agonist–mediated improvements in glucose
metabolism are solely driven by reduced food intake and highlight the importance of
developmental timing and nutritional context when evaluating safety and efficacy of GLP-1
receptor agonists. These results underscore the need for further studies to investigate the
integrated mechanisms of action of GLP-1 receptor agonists across physiological states and
life stages.

Acknowledgments

The authors thank Benjamin Gillet and Sandrine Hughes from the sequencing platform of
IGFL (PSI) for their help regarding the sequencing project. They also thank Raghav Galgali
for his help in setting up the bioinformatics pipeline for liver transcriptomic data analysis, the
animal facility of IGFL (PEHR) where animal experiments were conducted, and the
associated group for animal welfare (SBEA).

Data availability

The RNA sequencing dataset generated in this study has been deposited in the NCBI Gene

Expression Omnibus (GEO) under accession number GSE266477.

Some or all datasets generated during and/or analyzed during the current study are not

publicly available but are available from the corresponding author on reasonable request.

References

1.   Yao H, Zhang A, Li D, et al. Comparative effectiveness of GLP-1 receptor agonists on
glycaemic control, body weight, and lipid profile for type 2 diabetes: systematic review and network
meta-analysis. *BMJ*. 2024;384:e076410. doi:10.1136/bmj-2023-076410
2.   Lau J, Bloch P, Schäffer L, et al. Discovery of the Once-Weekly Glucagon-Like Peptide-1 (GLP-
1) Analogue Semaglutide. *J Med Chem*. 2015;58(18):7370-7380. doi:10.1021/acs.jmedchem.5b00726
3.   Kapitza C, Dahl K, Jacobsen JB, Axelsen MB, Flint A. Effects of semaglutide on beta cell
function and glycaemic control in participants with type 2 diabetes: a randomised, double-blind,
placebo-controlled trial. *Diabetologia*. 2017;60(8):1390-1399. doi:10.1007/s00125-017-4289-0
4.   Blundell J, Finlayson G, Axelsen M, et al. Effects of once-weekly semaglutide on appetite,
energy intake, control of eating, food preference and body weight in subjects with obesity. *Diabetes
Obes Metab*. 2017;19(9):1242-1251. doi:10.1111/dom.12932
5.   Wilding JPH, Batterham RL, Calanna S, et al. Once-Weekly Semaglutide in Adults with
Overweight or Obesity. *N Engl J Med*. 2021;384(11):989-1002. doi:10.1056/NEJMoa2032183
6.   Gao X, Hua X, Wang X, et al. Efficacy and safety of semaglutide on weight loss in obese or
overweight patients without diabetes: A systematic review and meta-analysis of randomized
controlled trials. *Front Pharmacol*. 2022;13. doi:10.3389/fphar.2022.935823
7.   Weghuber D, Barrett T, Barrientos-Pérez M, et al. Once-Weekly Semaglutide in Adolescents
with Obesity. *N Engl J Med*. 2022;387(24):2245-2257. doi:10.1056/NEJMoa2208601
8.   Fox CK, Barrientos-Pérez M, Bomberg EM, et al. Liraglutide for Children 6 to <12 Years of Age
with Obesity — A Randomized Trial. *N Engl J Med*. 2025;392(6):555-565.
doi:10.1056/NEJMoa2407379
9.   Zhang M, Xie Y, Zhou Y, et al. Exendin-4 enhances proliferation of senescent osteoblasts
through activation of the IGF-1/IGF-1R signaling pathway. *Biochem Biophys Res Commun*.
2019;516(1):300-306. doi:10.1016/j.bbrc.2019.06.112
10.   Ma X, Meng J, Jia M, et al. Exendin-4, a glucagon-like peptide-1 receptor agonist, prevents
osteopenia by promoting bone formation and suppressing bone resorption in aged ovariectomized
rats. *J Bone Miner Res Off J Am Soc Bone Miner Res*. 2013;28(7):1641-1652. doi:10.1002/jbmr.1898
11.   Cornu M, Yang JY, Jaccard E, Poussin C, Widmann C, Thorens B. Glucagon-like peptide-1
protects beta-cells against apoptosis by increasing the activity of an IGF-2/IGF-1 receptor autocrine
loop. *Diabetes*. 2009;58(8):1816-1825. doi:10.2337/db09-0063
12.   Arakawa M, Ebato C, Mita T, et al. Effects of exendin-4 on glucose tolerance, insulin
secretion, and beta-cell proliferation depend on treatment dose, treatment duration and meal
contents. *Biochem Biophys Res Commun*. 2009;390(3):809-814. doi:10.1016/j.bbrc.2009.10.054
13.   Yamada C, Yamada Y, Tsukiyama K, et al. The murine glucagon-like peptide-1 receptor is
essential for control of bone resorption. *Endocrinology*. 2008;149(2):574-579. doi:10.1210/en.2007-
1292
14.   Qa'aty N, Wang Y, Wang A, et al. The antidiabetic hormone glucagon-like peptide-1 induces
formation of new elastic fibers in human cardiac fibroblasts after cross-activation of IGF-IR.
Endocrinology. 2015;156(1):90-102. doi:10.1210/en.2014-1519
15.   Heinla K, Vasar E, Reppo I, Sedman T, Volke V. GLP-1 Receptor Agonists Induce Growth
Hormone Secretion in Healthy Volunteers. *Diabetes Ther Res Treat Educ Diabetes Relat Disord*.
2023;14(4):777-786. doi:10.1007/s13300-023-01381-w

16.   Cignarelli A, Genchi VA, Le Grazie G, et al. Mini Review: Effect of GLP-1 Receptor Agonists and
SGLT-2 Inhibitors on the Growth Hormone/IGF Axis. *Front Endocrinol*. 2022;13. Accessed September
7, 2023. https://www.frontiersin.org/articles/10.3389/fendo.2022.846903
17.   Cooper DM, Rothstein MA, Amin A, Hirsch JD, Cooper E. Unintended consequences of
glucagon-like peptide-1 receptor agonists medications in children and adolescents: A call to action. *J*
Clin Transl Sci. 7(1):e184. doi:10.1017/cts.2023.612
18.   Thoumas JL, Cavaroc A, Sery D, Leulier F, De Vadder F. Suboptimal Refeeding Compensates
Stunting in a Mouse Model of Juvenile Malnutrition. *J Nutr*. Published online January 3, 2025:S0022-
3166(25)00002-1. doi:10.1016/j.tjnut.2025.01.002
19.   Schwarzer M, Makki K, Storelli G, et al. Lactobacillus plantarum strain maintains growth of
infant mice during chronic undernutrition. *Science*. 2016;351(6275):854-857.
doi:10.1126/science.aad8588
20.   Fosang AJ, Colbran RJ. Transparency Is the Key to Quality. *J Biol Chem*. 2015;290(50):29692-
29694. doi:10.1074/jbc.E115.000002
21.   Bustin SA, Benes V, Garson JA, et al. The MIQE guidelines: minimum information for
publication of quantitative real-time PCR experiments. *Clin Chem*. 2009;55(4):611-622.
doi:10.1373/clinchem.2008.112797
22.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal. 2011;17(1):10-12. doi:10.14806/ej.17.1.200
23.   Dobin A, Davis CA, Schlesinger F, et al. STAR: ultrafast universal RNA-seq aligner. *Bioinforma*
Oxf Engl. 2013;29(1):15-21. doi:10.1093/bioinformatics/bts635
24.   Smith T, Heger A, Sudbery I. UMI-tools: modeling sequencing errors in Unique Molecular
Identifiers to improve quantification accuracy. *Genome Res*. 2017;27(3):491-499.
doi:10.1101/gr.209601.116
25.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-throughput
sequencing data. *Bioinforma Oxf Engl*. 2015;31(2):166-169. doi:10.1093/bioinformatics/btu638
26.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8
27.   EnhancedVolcano. Bioconductor. Accessed February 16, 2025.
http://bioconductor.org/packages/EnhancedVolcano/
28.   Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res*. 2002;30(1):207-210. doi:10.1093/nar/30.1.207
29.   Reeves PG, Nielsen FH, Fahey GC. AIN-93 Purified Diets for Laboratory Rodents: Final Report
of the American Institute of Nutrition Ad Hoc Writing Committee on the Reformulation of the AIN-
76A Rodent Diet. *J Nutr*. 1993;123(11):1939-1951. doi:10.1093/jn/123.11.1939
30.   Niu S, Chen S, Chen X, et al. Semaglutide ameliorates metabolism and hepatic outcomes in an
NAFLD mouse model. *Front Endocrinol*. 2022;13:1046130. doi:10.3389/fendo.2022.1046130
31.   Gabery S, Salinas CG, Paulsen SJ, et al. Semaglutide lowers body weight in rodents via
distributed neural pathways. *JCI Insight*. 2020;5(6):e133429, 133429. doi:10.1172/jci.insight.133429
32.   Shavlakadze T, Chai J, Maley K, et al. A growth stimulus is needed for IGF-1 to induce skeletal
muscle hypertrophy in vivo. *J Cell Sci*. 2010;123(Pt 6):960-971. doi:10.1242/jcs.061119
33.   Tadross JA, Steuernagel L, Dowsett GKC, et al. A comprehensive spatio-cellular map of the
human hypothalamus. *Nature*. Published online February 5, 2025:1-9. doi:10.1038/s41586-024-
08504-8
34.   Kelly AS, Auerbach P, Barrientos-Perez M, et al. A Randomized, Controlled Trial of Liraglutide
for Adolescents with Obesity. *N Engl J Med*. 2020;382(22):2117-2128. doi:10.1056/NEJMoa1916038
35.   Diene G, Angulo M, Hale PM, et al. Liraglutide for Weight Management in Children and
Adolescents With Prader-Willi Syndrome and Obesity. *J Clin Endocrinol Metab*. 2022;108(1):4-12.
doi:10.1210/clinem/dgac549
36.   Moiz A, Filion KB, Tsoukas MA, Yu OHY, Peters TM, Eisenberg MJ. Mechanisms of GLP-1
receptor agonist-induced weight loss: A review of central and peripheral pathways in appetite and
energy regulation. *Am J Med*. Published online January 31, 2025. doi:10.1016/j.amjmed.2025.01.021

37.   Kurtzhals P, Flindt Kreiner F, Singh Bindra R. The role of weight control in the management of
type 2 diabetes mellitus: Perspectives on semaglutide. *Diabetes Res Clin Pract*. 2023;203:110881.
doi:10.1016/j.diabres.2023.110881
38.   Tamayo-Trujillo R, Ruiz-Pozo VA, Cadena-Ullauri S, et al. Molecular mechanisms of
semaglutide and liraglutide as a therapeutic option for obesity. *Front Nutr*. 2024;11.
doi:10.3389/fnut.2024.1398059
446

Figure 1. **Semaglutide modulates glucose metabolism without affecting growth in control-fed juvenile mice**. (A) Fasting glycemia and (B) insulinemia. (C) Glycemia during an oral glucose tolerance test (OGTT). (D) HOMA-IR. (E) Body weight over time. (F) Cumulative food intake. (G) Body size. (H-I) Femur (H) and tibia (I) lengths at postnatal day 56 (P56). (J-K) Weights of subcutaneous white adipose tissue (WAT) (J) and gastrocnemius muscle (K) at P56. Veh, vehicle-treated; Sema, semaglutide-treated. Data are shown as mean ± 95% CI, except for cumulative food intake (mean ± SEM). $^*p < 0.05$, $^{**}p < 0.01$, $^{****}p < 0.0001$, ns: not significant.

Figure 2. **Semaglutide further impairs linear growth and alters body composition in stunted juvenile mice**. (A) Body size. Femur (B) and tibia (C) lengths at postnatal day 56 (P56). (D) Body weight over time. (E) Cumulative food intake. (F-K) Somatotropic axis markers: serum IGF-1 concentration (F), liver *Igf1* expression (G), liver IGF-1 concentration (H), liver *Ghr* expression (I), muscle *Igf1r* expression (J) and phosphorylated Akt (S473) to total Akt ratio (K) measured by Western blot and normalized to total protein content. (L-M) Relative weights of subcutaneous WAT (L) and gastrocnemius muscle (M) at P56. Veh, vehicle-treated; Sema, semaglutide-treated. Data are shown as mean ± 95% CI, except for cumulative food intake (mean ± SEM). $^*p < 0.05$, $^{**}p < 0.01$, $^{****}p < 0.0001$, ns: not significant.

Figure 3. **Semaglutide alters energy metabolism gene expression in metabolically active tissues in stunted juvenile mice**. (A) Volcano plot showing differential gene expression in the liver of SID-fed juveniles treated with semaglutide (N = 5) versus vehicle (N = 4). (B-D) Relative expression of energy metabolism-related genes in brown adipose tissue (BAT, B), subcutaneous WAT (C), and gastrocnemius muscle (D). The reference genes used to normalize gene expression are: *Rpl32* and *Rsp18* for BAT and WAT; *Gapdh* and *Tbp* for muscle. Veh, vehicle-treated; Sema, semaglutide-treated. Data are shown as mean ± 95% CI. $^*p < 0.05$, $^{**}p < 0.01$, $^{****}p < 0.0001$.

Supplementary figure captions

Figure S1. **Semaglutide treatment does not alter the somatotropic axis in control-fed juvenile mice**. (A) Schematic representation of the somatotropic GH/IGF-1 axis. (B-G) Somatotropic axis markers at P56: serum IGF-1 (B), liver *Igf1* expression (C), liver IGF-1

concentration (D), liver *Ghr* expression (E), muscle *Igf1r* expression (F), and phosphorylated

Akt (S473) to total Akt ratio (G) measured by Western blot and normalized to total protein

content. Veh, vehicle-treated; Sema, semaglutide-treated. Data are shown as mean ± 95%

CI. ns: not significant.

<u>Figure S2.</u> **Semaglutide treatment lowers blood glucose and enhances glucose**

clearance in stunted juvenile mice. (A) Glycemia during an oral glucose tolerance test

(OGTT). (B-C) Fasting glycemia (B) and insulinemia (C). (D) HOMA-IR. Veh, vehicle-treated;

Sema, semaglutide-treated. Data are shown as mean ± 95% CI. **p < 0.01, ***p < 0.001, ns:

not significant.