A systematic review of laboratory investigations into the pathogenesis of avian influenza viruses
in wild avifauna of North America

Matthew Gonnerman[1,2], Christina Leyson[3,4], Jeffery D. Sullivan[5], Mary J. Pantin-Jackwood[3],
Erica Spackman[3], Jennifer M. Mullinax[2], Diann J. Prosser[5]

[1] Post-doctoral affiliate with the U.S. Geological Survey, Eastern Ecological Science Center,
Laurel, MD, 20708

[2] Department of Environmental Science and Technology, University of Maryland, College Park,
MD 20742

[3] Exotic and Emerging Avian Viral Diseases unit, U.S. National Poultry Research Center, Agricultural
Research Service, U.S. Department of Agriculture, Athens, GA 30602

[4] Department of Microbiology and Immunology, School of Medicine, Emory University, Emory,
GA 30322

[5] U.S. Geological Survey, Eastern Ecological Science Center, Laurel, MD 20708

ABSTRACT

The lack of consolidated information regarding wild bird species' response to avian influenza
virus (AIV) infection is a challenge for conservation managers, researchers, and related sectors
such as public health and commercial poultry. Such information could be used to model complex
disease dynamics within communities, prioritize species for surveillance and conservation
efforts, or identify species more likely to facilitate spillover into domestic animals or humans.
Using two independent searches, we reviewed published literature for studies describing wild
bird species experimentally infected with avian influenza to assess host species' relative
susceptibility to AIVs. Additionally, we summarized broad-scale parameters for elements such
as shedding duration and minimum infectious dose that can be used in transmission modeling
efforts. Our synthesis documented that waterfowl (i.e., Anatidae) comprise the vast majority of
published AIV pathobiology studies, whereas gulls and passerines were less represented in
research despite evidence that they also are susceptible and contribute to high pathogenicity
avian influenza disease dynamics. This study represents the first comprehensive effort to compile
available literature regarding the pathobiology of AIV's in all wild birds in over a decade. This
database provides an opportunity to critically examine and assess what is known and identify
where further insight is needed.

Keywords: avian influenza, challenge study, pathogenicity, wild bird

INTRODUCTION

While low pathogenic avian influenza (LPAI) virus circulates naturally with minimal clinical
signs in waterbirds in the orders Anseriformes and Charadriiformes [1,2], high pathogenicity
avian influenza (HPAI) virus historically emerges following mutation or reassortments gained by
LPAI viruses circulating in gallinaceous poultry [3], often originally introduced by wild
waterfowl (typically Anatidae). The recently circulating and expanding HPAI H5 viruses of the

Goose/Guangdong/1/1996 (Gs/GD) lineage are unusual because they have adapted to be carried
by certain wild waterfowl species, thereby expanding exposure to poultry, wild birds, and more
recently some mammals [4,5]. The original H5 Gs/GD virus was isolated in 1996 in southern
China [6] and by 2013 had evolved into clade 2.3.4.4 with a notably improved ability to infect
wild waterfowl including species that are generally not carriers of LPAI viruses [7,8]. In October
2020, the global reach of H5 Gs/GD viruses was again highlighted when clade 2.3.4.4b viruses
emerged, before spreading to over 80 countries in the following years [5]. This current outbreak
has been especially notable for reaching new hemispheres as it moved into North America in
2021 [9], South America in late 2022 [10] and the Antarctic region in 2023 [11], causing extreme
impact on numerous populations of wild birds such as the Peruvian Pelican (*Pelecanus thagus*),
Sandwich Tern (*Thalasseus sandvicensis*), and Common Crane (*Grus grus*) [4].

During this recent panzootic of HPAI, the conservation community was challenged to anticipate
how diverse species might respond when exposed to novel viral strains. For instance, there has
been great concern about potential impacts on penguin colonies and condor populations should
AIV be introduced [12] and recent outbreaks in Antarctica and South America indicate the
vulnerability of these species [11,12]. However, an extensive body of work has been produced
by researchers focused on challenge studies with wild bird species in a laboratory setting with
both LPAI and HPAI to determine factors including infectiousness, severity of clinical signs, and
recovery potential [13,14]. However, given the limited resources available, such work has not
encompassed the broad spectrum of potential species and viral strains [15]. This has left a
substantial knowledge gap as AIVs continue to evolve and present risks for an ever-growing list
of species [16]. Furthermore, the available literature is diffuse with no comprehensive review for
wild species collated in nearly a decade [14,17].

The lack of consolidated information on the response of wild bird species to infection with AIV
is an identified challenge to both conservation managers and researchers alike [16], with
additional impacts on related sectors such as public health and commercial poultry [18,19]. For
wildlife managers, it is important to understand the possible implications of an HPAI outbreak,
especially for sensitive populations [4]. For instance, knowing if a population is at-risk to
exposure and resilient to infection may play an important role in how that species is prioritized in
allocation of resources for habitat protection, captive breeding programs, and other programs.
Similarly, the diffuse knowledge base makes identifying data-informed parameters for
transmission risk models more difficult [16,18]. While we can infer some information from wild
surveillance and mortality data, such as which species have tested positive and which are most
likely associated with mortality events, we cannot determine important metrics like length of
infection or infectivity from these databases. The goal of this study is to collate all previous
pathobiology studies on wild avifauna in North America, providing insight into current gaps in
the available literature that may be filled by future targeted research. Additionally, we aim to
develop broad scale parameters for elements such as shedding duration and minimum infectious
dose that can be used in transmission modeling efforts until species and strain specific results
become available.

METHODS

Literature Search

Using two independent searches, we reviewed the published literature for studies in which wild
bird species were experimentally infected with avian influenza to assess the host species' relative
susceptibility to AIVs and ability to shed viruses. To provide reasonable bounds on which
species we included in our search, we created a list of species from those reported to the U.S.
Department of Agriculture Animal and Plant Health Inspection Service (USDA-APHIS)
database [20] and the U.S. National Institute of Allergy and Infectious Diseases (NIAID)-
sponsored Bioinformatics Resource Center Influenza Research Database (IRD) [21,22] as of Oct
16, 2023. We then supplemented this initial list with species that were both a high-priority
conservation concern and at high risk of exposure to HPAI due to either shared habitat with or a
close genetic relationship to known susceptible species. We conducted an initial search using
Google Scholar during October and November of 2023 using the search criteria '"avian
influenza" AND "[species name]" AND "challenge study" OR "experimental infection"'. The
search was conducted twice for each species of interest, where [species name] was replaced with
either the common or scientific name. This initial search was intended to produce a
representative, but not exhaustive, sample of research available for use in modeling avian
influenza disease dynamics. As some species produced a disproportionately large number of
papers (+1000) compared to the average (0–50), we limited our search to the first 5 pages of
Google Scholar results to balance search time and capturing relevant information for all species.

While limiting search efforts likely contributed to missing papers, there are also concerns that
Google Scholar is not an appropriate method for reproducible reviews of scientific literature
[23,24]. So, after completion of this initial search, we identified several relevant studies
providing information on unrepresented HPAI lineages not identified by Google Scholar by
conducting a second systematic review using Web of Science. We expanded our search criteria
as using the same search terms as the original review proved too restrictive when applied to Web
of Science. We chose to simplify phrasing, remove quotations, and include additional language
previously used in reference to challenge studies, resulting in a search criteria of '(TS=("avian
influenza" AND "[species name]") AND TS=(challenge OR experiment OR characterization OR
pathogenesis))'. We conducted the search as initially described for our first review, repeating the
search for each common and scientific name for birds on our search list.

Identifying Relevant Studies

Articles reviewed herein were considered for further review if the title, abstract, or methods
discussed AIV challenge studies for wild bird species that range within North America. We
excluded information for non-avian influenza viruses (i.e., mammal-derived strains of AIV) as
well as information related to domestic and hybrid host bird species. There were multiple
instances when a study challenged an individual with multiple viral strains in succession. In such
cases, we recorded the viral subtype of the previous infection but otherwise, recorded this
information as usual, as many wild animals are likely to have previous AIV infections. However,
several studies examined the impact of vaccines or viral inhibitors on host birds or local

environments (i.e., water treatments), in which case we recorded these studies in the database but
did not include them in any summaries.

For studies that met our above criteria, we recorded a unique entry for each combination of host
bird species, virus subtype, and challenge method (e.g., direct inoculation, contact, meat-fed)
used within a study. For each entry, we recorded information regarding the host animal species,
family, and age at inoculation, as well as the viral subtype, origin, whether the virus was a HPAI
or LPAI pathotype, and HA genetic lineage. AIV lineages for subtypes other than
goose/Guangdong/1996 H5 were described based on the HA gene of the most closely related
grouping classified by naming conventions utilized in previous reports or general geographic and
species origin from which similar isolates have been collected. Subtype H5 isolates were
classified by clade if from the goose/Guangdong/1996 lineage [25].

To describe the experimental design of each study, we recorded inoculation method, inoculation
dose, and swabbing schedule, with any further details that differentiated experimental groups
within a study provided in a notes column. For each entry, we recorded the number of
individuals in the study, the number of individuals identified as infected with AIV (i.e., virus
identified in swabs, blood, or organs), and the number of mortalities. We separately counted
direct mortalities from illness versus euthanasia due to symptoms deemed indicative of imminent
mortality. In some studies, individuals were euthanized (hereafter "culled") at predetermined
time points throughout the study to assess viral titers in organs and tissue. We recorded these
individuals separately from the prior two mortality categories to adjust estimates of mortality rate
due to AIV infection. Information on mortality and culling did not include individuals culled at
the end of the study (i.e., survivors). If provided, or if it could be calculated from the provided
mortality data, we recorded the mean death time in days.

Summarizing Viral Response

Across studies, viral response is measured and reported in a variety of ways which limited
comparison among studies. Identification of viruses in swab samples can be performed using
virus isolation or real-time RT-PCR methods[26], which are reported as at least one of multiple
metrics that are not all comparable. Virus titer levels in isolates are commonly quantified and
reported as the $\log_{10}$ 50% egg infective dose per milliliter (EID_{50}/mL); however, alternative units
can be used that produce estimates that are not comparable between methods [27]. Real-time RT-
PCR results are readily reported as a cycle threshold (Ct) value, but additional steps need to be
conducted to produce a comparable equivalent to the EID_{50}/mL obtained from viral isolation
[28]. As such, we only used information measured as EID_{50}/mL from viral isolation or, if not
available, EID_{50}/mL equivalents from RT-PCR in our summaries. For simplicity, we hereafter
use "shedding" to refer to the greatest daily-average viral titers identified in swabs. Information
on shedding could be collected via swabbing the oropharyngeal (OP) and/or the cloacal (CL)
routes, which are important prerequisites of animal-to-animal and environmental HPAI
transmission respectively [13]. As shedding patterns between these pathways may have
important but different implications for virus transmission, we recorded this information
separately. We recorded information on viral shedding by the host animal, however this
information is assessed and summarized in multiple formats across studies that are not directly

comparable. For example, viral shedding quantities and durations were reported using a broad
range of summary statistics, including mean, median, and maximum values for a given
experimental group as well as by individual, sometimes for multiple time points throughout an
infection. Thus, to describe the average capacity a host animal has for spreading the virus, we
report the greatest mean daily shedding quantities for a given experimental group over the course
of a study (Figure S1).

For all entries, we recorded the mean daily shedding quantities for a given experimental group
and mean bird infectious dose (units in $\log_{10}$ 50% bird infectious doses per milliliter; BID_{50}) if
estimated. We also recorded shedding duration (days that an animal shed the virus), either as a
single duration estimate where provided, or as a series of time points describing when the virus
was first detected (D_1; Figure S1), which day the greatest mean shedding rate occurred (D_{Max}),
the last day that the virus was detected in swabs (D_L), and the next day animals were swabbed
and virus was not detected following completion of shedding (D_{Neg}). From this information and
information on when swabbing occurred, we can calculate a shedding duration as the (D_L +
D_{Neg})/2 – D_1. In multiple instances, we encountered studies where shedding information was
presented only in plot form. To interpret information within these papers, we used a web-based
figure reader (PlotDigitizer; plotdigitizer.com) which is a free option for such data mining from
scatter and line plots [29] and recorded its use.

Tables including entries for all inoculation doses, inoculation methods, and prior infection
statuses were generated. To produce a final set of directly comparable metrics describing a
species' viral response, we subset our dataset to entries where animals were directly inoculated
with a Gs/GD lineage HPAI of dose greater than 4 $\log_{10} EID_{50}$, which is a cutoff that consistently
resulted in infection in most studies that estimated infectious dose.

Following the course of a successful infection of challenged individuals, we can further describe
immune response for both animals that recovered and those that succumb to the disease. From
surviving animals or animals euthanized during the course of recovering from infection, antibody
production can be assessed via serological assays such as bELISA [30] to determine the strength
and length of antibody response. In our review, we recorded whether a study collected
serological information to describe antibody production following infection, as well as the
maximum number of individuals recorded as positive for antibodies at a given timepoint. We can
further study the severity and location of infection through necropsies of infected individuals,
either to identify antigens or histological lesions in tissue samples [31,32]. If only a local
infection has occurred, antigens or lesions will be isolated to the respiratory or digestive tract of
the animal, whereas systematic infections of HPAI viruses can be identified throughout the body,
indicating a more severe infection. Unfortunately, which organs were sampled varied amongst
studies and studies were inconsistent in how they reported data (i.e., ranging from individual-
level data to study-level summaries). Thus, we only present whether histopathology information
was collected, and whether a group of animals had any individuals with local (i.e., lungs, trachea,
digestive tract only) or systematic (i.e., any other organs) infections, identified by either lesions
or antigens in a tissue sample. We note that immune response will be temporally variable relative
to exposure as when any given individual was sampled will dramatically impact results.

  Comparison to Incidence of Avian Influenza in Wild Birds

To identify potential knowledge gaps in AIV research, we compared our review of the published research to the observed incidence of avian influenza in wild birds. Information on wild bird infections was obtained via the USDA-APHIS database [20] and accessed on Nov 16, 2023. The database provided information on encounter location, encounter date, viral subtype, host species, and how the animal was reported, and was summarized to the number of positives per species.

RESULTS

Summary of Relevant Studies

From our review database [52], 132 unique challenge studies for AIV in wild bird species were identified, and of these, 104 had at least one focal species from our list (Supplemental 1, Figure S2). Of these 104 studies, Anatidae was the most well-represented family identified (78 studies), followed by Laridae (11), Passeridae (8), Columbidae (6), and Scolopacidae (4; Figure 1, Table S1). Within Anatidae (Table S2), challenge studies were distributed across 18 species including mallard (*Anas platyrhynchos*; 54 studies), Muscovy duck (*Cairina moschata*; 11), Canada goose (*Branta candadensis*; 5), northern pintail (*Anas acuta*; 4), and wood duck (*Aix sponsa*; 4). Outside of Anatidae, house sparrow (*Passer domesticus*; 8), laughing gull (*Leucophaeus atricilla*; 7), and rock pigeon (*Columba livia*; 6) were the most studied. Of note, 6 of the top 10 families with the most reported AIV infections had no studies identified within our review (Figure 2), including Accipitridae (727 reported infections), Cathartidae (691), Strigidae (248), Pelecanidae (119), Corvidae (118), and Phalacrocoracidae (39). Forty-three studies challenged animals with LPAI subtypes and 74 with HPAI subtypes, of which 66 included viruses of the Gs/GD HPAI lineage (Figure 1). Only 5 of the 1282 animals challenged with LPAI in the identified studies perished over the course of the infection. Animals challenged with Gs/GD perished with greater frequency (67%, N = 1755) compared to those challenged with other lineages of HPAI (6%, N = 300). Information on viral susceptibility and shedding was only available from one species for 6 of the 10 families identified in the review, and for 3 of these families only 1 study was found (Table S3-S5). One of these, Ardeidae, had challenge data for only 3 individuals, of which 1 was infected and died.

Summarizing Viral Response

Articles were subsetted according to challenge type (i.e., directly inoculated seronegative animals with >4 $\log_{10}$ EID_{50}/mL of a Gs/GD HPAI viruses) then summarized (Figure 3, Table S3,S4). The resulting dataset contained information on only 24 species, of which 13 are from the Anatidae family. Within Anatidae, American black duck (*Anas rubripes*), blue-winged teal (*Spatula discors*), lesser scaup (*Aythya affinis*), Northern pintail, and redhead (*Aythya americana*) all exhibited 0% mortality when challenged with Gs/GD whereas Canada goose, mute swan (*Cygnus olor*), and trumpeter swan (*Cygnus buccinator*) all experienced 100% mortality. For waterfowl, oral shedding (measured by RNA presence from OP swabs) rates averaged 4.54 $\log_{10}$ EID_{50} and ranged from 1.1 $\log_{10}$ EID_{50} to 7.7 EID_{50} while cloacal shedding (measured by RNA presence from cloacal swabs (CL)) rates averaged 3.28 $\log_{10}$ EID_{50} and ranged from 0.9 $\log_{10}$ EID_{50} to 6.8 $\log_{10}$ EID_{50}. Shedding durations averaged 5.72 and 5.21 days

for oral and cloacal routes respectively. Trumpeter and mute swans exhibited the highest average
oral (6.14 $\log_{10}$ EID$_{50}$) and cloacal (4.46 $\log_{10}$ EID$_{50}$) shedding rates respectively. The top eight
oral shedders and seven of the top eight cloacal shedders were all from the Anatidae family. A
number of these species, such as mallards and surf scoter (*Melanitta perspicillata*), also
exhibited relatively low mortality rates (31% and 22 %) when challenged with an HPAI virus.
Mallards exhibited higher than average shedding rates (4.78 $\log_{10}$ EID$_{50}$ via OP, 3.52 $\log_{10}$ EID$_{50}$
via CL) and durations (6.42 days via OP, 5.83 days via CL) across studies.

We identified only a limited number of comparable studies for non-waterfowl species. For
example, within Laridae, we identified herring gull (*Larus argentatus*) and laughing gull as the
only species with data available. Herring gull exhibited high infection rates (100%) and moderate
mortality rates (67%) when challenged and shed more via the OP route (3.82 $\log_{10}$ EID$_{50}$ for 5.27
days) than the cloaca (2.01 $\log_{10}$ EID$_{50}$ for 2.93 days). Comparatively, laughing gull had lower
infection (64%) and mortality (29%) rates, longer shedding durations (9 days and 6.5 days for
OP and CL), but similar shedding rates (3.48 $\log_{10}$ EID$_{50}$ and 1.89 $\log_{10}$ EID$_{50}$). Considering
passerines, house sparrows had relatively high infection (78%) and mortality (58%) rates and
shed moderately via both the OP and CL routes (3.28 $\log_{10}$ EID$_{50}$ via OP, 2.7 $\log_{10}$ EID$_{50}$ via CL)
for approximately 4.8 days. Limited information was also available for American robins (*Turdus
migratorius*), which were readily infected (92%) but showed no indication of negative impacts to
survival (0% mortality). Rock pigeons, a similarly broadly abundant bird species, were much less
readily infected (30%) and had similarly low mortality rates (7%). Within Falconidae, two
studies challenged a total of 22 American kestrels (*Falco sparverius*), of which all perished.

When the dataset was expanded to include any animal challenged with any dose of Gs/GD HPAI
(Table S5), we found Canada goose exhibited the greatest average viral titers in cloacal swabs
(5.75 $\log_{10}$ EID$_{50}$/mL, range 3.8-7.3) and trumpeter swan in OP swabs (6.14 $\log_{10}$ EID$_{50}$/mL).
Laughing gulls had the longest observed shedding duration via the OP route (9 days post
challenge, range 8-10), whereas surf scoters shed the longest via the CL route (8 days post
challenge).

DISCUSSION

This study represents the first comprehensive effort to compile available literature regarding the
pathobiology of AIVs in all wild birds of North America in over a decade [13]. The database
produced through this effort can serve as a tool for all researchers, providing generalized
estimates of pathobiology parameters such as mortality and viral shedding for a variety of wild
avian families. This database also provides the opportunity to critically examine previous
research to assess what has been learned from these efforts as well as to identify where additional
targeted efforts can provide further insight. However, we acknowledge the limitations of this
resource, as the variability in methods used to assess viral response amongst studies limits our
ability to compare among them. It should also be noted that infections in wild versus laboratory
settings can produce divergent results [33], which should motivate further research when
encountered. Thus we urge users to be thoughtful in their application of this database, and for
future research to be exhaustive in their reporting and consistent with current common practices.

Our synthesis shows that waterfowl (i.e., Anatidae) comprise the vast majority of published AIV pathobiology studies. Such an emphasis is warranted given the widely recognized role of these species in the spread of AIVs [18,34]. Indeed, previous efforts have demonstrated that *Anatidae* are competent vectors for avian influenza viruses due to their high shedding rates [35,36], long duration of infection [37,38], and comparatively high prevalence in the wild [39,40]. However, while previous studies have provided us with valuable information on the general response to infection for Anseriformes, the importance of these species to the transmission of AIVs both across space [34,41] and across the wild waterfowl – domestic poultry interface [18,42] suggests that future work with these species could aim to develop better estimates of parameters needed for transmission risk modeling [43] such as minimum infectious dose which is currently only rarely included [44,45]. Similarly, the continued high representation of Anatidae in mortality events and surveillance sampling [20] emphasizes the importance of studies focused on including emerging viral strains to inform possible outcomes prior to outbreak events and to maintain current data.

There was a dramatic decrease in available research between Anseriformes and Laridae, the second most studied family group with 10 studies across 4 species. Laridae as a group had a high capacity for infection and shedding, consistent with the recognized importance of Laridae in the inter-continental spread and persistence of AIV [46,47]. Given the apparent susceptibility of these species in the current HPAI outbreak, their large geographic distribution, long-distance migrations, as well as frequent overlap with agriculture, Laridae seem to be prime candidates for continued pathobiology research [15]. For similar reasons, other sea birds, such as skuas [48], have been implicated as possible long-distance carriers of the virus, but a lack of available research limits our ability to infer potential dynamics.

While identified as having a role in the spatial movement of AIVs [49], predatory and scavenging species appear especially susceptible to infection and mortality based on data from the current outbreak [20], as well as limited pathobiology studies with species outside North America [50]. In fact, Accipitridae, Cathartidae, and Strigidae were three of the top four families with the most reported infections to USDA yet had no studies were identified by our review. However, Falconidae had among the highest infectious rates and mortality rates (100% for each), though this finding was based on a single species across two studies (the American Kestrel). Given the dramatic impacts of HPAI on these species there is a clear need to improve the understanding of all aspects of pathobiology of AIV in raptors and other predatory avian species. Additionally, efforts to model broader ecological transmission and persistence could benefit from studies that explore transmission through meat/carcass consumption instead of traditional inoculation [51].

Of course, many other species outside of those specifically mentioned above have been impacted by the ongoing outbreak of HPAI. Our summary demonstrates that seven families had over 100 reported HPAI infections to USDA [20], but only two of these families have been studied (Anatidae and Laridae). Of the 10 families identified in our review, only three reported challenge data from more than 1 species, and 1 of these only included three individuals in total. It should also be noted that these metrics rely on data from infections reported to USDA [20], so species

that are under-sampled due to low detectability of sick or dead birds will also be underrepresented in this effort. Therefore, the real number of species that lack sufficient pathobiology data is likely larger than demonstrated here. This is partly because challenge studies with HPAI require BSL-3 laboratories with specialized spaces for avian species, which are limited and often have overriding study priorities. Additionally, it can be difficult to obtain and care for wild birds or wild bird eggs needed for conducting these experiments. Working to close this large data gap is a notable potential avenue for future research, even for species that are not suffering from large-scale mass mortality events and that do not have recognized roles in AIV spread.

Acknowledgements

This work was funded by the U.S. Geological Survey, Ecosystem Mission Area. All data reported in this manuscript are publicly available [52]. Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Literature Cited

1. Kuiken, T. Is Low Pathogenic Avian Influenza Virus Virulent for Wild Waterbirds? *Proceedings of the Royal Society B: Biological Sciences* **2013**, *280*, 20130990, doi:10.1098/rspb.2013.0990.

2. Fouchier, R.A.M.; Munster, V.J. Epidemiology of Low Pathogenic Avian Influenza Viruses in Wild Birds. *Rev. sci. tech. int. Epiz.* **2009**, *28*, 49–58.

3. Sonnberg, S.; Webby, R.J.; Webster, R.G. Natural History of Highly Pathogenic Avian Influenza H5N1. *Virus Res* **2013**, *178*, 63–77.

4. Wille, M.; Waldenström, J. Weathering the Storm of High Pathogenicity Avian Influenza in Waterbirds. *Waterbirds* **2023**, *46*, doi:10.1675/063.046.0113.

5. Harvey, J.A.; Mullinax, J.M.; Runge, M.C.; Prosser, D.J. The Changing Dynamics of Highly Pathogenic Avian Influenza H5N1: Next Steps for Management & Science in North America. *Biol Conserv* **2023**, *282*, 110041, doi:10.1016/j.biocon.2023.110041.

6. Xu, X.; Subbarao, K.; Cox, N.J.; Guo, Y. Genetic Characterization of the Pathogenic Influenza A/Goose/Guangdong/1/96 (H5N1) Virus: Similarity of Its Hemagglutinin Gene to Those of H5N1 Viruses from the 1997 Outbreaks in Hong Kong. *Virology* **1999**, *261*, 15–19, doi:10.1006/viro.1999.9820.

7. Lee, D.H.; Bertran, K.; Kwon, J.H.; Swayne, D.E. Evolution, Global Spread, and Pathogenicity of Highly Pathogenic Avian Influenza H5Nx Clade 2.3.4.4. *J Vet Sci* 2017, *18*, 269–280.

8. Kang, H.-M.; Lee, E.-K.; Song, B.-M.; Jeong, J.; Choi, J.-G.; Jeong, J.; Moon, O.-K.; Yoon, H.; Cho, Y.; Kang, Y.-M.; et al. Novel Reassortant Influenza a(H5N8) Viruses among Inoculated Domestic and Wild Ducks, South Korea, 2014. *Emerg Infect Dis* **2015**, *21*, 298–304, doi:10.3201/eid2102.141268.

9. Caliendo, V.; Lewis, N.S.; Pohlmann, A.; Baillie, S.R.; Banyard, A.C.; Beer, M.; Brown, I.H.; Fouchier, R.A.M.; Hansen, R.D.E.; Lameris, T.K.; et al. Transatlantic Spread of Highly Pathogenic Avian Influenza H5N1 by Wild Birds from Europe to North America in 2021. *Sci Rep* **2022**, *12*, doi:10.1038/s41598-022-13447-z.

10. Gamarra-Toledo, V.; Plaza, P.I.; Gutiérrez, R.; Luyo, P.; Hernani, L.; Angulo, F.; Lambertucci, S.A. Avian Flu Threatens Neotropical Birds. *Science (1979)* **2023**, *379*, 246.

11. Bennison, A.; P Byrne, A.M.; Reid, S.M.; Lynton-Jenkins, J.G.; De Sliva, D.; Peers-Dent, J.; Finlayson, K.; Hall, R.; Blockley, F.; Blyth, M.; et al. Detection and Spread of High Pathogenicity Avian Influenza Virus H5N1 in the Antarctic Region. *bioRxiv* **2023**, doi:10.1101/2023.11.23.568045.

12. Leguia, M.; Garcia-Glaessner, A.; Muñoz-Saavedra, B.; Juarez, D.; Barrera, P.; Calvo-Mac, C.; Jara, J.; Silva, W.; Ploog, K.; Amaro, Lady; et al. Highly Pathogenic Avian Influenza A (H5N1) in Marine Mammals and Seabirds in Peru. *Nat Commun* **2023**, *14*, doi:10.1038/s41467-023-41182-0.

13. Hénaux, V.; Samuel, M.D. Avian Influenza Shedding Patterns in Waterfowl: Implications for Surveillance, Environmental Transmission, and Disease Spread. *J Wildl Dis* **2011**, *47*, 566–578, doi:10.7589/0090-3558-47.3.566.

14. Pantin-Jackwood, M.J.; Swayne, D.E. Pathogenesis and Pathobiology of Avian Influenza Virus Infection in Birds. *Rev. sci. tech. Off. int. Epiz* **2009**, *28*, 113–136.

15. Arnal, A.; Vittecoq, M.; Pearce-Duvet, J.; Gauthier-Clerc, M.; Boulinier, T.; Jourdain, E. Laridae: A Neglected Reservoir That Could Play a Major Role in Avian Influenza Virus Epidemiological Dynamics. *Crit Rev Microbiol* 2015, *41*, 508–519.

16. Ramey, A.M.; Hill, N.J.; Deliberto, T.J.; Gibbs, S.E.J.; Hopkins, M.C.; Lang, A.S.; Poulson, R.L.; Prosser, D.J.; Sleeman, J.M.; Stallknecht, D.E.; et al. Highly Pathogenic Avian Influenza Is an Emerging Disease Threat to Wild Birds in North America. *Journal of Wildlife Management* **2022**, *2022*, 1–21, doi:10.1002/jwmg.22171.

17. Abolnik, C. A Current Review of Avian Influenza in Pigeons and Doves (Columbidae). *Vet Microbiol* 2014, *170*, 181–196.

18. Blagodatski, A.; Trutneva, K.; Glazova, O.; Mityaeva, O.; Shevkova, L.; Kegeles, E.; Onyanov, N.; Fede, K.; Maznina, A.; Khavina, E.; et al. Avian Influenza in Wild Birds and Poultry: Dissemination Pathways, Monitoring Methods, and Virus Ecology. *Pathogens* **2021**, *10*, 1–23, doi:10.3390/pathogens10050630.

19. Philippon, D.A.M.; Wu, P.; Cowling, B.J.; Lau, E.H.Y. Avian Influenza Human Infections at the Human-Animal Interface. *Journal of Infectious Diseases* **2020**, *222*, 528–537, doi:10.1093/infdis/jiaa105.

20. USDA 2022-2023 Detections of Highly Pathogenic Avian Influenza in Wild Birds.

21. Zhang, Y.; Aevermann, B.D.; Anderson, T.K.; Burke, D.F.; Dauphin, G.; Gu, Z.; He, S.;
Kumar, S.; Larsen, C.N.; Lee, A.J.; et al. Influenza Research Database: An Integrated
Bioinformatics Resource for Influenza Virus Research. *Nucleic Acids Res* **2017**, *45*,
D466–D474, doi:10.1093/nar/gkw857.

22. Squires, R.B.; Noronha, J.; Hunt, V.; García-Sastre, A.; Macken, C.; Baumgarth, N.;
Suarez, D.; Pickett, B.E.; Zhang, Y.; Larsen, C.N.; et al. Influenza Research Database: An
Integrated Bioinformatics Resource for Influenza Research and Surveillance. *Influenza*
Other Respir Viruses **2012**, *6*, 404–416, doi:10.1111/j.1750-2659.2011.00331.x.

23. Haddaway, N.R.; Collins, A.M.; Coughlin, D.; Kirk, S. The Role of Google Scholar in
Evidence Reviews and Its Applicability to Grey Literature Searching. *PLoS One* **2015**, *10*,
doi:10.1371/journal.pone.0138237.

24. Gusenbauer, M.; Haddaway, N.R. Which Academic Search Systems Are Suitable for
Systematic Reviews or Meta-Analyses? Evaluating Retrieval Qualities of Google Scholar,
PubMed, and 26 Other Resources. *Res Synth Methods* **2020**, *11*, 181–217,
doi:10.1002/jrsm.1378.

25. Hnevolution, W.; Group, W. Toward a Unified Nomenclature System for Highly
Pathogenic Avian Influenza Virus (H5N1). *Emerg Infect Dis* **2008**, *14*, e1,
doi:10.3201/eid1407.071681.

26. Van Den Brand, J.M.A.; Verhagen, J.H.; Veldhuis Kroeze, E.J.B.; Van De Bildt, M.W.G.;
Bodewes, R.; Herfst, S.; Richard, M.; Lexmond, P.; Bestebroer, T.M.; Fouchier, R.A.M.;
et al. Wild Ducks Excrete Highly Pathogenic Avian Influenza Virus H5N8 (2014-2015)
without Clinical or Pathological Evidence of Disease Article. *Emerg Microbes Infect*
2018, *7*, doi:10.1038/s41426-018-0070-9.

27. Krauss, S.; Walker, D.; Webster, R.G. Influenza Virus Isolation. In *Influenza virus:*
methods in molecular biology; Kawaoka, Y., Neumann, G., Eds.; Humana Press, 2012;
Vol. 865.

28. Lee, C.-W.; Suarez, D.L. Application of Real-Time RT-PCR for the Quantitation and
Competitive Replication Study of H5 and H7 Subtype Avian Influenza Virus. *J Virol*
2004, *119*, 151–158.

29. Aydin, O.; Yassikaya, M.Y. Validity and Reliability Analysis of the PlotDigitizer
Software Program for Data Extraction from Single-Case Graphs. *Perspect Behav Sci*
2022, *45*, 239–257, doi:10.1007/s40614-021-00284-0.

30. Webster, R.; Cox, N.; Storh, K. *WHO Manual on Animal Influenza Diagnosis and*
Surveillance; Geneva, Switzerland, 2002;

31. Caliendo, V.; Leijten, L.; van de Bildt, M.W.G.; Poen, M.J.; Kok, A.; Bestebroer, T.;
Richard, M.; Fouchier, R.A.M.; Kuiken, T. Long-Term Protective Effect of Serial
Infections with H5N8 Highly Pathogenic Avian Influenza Virus in Wild Ducks. *J Virol*
2022, *96*, doi:10.1128/jvi.01233-22.

32. Soda, K.; Tomioka, Y.; Hidaka, C.; Matsushita, M.; Usui, T.; Yamaguchi, T. Susceptibility of Common Family Anatidae Bird Species to Clade 2.3.4.4e H5N6 High Pathogenicity Avian Influenza Virus: An Experimental Infection Study. *BMC Vet Res* **2022**, *18*, doi:10.1186/s12917-022-03222-7.

33. Umar, S.; Guerin, J.L.; Ducatez, M.F. Low Pathogenic Avian Influenza and Coinfecting Pathogens: A Review of Experimental Infections in Avian Models. *Avian Dis* **2017**, *61*, 3–15, doi:10.1637/11514-101316-Review.

34. Gorsich, E.E.; Webb, C.T.; Merton, A.A.; Hoeting, J.A.; Miller, R.S.; Farnsworth, M.L.; Swafford, S.R.; DeLiberto, T.J.; Pedersen, K.; Franklin, A.B.; et al. Continental-Scale Dynamics of Avian Influenza in U.S. Waterfowl Are Driven by Demography, Migration, and Temperature. *Ecological Applications* **2020**, *0*, 1–12, doi:10.1002/eap.2245.

35. Spackman, E.; Pantin-Jackwood, M.J.; Lee, S.A.; Prosser, D. The Pathogenesis of a 2022 North American Highly Pathogenic Clade 2.3.4.4b H5N1 Avian Influenza Virus in Mallards (*Anas Platyrhynchos*). *Avian Pathology* **2023**, 1–10, doi:10.1080/03079457.2023.2196258.

36. Brown, J.D.; Stallknecht, D.E.; Swayne, D.E. Experimental Infection of Swans and Geese with Highly Pathogenic Avian Influenza Virus (H5N1) of Asian Lineage. *Emerg Infect Dis* **2008**, *14*, 136–142, doi:10.3201/eid1401.070740.

37. Stephens, C.B.; Prosser, D.J.; Pantin-Jackwood, M.J.; Berlin, A.M.; Spackman, E. The Pathogenesis of H7 Highly Pathogenic Avian Influenza Viruses in Lesser Scaup (*Aythya Affinis*). *Avian Dis* **2019**, *63*, 230–234.

38. Van Den Brand, J.M.A.; Verhagen, J.H.; Veldhuis Kroeze, E.J.B.; Van De Bildt, M.W.G.; Bodewes, R.; Herfst, S.; Richard, M.; Lexmond, P.; Bestebroer, T.M.; Fouchier, R.A.M.; et al. Wild Ducks Excrete Highly Pathogenic Avian Influenza Virus H5N8 (2014-2015) without Clinical or Pathological Evidence of Disease. *Emerg Microbes Infect* **2018**, *7*, 7–67, doi:10.1038/s41426-018-0070-9.

39. Kent, C.M.; Ramy, A.M.; Ackerman, J.T.; Bahl, J.; Bevins, S.N.; Bowman, A.S.; Boyce, W.; Cardona, C.; Casazza, M.L.; Cline, T.D.; et al. Spatiotemporal Changes in Influenza A Virus Prevalence among Wild Waterfowl Inhabiting the Continental United States throughout the Annual Cycle. *Sci Rep* **2022**, *12*.

40. Kent, C.M.; Bevins, S.N.; Mullinax, J.M.; Sullivan, J.D.; Prosser, D.J. Waterfowl Show Spatiotemporal Trends in Influenza A H5 and H7 Infections but Limited Taxonomic Variation. *Ecological Applications* **2023**.

41. Dusek, R.J.; Hallgrimsson, G.T.; Ip, H.S.; Jónsson, J.E.; Sreevatsan, S.; Nashold, S.W.; TeSlaa, J.L.; Enomoto, S.; Halpin, R.A.; Lin, X.; et al. North Atlantic Migratory Bird Flyways Provide Routes for Intercontinental Movement of Avian Influenza Viruses. *PLoS One* **2014**, *9*, doi:10.1371/journal.pone.0092075.

42. Humphreys, J.M.; Ramey, A.M.; Douglas, D.C.; Mullinax, J.M.; Soos, C.; Link, P.;
Walther, P.; Prosser, D.J. Waterfowl Occurrence and Residence Time as Indicators of H5
and H7 Avian Influenza in North American Poultry. *Sci Rep* **2020**, *10*, 1–16,
doi:10.1038/s41598-020-59077-1.

43. Kirkeby, C.; Ward, M.P. A Review of Estimated Transmission Parameters for the Spread
of Avian Influenza Viruses. *Transbound Emerg Dis* **2022**, *69*, 3238–3246,
doi:10.1111/tbed.14675.

44. Spackman, E.; Prosser, D.J.; Pantin-Jackwood, M.J.; Berlin, A.M.; Stephens, C.B. The
Pathogenesis of Clade 2.3.4.4 H5 Highly Pathogenic Avian Influenza Viruses in Ruddy
Duck (Oxyura Jamaicensis) and Lesser Scaup (Aythya Affinis). *J Wildl Dis* **2017**, *53*,
832–842, doi:10.7589/2017-01-003.

45. Spackman, E.; Prosser, D.J.; Pantin-Jackwood, M.; Stephens, C.B.; Berlin, A.M. Clade
2.3.4.4 H5 North American Highly Pathogenic Avian Influenza Viruses Infect, but Do Not
Cause Clinical Signs in, American Black Ducks (Anas Rubripes). *Avian Dis* **2019**, *63*,
366–370, doi:10.1637/11950-081418-ResNote.1.

46. Wille, M.; Robertson, G.J.; Whitney, H.; Bishop, M.A.; Runstadler, J.A.; Lang, A.S.
Extensive Geographic Mosaicism in Avian Influenza Viruses from Gulls in the Northern
Hemisphere. *PLoS One* **2011**, *6*, 1–10, doi:10.1371/journal.pone.0020664.

47. Hall, J.S.; TeSlaa, J.L.; Nashold, S.W.; Halpin, R.A.; Stockwell, T.; Wentworth, D.E.;
Dugan, V.; Ip, H.S. Evolution of a Reassortant North American Gull Influenza Virus
Lineage: Drift, Shift and Stability. *Virol J* **2013**, *10*, 1–9, doi:10.1186/1743-422X-10-179.

48. Lean, F.Z.X.; Falchieri, M.; Furman, N.; Tyler, G.; Robinson, C.; Holmes, P.; Reid, S.M.;
Banyard, A.C.; Brown, I.H.; Man, C.; et al. Highly Pathogenic Avian Influenza Virus
H5N1 Infection in Skua and Gulls in the United Kingdom, 2022. *Vet Pathol* **2023**,
doi:10.1177/03009858231217224.

49. Bodewes, R.; Kuiken, T. Changing Role of Wild Birds in the Epidemiology of Avian
Influenza A Viruses. In *Advances in Virus Research*; Kielian, M., Mettenleiter, T.,
Roossinck, M.J., Eds.; Academic Press, 2018; pp. 279–307 ISBN 9780128152010.

50. Fujimoto, Y.; Ogaswara, K.; Isoda, N.; Hatai, H.; Okuya, K.; Watanabe, Y.; Takada, A.;
Sakoda, Y.; Saito, K.; Ozawa, M. Experimental and Natural Infections of White-Tailed
Sea Eagles (Haliaeetus Albicilla) with High Pathogenicity Aviancinfluenza Virus of H5
Subtype. *Front Microbiol* **2022**, *13*.

51. Brown, J.D.; Stallknecht, D.E.; Swayne, D.E. Experimental Infections of Herring Gulls
(Larus Argentatus) with H5N1 Highly Pathogenic Avian Influenza Viruses by Intranasal
Inoculation of Virus and Ingestion of Virus-Infected Chicken Meat. *Avian Pathology*
2008, *37*, 393–397, doi:10.1080/03079450802216595.

52. Gonnerman, M.G.; Leyson, C.; Sullivan, J.D.; Pantin-Jackwood, M.J.; Mullinax, J.M.;
Prosser, D.J. Database Collating Previous Laboratory Investigations into the Pathogenesis

of Avian Influenza Viruses in Wild Avifauna of North America: U.S. Geological Survey Data Release. *https://doi.org/10.5066/P14MOUAF* 2024.

TABLES AND FIGURES

Figure 1. Bubble plot depicting the number of avian influenza virus challenge studies by viral
lineage. We present information A) for all HPAI and LPAI lineages. Data is grouped by host
species family and viral lineage and B) by clade of HPAI H5 viruses of the
Goose/Guangdong/1/1996 (Gs/GD) lineage. Viral lineages identified include Anhui (Anh),
Australian (Aus), Classical Eurasian (Cl.EA), Eurasian (EA), Gs/GD, North American poultry
(NA.P), North American wild birds (NA.WB), Pakistan H7 (Pak), and South American (SA).

A

*USDA Only (N = Individuals):
Accipitridae (N = 727)
Alcidae (N = 20)
Cathartidae (N = 691)
Charadriidae (N = 4)
Ciconiidae (N = 2)
Corvidae (N = 118)
Gaviidae (N = 9)
Gruidae (N = 10)
Hirundinidae (N = 1)
Icteridae (N = 7)
Pandionidae (N = 2)
Passerellidae (N = 1)
Pelecanidae (N = 119)
Phalacrocoracidae (N = 39)
Podicipedidae (N = 18)
Procellariidae (N = 4)
Rallidae (N = 2)
Stercorariidae (N = 2)
Strigidae (N = 248)
Sulidae (N = 3)
Tytonidae (N = 3)

*Challenge Only (N = Studies):
Columbidae (N = 8)

Figure 2. Comparison of the proportional composition of families identified in HPAI infections
reported to the USDA-APHIS database [20] (accessed Nov 16, 2023) and HPAI challenge
studies identified in our review. Families represented in only one dataset are listed with their
respective sample size. We present A) a list of families which were only identified in one of the
two datasets (i.e., "Single Dataset" in B) and B) a Sankey plot comparing proportional
composition of families identified in each dataset.

Figure 3. Disease response of seronegative host animals inoculated with >4 $\log_{10}$ EID_{50}/mL of a Gs/GD HPAI. Across studies, we summarize and present A) infection (blue) and mortality (red) rates, B) shedding rate in $\log_{10}$ EID_{50} or EIDeq, and C) duration of shedding in days for oropharyngeal (purple) and cloacal (orange) routes. Abbreviations by common name, repeated in order across panels: Wood Duck (WODU), Northern Pintail (NOPI), Mallard (MALL), American Black Duck (ABDU), Lesser Scaup (LESC), Redhead (REDH), Canada Goose (CAGO), Cackling Goose (CACG), Muscovy Duck (MUDU), Trumpeter Swan (TRUS), Mute Swan (MUSW), Surf Scoter (SUSC), Ruddy Duck (RUDU), Blue-winged Teal (BWTE), Herring Gull (HERG), Laughing Gull (LAGU), Ruddy Turnstone (RUTU), Dunlin (DUNL), Rock Pigeon (ROPI), American Kestrel (AMKE), Ringed-neck pheasant (RNEP), House Sparrow (HOSP), American Robin (AMRO), Black-crowned Night Heron (BCNH)