Title

A novel critic signal in identified midbrain dopaminergic neurons of mice training in operant

tasks

Abbreviated title

VTA DA neurons predict rewards during training

Authors

Jumpei Matsumoto[1], Virginie J. Oberto [2,3], Marco N. Pompili[2], Ralitsa Todorova[2], Francesco

Papaleo[4], Hisao Nishijo[1], Laurent Venance[2], Marie Vandecasteele[2] and Sidney I. Wiener[2,*]

[1]System Emotional Science, University of Toyama, 930-0194 Toyama, Japan

[2]Center for Interdisciplinary Research in Biology (CIRB), Collège de France, CNRS, INSERM,

Université PSL, 75005 Paris, France

[3]Neuro-Electronics Research Flanders, 3001 Leuven, Belgium

[4]Genetics of Cognition Laboratory, Neuroscience Area, Istituto Italiano di Tecnologia, 16163

Genova, Italy

* Corresponding author: sidney.wiener@college-de-france.fr

38 pages

4 Figures, 1 Supplementary table, 4 Supplementary figures

Abstract: 250 words; Introduction: 631 words; Discussion: 1480 words

Conflict of interest statement

The authors declare no competing financial interests.

Acknowledgements

Thanks to France Maloumian for invaluable help with figures, to Yves Dupraz for constructing

the experimental chamber, Estelle Anceaume for microscopy training, Michaël Zugaro for

helpful suggestions and informatics support, Dr. Guillaume Dugué and NeuroFabLab for facil-

itating with 3D printing of headstages, Drs. Karim Benchenane and Liyang Xiang for help

setting up the optical stimulation, and Dr. Philippe Faure for advice on recording dopaminergic

neurons with chronically implanted octrodes.

<u>Keywords</u>

ventral tegmental area; basal ganglia; learning; temporal difference learning; actor-critic

model

Abstract

In the canonical interpretation of dopaminergic neuron activity during Pavlovian conditioning, initially cell firing is triggered by unexpected rewards. Upon learning, activation instead follows the reward-predictive conditioned stimulus, and when expected rewards are withheld, firing is inhibited. However, little is known about dopaminergic neuron activity during the actual learning process in operant tasks. Here, we recorded optogenetically identified dopaminergic neurons of ventral tegmental area (VTA) in mice training in multiple, successive operant sensory discrimination tasks. A delay between nose-poke choices and trial outcome signals (for reward or punishment) probed for instructional or predictive activity. During training, but prior to criterion performance, firing increased after correct, but not incorrect choices, but prior to outcome signals. Thus, the neurons predicted whether choices would be rewarded, despite the animals' subthreshold behavioral performance. Surprisingly, these neurons also fired after reward delivery, as if the rewards had been unexpected according to the canonical view, but activity was inhibited after punishment signals, as if the reward had been expected after all. These inconsistencies suggest revision of theoretical formulations of dopaminergic neuronal activity to embody multiple roles in temporal difference learning and actor-critic models. Furthermore, on training trials when these neurons predicted that a given choice was correct and would be rewarded, surprisingly, the mice adhered to other non-rewarded and untrained task strategies (e.g., alternation). The DA neurons' reward prediction activity could serve as critic signals for the choices just made. This consistent with the notion that the brain must reconcile multiple Bayesian belief representations during learning.

Significance statement

The canonical view of dopaminergic function based on classical conditioning studies evokes reward-prediction error (RPE) signaling. Here, in mice performing a series of novel operant tasks with a delay between behavioral responses and reward/punishment signals, some neurons began firing after correct (but not incorrect) responses, but prior to the trial outcome (reward/punishment) signal. Nevertheless, the animals performed at chance levels, employing behavioral strategies other than the one signaled by the neurons. Furthermore, these same neurons showed canonical RPE responses, increased firing after reward signals (interpreted as the reward being unexpected) and firing rate decreases punishment

71   (interpreted as a reward having been expected). These findings indicate that dopaminergic

72   neurons can participate in diverse functions underlying different behavioral strategies for

73   learning.

Introduction

Adaptation for survival requires associating predictive cues with appropriate behavioral patterns to favor positive outcomes and to avoid aversive ones. Dopaminergic activity is implicated in the modification of neural circuits making such associations. Early in Pavlovian conditioning, dopaminergic neurons fire phasically after unexpected rewards. The DA neuron responses to rewards persist on trials immediately following learning acquisition, and then gradually diminish on subsequent trials (Hollerman & Schultz, 1998). With learning, responses appear after presentations of a conditioned stimulus (CS) that predicts the reward, and inhibitory responses occur after the expected reward is withheld.

These responses are also interpreted as evidence that dopaminergic neurons signal "reward prediction errors" (RPEs) (Houk et al., 1995; Montague et al., 1996, Schultz et al., 1997). RPE activities are posited to code "motivational value" (Schultz et al., 1997; Wise, 2005). RPE responses in dopaminergic neurons would correspond to a neural substrate for the "temporal difference (TD)" machine-learning algorithm which attributes eventual rewards or punishments to modify the circuitry leading up to these outcomes. TD learning estimates the expected reward on the basis of the current "state", that is, environmental context and cues, as well as the agent's actions. The expected reward is compared between the current state, and preceding estimates. Any differences, positive or negative, referred to as the "TD error", can then be employed to improve estimates of the state value. This "critic" dopaminergic signal would then modify activity of striatal "actors" for adaptive action selection (e.g., Khamassi, et al., 2005). These formulations originated in studies of Pavlovian conditioning, and their extension to learning adaptive behaviors, as in instrumental conditioning, still requires further elaboration regarding how appropriate actions are developed to optimize the balance of positive and aversive outcomes.

Recently, Cazettes et al. (2023) recorded in mice performing tasks with multiple possible response policies, and observed representations of unused strategies in M2 cortex, which projects to VTA (Watabe-Uchida et al., 2012). We hypothesized that if VTA neurons could then represent the rule currently being acquired during training, even while the animals are performing other strategies, since this could eventually serve as a "teacher" signal.

To test for such activity, we recorded mice as they trained in and acquired several versions of odor discrimination and visual discrimination tasks within the same experimental chamber. Trial outcome signals (signaling reward or punishment) were delayed after the

nosepoke responses. The mice were a mutant strain, permitting optogenetic identification of

dopaminergic neurons with light stimulation (Eshel et al., 2015).

As anticipated, we observed phasic dopaminergic neuron activity in the 0.5-1.0 s

interval after nose-poke choices but prior to trial outcome signals. Interestingly, a subset of

neurons discharged in this interval following correct, but not incorrect choices. Thus, these

responses signaled reward prediction (RP). Furthermore, this activity occurred during training,

while the animals executed a variety of other strategies prior to ultimately reaching criterion

performance. This activity could thus serve as a timely teacher signal. Surprisingly, most of

these same neurons also had typical excitatory responses to reward delivery signals (as found

prior to Pavlovian learning; Schultz, et al., 1997), and inhibition to punishment signals (as

found after Pavlovian learning; Schultz, et al., 1997). Thus, the neurons signaled that the

nosepoke choice was correct, and thus predicted reward, but also responded as if the rewards

were unexpected, according to the canonical interpretation of dopamine's role in the RPE/TD

learning framework. This may be reconciled in the perspective of instrumental learning, where

the response policy could be instructed and reinforced by a dopaminergic teacher signal prior

to trial outcome. These representations of alternative strategies could reflect multiple

Bayesian belief representations which must be reconciled prior to reaching criterion

performance levels. Functionally, dopamine release from reward prediction could prime the

network for reinforcement by the subsequent RPE reward-related dopamine release, or

punishment-related dopamine absence.

Materials and Methods

Animals

All experiments were performed in accordance with EU guidelines (Directive

86/609/EEC). Subjects were four transgenic adult (2-9 months) male mice expressing a

channelrhodopsin2- yellow fluorescent protein fusion protein (ChR2 (H134R)-eYFP) on

dopamine neurons. These mice were obtained by mating mice expressing the CRE

recombinase under the control of the dopamine transporter (DAT-IRES-CRE mice, Stock

006660, Jackson Laboratory, ME, USA) with mice bearing a CRE-dependent ChR2(H134R)-eYFP

gene (Ai32 mice, Stock 012569, Jackson Laboratory). Animals were housed in temperature-

controlled rooms with standard 12-hour light/dark cycles and food and water were available

ad libitum. Each workday, animals were handled to habituate to human contact, and weighed.

During pre-training and experimental periods, food intake, including pellets provided in the

experimental apparatus, was restricted to a maximum of 3 g/day. Water access remained ad

lib. Supplemental food was provided if weight fell below 85% of the normal weight.

Surgery

Anesthesia was induced with a mixture of ketamine (66 mg/kg) and xylazine (13 mg/kg)

and sustained with isoflurane (0.5 – 1.0%). Mice were placed in a stereotaxic device (David

Kopf Instruments) and maintained at 37° C. The scalp was exposed and cleaned with hydrogen

peroxide solution. Miniature jeweler's screws were screwed into trephines, and attached with

dental cement. A custom electrode assembly (developed by JM; see Oberto, Matsumoto, et

al., 2023) was implanted into the left VTA and SN. Briefly, the assembly consisted of

microdrives holding four recording probes for recording neural activity and two optic fibers

for optogenetic identification with light stimulation. Each recording probe was twisted

bundles of 8 formvar coated 12 micron diameter nichrome wires ("octrodes") were inserted

in polyimide tubes. The wires were gold-plated to an impedance of about 350 kohm. The

stainless-steel screws implanted on the left and right cerebellum as ground and reference

electrodes, respectively.

The behavioral task

Experimental apparatus

The 'Operon' system developed by Scheggia et al (2014) was adapted as an

experimental chamber permitting the mice to perform olfactory and visual discrimination

tasks (Figure 1). Many components were purchased from Med Associates (Fairfax, VT, USA).

Each of the short sides of the rectangular (160L x 136W x 160H mm) plexiglass chamber has

two arrays of LEDs and two ports used for both odor sampling and nose pokes. An

olfactometer presented d- and l- stereoisomers of limonene at the two odor ports in a pseudo-

random sequence. All six white LEDs were lit above one nose-poke port and off above the

other, again in pseudo-random sequence. Between the ports is a central feeder port with an

overhead lamp which is lit for 7 s during the delay after error trials. This lamp emits only 55 lx,

which is not intrinsically aversive. A reward pellet (5TUL TestDiet, Richmond, IN, USA, 14 mg,)

was delivered by a dispenser (ENV-203-14P, Med Associates, Fairfax, VT, USA). A plexiglass

barrier divided the two sides of the chamber and was slid up from below via an automated

motor assembly. Behavioral procedures followed those of the "Extradimensional shift"

protocol of Scheggia and Papaleo (2016) but without textural cues, and further details can be

found there and in Scheggia et al. (2014).

Figure 1. The behavioral apparatus and task (after Scheggia, et al., 2014).

A) Top) Overhead view. 1- Hexagonal LED array; 2- Reward dispenser; 3- Nose-poke port; 4- Moveable barrier; 5-House light; 6- Photobeam. Bottom) Frontal view of one chamber (photobeam not shown here). **B)** The automated task sequence.

Pre-training

Mice were provided water ad libitum, but mildly food deprived to maintain them above

85% their baseline weight, as controlled with daily weighing. The behavioral apparatus was

turned on, and the olfactometer outputs were confirmed to present odors discriminable by

the experimenter. Then, the mouse was placed into one compartment of the conditioning

chamber. The room lights were dimmed and the behavioral task was started. Animals were

first pre-trained to shuttle between two boxes for reward. In the first stage of training, a nose

poke into either port triggered a pellet delivery. Then animals were pre-trained for simple

visual discrimination task (light-on vs light off). While the rewarded side was varied pseudo-

randomly, in cases of spatial persistence, the other side was rewarded more. Once they could

regularly perform sequential trials in the maze, mice were provided food ad lib prior to

implantation with optrode assemblies, and training was resumed along with recordings.

Behavioral task sequence

After the pre-training, animals performed the following behavioral training on each

recording day. At the beginning of each trial, the barrier was lowered. Once the mouse crossed

a photodetector on its way to the other half of the chamber, one of the two LED arrays was lit

and the two odors were released from the respective odor ports on that side. This

photodetector crossing is called the "Cue" event. Once the mouse cleared these

photodetectors, the barrier was slid shut. Nose-pokes to the port on the side of the

(preselected) odor or visual cue crossed a photodetector within the port that triggered food

pellet release, or an error signal, at a variable latency ranging from 0.5 to 1.0 s. The position

of the rewarded cue was pseudo-randomly varied on successive trials so that the same odor

or visual cue were not presented more than twice at the left or the right nose-poke port.

Furthermore, the same trajectory orientation across the chamber (diagonally crossing or

running along a side wall) was not rewarded on more than 2 successive trials. The task control

system permitted programming the cue contingencies and reward delivery, outputting time

stamps. Erroneous choices triggered illumination of an overhead light during a timeout period

of 5 or 8 s while the barrier remained up. Once mice attained criterion performance (8

rewarded choices out of 10 consecutive trials, or 6 consecutively rewarded trials) in the

olfactory or visual discrimination tasks, the system automatically triggered a switch to the

other task. This is the same, or even more stringent, criterion than used in this type of study

(e.g., Kaefer, et al., 2020; Lapiz-Bluhm, et al., 2009, Tait, et al., 2017).

As criterion performance was successive attained, the mice were cycled between the

two tasks. In some sessions, it was necessary to pretrain the mice, starting with only the visual

or olfactory cue discrimination before adding the other modality cues. When the mouse

stopped performing trials for more than 120 s, the behavioral session of the day was ended.

The weight of the rewards earned was calculated, and the required amount of chow for the

daily feeding was calculated and provided.

Recording procedure

Recordings of neural signals were made with an Ampliplex system (sampling rate: 20

kHz; cutoff frequencies of the analog low-pass and high-pass filters were 0.3 and 10 kHz,

respectively). After surgical implantation, electrodes were advanced as often as twice daily

until neurons appeared with responses to optical stimulation. Electrodes were further

advanced when discriminable units were no longer present.

In each recording day, the headstage was plugged in and the mouse was placed in the

Operon system. Then, the neural signals were recorded while the mouse was performing the

above behavioral task. The position of the mouse within the Operon box was detected and

tracked using a LED attached on the headstage, which was captured with a video camera (30

frames/sec) installed above the system. After the mouse stopped performing the task, the

headstage plug was changed, optic fibers were connected, and the animal was placed in a

large plastic beaker. Using a 475 nm-laser light source (DPSSL BL473T3-100FL, Shanghai Laser

and Optics Century, Shanghai, PRC), trains of 10 light pulses (pulse duration: 10 msec; light

power: 10 mW max; frequency: 5 Hz) were given for 20 times with 10 sec inter train intervals

under control of an Arduino-based system.

Data analyses

Spike sorting and neuron type classification

For single unit discrimination from the extracelluar signals recorded from the octrodes,

offline spike sorting was carried out with KiloSort (Pachitariu et al., 2016) followed by manual

curation using Klusters (L. Hazan, http://neurosuite.sourceforge.net).

243   To identify neurons as dopaminergic, we used the Stimulus-Associated spike Latency

244 Test (SALT; Kvitsiani et al., 2013; Eshel et al., 2015). The test determines whether light pulses

245 significantly changed a neuron's spike timing by comparing the distribution of first spike

246 latencies relative to the light pulse, assessed in a 10-msec window after light-stimulation, to

247 10-msec epochs in the baseline period (-150 to 0 msec from the onset of light-stimulation; see

248 Kvitsiani et al., 2013 for details). A significance level of $p<0.01$ was selected for this.

249   All neurons recorded from an octrode with at least one SALT+ response were

250 considered to be in a dopaminergic nucleus. If the octrode was not advanced, neurons on the

251 day before and the day after were also counted as in VTA or SNc, even if no SALT+ responses

252 were recorded on those days. Similarly, if the octrode had not been advanced, and SALT+

253 responses had been recorded on non-consecutive days, intervening days with no SALT+

254 responses were still considered as in a dopaminergic nucleus. Also, when electrodes had been

255 advanced, all recordings in days between those with SALT+ recordings were also considered

256 to be in VTA/SNc. Neurons with SALT+ responses are considered dopaminergic. Neurons in a

257 dopaminergic nucleus not demonstrating a significant SALT+ response are labelled SALT-,

258 although this negative result does not conclusively show that the neuron is not dopaminergic.

259 All neurons qualifying as in dopaminergic nuclei were categorized with clustering according to

260 the following parameters: spike width, mean firing rate. The criterion for fast spiking

261 interneurons (FSI) was firing rate > 15 Hz and spike width < 1.5 ms (cf., Ungless and Grace,

262 2012). No FSI's were SALT+. All other SALT- neurons in dopaminergic nuclei are referred to as

263 "Other".

265 *Behavioral correlates of neural activity*

266   The cells with mean firing rates <1 spike/s in a given epoch were excluded from

267 analyses. We defined that "Cue" as when the mouse first crossed the photobeam while

268 entering the next chamber, triggering cue onset; "Choice" is the time of the nose-poke

269 response, as detected by the photo-detectors in the ports; "Outcome" is either the instant the

270 dispenser released the reward pellet (which made a salient sound), or the onset of the

271 punishment period. For each neuron in each epoch, firing rate during baseline periods (0.5 s

272 before the cue), [cue, choice] periods, [choice, outcome signal] periods, and post-outcome

273 periods [0.5 s after the outcome signal] were calculated. The RP responsive neurons were

274 defined as a neuron that showed significant difference(s) between rewarded and non-

rewarded trials in either or both of [cue, choice] and [choice, outcome signal] periods

(unpaired t-test, p < 0.05). Similarly, the responses to the reward and punishment were tested

by comparing firing rate in the baseline period with the outcome periods of rewarded and

non-rewarded trials, respectively (unpaired t-test, p < 0.05).

Experimental design and statistical analyses

Subjects were four transgenic adult (2-9 months) male mice expressing a

channelrhodopsin2- yellow fluorescent protein fusion protein (ChR2 (H134R)-eYFP) on

dopamine neurons. Data samples were recordings of neurons from individual periods when

the mice were challenged with one of several sensory discrimination tasks. (The result show

that a single neuron could show different response properties in the different tasks.) In order

to respect "Reduction" of the 3R's, experiments were stopped when sample sizes were

decided to be sufficient, i.e., when significant effects could be determined reliably and SEM

error bars were minor.

Basic statistical methods such as paired and unpaired t-test, chi-square test, and

Pearson's correlation analysis were used for statistical comparisons. The statistical tests were

performed using MATLAB. The significance threshold was set to p<0.05. The values for n, p,

and the specific statistical test performed for each analysis are included in the corresponding

figure legend, table, or main text.

Histology

Once stable recordings were no longer possible, marking lesions were made with 10 s

of 30 μA cathodal current. After waiting for at least 90 min, mice were then killed with a lethal

intraperitoneal injection of sodium pentobarbital, and perfused intra-ventricularly with

phosphate buffered saline solution (PBS), followed by 10% phosphate buffered formalin. The

brain was removed, post-fixed overnight, and placed in phosphate buffered 30% sucrose

solution for 2-3 days. Frozen sections were cut at 80 μm, and permeabilized in 0.2% Triton in

PBS for 1 h at room temperature. Antibody staining was performed to confirm localization of

electrodes in dopaminergic nuclei. Sections were then treated with 3% bovine serum albumen

(BSA) and 0.2% Triton in PBS for 1 h with gentle agitation at room temperature to block non-

specific binding. Sections were then rinsed for 5 min in PBS at room temperature with gentle

agitation. Then the sections were left overnight with gentle agitation at 4° C in a solution of

the first antibody, mouse monoclonal anti-TH MAB318 (1:500), 0.067% Triton, and 1% BSA in

PBS. After three rinses for five minutes in PBS, the sections were treated with a second

antibody (1/200, anti-mouse, green), Nissl-red (1:250), and 0.067% Triton in PBS for two hours

at room temperature. Then sections were rinsed 3 times for five minutes in PBS, and mounted

with Fluoromount®. Sections were examined with fluorescence microscopy to verify electrode

tips and trajectories in the immunofluorescent stained zones (see Supp. Fig. 3D of Oberto,

Matsumoto, et al., 2023). Anatomical reconstruction with fluorescence microscopy revealed

RP neuron recording within the ventral tegmental area at mediolateral positions ranging from

375 to 517 μm.

For further details, see Oberto, Matsumoto, et al. (2023).

Results

The behavioral task

Mice with optogenetically responsive dopaminergic neurons were first trained in a visual discrimination task in a double chamber apparatus adapted from Scheggia, et al. (2014) (Figure 1). At the beginning of each trial, a central barrier was lowered. Once the mouse crossed a photodetector to enter the other chamber, a cue was presented (LED lights on at the left or the right nose-poke ports in a pseudo-random sequence) for the visual discrimination task. Nose-poke choices were followed by a delay at variable latencies ranging from 0.5 to 1.0 s prior to either the reward signal (sound of a food pellet dispenser), or, alternatively, a punishment signal (for 5 or 7 seconds, chamber lights on coupled with a delay for barrier removal permitting the next trial). Once criterion performance was achieved in the visual discrimination task, mice were then implanted with multiple electrodes and equipped for intracranial optogenetic stimulation.

The mice were then recorded as they re-acquired the visual discrimination task. When criterion performance was again reached, an olfactory discrimination task was introduced in the same maze, with two different odors emanating from the nose-poke ports, again in pseudo-random sequence. In practice, when a mouse achieved criterion performance (6 successive rewarded trials, or 8 out of 10), the task was changed (see Annex of Khamassi, et al., 2024, for justification of these criteria). If only one cue modality had been presented, the other was then added without changing the reward contingency. This challenged the animal to continue performing the task in the presence of a second inconsequential cue. If criterion performance was reached while the two cue modalities were presented simultaneously, the reward contingency was shifted to the other sensory modality, and both cue types were still presented. Should the mouse reach criterion again, the reward contingency was again shifted to the other cue modality discrimination task (visual to olfactory, or olfactory to visual). Each period with one or both cue types or with different reward contingencies is referred to below as a "task epoch", and these periods are analyzed separately since a given neuron could respond differently in different task epochs (detailed below). The reward contingency switching corresponds to the extra-dimensional attentional set-shifting task of Scheggia, et al (2014) which would require intact prefrontal cortical function (Dias, et al., 1996; Birrell & Brown, 2000; Bissonette, et al., 2008), which, in turn, depends upon dopaminergic input (see e.g., Vander Weele, et al., 2019; Di Domenico & Mapelli, 2023).

Behavior

Of the 53 task epochs when one or more reward-predictive optogenetically identified DA neurons were recorded, in 28, the mice ultimately achieved criterion performance in the task. In 18 others, the mice achieved criterion performance in another epoch in the session. Thus, learning was indeed taking place. The overall performance in these sessions was 56±1%, mean ± SEM; Supplementary Figure 1), consistent with reward contingencies being changed once criterion was reached (and thus this percentage is low since only a small fraction of trials were recorded during criterion performance).

Supplementary Figure 1. Distribution of mean performance levels (proportion of trials that were rewarded) in epochs with reward-predictive responses in identified dopaminergic neurons. N = 53 epochs. If more than one cell was recorded in a given epoch, it was only counted once.

Interestingly, as the animals were acquiring the tasks, their choices were not random. During epochs with RP responses, animals performed at criterion levels (i.e., at least 6 successive adherent trials) in strategies other than the current rule (i.e., go left, go right, spatial alternation, or choosing a currently unrewarded odor or lit/unlit nose-poke port; see Khamassi, et al., 2024, for justification of this selection of strategies analyzed). Of the 24 task epochs with RP responses and sufficient numbers of trials to analyze, all but one had runs of trials that were performed according to strategies other than the current rule (see Supplementary Figure 2).

Supplementary Figure 2. Incidences of the lengths of 247 runs of successive trials adherent to strategies other than the current rewarded rule for a sample of 24 epochs when DA reward predictive activity was recorded. (If more than one RP responding neuron was recorded within an epoch, the runs were only counted once.) These tallies included 1723 trials (See Materials and Methods for further details). Only epochs with 15 or more trials were tallied (not counting those trials qualifying for criterion performance at the currently rewarded task. For the strict "100% adherence" approach, we counted occurrences of at least successive six trials adhering to the strategy. For the more permissive "90%" adherence" approach, intermittent single non-adherent trials were allowed, as long as they were contained within a sequence of at least eight adherent trials. In cases where overlapping series adhering to two different strategies, shared trials were assigned to the longest sequence (or assigned randomly when sequences were of equal length).

Neuronal recordings

In 156 recording sessions in four mice, of total 967 neurons recorded under different task conditions, 272 were optogenetically identified as dopaminergic (DA) neurons, and 206 were fast-spiking interneurons (FSI), while 489 others could not be identified as DA or FSI. The latter are referred to as "OTHER" neurons (see Materials and Methods for criteria for category assignments). While mice were *performing at sub-criterion levels* in the sensory discrimination tasks, in the 272 task epochs with optogenetically identified DA neurons, 82 (30%) of these fired selectively according to whether the trials would be rewarded or not, but *prior to the signals indicating whether the choice would be rewarded or punished* (Figure 2). These are referred to as "reward prediction" (RP) cells. This selectivity also occurred in 106 OTHER neurons recorded in 489 epochs (21%; Supplementary Figure 3) and 31 FSI in 206 epochs (15%). The incidence of RP responses was greater in DA neurons than FSI ($\chi^2(1) = 14.8$, $p = 1.2 \times 10^{-4}$) or OTHER neurons ($\chi^2(1) = 6.74$, $p = 9.4 \times 10^{-3}$). The results reported below focus on the identified DA neurons, because of their theoretical interest. Most neurons significantly increased firing when predicting rewards than punishments (85% for DA, 55% for OTHER, and 68% for FSI; Supplementary Table 1) and only these are illustrated in the Figures and further

 characterized below. In the RP cells, when the target (i.e., currently rewarded) cue was visual,

79% of significant RP responses occurred in the [cue onset, nose-poke] interval but only 31%

in the [nosepoke, outcome] period in DA neurons (Figure 2, left). In OTHER neurons, 50% and

56% of significant RP responses occurred in the [cue onset, nose-poke] and the [nosepoke,

outcome] periods, respectively; Supplementary Figure 3, left). In contrast, when target cues

were odors, only 12% of DA (and 10% of OTHER) RP responses occurred in the [cue onset,

nose-poke] while 95% (and 93% of OTHER) were in the [nosepoke, outcome] period (Figure 2,

right and Supplementary Figure 3 right). In no case did the activity clearly ramp up until trial

outcome signals were presented (Figure 2 and Supplementary Figure 3), in contrast with the

ramping responses previously shown in dopaminergic neurons prior to rewards (e.g., Farrell

et al, 2022), and which have been assimilated to a "motivational incentive" signal associated

with reward seeking (Berridge and Robinson, 1998).

Supplementary Table 1. Relative incidence of RP responses in DA neurons by epoch.

DA neurons

	Visual target			Odor target			All targets		
	simple	complex	total	simple	complex	total	simple	complex	total
R>NR	16	13	29	12	29	41	28	42	70
R<NR	1	3	4	2	6	8	3	9	12
Both	0	0	0	0	0	0	0	0	0
n.s.	44	72	116	17	57	74	61	129	190
Total	61	88	149	31	92	123	92	180	272
% R>NR	26%	15%	19%	39%	32%	33%	30%	23%	26%

OTHER neurons

	Visual target			Odor target			All targets		
	simple	complex	total	simple	complex	total	simple	complex	total
R>NR	10	6	16	15	27	42	25	33	58
R<NR	11	9	20	6	19	25	17	28	45
Both	1	0	1	2	0	2	3	0	3
n.s.	71	137	208	38	137	175	109	274	383
Total	93	152	245	61	183	244	154	335	489
% R>NR	11%	4%	7%	25%	15%	17%	16%	10%	12%

R>NR (or R<NR) signifies that the average firing rate during rewarded trials was significantly higher (or lower) than during the punished (not rewarded) trials in either the [cue onset, nose-poke] or [nose-poke, outcome] period. Both means that the neuron exhibited opposite RP responses (R>NR and R<NR) in the two different periods ([cue onset, nose-poke] and [nose-poke, outcome]). n.s. – not significant.

DA and OTHER neurons with RP activity also had canonical responses to reward and

punishment signals. Firing rates significantly increased after reward signals in 54/70 (77%) of

DA and 40/58 (77%) of OTHER neuron epochs (unpaired t-test, $p < 0.05$) while firing rates

decreased significantly after punishment signals in 47/70 (67%) of DA and 20/58 (34%) of

OTHER neuron epochs (unpaired t-test, $p < 0.05$; columns of stars to the right of color rasters

in Figure 2 and Supplementary Figure 3). Note that when this firing rate reduction occurred, it

only lasted at most for 1 s after punishment onset, even though the punishment period lasted

5 to 7 s.

Figure 2. RP responses in DA neurons.
A, top two rows of panels) Color rasters showing z-scored firing rates in 0.5 s windows (framed by continuous vertical lines) around the three principal task events (dashed vertical lines). Each row represents firing of a neuron during a single task contingency epoch. The z-scores are calculated as bin value/[mean of all 60 bins for the cell in that epoch/SD for the 60 bins].

Right columns of stars) Canonical dopaminergic responses. In the 0.5 s periods after reward or punishment (outcome) signals, red stars indicate significant increases in firing rate relative to the 0.5 s baseline period prior to cue onset, while blue stars indicate significant firing rate decreases (unpaired two-tailed t-test, $p<0.05$). **A, bottom)** Reward prediction quantified as differences between values for rewarded and punished trials. Right) Left columns are for the [cue onset, nosepoke choice] period and right columns are for the [nosepoke choice, outcome signal] period. Here red stars indicate significant differences between firing in rewarded and punished trials (unpaired two-tailed t-test, $p<0.05$). Staircase plots are ordered based on the latency of the peak firing rate between cue and outcome in the rewarded trials in the displayed data. Note that for the statistical analyses of [cue, choice] and [choice, outcome signal] periods, all data from these periods were used, which are not necessarily represented in the figures visualizing the neural activity in 1 s windows around the events since these intervals could last longer than one second. **B)** Means of the data of the upper two rows of panels of A. Darker lines indicate mean and shaded areas show SEM. **C)** Means of differences in the third row of panels of A. N = 29 (Light) and 41 (Odor) epochs.

Supplementary Figure 3. OTHER neurons with RP responses.
Same format as Figure 2. N = 16 (Light) and 42 (Odor) epochs.

Figures 3 and 4 show examples of these RP responses in DA neurons. In addition to

their prominent RP responses, firing rates significantly increased (relative to baseline) after

reward onset (p<0.05, unpaired t-test), the canonical response previously shown in Pavlovian

conditioning prior to acquisition (Schultz, et al., 1997). The cells of Figures 3B, 4A and 4B also

had another canonical DA neuronal response, inhibition after the punishment onset (p<0.05,

unpaired t-test). In the literature, this inhibition has typically been observed after the animals

acquire Pavlovian conditioning, in those trials when an expected reward is withheld (Schultz,

et al., 1997). Thus, in the RPE framework, this response would indicate that a reward was

expected. However, the significantly lower firing rate prior to the punishment signals on these

trials (relative to activity in the same period in trials to be rewarded) would indicate that no

reward was expected.

In the sessions of Figures 4A and B, the mouse reached criterion twice, and, accordingly,

the task was changed each time. In Figure 4A, the RP response was significant in all three of

the task epochs, while the performance level averaged only 51%. But in the neuron of Figure

4B, the RP response was only significant in the third epoch (while performance was 52%). Thus,

in this paper, the incidence of RP responses is tallied by task epochs rather than over whole

sessions (which could dilute significant effects).

The numbers of successive trials in runs with adherence to non-rewarded strategies is

shown in Supplementary Figure 2. The legends of Figures 3A, 3B, 4A and 4B describe the

numerous runs of trials adhering to unrewarded strategies in the RP epochs, despite the RP

responses reflecting the correct strategy. This, in conjunction with the RP neuronal responses,

would be consistent with multiple functional networks underlying the respective strategies

operating in parallel, guiding the behavior of the animals.

Figure 3. Examples of RP responses in sessions with performance levels of 53% (A) and 46% (B).

In both A and B, the RP response is significant in the [nose-poke, outcome signal] period, but the [cue onset, nose-poke] RP activity is only significant in A (p<0.05, unpaired t-test). Also,

the canonical activity increase after post-reward signals is significant in both, but activity inhibition after the post-punishment signal is significant only in B ($p < 0.05$, unpaired t-test). Green circles indicate the times of nose-poke choices in left columns while at the right blue circles indicate timing of reward signals and red x's indicate punishment signals. Horizontal dashed lines in histograms indicate background firing rate prior to trial onset. These mice did not reach criterion performance in these sessions, and thus there is only one task epoch. For A, the target cue was an odor, but the mouse chose the lit port with >87% adherence on trials 3-11, and 23-45. For B, the target cue was an odor, and the mouse chose the unlit port on trials 3-12, the port emitting the non-target odor 4 on trials 18-24, alternated left and right ports on trials 42-50 (Alternation), then went to the lit port on trials 53-63 and 67-75, and to the left port on trials 78-89, again all with >87% adherence.

Figure 4. Examples of RP activity during the [nose-poke, outcome signal] period in two cells in different sessions.

(Same format as Figure 3.) Panels A and B show the response of two different cells. Rasters are synchronized with outcome signals while histograms in columns 2 and 4 are synchronized

with nose-poke responses (green circles in rasters). Bottom) Mean histograms from the entire
session. Dashed lines indicate background firing rates. **A)** In epochs 1 and 3 the target cue was
an odor, and no visual cues were presented. In epoch 2, the odor was also the target cue, but
with visual cues present. RP activity was significant in all three epochs ($p<0.05$, unpaired t-
test). Inhibition was significant after punishment signals in epochs 1 and 2 only ($p<0.05$,
unpaired t-test), but none had significant post-reward signal excitation ($p>0.05$, unpaired t-
test). Performance levels were 46%, 51% and 47% in the respective epochs. Unrewarded
strategies were: trials 20-29, 93-98 and 114-131 Left port; trials 29-35, Odor 4; trials 47-56,
Right port; trials 57-67 and 83-92, Unlit port; trials 72-77, Alternation. **B)** Reward predictive
responses significant only in epoch 3. ($p<0.05$, unpaired t-test. Same format as A.) The
inhibition after the punishment signal is significant in epoch 3 ($p<0.05$, unpaired t-test), but
the firing rate was not significant higher after the reward signal ($p>0.05$, unpaired t-test). In
epoch 3, unrewarded strategies were: trials 64-72 and 87-95, Unlit port; trials 75-85, 105-115
125-130, 133-150, and 153-159, Left port; Epoch 1- Visual target cue with no olfactory cues
present. Epoch 2-Visual target cue with olfactory cues present. Epoch 3- Olfactory target cue
with visual cues present.

We distinguished between two levels of difficulty in the sensory discrimination tasks.

When only the target cue modality was presented, the epoch was considered "simple". When

both cue modalities were presented, and thus only one had to be attended to for the

discrimination task, while the other had to be ignored, the epoch was considered "complex".

Overall, proportions of RP responses occurred in epochs with simple and complex

discriminations were not significantly different (34% vs 28%, $\chi^2(1) = 0.83$, $p = 0.36$;

Supplementary Table 1). There was a greater incidence of RP responses in epochs with

olfactory target cues than visual target cues (40% vs 22%; $\chi^2(1) = 10.0$, $p = 1.6 \times 10^{-3}$; see

Supplementary Table 1).

Concerning anatomical localization, most recordings, and RP responses were made in

the VTA (see Supplementary Figure 3D of Oberto, Matsumoto, et al., 2023) with only one

epoch for a dopaminergic neuron in SNc (which was more difficult for electrode placement

and the sample there is smaller) and eight epochs for OTHER SNc neurons.

Possible motor factors

We examined motor parameters in the [cue onset, nose-poke] period on rewarded and

punished trials as a potential indicator of impulsivity/deliberateness of the choices. There was

no significant difference for cue-to-nosepoke delay, distance traveled, or initial velocity for

rewarded vs. punished trials (Supplementary Figure 4). In the period after nose-poke

responses, the initial velocity was significantly greater on trials prior to punishment signals

was greater than prior to reward trials (Supplementary Figure 4). This raised the possibility

that the differential firing on rewarded vs. punished trials could be related to velocity

dependence of firing rate (Puryear et al, 2010). However, Pearson's correlation analyses of

firing rate and velocity were significant in only five of the 74 neurons (p<0.05, r = -0.43 to -

0.23).

Supplementary figure 4. Control comparison of motor parameters in rewarded (R) and non-rewarded (NR) trials in epochs with RP activity.

556 Each value was z-scored based on the distribution relative to the values for the respective

557 epoch. Gray dots represent individual epoch values. Black dots and error bars indicate means

558 and SEMs. p-values are from paired t-tests. N = 74 epochs with RP responses in identified DA

559 neurons. Of 82 RP DA neurons, 8 neurons were excluded from the following analysis due to

560 technical issues in animal tracking (e.g., videos were not recorded properly).

Discussion

In summary, in mice training in sensory discrimination tasks in an automated double

chamber, dopaminergic neurons fired phasically following correct nose-poke responses, but

prior to signals independently of whether the trial was to be rewarded or not. These reward-

prediction responses occurred while behavioral performance was at chance and sub-criterion

levels. Thus, the neurons accurately predicted the imminent reward, even though the animal's

low performance level did not reflect this information. Furthermore, following a delay after

the nose-poke responses, trial outcome signals evoked contradictory responses within the

"reward prediction error" (RPE) framework of interpreting dopaminergic neuronal activity

during Pavlovian conditioning (Schultz, 2016). There, excitatory activity after reward signals

are interpreted as the reward being unexpected. However, punishment signals could evoke

inhibitory responses in the same epoch, an indicator that an *expected* reward had been

withheld. These diverse responses in dopaminergic neurons could reflect inputs from distinct

brain systems that represent conflicting "belief states" (of expectation of reward; see

Gershman & Uchida, 2019; Rao, 2010), as expressed in a Bayesian theoretical framework.

Here, individual neurons displayed responses associated with rewards being both expected

(reward prediction and post-punishment inhibition) and unexpected (post-reward firing rate

increase) on successive trials. Crucially, the reward predictive firing occurred while the animals'

behavior was below criterion performance, and they were still learning the stimulus-response-

reward contingencies. Thus, these neurons signaled whether the choice was correct or not,

even though this information was not exploited in the animal's performance levels. But these

signals could have participated in the learning process. The presence (after correct choices)

and absence (on incorrect choices) of RP activity could be considered as a critic signal as

postulated in reinforcement learning algorithms. The earliest appearance of reward prediction

responses occurred primarily after cue presentation when the target was visual, but after

nose-pokes for odor targets. The early responses for visual targets demonstrate that the

reward prediction activity could precede the choice response, and potentially serve as an

instruction signal.

While there are several studies of dopaminergic activity during instrumental learning

and decision-making (Bayer & Glimcher, 2005; Morris, et al., 2006; Nishino, et al., 1987;

Phillips, et al., 2003; Roesch, et al., 2007; Roitman, et al., 2004; Stuber, et al., 2005; Jones et

al., 2010), in these experiments, CRs and trial outcome signals were concurrent, and phasic

responses to the CR could not be disambiguated from those to the reward signal.

The use of the term CR here is expanded from common usage. Firstly, the behavioral

responses occurred during training, prior to when the responses had been learned. Further,

the CRs being learned are not simply the motor acts of making a nose-poke, but rather making

an appropriate (left or right) choice based upon the previous history of rewarded task

contingencies. Since the tasks required shifting attention between one or more multi-modal

cues target cues in different spatial locations, there was an important contextual component

to choices. Thus, the dopaminergic activity in these VTA areas that receive input from the

hippocampal-accumbens (limbic) pathway, rather than the SNc regions whose input is from

more sensorimotor cortico-striatal pathways (Haber, 2014).

Comparison with published responses in operant tasks

Studies of dopamine release in the striatum and in recordings of midbrain

dopaminergic neurons have typically focused on animals undergoing classical conditioning or

instrumental conditioning with instructed choices. Studies of instrumental conditioning with

free choices have provided intriguing results pointing towards models other than actor/critic

TD learning. Similar to the present results, Syed et al (2016) found that in an instrumental

learning task, dopamine release in striatum was greater after initiation of correct than

incorrect choice responses, but prior to the reward outcome signal. When the animals made

the incorrect choice, the movement dynamics were slower than on rewarded trials. However,

after correct choices, the DA release dynamics were fundamentally different from the present

phasic responses: the dopamine levels gradually built up over the course of several seconds

to peak after the reward outcome signal. These dynamics were also observed in recordings of

VTA neurons under conditions of high uncertainty (Fiorillo, et al., 2003). This type of response

is generally assimilated with motivation incentive or drive for reward acquisition (Nishino, et

al., 1987; Howe, et al., 2013), rather than with short-latency phasic responses related to

motivation value for learning associations (Schultz et al., 1997; Wise, 2005). For example, in a

task requiring self-initiated sequences of bar presses, dopamine release preceded and

continued during these movements (Wassum et al., 2012). Again, the time course was over

several seconds and the responses were interpreted as representing incentive motivation. In

a study of non-human primates performing a two-arm bandit task, Morris et al. (2006)

observed that population activity of DA neurons corresponds to choice behaviors. However, the latter held only for instructed choice trials, not for free choice decision trials, as was the case here. They concluded that dopamine neurons receive information about upcoming actions, and argue for including Q-learning into algorithms describing this activity. A challenge to a role for RPE in learning comes from Coddington and Dudman's (2018) study of trace conditioning in mice. There, the canonical RPE correlates were found in VTA and SNc dopaminergic neurons, but these only appeared after learning. Also, Lak et al. (2016) studied monkeys learning the values of probabilistic rewards. Even though behavior indicated that acquisition occurred within a few trials, DA activity corresponding to learning (a late response appearing 0.4-0.65 s after the cue) only developed later, after the 4th trial.

Timing in relation to previously reported DA responses

The phasic reward prediction responses here are on the same time scale as found in Pavlovian conditioning, where responses to rewards or to reward-associated cues last for up to 200 ms (Schultz, 1997; Hollerman and Schultz, 1998). Lak et al (2016) distinguished a rapid time scale, at 0.1-0.2 s after cue presentation, but these corresponded to novelty responses which disappeared with familiarity. Later activity, 0.4-0.6 s after cue presentation reflected learned reward value, but appeared only after performance levels had improved. Thus, this does not correspond to the activity found here. Another response type at longer time scales is characterized by sustained gradually ramping from cue presentation until reward delivery. This is in contrast with the phasic RP activity here that immediately followed the behavioral response. For example, the RP responses here bear only superficial resemblance Goedhoop et al's (2023) recordings of DA release in the nucleus accumbens (which receives VTA inputs) in animals performing similar Pavlovian and operant conditioning tasks. Ramping of DA release occurred only in their operant task, starting with the onset of the reward-predictive cue, and leading up to the operant response. This was interpreted as anticipation of a rewarded action.

Reward prediction, and mismatch with behavior

Engelhard, et al (2019) recorded DA neurons in mice performing a visual discrimination task on a virtual T-maze. During the cue presentation, but prior to choice, some neurons selectively increased their discharge rate for trials where the mice made incorrect choices. Roesch et al (2007) found cue-triggered DA unit responses indicated the most valuable reward

choice available, even on trials when the animal subsequently did not select this. After cues
were turned off, but prior to movement for reward, the activity shifted to represent the actual
choice. Thus, this is not similar to the observations here. Furthermore, cue selectivity
developed with learning, while the responses here persisted over many trials, sometimes in
sessions where learning did not occur. However, note that Drieu, et al (2025) found in mouse
auditory cortex neurons a late component of responses that predicted rewards in an auditory
go/no-go discrimination task. These responses occurred during learning, prior to when the
animals achieved criterion performance, like those shown here.

Our results pose a conundrum: the neurons "know" the correct response, but this is
not reflected in the animal's behavior. One possible explanation would be that the DA signal
acts gradually upon the network until a critical amount of circuit changes are implemented to
finally modify behavior. Here only a fraction of recorded neurons predicted reward in these
sessions while the animals had not yet reached performance criterion. Perhaps further work
will reveal mechanisms by which RP activity could lead to rule acquisition.

During training, behavioral choices were not random. Rather, animals performed other
strategies while DA neurons predicted rewards in the visual and olfactory discrimination tasks.
This could reflect the presence of multiple Bayesian belief representations which must be
reconciled prior to reaching criterion performance levels. Different subsets of neurons or
neuronal circuit dynamics could underlie performance of the respective strategies. Such
representations were reported in mouse secondary motor (M2) cortex by Cazettes, et al.
(2023) where neural ensembles simultaneously encoded multiple strategies in foraging tasks.
Note that M2 projects to VTA in mice (Watabe-Uchida, et al., 2012). Functionally, dopamine
release during reward prediction could prime the network for reinforcement by the
subsequent reward prediction error reward-related dopamine release, or punishment-related
dopamine absence.

References

1. Bayer HM, Glimcher PW (2005) Midbrain dopamine neurons encode a quantitative
reward prediction error signal. Neuron 47(1):129-41. doi: 10.1016/j.neuron.2005.05.020.

2.   Berridge KC, Robinson TE. (1998) What is the role of dopamine in reward: Hedonic impact,

reward learning, or incentive salience? Brain Res Brain Res Rev. 28(3):309-69. doi:

10.1016/s0165-0173(98)00019-8.

3.   Birrell JM, Brown VJ (2000) Medial frontal cortex mediates perceptual attentional set

shifting in the rat. J Neurosci 20:4320–4324

4.   Bissonette GB, Martins GJ, Franz TM, Harper ES, Schoenbaum G, Powell EM (2008) Doub

le dissociation of the effects of medial and orbital prefrontal cortical lesions on attentional

and affective shifts in mice. J Neurosci 28:11124–11130.

5.   Cazettes F, Mazzucato L, Murakami M, Morais JP, Augusto E, Renart A, Mainen ZF (2023)

A reservoir of foraging decision variables in the mouse brain. Nat Neurosci 26, 840–849.

doi:10.1038/s41593-023-01305-8

6.   Coddington LT, Dudman JT (2018) The timing of action determines reward prediction

signals in identified midbrain dopamine neurons. Nat Neurosci. 21(11):1563-1573. doi:

10.1038/s41593-018-0245-7.

7.   Di Domenico D, Mapelli L. (2023) Dopaminergic modulation of prefrontal cortex inhibition.

Biomedicines. 11(5):1276. doi: 10.3390/biomedicines11051276.

8.   Dias R, Robbins TW, Roberts AC (1996) Dissociation in prefrontal cortex of affective and

attentional shifts. Nature 380:69–72

9.   Drieu C, Zhu Z, Wang Z, Fuller K, Wang A, Elnozahy S, Kuchibhotla K (2025) Rapid

emergence of latent knowledge in the sensory cortex drives learning. Nature.   Mar 19.

doi: 10.1038/s41586-025-08730-8. Epub ahead of print.

10. Engelhard B, Finkelstein J, Cox J, Fleming W, Jang HJ, Ornelas S, Koay SA, Thiberge SY, Daw

ND, Tank DW, Witten IB (2019) Specialized coding of sensory, motor and cognitive

variables in VTA dopamine neurons. Nature 570:509–513.

11. Eshel N, Bukwich M, Rao V, Hemmelder V, Tian J, Uchida N. (2015) Arithmetic and local

circuitry underlying dopamine prediction errors. Nature. 525(7568):243-6. doi:

10.1038/nature14855.

12. Farrell K, Lak A, Saleem AB (2022) Midbrain dopamine neurons signal phasic and ramping

reward prediction error during goal-directed navigation. Cell Rep. 41(2):111470. doi:

10.1016/j.celrep.2022.111470.

13. Fiorillo CD, Tobler PN, Schultz W (2003) Discrete coding of reward probability and

uncertainty by dopamine neurons. Science. 299(5614):1898-902. doi:

10.1126/science.1077349

14. Gershman SJ, Uchida N. (2019). Believing in dopamine. Nat. Rev. Neurosci. 20, 703–714.

doi.org/10.1038/s41583-019-0220-7

15. Goedhoop J, Arbab T, Willuhn I. (2023) Anticipation of appetitive operant action induces

sustained dopamine release in the nucleus accumbens. J Neurosci. 43(21):3922-3932. doi:

10.1523/JNEUROSCI.1527-22.2023.

16. Haber SN. (2014) The place of dopamine in the cortico-basal ganglia circuit. Neuroscience.

282:248-57. doi: 10.1016/j.neuroscience.2014.10.008.

17. Hollerman JR, Schultz W. (1998) Dopamine neurons report an error in the temporal

prediction of reward during learning. Nat Neurosci. 1(4):304-9. doi: 10.1038/1124.

18. Houk JC, Adams JL, Barto AG. (1995) A model of how the basal ganglia generate and use

neural signals that predict reinforcement. In: Models of Information Processing in the

Basal Ganglia, eds. J. C. Houk, J. L. Davis, and D. G. Beiser. Cambridge, MA: MIT Press, p.

249–270.

19. Howe MW, Tierney PL, Sandberg SG, Phillips PE, Graybiel AM. (2013) Prolonged dopamine

signalling in striatum signals proximity and value of distant rewards. Nature.

29;500(7464):575-9. doi: 10.1038/nature12475.

20. Jones JL, Day JJ, Aragona BJ, Wheeler RA, Wightman RM, Carelli RM (2010) Basolateral

amygdala modulates terminal dopamine release in the nucleus accumbens and

conditioned   responding.   Biol.   Psychiat.   67(8):737-44.   doi:

10.1016/j.biopsych.2009.11.006.

21. Kaefer K, Nardin M, Blahna K, Csicsvari J. (2020) Replay of behavioral sequences in the

medial prefrontal cortex during rule switching. Neuron 106(1):154-165.e6. doi:

10.1016/j.neuron.2020.01.015

22. Khamassi, M., Lachéze, L., Girard, B., Berthoz, A., & Guillot, A. (2005). Actor-critic models

of reinforcement learning in the basal ganglia: From natural to artificial rats. *Adaptive*

Behavior, 13(2), 131–148. doi:10.1177/105971230501300205

23. Khamassi M, Peyrache A, Benchenane K, Hopkins DA, Lebas N, Douchamps V, Droulez J,

Battaglia FP, Wiener SI. (2024) Rat anterior cingulate neurons responsive to rule or

strategy changes are modulated by the hippocampal theta rhythm and sharp-wave ripples.

Eur J Neurosci. doi: 10.1111/ejn.16496

24. Kvitsiani D, Ranade S, Hangya B, Taniguchi H, Huang JZ, Kepecs A. (2013) Distinct

behavioural and network correlates of two interneuron types in prefrontal cortex. Nature.

498(7454):363-6. doi: 10.1038/nature12176.

25. Lak A, Stauffer WR, Schultz W. (2016) Dopamine neurons learn relative chosen value from

probabilistic rewards. Elife. 5:e18044. doi: 10.7554/eLife.18044.

26. Lapiz-Bluhm MD, Soto-Piña AE, Hensler JG, Morilak DA. (2009) Chronic intermittent cold

stress and serotonin depletion induce deficits of reversal learning in an attentional set-

756 shifting test in rats. Psychopharmacol (Berl). 202(1-3):329-41. doi: 10.1007/s00213-008-

757 1224-6.

758 27. Montague PR, Dayan P, Sejnowski TJ. (1996) A framework for mesencephalic dopamine

759 systems based on predictive Hebbian learning. J Neurosci. 16(5):1936-47. doi:

760 10.1523/JNEUROSCI.16-05-01936.1996.

761 28. Morris G, Nevet A, Arkadir D, Vaadia E, Bergman H. (2006) Midbrain dopamine neurons

762 encode decisions for future action. Nat Neurosci. 9(8):1057-63. doi: 10.1038/nn1743.

763 29. Nishino H, Ono T, Muramoto K, Fukuda M, Sasaki K (1987) Neuronal activity in the ventral

764 tegmental area (VTA) during motivated bar press feeding in the monkey. Brain Res.

765 413(2):302-13. doi: 10.1016/0006-8993(87)91021-3.

766 30. Oberto VJ, Matsumoto J, Pompili MN, Todorova R, Papaleo F, Nishijo H, Venance L,

767 Vandecasteele M, Wiener SI. (2023) Rhythmic oscillations in the midbrain dopaminergic

768 nuclei in mice. Front Cell Neurosci. 17:1131313. doi: 10.3389/fncel.2023.1131313.

769 31. Phillips PE, Stuber GD, Heien ML, Wightman RM, Carelli RM. (2003) Subsecond dopamine

770 release promotes cocaine seeking. Nature 422(6932):614-8. doi: 10.1038/nature01476.

771 32. Pachitariu, M., Steinmetz, N., Kadir, S., Carandini, M., and Harris, K. (2016). Kilosort:

772 Realtime spike-sorting for extracellular electrophysiology with hundreds of channels.

773 bioRxiv [Preprint] doi: 10.1101/061481

774 33. Puryear CB, Kim MJ, Mizumori SJ. (2010) Conjunctive encoding of movement and reward

775 by ventral tegmental area neurons in the freely navigating rodent. Behav Neurosci.

776 124(2):234-47. doi: 10.1037/a0018865.

777 34. Rao RPN. (2010) Decision making under uncertainty: a neural model based on partially

778 observable Markov decision processes. Front Comput Neurosci. 4, 146.

35. Roesch MR, Calu DJ, Schoenbaum G. (2007) Dopamine neurons encode the better option

in rats deciding between differently delayed or sized rewards. Nat Neurosci. 10(12):1615-

24. doi: 10.1038/nn2013.

36. Roitman MF, Stuber GD, Phillips PE, Wightman RM, Carelli RM (2004) Dopamine operates

as a subsecond modulator of food seeking. J Neurosci. 24(6):1265-71. doi:

10.1523/JNEUROSCI.3823-03.2004.

37. Scheggia D, Bebensee A, Weinberger DR, Papaleo F. (2014) The ultimate intra-/extra-

dimensional attentional set-shifting task for mice. Biol Psychiat. 75:660–670. doi:

10.1016/j.biopsych.2013.05.021

38. Scheggia D, Papaleo F. (2016) An operant intra-/extra-dimensional set-shift task for mice.

J Vis Exp. 107:e53503. doi: 10.3791/53503.

39. Schultz W. (2016) Dopamine reward prediction-error signalling: a two-component

response. Nat Rev Neurosci. 2016 17(3):183-95. doi: 10.1038/nrn.2015.26.

40. Schultz W, Dayan P, Montague PR. (1997) A neural substrate of prediction and reward.

Science. 275(5306):1593-9. doi: 10.1126/science.275.5306.1593.

41. Stuber GD, Roitman MF, Phillips PE, Carelli RM, Wightman RM. (2005) Rapid dopamine

signaling in the nucleus accumbens during contingent and noncontingent cocaine

administration. Neuropsychopharmacology. 30(5):853-63. doi: 10.1038/sj.npp.1300619.

42. Syed EC, Grima LL, Magill PJ, Bogacz R, Brown P, Walton ME. (2016) Action initiation

shapes mesolimbic dopamine encoding of future rewards. Nat Neurosci. 19(1):34-6. doi:

10.1038/nn.4187.

43. Tait DS, Phillips JM, Blackwell AD, Brown VJ. (2017) Effects of lesions of the subthalamic

nucleus/zona incerta area and dorsomedial striatum on attentional set-shifting in the rat.

Neurosci. 345:287-296. doi: 10.1016/j.neuroscience.2016.08.008.

44. Ungless MA, Grace AA. (2012) Are you or aren't you? Challenges associated with

physiologically identifying dopamine neurons. Trends Neurosci. 35(7):422-30. doi:

10.1016/j.tins.2012.02.003.

45. Vander Weele CM, Siciliano CA, Tye KM. (2019) Dopamine tunes prefrontal outputs to

orchestrate   aversive   processing.   Brain   Res.   1713:16-31.   doi:

10.1016/j.brainres.2018.11.044.

46. Wassum KM, Ostlund SB, Maidment NT. (2012) Phasic mesolimbic dopamine signaling

precedes and predicts performance of a self-initiated action sequence task. Biol

Psychiatry. 71(10):846-54. doi: 10.1016/j.biopsych.2011.12.019.

47. Watabe-Uchida M, Zhu L, Ogawa SK, Vamanrao A, Uchida N. (2012) Whole-brain mapping

of direct inputs to midbrain dopamine neurons. Neuron. 74(5):858-73. doi:

10.1016/j.neuron.2012.03.017.

48. Wise RA. (2005) Forebrain substrates of reward and motivation. J Comp Neurol.

493(1):115-21. doi: 10.1002/cne.20689.

Author contributions

JM and SIW designed and performed the experiments and obtained funding support. MNP performed experiments. JM designed the carrier for optic fibers and octrode driver assemblies, constructed the experimental apparatus, the optrodes and optogenetic stimulation apparatus, informatics control, and data acquisition systems with support from HN. MV and LV maintained and genotyped the mouse line and guided immunohistochemical processing. FP guided adaptation of the maze and behavioral protocols for training and recording. JM, VJO, and SIW analyzed the data with support from RT and HN, and wrote the manuscript. All authors approved of the manuscript.

Funding

Grants from Uehara Memorial Foundation (to JM), the Takeda Science Foundation (to JM and HN), the Labex Memolife, Fondation pour la Recherche Médical, Fondation Bettencourt Schueller, and International Research Laboratory DALoops (to SW) provided support.

Ethics

All procedures were in accord with local (autorisation d'experimenter no. 75-1328-R; Comite d'Ethique pour l'Experimentation Animale no. 59, dossier 2012-0007) and international (European Directive 2010/63/EU; US National Institutes of Health guidelines) standards and legal regulations regarding the use and care of animals.