Species interactions and food-web context drive temperature-dependent prey evolution

Ze-Yi Han[1], Yaning Yuan[1], Katrina DeWitt[1], Andrea Yammine[1], Daniel J. Wieczynski[1], Masayuki Onishi[1], Jean P. Gibert[1]

[1]Department of Biology, Duke University, Durham, NC, USA

To whom correspondence should be addressed: Ze-Yi Han, Department of Biology, Duke University, Durham, NC, USA, 402-405-3888, zeyihanw@gmail.com

Keywords:
Thermal evolution, Food-web eco-evolutionary dynamics, Food-web temperature response, Trait-base ecology, plasticity and evolutionary response

Data and code availability: Data and code will be archived upon publication.

Acknowledgments: This work was supported by a U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomic Science Program Grant under Award Number DE-SC0020362 to JPG.

ABSTRACT

Understanding how global warming shapes species evolution within communities is a pressing

goal of ecology. Temperature affects interacting species and can lead to changes in species

interactions, but how that will alter species evolutionary trajectories within complex food webs is

poorly understood. Here we address 1) whether different predators affect prey evolution

differentially, 2) whether the food web context in which this happens influences prey evolution,

3) whether temperature affects prey evolution directly, and 4) whether ecological interactions

mediate how temperature affects prey evolution. We use a combination of mathematical

modeling and experimental evolution assays in microbial food webs composed of prey algae and

their protists predators. We found that temperature alone doesn't drive prey evolution unless

predators are involved. Importantly, the influence of temperature through predation is contingent

on the food web structure. This leads to distinct evolutionary trajectories when prey evolves with

predators alone or with a competing predator present. Our findings indicate that the species

evolution to warming is likely contingent on their specific ecological contexts, suggesting that

similar species across different food webs could exhibit diverse evolutionary responses to new

climates.

INTRODUCTION

Global warming is reshaping natural communities worldwide (Parmesan 2006) as rising
temperatures affect population growth (Dell *et al.* 2011; Frazier *et al.* 2006; Kontopoulos *et al.*
2020; Savage *et al.* 2004; Wieczynski *et al.* 2021) and species interactions (Blois *et al.* 2013;
Dell *et al.* 2014; Gilman *et al.* 2010; Suttle *et al.* 2007; Tylianakis *et al.* 2008). Temperature, in
particular, increases metabolic costs (Clarke 2006; Clarke & Fraser 2004), leading predators to
consume more prey (Sheridan & Bickford 2011) while reaping ever decreasing energetic benefits
(Barneche *et al.* 2021). This decreasing energetic intake results in the energetic choking of upper
food-web trophic levels, subsequent trophic collapse (Ullah *et al.* 2018; Voigt *et al.* 2003;
Zarnetske *et al.* 2012), and food web rewiring (Barbour & Gibert 2021; Bartley *et al.* 2019). This
has grave implications, contributing to large changes in community composition and
structure(Komatsu *et al.* 2019) as well as the marked loss of top predators throughout the
Anthropocene (Estes *et al.* 2011; Voigt *et al.* 2003).

Although theory predicts that rapid prey evolution and predator-prey co-evolution may mitigate
these temperature impacts, empirical evidence of rapid adaptation under global warming in a
food web context remains scarce (De Meester *et al.* 2019; Tseng & O'Connor 2015). On one
hand, temperature can directly influence evolution through effects on organismal metabolism
(Alton *et al.* 2024; Clarke 2003; Gillooly 2000; Terblanche *et al.* 2009), morphology (Brans *et*
al. 2017; Diamond *et al.* 2017; Yampolsky *et al.* 2014), and fitness (Diamond *et al.* 2018; Norin
& Metcalfe 2019; Padfield *et al.* 2016). On the other hand, predation within complex food webs
also drives species evolution (Abrams 2000; De Meester *et al.* 2019; Frickel *et al.* 2017; Hairston
et al. 2005; Palkovacs & Post 2009) and often selects for more defensive but less competitive

prey (Yoshida *et al.* 2004). The link between abiotic drivers of selection, like temperature, and

biotic drivers of selection, like predation, is established through the often asymmetric

temperature effects on key demographic and ecological predator traits such as growth or attack

rates (Gibert *et al.* 2022; Robertson & Hammill 2021; Wieczynski *et al.* 2021). This suggests that

prey evolutionary responses are likely jointly influenced by temperature, as well as predation and

competition among predators –i.e., the broader food web context the interactions are embedded

in. But whether or how this happens in nature is poorly understood.

To disentangle this potentially complex interplay between biotic and abiotic drivers of species

evolution and understand its implications within broader food webs under climate change, we

focus on a model system of global distribution and relevance (Bar-On *et al.* 2018; Herron *et al.*

2019; Wieczynski *et al.* 2021): the interaction between the unicellular green algae

Chlamydomonas reinhardtii and its ciliate predators. We assembled tractable microcosm food

webs of organisms that co-occur across ecosystems (from soils to wetlands, Foissner *et al.* 2009),

and studied the joint effects of temperature, predation, and predator competition (as a proxy for

food web context), on prey evolution. We tracked population dynamics, phenotypic changes, and

rapid shifts in prey genetic makeup (i.e., rapid evolution), to understand the eco-evolutionary

dynamics that ensue within these microbial food webs across temperatures and the processes that

determine them.

Specifically, we address the following questions: 1) do different predators affect prey evolution

differentially? 2) Does competition between predators (i.e., food web context) influence prey

evolution, and, how? 3) Does temperature affect prey evolution directly? And, 4) Do ecological

interactions mediate how temperature affects prey evolution? We show that temperature does not

directly affect prey evolution in the absence of predators, but does so in their presence through

direct temperature effects on the ecological interactions. Moreover, this mediation effect of

temperature is determined by the food web context, such that evolution under predation with and

without a predator competitor leads to different evolutionary outcomes. Our results emphasize

that organismal evolutionary responses to a warming world likely are strongly dependent on the

biotic context in which they are embedded, so that similar species in different food webs might

show different eco-evolutionary responses to novel climates.

RESULTS

Predation maintains genetic diversity

To understand how predation influences prey evolution, we established a tractable model of

rapid evolution under predation that keeps track of two genetically distinct prey strains that share

a common predator (see Methods). The model suggests that two different prey strains cannot

coexist in the absence of predators –classically, unless inter-strain competition is lower than

intra-strain competition– (Appendix I Equation 1-2). Predation, however, facilitates the invasion

of the strain that would otherwise be lost (Appendix I Equations 1,3), implying that predation

can maintain genetic diversity. We tested these theoretical predictions using an experimental

predator-prey microbial system where the alga *Chlamydomonas reinhardtii* is preyed upon by

one of three possible ciliate protist species (*Glaucoma sp.*, *Tetrahymena pyriformis*, *Paramecium*

caudatum, see Methods). The algal population is composed of two genetically and

phenotypically distinct, algal prey strains – fluorescently tagged wild type (*wt*) and untagged

vfl1-1 (i.e., 'variable flagellar number1-1')– as our prey population. We kept track of their

abundances, phenotypes, and shifts in clonal frequencies (using their fluorescent spectrum) over

time as means to quantify evolution by clonal sorting (see Methods). Our empirical results

supported our model predictions: while *wt* consistently outcompeted *vfl1-1* in the absence of

predators, predation facilitated the persistence of the inferior competitor in all predation

treatments, regardless of the presence of the predator competitor (Fig 2a-d; Appendix II Figure

1-2). Interestingly, *C. reinhardtii* grew faster in all predation treatments, perhaps due to

increased fertilization by predator waste (Cech *et al.* 2008; McNaughton *et al.* 1997). We

hypothesized that predation would select for *wt* and *vfl1-1* differently (Fig 2a-d; Appendix IV

Table 1-2). *vfl1-1* had the highest frequency under predation from *Glaucoma sp.*, but lowest

frequency under predation from *P. caudatum* (Fig 2d; Appendix IV Table 2).

Prey traits determine eco-evolutionary outcomes

Our model suggested that variations in ciliate predator attack rates are sufficient to explain the

different evolutionary outcomes across treatments (Fig 2a-b, model parameter see Appendix I

Table 1), thus offering a simple explanation for an otherwise seemingly complex dynamic. In our

experimental system, predation selects on two key functional traits of the prey: defensive

clumping and motility. Strain *vfl1-1* has impaired swimming ability compared to *wt*, due to

pleiotropic effects of mutations on *VFL1* gene, (Adams *et al.* 1985; Silflow *et al.* 2001)

Appendix IV videos) and is therefore predicted to experience lower predation rates relative to the

more motile *wt*, all else being equal (Aljetlawi *et al.* 2004; González *et al.* 1993; Pawar *et al.*

2012; Visser 2007). Our data supports this prediction as *vfl1-1* is selected against in the absence

of predators (Fig 2 c-d), and selected for in the presence of most predators, with frequencies

higher relative to control, likely owing to decreased predation through impaired motility relative

to *wt*. However, in addition to increased mobility, *wt* can form large defensive clumps (Lurling &

Beekman 2006), while *vfl1-1* shows impaired clumping ability under predation in lab. Through

cytometry, we measured the particle size of the algal population as a proxy for the onset of

clumping, with larger average particle size and maximum particle size indicating clumping (see

Methods). Although, *vfl1-1* showed very small size changes in response to predation (Fig 2e,

Appendix IV Table 3), *wt* showed strong plastic clumping defense against predation with an

increase in both average and maximum particle size (Fig 2e green; Appendix II Fig 3-4;

Appendix IV Table 4). The distribution of *wt* particle size additionally vary by predators (Fig 2e,

Appendix IV Table 4), suggesting species-specific predation rates, in support of our model

prediction. Finally, the relative fitness of motility and clumping traits is predator-dependent:

Glaucoma sp. selected for *vfl1-1* while *P. caudatum* drove *vfl1-1* to extinction (Fig 2c-d;

Appendix IV Table 2).

Temperature directly affects ecological dynamics and prey plasticity, but indirectly affects prey

evolution

Although temperature influenced the growth rate of *wt* and *vfl1-1* (Appendix II Fig 6),

temperature alone did not affect prey genetic frequencies, and hence, prey evolution (Fig 3a

Control; Appendix II Figure 2). Only in the presence of predation did temperature significantly

affect prey evolution, suggesting that temperature effects on prey evolution are mediated by

predation. This likely happened through a combination of differential predator thermal

performance and prey plastic responses (Fig 3; Appendix IV Table 2,5). Indeed, temperature

significantly affected the demographic parameters that underpin the ecological dynamics of each

predator: the intrinsic growth rate, *r*, and their maximum abundance (a measure of their carrying

capacity, K; Fig 3c, Appendix II Fig 7, Appendix IV Table 6-7). These temperature-dependence

of key predator demographic parameters likely determined predation intensity, leading to

significant differences in the mean, median, variance, skewness and kurtosis of *wt* particle size

(Fig 3b; Appendix IV Table 8). As temperature indirectly affects *wt* plastic response through the

predator-prey interaction, predator strain-specific attack rate on *wt* and *vfl1-1* will also likely

change over time depending on predator preference.

Interestingly, the fitness of *vfl1-1* and *wt* under predation by *P. caudatum* shifted dynamically

over time, and these shifts were most pronounced the higher temperature treatment (Fig 3c).

Indeed, *vfl1-1* frequency increased with *P. caudatum* density by day 5 compare to control, but

dropped to near extinction by day 15, and this happened more rapidly at higher temperatures (Fig

3c). This indicates that the low motility of *vfl1-1* was favored at first through differential

predation (rapid fast evolutionary response, Fig 3c, left) while the plastic clumping defense of *wt*

was favored later on (Fig 3c, right). In other words, intermittent rapid evolutionary and plastic

responses ultimately determined the eco-evolutionary response of the prey. Additionally, the

boom-bust change in *vfl1-1* frequency, with initial increase that followed by a steep drop, at

higher temperatures (Fig 3c) likely indicates that the faster increase in *P. caudatum* density at

higher temperature (Fig 3c)   led to a faster onset of defensive the clumping response by *wt* (Fig

3b). Therefore, direct temperature effects on predator ecological dynamics and prey plastic

responses sped up changes in strain relative fitness that resulted in changes in genetic

frequencies and prey evolution (Fig 3c). Altogether, our results reveal the mechanisms by which

temperature indirectly affect prey evolution through its direct and differential effects on predator

ecological dynamics and prey plastic responses, which ultimately shape prey functional traits and

evol-evolutionary dynamics.

Temperature-mediated predator competition (i.e., food web context) affects prey evolution

The presence of a second predator independently and interactively affected prey evolution and

prey plastic clumping with temperature, predator species, and time (Fig 4; Appendix IV Table 9-

12). Specifically, we observed a combined effect from both predators on prey evolution in

competition treatments (Fig 4a; Appendix III Fig 1). Indeed, when we analyzed the evolutionary

dynamics of the *C. reinharditt* (i.e. genetic frequencies of *vfl1-1* over time) using Earth Mover's

Distance approach (see Methods), we found that the similarity of *vfl1-1* genetic frequencies over

time between competition treatments and single predator treatments (for example, between

Glaucoma sp + P. caudatum and *P. caudatum* or *Glaucoma sp*) were higher than the similarity

between those single predator treatments (for example, between *Glaucoma sp.* and *P. caudatum*;

Appendix III Fig 1).

Meanwhile, temperature also indirectly affected the relative contribution of each predator on

prey evolution within predator competition treatments (Fig 4a) through its effect on predator

performance. *P. caudatum* had increased initial population growth rate, *r*, and maximum

population density in higher temperature while those of *T. pyriformis* and *Glaucoma sp.*

decreased in higher temperature (Fig 4b; Append II Fig 7), and *vfl1-1* genetic frequencies over

time in competition treatments that include *P. caudatum* resemble most to those in which *P.*

caudatum is the only predator at high temperature (lowest EMD; Appendix III Fig 1).

Additionally, the size distribution of *wt* in competition treatments also showed combined effects

of both predators and temperature mediated the predator-specific effects on *wt* plastic responses

(Fig 4c; Appendix III Fig 2; Appendix IV Table 11). Altogether, our result show how

temperature mediates the effect of food web contexton these eco-evolutionary dynamics, through

differences in predator thermal performance, predator competition, and prey plasticity.

DISCUSSION

Our results show that temperature strongly but indirectly affects prey evolution in the presence of

predation through its direct effects on predator population dynamics and species interactions.

Predator-specific selection on prey traits and predator thermal performance together determine

prey plastic responses, which in return led to changes in fitness of prey functional traits overtime

as ecological dynamics unfold (Fig 2-3). Moreover, multiple predators can concurrently respond

to temperature and jointly affect the prey phenotypic responses that underpin prey evolution (Fig

4), thus highlighting the importance of the food web context in indirectly shaping species

evolution under global warming.

Prey motility, defensive clumping, and predator identity, jointly determine prey rapid evolution

Our study shows that predator differential selection on multiple prey functional traits (i.e.

motility and clumping) allow the less dominant strain to persist (Siepielski *et al.* 2020), thus

maintaining higher additive genetic diversity (Fig 2). More so, we revealed the reciprocal nature

of the effects of species functional traits and species interactions, and how these underpin the

dynamics of species evolution within food webs (Fig 3).While previous studies showed that

more motile prey experience higher predation rates than slow or immotile prey (Andersen &

Dölger 2019; Brodin & Johansson 2004; González *et al.* 1993; Visser 2007), the relative fitness

of low motility of *vfl1-1* under predation is ultimately controlled by the onset of *wt* plastic

clumping and the predator response to *wt* clumping defense. In other words, dynamic plasticity

in one strain alters the course of the evolutionary process within food webs by influencing

relative fitness across strains.

Predator identity affects prey plastic responses

Chlamydomonas reinhardtii is known to develop clumping in response to abiotic and biotic

stressors (Fischer *et al.* 2016; Herron *et al.* 2019). However, our study showed predator

preferential selection led to a novel predator-specificity in plastic clumping distribution of *C.*

reinhardtii (Fig 3). Moreover, multiple predators can have coordinated effects in shaping prey

plastic traits distribution in predictable ways (Fig 4; Appendix III Fig 2). Prey evolutionary

dynamics under selection by multiple predators reflected the combined effects of selection

imposed by individual predator species (Fig 4; Appendix III Fig 1). These results emphasize the

interesting but unexplored consequences of species phenotypic plasticity: shifts in one functional

trait in response to one predator may affect the interaction with another predator, which could

cascade down through the entire complex community, as previously suggested theoretically

(Cosmo *et al.* 2023; Guimarães *et al.* 2017). Our study thus highlights the importance of a

mechanistic understanding of the interplay between species functional trait dynamics and biotic

interactions across trophic levels in complex communities, not just pairwise interactions (Henn *et*

al. 2018).

Food web context and temperature interactively drive prey evolution

Our study shows the underlying mechanisms through which reciprocal effects between food web

context and trait dynamics together drive prey evolutionary trajectories across temperatures (Fig

3-4). We observed that asymmetric warming effects on species at different trophic levels

influence predator-prey interactions (Gibert *et al.* 2022; Kordas *et al.* 2011; Lindmark *et al.*

2019), which had been previously suggested to be an important aspect of food web dynamical

responses to rising temperatures (Gibert *et al.* 2022; Parain *et al.* 2016; Thakur *et al.* 2018;

Vasseur & McCann 2005). We also show that different predator thermal responses mediate

competition among predators at the same trophic level, allowing the more dominant predators to

shape prey traits and evolutionary dynamics more strongly (Fig 4; Appendix III Fig 1-2).

Temperature-mediated changes in species interactions drive changes in species plastic and

evolutionary functional traits (Barbour & Gibert 2021; Fischer *et al.* 2016; Fordyce 2006), which

in turn can reshape food web dynamics, structure, and function (Bartley *et al.* 2019; DeLong *et*

al. 2023; Gibert 2019; Han *et al.* 2023; Lister & Garcia 2018; Petchey *et al.* 1999; Thakur *et al.*

2018). As warming increases extinction rates among predators (Thunell *et al.* 2021; Voigt *et al.*

2003; Zarnetske *et al.* 2012), a decrease in predator diversity might also lead to decreases in prey

trait diversity that may result in disproportionate effects from more dominant predators and

increased loss of genetic diversity within prey populations. Our results have a major implication:

species evolution in novel climates will be shaped by the food web context within which they are

embedded, such that both the composition of the food web, but also their network complexity,

likely will determine the evolutionary outcome, and hence the fate of these species (Barbour &

Gibert 2021).

Concluding remarks

Overall, we showed that biotic interactions mediate temperature effects on species rapid thermal

adaptation. We showed that predator-specific selection can change over time depending on the

onset of plastic responses in other traits, which determines the evolutionary outcome.

Additionally, predation triggers prey thermal adaptation, and this is in turn co-determined by

predator thermal performance, prey plasticity, and temperature effects on competition among

predators (food web context). Our results emphasize that, by directly affecting biotic interactions

and species plastic traits, temperature can play an indirect –but key– role, in determining species

evolution in a warming globe.

METHODS

Experimental work

Unicellular microalgae can be found in diverse habitats, from soil to freshwater ecosystems

across the world (Arora & Sahoo 2015; Falkowski 1994). These organisms are at the base of all

food webs, fueling both green and brown food webs (Brito *et al.* 2006; Descroix *et al.* 2010; Guo

et al. 2016), and are routinely preyed upon by microbes (e.g. ciliate protists) and metazoans (e.g.,

rotifers, cladocerans) alike (Calatrava *et al.* 2023). Our focal prey species, the unicellular green

algae *Chlamydomonas reinhardtii*, is a well-established model organism with known mutations

linked to functional traits (Sasso *et al.* 2018; Calatrava *et al.* 2023). Here we used two genetically

and phenotypically distinct strains of *C. reinhardtii*: wild type (*wt*) and *vfl1-1* (i.e., 'variable

flagellar number1-1') as our prey population, and kept track of their abundances, phenotypes,

and shifts in genetic frequencies over time as means to quantify evolution by clonal sorting.

Wild type (*wt*) possesses two flagella that allow the entire range of normal locomotive behaviors

(Huang 1986) and can form large cell clumps, a common form of defense against predation

(Herron *et al.* 2019; Lurling & Beekman 2006). Strain *vfl1-1* produces individuals with variable

numbers of flagella from 0 to 10 and defective swimming ability (Adams *et al.* 1985; Silflow *et*

al. 2001). Additionally, *vfl1-1* seems to be unable to clump under predation. To distinguish these

two strains and keep track of their relative frequencies in mixture populations, we used

mNeonGreen fluorescent protein tagged *wt* populations and used flow cytometry to count tagged

wt vs non-tagged *vfl1-1* individuals over time (Fig 1a). Both strains can autfluoresce (Fig 1a) and

be detected by the cytometer, but only one (*wt*) will show fluorescence in the near green spectrum

(Fig 1a). We used two protist predators of similar body size, *Tetrahymena pyriformis* and

Glaucoma sp., as the focal predators, and a larger protist *Paramecium caudatum*, as a competitor

for the predators (Fig 1b). All four species are commonly found in freshwater and soil systems

(Cornwallis *et al.* 2023; Foissner & Berger 1996).

To understand how temperature and ecological interactions (predation and competition among

predators, as a proxy for food web context) jointly influence *C. reinhardtii* evolution, we set up

experimental microcosms in autoclaved 250ml borosilicate jars filled with 100ml of 9:1

COMBO media:timothy hay infusion and one wheat seed as a carbon source. Each microcosm

was assigned to one of three possible predation treatments (No predation/Control, + *T.*

pyriformis, + *Glaucoma sp.*), one of two possible predator competition treatments (No

competition/Control, + *P. caudatum*), and one of three possible temperatures (19°C, 22°C, and

25°C). The manipulations produced a factorial design with 18 combinations of treatments (Fig

1b), each replicated six times, yielding a total of 108 microcosms.

Prior to experimentation, the algae strains were maintained in TAP agar (Rajam & Kumar 2007)

at room temperature. Protist cultures were maintained in bacterized timothy hay protist media at

22°C and a 16:8-hour light-dark cycle. All cultures were transferred to 9:1 COMBO media:

timothy hay protists media (Brans *et al.* 2022) and cultured under the same light and temperature

regime as protist stock cultures 2 weeks prior to experimental work. We carried out the

experiment in two blocks on two consecutive days, each block had half of the replicates in all

treatments. We started *wt* and *vfl1-1* strains at equal densities of 2000 individuals/ml in all

microcosms and initialized *T. pyriformis*, *Glaucoma sp.*, and *P. caudatum* populations at density

of 5 ind/ml, 5 ind/ml, and 0.5 ind/ml, respectively. The experiment was carried out for 15 days,

or ~ 30-45 *C. reinhardtii* generations.

We recorded all species densities on days 5, 10, and 15. We used flow cytometry (NovoCyte

2000R, Agilent, CA, USA) to distinguish mNeonGreen tagged *wt* individuals from untagged

vfl1-1 individuals. This allowed us to track the abundance and frequencies of both strains over

time. We used forward scatter height (FSC-H) as a measure of cell/cell clump size of *C.*

reinhardtii (Adan *et al.* 2017) to track plastic morphological change in response to protist

predation, thus providing a window into both rapid plastic change (i.e., within strains), and rapid

evolutionary change (i.e., change in genetic frequencies; Fig 1c). We recorded the density of all

protist species through fluid imaging (FlowCam; Yokogawa Fluid Imaging Technologies,

Portland, ME, USA) at a magnification of 10x (Fig 1c).

We hypothesized that 1) different predators select for different traits of the prey, 2) competition

between predators might weaken directional selection on prey, and 3) temperature affects species

performance differently and may lead to changes in the strength of predator-prey interactions,

which can mediate prey evolution.

Data analysis

To analyze how temperature and ecological interactions influenced prey evolution, we used a

combination of autoregressive moving average linear mixed models (ARMA-LMMs; 'nlme'

package, v. 3.1-162) and classic linear mixed models (LMM) using the 'lme4'package (version

1.1-3 in R v. 4.3.1). We calculated the relative frequencies of *vfl1-1* in total *C. reinhardtii*

population as the measure of prey evolution. To better evaluate changes in prey evolution

affected by temperature, we also calculated the changes in *vfl1-1* frequencies across temperatures

(Δ *vfl1-1* frequency) by deducting the mean *vfl1-1* frequencies at 19°C from each replicate at

22°C and 25°C in each treatment.

We first performed LMM on the single predator treatments and the control (Fig 1b) to

understand the individual effects of each predator on prey evolution. We analyzed the fixed

effects of predator species identity, temperature, and time on the prey evolution and temperature

effects on prey evolution, by using *vfl1-1* frequency and Δ *vfl1-1* frequency as response variables

respectively. We then added in the densities of the single predators as a fixed effect to the

previous model to analyze the effects of predation pressure of prey evolution. We then

performed LMM on *T. pyriformis* and *Glaucoma sp.* single predator treatments and the

competition treatments (Fig 1b) to test whether the presence of a second predator had joint

effects with predator species identity, predator density, and temperature, on *vfl1-1* genetic

frequency and Δ *vfl1-1* frequency using previous linear mixed models.

Additionally, to quantify the relative importance of each predator on prey evolution in

competition treatments, we calculated Earth Mover's Distance (EMD from now on) with

'emdist' (R package, version 0.3-3), which quantifies the similarity of prey evolution patterns

over time between different predation treatments (higher EMD means less similarity).

Specifically, within each temperature, we calculated the EMD of *vfl1-1* frequencies 1) between

the single predator treatments in each competition pair and 2) between each of the single

predator treatments and the corresponding competition treatment. Similarly, to quantify the

relative effects of each predator on prey plasticity, we also calculated the EMD of *wt* particle size

distribution between: 1) the single predator treatments in each competition pair, and 2) each of

the single predator treatments and the corresponding competition treatment.

To understand the mechanism through which predation and temperature jointly affect prey

evolution, we additionally quantified demographic parameter that govern species ecological

dynamics. Specifically, we calculated 1) the initial growth rate, r, at each temperature, as

$\ln(N_t) - \ln(N_0)$]/time using day 5 data and their initial densities on day 0, and, 2) maximum

density, N_max, by measuring the highest daily average across replicates. We used linear models

('stats' v4.3.1) and stepwise model selection ('stepAIC' in R package 'MASS' v7.3-60) to test

the effects of temperature and competition on each of the predators.

We also used linear mixed models to understand how abiotic and biotic factors affect prey

plasticity. Similar to previous LMMs, we tested how temperature, predator species, predator

density, and time, affect wt and vfl1-1 particle sizes in single predator treatments. In addition to

those factors, we also tested how the presence/absence of a predator competitor affects prey

particle size in the competition treatment.

Mathematical modeling

To understand the processes that drive rapid evolution in the prey population across temperatures

and in the presence/absence of a predator, we mathematically kept track the population dynamics

of a system with two genetically distinct prey strains under predation by a shared predator. We

assumed that *wt* (W) and *vfl1-1* (V) grow logistically and compete for resources at different rates

using a classic Lotka-Volterra competition framework. Further, we assumed that the predator (P)

has a multispecies type II functional response and dies at a background mortality rate, *m*. Our

model allows for the predator to prey on each strain at different rates. Taken together, the

equations modeling the strain dynamics and predator population are:

396
$$\frac{dW}{dt} = r_w W \left(1 - \frac{W}{K_w} - \frac{a_{wv}V}{K_w}\right) - \frac{d_{wp}WP}{1+d_{vp}\eta_{vp}V+d_{wp}\eta_{wp}W}$$

397
$$\frac{dV}{dt} = r_v V \left(1 - \frac{V}{K_v} - \frac{a_{vw}W}{K_v}\right) - \frac{d_{vp}VP}{1+d_{vp}\eta_{vp}V+d_{wp}\eta_{wp}W}$$

398
$$\frac{dP}{dt} = \frac{e_v d_{vp}VP}{1+d_{vp}\eta_{vp}V+d_{wp}\eta_{wp}W} + \frac{e_w d_{wp}WP}{1+d\eta_{vp}V+d_{wp}\eta_{wp}W} - mP \,,$$

where r is intrinsic growth rate, K is the carrying capacity, a_{wv} is the competition coefficient *vfl1-*

1 has on *wt* (vice versa for a_{vw}), d_{wp} and η_{wp} are the attack rate and the handling time of

predator on *wt*, a_{vp} and η_{vp} are those of predator on *vfl1-1*, e_w and e_v are the conversion

efficiencies of *wt* and *vfl1-1* to predator. We used the growth rate of *wt* and *vfl1-1* in control as r_w

and r_v and then explored parameter space with the remaining model parameters to find dynamics

that qualitatively reproduced the observed dynamics. In the appendix we also include alternative

model formulations (e.g., treating predation as a constant mortality rate, predators with a type I

functional response) and provide an analytical treatment of the model and associated predictions

(Appendix I).

Figure 1. Experimental organisms and designs. a) Images of *Chlamydomonas reinhardtii* wild

type with mNeonGreen tagged on EB1 protein and *vfl1-1* mutant strain. b) Factorial design of the

experiment, showing 3 temperatures by 3 predations by 2 competition treatments. C)

Experimental procedures of sampling methods.

Figure 2. Model prediction and empirical data of prey clonal dynamics and prey body size.

Chlamydomonas reinhardtii population density and genetic frequencies predicted by

mathematical models (a-b) and from empirical data (c-d; Specific parameter see Appendix I

XX). Panel e shows the partical sizes of *C. reinhardtii* over time. Yellow represents *vfl1-1*

mutant strain and green represents *wt*.

Figure 3. Temperature affects prey evolution through predator thermal performance and prey pasticity. a) To understand temperature effects on prey evolution, we calculated the changes in *vfl1-1* across temperature in control and all single predator treatments, using *vfl1-1* frequency at 19°C as the baseline (representing as blue horizontal line at 0). b) *wt* particle size distribution in *P. caudatum* treatment across temperature. c) shows the *P. caudatum* density (left axis) and *vfl1-1* frequency (right axis) across temperature in *P. caudatum* single predator treatment. Circles, triangles, and rectangles represent 19°C, 22°C, and 25°C respectively. Yellow and green bands on top indicate the favored functional trait at different time while the grey dash line represents the switch in selection direction after strong plastic response of *wt*.

Figure 4. Prey evolution under predator competition. a) Frequencies of *vfl1-1* in competition

treatments with two predators. Solid color bars represent the frequencies of *vfl1-1* in control and

two predator-competition treatments at different temperature. Bars with patterns represents the

frequencies of *vfl1-1* in the single predator treatments at each temperature. b) *vfl1-1* frequencies

and all predator log density with error bars at different temperature. Triangles, squares, and

circles represent *T. pyriformis*, *Glaucoma sp.*, and *P. caudatum* respectively. Open shapes

represent each replicate while the solid shapes represent the mean within each treatment. c) Wild

type size distribution measured as log FSC-H at different temperature over time. White triangles

and grey open circles represent the mean and the median.

References

Abrams, P.A. (2000). The Evolution of Predator-Prey Interactions: Theory and Evidence. *Annu. Rev. Ecol. Syst.*, 31, 79–105.

Adams, G.M., Wright, R.L. & Jarvik, J.W. (1985). Defective temporal and spatial control of flagellar assembly in a mutant of Chlamydomonas reinhardtii with variable flagellar number. *J. Cell Biol.*, 100, 955–964.

Adan, A., Alizada, G., Kiraz, Y., Baran, Y. & Nalbant, A. (2017). Flow cytometry: basic principles and applications. *Crit. Rev. Biotechnol.*, 37, 163–176.

Aljetlawi, A.A., Sparrevik, E. & Leonardsson, K. (2004). Prey–predator size-dependent functional response: derivation and rescaling to the real world. *J. Anim. Ecol.*, 73, 239–252.

Alton, L.A., Kutz, T., Bywater, C.L., Lombardi, E., Cockerell, F.E., Layh, S., *et al.* (2024). Temperature and nutrition do not interact to shape the evolution of metabolic rate. *Philos. Trans. R. Soc. B Biol. Sci.*, 379, 20220484.

Andersen, A. & Dölger, J. (2019). Planktonic encounter rates with non-spherical encounter zones. *J. R. Soc. Interface*, 16, 20190398.

Arora, M. & Sahoo, D. (2015). Green Algae. In: *The Algae World*, Cellular Origin, Life in Extreme Habitats and Astrobiology (eds. Sahoo, D. & Seckbach, J.). Springer Netherlands, Dordrecht, pp. 91–120.

Barbour, M.A. & Gibert, J.P. (2021). Genetic and plastic rewiring of food webs under climate change. *J. Anim. Ecol.*, 1365-2656.13541.

Barneche, D.R., Hulatt, C.J., Dossena, M., Padfield, D., Woodward, G., Trimmer, M., *et al.* (2021). Warming impairs trophic transfer efficiency in a long-term field experiment. *Nature*.

Bar-On, Y.M., Phillips, R. & Milo, R. (2018). The biomass distribution on Earth. *Proc. Natl. Acad. Sci.*, 115, 6506–6511.

Bartley, T.J., McCann, K.S., Bieg, C., Cazelles, K., Granados, M., Guzzo, M.M., *et al.* (2019). Food web rewiring in a changing world. *Nat. Ecol. Evol.*, 3, 345–354.

Blois, J.L., Zarnetske, P.L., Fitzpatrick, M.C. & Finnegan, S. (2013). Climate Change and the Past, Present, and Future of Biotic Interactions. *Science*, 341, 499–504.

Brans, K.I., Govaert, L., Engelen, J.M.T., Gianuca, A.T., Souffreau, C. & De Meester, L. (2017). Eco-evolutionary dynamics in urbanized landscapes: evolution, species sorting and the change in zooplankton body size along urbanization gradients. *Philos. Trans. R. Soc. B Biol. Sci.*, 372, 20160030.

Brito, E.F., Moulton, T.P., De Souza, M.L. & Bunn, S.E. (2006). Stable isotope analysis indicates microalgae as the predominant food source of fauna in a coastal forest stream, south-east Brazil. *Austral Ecol.*, 31, 623–633.

Brodin, T. & Johansson, F. (2004). CONFLICTING SELECTION PRESSURES ON THE GROWTH/PREDATION-RISK TRADE-OFF IN A DAMSELFLY. *Ecology*, 85, 2927–2932.

Calatrava, V., Tejada-Jimenez, M., Sanz-Luque, E., Fernandez, E., Galvan, A. & Llamas, A. (2023). Chlamydomonas reinhardtii, a Reference Organism to Study Algal–Microbial Interactions: Why Can't They Be Friends? *Plants*, 12, 788.

Cech, P.G., Kuster, T., Edwards, P.J. & Olde Venterink, H. (2008). Effects of Herbivory, Fire and N2-fixation on Nutrient Limitation in a Humid African Savanna. *Ecosystems*, 11, 991–1004.

Clarke, A. (2003). Costs and consequences of evolutionary temperature adaptation. *Trends Ecol. Evol.*, 18, 573–581.

Clarke, A. (2006). Temperature and the metabolic theory of ecology. *Funct. Ecol.*, 20, 405–412.

Clarke, A. & Fraser, K.P.P. (2004). Why does metabolism scale with temperature? *Funct. Ecol.*, 18, 243–251.

Cornwallis, C.K., Svensson-Coelho, M., Lindh, M., Li, Q., Stábile, F., Hansson, L.-A., *et al.* (2023). Single-cell adaptations shape evolutionary transitions to multicellularity in green algae. *Nat. Ecol. Evol.*, 7, 889–902.

Cosmo, L.G., Assis, A.P.A., de Aguiar, M.A.M., Pires, M.M., Valido, A., Jordano, P., *et al.* (2023). Indirect effects shape species fitness in coevolved mutualistic networks. *Nature*, 619, 788–792.

De Meester, L., Brans, K.I., Govaert, L., Souffreau, C., Mukherjee, S., Vanvelk, H., *et al.* (2019). Analysing eco-evolutionary dynamics—The challenging complexity of the real world. *Funct. Ecol.*, 33, 43–59.

Dell, A.I., Pawar, S. & Savage, V.M. (2011). Systematic variation in the temperature dependence of physiological and ecological traits. *Proc. Natl. Acad. Sci.*, 108, 10591–10596.

Dell, A.I., Pawar, S. & Savage, V.M. (2014). Temperature dependence of trophic interactions are driven by asymmetry of species responses and foraging strategy. *J. Anim. Ecol.*, 83, 70–84.

DeLong, J.P., Coblentz, K.E., Uiterwaal, S.F., Akwani, C. & Salsbery, M.E. (2023). Temperature and predators as interactive drivers of community properties. *Ecol. Evol.*, 13, e10665.

Descroix, A., Bec, A., Bourdier, G., Sargos, D., Sauvanet, J., Misson, B., *et al.* (2010). Fatty acids as biomarkers to indicate main carbon sources of four major invertebrate families in a large River (the Allier, France). *Fundam. Appl. Limnol.*, 177, 39–55.

Diamond, S.E., Chick, L., Perez, A., Strickler, S.A. & Martin, R.A. (2017). Rapid evolution of ant thermal tolerance across an urban-rural temperature cline. *Biol. J. Linn. Soc.*, 121, 248–257.

Diamond, S.E., Chick, L.D., Perez, A., Strickler, S.A. & Martin, R.A. (2018). Evolution of thermal tolerance and its fitness consequences: parallel and non-parallel responses to urban heat islands across three cities. *Proc. R. Soc. B Biol. Sci.*, 285, 20180036.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., *et al.* (2011). Trophic Downgrading of Planet Earth. *Science*, 333, 301–306.

Falkowski, P.G. (1994). The role of phytoplankton photosynthesis in global biogeochemical cycles. *Photosynth. Res.*, 39, 235–258.

Fischer, B.B., Kwiatkowski, M., Ackermann, M., Krismer, J., Roffler, S., Suter, M.J.F., *et al.* (2016). Phenotypic plasticity influences the eco-evolutionary dynamics of a predator–prey system.

Foissner, W. & Berger, H. (1996). A user-friendly guide to the ciliates (Protozoa, Ciliophora) commonly used by hydrobiologists as bioindicators in rivers, lakes, and waste waters, with notes on their ecology. *Freshw. Biol.*, 35, 375–482.

Foissner, W., Chao, A. & Katz, L.A. (2009). Diversity and geographic distribution of ciliates (Protista: Ciliophora). In: *Protist Diversity and Geographical Distribution*, Topics in

Biodiversity and Conservation (eds. Foissner, W. & Hawksworth, D.L.). Springer Netherlands, Dordrecht, pp. 111–129.

Fordyce, J.A. (2006). The evolutionary consequences of ecological interactions mediated through phenotypic plasticity. *J. Exp. Biol.*, 209, 2377–2383.

Frazier, M.R., Huey, R.B. & Berrigan, D. (2006). Thermodynamics Constrains the Evolution of Insect Population Growth Rates: "Warmer Is Better." *Am. Nat.*, 168, 512–520.

Frickel, J., Theodosiou, L. & Becks, L. (2017). Rapid evolution of hosts begets species diversity at the cost of intraspecific diversity. *Proc. Natl. Acad. Sci.*, 114, 11193–11198.

Gibert, J.P. (2019). Temperature directly and indirectly influences food web structure. *Sci. Rep.*, 9, 5312.

Gibert, J.P., Grady, J.M. & Dell, A.I. (2022). Food web consequences of thermal asymmetries. *Funct. Ecol.*, 1365-2435.14091.

Gillooly, J.F. (2000). Effect of body size and temperature on generation time in zooplankton. *J. Plankton Res.*, 22, 241–251.

Gilman, S.E., Urban, M.C., Tewksbury, J., Gilchrist, G.W. & Holt, R.D. (2010). A framework for community interactions under climate change. *Trends Ecol. Evol.*, 25, 325–331.

González, J., Sherr, E. & Sherr, B. (1993). Differential feeding by marine flagellates on growing versus starving, and on motile versus nonmotile, bacterial prey. *Mar. Ecol. Prog. Ser.*, 102, 257–267.

Guimarães, P.R., Pires, M.M., Jordano, P., Bascompte, J. & Thompson, J.N. (2017). Indirect effects drive coevolution in mutualistic networks. *Nature*, 550, 511–514.

Guo, F., Kainz, M.J., Sheldon, F. & Bunn, S.E. (2016). The importance of high-quality algal food sources in stream food webs – current status and future perspectives. *Freshw. Biol.*, 61, 815–831.

Hairston, N.G., Ellner, S.P., Geber, M.A., Yoshida, T. & Fox, J.A. (2005). Rapid evolution and the convergence of ecological and evolutionary time: Rapid evolution and the convergence of ecological and evolutionary time. *Ecol. Lett.*, 8, 1114–1127.

Han, Z.-Y., Wieczynski, D.J., Yammine, A. & Gibert, J.P. (2023). Temperature and nutrients drive eco-phenotypic dynamics in a microbial food web. *Proc. R. Soc. B Biol. Sci.*, 290, 20222263.

Henn, J.J., Buzzard, V., Enquist, B.J., Halbritter, A.H., Klanderud, K., Maitner, B.S., *et al.* (2018). Intraspecific Trait Variation and Phenotypic Plasticity Mediate Alpine Plant Species Response to Climate Change. *Front. Plant Sci.*, 9.

Herron, M.D., Borin, J.M., Boswell, J.C., Walker, J., Chen, I.-C.K., Knox, C.A., *et al.* (2019). De novo origins of multicellularity in response to predation. *Sci. Rep.*, 9, 2328.

Huang, B.P.-H. (1986). *Chlamydomonas reinhardtii:* A Model System for the Genetic Analysis of Flagellar Structure and Motility. In: *International Review of Cytology*, Molecular Approaches to the Study of Protozoan Cells (eds. Bourne, G.H., Danielli, J.F. & Jeon, K.W.). Academic Press, pp. 181–215.

Kilham, S.S., Kreeger, D.A., Lynn, S.G., Goulden, C.E. & Herrera, L. (1998). COMBO: a defined freshwater culture medium for algae and zooplankton. *Hydrobiologia*, 377, 147–159.

Komatsu, K.J., Avolio, M.L., Lemoine, N.P., Isbell, F., Grman, E., Houseman, G.R., *et al.* (2019). Global change effects on plant communities are magnified by time and the number of global change factors imposed. *Proc. Natl. Acad. Sci.*, 116, 17867–17873.

Kontopoulos, D.-G., Smith, T.P., Barraclough, T.G. & Pawar, S. (2020). Adaptive evolution shapes the present-day distribution of the thermal sensitivity of population growth rate. *PLOS Biol.*, 18, e3000894.

Kordas, R.L., Harley, C.D.G. & O'Connor, M.I. (2011). Community ecology in a warming world: The influence of temperature on interspecific interactions in marine systems. *J. Exp. Mar. Biol. Ecol.*, Global change in marine ecosystems, 400, 218–226.

Lindmark, M., Ohlberger, J., Huss, M. & Gårdmark, A. (2019). Size-based ecological interactions drive food web responses to climate warming. *Ecol. Lett.*, 22, 778–786.

Lister, B.C. & Garcia, A. (2018). Climate-driven declines in arthropod abundance restructure a rainforest food web. *Proc. Natl. Acad. Sci.*, 115, E10397–E10406.

Lurling, M. & Beekman, W. (2006). Palmelloids formation in Chlamydomonas reinhardtii : defence against rotifer predators? *Ann. Limnol. - Int. J. Limnol.*, 42, 65–72.

McNaughton, S.J., Banyikwa, F.F. & McNaughton, M.M. (1997). Promotion of the Cycling of Diet-Enhancing Nutrients by African Grazers. *Science*, 278, 1798–1800.

Norin, T. & Metcalfe, N.B. (2019). Ecological and evolutionary consequences of metabolic rate plasticity in response to environmental change. *Philos. Trans. R. Soc. B Biol. Sci.*, 374, 20180180.

Padfield, D., Yvon-Durocher, G., Buckling, A., Jennings, S. & Yvon-Durocher, G. (2016). Rapid evolution of metabolic traits explains thermal adaptation in phytoplankton. *Ecol. Lett.*, 19, 133–142.

Palkovacs, E.P. & Post, D.M. (2009). Experimental evidence that phenotypic divergence in predators drives community divergence in prey. *Ecology*, 90, 300–305.

Parain, E.C., Gravel, D., Rohr, R.P., Bersier, L.-F. & Gray, S.M. (2016). Mismatch in microbial food webs: predators but not prey perform better in their local biotic and abiotic conditions. *Ecol. Evol.*, 6, 4885–4897.

Parmesan, C. (2006). Ecological and Evolutionary Responses to Recent Climate Change. *Annu. Rev. Ecol. Evol. Syst.*, 37, 637–669.

Pawar, S., Dell, A.I., & Van M. Savage. (2012). Dimensionality of consumer search space drives trophic interaction strengths. *Nature*, 486, 485–489.

Petchey, O.L., McPhearson, P.T., Casey, T.M. & Morin, P.J. (1999). Environmental warming alters food-web structure and ecosystem function. *Nature*, 402, 69–72.

Rajam, M.V. & Kumar, S.V. (2007). Green Alga (Chlamydomonas reinhardtii). In: *Agrobacterium Protocols Volume 2*, Methods in Molecular Biology (ed. Wang, K.). Humana Press, Totowa, NJ, pp. 421–433.

Robertson, M.L. & Hammill, E. (2021). Temperature and prey morphology influence attack rate and handling time in a predator–prey interaction. *Hydrobiologia*, 848, 4637–4646.

Sasso, S., Stibor, H., Mittag, M. & Grossman, A.R. (2018). From molecular manipulation of domesticated Chlamydomonas reinhardtii to survival in nature. *eLife*, 7, e39233.

Savage, V.M., Gillooly, J.F., Brown, J.H., West, G.B. & Charnov, E.L. (2004). Effects of Body Size and Temperature on Population Growth. *Am. Nat.*, 163, 429–441.

Sheridan, J.A. & Bickford, D. (2011). Shrinking body size as an ecological response to climate change. *Nat. Clim. Change*, 1, 401–406.

Siepielski, A.M., Hasik, A.Z., Ping, T., Serrano, M., Strayhorn, K. & Tye, S.P. (2020). Predators weaken prey intraspecific competition through phenotypic selection. *Ecol. Lett.*, 23, 951–961.

Silflow, C.D., LaVoie, M., Tam, L.-W., Tousey, S., Sanders, M., Wu, W., *et al.* (2001). The Vfl1
Protein in Chlamydomonas Localizes in a Rotationally Asymmetric Pattern at the Distal
Ends of the Basal Bodies. *J. Cell Biol.*, 153, 63–74.

Suttle, K.B., Thomsen, M.A. & Power, M.E. (2007). Species interactions reverse grassland
responses to changing climate. *Science*, 315, 640–642.

Terblanche, J.S., Clusella-Trullas, S., Deere, J.A., Van Vuuren, B.J. & Chown, S.L. (2009).
Directional Evolution of the Slope of the Metabolic Rate–Temperature Relationship Is
Correlated with Climate. *Physiol. Biochem. Zool.*, 82, 495–503.

Thakur, M.P., Griffin, J.N., Künne, T., Dunker, S., Fanesi, A. & Eisenhauer, N. (2018).
Temperature effects on prey and basal resources exceed that of predators in an
experimental community. *Ecol. Evol.*, 8, 12670–12680.

Thunell, V., Lindmark, M., Huss, M. & Gårdmark, A. (2021). Effects of Warming on Intraguild
Predator Communities with Ontogenetic Diet Shifts. *Am. Nat.*, 198, 706–718.

Tseng, M. & O'Connor, M.I. (2015). Predators modify the evolutionary response of prey to
temperature change. *Biol. Lett.*, 11, 20150798.

Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and
species interactions in terrestrial ecosystems. *Ecol. Lett.*, 11, 1351–1363.

Ullah, H., Nagelkerken, I., Goldenberg, S.U. & Fordham, D.A. (2018). Climate change could
drive marine food web collapse through altered trophic flows and cyanobacterial
proliferation. *PLOS Biol.*, 16, e2003446.

Vasseur, D.A. & McCann, K.S. (2005). A Mechanistic Approach for Modeling Temperature-
Dependent Consumer-Resource Dynamics. *Am. Nat.*, 166, 184–198.

Visser, A.W. (2007). Motility of zooplankton: fitness, foraging and predation. *J. Plankton Res.*,
29, 447–461.

Voigt, W., Perner, J., Davis, A.J., Eggers, T., Schumacher, J., Bährmann, R., *et al.* (2003).
TROPHIC LEVELS ARE DIFFERENTIALLY SENSITIVE TO CLIMATE. *Ecology*,
84, 2444–2453.

Wieczynski, D.J., Singla, P., Doan, A., Singleton, A., Han, Z.-Y., Votzke, S., *et al.* (2021).
Linking species traits and demography to explain complex temperature responses across
levels of organization. *Proc. Natl. Acad. Sci.*, 118, e2104863118.

Yampolsky, L.Y., Schaer, T.M.M. & Ebert, D. (2014). Adaptive phenotypic plasticity and local
adaptation for temperature tolerance in freshwater zooplankton. *Proc. R. Soc. B Biol. Sci.*,
281, 20132744.

Yoshida, T., Hairston, N.G. & Ellner, S.P. (2004). Evolutionary trade–off between defence
against grazing and competitive ability in a simple unicellular alga, Chlorella vulgaris.
Proc. R. Soc. Lond. B Biol. Sci., 271, 1947–1953.

Zarnetske, P.L., Skelly, D.K. & Urban, M.C. (2012). Biotic Multipliers of Climate Change.
Science, 336, 1516–1518.

667