A transcriptional dissection of the petty spurge (*Euphorbia peplus* L.) reproductive structure

Authors: Arielle Rose Johnson[1*], Ashley Bao[1], Margaret Hannah Frank[1]
Affiliation: [1]Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca, NY, USA
*Contact: arj66@stanford.edu

Abstract: The unique reproductive structure of *Euphorbia* species, the cyathium, is an

inflorescence that resembles a single complete flower. This structure has not yet been

investigated using transcriptomics at a genomic scale. Enabled by the petty spurge (*Euphorbia*

peplus L.) genome and guided by the ABCDE model of floral development, we dissected petty

spurge cyathia and carried out a transcriptomic analysis of the different organs. We also

constructed gene phylogenies and performed dN/dS analysis on select floral genes. Unlike in

model species in the Asteraceae, which also have pseudanthia (false flowers), we did not find

evidence for lineage-specific duplications of E class genes in *Euphorbia*. Rather, our

transcriptomic and genomic analyses indicate that SEP1 paralogs have distinct expression

patterns within the cyathium. *E. peplus* filiform structures (sterile organs within the cyathium)

show upregulation of B and E class genes and transcriptomic signatures of heterochromatin

formation, consistent with some form of floral identity and potentially consistent with their being

reduced flowers in an inflorescence. The gene *CRABS CLAW* (*CRC*), which is involved in

nectary development in both arabidopsis and petunia, is not upregulated in *E. peplus* nectaries,

indicating an alternative nectary development pathway in *Euphorbia.* Finally, *Euphorbia*

homologs for the inflorescence/floral meristem genes *UFO* and *LFY* and the B class genes *AP3*

and *PI* have highly diverged sequences relative to other Euphorbiaceae species.

18
19   **Key words**: *Euphorbia peplus*, *Euphorbia*, cyathium, spurge, inflorescence
20
21

INTRODUCTION

There are approximately 2000 species of *Euphorbia*, including horticulturally important plants such as poinsettia (*Euphorbia pulcherrima* L). The reproductive structure of *Euphorbia* species is called a cyathium. A cyathium resembles a complete flower with pistils, stamens, and a perianth; however, it is a pseudanthium (false flower) that is a condensed inflorescence consisting of a pistillate flower surrounded by staminate flowers and an involucral bract (Figure 1).

Figure 1

Euphorbia peplus L., commonly known as petty spurge, is an emerging model species for *Euphorbia* with a publicly available genome assembly (Czechowski et al., 2022; Johnson et al., 2023). In *E. peplus*, each immature cyathium is enclosed within two bracts, called cyathophylls. The cyathophyll contains the immature cyathium as well as multiple smaller cyathium primordia

(enclosed in their own cyathophylls). The total number of primordia can vary by ecotype.

Although *E. peplus* is primarily self-pollinated, the cyathium remains enclosed by cyathophylls

until the stigma is receptive to pollen (indicated by maturation of stigma lobe morphology), which

brings the cyathium into the open for pollination and later seed dispersal (Li et al., 2014).

The *E. peplus* cyathium itself, as in other *Euphorbia* species, is composed of a single pistillate

flower in the center, surrounded by five mature staminate flowers. Each of the five mature

staminate flowers is subtended by immature or nonfunctional staminate flowers in a zigzag

pattern. Interspersed with the staminate flowers in the *E. peplus* cyathium are fringed filiform

(threadlike) structures. These filiform structures have been referred to by various terms

including "filamentous structures" (Prenner and Rudall, 2007) and "filiform bracteoles" (Bruyns,

2022). For the sake of clarity, we will refer to these structures as "filiform structures" in this

paper, because "filamentous" implies a developmental homology between these structures and

filaments, and because Prenner and Rudall 2007 also refer to cyathophylls as "bracteoles"

(Prenner and Rudall, 2007). The pistillate flower, staminate flowers, and filiform structures are

surrounded by a fused cuplike structure of involucral bracts, also known as the involucre. It is

this cup-shaped structure that gives cyathia their name, as "cyathium" comes from the Greek

word for cup, "kyathos". In *E. peplus*, 4 or 5 crescent-shaped nectaries are present near the top

of the involucre.

Although the cyathium has the superficial organization of a single flower (i.e., a pistillate

structure surrounded by staminate structures that are surrounded by sterile organs), it is

generally interpreted as an inflorescence based on several criteria (reviewed in Prenner and

Rudall, 2007). There is a line of constriction on the staminate flower that is interpreted as the

flower-stamen boundary, rather than individual stamens. Other genera in the Euphorbiaceae

(e.g. *Anthostema*, Figure 2) form distinct staminate flowers that are clustered around a single

pistillate flower. In addition, species within the Euphorbiaceae typically have unisexual flowers,

and the bisexual composition of the cyathium would be an exception to this trend (Kaplan and

Specht, 2022). Other genera within the Euphorbiaceae, such as *Dalechampia* and *Pera* have

reproductive structures that have long been established to be pseudanthia (i.e., inflorescences

that resemble single flowers; Gagliardi et al., 2018). Prenner and Rudall 2007 speculate that

because the cyathium is an inflorescence, the sterile filiform structures that surround the

staminate flowers could be interpreted either as prophylls or as highly reduced staminate

flowers from a dichasial inflorescence (Prenner and Rudall, 2007). They interpret the involucre

as a non-floral organ.

As compared with its relatives such as *Anthostema* (Figure 2), the *Euphorbia* inflorescence is

very compact structurally. Cyathia have a reduced axis, different placement of their flowers,

and an earlier emergence of pistillate flower primordia compared with staminate flower

primordia than inflorescences of related genera (Gagliardi et al., 2018). The Australian genus

Calycopeplus and the New Caledonia tree *Neoguillauminia cleopatra* have been proposed as

"missing links" towards the evolution of the *Euphorbia* cyathium from separate pistillate and

staminate flowers; these close relatives to *Euphorbia* both have an inflorescence where a

pistillate flower has its own small perianth that is then surrounded by staminate flowers with their

own perianth whorls (Figure 2) (Western Australian Herbarium et al., n.d.; Prenner and Rudall,

2007; Prenner et al., 2008; Dressler et al., 2014; Gilbert, 1994).

| *Calycopeplus paucifolius* | *Neoguillauminia cleopatra* | *Anthostema senegalense* |
| © overlander (Gerald Krygsman), iNaturalist | © Benoit Henry, Pl@ntNet | © Philippe Birnbaum, African Plants-- A Photo Guide |

Figure 2

Now that a high-quality genome is publicly available for two *Euphorbia* species, the cyathium

can be investigated from a comparative molecular developmental perspective. This paper will

use transcriptomics and evolutionary analysis to shed more light on the cyathium structure.

Transcriptomics of floral development of another species in the Euphorbiaceae, *Jatropha curcas*

L., showed that the ABCDE model of floral developmental regulation (Irish, 2017) is mostly

conserved in *J. curcas* (Hui et al., 2017); therefore, we decided to investigate the ABCDE genes

in *Euphorbia* as well.

Under the ABCDE model, different combinations of MADS-box transcription factors combine to

specify the four floral whorls: the sepals, petals, stamens, and carpel (Figure 1E). E class

genes are present throughout the flower and help specify all four whorls. C class transcription

factors specify the carpel; B and C genes together specify stamens; B and A genes together

specify the petal; and A genes alone specify the sepal. D class genes specify ovules.

Generally, B and C class genes are conserved between taxa, while A class genes are not highly

conserved (Litt and Kramer, 2010). Complete flowers contain all of the floral whorls, while

unisexual flowers have either functional stamens or functional carpels but not both. As defined

by Mitchell and Diggle 2005, unisexual flowers can be classified based on the presence or

absence of rudiments of the gynoecium or androecium (Type I = rudiments present; Type II =

rudiments absent) (Mitchell and Diggle, 2005). The *Euphorbia* cyathium has Type II unisexual

flowers, with no evidence of sterile carpels/stamens present. The complete disappearance of a

reproductive whorl in Type II unisexual flowers is hypothesized to result from differential

regulation of the MADS-box transcription factors (Feng et al., 2020). The flowers of the

Euphorbia cyathium also lack a perianth (sepals and petals). Loss of the perianth whorls can

result from edits to MADS-box transcription factors or to other transcription factors not related to

whorl identity, as is seen in the arabidopsis *petal loss* (*ptl*) mutant (Brewer et al., 2004).

We also wanted to contextualize the *Euphorbia* cyathium as a pseudanthium (false flower).

False flowers can form from multiple types of meristems. Claßen-Bockhoff and Bull-Hereñu

2013 use an ontogenetic framework to draw distinctions between two different types of flower-

producing meristems: 'flower unit meristems' (FUMs) which are determinate and lack apical

growth, and inflorescence meristems (IMs) which are initially indeterminate and show apical

growth (Claßen-Bockhoff and Bull-Hereñu, 2013). IMs produce the basic branched

inflorescence types, e.g. racemes and panicles, whereas FUMs are relatively similar to floral

meristems and generate dense heads of flowers. Head meristems in Asteraceae are considered

FUMs based on their mantle-core structure and lack of apical growth (Zhang and Elomaa,

2021). Morphological work on *Ricinus communis* (Euphorbiaceae), reinterpreted its "staminate

flower" which forms many-branched "staminal trees" as a multi-flower unit with each flower

reduced to a single anther, forming from an FUM (Claßen-Bockhoff and Frankenhäuser, 2020).

The *Euphorbia* cyathium similarly has a meristem that could be interpreted as an FUM, as it

initially lacks visible primordia (Prenner and Rudall, 2007). Because of their ontogenetic

similarity, we compared the *Euphorbia* cyathium with the Asteraceae pseudanthium.

MATERIALS AND METHODS

Plant growth conditions—

E. peplus plants were grown under long day conditions with 25 °C day/16 °C night. Seeds

were germinated without any pretreatment in closed Phytatrays (P5929; Sigma-Aldrich, Saint

Louis, MO, USA) in LM 1-1-1 soil mix, then week-old plants were uppotted to 4-inch pots

containing LM 1-1-1 soil mix.

Imaging—

Images of cyathia were taken using a Leica M205 FCA stereo microscope with a DMC6200

camera.

Fixation and dissection for RNA sequencing—

Cyathia from twelve-week-old *E. peplus* plants were collected into ice-cold acetone then stored

at -20°C until dissection. Cyathia were dissected under the dissecting scope in acetone on a

cold block and dissected organs were put immediately into chilled acetone tubes. 5 biological

replicates of 10 different organs were collected, with 10 cyathia dissected for each biological

replicate.

Stages used for RNA sequencing—

We used primordia approximately 200 μm tall for our "small primordium" samples. We used

cyathium primordia approximately 500 μm tall for our "large primordium" samples. We used

staminate flower primordia approximately 200 μm tall for our "young staminate flowers" and

pistillate flower primordia approximately 400 μm tall for our "young pistillate flowers". For the

remaining samples, we dissected them from nearly-mature cyathia still enclosed within

unopened cyathophylls when cyathia were approximately 1.5 mm tall not including cyathophylls:

nectary, involucre, mature staminate flower, mature pistillate flower, filiform structure. In our

organ stages, 'mature' here refers to developmental maturity of form, not reproductive maturity,

which roughly coincides with the opening of the cyathophylls. At reproductive maturity, cyathia

are approximately 2mm tall.

RNA extraction—

RNA was extracted using the New England Biolabs Monarch Total RNA Miniprep Kit (NEB

#T2010, Ipswich, MA, USA). Acetone was removed from the tubes under the dissecting scope,

2-3 beads (BioSpec Products #11079124, Bartlesville, OK, USA) were added, the tubes were

placed into liquid nitrogen, and the tissue was ground in a bead beater (Qiagen TissueLyser II,

Germantown, MD, USA). 400uL of 1x protection reagent from the kit was added to the ground

tissue and beads, and the tissue was ground again in the protection reagent. The liquid was

transferred to a new tube and the rest of the kit protocol was followed. RNA was eluted with

100uL nuclease-free water.

RNA purification—

The 100uL of RNA was purified by adding 10uL 3M sodium acetate, 0.75uL GlycoBlue

(ThermoFisher #AM9515, Waltham, MA USA) and 275uL ice-cold 100% ethanol, vortexing to

mix, precipitating at -80°C overnight, centrifuging at 13000 rpm at 4°C for 30 minutes, washing

the pellet twice with 0.5mL ice-cold 75% ethanol, air-drying the pellet for 5 minutes, and

resuspending the RNA in 20uL of nuclease-free water.

3' sequencing—

RNA samples were submitted to the Cornell Institute for Biotechnology Genomics Center for 3'

mRNA-seq library construction using a QuantSeq FWD kit (Lexogen, Vienna, Austria) followed

by 1x75 bp sequencing on an Illumina NextSeq500 (Illumina, Hayward, CA, USA). Sequencing

statistics were calculated with v0.15.0 seqkit stats with parameters -j 10 -T(Shen et al., 2016).

(Supplementary File 1).

Orthogroup inference—

To assemble orthogroups, we ran OrthoFinder v2.5.4 (Emms and Kelly, 2019) on the longest

isoform protein sequences from *E. peplus*, *E. lathyris*, arabidopsis, soybean, Pará rubber tree,

cassava, poplar, and castor bean using the parameters -S diamond -t 40 -a 8(Buchfink et al.,

2021). Orthogroups are included in Supplementary File 2.

Identification of ABCDE floral identity genes—

In order to identify floral identity genes of interest, we retrieved the sequences of genes of

interest from Uniprot accessed through TAIR and ran blastp with our *E. peplus* proteins file as

the query and the parameters -qcov_hsp_perc 80 e-value 1e-10(Camacho et al., 2009). Where

BLAST results surfaced multiple candidates with low e-values and high percent identity, a

reciprocal BLAST to arabidopsis was used to select the best candidate(s). We further

confirmed the results through comparison to the clades identified by OrthoFinder. Gene IDs

accessed through TAIR: AT1G69120(AP1)(Bowman et al., 1993); AT4G36920(AP2)(Jofuku et

al., 1994); AT3G54340(AP3)(Jack et al., 1992); AT4G18960(AG)(Mizukami and Ma, 1997);

AT2G45650(AGL6)(Ma et al., 1991); AT1G69180(CRC)(Alvarez and Smyth, 1999);

AT1G65480(FT)(Kobayashi et al., 1999); AT5G60910(AGL8)(Mandel and Yanofsky, 1995);

AT5G61850(LFY)(Schultz and Haughn, 1991); AT5G20240(PI)(Goto and Meyerowitz, 1994);

AT5G15800(SEP1)(Pelaz et al., 2000); AT1G24260(SEP3)(Pelaz et al., 2000);

AT3G58780(SHP1)(Liljegren et al., 2000); AT2G42830(SHP2)(Liljegren et al., 2000);

AT2G45660(AGL20)(Lee et al., 2000); AT4G09960(STK)(Favaro et al., 2003);

AT5G03840(TFL1)(Kobayashi et al., 1999); AT1G30950(UFO)(Levin and Meyerowitz, 1995).

***Transcriptomic analysis*—**

Raw read files were filtered with fastp v0.23.2 using parameters --cut_right --cut_window_size 5

--cut_mean_quality 20 --length_required 50 (Chen et al., 2018). Transcript abundance was

generated using *E. peplus* coding sequences with kallisto quant v0.42.4 with parameters --

single -b 100 -t 10 (Bray et al., 2016). The resulting data was processed in R using the DESeq2

pipeline; all genes with >20 estimated counts across all samples were considered to be

expressed and were included in the analysis (Love et al., 2014). Variance-stabilized-

transformed data were visualized in heatmaps using the package pheatmap with expression

normalized by row and genes clustered by expression pattern (Kolde, n.d.). Of the floral genes

of interest investigated (Supplementary File 3), the *E. peplus* homologs for AGL8/FUL

(Ep_chr6_g18041), TFL1 (Ep_chr7_g23231) and two paralogs of FT (Ep_chr8_g25489,

Ep_chr6_g20489) had expression too low (i.e. 20 or fewer counts across all samples) to be

analyzed. Genes are said to be differentially expressed in this paper if padj<0.05. All DESeq

filtered untransformed data is included in Supplementary File 4. Comparisons of differentially

expressed genes between different tissues are included in Supplementary File 5.

***Biological Process and Molecular Function GO term enrichment analysis*—**

GO terms were assigned to *E. peplus* genes as specified in Johnson et al 2023 (Johnson et al.,

2023); GO terms assigned to each gene are included in Supplementary File 6. In order to find

which GO terms were enriched in differentially expressed genes in different organs as

compared with cyathophylls, we ran TopGO v.2.46.0 using the "classicCount" option, using

Fisher's test to determine significance (Alexa and Rahnenfuhrer, 2021). TopGO results are

included in Supplementary File 7. We summarized and visualized the GO results using

REVIGO and downloaded the R scripts to edit the color scale and axis limits (Supek et al.,

2011). Terms with a dispensability higher than 0.9 were removed and clustered with other

terms for ease of visualization. Only terms with a dispensability less than 0.2 were labeled with

text in the visualization.

Manual curation of OrthoFinder clades—

Genes with highly truncated annotations were removed from the analysis (LFY: rubber tree

KAF2311631.1, soybean Glyma.04G202002.1.p; PHYA: soybean Glyma.03G227333.1.p; PI: *E.*

lathyris Casp05761.t1; SEP1: rubber tree KAF2325472.1). For PI, *E. lathyris* Casp33119.t1 and

E. peplus Ep_chr8_g25689.t1 were included in the PI phylogenetic analysis despite being

classified into their own orthogroup because they clearly were a result of a tandem duplication

(Casp33120.1 and Ep_chr8_g25690.t1 were included in the orthogroup with Arabidopsis PI,

AT5G20240.1) and because of high reciprocal BLAST scores with Arabidopsis PI.

Gene alignment, gene tree construction and selection analysis—

For each clade of interest (generated by OrthoFinder then manually curated as specified above;

see Supplementary File 3 for gene names), a preliminary alignment was generated using

MAFFT v7.522 with parameters --maxiterate 5000 --auto --adjustdirectionaccurately --thread 8 --

op 3. Then macse v2.07 alignSequences was used to generate a codon-aware alignment with

default parameters. For a few alignments, macse generated exclamation points due to

sequencer errors; these were converted to N. Then alignment trimming was performed in trimAl

v1.4 with parameter -gappyout. Finally, sequences were exported for phylogeny construction

using macse v2.07 exportAlignment -codonForFinalStop NNN -codonForInternalStop NNN -

keep_gap_only_sites_ON. To generate phylogenetic trees, RAxML NG v1.2.0 was run with

parameters --all --model GTR+G --threads 12 --bs-trees 1000(Kozlov et al., 2019). In order to

generate dN/dS values for each branch of the tree, PAML v4.10.6 was run on the best tree from

RAxML with model=1 and NSSites=0.   In order to test for relaxed selection at sites within

genes, PAML v4.10.6 was run on the best tree from RAxML with model=2 and NSSites=2.

Alignments were visualized in ESPript 3.0 online (https://espript.ibcp.fr/ESPript/cgi-

bin/ESPript.cgi)(Robert and Gouet, 2014). Sites flagged for positive selection based on the

Bayes empirical Bayes method (BEB) are included in Supplementary File 8.

Motif analysis—

In order to retrieve a dataset that could be used to detect the canonical LFY binding motif, a bed

file of LFY ChIPseq targets (ChIPseq MACS2 q-value 10-10, DANPOS Mnase-seq q-value 10-

50) was taken from Supplementary Data 1 from Jin et al 2021 (Jin et al., 2021). This was

converted to a fasta file of the target sequences using bedtools getfasta (Quinlan and Hall,

2010). Then MEME-ChIP from MEME v5.5.2 was used to detect the LFY canonical binding

motif using default parameters except -maxw 25 (Machanick and Bailey, 2011). The dLUBS

and mLUBS motifs were acquired from the authors of Rieu et al 2023 in pfm format and

converted to meme format using jaspar2meme -pfm from MEME v4.11.2 (Bailey et al., 2015;

Rieu et al., 2023b). Then FIMO from MEME v5.5.2 was run with default parameters on a fasta

file of the 3kb upstream of the start of the LFY target genes AP1, AG, AP3, and PI in order to

assess whether the motifs were present or not (Grant et al., 2011). MEME FIMO motif search

results are included in Supplementary File 9.

Genome assemblies used for analysis—

The Johnson et al *Euphorbia peplus* genome assembly and annotation was accessed through

Zenodo (Johnson et al., 2023): https://zenodo.org/records/8033914 The Wang et al. *Euphorbia*

lathyris assembly and annotation was accessed through figshare (Wang et al., 2021):

https://figshare.com/articles/dataset/High-

quality_genome_assembly_of_the_biodiesel_plant_Euphorbia_lathyris/14909913/1 The Liu et

al. *Hevea brasiliensis* (rubber tree) assembly and annotation was accessed through NCBI (Liu

et al., 2020): https://www.ncbi.nlm.nih.gov/assembly/GCA_010458925.1/. The Xu et al. wild

Ricinus communis assembly and annotation was accessed through NCBI (Xu et al., 2021):

https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_019578655.1/. The Bredeson et al.

Manihot esculenta v8.1 assembly and annotation was accessed through Phytozome:

https://phytozome-next.jgi.doe.gov/info/Mesculenta_v8_1 The arabidopsis Araport 11 genome

release was accessed through TAIR (Cheng et al., 2017):

https://www.arabidopsis.org/download/index-

auto.jsp?dir=%2Fdownload_files%2FGenes%2FAraport11_genome_release%2FAraport11_bla

stsets. The *Glycine max* (soybean) Wm82.a4.v1 assembly and annotation was accessed

through Phytozome (Valliyodan et al., 2019): https://phytozome-

next.jgi.doe.gov/info/Gmax_Wm82_a4_v1. The *Populus trichocarpa* v4.1 assembly and

annotation was accessed through Phytozome (Tuskan et al., 2006): https://phytozome-

next.jgi.doe.gov/info/Ptrichocarpa_v4_1

RESULTS

Analysis of ABCDE gene expression in cyathium components—

We dissected the different organs of the cyathium and sequenced their RNA (Figure 3 A-D). Of

25,473 annotated *E. peplus* genes, 17,164 were expressed in the cyathium. Staminate flowers

had the most distinct expression pattern from other organs; a clear separation is visible between

the mature staminate flowers, young staminate flowers, and other cyathium organs in the first

principal component (PC) of the data (Figure 3E). This data is consistent with many other

flowering plant species' transcriptomic pattern of distinct staminate flowers compared with other

organs, or distinct stamens compared with other floral organs in complete flowers (Klepikova et

al., 2015; Xanthopoulou et al., 2021).

Figure 3

Using BLAST and OrthoFinder, we identified *E. peplus* homologs for all classes of the ABCDE

genes and visualized their expression pattern across different organs (Figure 4).

Figure 4

In the ABCDE model, the E class genes SEPALLATA1 (SEP1) and SEPALLATA3 (SEP3) are

partially redundant in function and help specify all four floral whorls; a closely related E class

gene, AGAMOUS-LIKE 6 (AGL6), also helps specify all of the floral organs and is also involved

floral meristem regulation (Dreni and Zhang, 2016; Morel et al., 2019). There are three

orthologs of SEP1 in *Euphorbia*, which is consistent with other Euphorbiaceae (Supplementary

Figure 1). *E. peplus* SEP1_1 is significantly upregulated in both stages of pistillate flowers

compared with cyathophylls, SEP1_2 is significantly upregulated in staminate flowers as

compared with pistillate flowers, and SEP 1_3 is significantly upregulated in the filiform

structures as compared with cyathophylls (Figure 4, Supplementary File 5). *Euphorbia*'s sole

SEP3 ortholog is significantly upregulated in both stages of the pistillate and staminate flowers,

both stages of the primordia, and the filiform structures as compared with cyathophylls. AGL6 is

not significantly expressed in any tissue as compared with the cyathophylls due to low

expression overall (Supplementary File 4, Supplementary File 5).

The B class genes APETALA3 (AP3) and PISTILLATA (PI) specify petal and stamen identity;

this role is conserved across bisexual and unisexual flowers (Litt and Kramer, 2010; Zhang et

al., 2022). All three *E. peplus* AP3 paralogs and both PI paralogs are significantly upregulated

in staminate flowers compared with pistillate flowers at both stages studied, in mature staminate

flowers as compared with cyathophylls, in young staminate flowers as compared with

cyathophylls (except AP3_3), and in filiform structures as compared with cyathophylls (Figure 4,

Supplementary File 5). Of the AP3 and PI homologs, only AP3_2 is significantly expressed in

the involucres as compared with the cyathophylls.

The C class gene AGAMOUS (AG) is deeply conserved across evolution and helps specify both

pistillate and staminate development (Dreni and Kater, 2014). The *E. peplus* ortholog of AG is

significantly upregulated in both stages of both pistillate flowers and staminate flowers as

compared with cyathophylls, and not significantly upregulated in any other parts of the cyathium

(Figure 4, Supplementary File 5).

In arabidopsis, the A class genes APETALA1 (AP1) and APETALA2 (AP2) help specify sepals

and help specify petals in conjunction with B class genes, in addition to other functions in floral

meristem identity and determinacy (Han et al, 2014). Notably, these genes' functions are not

conserved outside the Brassicaceae and are unlikely to be the same in *Euphorbia* as in

arabidopsis (Litt and Kramer, 2010; Zumajo-Cardona and Pabón-Mora, 2016). There are two

homologs of arabidopsis AP1 in *E. peplus*. AP1_1, which is most similar in sequence to

arabidopsis AP1 and CAULIFLOWER (Supplementary Figure 2), is upregulated in the nectary,

involucre, filiform structures, both stages of primordia, both stages of staminate flowers, and

young pistillate flowers as compared with cyathophylls (Figure 4, Supplementary File 5). *E.*

peplus AP1_2 and AP2 are not significantly differentially expressed in any cyathium part as

compared with cyathophylls, likely due to low gene expression (Supplementary File 4,

Supplementary File 5).

We were also interested in the genes LEAFY (LFY) and UNUSUAL FLORAL ORGANS (UFO)

which are known to directly activate many of the ABCDE genes. LFY is a transcription factor

that is a key regulator of floral meristem and floral organ identity (Moyroud et al., 2010); it opens

up the chromatin around the ABCDE genes, enabling their transcription (Jin et al., 2021). In

arabidopsis, UFO interacts with LFY to activate the B class genes AP3 and PI as well as other

targets (Rieu et al., 2023b). In some species such as petunia, UFO is also required in addition

to LFY to promote floral meristem identity (Souer et al., 2008). *E. peplus* has two homologs of

LFY: LFY1_1 is significantly upregulated in both stages of primordia as compared with

cyathophylls, and LFY_2 is significantly upregulated in small primordia and filiform structures as

compared with cyathophylls (Figure 4, Supplementary File 5). *E. peplus* UFO is upregulated in

both stages of primordia, both stages of staminate flowers, young pistillate flowers, and in

filiform structures, as compared with cyathophylls. It is significantly upregulated in mature

staminate flowers as compared with mature pistillate flowers.

The gene CRABS CLAW (CRC), which is necessary for nectary development in both

arabidopsis and petunia (Morel et al., 2018; Slavković et al., 2021), is not upregulated in *E.*

peplus nectaries as compared with cyathophylls (Figure 4, Supplementary File 5), suggesting

an alternative nectary development pathway in *E. peplus*. It is significantly upregulated in both

stages of pistillate flowers as well as in filiform structures compared with cyathophylls, and

significantly upregulated in both stages of pistillate flowers compared with staminate flowers.

The upregulation of CRC in pistillate flowers in *E. peplus* is consistent with CRC's other

functions in carpel development and meristem termination and with results from *J. curcas*,

where CRC is highly upregulated during developmental stages of pistillate flowers (Xu et al.,

2016).

Transcriptomic analysis of differentially expressed genes in cyathium components

We also wanted to investigate what made different parts of the cyathium unique beyond the

curated list of floral genes we had selected. Therefore, we performed a gene ontology (GO)

term analysis on the genes significantly upregulated in mature staminate flowers, mature

pistillate flowers, young staminate flowers, young pistillate flowers, involucres, filiform structures,

and nectaries as compared with cyathophylls (Supplementary File 7). We visualized the result

for nectaries, involucres, and filiform structures based on semantic similarity of the GO terms

(Figure 5).

Figure 5

We found that genes upregulated in the nectary as compared with cyathophylls were

significantly enriched for multiple Biological Process GO terms relating to transmembrane

transport and specialized metabolism (Figure 5, Supplementary File 7). GO terms relating to

gene silencing were also enriched. The Molecular Function GO terms enriched in the nectary

included multiple significant terms relating to transmembrane transporter activity and

glycosyltransferase activity, similarly to the Biological Process GO terms. ATP and NADH

metabolism-related terms were also significantly enriched in the nectaries, suggesting an

enrichment in energy-generating redox activities.

Genes upregulated in involucres as compared with cyathophylls were also significantly enriched

in genes with Biological Process GO terms relating to specialized metabolite biosynthesis and

transmembrane transport (Figure 5, Supplementary File 7). This could potentially indicate that

some of the compounds present in the nectaries are at least partially biosynthesized in the

involucre tissue connected to the nectaries. The involucre also had several significant GO

terms relating to stress, immune responses, and defense responses. The defense GO category

may be enriched because of shared hormonal signaling between defense and development; for

example, the gene Ep_chr4_g10976, which is upregulated in involucres, is in an orthogroup with

AT1G31770(ABCG14), a transmembrane cytokinin transporter. Cytokinin is involved in

responses to osmotic stress, salt stress, and other environmental factors but also regulates

developmental processes such as inflorescence branching in *J. curcas* (Chen et al., 2019).

The involucre-upregulated gene with the highest p-value, Ep_chr5_g16200, is in an orthogroup

with AT5G03680, PETAL LOSS (PTL), which generally represses tissue growth by limiting cell

division (Supplementary File 2). (Despite its name, PTL does not regulate petal organ identity,

and its expression is not limited to petals– *ptl* mutants' lack of petals may be due to indirect

effects of tissue overgrowth disrupting auxin maxima (O'Brien et al., 2015; Takeda et al., 2022).)

This gene is upregulated in filiform structures, young pistillate flowers, mature pistillate flowers,

nectaries, and both stages of primordia as compared with cyathophylls (Supplementary File 5),

suggesting functions in patterning across multiple cyathium components.

For the genes upregulated in filiform structures as compared with cyathophylls, stamen

development Biological Process GO terms were significantly enriched (Figure 5, Supplementary

File 7).   Heterochromatin formation, gene silencing, and DNA methylation-related terms were

also all significantly enriched.

The mature staminate flowers had multiple significant Biological Process GO terms for stamen

identity and morphogenesis as well as ATP and NADPH related terms (Supplementary File 7).

The young staminate flowers had multiple significant GO terms associated with stamen and

pollen development but did not have significant ATP or NADPH related terms. The mature

pistillate flowers had two GO terms associated with petal development; the pistillate flower-

enriched gene causing this significant GO hit, Ep_chr1_g00250, is in an orthogroup with

arabidopsis AT4G16850, KronbladStorleK (KSK; the name is Swedish for petal size) which

influences auxin responses during petal development (Miller et al., 2020). Compared with the

mature pistillate flower, the younger pistillate flower was enriched for more GO terms relating to

reproductive structure development, suggesting that the patterning of the pistillate flower is

more active in this earlier stage. Both stages of both the pistillate and staminate flowers had

significant GO terms relating to gene silencing; the top five most significant mature pistillate

flower terms were all related to gene silencing.

The zinc finger motif-containing gene Ep_chr4_g09148 was in the top 10 most significant

upregulated genes in the involucre, nectary, and filiform structures, and was significantly

upregulated in every other cyathium tissue studied compared with the involucre (Supplementary

File 5). This gene shares an orthogroup with AT2G33835, FRIGIDA-ESSENTIAL 1 (FES1)

which is part of a complex that represses flowering in arabidopsis (Schmitz et al., 2005).

Evolutionary analysis of ABCDE floral identity gene sequences—

Changes in MADS-box binding partners are a driver of floral morphological diversity across

many plant families (Bartlett, 2017). For example, duplication and neofunctionalization of SEP

genes is involved in the generation of pseudanthia in the Asteraceae (Zhang et al., 2017).

Therefore, in addition to analyzing gene expression, we analyzed gene sequence evolution in

order to see whether there were changes in *Euphorbia* gene sequence and/or copy number that

might reflect changes in binding and function. We performed a codon-aware alignment, tree

construction, and dN/dS analysis using the gene sequences from *E. peplus*, *Euphorbia lathyris*,

several other Euphorbiaceae species, and the model systems arabidopsis, soybean, and poplar

(Figure 6, Supplementary Figure 1, Supplementary Figure 2, Supplementary Figure 3).

Figure 6

LFY, UFO, AP3, and PI show high sequence divergence in *Euphorbia* (Figure 6A). Compared

with the non-floral gene PHYTOCHROME A (PHYA) and with the C class gene AG, the genes

LFY, AP3, and PI have high dN/dS on their *Euphorbia* branches. The *E. peplus* and *E. lathyris*

homologs for LFY have an amino acid change from a positive amino acid to a hydrophobic

amino acid in a DNA-binding site that is otherwise deeply conserved in vascular plant lineages

dating back to the lycophytes (Figure 6B) (Rodríguez-Pelayo et al., 2022). The *E. peplus* and *E.*

lathyris homologs for UFO also differ greatly in sequence from other Euphorbiaceae homologs

of UFO, and one reason is that the *Euphorbia* versions have deletions near the start of the

protein (Figure 6C).   Interestingly, many of the AP3 residues flagged as "highly significant" for

positive selection in *Euphorbia* are in the K domain in a region involved in tetramer formation

(Lai et al., 2019), perhaps indicating a change in binding partner (Supplementary Figure 3).

Because LFY and UFO showed a high level of sequence divergence, in order to investigate

whether LFY and UFO binding targets were conserved, we looked for LFY binding motifs and

LFY-UFO binding motifs in the putative promoters of homologs of the LFY targets AP3, PI, AG,

and AP1 in *E. peplus*, *H. brasiliensis*, and arabidopsis. (AP3 and PI are targets of the LFY-UFO

complex as well.) We hypothesized that the binding motifs might not be conserved in *E. peplus*.

Both canonical LFY binding motifs and UFO-LFY motifs (mLUBS and dLUBS) are present in the

3kb upstream of these target genes in *E. peplus* (Supplementary File 9), suggesting that at least

some of the binding motifs of LFY and LFY-UFO are conserved despite the high divergence in

sequence.

DISCUSSION

Few pseudanthia have been well-characterized at the molecular level. Several studies have

examined transcriptional patterns and key developmental genes in the pseudanthium of gerbera

(*Gerbera hybrida*, Asteraceae), which has a flower unit meristem (FUM) that lacks apical

growth, initially lacks visible primordia, and fractionates into many flowers in a short

developmental span—similarly to *Euphorbia*'s cyathium FUM. In the ABCDE model, E-function

SEP genes generally promote floral identity. Gerbera has eight SEP homologs due to lineage-

specific gene duplications in both the SEP1 and SEP3 subclades; and genes in both subclades

have functionally diversified in terms of which floral organs they help specify in the gerbera

pseudanthium (Zhang et al, 2017). Interestingly, *Euphorbia* only has four SEP homologs and

they are not the result of *Euphorbia*-specific duplications (Supplementary Figure 1), indicating

that SEP gene family duplication may not have contributed as much to the development of a

pseudanthium in *Euphorbia* as in the Asteraceae. However, the *Euphorbia* SEP homologs do

show distinct expression patterns within the cyathium, with the three *Euphorbia* SEP1 homologs

all showing different expression patterns from each other. This is consistent with a broader

trend that SEP1 orthologs have been repurposed in many species to regulate bract identity and

inflorescence architecture (Dreni and Ferrándiz, 2022).

LFY and UFO are generally important in specifying floral meristem identity (Rieu et al., 2023a;

Weigel et al., 1992; Levin and Meyerowitz, 1995, Lippman et al., 2008), although this is not

necessarily true for pseudanthia. in gerbera, GhLFY is uniformly expressed in the

undifferentiated FUM and regulates its determinacy, while GhUFO is expressed in emerging

flower primordia and regulates floral meristem identity (Zhao et al, 2016). The LFY homolog in

J. curcas (Euphorbiaceae) does promote floral meristem identity; its overexpression causes

early flowering, and its co-suppression delays flower formation, promotes more inflorescence

branching, and the formation of sepal-like organs (Tang et al., 2016). Although our dataset

lacks the spatial and temporal resolution needed at the earliest time points to distinguish which

genes are expressed in the cyathium FUM as opposed to the floral meristems, both LFY and

UFO are expressed highly in our small primordium and large primordium stage suggesting that

they are both important in floral meristem and/or cyathium FUM identity.

In our analysis of gene sequence evolution, *Euphorbia* LFY, UFO, and the B-class genes AP3

and PI show signs of divergence based on phylogenetic distance and relaxed selection. Except

for UFO (which may have lacked signs of relaxed selection due to the gene tree topology),

these genes also had multiple sites that were classified as highly significant for positive

selection in Bayes Empirical Bayes (BEB) analysis (Supplementary File 8). The increased rates

of molecular evolution of AP3 and PI in *Euphorbia* is consistent with increased rates of

molecular evolution of B class genes in other lineages that have lost petals, for example the

Piperaceae and Saururaceae compared with other Piperales (Jaramillo and Kramer, 2007).

This likely reflects a release of evolutionary constraints when B class genes no longer need to

promote petal identity.

One interesting observation regarding the expression of UFO is that in *J. curcas*, UFO is

downregulated in both mature staminate flowers and mature pistillate flowers compared to

younger stages (Chen et al., 2016), whereas in our *E. peplus* dataset we saw the highest

expression of UFO in mature staminate flowers. This either suggests a repurposing of UFO, or

that developmentally mature but not reproductively mature (i.e. not elongated) staminate flowers

in the *E. peplus* cyathium aren't yet transcriptomically mature compared with pistillate flowers.

As UFO functions in the development of individual perianth organs as well as in the broader

activation of the B class genes (Takeda et al., 2022), some of the changes in UFO expression

and sequence in *Euphorbia* could be related to the suppression of perianth development in

addition to potential functions in defining the floral meristem.

Genes in the AP1 clade can also be involved in floral meristem specification– and are also

highly diverged in *Euphorbia*. In arabidopsis, AGL79 interacts with SUPPRESSOR OF

OVEREXPRESSION OF CONSTANS1 (SOC1) to regulate the expression of TERMINAL

FLOWER 1 (TFL1) which controls inflorescence meristem identity (Yang et al., 2023). This AP1

(AGL79) homolog in *Euphorbia* appears to have diverged, and now groups in a separate clade

within the AP1 clade "missing" its AP1-CAL homolog in the Euphorbiaceae AGL79 clade in our

gene tree because that gene, AP1_2, is so diverged that it appears elsewhere in the tree

(Supplementary Figure 2).

One question we had about the cyathium was which other organs might have homology with the

filiform structures. In the filiform structures, we observed significant expression of B-class and

E-class genes, as well as one homolog of LFY, potentially consistent with floral identity. The

filiform structures had low levels of RNA overall compared with other structures in the cyathium

(Supplementary File 1), which could be explained by the gene ontology analysis showing

significant GO terms relating to heterochromatin formation. Our data therefore agrees with a

potential interpretation of the filiform structures as reduced flowers in an inflorescence. Overall,

our GO term analysis showed that gene silencing seems to be involved in the creation of

multiple cyathium organs.

The C class gene AGAMOUS is highly expressed in both pistillate flowers and staminate

flowers, as would be expected given its function in pistillate and staminate development. We

did not see transcriptomic signs of abortion of other whorls in the pistillate and staminate flowers

(e.g. hypothetically, seeing B class expression in the young pistillate flower could have been an

example of that), besides the upregulation of the putative homolog of the arabidopsis perianth

gene KSK in the pistillate flower. Organ abortion can happen at many different stages,

however, and unisexual flowers do not always show evidence of their missing whorls (Mitchell

and Diggle, 2005).

In future studies, it would be fruitful to expand this analysis to closely related species given

Prenner and Rudall's interpretation of the Australian genus *Calycopeplus* and the New

Caledonia tree *Neoguillauminia cleopatra* as "missing links" towards the *Euphorbia* cyathium;

future studies could sequence the floral genes of *Calycopeplus*, *Neoguillauminia*, and

Anthostema to study the sequence evolution of LFY, UFO, AP1 and B class genes with those

taxa included to see how much of the sequence divergence occurred in *Euphorbia* alone and

how much is shared with these other taxa.

In addition to expanding our analyses to more taxa, future studies could experimentally

investigate the function of the UFO-LFY complex and its DNA binding targets in *Euphorbia*, e.g.

through electrophoretic mobility shift assays (EMSA), microscale thermophoresis (MST), or

DAP-seq. Additionally, yeast two-hybrid or similar techniques could be used to check the

transcription factor binding partners of *Euphorbia*'s B-class genes. It would also be interesting

to use ATAC-seq or another method to determine chromatin accessibility across the genome

and speculate on what might be being silenced in the different cyathium structures. *In situ*

hybridization, if optimized in this system, could also potentially reveal more spatially and

temporally resolved patterns of gene expression, including more detailed expression patterns of

LFY and UFO. We are excited to see what future work reveals about the developmental

genetics involved in creating the complex structure of the cyathium.

ACKNOWLEDGEMENTS

Thank you to Jacob Landis for extremely helpful advice on running PAML. Thank you to

François Parcy, Jérémy Lucas and Laura Turchi for graciously sending the motif files for

mLUBS and dLUBS. Thank you to the Cornell Weill growth chamber plant maintenance staff,

especially Corbin Lay, for their excellent upkeep of the plants used. Thank you to the Frank Lab

for very helpful comments on the manuscript. This work was funded by a Cornell Institute of

Biotechnology Seed Grant, National Science Foundation Integrative Organismal Systems (NSF

IOS) award 1942437, and startup funds from the College of Agriculture and Life Sciences at

Cornell University to Margaret Frank.

AUTHOR CONTRIBUTIONS
603

A.R.J., A.B., and M.H.F. all conceived of the study. Under the supervision of A.R.J., A.B.

collected the images of cyathia and dissected cyathia components, and performed preliminary

dissections and RNA extractions that contributed to the experimental design for the RNAseq

experiment. A.R.J. performed the dissection for the RNAseq experiment. A.B. and A.R.J. both

contributed to the RNA extraction for the RNAseq experiment. A.B. and A.R.J. collaborated on

the initial sequence alignment and phylogeny construction to assess conservation of sequence

similarity in floral genes. A.R.J. performed all other data analysis. A.R.J. wrote the first draft of

the manuscript, after which all authors contributed to reviewing and editing the manuscript.

DATA AVAILABILITY
615

The raw RNAseq data generated in this study are available from NCBI GenBank under the

accession PRJNA1063728. All code used in the data analysis is available on Github at

https://github.com/ariellerjohnson/Cyathium-2024.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

LITERATURE CITED

Alexa, A., and J. Rahnenfuhrer. 2021. topGO: enrichment analysis for gene ontology. *R*

package version 2.46: 2021.

Alvarez, J., and D. R. Smyth. 1999. CRABS CLAW and SPATULA, two Arabidopsis genes that

control carpel development in parallel with AGAMOUS. *Development* 126: 2377–2386.

Bailey, T. L., J. Johnson, C. E. Grant, and W. S. Noble. 2015. The MEME Suite. *Nucleic acids*

research 43: W39–49.

Bartlett, M. E. 2017. Changing MADS-Box Transcription Factor Protein-Protein Interactions as a

Mechanism for Generating Floral Morphological Diversity. *Integrative and comparative*

biology 57: 1312–1321.

Bowman, J. L., J. Alvarez, D. Weigel, E. M. Meyerowitz, and D. R. Smyth. 1993. Control of

flower development in Arabidopsis thaliana by APETALA1 and interacting genes.

Development 119: 721–743.

Bray, N. L., H. Pimentel, P. Melsted, and L. Pachter. 2016. Near-optimal probabilistic RNA-seq

quantification. *Nature biotechnology* 34: 525–527.

Brewer, P. B., P. A. Howles, K. Dorian, M. E. Griffith, T. Ishida, R. N. Kaplan-Levy, A. Kilinc, and

D. R. Smyth. 2004. PETAL LOSS, a trihelix transcription factor gene, regulates perianth

architecture in the Arabidopsis flower. *Development* 131: 4035–4045.

Bruyns, P. V. 2022. Euphorbia in Southern Africa: Volume 1. Springer Nature.

Buchfink, B., K. Reuter, and H.-G. Drost. 2021. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nature methods* 18: 366–368.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L. Madden.

2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421.

Cheng, C.-Y., V. Krishnakumar, A. P. Chan, F. Thibaud-Nissen, S. Schobel, and C. D. Town.

2017. Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.

The Plant journal: for cell and molecular biology 89: 789–804.

Chen, M.-S., B.-Z. Pan, Q. Fu, Y.-B. Tao, J. Martínez-Herrera, L. Niu, J. Ni, et al. 2016.

Comparative Transcriptome Analysis between Gynoecious and Monoecious Plants

Identifies Regulatory Networks Controlling Sex Determination in Jatropha curcas. *Frontiers*

in plant science 7: 1953.

Chen, M.-S., M.-L. Zhao, G.-J. Wang, H.-Y. He, X. Bai, B.-Z. Pan, Q.-T. Fu, et al. 2019.

Transcriptome analysis of two inflorescence branching mutants reveals cytokinin is an

important regulator in controlling inflorescence architecture in the woody plant Jatropha

curcas. *BMC plant biology* 19: 468.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* 34: i884–i890.

Claßen-Bockhoff, R., and H. Frankenhäuser. 2020. The 'Male Flower' of Ricinus communis

(Euphorbiaceae) Interpreted as a Multi-Flowered Unit. *Frontiers in cell and developmental*

biology 8: 313.

Claßen-Bockhoff, R., and K. Bull-Hereñu. 2013. Towards an ontogenetic understanding of

inflorescence diversity. *Ann. Bot.* 112: 1523-1542.

Czechowski, T., E. Forestier, S. H. Swamidatta, A. D. Gilday, A. Cording, T. R. Larson, D.

Harvey, et al. 2022. Gene discovery and virus-induced gene silencing reveal branched

pathways to major classes of bioactive diterpenoids in Euphorbia peplus. *Proceedings of*

the National Academy of Sciences of the United States of America 119: e2203890119.

Dreni, L., and C. Ferrándiz. 2022. Tracing the Evolution of the SEPALLATA Subfamily across

Angiosperms Associated with Neo- and Sub-Functionalization for Reproductive and

Agronomically Relevant Traits. *Plants* 11.

Dreni, L., and M. M. Kater. 2014. MADS reloaded: evolution of the AGAMOUS subfamily genes.

The New phytologist 201: 717–732.

Dreni, L., and D. Zhang. 2016. Flower development: the evolutionary history and functions of

the AGL6 subfamily MADS-box genes. *Journal of experimental botany* 67: 1625–1638.

Dressler, S., M. Schmidt, and G. Zizka. 2014. Introducing African plants—A photo guide—an

interactive photo data-base and rapid identification tool for continental Africa. *Taxon* 63:

1159–1161.

Emms, D. M., and S. Kelly. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome biology* 20: 238.

Favaro, R., A. Pinyopich, R. Battaglia, M. Kooiker, L. Borghi, G. Ditta, M. F. Yanofsky, et al.

2003. MADS-box protein complexes control carpel and ovule development in Arabidopsis.

The Plant cell 15: 2603–2611.

Feng, G., B. J. Sanderson, K. Keefover-Ring, J. Liu, T. Ma, T. Yin, L. B. Smart, et al. 2020.

Pathways to sex determination in plants: how many roads lead to Rome? *Current opinion in*

plant biology 54: 61–68.

Gagliardi, K. B., I. Cordeiro, and D. Demarco. 2018. Structure and development of flowers and

inflorescences in Peraceae and Euphorbiaceae and the evolution of pseudanthia in

Malpighiales. *PloS one* 13: e0203954.

Gilbert, M. G. 1994. The relationships of the Euphorbieae (Euphorbiaceae). *Annals of the*

Missouri Botanical Garden. Missouri Botanical Garden 81: 283.

Goto, K., and E. M. Meyerowitz. 1994. Function and regulation of the Arabidopsis floral

homeotic gene PISTILLATA. *Genes & development* 8: 1548–1560.

Grant, C. E., T. L. Bailey, and W. S. Noble. 2011. FIMO: scanning for occurrences of a given

motif. *Bioinformatics* 27: 1017–1018.

Han, Y., C. Zhang, H. Yang, and Y. Jiao. 2014. Cytokinin pathway mediates *APETALA1*

function in the establishment of determinate floral meristems in *Arabidopsis*. *PNAS* 111:

6840-6845.

Hui, W., Y. Yang, G. Wu, C. Peng, X. Chen, and M. Z. Zayed. 2017. Transcriptome profile

analysis reveals the regulation mechanism of floral sex differentiation in Jatropha curcas L.

Scientific reports 7: 16421.

Irish, V. 2017. The ABC model of floral development. *Current biology: CB* 27: R887–R890.

Jack, T., L. L. Brockman, and E. M. Meyerowitz. 1992. The homeotic gene APETALA3 of

Arabidopsis thaliana encodes a MADS box and is expressed in petals and stamens. *Cell*

68: 683–697.

Jaramillo, M.A., and E. M. Kramer. 2007. Molecular evolution of the petal and stamen identity

genes, *APETALA3* and *PISTILLATA*, after petal loss in the Piperales. *Molecular*

phylogenetics and evolution 44: 598-609.

Jin, R., S. Klasfeld, Y. Zhu, M. Fernandez Garcia, J. Xiao, S.-K. Han, A. Konkol, and D. Wagner.

2021. LEAFY is a pioneer transcription factor and licenses cell reprogramming to floral fate.

Nature communications 12: 626.

Jofuku, K. D., B. G. den Boer, M. Van Montagu, and J. K. Okamuro. 1994. Control of

Arabidopsis flower and seed development by the homeotic gene APETALA2. *The Plant cell*

6: 1211–1225.

Johnson, A. R., Y. Yue, S. B. Carey, S. J. Park, L. H. Kruse, A. Bao, A. Pasha, et al. 2023.

Chromosome-level genome assembly of Euphorbia peplus, a model system for plant latex,

reveals that relative lack of Ty3 transposons contributed to its small genome size. *Genome*

biology and evolution 15: evad018.

Kaplan, D., and C. D. Specht. 2022. Kaplan's Principles of Plant Morphology. CRC Press.

Klepikova, A. V., M. D. Logacheva, S. E. Dmitriev, and A. A. Penin. 2015. RNA-seq analysis of

an apical meristem time series reveals a critical point in Arabidopsis thaliana flower

initiation. *BMC genomics* 16: 466.

Kobayashi, Y., H. Kaya, K. Goto, M. Iwabuchi, and T. Araki. 1999. A pair of related genes with

antagonistic roles in mediating flowering signals. *Science* 286: 1960–1962.

Kolde. Pheatmap: pretty heatmaps. *R package version*.

Kozlov, A. M., D. Darriba, T. Flouri, B. Morel, and A. Stamatakis. 2019. RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics 35: 4453–4455.

Lai, X., H. Daher, A. Galien, V. Hugouvieux, and C. Zubieta. 2019. Structural Basis for Plant

MADS Transcription Factor Oligomerization. *Computational and structural biotechnology*

journal 17: 946–953.

Lee, H., S. S. Suh, E. Park, E. Cho, J. H. Ahn, S. G. Kim, J. S. Lee, et al. 2000. The

AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in

Arabidopsis. *Genes & development* 14: 2366–2376.

Levin, J. Z., and E. M. Meyerowitz. 1995. UFO: an Arabidopsis gene involved in both floral

meristem and floral organ development. *The Plant cell* 7: 529–548.

Li, C., J. Su, X. Liu, S. Chen, and L. He. 2014. Pistillate Flower Development and Stigma

Receptivity of Euphorbia pulcherrima. *Agricultural Science & Technology* 15: 1671–1675.

Liljegren, S. J., G. S. Ditta, Y. Eshed, B. Savidge, J. L. Bowman, and M. F. Yanofsky. 2000.

SHATTERPROOF MADS-box genes control seed dispersal in Arabidopsis. *Nature* 404:

766–770.

Lippman, Z. B., O. Cohen, J. P. Alvarez, M. Abu-Abied, I. Pekker, I. Paran, Y. Eshed, and D.

Zamir. 2008. The making of a compound inflorescence in tomato and related nightshades.

PLoS biology 6: e288.

Litt, A., and E. M. Kramer. 2010. The ABC model and the diversification of floral organ identity.

Seminars in cell & developmental biology 21: 129–137.

Liu, J., C. Shi, C.-C. Shi, W. Li, Q.-J. Zhang, Y. Zhang, K. Li, et al. 2020. The Chromosome-

Based Rubber Tree Genome Provides New Insights into Spurge Genome Evolution and

Rubber Biosynthesis. *Molecular plant* 13: 336–350.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome biology* 15: 550.

Machanick, P., and T. L. Bailey. 2011. MEME-ChIP: motif analysis of large DNA datasets.

Bioinformatics 27: 1696–1697.

Ma, H., M. F. Yanofsky, and E. M. Meyerowitz. 1991. AGL1-AGL6, an Arabidopsis gene family

with similarity to floral homeotic and transcription factor genes. *Genes & development* 5:

484–495.

Mandel, M. A., and M. F. Yanofsky. 1995. The Arabidopsis AGL8 MADS box gene is expressed

in inflorescence meristems and is negatively regulated by APETALA1. *The Plant cell* 7:

1763–1771.

Miller, C. N., J. Dumenil, F. H. Lu, C. Smith, N. McKenzie, V. Chapman, J. Ball, et al. 2020.

Variation in the expression of a transmembrane protein influences cell growth in

Arabidopsis thaliana petals by altering auxin responses. *BMC plant biology* 20: 482.

Mitchell, C. H., and P. K. Diggle. 2005. The evolution of unisexual flowers: morphological and

functional convergence results from diverse developmental transitions. *American journal of*

botany 92: 1068–1076.

Mizukami, Y., and H. Ma. 1997. Determination of Arabidopsis floral meristem identity by

AGAMOUS. *The Plant cell* 9: 393–408.

Morel, P., P. Chambrier, V. Boltz, S. Chamot, F. Rozier, S. Rodrigues Bento, C. Trehin, et al.

2019. Divergent Functional Diversification Patterns in the SEP/AGL6/AP1 MADS-Box

Transcription Factor Superclade. *The Plant cell* 31: 3033–3056.

Morel, P., K. Heijmans, K. Ament, M. Chopy, C. Trehin, P. Chambrier, S. Rodrigues Bento, et al.

2018. The Floral C-Lineage Genes Trigger Nectary Development in Petunia and

Arabidopsis. *Plant Cell* 30: 2020–2037.

Moyroud, E., E. Kusters, M. Monniaux, R. Koes, and F. Parcy. 2010. LEAFY blossoms. *Trends*

in plant science 15: 346–352.

O'Brien, M., R. N. Kaplan-Levy, T. Quon, P. G. Sappl, and D. R. Smyth. 2015. PETAL LOSS, a

trihelix transcription factor that represses growth in Arabidopsis thaliana, binds the energy-

sensing SnRK1 kinase AKIN10. *Journal of experimental botany* 66: 2475–2485.

Pelaz, S., G. S. Ditta, E. Baumann, E. Wisman, and M. F. Yanofsky. 2000. B and C floral organ

identity functions require SEPALLATA MADS-box genes. *Nature* 405: 200–203.

Prenner, G., N. I. Cacho, D. Baum, and P. J. Rudall. 2010. Is LEAFY a useful marker gene for

the flower–inflorescence boundary in the Euphorbia cyathium? *Journal of experimental*

botany 62: 345–350.

Prenner, G., S. D. Hopper, and P. J. Rudall. 2008. Pseudanthium development in Calycopeplus

paucifolius, with particular reference to the evolution of the cyathium in Euphorbieae

(Euphorbiaceae–Malpighiales). *Australian systematic botany* 21: 153–161.

Prenner, G., and P. J. Rudall. 2007. Comparative ontogeny of the cyathium in Euphorbia

(Euphorbiaceae) and its allies: exploring the organ-flower-inflorescence boundary.

American journal of botany 94: 1612–1629.

Quinlan, A. R., and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. *Bioinformatics* 26: 841–842.

Rieu, P., M. Arnoux-Courseaux, G. Tichtinsky, and F. Parcy. 2023a. Thinking outside the F-box:

how UFO controls angiosperm development. *The New phytologist* 240: 945–959.

Rieu, P., L. Turchi, E. Thévenon, E. Zarkadas, M. Nanao, H. Chahtane, G. Tichtinsky, et al.

2023b. The F-box protein UFO controls flower development by redirecting the master

transcription factor LEAFY to new cis-elements. *Nature plants* 9: 315–329.

Robert, X., and P. Gouet. 2014. Deciphering key features in protein structures with the new

ENDscript server. *Nucleic acids research* 42: W320–4.

Rodríguez-Pelayo, C., B. A. Ambrose, A. Vasco, J. F. Alzate, and N. Pabón-Mora. 2022.

Evolution and expression of LEAFY genes in ferns and lycophytes. *EvoDevo* 13: 2.

Schmitz, R. J., L. Hong, S. Michaels, and R. M. Amasino. 2005. FRIGIDA-ESSENTIAL 1

interacts genetically with FRIGIDA and FRIGIDA-LIKE 1 to promote the winter-annual habit

of Arabidopsis thaliana. *Development* 132: 5471–5478.

Schultz, E. A., and G. W. Haughn. 1991. LEAFY, a Homeotic Gene That Regulates

Inflorescence Development in Arabidopsis. *The Plant cell* 3: 771–781.

Shen, W., S. Le, Y. Li, and F. Hu. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for

FASTA/Q File Manipulation. *PloS one* 11: e0163962.

Slavković, F., C. Dogimont, H. Morin, A. Boualem, and A. Bendahmane. 2021. The Genetic

Control of Nectary Development. *Trends in plant science* 26: 260–271.

Souer, E., A. B. Rebocho, M. Bliek, E. Kusters, R. A. M. de Bruin, and Ronald Koes. 2008.

Patterning of Inflorescences and Flowers by the F-Box Protein DOUBLE TOP and the

LEAFY Homolog ABERRANT LEAF AND FLOWER of Petunia. *The Plant Cell* 20: 2033-

2048.

Supek, F., M. Bošnjak, N. Škunca, and T. Šmuc. 2011. REVIGO summarizes and visualizes

long lists of gene ontology terms. *PloS one* 6: e21800.

Takeda, S., Y. Hamamura, T. Sakamoto, S. Kimura, M. Aida, and T. Higashiyama. 2022. Non-

cell-autonomous regulation of petal initiation in Arabidopsis thaliana. *Development* 149.

Tang, M., Y.-B. Tao, Q. Fu, Y. Song, L. Niu, and Z.-F. Xu. 2016. An ortholog of LEAFY in

Jatropha curcas regulates flowering time and floral organ development. *Scientific reports* 6:

37306.

Tuskan, G. A., S. Difazio, S. Jansson, J. Bohlmann, I. Grigoriev, U. Hellsten, N. Putnam, et al.

2006. The genome of black cottonwood, Populus trichocarpa (Torr. & Gray). *Science* 313:

1596–1604.

Valliyodan, B., S. B. Cannon, P. E. Bayer, S. Shu, A. V. Brown, L. Ren, J. Jenkins, et al. 2019.

Construction and comparison of three reference-quality genome assemblies for soybean.

The Plant journal: for cell and molecular biology 100: 1066–1082.

Wang, M., Z. Gu, Z. Fu, and D. Jiang. 2021. High-quality genome assembly of an important

biodiesel plant, Euphorbia lathyris L. *DNA research: an international journal for rapid*

publication of reports on genes and genomes 28.

Weigel, D., J. Alvarez, D. R. Smyth, M. F. Yanofsky, and E. M. Meyerowitz. 1992. LEAFY

controls floral meristem identity in Arabidopsis. *Cell* 69: 843–859.

Western Australian Herbarium, Biodiversity, Conservation Science, Department of Biodiversity,

Conservation, and Attractions. Calycopeplus Planch. Website

https://florabase.dbca.wa.gov.au/browse/profile/21684 [accessed 25 January 2024].

Xanthopoulou, A., J. Montero-Pau, B. Picó, P. Boumpas, E. Tsaliki, H. S. Paris, A. Tsaftaris, et

al. 2021. A comprehensive RNA-Seq-based gene expression atlas of the summer squash

(Cucurbita pepo) provides insights into fruit morphology and ripening mechanisms. *BMC*

genomics 22: 341.

Xu, G., J. Huang, Y. Yang, and Y.-A. Yao. 2016. Transcriptome Analysis of Flower Sex

Differentiation in Jatropha curcas L. Using RNA Sequencing. *PloS one* 11: e0145613.

Xu, W., D. Wu, T. Yang, C. Sun, Z. Wang, B. Han, S. Wu, et al. 2021. Genomic insights into the

origin, domestication and genetic basis of agronomic traits of castor bean. *Genome biology*

22: 113.

Yang, H., P. Zhang, D. Guo, N. Wang, H. Lin, X. Wang, and L. Niu. 2023. Transcriptional

repressor AGL79 positively regulates flowering time in Arabidopsis. *Journal of plant*

physiology 285: 153985.

Zhang, T. and P. Elomaa. 2021. Don't be fooled: false flowers in Asteraceae. *Current opinion*

in plant biology 59: 101972.

Zhang, T., Y. Zhao, I. Juntheikki, K. Mouhu, S. K. Broholm, A. S. Rijpkema, L. Kins, T. Lan, V.

A. Albert, T. H. Teeria, and P. Elomaa. 2017. Dissecting functions of *SEPALLATA*-like

MADS box genes in patterning of the pseudanthial inflorescence of *Gerbera hybrida*. *New*

phytologist 216: 939-954.

Zhang, X., L. Pan, W. Guo, Y. Li, and W. Wang. 2022. A convergent mechanism of sex

determination in dioecious plants: Distinct sex-determining genes display converged

regulation on floral B-class genes. *Frontiers in plant science* 13: 953445.

Zhao, Y., T. Zhang, S. K. Broholm, S. Tähtiharju, K. Mouhu, V. A. Albert, T. H. Teeri, and P.

Elomaa. 2016. Evolutionary Co-Option of Floral Meristem Identity Genes for Patterning of

the Flower-Like Asteraceae Inflorescence. *Plant physiology* 172: 284-296.

Zumajo-Cardona, C., and N. Pabón-Mora. 2016. Evolution of the APETALA2 Gene Lineage in

Seed Plants. *Molecular biology and evolution* 33: 1818–1832.

FIGURE LEGENDS

Figure 1: The *Euphorbia* cyathium is a unique reproductive structure. A. Diagram of a *Euphorbia*

peplus cyathium; involucre is cut away to reveal filiform structures and staminate flowers inside.

B. *E. peplus* cyathium and adjacent primordia and cyathophylls. Scale=2mm. C. *E. peplus*

cyathium and adjacent primordia with adjacent cyathophylls removed. Scale=1mm. D. *E.*

peplus cyathium with involucre and nectaries removed. Scale=500 microns. E. Diagram of the

ABCDE model of floral development and the four whorls of a traditional flower; D class genes,

which control ovule development, are not shown.

Figure 2: Close relatives of *Euphorbia*, *Calycopeplus* and *Neoguillauminia*, have perianths

around their pistillate and staminate flowers and a condensed inflorescence similar to that of

Euphorbia. *Anthostema*, which is less closely related, has perianths around its pistillate and

staminate flowers and a less-condensed inflorescence.

Figure 3: Staminate flowers have a distinct expression pattern. A-D. Illustration of cyathium

parts used for RNAseq sampling. A. Small primordium. B. Large primordium. C. Young

staminate flower and young pistillate flower. D. Mature organs: nectary, involucre, staminate

flower, pistillate flower, filiform structure, involucre, nectary. (Duplicates of organs are omitted

for simplicity. Cyathophyll is not shown.) E. PCA plot of variance-stabilized-transformed data

for the 500 genes with the most variance. PC1 explains 22.4% of the variance, PC2 explains

18.9% of the variance, PC3 explains 16.7% of the variance, and PC4 explains 10.6% of the

variance.

Figure 4: E class gene SEP3 is highly expressed in the pistillate and staminate flowers. A.

Heatmap shows expression of ABCDE homologs and other floral identity genes in different

organs of the *E. peplus* cyathium. B-N. Images of the organs sampled. B. Young pistillate

flower. C. Young staminate flower. D. Nectary. E. Filiform structure. F. Mature pistillate flower.

G. Mature staminate flower. Scale bars = 250 microns.

Figure 5: Biological Process GO terms for genes significantly upregulated in the nectary,

involucre, and filiform structures as compared with the cyathophyll and Molecular Function GO

terms for genes significantly upregulated in the nectary as compared with the cyathophyll are

summarized using REVIGO. "Value" is the log10(p-value); a lower value, towards the red side

of the color scale, is more significant. The size of the points, the dispensability, represents the

semantic similarity to other terms in the dataset.

Figure 6: LFY, UFO, and B class genes show signs of divergence and relaxed selection in

Euphorbia. A. dN/dS values shown on gene trees of floral genes of interest; RAxML support

value from 1000 bootstraps is shown at nodes.   PHYA is shown as a non-floral "control".   B.

Visualization of amino acid alignment in a highly conserved region of LFY; site 279 is a DNA

binding site that is highly conserved (except in *Euphorbia*). Cyan color shows sites in the LFY

protein alignment rated as "highly significant" for positive selection in the *Euphorbia* clade by

PAML. C. Visualization of amino acid alignment at the start of the UFO protein showing

deletions in *Euphorbia* ("Ep" and "Casp" genes). Gene names and their gene ID equivalents are

listed in Supplementary File 3.

SUPPLEMENTAL DATA FILE TITLES

Supplementary Figures (combined into a single PDF)

Supplementary File 1—Sequence statistics

Supplementary File 2—Orthogroups

Supplementary File 3—Gene names

Supplementary File 4—DESeq data

Supplementary File 5—Differential expression

Supplementary File 6—E. peplus GO terms

Supplementary File 7—TopGO results

Supplementary File 8—Positive selection sites

Supplementary File 9—Motif search results