1 Transcriptional dissection of petty spurge (*Euphorbia peplus* L.) reproductive structures
2 supports the *Euphorbia* cyathium as an inflorescence rather than a complete flower
3

Authors: Arielle Rose Johnson[1], Ashley Bao[1], Margaret Hannah Frank[1]
Affiliation: [1]Plant Biology Section, School of Integrative Plant Science, Cornell University, Ithaca,
NY, USA

Abstract: The unique reproductive structure of *Euphorbia* species, the cyathium, has historically

been considered an inflorescence despite its visual resemblance to a single complete flower.

However, multiple other models have recently been proposed, including the idea that the

cyathium is a flower or has some degree of floral identity. Enabled by the petty spurge

(*Euphorbia peplus* L.) genome and guided by the ABCDE model of floral development, we

dissected petty spurge cyathia and carried out a transcriptomic analysis of the different organs.

We also constructed gene phylogenies and performed dN/dS analysis on select floral genes.

The E class gene SEP3 was not upregulated in the involucre as compared with the cyathophyll,

indicating that the cyathium is not a complete flower. The filiform structures show upregulation

of B and E class genes and transcriptomic signatures of heterochromatin formation, consistent

with the hypothesis that they are reduced flowers. *E. peplus* homologs for the

inflorescence/floral meristem genes UFO and LFY and the B class genes AP3 and PI have

highly diverged sequences relative to other Euphorbiaceae species, suggesting that these

reproductive patterning genes may have evolved divergent functions. We propose a new model

in which the evolution of the cyathium involves two innovations: 1. altering LFY function to

influence the timing of floral meristem development to create a compact flower-like

inflorescence, and 2. altering the B class genes so that no perianth is formed. Our new

hypothesis includes testable predictions about protein interactions and gene sequence

evolution.

29
30

There are approximately 2000 species of *Euphorbia*, including horticulturally important plants

such as poinsettia (*Euphorbia pulcherrima* L). The reproductive structure of *Euphorbia* species

is called a cyathium. A cyathium looks similar to a complete flower with pistils, stamens, and a

perianth; however, it has historically been interpreted as a pseudanthium (false flower) that is a

condensed inflorescence consisting of a pistillate flower surrounded by staminate flowers and

an involucral bract (Figure 1).

Figure 1

Euphorbia peplus L., commonly known as petty spurge, is an emerging model species for

Euphorbia with a publicly available genome assembly (Czechowski et al., 2022; Johnson et al.,

2023). In *E. peplus*, each immature cyathium is enclosed within two bracts, called cyathophylls.

The cyathophyll contains the immature cyathium as well as multiple smaller cyathium primordia

(enclosed in their own cyathophylls). The number of primordia varies by *E. peplus* ecotype. The

cyathophylls open at approximately the stage of stigma lobe morphology which corresponds to

high stigma receptivity (Li et al., 2014), bringing the cyathium into the open for pollination and

later seed dispersal. *E. peplus* is primarily self-pollinated.

The *E. peplus* cyathium itself, as in other *Euphorbia* species, is composed of a single pistillate

flower in the center, surrounded by five mature staminate flowers. Each of the five mature

staminate flowers is subtended by immature or nonfunctional staminate flowers in a zigzag

pattern. Interspersed with the staminate flowers in the *E. peplus* cyathium are fringed filiform

(threadlike) structures. These filiform structures have been referred to by various terms

including "filamentous structures" (Prenner and Rudall, 2007) and "filiform bracteoles" (Bruyns,

2022). For the sake of clarity we will refer to these structures as "filiform structures'' in this

paper, because "filamentous" implies a developmental homology between these structures and

filaments, and because Prenner and Rudall 2007 also refer to cyathophylls as "bracteoles"

(Prenner and Rudall, 2007). The pistillate flower, staminate flowers, and filiform structures are

surrounded by a fused cuplike structure of involucral bracts, also known as the involucre. It is

this cup-shaped structure that gives cyathia their name, as "cyathium" comes from the Greek

word for cup, "kyathos". Crescent-shaped glands or nectaries are present near the top of the

involucre. In *E. peplus*, there are 4 or 5 nectaries present; this number varies by species.

Historically, the cyathium has been interpreted as an inflorescence for several reasons that are

well reviewed in Prenner and Rudall 2007 (Prenner and Rudall, 2007): the cyathium has a line

of constriction on the staminate flower that is interpreted as the flower-stamen boundary; other

genera in the Euphorbiaceae (e.g. *Anthostema*, Figure 2) have staminate flowers clustered

around a pistillate flower; and flowers are usually unisexual in the Euphorbiaceae (Kaplan and

Specht, 2022). Euphorbiaceae genera such as *Dalechampia* and *Pera* have reproductive

structures that have long been established to be pseudanthia (Gagliardi et al., 2018), and

morphological work on which Euphorbiaceae have pseudanthia is still ongoing, e.g. recent work

interpreting the "staminate flower" of *Ricinus communis* as a multi-flower unit(Claßen-Bockhoff

and Frankenhäuser, 2020). Prenner and Rudall 2007 speculate that if the cyathium is an

inflorescence, the filiform structures could be interpreted either as prophylls or as "highly

reduced male flowers that represent remnants of the original dichasial male inflorescence. This

would explain the variable number of [filiform] structures and the fact that they are not

associated with particular male flowers"(Prenner and Rudall, 2007). The involucre would then

be interpreted as a non-floral organ.

The Australian genus *Calycopeplus* and the New Caledonia tree *Neoguillauminia cleopatra*

have been proposed as "missing links" towards the evolution of the *Euphorbia* cyathium from

separate pistillate and staminate flowers; these close relatives to *Euphorbia* both have an

inflorescence where a pistillate flower with a small perianth is surrounded by staminate flowers

with small perianths (Figure 2) (Western Australian Herbarium et al., n.d.; Prenner and Rudall,

2007; Prenner et al., 2008; Dressler et al., 2014).

Figure 2

However, Prenner and Rudall posit that the case for *Euphorbia* as a pseudanthium is not clear-

cut: other Euphorbiaceae have abscission zones that are not at the base of an organ;

transitioning between an inflorescence arrangement like *Anthostema*'s and the inflorescence

arrangement of the *Euphorbia* cyathium would require a large change in the orientation of the

flowers, so a more complicated evolutionary history would be necessary to explain the cyathium

(Gilbert, 1994); and bisexual flowers are known to occur in ~15 genera in the Euphorbiaceae so

it is not implausible that they could occur in *Euphorbia(Prenner and Rudall, 2007)*. Subsequent

immunohistology assays showing expression of LEAFY (LFY), a gene associated with floral

identity, in cyathium primordia as well as individual floral primordia were interpreted as evidence

of the entire cyathium having some degree of floral identity (Prenner et al., 2010). If the

cyathium were to be interpreted as a complete flower, the pistillate flower could be interpreted

as a pistil and the staminate flowers could be interpreted as stamens. The filiform structures and

the involucre could be interpreted as the petals and a connately fused sepal respectively.

Now that a high-quality genome is publicly available for two *Euphorbia* species, the cyathium

can be investigated from a comparative molecular developmental perspective. This paper uses

transcriptomics and evolutionary analysis to shed more light on the controversial cyathium

structure and whether it is homologous to an inflorescence and/or a flower. Transcriptomics of

floral development of another species in the Euphorbiaceae, *Jatropha curcas* L., showed that

the ABCDE model of floral developmental regulation is mostly conserved in *J. curcas* (Hui et al.,

2017). Therefore we can make transcriptomic predictions in *Euphorbia* based on the ABCDE

model of floral development.

Under the ABCDE model, different combinations of MADS-box transcription factors combine to

specify the four floral whorls: the sepals, petals, stamens, and carpel (Figure 1E). E class

genes are present throughout the flower and help specify all four whorls. C class transcription

factors specify the carpel; B and C genes together specify stamens; B and A genes together

specify the petal; and A genes alone specify the sepal. D class genes specify ovules.

Generally, B and C class genes are conserved between taxa, while A class genes are not highly

conserved(Litt and Kramer, 2010). Complete flowers contain all of the floral whorls, while

unisexual flowers have either functional stamens or functional carpels but not both. Type I

unisexual flowers have sterile carpels/stamens, whereas Type II unisexual flowers, the type

posited to be present in the *Euphorbia* cyathium, do not have sterile carpels/stamens. The

disappearance of a reproductive whorl in Type II unisexual flowers is hypothesized to result

from differential regulation of the MADS-box transcription factors (Feng et al., 2020). Another

type of incomplete flowers, achlamydeous flowers, lack a perianth (sepals and petals). Loss of

the perianth whorls can result from edits to MADS-box transcription factors or to other

transcription factors not related to whorl identity, as in the arabidopsis *petal loss* (*ptl*)

mutant(Brewer et al., 2004).

Based on the work of Prenner and Rudall in *Euphorbia spp.* and the ABCDE model, we

examine two conflicting hypotheses: the cyathium-as-inflorescence model and the cyathium-as-

complete-flower model. Under the cyathium-as-inflorescence model, we expect to see E class

genes highly expressed in the pistillate and staminate flowers, potentially expressed in the

filiform structures if they constitute reduced flowers, and not expressed in the involucre as it

would not be a floral organ. There may or may not be transcriptomic signs of floral organs being

initiated that are later aborted (e.g. B class genes expressed in young pistillate flowers and/or

perianth-related genes expressed in pistillate or staminate flowers). Under the cyathium-as-

complete-flower model, we expect to see expression of E class genes in all of the other "whorls"

that would constitute part of the flower including both the filiform structures and involucre which

would constitute the "petals" and "sepals".

MATERIALS AND METHODS

Plant growth conditions—

E. peplus plants were grown under long day conditions with 25 °C day/16 °C night.   Seeds

were germinated without any pretreatment in closed Phytatrays (P5929; Sigma-Aldrich, Saint

Louis, MO, USA) in LM 1-1-1 soil mix, then week-old plants were uppotted to 4-inch pots

containing LM 1-1-1 soil mix.

Imaging—

Images of cyathia were taken using a Leica M205 FCA stereo microscope with a DMC6200

camera.

Fixation and dissection for RNA sequencing—

Cyathia from twelve-week-old *E. peplus* plants were collected into ice-cold acetone then stored

at -20°C until dissection. Cyathia were dissected under the dissecting scope in acetone on a

cold block and dissected organs were put immediately into chilled acetone tubes. 5 biological

replicates of 10 different organs were collected, with 10 cyathia dissected for each biological

replicate.

Stages used for RNA sequencing—

We used primordia approximately 200 µm tall for our "small primordium" samples. We used

cyathium primordia approximately 500 µm tall for our "large primordium" samples. We used

staminate flower primordia approximately 200 µm tall for our "young staminate flowers" and

pistillate flower primordia approximately 400 µm tall for our "young pistillate flowers". For the

remaining samples, we dissected them from nearly-mature cyathia still enclosed within

unopened cyathophylls when cyathia were approximately 1.5 mm tall not including cyathophylls:

nectary, involucre, mature staminate flower, mature pistillate flower, filiform structure. In our

organ stages, 'mature' here refers to developmental maturity of form, not reproductive maturity,

which roughly coincides with the opening of the cyathophylls.

RNA extraction—

RNA was extracted using the New England Biolabs Monarch Total RNA Miniprep Kit (NEB

#T2010, Ipswich, MA, USA). Acetone was removed from the tubes under the dissecting scope,

2-3 beads (BioSpec Products #11079124, Bartlesville, OK, USA) were added, the tubes were

placed into liquid nitrogen, and the tissue was ground in a bead beater (Qiagen TissueLyser II,

Germantown, MD, USA). 400uL of 1x protection reagent from the kit was added to the ground

tissue and beads, and the tissue was ground again in the protection reagent. The liquid was

transferred to a new tube and the rest of the kit protocol was followed. RNA was eluted with

100uL nuclease-free water.

RNA purification—

The 100uL of RNA was purified by adding 10uL 3M sodium acetate, 0.75uL GlycoBlue

(ThermoFisher #AM9515, Waltham, MA USA) and 275uL ice-cold 100% ethanol, vortexing to

mix, precipitating at -80°C overnight, centrifuging at 13000 rpm at 4°C for 30 minutes, washing

the pellet twice with 0.5mL ice-cold 75% ethanol, air-drying the pellet for 5 minutes, and

resuspending the RNA in 20uL of nuclease-free water.

3' sequencing—

RNA samples were submitted to the Cornell Institute for Biotechnology Genomics Center for 3'

mRNA-seq library construction using a QuantSeq FWD kit (Lexogen, Vienna, Austria) followed

by 1x75 bp sequencing on an Illumina NextSeq500 (Illumina, Hayward, CA, USA). Sequencing

statistics were calculated with v0.15.0 seqkit stats with parameters -j 10 -T(Shen et al., 2016).

(Supplementary File 1).

Orthogroup inference—

To assemble orthogroups, we ran OrthoFinder v2.5.4 (Emms and Kelly, 2019) on the longest

isoform protein sequences from *E. peplus*, *E. lathyris*, arabidopsis, soybean, Pará rubber tree,

cassava, poplar, and castor bean using the parameters -S diamond -t 40 -a 8(Buchfink et al.,

2021). Orthogroups are included in Supplementary File 2.

Identification of ABCDE floral identity genes—

In order to identify floral identity genes of interest, we retrieved the sequences of genes of

interest from Uniprot accessed through TAIR and ran blastp with our *E. peplus* proteins file as

the query and the parameters -qcov_hsp_perc 80 e-value 1e-10(Camacho et al., 2009). Where

BLAST results surfaced multiple candidates with low e-values and high percent identity, a

reciprocal BLAST to arabidopsis was used to select the best candidate(s). We further

confirmed the results through comparison to the clades identified by OrthoFinder. Gene IDs

accessed through TAIR: AT1G69120(AP1)(Bowman et al., 1993); AT4G36920(AP2)(Jofuku et

al., 1994); AT3G54340(AP3)(Jack et al., 1992); AT4G18960(AG)(Mizukami and Ma, 1997);

AT2G45650(AGL6)(Ma et al., 1991); AT1G69180(CRC)(Alvarez and Smyth, 1999);

AT1G65480(FT)(Kobayashi et al., 1999); AT5G60910(AGL8)(Mandel and Yanofsky, 1995);

AT5G61850(LFY)(Schultz and Haughn, 1991); AT5G20240(PI)(Goto and Meyerowitz, 1994);

AT5G15800(SEP1)(Pelaz et al., 2000); AT1G24260(SEP3)(Pelaz et al., 2000);

AT3G58780(SHP1)(Liljegren et al., 2000); AT2G42830(SHP2)(Liljegren et al., 2000);

AT2G45660(AGL20)(Lee et al., 2000); AT4G09960(STK)(Favaro et al., 2003);

AT5G03840(TFL1)(Kobayashi et al., 1999); AT1G30950(UFO)(Levin and Meyerowitz, 1995).

Transcriptomic analysis—

Raw read files were filtered with fastp v0.23.2 using parameters --cut_right --cut_window_size 5

--cut_mean_quality 20 --length_required 50(Chen et al., 2018). Transcript abundance was

generated using *E. peplus* coding sequences with kallisto quant v0.42.4 with parameters --

single -b 100 -t 10(Bray et al., 2016). The resulting data was processed in R using the DESeq2

pipeline; all genes with >20 estimated counts across all samples were considered to be

expressed and were included in the analysis (Love et al., 2014). Variance-stabilized-

transformed data were visualized in heatmaps using the package pheatmap with expression

normalized by row and genes clustered by expression pattern(Kolde, n.d.). Of the floral genes

of interest investigated (Supplementary File 3), the *E. peplus* homologs for AGL8/FUL

(Ep_chr6_g18041), TFL1 (Ep_chr7_g23231) and two paralogs of FT (Ep_chr8_g25489,

Ep_chr6_g20489) had expression too low (i.e. 20 or fewer counts across all samples) to be

analyzed. Genes are said to be differentially expressed in this paper if padj<0.05. All DESeq

filtered untransformed data is included in Supplementary File 4. Comparisons of differentially

expressed genes between different tissues are included in Supplementary File 5.

Transcriptomic comparison between orthologs in arabidopsis and **E. peplus**—

Klepikova floral atlas data for Arabidopsis was downloaded through NCBI PRJNA314076 for the

mature sepal, mature anther, mature filament, mature petal, mature carpel, and leaf samples

(Klepikova et al., 2015). The raw reads were processed in the fastp/kallisto/DESeq2 pipeline

using the same parameters as for the *E. peplus* data. For each orthogroup containing an *E.*

peplus gene that was upregulated in the cyathium whorl, we asked whether there were one or

more arabidopsis genes in that orthogroup that were upregulated in the arabidopsis flower whorl

corresponding to the cyathium whorl.

Biological Process and Molecular Function GO term enrichment analysis—

GO terms were assigned to *E. peplus* genes as specified in Johnson et al 2023 (Johnson et al.,

2023); GO terms assigned to each gene are included in Supplementary File 6. In order to find

which GO terms were enriched in differentially expressed genes in different organs as

compared with cyathophylls, we ran TopGO v.2.46.0 using the "classicCount" option, using

Fisher's test to determine significance(Alexa and Rahnenfuhrer, 2021). TopGO results are

included in Supplementary File 7. We summarized and visualized the GO results using

REVIGO and downloaded the R scripts to edit the color scale and axis limits (Supek et al.,

2011). Terms with a dispensability higher than 0.9 were removed and clustered with other

terms for ease of visualization. Only terms with a dispensability less than 0.2 were labeled with

text in the visualization.

Manual curation of OrthoFinder clades—

Genes with highly truncated annotations were removed from the analysis (LFY: rubber tree

KAF2311631.1, soybean Glyma.04G202002.1.p; PHYA: soybean Glyma.03G227333.1.p; PI: *E.*

lathyris Casp05761.t1; SEP1: rubber tree KAF2325472.1). For PI, *E. lathyris* Casp33119.t1 and

E. peplus Ep_chr8_g25689.t1 were included in the PI phylogenetic analysis despite being

classified into their own orthogroup because they clearly were a result of a tandem duplication

(Casp33120.1 and Ep_chr8_g25690.t1 were included in the orthogroup with Arabidopsis PI,

AT5G20240.1) and because of high reciprocal BLAST scores with Arabidopsis PI.

Gene alignment, gene tree construction and selection analysis—

For each clade of interest (generated by OrthoFinder then manually curated as specified above;

see Supplementary File 3 for gene names), a preliminary alignment was generated using

MAFFT v7.522 with parameters --maxiterate 5000 --auto --adjustdirectionaccurately --thread 8 --

op 3. Then macse v2.07 alignSequences was used to generate a codon-aware alignment with

default parameters. For a few alignments, macse generated exclamation points due to

sequencer errors; these were converted to N. Then alignment trimming was performed in trimAl

v1.4 with parameter -gappyout. Finally, sequences were exported for phylogeny construction

using macse v2.07 exportAlignment -codonForFinalStop NNN -codonForInternalStop NNN -

keep_gap_only_sites_ON. To generate phylogenetic trees, RAxML NG v1.2.0 was run with

parameters --all --model GTR+G --threads 12 --bs-trees 1000(Kozlov et al., 2019). In order to

generate dN/dS values for each branch of the tree, PAML v4.10.6 was run on the best tree from

RAxML with model=1 and NSSites=0. In order to test for relaxed selection at sites within

genes, PAML v4.10.6 was run on the best tree from RAxML with model=2 and NSSites=2.

Alignments were visualized in ESPript 3.0 online (https://espript.ibcp.fr/ESPript/cgi-

bin/ESPript.cgi)(Robert and Gouet, 2014). Sites flagged for positive selection based on the

Bayes empirical Bayes method (BEB) are included in Supplementary File 8.

Motif analysis—

In order to retrieve a dataset that could be used to detect the canonical LFY binding motif, a bed

file of LFY ChIPseq targets (ChIPseq MACS2 q-value 10-10, DANPOS Mnase-seq q-value 10-

50) was taken from Supplementary Data 1 from Jin et al 2021(Jin et al., 2021). This was

converted to a fasta file of the target sequences using bedtools getfasta(Quinlan and Hall,

2010). Then MEME-ChIP from MEME v5.5.2 was used to detect the LFY canonical binding

motif using default parameters except -maxw 25(Machanick and Bailey, 2011). The dLUBS and

mLUBS motifs were acquired from the authors of Rieu et al 2023 in pfm format and converted to

meme format using jaspar2meme -pfm from MEME v4.11.2(Bailey et al., 2015; Rieu et al.,

2023b). Then FIMO from MEME v5.5.2 was run with default parameters on a fasta file of the

3kb upstream of the start of the LFY target genes AP1, AG, AP3, and PI in order to assess

whether the motifs were present or not(Grant et al., 2011). MEME FIMO motif search results

are included in Supplementary File 9.

Genome assemblies used for analysis—

The Johnson et al *Euphorbia peplus* genome assembly and annotation was accessed through

Zenodo(Johnson et al., 2023): https://zenodo.org/records/8033914 The Wang et al. *Euphorbia*

lathyris assembly and annotation was accessed through figshare(Wang et al., 2021):

https://figshare.com/articles/dataset/High-

quality_genome_assembly_of_the_biodiesel_plant_Euphorbia_lathyris/14909913/1 The Liu et

al. *Hevea brasiliensis* (rubber tree) assembly and annotation was accessed through NCBI(Liu et

al., 2020): https://www.ncbi.nlm.nih.gov/assembly/GCA_010458925.1/. The Xu et al. wild

Ricinus communis assembly and annotation was accessed through NCBI (Xu et al., 2021):

https://www.ncbi.nlm.nih.gov/datasets/genome/GCF_019578655.1/. The Bredeson et al.

Manihot esculenta v8.1 assembly and annotation was accessed through Phytozome:

https://phytozome-next.jgi.doe.gov/info/Mesculenta_v8_1 The arabidopsis Araport 11 genome

release was accessed through TAIR(Cheng et al., 2017):

https://www.arabidopsis.org/download/index-

auto.jsp?dir=%2Fdownload_files%2FGenes%2FAraport11_genome_release%2FAraport11_bla

stsets. The *Glycine max* (soybean) Wm82.a4.v1 assembly and annotation was accessed

through Phytozome(Valliyodan et al., 2019): https://phytozome-

next.jgi.doe.gov/info/Gmax_Wm82_a4_v1. The *Populus trichocarpa* v4.1 assembly and

annotation was accessed through Phytozome(Tuskan et al., 2006): https://phytozome-

next.jgi.doe.gov/info/Ptrichocarpa_v4_1

RESULTS

Analysis of ABCDE gene expression in cyathium components—

We dissected the different organs of the cyathium and sequenced their RNA (Figure 3 A-D). Of

25,473 annotated *E. peplus* genes, 17,164 were expressed in the cyathium. Staminate flowers

had the most distinct expression pattern from other organs; a clear separation is visible between

the mature staminate flowers, young staminate flowers, and other cyathium organs in the first

principal component (PC) of the data (Figure 3E). This data is consistent with many other

flowering plant species' transcriptomic pattern of distinct staminate flowers compared with other

organs, or distinct stamens compared with other floral organs in complete flowers (Klepikova et

al., 2015; Xanthopoulou et al., 2021).

Figure 3

Using BLAST and OrthoFinder, we identified *E. peplus* homologs for all classes of the ABCDE

genes and visualized their expression pattern across different organs (Figure 4).

Figure 4

In the ABCDE model, the E class genes SEPALLATA1 (SEP1) and SEPALLATA3 (SEP3) are

partially redundant in function and help specify all four floral whorls; a closely related E class

gene, AGAMOUS-LIKE 6 (AGL6), also helps specify all of the floral organs and is also involved

floral meristem regulation (Dreni and Zhang, 2016; Morel et al., 2019). There are three

orthologs of SEP1 in *Euphorbia*, which is consistent with other Euphorbiaceae (Supplementary

Figure 1). *E. peplus* SEP1_1 is significantly upregulated in both stages of pistillate flowers

compared with cyathophylls, SEP1_2 is significantly upregulated in staminate flowers as

compared with pistillate flowers, and SEP 1_3 is significantly upregulated in the filiform

structures as compared with cyathophylls. *Euphorbia*'s sole SEP3 ortholog is significantly

upregulated in both stages of the pistillate and staminate flowers, both stages of the primordia,

and the filiform structures as compared with cyathophylls. AGL6 is not significantly expressed

in any tissue as compared with the cyathophylls due to low expression overall.

The B class genes APETALA3 (AP3) and PISTILLATA (PI) specify petal and stamen identity;

this role is conserved across bisexual and unisexual flowers (Zhang et al., 2022). All three *E.*

peplus AP3 paralogs and both PI paralogs are significantly upregulated in staminate flowers

compared with pistillate flowers at both stages studied, in mature staminate flowers as

compared with cyathophylls, in young staminate flowers as compared with cyathophylls (except

AP3_3), and in filiform structures as compared with cyathophylls. Of the AP3 and PI homologs,

only AP3_2 is significantly expressed in the involucres as compared with the cyathophylls.

The C class gene AGAMOUS (AG) is deeply conserved across evolution and helps specify both

pistillate and staminate development (Dreni and Kater, 2014). The *E. peplus* ortholog of AG is

significantly upregulated in both stages of both pistillate flowers and staminate flowers as

compared with cyathophylls, and not significantly upregulated in any other parts of the cyathium.

In arabidopsis, the A class genes APETALA1 (AP1) and APETALA2 (AP2) help specify sepals

and help specify petals in conjunction with B class genes, in addition to other functions in floral

meristem identity and determinacy; however these genes' functions are not conserved outside

the Brassicaceae and are unlikely to be the same in *Euphorbia* as arabidopsis (Litt and Kramer,

2010; Zumajo-Cardona and Pabón-Mora, 2016). There are two homologs of arabidopsis AP1 in

E. peplus. AP1_1, which is most similar in sequence to arabidopsis AP1 and CAULIFLOWER,

is upregulated in the nectary, involucre, filiform structures, both stages of primordia, both stages

of staminate flowers, and young pistillate flowers as compared with cyathophylls. *E. peplus*

AP1_2 and AP2 are not significantly differentially expressed in any cyathium part as compared

with cyathophylls, likely due to low gene expression.

We were also interested in the genes LEAFY (LFY) and UNUSUAL FLORAL ORGANS (UFO)

which are known to directly activate many of the ABCDE genes. LFY is a transcription factor

that is a key regulator of floral meristem and floral organ identity(Moyroud et al., 2010); it opens

up the chromatin around the ABCDE genes, enabling their transcription(Jin et al., 2021). In

arabidopsis, UFO interacts with LFY to activate the B class genes AP3 and PI as well as other

targets (Rieu et al., 2023b). *E. peplus* has two homologs of LFY: LFY1_1 is significantly

upregulated in both stages of primordia as compared with cyathophylls, and LFY_2 is

significantly upregulated in small primordia and filiform structures as compared with

cyathophylls. *E. peplus* UFO is upregulated in both stages of primordia, both stages of

staminate flowers, young pistillate flowers, and in filiform structures, as compared with

cyathophylls. It is significantly upregulated in mature staminate flowers as compared with

mature pistillate flowers.

The gene CRABS CLAW (CRC), which is necessary for nectary development in both

arabidopsis and petunia (Morel et al., 2018; Slavković et al., 2021), is not upregulated in *E.*

peplus nectaries as compared with cyathophylls, suggesting an alternative nectary development

pathway in *E. peplus*. It is significantly upregulated in both stages of pistillate flowers as well as

in filiform structures compared with cyathophylls, and significantly upregulated in both stages of

pistillate flowers compared with staminate flowers. The upregulation of CRC in pistillate flowers

in *E. peplus* is consistent with CRC's other functions in carpel development and meristem

termination and with results from *J. curcas*, where CRC is highly upregulated during

developmental stages of pistillate flowers(Xu et al., 2016).

Transcriptomic analysis of differentially expressed genes in cyathium components—

Further investigating whether the *E. peplus* cyathium could be homologous to a complete

flower, we asked whether the orthogroups containing at least one gene significantly upregulated

in *E. peplus* pistillate flowers as compared with cyathophylls also contained at least one gene

significantly upregulated in the arabidopsis carpels compared with leaves; whether orthogroups

with *E. peplus* genes upregulated in staminate flowers also contained arabidopsis genes

upregulated in stamens; whether orthogroups with *E. peplus* genes upregulated in filiform

structures also contained arabidopsis genes upregulated in petals; and whether orthogroups

with *E. peplus* genes upregulated in involucres also contained arabidopsis genes upregulated in

sepals.

Figure 5

We found that 60% of genes significantly upregulated in *E. peplus* pistillate flowers had an

ortholog significantly upregulated in arabidopsis carpels, and 65% of genes significantly

upregulated in staminate flowers had an ortholog significantly upregulated in arabidopsis

stamens. The filiform structures and involucre both had a much lower level of conservation of

expression with arabidopsis floral whorls. 38% of genes significantly upregulated in *E. peplus*

filiform structures had an ortholog significantly upregulated in arabidopsis petals, and 38% of

genes significantly upregulated in involucres had an ortholog significantly upregulated in

arabidopsis sepals.

Because of the lack of shared upregulated orthogroups between involucre and filiform structures

with sepal and petal whorls, respectively, and the low level of expression of CRABS CLAW in

the nectary, we wanted to investigate what made these parts of the cyathium unique. We

performed a gene ontology (GO) term analysis on the genes significantly upregulated in mature

staminate flowers, mature pistillate flowers, young staminate flowers, young pistillate flowers,

involucres, filiform structures, and nectaries as compared with cyathophylls (Supplementary File

7). We visualized the result for nectaries, involucres, and filiform structures based on semantic

similarity of the GO terms (Figure 6).

Figure 6

We found that genes upregulated in the nectary as compared with cyathophylls were

significantly enriched for multiple Biological Process GO terms relating to transmembrane

transport and specialized metabolism. GO terms relating to gene silencing were also enriched.

The Molecular Function GO terms enriched in the nectary included multiple significant terms

relating to transmembrane transporter activity and glycosyltransferase activity, similarly to the

Biological Process GO terms. ATP and NADH metabolism-related terms were also significantly

enriched in the nectaries, suggesting an enrichment in energy-generating redox activities.

Genes upregulated in involucres as compared with cyathophylls were also significantly enriched

in genes with Biological Process GO terms relating to specialized metabolite biosynthesis and

transmembrane transport. This could potentially indicate that some of the compounds present

in the nectaries are at least partially biosynthesized in the involucre tissue connected to the

nectaries. The involucre also had several significant GO terms relating to stress, immune

responses, and defense responses. Some of these may be related to cytokinin levels; the gene

Ep_chr4_g10976, which is upregulated in involucres, is in an orthogroup with

AT1G31770(ABCG14), a transmembrane cytokinin transporter. Cytokinin is involved in

responses to osmotic stress, salt stress, and other environmental factors but also regulates

developmental processes such as inflorescence branching in *J. curcas*(Chen et al., 2019).

The involucre-upregulated gene with the highest p-value, Ep_chr5_g16200, is in an orthogroup

with AT5G03680, PETAL LOSS (PTL), which generally represses tissue growth by limiting cell

division. (Despite its name, PTL does not regulate petal organ identity, and its expression is not

limited to petals– *ptl* mutants' lack of petals may be due to indirect effects of tissue overgrowth

disrupting auxin maxima (O'Brien et al., 2015; Takeda et al., 2022).) This gene is upregulated

in filiform structures, young pistillate flowers, mature pistillate flowers, nectaries, and both

stages of primordia as compared with cyathophylls, suggesting functions in patterning across

multiple cyathium components.

For the genes upregulated in filiform structures as compared with cyathophylls, stamen

development Biological Process GO terms were significantly enriched. Heterochromatin

formation, gene silencing, and DNA methylation-related terms were also all significantly

enriched.

The mature staminate flowers had multiple significant Biological Process GO terms for stamen

identity and morphogenesis as well as ATP and NADPH related terms. The young staminate

flowers had multiple significant GO terms associated with stamen and pollen development but

did not have significant ATP or NADPH related terms. The mature pistillate flowers had two GO

terms associated with petal development; the pistillate flower-enriched gene causing this

significant GO hit, Ep_chr1_g00250, is in an orthogroup with arabidopsis AT4G16850,

KronbladStorleK (KSK; the name is Swedish for petal size) which influences auxin responses

during petal development(Miller et al., 2020). Compared with the mature pistillate flower, the

younger pistillate flower was enriched for more GO terms relating to reproductive structure

development, suggesting that the patterning of the pistillate flower is more active in this earlier

stage. Both stages of both the pistillate and staminate flowers had significant GO terms relating

to gene silencing; the top five most significant mature pistillate flower terms were all related to

gene silencing.

The zinc finger motif-containing gene Ep_chr4_g09148 was in the top 10 most significant

upregulated genes in the involucre, nectary, and filiform structures, and was significantly

upregulated in every other cyathium tissue studied compared with the involucre. This gene

shares an orthogroup with AT2G33835, FRIGIDA-ESSENTIAL 1 (FES1) which is part of a

complex that represses flowering in arabidopsis(Schmitz et al., 2005).

Evolutionary analysis of ABCDE floral identity gene sequences—

Changes in MADS-box binding partners are a driver of floral morphological diversity across

many plant families(Bartlett, 2017). For example, duplication and neofunctionalization of SEP

genes is involved in the generation of pseudanthia in the Asteraceae(Zhang et al., 2017).

Therefore, in addition to analyzing gene expression, we analyzed gene sequence evolution in

order to see whether there were changes in *Euphorbia* gene sequence and/or copy number that

might reflect changes in binding and function. We performed a codon-aware alignment, tree

construction, and dN/dS analysis using the gene sequences from *E. peplus*, *Euphorbia lathyris*,

several other Euphorbiaceae species, and the model systems arabidopsis, soybean, and poplar

(Figure 7, Supplementary Figure 1, Supplementary Figure 2, Supplementary Figure 3).

Figure 7

LFY, UFO, AP3, and PI show high sequence divergence in *Euphorbia* (Figure 7A). Compared

with the non-floral gene PHYTOCHROME A (PHYA) and with the C class gene AG, the genes

LFY, AP3, and PI have high dN/dS on their *Euphorbia* branches. The *E. peplus* and *E. lathyris*

homologs for LFY have an amino acid change from a positive amino acid to a hydrophobic

amino acid in a DNA-binding site that is otherwise deeply conserved in vascular plant lineages

dating back to the lycophytes (Figure 7B) (Rodríguez-Pelayo et al., 2022). The *E. peplus* and *E.*

lathyris homologs for UFO also differ greatly in sequence from other Euphorbiaceae homologs

of UFO, and one reason is that the *Euphorbia* versions have deletions near the start of the

protein (Figure 7C).   Interestingly, many of the AP3 residues flagged as "highly significant" for

positive selection in *Euphorbia* are in the K domain in a region involved in tetramer formation(Lai

et al., 2019), perhaps indicating a change in binding partner (Supplementary Figure 3).

Because LFY and UFO showed a high level of sequence divergence, in order to investigate

whether LFY and UFO binding targets were conserved, we looked for LFY binding motifs and

LFY-UFO binding motifs in the putative promoters of homologs of the LFY targets AP3, PI, AG,

and AP1 in *E. peplus*, *H. brasiliensis*, and arabidopsis. (AP3 and PI are targets of the LFY-UFO

complex as well.) We hypothesized that the binding motifs might not be conserved in *E. peplus*.

Both canonical LFY binding motifs and UFO-LFY motifs (mLUBS and dLUBS) are present in the

3kb upstream of these target genes in *E. peplus* (Supplementary File 9), suggesting that at least

some of the binding motifs of LFY and LFY-UFO are conserved despite the high divergence in

sequence.

DISCUSSION

Across angiosperm evolution, the E class gene SEP3 is generally highly conserved in terms of

sequence and expression pattern and usually helps define floral whorls, whereas SEP1

orthologs have been repurposed in different species to regulate bract identity and inflorescence

architecture(Dreni and Ferrándiz, 2022). The fact that *Euphorbia*'s SEP3 ortholog is not

upregulated in the involucre compared with cyathophylls is inconsistent with the cyathium-as-

complete-flower hypothesis.

In the filiform structures, we observed significant expression of B-class and E-class genes, as

well as one homolog of LFY, suggesting floral identity. The filiform structures had low levels of

RNA overall compared with other structures in the cyathium (Supplementary File 1), which could

be explained by the gene ontology analysis showing significant GO terms relating to

heterochromatin formation. Our data therefore agrees with the interpretation of the filiform

structures as reduced flowers in an inflorescence rather than as petals in a complete flower.

Overall, our GO term analysis showed that gene silencing seems to be involved in the creation

of multiple cyathium organs.

We did not see transcriptomic signs of abortion of other whorls in the pistillate and staminate

flowers (e.g. hypothetically, seeing B class expression in the young pistillate flower could have

been an example of that), besides the upregulation of the putative homolog of the arabidopsis

perianth gene KSK in the pistillate flower. Organ abortion can happen at many different stages,

however, and unisexual flowers do not always show evidence of their missing whorls(Mitchell

and Diggle, 2005). The C class gene AGAMOUS is highly expressed in both pistillate flowers

and staminate flowers, as would be expected under both of our hypotheses/models.

Euphorbia LFY, UFO, and the B-class genes AP3 and PI show signs of divergence based on

phylogenetic distance and relaxed selection. Except for UFO (which may have lacked signs of

relaxed selection due to the gene tree topology), these genes also had multiple sites that were

classified as highly significant for positive selection in Bayes Empirical Bayes (BEB) analysis

(Supplementary File 8).

LFY and UFO are critically important in specifying floral meristem identity(Rieu et al., 2023a).

Many of the originally described loss-of-function or dominant negative *ufo* and *lfy* mutants in

arabidopsis had an inflorescence subtended by a bract in place of a flower(Weigel et al., 1992;

Levin and Meyerowitz, 1995). UFO mutants in other species also have an inflorescence in

place of a flower, e.g. *anantha* mutants in tomato(Lippman et al., 2008). In the Euphorbiaceae,

the LFY homolog in *J. curcas* promotes the transition from inflorescence to floral meristem

identity. Overexpression of JcLFY causes early flowering, and its co-suppression delays flower

formation, causes more inflorescence branches, and causes sepal-like organs(Tang et al.,

2016). As compared with its relatives such as *Anthostema* (Figure 2), the *Euphorbia*

inflorescence is very compact structurally. Cyathia have a reduced axis, different placement of

their flowers, and an earlier emergence of pistillate flower primordia compared with staminate

flower primordia than inflorescences of related genera (Gagliardi et al., 2018). These

differences could not be explained by differential elongation of organs. This suggests that a

difference in timing or spatial location of the specification and/or termination of the floral

meristems led to the cyathium structure. The drastic changes in sequence of LFY and UFO in

Euphorbia could contribute to those functional changes in floral meristem specification.

One interesting side note on the expression of UFO is that in *J. curcas*, UFO is downregulated

in both mature staminate flowers and mature pistillate flowers compared to younger

stages(Chen et al., 2016), whereas in our *E. peplus* dataset we saw the highest expression of

UFO in mature staminate flowers. This either suggests a repurposing of UFO, or that

developmentally mature but not reproductively mature (i.e. not elongated) staminate flowers

aren't yet mature transcriptomically compared with pistillate flowers. As UFO functions in the

development of individual perianth organs as well as in the broader activation of the B class

genes(Takeda et al., 2022), some of the changes in UFO expression and sequence in

Euphorbia could be related to the suppression of perianth development in addition to potential

functions in defining the floral meristem.

Genes in the AP1 clade can also be involved in floral meristem specification– and are also

highly diverged in *Euphorbia*. In arabidopsis, AGL79 interacts with SUPPRESSOR OF

OVEREXPRESSION OF CONSTANS1 (SOC1) to regulate the expression of TERMINAL

FLOWER 1 (TFL1) which controls inflorescence meristem identity (Yang et al., 2023).

Euphorbia appears to be "missing" its AP1-CAL homolog in the Euphorbiaceae AGL79 clade in

our gene tree because that gene, AP1_2, is so diverged that it appears elsewhere in the tree

(Supplementary Figure 2).

Based on our results, we hypothesize that the cyathium is indeed an inflorescence, not a

complete flower, and that there are two key genetic innovations that led to the cyathium: 1.

Alteration of LFY function to change the timing of floral meristem specification and floral

development to create a compact inflorescence; and 2. Changing the transcription factor binding

partners of AP3 and PI so that they are no longer involved in perianth formation, deleting the

structure of the perianth.

The modification of AP3 and PI to change their binding partners and function in order to delete

the perianth would constitute a destruction of the system where overlapping ABC genes create

four whorls of a flower; in that case, Euphorbia would be similar to gymnosperms where B and

C class expression together specifies pollenate organs and C class alone specifies ovulate

organs but there is no perianth (Figure 8)(Irish, 2017).   Hypothetically, because the B class

proteins would no longer be required to work together with A class members, the B class genes

would be freed to take on new functions, which could help explain some of the high diversity in

cyathium form in *Euphorbia*.

Figure 8

Our hypothesis could be consistent with Prenner and Rudall's interpretation of the Australian

genus *Calycopeplus* and the New Caledonia tree *Neoguillauminia cleopatra* as "missing links"

towards the *Euphorbia* cyathium. Because *Calycopeplus* and *Neoguillauminia* have a similarly

compact inflorescence structure to that of *Euphorbia* (compared with e.g. *Anthostema*) but still

have perianths, we hypothesize that a step-wise evolutionary process occurred where an

inflorescence-disrupting event (i.e. disruption of LFY function) happened in the common

ancestor of *Calycopeplus*, *Neoguillauminia,* and *Euphorbia* but the alteration of B class genes to

delete the perianth happened only in *Euphorbia*. Future studies could sequence the floral

genes of *Calycopeplus*, *Neoguillauminia*, and *Anthostema* to test this hypothesis.

In addition to expanding our analyses to more taxa, future studies could test our model through

experimental investigations of the function of the UFO-LFY complex and its DNA binding targets

in *Euphorbia*, e.g. through electrophoretic mobility shift assays (EMSA), microscale

thermophoresis (MST), or DAP-seq. Additionally, yeast two-hybrid or similar techniques could

be used to check the transcription factor binding partners of *Euphorbia*'s B-class genes. It

would also be interesting to use ATAC-seq or another method to determine chromatin

accessibility across the genome and speculate on what might be being silenced in the different

cyathium structures. *In situ* hybridization, if optimized in this system, could also potentially

reveal more spatially and temporally resolved patterns of gene expression. We are excited to

see what future work reveals about the developmental genetics involved in creating the complex

structure of the cyathium.

ACKNOWLEDGEMENTS

Thank you to Jacob Landis for extremely helpful advice on running PAML. A warm thank you to

François Parcy, Jérémy Lucas and Laura Turchi for sending the motif files for mLUBS and

dLUBS. Thank you to the Cornell Weill growth chamber plant maintenance staff, especially

Corbin Lay, for their excellent upkeep of the plants used. Thank you to the Frank Lab for helpful

comments on the manuscript. This work was funded by a Cornell Institute of Biotechnology

Seed Grant, National Science Foundation Integrative Organismal Systems (NSF IOS) award

1942437, and startup funds from the College of Agriculture and Life Sciences at Cornell

University to Margaret Frank.

AUTHOR CONTRIBUTIONS

A.R.J., A.B., and M.H.F. all conceived of the study. Under the supervision of A.R.J., A.B.

collected the images of cyathia and dissected cyathia components, and performed preliminary

dissections and RNA extractions that contributed to the experimental design for the RNAseq

experiment. A.R.J. performed the dissection for the RNAseq experiment. A.B. and A.R.J. both

contributed to the RNA extraction for the RNAseq experiment. A.B. and A.R.J. collaborated on

the initial sequence alignment and phylogeny construction to assess conservation of sequence

similarity in floral genes. A.R.J. performed all other data analysis. A.R.J. wrote the first draft of

the manuscript, after which all authors contributed to reviewing and editing the manuscript.

DATA AVAILABILITY

The raw RNAseq data generated in this study are available from NCBI GenBank under the

accession PRJNA1063728. All code used in the data analysis is available on Github at

https://github.com/ariellerjohnson/Cyathium-2024.

LITERATURE CITED

Alexa, A., and J. Rahnenfuhrer. 2021. topGO: enrichment analysis for gene ontology. *R*

package version 2.46: 2021.

Alvarez, J., and D. R. Smyth. 1999. CRABS CLAW and SPATULA, two Arabidopsis genes that

control carpel development in parallel with AGAMOUS. *Development* 126: 2377–2386.

Bailey, T. L., J. Johnson, C. E. Grant, and W. S. Noble. 2015. The MEME Suite. *Nucleic acids*

research 43: W39–49.

Bartlett, M. E. 2017. Changing MADS-Box Transcription Factor Protein-Protein Interactions as a

Mechanism for Generating Floral Morphological Diversity. *Integrative and comparative*

biology 57: 1312–1321.

Bowman, J. L., J. Alvarez, D. Weigel, E. M. Meyerowitz, and D. R. Smyth. 1993. Control of

flower development in Arabidopsis thaliana by APETALA1 and interacting genes.

Development 119: 721–743.

Bray, N. L., H. Pimentel, P. Melsted, and L. Pachter. 2016. Near-optimal probabilistic RNA-seq

quantification. *Nature biotechnology* 34: 525–527.

Brewer, P. B., P. A. Howles, K. Dorian, M. E. Griffith, T. Ishida, R. N. Kaplan-Levy, A. Kilinc, and

D. R. Smyth. 2004. PETAL LOSS, a trihelix transcription factor gene, regulates perianth

architecture in the Arabidopsis flower. *Development* 131: 4035–4045.

Bruyns, P. V. 2022. Euphorbia in Southern Africa: Volume 1. Springer Nature.

Buchfink, B., K. Reuter, and H.-G. Drost. 2021. Sensitive protein alignments at tree-of-life scale

using DIAMOND. *Nature methods* 18: 366–368.

Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos, K. Bealer, and T. L. Madden.

2009. BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421.

Cheng, C.-Y., V. Krishnakumar, A. P. Chan, F. Thibaud-Nissen, S. Schobel, and C. D. Town.

2017. Araport11: a complete reannotation of the Arabidopsis thaliana reference genome.

The Plant journal: for cell and molecular biology 89: 789–804.

Chen, M.-S., B.-Z. Pan, Q. Fu, Y.-B. Tao, J. Martínez-Herrera, L. Niu, J. Ni, et al. 2016.

Comparative Transcriptome Analysis between Gynoecious and Monoecious Plants

Identifies Regulatory Networks Controlling Sex Determination in Jatropha curcas. *Frontiers*

in plant science 7: 1953.

Chen, M.-S., M.-L. Zhao, G.-J. Wang, H.-Y. He, X. Bai, B.-Z. Pan, Q.-T. Fu, et al. 2019.

Transcriptome analysis of two inflorescence branching mutants reveals cytokinin is an

important regulator in controlling inflorescence architecture in the woody plant Jatropha

curcas. *BMC plant biology* 19: 468.

Chen, S., Y. Zhou, Y. Chen, and J. Gu. 2018. fastp: an ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* 34: i884–i890.

Claßen-Bockhoff, R., and H. Frankenhäuser. 2020. The 'Male Flower' of Ricinus communis

(Euphorbiaceae) Interpreted as a Multi-Flowered Unit. *Frontiers in cell and developmental*

biology 8: 313.

Czechowski, T., E. Forestier, S. H. Swamidatta, A. D. Gilday, A. Cording, T. R. Larson, D.

Harvey, et al. 2022. Gene discovery and virus-induced gene silencing reveal branched

pathways to major classes of bioactive diterpenoids in Euphorbia peplus. *Proceedings of*

the National Academy of Sciences of the United States of America 119: e2203890119.

Dreni, L., and C. Ferrándiz. 2022. Tracing the Evolution of the SEPALLATA Subfamily across

Angiosperms Associated with Neo- and Sub-Functionalization for Reproductive and

Agronomically Relevant Traits. *Plants* 11.

Dreni, L., and M. M. Kater. 2014. MADS reloaded: evolution of the AGAMOUS subfamily genes.

The New phytologist 201: 717–732.

Dreni, L., and D. Zhang. 2016. Flower development: the evolutionary history and functions of

the AGL6 subfamily MADS-box genes. *Journal of experimental botany* 67: 1625–1638.

Dressler, S., M. Schmidt, and G. Zizka. 2014. Introducing African plants—A photo guide—an

interactive photo data☐base and rapid identification tool for continental Africa. *Taxon* 63:

1159–1161.

Emms, D. M., and S. Kelly. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome biology* 20: 238.

Favaro, R., A. Pinyopich, R. Battaglia, M. Kooiker, L. Borghi, G. Ditta, M. F. Yanofsky, et al.

2003. MADS-box protein complexes control carpel and ovule development in Arabidopsis.

The Plant cell 15: 2603–2611.

Feng, G., B. J. Sanderson, K. Keefover-Ring, J. Liu, T. Ma, T. Yin, L. B. Smart, et al. 2020.

Pathways to sex determination in plants: how many roads lead to Rome? *Current opinion in*

plant biology 54: 61–68.

Gagliardi, K. B., I. Cordeiro, and D. Demarco. 2018. Structure and development of flowers and

inflorescences in Peraceae and Euphorbiaceae and the evolution of pseudanthia in

Malpighiales. *PloS one* 13: e0203954.

Gilbert, M. G. 1994. The relationships of the Euphorbieae (Euphorbiaceae). *Annals of the*

Missouri Botanical Garden. Missouri Botanical Garden 81: 283.

Goto, K., and E. M. Meyerowitz. 1994. Function and regulation of the Arabidopsis floral

homeotic gene PISTILLATA. *Genes & development* 8: 1548–1560.

Grant, C. E., T. L. Bailey, and W. S. Noble. 2011. FIMO: scanning for occurrences of a given

motif. *Bioinformatics* 27: 1017–1018.

Hui, W., Y. Yang, G. Wu, C. Peng, X. Chen, and M. Z. Zayed. 2017. Transcriptome profile

analysis reveals the regulation mechanism of floral sex differentiation in Jatropha curcas L.

Scientific reports 7: 16421.

Irish, V. 2017. The ABC model of floral development. *Current biology: CB* 27: R887–R890.

Jack, T., L. L. Brockman, and E. M. Meyerowitz. 1992. The homeotic gene APETALA3 of

Arabidopsis thaliana encodes a MADS box and is expressed in petals and stamens. *Cell*

68: 683–697.

Jin, R., S. Klasfeld, Y. Zhu, M. Fernandez Garcia, J. Xiao, S.-K. Han, A. Konkol, and D. Wagner.

2021. LEAFY is a pioneer transcription factor and licenses cell reprogramming to floral fate.

Nature communications 12: 626.

Jofuku, K. D., B. G. den Boer, M. Van Montagu, and J. K. Okamuro. 1994. Control of

Arabidopsis flower and seed development by the homeotic gene APETALA2. *The Plant cell*

6: 1211–1225.

Johnson, A. R., Y. Yue, S. B. Carey, S. J. Park, L. H. Kruse, A. Bao, A. Pasha, et al. 2023.

Chromosome-level genome assembly of Euphorbia peplus, a model system for plant latex,

reveals that relative lack of Ty3 transposons contributed to its small genome size. *Genome*

biology and evolution 15: evad018.

Kaplan, D., and C. D. Specht. 2022. Kaplan's Principles of Plant Morphology. CRC Press.

Klepikova, A. V., M. D. Logacheva, S. E. Dmitriev, and A. A. Penin. 2015. RNA-seq analysis of

an apical meristem time series reveals a critical point in Arabidopsis thaliana flower

initiation. *BMC genomics* 16: 466.

Kobayashi, Y., H. Kaya, K. Goto, M. Iwabuchi, and T. Araki. 1999. A pair of related genes with

antagonistic roles in mediating flowering signals. *Science* 286: 1960–1962.

Kolde. Pheatmap: pretty heatmaps. *R package version*.

Kozlov, A. M., D. Darriba, T. Flouri, B. Morel, and A. Stamatakis. 2019. RAxML-NG: a fast,

scalable and user-friendly tool for maximum likelihood phylogenetic inference.

Bioinformatics 35: 4453–4455.

Lai, X., H. Daher, A. Galien, V. Hugouvieux, and C. Zubieta. 2019. Structural Basis for Plant

MADS Transcription Factor Oligomerization. *Computational and structural biotechnology*

journal 17: 946–953.

Lee, H., S. S. Suh, E. Park, E. Cho, J. H. Ahn, S. G. Kim, J. S. Lee, et al. 2000. The

AGAMOUS-LIKE 20 MADS domain protein integrates floral inductive pathways in

Arabidopsis. *Genes & development* 14: 2366–2376.

Levin, J. Z., and E. M. Meyerowitz. 1995. UFO: an Arabidopsis gene involved in both floral

meristem and floral organ development. *The Plant cell* 7: 529–548.

Li, C., J. Su, X. Liu, S. Chen, and L. He. 2014. Pistillate Flower Development and Stigma

Receptivity of Euphorbia pulcherrima. *Agricultural Science & Technology* 15: 1671–1675.

Liljegren, S. J., G. S. Ditta, Y. Eshed, B. Savidge, J. L. Bowman, and M. F. Yanofsky. 2000.

SHATTERPROOF MADS-box genes control seed dispersal in Arabidopsis. *Nature* 404:

766–770.

Lippman, Z. B., O. Cohen, J. P. Alvarez, M. Abu-Abied, I. Pekker, I. Paran, Y. Eshed, and D.

Zamir. 2008. The making of a compound inflorescence in tomato and related nightshades.

PLoS biology 6: e288.

Litt, A., and E. M. Kramer. 2010. The ABC model and the diversification of floral organ identity.

Seminars in cell & developmental biology 21: 129–137.

Liu, J., C. Shi, C.-C. Shi, W. Li, Q.-J. Zhang, Y. Zhang, K. Li, et al. 2020. The Chromosome-

Based Rubber Tree Genome Provides New Insights into Spurge Genome Evolution and

Rubber Biosynthesis. *Molecular plant* 13: 336–350.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome biology* 15: 550.

Machanick, P., and T. L. Bailey. 2011. MEME-ChIP: motif analysis of large DNA datasets.

Bioinformatics 27: 1696–1697.

Ma, H., M. F. Yanofsky, and E. M. Meyerowitz. 1991. AGL1-AGL6, an Arabidopsis gene family

with similarity to floral homeotic and transcription factor genes. *Genes & development* 5:

484–495.

Mandel, M. A., and M. F. Yanofsky. 1995. The Arabidopsis AGL8 MADS box gene is expressed

in inflorescence meristems and is negatively regulated by APETALA1. *The Plant cell* 7:

1763–1771.

Miller, C. N., J. Dumenil, F. H. Lu, C. Smith, N. McKenzie, V. Chapman, J. Ball, et al. 2020.

Variation in the expression of a transmembrane protein influences cell growth in

Arabidopsis thaliana petals by altering auxin responses. *BMC plant biology* 20: 482.

Mitchell, C. H., and P. K. Diggle. 2005. The evolution of unisexual flowers: morphological and

functional convergence results from diverse developmental transitions. *American journal of*

botany 92: 1068–1076.

Mizukami, Y., and H. Ma. 1997. Determination of Arabidopsis floral meristem identity by

AGAMOUS. *The Plant cell* 9: 393–408.

Morel, P., P. Chambrier, V. Boltz, S. Chamot, F. Rozier, S. Rodrigues Bento, C. Trehin, et al.

2019. Divergent Functional Diversification Patterns in the SEP/AGL6/AP1 MADS-Box

Transcription Factor Superclade. *The Plant cell* 31: 3033–3056.

Morel, P., K. Heijmans, K. Ament, M. Chopy, C. Trehin, P. Chambrier, S. Rodrigues Bento, et al.

2018. The Floral C-Lineage Genes Trigger Nectary Development in Petunia and

Arabidopsis. *Plant Cell* 30: 2020–2037.

Moyroud, E., E. Kusters, M. Monniaux, R. Koes, and F. Parcy. 2010. LEAFY blossoms. *Trends*
in plant science 15: 346–352.

O'Brien, M., R. N. Kaplan-Levy, T. Quon, P. G. Sappl, and D. R. Smyth. 2015. PETAL LOSS, a
trihelix transcription factor that represses growth in Arabidopsis thaliana, binds the energy-
sensing SnRK1 kinase AKIN10. *Journal of experimental botany* 66: 2475–2485.

Pelaz, S., G. S. Ditta, E. Baumann, E. Wisman, and M. F. Yanofsky. 2000. B and C floral organ
identity functions require SEPALLATA MADS-box genes. *Nature* 405: 200–203.

Prenner, G., N. I. Cacho, D. Baum, and P. J. Rudall. 2010. Is LEAFY a useful marker gene for
the flower–inflorescence boundary in the Euphorbia cyathium? *Journal of experimental*
botany 62: 345–350.

Prenner, G., S. D. Hopper, and P. J. Rudall. 2008. Pseudanthium development in Calycopeplus
paucifolius, with particular reference to the evolution of the cyathium in Euphorbieae
(Euphorbiaceae–Malpighiales). *Australian systematic botany* 21: 153–161.

Prenner, G., and P. J. Rudall. 2007. Comparative ontogeny of the cyathium in Euphorbia
(Euphorbiaceae) and its allies: exploring the organ-flower-inflorescence boundary.
American journal of botany 94: 1612–1629.

Quinlan, A. R., and I. M. Hall. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26: 841–842.

Rieu, P., M. Arnoux-Courseaux, G. Tichtinsky, and F. Parcy. 2023a. Thinking outside the F-box:
how UFO controls angiosperm development. *The New phytologist* 240: 945–959.

Rieu, P., L. Turchi, E. Thévenon, E. Zarkadas, M. Nanao, H. Chahtane, G. Tichtinsky, et al.

2023b. The F-box protein UFO controls flower development by redirecting the master

transcription factor LEAFY to new cis-elements. *Nature plants* 9: 315–329.

Robert, X., and P. Gouet. 2014. Deciphering key features in protein structures with the new

ENDscript server. *Nucleic acids research* 42: W320–4.

Rodríguez-Pelayo, C., B. A. Ambrose, A. Vasco, J. F. Alzate, and N. Pabón-Mora. 2022.

Evolution and expression of LEAFY genes in ferns and lycophytes. *EvoDevo* 13: 2.

Schmitz, R. J., L. Hong, S. Michaels, and R. M. Amasino. 2005. FRIGIDA-ESSENTIAL 1

interacts genetically with FRIGIDA and FRIGIDA-LIKE 1 to promote the winter-annual habit

of Arabidopsis thaliana. *Development* 132: 5471–5478.

Schultz, E. A., and G. W. Haughn. 1991. LEAFY, a Homeotic Gene That Regulates

Inflorescence Development in Arabidopsis. *The Plant cell* 3: 771–781.

Shen, W., S. Le, Y. Li, and F. Hu. 2016. SeqKit: A Cross-Platform and Ultrafast Toolkit for

FASTA/Q File Manipulation. *PloS one* 11: e0163962.

Slavković, F., C. Dogimont, H. Morin, A. Boualem, and A. Bendahmane. 2021. The Genetic

Control of Nectary Development. *Trends in plant science* 26: 260–271.

Supek, F., M. Bošnjak, N. Škunca, and T. Šmuc. 2011. REVIGO summarizes and visualizes

long lists of gene ontology terms. *PloS one* 6: e21800.

Takeda, S., Y. Hamamura, T. Sakamoto, S. Kimura, M. Aida, and T. Higashiyama. 2022. Non-

cell-autonomous regulation of petal initiation in Arabidopsis thaliana. *Development* 149.

Tang, M., Y.-B. Tao, Q. Fu, Y. Song, L. Niu, and Z.-F. Xu. 2016. An ortholog of LEAFY in

Jatropha curcas regulates flowering time and floral organ development. *Scientific reports* 6:

37306.

Tuskan, G. A., S. Difazio, S. Jansson, J. Bohlmann, I. Grigoriev, U. Hellsten, N. Putnam, et al.

2006. The genome of black cottonwood, Populus trichocarpa (Torr. & Gray). *Science* 313:

1596–1604.

Valliyodan, B., S. B. Cannon, P. E. Bayer, S. Shu, A. V. Brown, L. Ren, J. Jenkins, et al. 2019.

Construction and comparison of three reference-quality genome assemblies for soybean.

The Plant journal: for cell and molecular biology 100: 1066–1082.

Wang, M., Z. Gu, Z. Fu, and D. Jiang. 2021. High-quality genome assembly of an important

biodiesel plant, Euphorbia lathyris L. *DNA research: an international journal for rapid*

publication of reports on genes and genomes 28.

Weigel, D., J. Alvarez, D. R. Smyth, M. F. Yanofsky, and E. M. Meyerowitz. 1992. LEAFY

controls floral meristem identity in Arabidopsis. *Cell* 69: 843–859.

Western Australian Herbarium, Biodiversity, Conservation Science, Department of Biodiversity,

Conservation, and Attractions. Calycopeplus Planch. Website

https://florabase.dbca.wa.gov.au/browse/profile/21684 [accessed 25 January 2024].

Xanthopoulou, A., J. Montero-Pau, B. Picó, P. Boumpas, E. Tsaliki, H. S. Paris, A. Tsaftaris, et

al. 2021. A comprehensive RNA-Seq-based gene expression atlas of the summer squash

(Cucurbita pepo) provides insights into fruit morphology and ripening mechanisms. *BMC*

genomics 22: 341.

Xu, G., J. Huang, Y. Yang, and Y.-A. Yao. 2016. Transcriptome Analysis of Flower Sex

Differentiation in Jatropha curcas L. Using RNA Sequencing. *PloS one* 11: e0145613.

Xu, W., D. Wu, T. Yang, C. Sun, Z. Wang, B. Han, S. Wu, et al. 2021. Genomic insights into the

origin, domestication and genetic basis of agronomic traits of castor bean. *Genome biology*

22: 113.

Yang, H., P. Zhang, D. Guo, N. Wang, H. Lin, X. Wang, and L. Niu. 2023. Transcriptional

repressor AGL79 positively regulates flowering time in Arabidopsis. *Journal of plant*

physiology 285: 153985.

Zhang, X., L. Pan, W. Guo, Y. Li, and W. Wang. 2022. A convergent mechanism of sex

determination in dioecious plants: Distinct sex-determining genes display converged

regulation on floral B-class genes. *Frontiers in plant science* 13: 953445.

Zumajo-Cardona, C., and N. Pabón-Mora. 2016. Evolution of the APETALA2 Gene Lineage in

Seed Plants. *Molecular biology and evolution* 33: 1818–1832.

FIGURE LEGENDS

Figure 1: The *Euphorbia* cyathium is a unique reproductive structure. A. Diagram of a *Euphorbia*

peplus cyathium. B. *E. peplus* cyathium and adjacent primordia and cyathophylls.

Scale=2mm. C. *E. peplus* cyathium and adjacent primordia with adjacent cyathophylls

removed. Scale=1mm. D. *E. peplus* cyathium with involucre and nectaries removed.

Scale=500 microns. E. Diagram of the ABCDE model of floral development and the four whorls

of a traditional flower; D class genes, which control ovule development, are not shown.

Figure 2: Close relatives of *Euphorbia*, *Calycopeplus* and *Neoguillauminia*, have perianths

around their pistillate and staminate flowers and a condensed inflorescence similar to that of

Euphorbia. *Anthostema*, which is less closely related, has perianths around its pistillate and

staminate flowers and a less-condensed inflorescence.

Figure 3: Staminate flowers have a distinct expression pattern. A-D. Illustration of cyathium

parts used for RNAseq sampling. A. Small primordium. B. Large primordium. C. Young

staminate flower and young pistillate flower. D. Mature organs: nectary, involucre, staminate

flower, pistillate flower, filiform structure, involucre, nectary. (Duplicates of organs are omitted

for simplicity. Cyathophyll is not shown.) E. PCA plot of variance-stabilized-transformed data

for the 500 genes with the most variance. PC1 explains 22.4% of the variance, PC2 explains

18.9% of the variance, PC3 explains 16.7% of the variance, and PC4 explains 10.6% of the

variance.

Figure 4: E class gene SEP3 is highly expressed in the pistillate and staminate flowers. A.

Heatmap shows expression of ABCDE homologs and other floral identity genes in different

organs of the *E. peplus* cyathium. B-N. Images of the organs sampled. B. Young pistillate

flower. C. Young staminate flower. D. Nectary. E. Filiform structure. F. Mature pistillate flower.

G. Mature staminate flower. Scale bars = 250 microns.

Figure 5: *E. peplus* staminate flowers and pistillate flowers have higher transcriptomic

conservation with arabidopsis stamens and carpels than *E. peplus* filiform structures and

involucres have with arabidopsis petals and sepals. Total number of orthogroups with

upregulated genes in each *E. peplus* structure (compared with *E. peplus* cyathophylls) shown in

plot: involucre (vs. sepal): 649; filiform structure (vs. petal): 1422; staminate flower (vs. stamen):

1424; pistillate flower (vs. carpel): 473.

Figure 6: Biological Process GO terms for genes significantly upregulated in the nectary,

involucre, and filiform structures as compared with the cyathophyll and Molecular Function GO

terms for genes significantly upregulated in the nectary as compared with the cyathophyll are

summarized using REVIGO. "Value" is the log10(p-value); a lower value, towards the red side

of the color scale, is more significant. The size of the points, the dispensability, represents the

semantic similarity to other terms in the dataset.

Figure 7: LFY, UFO, and B class genes show signs of divergence and relaxed selection in

Euphorbia. A. dN/dS values shown on gene trees of floral genes of interest; RAxML support

value from 1000 bootstraps is shown at nodes.   PHYA is shown as a non-floral "control".   B.

Visualization of amino acid alignment in a highly conserved region of LFY; site 279 is a DNA

binding site that is highly conserved (except in *Euphorbia*). Cyan color shows sites in the LFY

protein alignment rated as "highly significant" for positive selection in the *Euphorbia* clade by

PAML. C. Visualization of amino acid alignment at the start of the UFO protein showing

deletions in *Euphorbia* ("Ep" and "Casp" genes). Gene names and their gene ID equivalents are

listed in Supplementary File 3.

Figure 8: Proposed new model of *Euphorbia* cyathium development in relation to angiosperm

and gymnosperm reproductive development.