Altered markers of brain metabolism and excitability are associated with executive functioning in young children exposed to alcohol *in utero*

Meaghan V. Perdue[1,2,3], Mohammad Ghasoub[1,2,3], Madison Long[1,2,3], Marilena M. DeMayo[1,2,3,4,5], Tiffany K. Bell[1,2,3], Carly A. McMorris[2,3,6], Deborah Dewey[2,3,7,8], W. Ben Gibbard[2,7], Christina Tortorelli[9], Ashley D. Harris[1,2,3] & Catherine Lebel[1,2,3]

[1]University of Calgary, Department of Radiology, Calgary, Alberta, Canada

[2]Alberta Children's Hospital Research Institute, Calgary, Alberta, Canada

[3]University of Calgary, Hotchkiss Brain Institute, Calgary, Alberta, Canada

[4]Mathison Centre for Mental Health Research and Education, Calgary, Alberta, Canada

[5]University of Calgary, Department of Psychiatry, Calgary, Alberta, Canada

[6]University of Calgary, Werklund School of Education, Calgary, Alberta, Canada

[7]University of Calgary, Department of Pediatrics, Calgary, Alberta, Canada

[8]University of Calgary, Department of Community Health Sciences, Calgary, Alberta, Canada

[9]Mount Royal University, Department of Social Work, Calgary, Alberta, Canada

Corresponding Author: Meaghan V. Perdue - meaghan.perdue@ucalgary.ca

Alberta Children's Hospital, 28 Oki Drive NW, Calgary, AB T3B 6A8, Canada

Abstract

Prenatal alcohol exposure (PAE) is the leading known cause of birth defects and cognitive disabilities, with impacts on brain development and executive functioning. Abnormalities in structural and functional brain features are well-documented in children with PAE, but the effects of PAE on brain metabolism in children have received less attention. Levels of brain metabolites can be measured non-invasively using magnetic resonance spectroscopy (MRS). Here, we present the first study of PAE-related brain metabolite differences in early childhood (ages 3-8 years) and their associations with cognitive performance, including executive functioning (EF) and pre-reading skills. We measured metabolites in two cohorts of children with PAE and unexposed children using MRS in the anterior cingulate cortex (ACC; cohort 1) and left temporo-parietal cortex (LTP; cohort 2). Total choline (tCho), a marker of membrane/myelin metabolism, was elevated in both regions in children with PAE compared to UC, and glutamate+glutamine (Glx), a marker of excitability, was elevated in the ACC. The PAE group exhibited more difficulties with EF, and higher tCho was associated with better EF in both PAE and unexposed groups. In addition, elevated Glx in the ACC was associated with poorer inhibitory control within the PAE group only. LTP metabolites were not significantly associated with pre-reading skills in PAE or unexposed groups. Together, these findings point to altered membrane metabolism and excitability in young children with PAE. These findings provide new insight to potential mechanisms by which PAE disrupts brain development and cognitive functioning in early childhood.

Keywords: magnetic resonance spectroscopy (MRS), alcohol, metabolism, childhood, cognition

Statements and Declarations

Competing Interests: None

Acknowledgements: This work was supported by the Alberta Children's Hospital Research Institute and the Canadian Institutes of Health Research (IHD-134090, MOP-136797). Salary support was provided by the Canada Research Chair Program (ADH, CL), the Hotchkiss Brain Institute (MMD), the Cumming School of Medicine (MVP, MMD), and the Killam Trusts (MVP).

Author Contributions:

Meaghan V. Perdue: conceptualization, methodology, formal analysis, writing – original draft, writing – review & editing, data visualization; **Mohammad Ghasoub:** data curation, formal analysis, writing – original draft, writing – review & editing; **Madison Long:** data curation, writing – original draft, writing – review & editing; **Marilena M. DeMayo:** methodology, writing – review & editing; **Tiffany K. Bell:** methodology, writing – review & editing; **Carly A. McMorris:** conceptualization, writing – review & editing; **Deborah Dewey:** conceptualization, writing – review & editing, funding acquisition; **W. Ben Gibbard:** writing – review & editing; **Christina Tortorelli:** writing – review & editing; **Ashley D. Harris:** methodology, writing – review & editing, supervision; **Catherine Lebel:** conceptualization, methodology, writing – review & editing, funding acquisition, supervision.

Data Availability: Code used for pre-processing and quantification of MRS data is available at https://github.com/developmental-neuroimaging-lab/mrs; De-identified participant demographics, metabolite concentration values, cognitive scores, and data processing and analysis code are available at https://osf.io/4zkd8/?view_only=f7526d82205044feb07796410a937ca5

Introduction

Alcohol consumption during pregnancy can lead to a range of cognitive, behavioral, and neurological difficulties in offspring (Popova et al., 2023). Prenatal alcohol exposure (PAE) is the leading known cause of developmental disabilities, with approximately 10% of children exposed worldwide (Popova et al., 2023). Alcohol is a teratogen that can easily cross the placenta, disrupting the fetus' neurodevelopment at any stage. Disruptions include, but are not limited to, altered neuronal proliferation and migration (Almeida et al., 2020; Veazey et al., 2013), synaptogenesis (Adams et al., 2023; Basavarajappa & Subbanna, 2023), dendritic spine morphology (Cui et al., 2010; Lee et al., 2021), and gliogenesis and myelination (Cantacorps et al., 2017; Darbinian & Selzer, 2022), including phospholipid metabolism (Darbinian et al., 2023; Hwang et al., 2023).

The teratogenic effects of alcohol on brain development can lead to long-term cognitive and behavioral difficulties (Mattson et al., 2019; Subramoney et al., 2018). Children with PAE may be diagnosed with Fetal Alcohol Spectrum Disorder (FASD), which is characterized by physical and neurodevelopmental impairments, though diagnostic criteria vary (Cook et al., 2016; Popova et al., 2023). Children with PAE often exhibit attention-deficit/hyperactivity disorder (ADHD)-like behavior (Pinner et al., 2023), along with executive and social functioning challenges (Khoury et al., 2015; Kodituwakku et al., 2001; Mattson et al., 2019; Mattson & Riley, 1999; Rasmussen et al., 2007, 2013; Rasmussen & Bisanz, 2009; Rockhold et al., 2021). In addition, children with PAE are more likely than their unexposed peers to have difficulties in academic performance, including reading and math (Kodituwakku et al., 2011; O'Leary et al., 2013). Academic difficulties experienced by children with PAE may arise from domain-general cognitive difficulties, including problems in memory, attention, processing speed and executive functions (Kodituwakku et al., 2011).

These behavioral difficulties emerge in the context of PAE-related brain differences (Donald et al., 2015; Lebel et al., 2011; Moore & Xia, 2022). Reduced white matter and gray matter volume (Nardelli et al., 2011; Subramoney et al., 2022), altered microstructure and developmental trajectories (Donald et al., 2024; Gimbel et al., 2023; Kar et al., 2021, 2022; M. Long et al., 2024; X. Long & Lebel, 2022), and disrupted structural and functional connectivity (X. Long et al., 2019, 2020; Roos et al., 2021; Wozniak et al., 2013) have been reported in children and adolescents with PAE. Moreover, neural alterations are observed even with relatively low levels of PAE, around 1 drink/week (X. Long & Lebel, 2022).

Despite known impacts of PAE on brain development, the neurobiological mechanisms underlying cognitive difficulties in children with PAE remain poorly understood. Examining features of brain metabolism and chemistry provides complementary evidence to structural and functional neuroimaging data, offering a window to biochemical mechanisms underlying altered brain development, metabolism, and functioning in those with PAE. Brain metabolites are naturally occurring chemicals that are crucial for development, metabolism, and neural signalling that can be measured non-invasively using magnetic resonance spectroscopy (MRS). Five key brain metabolites of interest that are measured using standard MRS methods include total NAA (N-acetylaspartate + N-acetylaspartylglutamate; tNAA), which reflects neuronal health and metabolism, total creatine (creatine + phosphocreatine; tCr), which reflects energy homeostasis, total choline (phosphocholine + glycerophosphocholine + choline, tCho), which reflects phospholipid metabolism; glutamate +

glutamine (Glx), which reflects neural excitability, and myo-inositol (Ins) a marker of glial tissue. For an in-depth discussion of the roles of each metabolite in the brain, see Rae (2014).

MRS offers insight to potential neurochemical and metabolic mechanisms by which alcohol disrupts brain structure development and functioning. Specifically, it has been posited that alcohol disrupts the cortico-striatal excitation/inhibition balance by increasing glutamatergic activity and supressing migration of inhibitory neurons to their cortical targets (Bariselli & Lovinger, 2021). This system can be investigated in human using MRS to measurement Glx and g-aminobutyric acid (GABA). In addition, myelination is impaired in PAE, and preclinical models point to disruption of oligodendrocyte proliferation, migration, and maturation (Darbinian & Selzer, 2022). However, the mechanism by which alcohol impacts myelination is not fully understood. The tCho signal in MRS is a marker of phospholipid metabolism, including the synthesis/repair and break-down of neuronal cell membranes and myelin (Rae, 2014), which may be impacted by PAE.

Despite the potential of MRS to provide insight to neurometabolic brain alterations, only six studies have applied MRS to investigate PAE, and half of these included sample sizes of 10 or fewer in each group. Nonetheless, several trends have emerged in this small body of literature, indicating altered brain metabolism and excitability in infants and youth with PAE relative to controls. Reduced tCho in children and adolescents with PAE has been reported in three studies that measured metabolites in different regions, with effects shown in frontal parietal white matter, left striatum, and cerebellum, indicating that tCho may be affected globally in PAE (Astley et al., 2009; du Plessis et al., 2014; Gonçalves et al., 2009). Lower tCho in PAE is thought to indicate altered phospholipid metabolism, impaired myelin maintenance, and reduced cell density/arborization relative to unexposed groups. One study linked higher rates of alcohol exposure during pregnancy with lower tCho and tNAA, and higher Glx, in children ages 8-17 years, highlighting the persistent effects of PAE on neurometabolic outcomes into adolescence (du Plessis et al., 2014). Glx differences in PAE are mixed across studies: du Plessis and colleagues (2014) found that elevated cerebellar Glx levels in children/adolescents were associated with increased rates of alcohol consumption around conception and during pregnancy, while Howells and colleagues (2016) found that parietal white matter Glx levels were reduced in newborns with PAE relative to controls. These contrasting patterns may reflect differences across studies in developmental stages examined, brain regions and tissue types from which metabolites were measured, and relative contributions of glutamate and glutamine to the Glx signal. These varied methods and findings highlight the need for further investigation across different age groups and brain regions, to better characterize the impacts of PAE on neurochemistry across development. Moreover, no study to date has investigated associations between metabolites and cognitive traits in participants with PAE.

Here, we present the first study of neurometabolic differences associated with PAE in early childhood, and report novel links between brain metabolites and cognition in this population. We measured brain metabolites and cognitive traits in two cohorts of young children with PAE in comparison to unexposed controls. Metabolite levels (tCho, tNAA, tCr, Glx, & mI) were measured in the anterior cingulate cortex (ACC; Cohort 1) and left temporo-parietal cortex (LTP; Cohort 2). Children's EF (Cohort 1) and pre-reading skills (Cohort 2) were assessed using age-appropriate standardized measures. We predicted that children with PAE would have lower tCho and higher Glx relative to controls; our analyses of tNAA, tCr and

ml were exploratory given the lack of prior research in this population. In addition, we predicted that tCho and Glx in the ACC would be associated with EF, while tCho and Glx levels in the LTP would be associated with pre-reading skills.

Methods

Participants

This study included two cohorts of children with PAE from a longitudinal study of early childhood brain development in children with PAE, and unexposed children drawn from a longitudinal study of typical brain development with matching brain imaging protocols. Parent/guardian written informed consent and child verbal assent were obtained for each child. Both studies were approved by the University of Calgary's Conjoint Health Research Ethics Board (REB14-2266, REB13-0020).

Children with PAE

Cohort 1 (ACC) consisted of 21 datasets collected from 20 children (11 females) with PAE (5.38 ± 1.11 years). Cohort 2 (LTP) consisted of 28 datasets collected from 24 subjects (16 females) with PAE (6.45 ± 0.764 years). Ten children with PAE had MRS data from both regions and were included in both cohorts; in these children, MRS data were acquired from each region at different study visits (6 months - 1 year apart). These numbers reflect the participants included in analysis after exclusion for data quality issues (detailed below). Children from both cohorts were recruited from Alberta Children's Services, caregiver support groups, and early intervention services. PAE was confirmed from the children's welfare file and/or from caregivers, caseworkers, and family members. All children were in adoptive, foster, or kinship care. None of the children had a formal FASD diagnosis at the time of assessment given their young age, as FASD diagnoses in some parts of Alberta typically begins at 7 years of age. All children were born at ≥34 weeks' gestation and spoke English as their primary language. Children with epilepsy, autism spectrum disorder, history of head injury, MRI contraindications, and any other genetic or medical conditions associated with cognitive disabilities were excluded from the study. Given their high co-occurrence with PAE, children with conditions such as ADHD, learning disabilities, and other mental health diagnoses were included. Further details on the children in this study are reported in (Kar et al., 2021).

Unexposed children

Cohort 1 (ACC) consisted of 76 datasets collected from 72 children (33 females; 4.51 ± 0.950 years). Cohort 2 (LTP) consisted of 128 datasets from 71 children (34 females; 6.45 ± 0.925 years). 39 unexposed children were included in both cohorts, but MRS data from the ACC and LTP were collected at different study visits (6 months - 1 year apart). These numbers reflect the participants included in analysis after exclusion for data quality issues (detailed below). Children from both cohorts were recruited from the community in Calgary, Alberta as well as the Alberta Pregnancy Outcomes and Nutrition (APrON) study. All children were born at ≥35 weeks gestation and spoke English as their primary language. Inclusion criteria included no MRI contraindications, history of brain injury, or genetic, motor, or developmental disorders at the time of enrollment. These children had no PAE as verified by maternal report. Further details on this study are reported in (Reynolds et al., 2020). The sample of unexposed children in this study were included in a larger study of metabolite development across

childhood (Perdue et al., 2023); the present study includes a subset of these children within age ranges restricted to match the age ranges of the participants with PAE for each cohort.

Metabolite data acquisition

Brain metabolites were measured using magnetic resonance spectroscopy (MRS) included in a multimodal magnetic resonance imaging (MRI) protocol. MRI and MRS data were acquired with a 3T GE MR750w MR system with a 32-channel head coil at the Alberta Children's Hospital (Calgary, AB, Canada) by staff who are highly skilled in pediatric neuroimaging. Children were scanned while watching a movie or during natural sleep; no sedation was used. T1-weighted anatomical images were acquired using a spoiled gradient echo sequence (210 axial slices; 0.9 x 0.9 x 0.9mm resolution, TR = 8.23 ms, TE = 3.76 ms, flip angle = 12º, matrix size = 512 x 512, inversion time = 540 ms). These images were reconstructed to provide axial, sagittal, and coronal views at the scanner which were used for placement of spectroscopy voxels. MRS data were acquired using short echo time Point RESolved spectroscopy (PRESS; TE = 30 ms, TR = 2000 ms, 96 averages, 20 x 20 x 15 mm voxel size). A strong body of literature demonstrates the test-retest reliability and stability of PRESS sequences and supports the validity of MRS measurements for longitudinal research (Baeshen et al., 2020; Fayed et al., 2009; Gasparovic et al., 2011; Soreni et al., 2010; Volk et al., 2018, 2019). MRS voxels were placed in either the ACC or in the LTP by trained members of the research team according to detailed instructions and reference images. The ACC voxel was localized anterior to and at approximately the same level as the genu of the corpus callosum viewed on a midsagittal slice and consisted almost entirely of gray matter (Figure 1). The LTP voxel was localized to capture the left angular gyrus based on all three image planes, and included portions of the supramarginal gyrus, parietal operculum, and posterior superior temporal gyrus due to the extent of the voxel (Figure 1). The LTP voxel consisted of gray and white matter, excluding CSF to the extent possible.

Figure 1. MRS voxel positions and sample spectra. ACC shown on top row, LTP shown on bottom row, participants with PAE shown on left, unexposed participants shown on right. Sample participant age and sex are as follows: Cohort 1 PAE: 4.87 year-old male, Cohort 1 unexposed: 4.89 year-old male, Cohort 2 PAE: 6.14 year-old female, Cohort 2 unexposed: 6.25 year-old female

Metabolite data processing and quantification

The PRESS acquisition was pre-processed with the FID-A toolbox (Simpson et al., 2017). Pre-processing included coil combination, removal of bad averages, frequency alignment, and phase correction, according to recent recommendations (Near et al., 2020). Metabolites were then fit using LCModel v6.3 (Provencher, 1993). The metabolites of interest were: tNAA, tCr, tCho, mI and Glx. The basis set used for quantification included alanine, ascorbate, aspartate, choline, citrate, creatine, ethanol, GABA, glycerophosphocholine, glutathione, glucose, glycine, glutamine, glutamate, water, mI, lactate, NAA, NAAG, phosphocholine, phosphocreatine, phosphoryl ethanolamine, scyllo-inositol, taurine and beta-Hydroxybutyrate, simulated using the FID-A toolbox (Simpson et al., 2017) with sequence specific timings and RF pulse shapes. Fitted spectra were visually inspected for quality assurance and datasets with poor spectra were excluded from analysis (ACC [Cohort 1]: PAE n=5, unexposed n=3; LTP [Cohort 2]: PAE n=0, unexposed n=3). Quantitative quality metrics (signal-to-noise ratio [SNR] and linewidth [LW] for NAA) were obtained using the "op_getLW.m" and "op_getSNR.m" functions in FID-A. The Gannet CoRegStandAlone function (Harris et al., 2015), which calls SPM12 segmentation (Ashburner & Friston, 2005), was used to quantify tissue content within the MRS voxels by co-registering them to the T1-weighted image acquired during the same scanning session. Co-registration and segmentation outputs were visually inspected to ensure accurate localization and segmentation of the MRS voxels. Absolute metabolite values are expressed

in molal units (moles/kg), as per the approach recommended in the MRS consensus statement on metabolite quantification (Near et al 2020) and described by Gasparovic et al (2006). This method accounts for the differential water T1- and T2-relaxation in white matter, gray matter and CSF in the voxel to enable absolute metabolite measurements. The scripts for the correction were developed by (DeMayo et al., 2023), and are publicly available (https://github.com/HarrisBrainLab/TissueCorrections). However, this approach does not address differences in metabolite concentrations between white matter and gray matter (as suggested in (Gasparovic et al., 2018)) as this ratio is not agreed upon for all metabolites (and may change with development). For this reason, the gray matter tissue fraction was included as a covariate in analyses.

The gray matter tissue fraction of each MRS voxel was calculated as the fraction of voxel volume composed of gray matter divided by total tissue within the voxel (sum of the fractions of voxel volume for gray matter (GM) and white matter (WM; formula: fGM/[fGM + fWM]). The midline location of the ACC voxel was set to exclude white matter to the extent possible. To ensure minimal white matter in the ACC voxel, tissue fraction outliers were identified using boxplots and Rosner's test using the *EnvStats R* package (Millard, 2013), and two datasets with tissue fraction <.88 were excluded from analysis (both from the unexposed group). Given that the LTP voxel included a substantial mix of gray and white matter in all participants, tissue fraction was applied as a covariate in analyses to account for individual differences in tissue composition of the voxels to ensure that group differences reflect differences in metabolites and not in tissue content of the voxels.

Code for processing and quantification of MRS is available here: https://github.com/developmental-neuroimaging-lab/mrs

Cognitive assessments

Participants' executive function and pre-reading abilities were measured with the NEuroPSYchological Assessment - Second Edition (NEPSY-II; Korkman et al., 2007), and the caregiver-report Behavior Rating Inventory of Executive Function (BRIEF; Gioia et al., 2000) or BRIEF-Preschool; (BRIEF-P; Gioia et al., 2003) using age-appropriate forms (BRIEF-P: ages 2-5 years, BRIEF: ages ≥6 years), most often administered on the same day or within two weeks of the MRI scan and always within six months. Cognitive data was available for a sub-set of participants in each cohort; EF skills (BRIEF/BRIEF-P and NEPSY-II Statue subtest) were examined in Cohort 1, and pre-reading skills (NEPSY-II Phonological Processing and Speeded Naming subtests) were examined in Cohort 2.

NEPSY-II

Inhibitory control (motor inhibition) was assessed via the Statue subtest of the NEPSY-II. Participants were asked to stand still with their eyes closed and inhibit responses to sound distractors. A researcher observed the child while they tried to stand motionless with eyes closed for a 75 second period. A perfect score would indicate that the child remained still with eyes closed the entire time with increasingly lower scores indicating more errors (movement, eyes opened) during the task.

Key pre-reading skills were assessed via the Phonological Processing (PP) and Speeded Naming (SN) subtests of the NEPSY-II. PP measured the child's ability to segment and manipulate phonemes, and SN measured the child's ability to rapidly identify pictured objects and colors.

Performance on NEPSY-II sub-tests is indexed by scaled scores based on population norms (mean=10, SD=3) for child age. Higher scores indicate better performance.

<u>BRIEF and BRIEF-P</u>

Global executive functioning was assessed using the BRIEF or BRIEF-P as appropriate for each child's age. These measures are parent-report measures of naturalistic executive function. Children's executive functioning abilities across various sub-domains are summarized in the Global Executive Composite (GEC) T-score, normed for child age. GEC T-scores are scaled based on population norms (mean=50, SD=10) for child age. Higher GEC T-scores indicate poorer performance.

Data analysis

Group differences in metabolite levels

Statistical analysis was performed in RStudio (RStudio Team, 2020) using *R* version 4.3.1 (R Core Team, 2016). Group differences in metabolite levels were tested using linear mixed effects models (*lme4* and *lmerTest* packages; Bates et al., 2015; Kuznetsova et al., 2017) with random intercept by participant to account for repeated measures within participants. Metabolite differences between exposed and unexposed children were tested with the formula:

$$Y_{ij} = B_1 \cdot p_i + B_2 \cdot a_{ij} + B_{0i} + \epsilon_{ij}$$

Where Y_{ij} = jth metabolite measurement for the ith subject, B_1 = coefficient for PAE status, p_i = the ith subject's PAE status (0 or 1), B_2 = coefficient for age, a_{ij} = ith subject's age at time of jth scan, B_{0i}= subject-specific y-intercept, and ϵ_{ij}= random error.

Age was included as a covariate in models in which a significant main effect of age on metabolite level was observed (tCho, tNAA, both cohorts). Tissue fraction was included as a covariate for LTP analyses only. Random effects were dropped from models in which random effects variance = 0 (Glx, both cohorts). False discovery rate (FDR; Benjamini & Hochberg, 1995) correction was applied separately in each cohort to correct for multiple comparisons (testing five metabolites); we report both uncorrected and corrected results.

Metabolite associations with EF and pre-reading skills

Associations between metabolites and cognitive scores were tested for metabolites that showed significant group differences, with separate models for each metabolite.

For Cohort 1 (ACC), associations between metabolites (tCho and Glx) and age-normed GEC T-scores and PAE-by-metabolite interactions were tested using linear models with the formula:

$$Y_i = B_1 \cdot p_i + B_2 \cdot a_i + B_3 \cdot m_i + (B_4 \cdot p_i \cdot m_i) + B_0 + \epsilon_i$$

Where Y_i = cognitive score for the ith subject, B_1 = coefficient for PAE status, p_i = the ith subject's PAE status (0 or 1), B_2 = coefficient for age, a_i = ith subject's age, B_3 = the coefficient for metabolite measurement, m_i = the ith subject's metabolite measurement, B_4 = the coefficient for the interaction between PAE status and metabolite measurement, B_0 = intercept, and ϵ_i = random error.

For Cohort 2 (LTP), associations between tCho and age-normed pre-reading scores (PP or SN) and PAE-by-metabolite interactions were tested using linear mixed effects models with the formula:

$$Y_i = B_1 \cdot p_i + B_2 \cdot a_{ij} + B_3 \cdot m_i + (B_4 \cdot p_{ij} \cdot m_{ij}) + B_{0i} + \epsilon_{ij}$$

Where Y_i = pre-reading score for the ith subject, B_1 = coefficient for PAE status, p_i = the ith subject's PAE status (0 or 1), B_2 = coefficient for age, a_{ij} = ith subject's age at time of jth scan, B_3 = the coefficient for metabolite measurement, m_{ij} = the ith subject's metabolite measurement (adjusted for tissue fraction by residualization) at time of jth scan, B_4 = the coefficient for the interaction between PAE status and metabolite measurement, B_{0i}= subject-specific y-intercept, and ϵ_{ij}= random error.

Data and code for analyses are available at: https://osf.io/4zkd8/?view_only=f7526d82205044feb07796410a937ca5

Results

Metabolite differences in PAE vs. unexposed children

Cohort 1: The PAE group had elevated levels of tCho (*beta* = .226, 95% CI [.04, .41], *p* = .016, p_{FDR} = .063) and Glx (*beta* = 1.474, 95% CI [.19, 2.76], *p* = .025, p_{FDR} = .063) in the ACC relative to unexposed children (Table 1, Figure 2), though these effects do not pass FDR correction for multiple comparisons. No significant group differences were found for tNAA, tCr or mI in the ACC. Measures of MRS data quality (Linewidth and SNR) differed significantly between groups, with children with PAE generally having lower quality data, but post-hoc analyses including linewidth and SNR as covariates in the models showed no significant effects of these metrics on tCho and Glx, and the pattern of elevated tCho and Glx in the PAE group was maintained in these models, see Supplementary Information for details (Table S1). Moreover, poorer data quality (greater linewidth and lower SNR) is expected to result in *lower* (rather than higher) estimates of metabolite concentrations, which is opposite to the observed pattern in our data, further supporting the robustness of these results.

Cohort 2: The PAE group had elevated levels of tCho (*beta* = .135, 95% CI [.02, .25], *p* = .02, p_{FDR} = .10) in the LTP relative to the unexposed group (Table 1, Figure 2), though this effect did not pass FDR correction for multiple comparisons. No significant group differences were found for tNAA, tCr, Glx or mI in the LTP.

Table 1. Descriptive statistics and group differences in metabolites, quality metrics, and cognitive skills.

Cohort 1: ACC	Unexposed Children			Prenatal Alcohol Exposure			Group Difference	
MRS Data	N (obs.)	Mean	SD	N (obs.)	Mean	SD	*beta*	*p*
Age at MRI (years)	76	4.51	0.95	21	5.34	1.12	**0.889**	**<.001**
tNAA (mmol/kg)	76	15.84	1.16	21	15.89	1.1	-0.179	0.543
tCr(mmol/kg)	76	12.8	1.1	21	12.83	1.16	0.095	0.757
tCho(mmol/kg)	75	2.58	0.35	21	2.73	0.38	**0.226**	**0.016**
Glx (mmol/kg)	76	27.02	2.57	21	28.46	2.52	**1.474**	**0.025**
mI (mmol/kg)	74	7.57	1.56	19	7.55	1.95	-.010	0.982
Tissue Fraction (GM/[GM+WM])	76	0.96	0.02	21	0.97	0.03	0.009	0.126
Linewidth (NAA)	76	7.39	3.04	21	10.25	3.01	**2.766**	**<.001**
SNR (NAA)	76	56.67	9.62	21	46.72	7.44	**-9.616**	**<.001**
Cognitive Assessment Data	N (obs.)	Mean	SD	N (obs.)	Mean	SD	*beta*	*p*
Age at Assessment (years)	25	4.37	0.76	17	5.54	1.15	**1.168**	**<.001**
NEPSY-II Statue Total Scaled Score	25	10.56	2.72	17	9.76	4.09	-0.795	0.453
BRIEF/BRIEF-P GEC T-score	22	49.23	11.56	16	73.44	17.29	**24.062**	**<.001**

Cohort 2: LTP	Unexposed Children			Prenatal Alcohol Exposure			Group Difference	
MRS Data	N (obs.)	Mean	SD	N (obs.)	Mean	SD	*beta*	*p*
Age at MRI (years)	128	6.45	0.93	28	6.45	0.76	-0.001	0.996
tNAA (mmol/kg)	128	15.2	0.91	28	14.93	1.13	-0.142	0.483
tCr(mmol/kg)	128	10.9	0.83	28	10.87	1.08	0.2	0.265
tCho(mmol/kg)	128	2.21	0.25	28	2.39	0.3	**0.135**	**0.02**
Glx (mmol/kg)	128	22.21	1.86	28	21.56	1.95	-0.099	0.769
mI (mmol/kg)	126	6.39	1.18	28	6.6	1.15	0.271	0.284
Tissue Fraction (GM/[GM+WM])	128	0.66	0.08	28	0.61	0.09	**-0.047**	**0.01**
Linewidth (NAA)	128	6.68	2.15	28	7.39	2.61	0.69	0.15
SNR (NAA)	128	72.52	13.24	28	69.32	12.79	-3.15	0.28
Cognitive Assessment Data	N (obs.)	Mean	SD	N (obs.)	Mean	SD	*beta*	*p*
Age at Assessment (years)	120	6.48	.93	24	6.37	.76	-.118	.560
NEPSY-II PP Total Scaled Score	120	11.78	2.77	24	10.96	2.27	-0.779	0.226
NEPSY-II SN Total Scaled Score	109	11.82	2.46	24	11.33	2.91	-0.382	0.523

Figure 2. Metabolite differences by alcohol exposure group. Significant group differences in metabolite levels in the ACC (Cohort 1; left and middle panels) and LTP (Cohort 2; right panel). Individual data points plotted (purple=unexposed, orange=PAE).

Executive functioning and pre-reading skill associations with PAE and metabolites

Cohort 1: EF and Metabolites in the ACC

A regression model examining the effects of PAE status and Glx on inhibitory control measured via the NEPSY-II Statue Total Scaled Score revealed a significant interaction between PAE status and Glx (*beta* = -.83, 95% CI[-1.55, -.11], *p*=.026). Post-hoc analysis indicated a negative association between Glx and inhibitory control in children with PAE, but not unexposed children (Figure 3A). No significant associations between inhibitory control and tCho were found.

A regression model examining the effects of PAE status and tCho on EF indexed by the BRIEF-II/BRIEF-P GEC T-score indicated that children with PAE had worse executive functioning relative to unexposed children (*beta* = 24.92, 95%CI[16.08, 33.76], *p*<.001), and lower tCho was associated with worse executive functioning in both groups (*beta*=-14.83, 95%CI[-28.54, -1.12], *p*=.035) (Table 1; Figure 3B). No significant group-by-metabolite interactions were found, and BRIEF/BRIEF-P GEC scores were not significantly associated with Glx.

Figure 3. Associations between EF measures and metabolites in ACC (Cohort 1). A) Inhibitory control was associated with Glx in the ACC (Cohort 1, n=42) in the PAE group (orange) but not the unexposed group (purple). Inhibitory control indexed by NEPSY-II Statue Total Scaled Score. B) Associations between PAE status, ACC tCho levels, and EF (Cohort 1, n=39). Fit lines represent negative association between tCho and EF within each group for unexposed (red) and PAE (yellow). EF indexed by BRIEF/BRIEF-P GEC T-score: higher scores indicate more difficulties in executive functioning.

Cohort 2: Pre-reading skills and metabolites in the LTP

Pre-reading skills (PP and SN) did not show any significant associations with tCho in the LTP, or alcohol exposure status. Groups performed similarly on PP and SN measures and group means were in the average to above-average range (Table 1).

Discussion

In this study, we report the first evidence of PAE-related brain metabolite differences in early childhood, showing elevated levels of tCho and Glx in children with PAE relative to unexposed children. These findings offer insight to neurochemical disruptions that correspond to differences in brain structure and function observed in PAE across childhood. Further, we show associations between brain metabolites and measures of executive function, one of the most commonly observed areas of cognitive/behavioral difficulty in individuals with PAE.

Metabolite differences in PAE vs. unexposed children

We found higher tCho in both the ACC and LTP of children with PAE compared to unexposed children, indicating widespread alterations in tissue microstructure and phospholipid metabolism across brain regions and tissue types (gray and white matter). Although the effects did not pass correction for multiple comparisons, the replication of this finding across two cohorts of children with PAE increases the robustness of this result. tCho levels are sensitive to the presence of all types of cell membranes and are associated with myelin content and fiber density (Perdue et al., *in prep*). The ACC voxel examined

in Cohort 1 included >90% gray matter, so our results likely reflect more complex cortical architecture in the children with PAE, possibly as a result of reduced/delayed pruning of synapses and more dense packing of neurites. In Cohort 2, the LTP voxel contained a mix of gray and white matter, and elevated tCho was present, even after accounting for individual differences in tissue composition of the MRS voxels. The tCho effects likely reflect differences in both gray and white matter microstructure, including neurite density and myelin content.

Notably, our finding of *higher* tCho in PAE groups contrasts with prior evidence that showed *lower* tCho in older children and adolescents with PAE. This may point to altered developmental trajectories of phospholipid metabolism, myelination, and cortical development in PAE, consistent with findings from structural brain imaging studies (Gimbel et al., 2023; Kar et al., 2022). In fact, no differences in tCho were found in an MRS study of neonates with PAE (Howells et al., 2016), suggesting that alterations in tCho may emerge later in postnatal development and change over childhood. We recently showed that tCho levels decrease in early childhood until about 9 years of age in unexposed children (Perdue et al., 2023), so elevated levels of tCho in early childhood in our cohorts with PAE likely reflect a developmental delay in the tCho decline that accompanies myelination and cortical refinement. Our present results show larger differences between tCho levels in Cohort 1, which includes children ages 3-6 years old, compared to Cohort 2, which includes children ages 5-8 years old. Although we cannot fully compare the two cohorts due to differences in brain regions examined, this suggests that elevated tCho observed in our study may be driven by younger children, and a reversal of this effect may occur in middle childhood when tCho levels tend to stabilize. This fits with studies in older children, where *reduced* tCho is observed in groups with PAE (Astley et al., 2009; du Plessis et al., 2014; Gonçalves et al., 2009). Longitudinal research tracking tCho across childhood in children with PAE is needed to confirm this hypothesis.

Glx was elevated in the ACC (Cohort 1), but not the LTP (Cohort 2). Given that glutamate is the primary excitatory neurotransmitter of the central nervous system and primary contributor to the Glx signal in MRS, over-abundance of Glx in the ACC is consistent with hypothesized hyperexcitability in fronto-striatal circuits in PAE (Bariselli & Lovinger, 2021). This hypothesis is supported by findings of altered functional activity in frontal brain regions, including the ACC, which are thought to be associated with EF difficulties in individuals with PAE (Fryer et al., 2007; Kodali et al., 2017; Ware et al., 2015). Concerning mechanism, higher measurements of Glx by MRS likely reflect an excess of excitatory synapses, indicating a delay or failure of synaptic pruning in the ACC and potential for hyperexcitability. Prior results for Glx in individuals with PAE have been somewhat mixed, with higher Glx in the cerebellum associated with PAE in older children (du Plessis et al., 2014), but lower Glx in parietal white matter in newborns with PAE (Howells et al., 2016). These discrepancies may reflect age- and/or tissue-dependent effects.

Associations between metabolites and executive functioning

Our results showing that increased Glx is associated with EF difficulties in children with PAE support the role of hyperexcitability in altered behavior and cognition in the context of PAE. In a subset of participants who completed the NEPSY-II Statue task, higher Glx (i.e., higher excitability) in the ACC was associated with poorer motor inhibitory control. This effect was specific to the PAE group, which indicates a particular role of Glx and hyperexcitability and inefficient

inhibitory control in children with PAE. Alterations in Glx may reflect dysfunction in the cycling of glutamate, glutamine, and the inhibitory neurotransmitter, GABA, which is needed to maintain the excitatory-inhibitory balance for efficient functioning (Bak et al., 2006; Sears & Hewett, 2021). The absence of such an effect in the unexposed group may indicate that unexposed children have more optimal Glx levels and intact glutamate-glutamine-GABA cycling. Thus, it may be that intact inhibitory systems and glutamate-glutamine-GABA cycling in unexposed individuals with relatively high Glx (similar levels to the PAE group) are better able to maintain the excitatory-inhibitory balance, resulting in no net effect of Glx on inhibitory control task performance. However, as the inhibitory neurotransmitter, GABA, was not measured in this study, nor in any prior human study of PAE, future research is needed to explicitly compare inhibitory systems in groups with and without PAE.

In addition to the association between Glx and inhibitory control, we observed a main effect of tCho on global EF indexed by the BRIEF/BRIEF-P GEC, such that higher tCho in the ACC was associated with better global EF. This finding was paired with a strong main effect of PAE status, such that children with PAE displayed more difficulties with EF relative to unexposed children. tCho levels in gray matter likely reflect packing density of cell bodies and neurites as well as ongoing phospholipid metabolism to maintain healthy tissue (Rae, 2014). Thus, higher tCho in the ACC may indicate that there are more neurons and synaptic connections available for top-down cognitive control, and that this architecture confers a benefit for EF within groups with and without PAE, despite overall EF and tCho differences between groups. This interpretation is supported by evidence that linked cortical morphometry of the ACC (greater surface area) to faster reaction time on a cognitive control task in typically developing children (Fjell et al., 2012), and another study that linked poorer inhibition to reduced ACC surface area in adolescents with PAE (Migliorini et al., 2015).

Together, the observed associations between metabolite levels in the ACC and EF suggest that PAE-related alterations in excitability, metabolism, and cortical microstructure are related to cognitive and behavioral difficulties in children with PAE. In contrast to EF, prereading skills (phonological processing or speeded naming) were spared in our sample (Cohort 2) and not associated with metabolite levels in the LTP. This is not surprising, as EF is known to be particularly impaired in PAE and findings related to language and reading skills in PAE have been mixed (Subramoney et al., 2018). Language and literacy impairments may depend on the amount and timing of alcohol exposure (O'Leary et al., 2013), and difficulties in this domain become apparent later in the acquisition of reading skills, so further research is needed to fully characterize PAE-related associations between brain metabolites and cognition.

Conclusion

Alcohol is known to disrupt multiple aspects of brain development and functioning. Here we show persistent effects of PAE on brain metabolites in early childhood, with implications for cognition and behavior. This evidence can support the development of interventions to modulate cortical hyperexcitability and improve EF in children with PAE. Future research combining multimodal neuroimaging techniques with MRS in PAE is needed to provide further insight to the metabolic mechanisms underlying altered brain structure and function in PAE.

References

Adams, J. W., Negraes, P. D., Truong, J., Tran, T., Szeto, R. A., Guerra, B. S., Herai, R. H., Teodorof-Diedrich, C., Spector, S. A., Del Campo, M., Jones, K. L., Muotri, A. R., & Trujillo, C. A. (2023). Impact of alcohol exposure on neural development and network formation in human cortical organoids. *Mol. Psychiatry*, *28*(4), 1571–1584.

Almeida, L., Andreu-Fernández, V., Navarro-Tapia, E., Aras-López, R., Serra-Delgado, M., Martínez, L., García-Algar, O., & Gómez-Roig, M. D. (2020). Murine models for the study of fetal alcohol spectrum disorders: An overview. *Front. Pediatr.*, *8*, 359.

Ashburner, J., & Friston, K. J. (2005). Unified segmentation. *NeuroImage*, *26*(3), 839–851. https://doi.org/10.1016/J.NEUROIMAGE.2005.02.018

Astley, S. J., Richards, T., Aylward, E. H., Olson, H. C., Kerns, K., Brooks, A., Coggins, T. E., Davies, J., Dorn, S., Gendler, B., Jirikowic, T., Kraegel, P., & Maravilla, K. (2009). Magnetic resonance spectroscopy outcomes from a comprehensive magnetic resonance study of children with fetal alcohol spectrum disorders. *Magnetic Resonance Imaging*, *27*(6), 760–778. https://doi.org/10.1016/j.mri.2009.01.003

Baeshen, A., Wyss, P. O., Henning, A., O'Gorman, R. L., Piccirelli, M., Kollias, S., & Michels, L. (2020). Test–Retest Reliability of the Brain Metabolites GABA and Glx With JPRESS, PRESS, and MEGA-PRESS MRS Sequences in vivo at 3T. *Journal of Magnetic Resonance Imaging*, *51*(4), 1181–1191. https://doi.org/10.1002/JMRI.26921

Bak, L. K., Schousboe, A., & Waagepetersen, H. S. (2006). The glutamate/GABA-glutamine cycle: Aspects of transport, neurotransmitter homeostasis and ammonia transfer. *Journal of Neurochemistry*, *98*(3), 641–653. https://doi.org/10.1111/j.1471-4159.2006.03913.x

Bariselli, S., & Lovinger, D. M. (2021). Corticostriatal circuit models of cognitive impairments induced by fetal exposure to alcohol. *Biological Psychiatry*, *90*(8), 516–528. https://doi.org/10.1016/j.biopsych.2021.05.014

Basavarajappa, B. S., & Subbanna, S. (2023). Synaptic plasticity abnormalities in fetal alcohol spectrum disorders. *Cells*, *12*(3), 442.

Bates, D., Mächler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, *067*(i01). https://doi.org/10.18637/JSS.V067.I01

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society*, *57*(1), 289–300.

Cantacorps, L., Alfonso-Loeches, S., Moscoso-Castro, M., Cuitavi, J., Gracia-Rubio, I., López-Arnau, R., Escubedo, E., Guerri, C., & Valverde, O. (2017). Maternal alcohol binge drinking induces persistent neuroinflammation associated with myelin damage and behavioural dysfunctions in offspring mice. *Neuropharmacology*, *123*, 368–384.

Cook, J. L., Green, C. R., Lilley, C. M., Anderson, S. M., Baldwin, M., Chudley, A. E., Conry, J. L., LeBlanc, N., Loock, C. A.,

Lutke, J., Mallon, B. F., McFarlane, A. A., Temple, V. K., & Rosales, T. (2016). Fetal alcohol spectrum disorder: Reconsidering blame. *CMAJ*, *188*(3), 191–197. https://doi.org/10.1503/cmaj.151425

Cui, Z.-J., Zhao, K.-B., Zhao, H.-J., Yu, D.-M., Niu, Y.-L., Zhang, J.-S., & Deng, J.-B. (2010). Prenatal alcohol exposure induces long-term changes in dendritic spines and synapses in the mouse visual cortex. *Alcohol Alcohol*, *45*(4), 312–319.

Darbinian, N., C. Sparks, E., Darbinyan, A., Merabova, N., Tatevosian, G., Vadachkoria, E., Zhao, H., Amini, S., Goetzl, L., & E. Selzer, M. (2023). Maternal blood lipid biomarkers of oligodendrocyte pathology to predict fetal alcohol spectrum disorders. *Obstetrics and Gynecology Research*, *06*(02), 127–138. https://doi.org/10.26502/ogr0122

Darbinian, N., & Selzer, M. (2022). Oligodendrocyte pathology in fetal alcohol spectrum disorders. *Neural Regeneration Research*, *17*(3), 497–502. https://doi.org/10.4103/1673-5374.314294

DeMayo, M. M., McGirr, A., Selby, B., MacMaster, F. P., Debert, C. T., & Harris, A. D. (2023). Consistency of frontal cortex metabolites quantified by magnetic resonance spectroscopy within overlapping small and large voxels. *Scientific Reports*, *13*(2246), 1–14. https://doi.org/10.1038/s41598-023-29190-y

Donald, K. A., Eastman, E., Howells, F. M., Adnams, C., Riley, E. P., Woods, R. P., Narr, K. L., & Stein, D. J. (2015). Neuroimaging effects of prenatal alcohol exposure on the developing human brain: A magnetic resonance imaging review. *Acta Neuropsychiatrica*, *27*(5), 251–269. https://doi.org/10.1017/neu.2015.12

Donald, K. A., Hendrikse, C. J., Roos, A., Wedderburn, C. J., Subramoney, S., Ringshaw, J. E., Bradford, L., Hoffman, N., Burd, T., Narr, K. L., Woods, R. P., Zar, H. J., Joshi, S. H., & Stein, D. J. (2024). Prenatal alcohol exposure and white matter microstructural changes across the first 6-7 years of life: A longitudinal diffusion tensor imaging study of a South African birth cohort. *NeuroImage Clin.*, *103572*, 103572.

du Plessis, L., Jacobson, J. L., Jacobson, S. W., Hess, A. T., van der Kouwe, A., Avison, M. J., Molteno, C. D., Stanton, M. E., Stanley, J. A., Peterson, B. S., & Meintjes, E. M. (2014). An in vivo 1H magnetic resonance spectroscopy study of the deep cerebellar nuclei in children with fetal alcohol spectrum disorders. *Alcoholism: Clinical and Experimental Research*, *38*(5), 1330–1338. https://doi.org/10.1111/acer.12380

Fayed, N., Modrego, P. J., & Medrano, J. (2009). Comparative test-retest reliability of metabolite values assessed with magnetic resonance spectroscopy of the brain. The LCModel versus the manufacturer software. *Neurological Research*, *31*(5), 472–477. https://doi.org/10.1179/174313209X395481

Fjell, A. M., Walhovd, K. B., Brown, T. T., Kuperman, J. M., Chung, Y., Hagler, D. J., Venkatraman, V., Cooper Roddey, J., Erhart, M., McCabe, C., Akshoomoff, N., Amaral, D. G., Bloss, C. S., Libiger, O., Darst, B. F., Schork, N. J., Casey, B. J., Chang, L., Ernst, T. M., … Dale, A. M. (2012). Multimodal imaging of the self-regulating developing brain. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(48), 19620–19625. https://doi.org/10.1073/pnas.1208243109

Fryer, S. L., Tapert, S. F., Mattson, S. N., Paulus, M. P., Spadoni, A. D., & Riley, E. P. (2007). Prenatal alcohol exposure

affects frontal-striatal BOLD response during inhibitory control. *Alcoholism: Clinical and Experimental Research*, *31*(8), 1415–1424. https://doi.org/10.1111/j.1530-0277.2007.00443.x

Gasparovic, C., Bedrick, E. J., Mayer, A. R., Yeo, R. A., Chen, H., Damaraju, E., Calhoun, V. D., & Jung, R. E. (2011). Test-retest reliability and reproducibility of short-echo-time spectroscopic imaging of human brain at 3T. *Magnetic Resonance in Medicine*, *66*(2), 324–332. https://doi.org/10.1002/mrm.22858

Gasparovic, C., Chen, H., & Mullins, P. G. (2018). Errors in 1H-MRS estimates of brain metabolite concentrations caused by failing to take into account tissue-specific signal relaxation. *NMR in Biomedicine*, *31*(6), e3914. https://doi.org/10.1002/NBM.3914

Gimbel, B. A., Roediger, D. J., Ernst, A. M., Anthony, M. E., de Water, E., Mueller, B. A., Rockhold, M. N., Schumacher, M. J., Mattson, S. N., Jones, K. L., Lim, K. O., & Wozniak, J. R. (2023). Delayed cortical thinning in children and adolescents with prenatal alcohol exposure. *Alcohol: Clinical and Experimental Research*, *47*(7), 1312–1326. https://doi.org/10.1111/acer.15096

Gioia, G. A., Andrews Espy, K., & Isquith, P. K. (2003). *Behavior Rating Inventory of Executive Function - Preschool Version*. Pearson Assessment Resources.

Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). *Behavior Rating Inventory of Executive Function*. Pearson Assessment Resources.

Gonçalves, R. D. C. F., Vasconcelos, M. M., Faleiros, L. O., Da Cruz, L. C. H., Domingues, R. C., Brito, A. R., Werner, J., & Herdy, G. V. H. (2009). Proton magnetic resonance spectroscopy in children with fetal alcohol spectrum disorders. *Arquivos de Neuro-Psiquiatria*, *67*(2 A), 254–261. https://doi.org/10.1590/S0004-282X2009000200015

Harris, A. D., Puts, N. A. J., & Edden, R. A. E. (2015). Tissue correction for GABA-edited MRS: Considerations of voxel composition, tissue segmentation, and tissue relaxations. *Journal of Magnetic Resonance Imaging : JMRI*, *42*(5), 1431–1440. https://doi.org/10.1002/JMRI.24903

Howells, F. M., Donald, K. A., Roos, A., Woods, R. P., Zar, H. J., Narr, K. L., & Stein, D. J. (2016). Reduced glutamate in white matter of male neonates exposed to alcohol in utero: a 1H-magnetic resonance spectroscopy study. *Metabolic Brain Disease*, *31*(5), 1105–1112. https://doi.org/10.1007/s11011-016-9850-x

Hwang, H. M., Kawasawa, Y. I., Basha, A., Mohammad, S., Ito, M., & Hashimoto-Torii, K. (2023). Fatty acid metabolism changes in association with neurobehavioral deficits in animal models of fetal alcohol spectrum disorders. *Communications Biology*, *6*(1), 1–12. https://doi.org/10.1038/s42003-023-05127-z

Kar, P., Reynolds, J. E., Gibbard, W. Ben, McMorris, C., Tortorelli, C., & Lebel, C. (2022). Trajectories of brain white matter development in young children with prenatal alcohol exposure. *Human Brain Mapping*, *43*(13), 4145–4157. https://doi.org/10.1002/hbm.25944

Kar, P., Reynolds, J. E., Grohs, M. N., Gibbard, W. Ben, McMorris, C., Tortorelli, C., & Lebel, C. (2021). White matter

alterations in young children with prenatal alcohol exposure. *Developmental Neurobiology*, *81*(4), 400–410. https://doi.org/10.1002/dneu.22821

Khoury, J. E., Milligan, K., & Girard, T. A. (2015). Executive functioning in children and adolescents prenatally exposed to alcohol: A meta-analytic review. *Neuropsychol. Rev.*, *25*(2), 149–170.

Kodali, V. N., Jacobson, J. L., Lindinger, N. M., Dodge, N. C., Molteno, C. D., Meintjes, E. M., & Jacobson, S. W. (2017). Differential recruitment of brain regions during response inhibition in children prenatally exposed to zlcohol. *Alcoholism: Clinical and Experimental Research*, *41*(2), 334–344. https://doi.org/10.1111/acer.13307

Kodituwakku, P. W., Kalberg, W., & May, P. A. (2001). The effects of prenatal alcohol exposure on executive functioning. *Alcohol Research & Health*, *25*(3), 192–198.

Kodituwakku, P. W., Segall, J. M., & Beatty, G. K. (2011). Cognitive and behavioral effects of prenatal alcohol exposure. *Future Neurology*, *6*(2), 237–259. https://doi.org/10.2217/fnl.11.4

Korkman, M., Kirk, U., & Kemp, S. (2007). *NEPSY - Second Edition (NEPSY-II)*. Harcourt Assessment.

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13), 1–26. https://doi.org/10.18637/JSS.V082.I13

Lebel, C., Roussotte, F., & Sowell, E. R. (2011). Imaging the impact of prenatal alcohol exposure on the structure of the developing human brain. *Neuropsychology Review*, *21*(2), 102–118. https://doi.org/10.1007/s11065-011-9163-0

Lee, J., Naik, V., Orzabal, M., Lunde-Young, R., & Ramadoss, J. (2021). Morphological alteration in rat hippocampal neuronal dendrites following chronic binge prenatal alcohol exposure. *Brain Res.*, *1768*(147587), 147587.

Long, M., Kar, P., Forkert, N. D., Landman, B. A., Gibbard, W. Ben, Tortorelli, C., Mcmorris, C. A., Lebel, C. A., & Louth, E. L. (2024). Sex and age effects on gray matter volume trajectories in young children with prenatal alcohol exposure. *Frontiers in Human Neuroscience*, *April*, 1–16. https://doi.org/10.3389/fnhum.2024.1379959

Long, X., Kar, P., Gibbard, B., Tortorelli, C., & Lebel, C. (2019). The brain's functional connectome in young children with prenatal alcohol exposure. *NeuroImage: Clinical*, *24*(June), 0–7. https://doi.org/10.1016/j.nicl.2019.102082

Long, X., & Lebel, C. (2022). Evaluation of brain alterations and behavior in children with low levels of prenatal alcohol exposure. *JAMA Network Open*, *5*(4), E225972. https://doi.org/10.1001/jamanetworkopen.2022.5972

Long, X., Little, G., Treit, S., Beaulieu, C., Gong, G., & Lebel, C. (2020). Altered brain white matter connectome in children and adolescents with prenatal alcohol exposure. *Brain Structure and Function*, *225*(3), 1123–1133. https://doi.org/10.1007/s00429-020-02064-z

Mattson, S. N., Bernes, G. A., & Doyle, L. R. (2019). Fetal alcohol spectrum disorders: A review of the neurobehavioral deficits associated with prenatal alcohol exposure. *Alcoholism: Clinical and Experimental Research*, *43*(6), 1046–1062. https://doi.org/10.1111/acer.14040

Mattson, S. N., & Riley, E. P. (1999). Implicit and explicit memory functioning in children with heavy prenatal alcohol exposure. *Journal of the International Neuropsychological Society*, *5*(5), 462–471. https://doi.org/10.1017/S1355617799555082

Migliorini, R., Moore, E. M., Glass, L., Infante, M. A., Tapert, S. F., Jones, K. L., Mattson, S. N., & Riley, E. P. (2015). Anterior cingulate cortex surface area relates to behavioral inhibition in adolescents with and without heavy prenatal alcohol exposure. *Behavioural Brain Research*, *292*, 26–35. https://doi.org/10.1016/j.bbr.2015.05.037

Millard, S. P. (2013). *EnvStats: An R Package for Environmental Statistics*. Springer. https://www.springer.com

Moore, E. M., & Xia, Y. (2022). Neurodevelopmental trajectories following prenatal alcohol exposure. *Frontiers in Human Neuroscience*, *15*(January), 1–11. https://doi.org/10.3389/fnhum.2021.695855

Nardelli, A., Lebel, C., Rasmussen, C., Andrew, G., & Beaulieu, C. (2011). Extensive deep gray matter volume reductions in children and adolescents with fetal alcohol spectrum disorders. *Alcohol. Clin. Exp. Res.*, *35*(8), 1404–1417.

Near, J., Harris, A. D., Juchem, C., Kreis, R., Marjańska, M., Öz, G., Slotboom, J., Wilson, M., & Gasparovic, C. (2020). Preprocessing, analysis and quantification in single-voxel magnetic resonance spectroscopy: experts' consensus recommendations. *NMR in Biomedicine*, *34*(5), 1–23. https://doi.org/10.1002/nbm.4257

O'Leary, C. M., Taylor, C., Zubrick, S. R., Kurinczuk, J. J., & Bower, C. (2013). Prenatal alcohol exposure and educational achievement in children aged 8-9 years. *Pediatrics*, *132*(2). https://doi.org/10.1542/peds.2012-3002

Perdue, M. V., DeMayo, M. M., Bell, T. K., Boudes, E., Bagshawe, M., Harris, A. D., & Lebel, C. (2023). Changes in brain metabolite levels across childhood. *NeuroImage*, *274*, 120087. https://doi.org/10.1016/j.neuroimage.2023.120087

Perdue, M. V., DeMayo, M. M., Bell, T. K., Warriyar K.V., V., Alsop, D., Varma, G., Lebel, R.M., Harris, A.D., & Lebel, C. (in prep). Associations between brain metabolites and tissue microstructure in children: A multimodal imaging study

Pinner, J. F. L., Collishaw, W., Schendel, M. E., Flynn, L., Candelaria-Cook, F. T., Cerros, C. M., Williams, M., Hill, D. E., & Stephen, J. M. (2023). Examining the effects of prenatal alcohol exposure on performance of the sustained attention to response task in children with an FASD. *Human Brain Mapping*, *44*(17), 6120–6138. https://doi.org/10.1002/hbm.26501

Popova, S., Charness, M. E., Burd, L., Crawford, A., Hoyme, H. E., Mukherjee, R. A. S., Riley, E. P., & Elliott, E. J. (2023). Fetal alcohol spectrum disorders. *Nature Reviews Disease Primers*, *9*(1), 1–21. https://doi.org/10.1038/s41572-023-00420-x

Provencher, S. W. (1993). Estimation of metabolite concentrations from localized in vivo proton NMR spectra. *Magnetic Resonance in Medicine*, *30*(6), 672–679. https://doi.org/10.1002/MRM.1910300604

Rae, C. D. (2014). A guide to the metabolic pathways and function of metabolites observed in human brain 1 H magnetic resonance spectra. *Neurochem Res*, *39*, 1–36. https://doi.org/10.1007/s11064-013-1199-5

Rasmussen, C., & Bisanz, J. (2009). Executive functioning in children with Fetal Alcohol Spectrum Disorders: profiles and age-related differences. *Child Neuropsychol.*, *15*(3), 201–215.

Rasmussen, C., Mcauley, R., Andrew, G., & Hospital, G. R. (2007). Parental Ratings of Children With Fetal Alcohol Spectrum. *Children*, *5*, 1–8.

Rasmussen, C., Tamana, S., Baugh, L., Andrew, G., Tough, S., & Zwaigenbaum, L. (2013). Neuropsychological impairments on the NEPSY-II among children with FASD. *Child Neuropsychology*, *19*(4), 337–349. https://doi.org/10.1080/09297049.2012.658768

Reynolds, J. E., Long, X., Paniukov, D., Bagshawe, M., & Lebel, C. (2020). Calgary Preschool magnetic resonance imaging (MRI) dataset. *Data in Brief*, *29*, 105224. https://doi.org/10.1016/J.DIB.2020.105224

Rockhold, M. N., Krueger, A. M., de Water, E., Lindgren, C. W., Sandness, K. E., Eckerle, J. K., Schumacher, M. J., Fink, B. A., Boys, C. J., Carlson, S. M., Fuglestad, A. J., Mattson, S. N., Jones, K. L., Riley, E. P., & Wozniak, J. R. (2021). Executive and social functioning across development in children and adolescents with prenatal alcohol exposure. *Alcoholism: Clinical and Experimental Research*, *45*(2), 457–469. https://doi.org/10.1111/acer.14538

Roos, A., Fouche, J.-P., Ipser, J. C., Narr, K. L., Woods, R. P., Zar, H. J., Stein, D. J., & Donald, K. A. (2021). Structural and functional brain network alterations in prenatal alcohol exposed neonates. *Brain Imaging Behav.*, *15*(2), 689–699.

RStudio Team. (2020). *RStudio: Integrated Development Environment for R*. http://www.rstudio.com/

Sears, S. M. S., & Hewett, S. J. (2021). Influence of glutamate and GABA transport on brain excitatory/inhibitory balance. *Experimental Biology and Medicine*, *246*(9), 1069–1083. https://doi.org/10.1177/1535370221989263

Simpson, R., Devenyi, G. A., Jezzard, P., Hennessy, T. J., & Near, J. (2017). Advanced processing and simulation of MRS data using the FID appliance (FID-A)—An open source, MATLAB-based toolkit. *Magnetic Resonance in Medicine*, *77*(1), 23–33. https://doi.org/10.1002/MRM.26091

Soreni, N., Noseworthy, M. D., Konyer, N. B., Pullenayegum, E., & Schachar, R. (2010). Interindividual, repositioning, and time-of-day effects on single voxel proton MR spectroscopy of the anterior cingulate cortex. *Journal of Magnetic Resonance Imaging*, *32*(2), 276–282. https://doi.org/10.1002/JMRI.22264

Subramoney, S., Eastman, E., Adnams, C., Stein, D. J., & Donald, K. A. (2018). The early developmental outcomes of prenatal alcohol exposure: A review. *Front. Neurol.*, *9*, 1108.

Subramoney, S., Joshi, S. H., Wedderburn, C. J., Lee, D., Roos, A., Woods, R. P., Zar, H. J., Narr, K. L., Stein, D. J., & Donald, K. A. (2022). The impact of prenatal alcohol exposure on gray matter volume and cortical surface area of 2 to 3-year-old children in a South African birth cohort. *Alcoholism: Clinical and Experimental Research*, *46*(7), 1233–1247. https://doi.org/10.1111/acer.14873

Team, R. C. (2016). *R: A language and environment for statistical computing*. R Foundation for Statistical Computing.

Veazey, K. J., Muller, D., & Golding, M. C. (2013). Prenatal alcohol exposure and cellular differentiation. *Alcohol Research: Current Reviews*, *35*(1), 77–85. https://ahs.idm.oclc.org/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=ccm&AN=90403011&site=ehost-live&scope=site

Volk, C., Jaramillo, V., Merki, R., Tuura, G., & Huber, R. (2018). Diurnal changes in glutamate & glutamine levels of healthy young adults assessed by proton magnetic resonance spectroscopy. *Human Brain Mapping*, *39*, 3984–3992. https://doi.org/10.1002/hbm.24225

Volk, C., Jaramillo, V., Studler, M., Furrer, M., O'gorman Tuura, R. L., & Huber, R. (2019). Diurnal changes in human brain glutamate þ glutamine levels in the course of development and their relationship to sleep. *NeuroImage*, *196*, 269–275. https://doi.org/10.1016/j.neuroimage.2019.04.040

Ware, A. L., Infante, M. A., O'Brien, J. W., Tapert, S. F., Jones, K. L., Riley, E. P., & Mattson, S. N. (2015). An fMRI study of behavioral response inhibition in adolescents with and without histories of heavy prenatal alcohol exposure. *Behavioural Brain Research*, *278*, 137–146. https://doi.org/10.1016/j.bbr.2014.09.037

Wozniak, J. R., Mueller, B. A., Bell, C. J., Muetzel, R. L., Hoecker, H. L., Boys, C. J., & Lim, K. O. (2013). Global functional connectivity abnormalities in children with fetal alcohol spectrum disorders. *Alcohol. Clin. Exp. Res.*, *37*(5), 748–756.

Supplementary Information

Table S1. Model summaries for ACC (Cohort 1) including MRS quality metrics (SNR and linewidth [LW] of the NAA peak) as covariates.

Predictors	tCho (molal)			Glx (molal)		
	Estimates	*CI*	*p*	*Estimates*	*CI*	*p*
(Intercept)	2.52	1.77 – 3.26	**<0.001**	26.04	21.01 – 31.08	**<0.001**
PAE	0.27	0.08 – 0.47	**0.007**	1.42	-0.01 – 2.85	0.051
SNR (NAA)	0.01	-0.00 – 0.02	0.183	0.01	-0.06 – 0.08	0.772
LW (NAA)	0.01	-0.02 – 0.04	0.690	0.05	-0.16 – 0.26	0.608
Age (years)	-0.08	-0.15 – -0.01	**0.036**			
N	89 subj_id			90 subj_id		
Observations	96			97		