Title

STOP1 dominates Arabidopsis tolerance to ammonium over NRT1.1/NPF6.3/CHL1

Running title

STOP1 overrides NRT1.1 in NH_4^+ tolerance

Authors and addresses

Takushi Hachiya[a,b,†], Mako Sakai[a], Tsuyoshi Nakagawa[a], and Hitoshi Sakakibara[b,c]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for

Science Research, Shimane University, Matsue 690-8504, Japan

[b]RIKEN Center for Sustainable Resource Science, Yokohama 230-0045, Japan

[c]Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8601,

Japan

Corresponding Author

Takushi Hachiya

Department of Molecular and Functional Genomics, Interdisciplinary Center for

Science Research, Shimane University, Matsue 690-8504, Japan

Email: takushi.hachiya@life.shimane-u.ac.jp, Tel: +81-852-32-6288

Type of contribution

Short paper

Division

Plant Nutrition

Abstract

The Arabidopsis nitrate transceptor NITRATE TRANSPORTER 1.1

(NRT1.1/NPF6.3/CHL1) plays significant roles even in the absence of nitrate. The

loss-of-function of NRT1.1 alleviates growth suppression and chlorosis under toxic

levels of ammonium as the sole nitrogen source. Recently, we reported that acidic

stress-inducible genes are downregulated by *NRT1.1* deficiency under ammonium

toxicity conditions, implying that NRT1.1 may exacerbate ammonium-dependent acidic

stress. The transcription factor SENSITIVE TO PROTON RHIZOTOXICITY 1

(STOP1) enhances Arabidopsis tolerance to ammonium and acidic stresses.

Furthermore, STOP1 directly activates *NRT1.1* transcription. These previous findings

prompted us to analyze the association between NRT1.1 and STOP1 on ammonium

tolerance. Here, we show that ammonium-inducible STOP1 plays a dominant role over

NRT1.1 in ammonium tolerance and expression of direct target genes of STOP1. We

present a novel scheme in which NRT1.1 and STOP1 jointly regulate common target(s)

associated with ammonium tolerance.

Keywords: ammonium toxicity, CHL1, NPF6.3, NRT1.1, STOP1

1. Introduction

Nitrate serves not only as a substrate for nitrogen (N) assimilation but also as a signaling molecule. The Arabidopsis NITRATE TRANSPORTER 1.1 (NRT1.1/NPF6.3/CHL1) acts as a nitrate sensor and orchestrates various plant responses to the nitrate signal, including gene expression and root system development (Bouguyon et al. 2015; Ho et al. 2009). Apart from its nitrate-dependent functions, NRT1.1 exhibits significant roles in the absence of nitrate (Hachiya et al. 2024). Previous studies have shown that the loss-of-function of NRT1.1 alleviates growth suppression and chlorosis under toxic levels of ammonium as the sole N source, i.e., ammonium toxicity (Hachiya et al. 2011; Jian et al. 2018; Liu et al. 2020). Recently, we reported that in early seedlings grown under 10 mM ammonium, the genes induced by acidic stress are downregulated in *NRT1.1*-deficient mutants relative to the wild-type, whereas those repressed by acidic stress are upregulated in the mutants (Hachiya et al. 2024). Moreover, an increase in medium pH promotes shoot growth of the wild-type more intensely than that of the *NRT1.1*-deficient mutant under 10 mM ammonium. Given that acidic stress is one of the primary causes of ammonium toxicity (Hachiya et al. 2021), NRT1.1 may exacerbate the ammonium-dependent acidic stress.

In *Arabidopsis*, the zinc finger transcription factor SENSITIVE TO PROTON RHIZOTOXICITY 1 (STOP1) positively regulates genes involved in the tolerance to ammonium and acidic stresses, as well as abiotic stresses, including aluminum, drought, salt, low phosphate, and low oxygen (Balzergue et al. 2017; Enomoto et al. 2019; Hachiya et al. 2021; Sadhukhan et al. 2019; Sawaki et al. 2009; Tian et al. 2021).

Furthermore, STOP1 directly activates *NRT1.1* expression under an acidic rhizosphere

containing nitrate as the primary N source (Ye et al. 2021). These findings prompted us

to analyze the association between NRT1.1 and STOP1 under toxic levels of

ammonium as the sole N source. Here, we show that STOP1 dominates ammonium

tolerance and expression of direct target genes of STOP1 over NRT1.1.

2. Materials and Methods

2.1. Plant materials and growth conditions

Arabidopsis thaliana accession Columbia (Col-0) was used as a control. T-DNA

insertion mutants *nrt1.1* (SALK_097431, Hachiya et al. 2011), *stop1^{KO}* (SALK_114108,

Sawaki et al. 2007), and gamma ray-mutagenized mutant c*hl1-5* (Tsay et al. 1993) were

obtained from the European Arabidopsis Stock Center. *stop1^{KO}* was backcrossed twice

with Col-0 before use. The line harboring *pSTOP1:GFP-STOP1* in the *stop1^{KO}*

background (Balzergue et al. 2017) was provided by Dr. Thierry Desnos (Aix-Marseille

University, France).

The seeds were surface sterilized, placed in plastic Petri dishes (diameter: 90 mm;

depth: 20 mm; Iwaki, Tokyo, Japan), and grown on a solid medium containing 30 mL

of Murashige and Skoog (MS) salts without N sources supplemented with 4.7 mM

MES-KOH (pH 5.7), 2% (w/v) sucrose, and 0.25% (w/v) gellan gum (Wako, Osaka,

Japan) with or without 100 nM indole-3-acetic acid (IAA). Two different N and K

sources were used: 10 mM KNO_3 (10 mM NO_3^- condition) or 5 mM $(NH_4)_2SO_4$ with 10

mM KCl (10 mM NH_4^+ condition). The plants were kept in the dark at 4°C for 3 days,

and then grown horizontally under a photosynthetic photon flux density of 100–130

µmol m^{-2} s^{-1} (16-h light/8-h dark cycle) at 23°C.

2.2. Extraction of RNA

Frozen samples were ground using TissueLyser II (Qiagen, Tokyo, Japan) using

zirconia beads. Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen),

following the manufacturer's instructions.

2.3. Quantitative reverse transcription polymerase chain reaction

Reverse transcription (RT) was conducted using ReverTra Ace qPCR RT Master Mix

with gDNA Remover (Toyobo, Tokyo, Japan), following the manufacturer's

instructions. The obtained cDNA was diluted 10-fold with distilled water for

quantitative PCR (qPCR). Transcript levels were evaluated using StepOnePlus

Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA, USA). The diluted

cDNA (2 µL) was amplified in the presence of 10 µL of KAPA SYBR FAST qPCR Kit

(Nippon Genetics, Tokyo, Japan), 0.4 µL of specific primers (0.2 µM final

concentration), and 7.2 µL of sterile water. Relative transcript levels were calculated

using the comparative cycle threshold method with *ACTIN3* as an internal standard.

Primer sequences are shown in Supplementary Table 1.

2.4. Observation of fluorescent signal

Confocal imaging was performed using a Leica SP5 Confocal Microscope (Leica

Microsystems, Wetzlar, Germany). GFP-STOP1 was excited with a 488-nm Ar laser,

and emission was detected from 500–530 nm using a Leica 25× objective (HCX IRAPO

L 25×/0.95 WATER).

3. Results and Discussion

3.1. STOP1 dominates ammonium tolerance over NRT1.1

To analyze the genetic relationship between *NRT1.1* and *STOP1*, *chl1-5* and *nrt1.1* were

crossed with *stop1^{KO}*, and their double homozygous mutants were obtained (Figure 1a).

The growth of Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}* under

10 mM ammonium supplemented with 100 nM IAA was compared. It should be noted

that appropriate IAA concentrations stabilize the ammonium tolerance of

NRT1.1-deficient mutants (Hachiya and Noguchi 2011). The shoots of *chl1-5* and *nrt1.1*

were larger and greener than those of Col-0 (Figure 1b). The shoot growth enhancement

in *NRT1.1*-deficient mutants was almost completely suppressed by *STOP1* deficiency.

Shoot fresh weights of *chl1-5* and *nrt1.1* were much larger than those of Col-0, whereas

those of *stop1^{KO}* were slightly smaller (Figure 1c). Fresh weights of the shoots of *chl1-5*

stop1^{KO} and *nrt1.1 stop1^{KO}* were comparable to those of *stop1^{KO}*. Also, under 10 mM

ammonium without IAA, the shoot fresh weights were increased by *NRT1.1* deficiency,

but they were reduced by *STOP1* deficiency to a level comparable to those of *stop1^{KO}*

(Figure 1d). Under 10 mM nitrate without IAA, the shoot fresh weights of

NRT1.1-deficient mutants were significantly smaller than those of other lines with or

without *STOP1*. Under both N conditions, the fresh weights of the root showed a similar

tendency to those of the shoot. Thus, our genetic analysis indicates that STOP1 is

epistatic to NRT1.1 regarding ammonium tolerance.

3.2 NRT1.1 does not downregulate* STOP1*/STOP1 and STOP1 dominates expression

of direct target genes of STOP1 over NRT1.1

The above growth analysis suggests that NRT1.1 could act upstream of STOP1 and

downregulate *STOP1*/STOP1 activities under ammonium. To verify this, we checked

the transcript levels of known direct target genes of STOP1, i.e.,

ALUMINUM-ACTIVATED MALATE TRANSPORTER 1 (*ALMT1*), *CALCINEURIN*

B-LIKE INTERACTING PROTEIN KINASE 23 (*CIPK23*), *GLUTAMATE*

DEHYDROGENASE 2 (*GDH2*), and *SENSITIVE TO PROTON RHIZOTOXICITY 2*

(*STOP2*) (Balzergue et al. 2017; Enomoto et al. 2019; Sadhukhan et al. 2019, Tokizawa

et al. 2021) in the absence of IAA. In five-day-old Col-0 shoots and roots, the

expression of *ALMT1*, *CIPK23*, *GDH2*, and *STOP2* was significantly higher under 10

mM ammonium than under 10 mM nitrate (Figure 2a). However, this

ammonium-dependent transcriptional induction was consistently decreased by *NRT1.1*

deficiency (Figure 2a). Also, *STOP1* expression in shoots and roots was not

significantly increased by *NRT1.1* deficiency under ammonium (Figure 2a). Moreover,

our microarray analysis using three- and five-day-old seedlings under 10 mM

ammonium without IAA (Hachiya et al. 2024) revealed that *NRT1.1* deficiency did not

generally induce genes whose expression is upregulated via STOP1 under acidic pH

(Sawaki et al. 2009) (Figure 2b and Supplementary Table S2). It is widely accepted that

the F-box protein REGULATION OF ATALMT1 EXPRESSION 1 (RAE1) interacts

with STOP1 and promotes its degradation via the ubiquitin-26S proteasome pathway

(Zhang et al. 2019), i.e., the post-translational regulation of STOP1 activity. Recent

studies have shown that aluminum stress induces phosphorylation of STOP1 by MAP

KINASE 4 and oxidative modification of RAE1 by reactive oxygen species (Zhou et al.

2023; Ding et al. 2024). These modifications inhibit RAE1-dependent proteolysis of

STOP1, thereby enhancing aluminum tolerance. Thus, to analyze the effect of NRT1.1

on STOP1 expression, *chl1-5* was crossed with *pSTOP1:GFP-STOP1* line in the

stop1^{KO} background (Balzergue et al. 2017) and produced the *pSTOP1:GFP-STOP1*

line in the *chl1-5 stop1^{KO}* background. The GFP-STOP1-derived fluorescence signal

was promoted in the roots under 10 mM ammonium compared to 10 mM nitrate (Figure

2c), similar to a previous result (Tian et al. 2021). However, the GFP signal was little

changed by *NRT1.1* deficiency. Meanwhile, in five-day-old seedlings grown under 10

mM ammonium, the transcript levels of *ALMT1*, *CIPK23*, and *GDH2* were similar

among *stop1^{KO}*, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}*, all of which were lower than those

observed in *chl1-5* and *nrt1.1* (Figure 2d). These observations suggest that NRT1.1 does

not downregulate *STOP1*/STOP1 activities and that STOP1 dominates expression of

direct target genes of STOP1 over NRT1.1.

3.3 Possible scheme for regulation of ammonium tolerance by NRT1.1 and STOP1

Ye et al. (2021) demonstrated that, under acidic conditions with nitrate as the primary N

source, STOP1 directly upregulates *NRT1.1* expression in the root, which facilitates

co-uptake of proton and nitrate, elevating rhizosphere pH to a favorable level for

Arabidopsis: therefore, NRT1.1 was epistatic to STOP1 on root nitrate uptake and root

elongation. Meanwhile, the present study showed that, under toxic levels of ammonium

without nitrate, ammonium-inducible STOP1 dominates ammonium tolerance and

target gene expression of STOP1 over NRT1.1 (Figure 1 and 2). Moreover, we found

that ammonium-dependent transcriptional induction of shoot *NRT1.1* was completely

suppressed by *STOP1* deficiency (Supplementary Figure 1), supporting that *NRT1.1* is

under the control of STOP1. To reconcile these observations, a possible scheme for

regulation of ammonium tolerance by NRT1.1 and STOP1 is presented (Figure 2e). The

STOP1 upregulated by ammonium induces the expression of *NRT1.1* and other target

gene(s). The function(s) of some target gene product(s) (i.e., Target X in Figure 2e)

could enhance ammonium tolerance but be suppressed by NRT1.1, possibly via

protein–protein interaction(s).

In *A. thaliana*, STOP1 directly induces *CIPK23* expression under toxic levels

of ammonium as the sole N source (Tian et al. 2021). CIPK23 inhibits ammonium

transport via phosphorylation of high-affinity AMMONIUM TRANSPORTER 1;1

(AMT1;1) and 1;2 (AMT1;2) (Straub et al. 2017). Since the loss-of-function of *CIPK23*

causes excessive ammonium uptake and thus ammonium hypersensitivity (Hachiya et al.

2024; Straub et al. 2017), STOP1 improves ammonium tolerance by upregulating

CIPK23 (Tian et al. 2021). Notably, under low nitrate conditions, CIPK23 physically

interacts with NRT1.1 and phosphorylates it, increasing the affinity of NRT1.1 for

nitrate (Ho et al. 2009). It remains unclear whether NRT1.1 reduces ammonium

tolerance through its interaction with CIPK23 as a candidate of Target X (Figure 2e).

Meanwhile, a recent study reported that STOP1 indirectly activates the ARABIDOPSIS

PLASMA-MEMBRANE H^+-ATPase 2 (AHA2) by upregulating *SMALL AUXIN-UP*

RNA 55 (*SAUR55*) whose translational product prevents PP2C.D2/5, negative regulators

of AHA2 (Agrahari et al. 2023). The plant Cistrome database implies that STOP1 could

directly bind to the promoter of *AHA2*. By contrast, NRT1.1 and receptor-kinase QIAN

SHOU KINASE 1 (QSK1) inactivate the AHA2 via phosphorylation on S899 under low

nitrate (Zhu et al. 2024). Hence, STOP1 and NRT1.1 may affect ammonium-derived

acidic stress via AHA2 in opposite directions and therefore, AHA2 could also be a

possible candidate for Target X in Figure 2e. It would be valuable to explore the

common target(s) of NRT1.1 and STOP1 for better understanding ammonium toxicity.

Acknowledgments

We would like to thank Dr. Thierry Desnos (Aix-Marseille University) for providing the

Arabidopsis seeds, Dr. Alain Gojon for valuable discussion, and Mr. Shun Ohno for

supporting experiments. The authors would like to thank Enago (www.enago.jp) for the

English language review.

Disclosure statement

The authors report there are no competing interests to declare.

Funding

This study was supported by the Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan under Grant [17K15237/20K05771/23K04978], the Inamori Foundation, the Agropolis Foundation Grant [1502-405], and the Nagase Science and Technology Foundation, and the Yanmar Joint Research to TH.

References

Agrahari, R. K., Y. Kobayashi, T. Enomoto, T. Miyachi, M. Sakuma, M. Fujita, T. Ogata, et al. 2023. "STOP1-regulated *SMALL AUXIN UP RNA55* (*SAUR55*) is involved in proton/malate co-secretion for Al tolerance in Arabidopsis." *Plant Direct* 8(1): e557. doi: 10.1002/pld3.557.

Balzergue, C., T. Dartevelle, C. Godon, E. Laugier, C. Meisrimler, J. M. Teulon, A. Creff, et al. 2017. "Low phosphate activates STOP1-ALMT1 to rapidly inhibit root cell elongation." *Nature Communications* 8 (1): 15300. doi: 10.1038/ncomms15300.

Bouguyon, E., F. Brun, D. Meynard, M. Kubeš, M. Pervent, S. Leran, B. Lacombe, et al. 2015. "Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor NRT1.1." *Nature Plants* 1 (3): 15015. doi:10.1038/nplants.2015.15.

Ding, Z. J., C. Xu, J. Y. Yan, Y. X. Wang, M. Q. Cui, J. J. Yuan, Y. N. Wang, et al.

(2024). "The LRR receptor-like kinase ALR1 is a plant aluminum ion sensor." *Cell*

Research, 34 (4): 281−294. doi: 10.1038/s41422-023-00915-y.

Enomoto, T., M. Tokizawa, H. Ito, S. Iuchi, M. Kobayashi, Y. Y. Yamamoto, Y.

Kobayashi, et al. 2019. "STOP1 regulates the expression of *HsfA2* and *GDHs* that are

critical for low-oxygen tolerance in Arabidopsis." *Journal of Experimental Botany* 70

(12): 3297−3311. doi:10.1093/jxb/erz124.

Hachiya, T., Y. Mizokami, K. Mizokami, D. Tholen, C. K. Watanabe, and K. Noguchi.

2011. "Evidence for a nitrate-independent function of the nitrate sensor NRT1.1 in

Arabidopsis thaliana." *Journal of Plant Research* 124 (3): 425−430.

doi:10.1007/s10265-010-0385-7.

Hachiya, T. and K. Noguchi. 2011. "Mutation of NRT1.1 enhances ammonium/low

pH-tolerance in *Arabidopsis thaliana*." *Plant Signaling & Behavior* 6 (5): 706−708.

doi:10.4161/psb.6.5.15068.

Hachiya, T., J. Inaba, M. Wakazaki, M. Sato, K. Toyooka, A. Miyagi, M.

Kawai-Yamada et al. 2021. "Excessive ammonium assimilation by plastidic glutamine

synthetase causes ammonium toxicity in *Arabidopsis thaliana*." *Nature*

Communications 12 (1): 4944. doi:10.1038/s41467-021-25238-7.

Hachiya, T., N. Makita, L. Bach, A. Gojon, T. Nakagawa, and H. Sakakibara. 2024. "

Genetic and transcriptomic dissection of nitrate- independent function of Arabidopsis

NRT1.1/ NPF6.3/CHL1 under high ammonium condition." *Soil Science and Plant*

Nutrition in press. doi: 10.1080/00380768.2024.2326086.

Ho, C. H., S. H. Lin, H. C. Hu, and Y. F. Tsay. 2009. "CHL1 functions as a nitrate

sensor in plants." *Cell* 138 (6): 1184−1194. doi:10.1016/j.cell.2009.07.004.

Jian, S., Q. Liao, H. Song, Q. Liu, J. E. Lepo, C. Guan, J. Zhang, et al. 2018.

"NRT1.1-related NH_4^+ toxicity is associated with a disturbed balance between NH_4^+

uptake and assimilation." *Plant Physiology* 178 (4): 1473−1488.

doi:10.1104/pp.18.00410.

Liu, X. X., Y. X. Zhu, X. Z. Fang, J. Y. Ye, W. X. Du, Q. Y. Zhu, X. Y. Lin, et al., 2020.

"Ammonium aggravates salt stress in plants by entrapping them in a chloride

over-accumulation state in an NRT1.1-dependent manner." *Science of The Total*

Environment 746: 141244. doi:10.1016/j.scitotenv.2020.141244.

Sawaki, Y., S. Iuchi, Y. Kobayashi, Y. Kobayashi, T. Ikka, N. Sakurai, M. Fujita, K., et

al. 2009. "STOP1 regulates multiple genes that protect Arabidopsis from proton and

aluminum toxicities." *Plant Physiology* 150 (1): 281−294. doi:10.1104/pp.108.134700.

Sadhukhan, A., T. Enomoto, Y. Kobayashi, T. Watanabe, S. Iuchi, M. Kobayashi, L.

Sahoo, et al. 2019. "Sensitive to proton rhizotoxicity1 regulates salt and drought

tolerance of *Arabidopsis thaliana* through transcriptional regulation of *CIPK23*." *Plant*

and Cell Physiology 60 (9): 2113−2126. doi:10.1093/pcp/pcz120.

Straub, T., U. Ludewig, and B. Neuhäuser. 2017. "The kinase CIPK23 inhibits

ammonium transport in *Arabidopsis thaliana*." *The Plant Cell* 29 (2): 409−422. doi:

10.1105/tpc.16.00806.

Tian, W. H., J. Y. Ye, M. Q. Cui, J. B. Chang, Y. Liu, G. X. Li, Y. R. Wu, et al. 2021. "A

transcription factor STOP1-centered pathway coordinates ammonium and phosphate

acquisition in *Arabidopsis*." *Molecular Plant* 14 (9): 1554−1568.

doi:10.1016/j.molp.2021.06.024.

Tokizawa, M., T. Enomoto, H. Ito, L. Wu, Y. Kobayashi, J. Mora-Macías, D.

Armenta-Medina, et al. 2021. "High affinity promoter binding of STOP1 is essential for

early expression of novel aluminum-induced resistance genes *GDH1* and *GDH2* in

Arabidopsis." *Journal of Experimental Botany* 72 (7): 2769−2789.

doi:10.1093/jxb/erab031.

Tsay, Y. F., J. I. Schroeder, K. A. Feldmann, and N. M. Crawford. 1993. "The herbicide

sensitivity gene *CHL1* of Arabidopsis encodes a nitrate-inducible nitrate transporter."

Cell 72 (5): 705−713. doi:10.1016/0092-8674(93)90399-B.

Ye, J. Y., W. H. Tian, M. Zhou, Q. Y. Zhu, W. X. Du, Y. X. Zhu, X. X. Liu, et al. 2021.

"STOP1 activates NRT1.1-mediated nitrate uptake to create a favorable rhizospheric pH

for plant adaptation to acidity." *The Plant Cell* 33 (12): 3658−3674.

doi:10.1093/plcell/koab226.

Zhang, Y., J. Zhang, J. Guo, F. Zhou, S. Singh, X. Xu, Q. Xie, et al. "F-box protein

RAE1 regulates the stability of the aluminum-resistance transcription factor STOP1 in

Arabidopsis." *Proceedings of the National Academy of Sciences USA* 116 (1): 319−327.

doi: 10.1073/pnas.181442611.

Zhou, F., S. Singh, J. Zhang, Q. Fang, C. Li, J. Wang, C. Zhao, et al. "The

MEKK1-MKK1/2-MPK4 cascade phosphorylates and stabilizes STOP1 to confer

aluminum resistance in Arabidopsis." *Molecular Plant* 16 (2): 337−353. doi:

10.1016/j.molp.2022.11.010.

Zhu, Z., L. Krall, Z. Li, L. Xi, H. Luo, S. Li, M. He, et al. "Transceptor NRT1.1 and

receptor-kinase OSK1 complex controls PM H^+-ATPase activity under low nitrate."

Current Biology 34(7): 1479−1491. doi: 10.1016/j.cub.2024.02.066.

Figure legends

Figure 1. *STOP1* deficiency suppresses enhanced ammonium tolerance in

NRT1.1-deficient mutants. (a) RT-PCR and agarose gel electrophoresis for *NRT1.1*,

STOP1, and *EF1α* using five-day-old seedlings of Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5*

stop1^{KO}, and *nrt1.1 stop1^{KO}* grown under 10 mM ammonium. (b,c) Representative

photographs (b) and fresh weights (c) of shoots from 11-day-old Col-0, *chl1-5*, *nrt1.1*,

stop1^{KO}, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}* seedlings grown under 10 mM ammonium

with 100 nM IAA (using 100 μM IAA (in dimethyl sulfoxide) stock solution). (d) Fresh

weights of shoots and roots from 11-day-old Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5*

stop1^{KO}, and *nrt1.1 stop1^{KO}* seedlings grown under 10 mM ammonium without IAA.

(b-d) In one dish, nine seedlings of each line were regarded as a single biological

replicate (Mean ± SD; n = 3–4). Different lowercase letters indicate significant

differences evaluated by the Tukey–Kramer multiple comparison test conducted at a

significance level of $P < 0.05$.

Figure 2. NRT1.1 does not downregulate *STOP1*/STOP1 and STOP1 dominates

expression of direct target genes of STOP1 over NRT1.1. (a) Relative transcript levels

of *ALMT1*, *CIPK23*, *GDH2*, *STOP1*, and *STOP2* in shoots and roots of five-day-old

seedlings from Col-0 and *chl1-5* grown under 10 mM ammonium or 10 mM nitrate

conditions without IAA. (b) Comparisons of the expression of STOP1-upregulated

genes under acidic pH (Sawaki et al. 2009) between Col-0 and *NRT1.1*-deficient

mutants (*chl1-5*, *nrt1.1*) grown under 10 mM ammonium without IAA based on the

microarray results (Hachiya et al. 2024). Changes in gene expression levels were shown

as logarithms to base 2 of fold change (Mean ± SD). (c) GFP-STOP1 signals from the

primary root tips of five-day-old seedlings grown under 10 mM ammonium or 10 mM

nitrate without IAA. The left four images and lower right image were generated by the

maximum intensity projection of multiple slices of the range where fluorescence is

clearly observed. The single cell image in the upper right is a composite of bright and

fluorescent images. Scale bars: 50 μm (left), 5 μm (upper right), and 20 μm (lower

right). (d) Relative transcript levels of *ALMT1*, *CIPK23*, and *GDH2* in five-day-old

seedlings from Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}* grown

under 10 mM ammonium without IAA. (a, d) Thirty-seven seedlings from one plate

were regarded as a single biological replicate (Mean ± SD; n = 3–5). (e) A possible

scheme for ammonium tolerance regulation by NRT1.1 and STOP1.

Figure 1

a

NRT1.1
STOP1
EF1α

Col
chl1-5
nrt1.1
stop1^{KO}
chl1-5 stop1^{KO}
nrt1.1 stop1^{KO}

NH_4^+

b

| Col | chl1-5 |
| stop1^{KO} | chl1-5 stop1^{KO} |

NH_4^+

| Col | nrt1.1 |
| stop1^{KO} | nrt1.1 stop1^{KO} |

NH_4^+

c

Shoot FW

% of Col

Col   chl1-5   stop1^{KO}   chl1.5 stop1^{KO}

NH_4^+

Shoot FW

% of Col

Col   nrt1.1   stop1^{KO}   nrt1.1 stop1^{KO}

NH_4^+

d

Shoot FW

% of Col

Col   chl1-5   stop1^{KO}   chl1.5 stop1^{KO}   Col   chl1-5   stop1^{KO}   chl1.5 stop1^{KO}

NH_4^+ NO_3^-

Shoot FW

% of Col

Col   nrt1.1   stop1^{KO}   nrt1.1 stop1^{KO}   Col   nrt1.1   stop1^{KO}   nrt1.1 stop1^{KO}

NH_4^+ NO_3^-

Root FW

% of Col

Col   chl1-5   stop1^{KO}   chl1.5 stop1^{KO}   Col   chl1-5   stop1^{KO}   chl1.5 stop1^{KO}

NH_4^+ NO_3^-

Root FW

% of Col

Col   nrt1.1   stop1^{KO}   nrt1.1 stop1^{KO}   Col   nrt1.1   stop1^{KO}   nrt1.1 stop1^{KO}

NH_4^+ NO_3^-

Figure 2

a

ALMT1
fold change (0–1.5)
a, b, c, c, d, de, ce, ce

CIPK23
fold change (0–3)
a, b, b, b, c, de, d, ce

GDH2
fold change (0–1.5)
a, b, c, c, d, e, f, f

STOP1
fold change (0–4)
a, ab, b, ab, c, cd, e, de

STOP2
fold change (0–15)
a, ab, b, b, c, d, e, f

Col, chl1-5, Col, chl1-5, Col, chl1-5, Col, chl1-5
NH₄⁺, NO₃⁻, NH₄⁺, NO₃⁻
Shoot, Root

b STOP1-up-genes

Log_2(*NRT1.1*-deficient/Col) (−4 to 2)

1st 2nd 1st 2nd 1st 2nd 1st 2nd
chl1-5 *nrt1.1* *chl1-5* *nrt1.1*
3-day, 5-day

d

ALMT1
fold change (0–1.5)
a, b, b, c, c, c

CIPK23
fold change (0–1.5)
a, b, b, c, c, c

GDH2
fold change (0–1.5)
a, b, b, c, c, c

Col, chl1-5, nrt1.1, stop1^KO, chl1.5 stop1^KO, nrt1.1 stop1^KO
NH₄⁺ · Seedling

c GFP-STOP1

stop1^KO, chl1-5 stop1^KO, stop1^KO, chl1-5 stop1^KO, stop1^KO
NH₄⁺, NO₃⁻, NH₄⁺

e

High NH₄⁺
↓
STOP1
↓ ↓ ↓
NRT1.1 ⊣ Target X CIPK23? AHA2? → Target Y
↓
NH₄⁺ tolerance