Title

STOP1 dominates Arabidopsis tolerance to ammonium over NRT1.1/NPF6.3/CHL1

Running title

STOP1 dominates NH_4^+ tolerance over NRT1.1

Authors and addresses

Takushi Hachiya[a,b,†], Mako Sakai[a], Tsuyoshi Nakagawa[a], and Hitoshi Sakakibara[b,c]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for

Science Research, Shimane University, Matsue 690-8504, Japan

[b]RIKEN Center for Sustainable Resource Science, Yokohama 230-0045, Japan

[c]Graduate School of Bioagricultural Sciences, Nagoya University, Nagoya 464-8601,

Japan

Corresponding Author

Takushi Hachiya

Department of Molecular and Functional Genomics, Interdisciplinary Center for

Science Research, Shimane University, Matsue 690-8504, Japan

Email: takushi.hachiya@life.shimane-u.ac.jp, Tel: +81-852-32-6288

Abstract

The Arabidopsis nitrate transporter/sensor NRT1.1/NPF6.3/CHL1 plays significant roles even in the absence of nitrate. The loss-of-function of NRT1.1 alleviates growth suppression and chlorosis under toxic levels of ammonium as the sole nitrogen source. Recently, we reported that acidic stress-inducible genes are downregulated by *NRT1.1* deficiency under ammonium toxicity conditions, implying that NRT1.1 may exacerbate ammonium-dependent acidic stress. The transcription factor SENSITIVE TO PROTON RHIZOTOXICITY (STOP1) enhances Arabidopsis tolerance to ammonium and acidic stresses. Furthermore, STOP1 directly activates *NRT1.1* transcription. These previous findings prompted us to analyze the association between NRT1.1 and STOP1 on ammonium tolerance. Here, we show that ammonium-inducible STOP1 plays a dominant role over NRT1.1 in ammonium tolerance and expression of direct target genes of STOP1. We present a novel scheme in which NRT1.1 and STOP1 jointly regulate common target(s) associated with ammonium tolerance.

Keywords: ammonium toxicity, CHL1, NPF6.3, NRT1.1, STOP1

1. Introduction

Nitrate serves not only as a substrate for nitrogen (N) assimilation but also as a signaling molecule. The Arabidopsis nitrate transporter NRT1.1/NPF6.3/CHL1 acts as a nitrate sensor and orchestrates various plant responses to the nitrate signal, including gene expression and root system development (Bouguyon et al. 2015; Ho et al. 2009). Apart from its nitrate-dependent functions, NRT1.1 exhibits significant roles in the absence of nitrate (Hachiya et al. 2024). Previous studies have shown that the loss-of-function of NRT1.1 alleviates growth suppression and chlorosis under toxic levels of ammonium as the sole N source, i.e., ammonium toxicity (Hachiya et al. 2011; Jian et al. 2018; Liu et al. 2020). Recently, we reported that in early seedlings grown under 10 mM ammonium, the genes induced by acidic stress are downregulated in *NRT1.1*-deficient mutants relative to the wild-type, whereas those repressed by acidic stress are upregulated in the mutants (Hachiya et al. 2024). Moreover, an increase in medium pH promotes shoot growth of the wild-type more intensely than that of the *NRT1.1*-deficient mutant under 10 mM ammonium. Given that acidic stress is one of the primary causes of ammonium toxicity (Hachiya et al. 2021), NRT1.1 may exacerbate the ammonium-dependent acidic stress.

In *Arabidopsis*, the zinc finger transcription factor SENSITIVE TO PROTON RHIZOTOXICITY (STOP1) positively regulates genes involved in the tolerance to ammonium and acidic stresses, as well as abiotic stresses, including aluminum, drought, salt, low phosphate, and low oxygen (Balzergue et al. 2017; Enomoto et al. 2019; Hachiya et al. 2021; Sadhukhan et al. 2019; Sawaki et al. 2009; Tian et al. 2021).

Furthermore, STOP1 directly activates *NRT1.1* expression under an acidic rhizosphere

containing nitrate (Ye et al. 2021). These findings prompted us to analyze the

association between NRT1.1 and STOP1 under toxic levels of ammonium as the sole N

source. Here, we show that STOP1 dominates ammonium tolerance and expression of

direct target genes of STOP1 over NRT1.1.

2. Materials and Methods

2.1. Plant materials and growth conditions

Arabidopsis thaliana accession Columbia (Col-0) was used as a control. T-DNA

insertion mutants *nrt1.1* (SALK_097431, Hachiya et al. 2011), *stop1^{KO}* (SALK_114108,

Sawaki et al. 2007), and gamma ray-mutagenized mutant c*hl1-5* (Tsay et al. 1993) were

obtained from the European Arabidopsis Stock Center. *stop1^{KO}* was backcrossed twice

with Col-0 before use. The line harboring *pSTOP1:GFP-STOP1* in the *stop1^{KO}*

background (Balzergue et al. 2017) was provided by Dr. Thierry Desnos (Aix-Marseille

University, France).

The seeds were surface sterilized, placed in plastic Petri dishes (diameter: 90 mm;

depth: 20 mm; Iwaki, Tokyo, Japan), and grown on a solid medium containing 30 mL

of Murashige and Skoog (MS) salts without N sources supplemented with 4.7 mM

MES-KOH (pH 5.7), 2% (w/v) sucrose, and 0.25% (w/v) gellan gum (Wako, Osaka,

Japan) with or without 100 nM indole-3-acetic acid (IAA). Two different N and K

sources were used: 10 mM KNO$_3$ (10 mM NO$_3^-$ condition) or 5 mM (NH$_4$)$_2$SO$_4$ with 10

mM KCl (10 mM NH$_4^+$ condition). The plants were kept in the dark at 4°C for 3 days,

and then grown horizontally under a photosynthetic photon flux density of 100–130

μmol m^{-2} s^{-1} (16-h light/8-h dark cycle) at 23°C.

2.2. Extraction of RNA

Frozen samples were ground using TissueLyser II (Qiagen, Tokyo, Japan) using

zirconia beads. Total RNA was extracted using an RNeasy Plant Mini Kit (Qiagen),

following the manufacturer's instructions.

2.3. Quantitative reverse transcription polymerase chain reaction

Reverse transcription (RT) was conducted using ReverTra Ace qPCR RT Master Mix

with gDNA Remover (Toyobo, Tokyo, Japan), following the manufacturer's

instructions. The obtained cDNA was diluted 10-fold with distilled water for

quantitative PCR (qPCR). Transcript levels were evaluated using StepOnePlus

Real-Time PCR System (Thermo Fisher Scientific, Waltham, MA, USA). The diluted

cDNA (2 μL) was amplified in the presence of 10 μL KAPA SYBR FAST qPCR Kit

(Nippon Genetics, Tokyo, Japan), 0.4 μL specific primers (0.2 μM final concentration),

and 7.2 μL of sterile water. Relative transcript levels were calculated using the

comparative cycle threshold method with *ACTIN3* as an internal standard. Primer

sequences are shown in Supplementary Table 1.

2.4. Observation of fluorescent signal

Confocal imaging was performed using a Leica SP5 Confocal Microscope (Leica

Microsystems, Wetzlar, Germany). GFP-STOP1 was excited with a 488-nm Ar laser,

and emission was detected from 500–530 nm using a Leica 25× objective (HCX IRAPO

L 25×/0.95 WATER).

3. Results and Discussion

3.1. STOP1 dominates ammonium tolerance over NRT1.1

To analyze the genetic relationship between *NRT1.1* and *STOP1*, *chl1-5* and *nrt1.1* were

crossed with *stop1^{KO}*, and their double homozygous mutants were obtained (Figure 1a).

The growth of Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}* under

10 mM ammonium supplemented with 100 nM IAA was compared. It should be noted

that appropriate IAA concentrations stabilize the ammonium tolerance of

NRT1.1-deficient mutants (Hachiya and Noguchi 2011). The shoots of *chl1-5* and *nrt1.1*

were larger and greener than those of Col-0 (Figure 1b). The growth enhancement in

NRT1.1-deficient mutants was almost completely suppressed by *STOP1* deficiency.

Shoot fresh weights of *chl1-5* and *nrt1.1* were much larger than those of Col-0, whereas

those of *stop1^{KO}* were slightly smaller (Figure 1c). Fresh weights of the shoots of *chl1-5*

stop1^{KO} and *nrt1.1 stop1^{KO}* were comparable to those of *stop1^{KO}*. Also, under 10 mM

ammonium without IAA, the shoot fresh weights were increased by *NRT1.1* deficiency,

but they were reduced by *STOP1* deficiency to a level comparable to those of *stop1^{KO}*

(Figure 1d). Under 10 mM nitrate without IAA, the shoot fresh weights of

NRT1.1-deficient mutants were significantly smaller than those of other lines with or

without *STOP1*. Under both N conditions, the fresh weights of the root showed a similar

tendency to those of the shoot. Thus, our genetic analysis indicates that STOP1

dominates ammonium tolerance over NRT1.1.

3.2 STOP1 dominates expression of direct target genes of STOP1 over NRT1.1

The above growth analysis suggests that NRT1.1 could act upstream of STOP1 and

downregulate *STOP1*/STOP1 activities under ammonium. To verify this, we checked

the transcript levels of known direct target genes of STOP1, i.e., *ALMT1*, *CIPK23*,

GDH2, and *STOP2* (Balzergue et al. 2017; Enomoto et al. 2019; Sadhukhan et al. 2019,

Tokizawa et al. 2021) in the absence of IAA. In five-day-old Col-0 shoots and roots, the

expression of *ALMT1*, *CIPK23*, *GDH2*, and *STOP2* was significantly higher under 10

mM ammonium than under 10 mM nitrate (Figure 2a). This ammonium-dependent

transcriptional induction was consistently decreased by *NRT1.1* deficiency (Figure 2a).

Also, *STOP1* expression in shoots and roots was not significantly increased by *NRT1.1*

deficiency under ammonium (Figure 2a). Moreover, in five-day-old seedlings grown

under 10 mM ammonium, the transcript levels of *ALMT1*, *CIPK23*, and *GDH2* were

lowest in *STOP1*-deficient mutants with or without *NRT1.1* (Figure 2b). Microarray

analysis using three- and five-day-old seedlings under 10 mM ammonium without IAA

revealed that *NRT1.1* deficiency did not generally induce genes whose expression is

upregulated via STOP1 under acidic pH (Sawaki et al. 2009) (Figure 2c and

Supplementary Table S2). The F-box protein RAE1 interacts with STOP1 and promotes

its degradation via the ubiquitin-26S proteasome pathway (Zhang et al. 2019),

suggesting the post-translational regulation of STOP1 activity. To analyze the effect of

NRT1.1 on STOP1 expression, *chl1-5* was crossed with *pSTOP1:GFP-STOP1* line in

the *stop1^{KO}* background (Balzergue et al. 2017) and produced the *pSTOP1:GFP-STOP1*

line in the *chl1-5 stop1^{KO}* background. The GFP-STOP1-derived fluorescence signal

was promoted in the roots under 10 mM ammonium, similar to a previous result (Tian

et al. 2021). Meanwhile, the GFP signal was little changed by *NRT1.1* deficiency under

ammonium (Figure 2d). These observations suggest that NRT1.1 does not downregulate

STOP1/STOP1 activities and that STOP1 dominates expression of direct target genes of

STOP1 over NRT1.1.

3.3 Possible scheme for regulation of ammonium tolerance by NRT1.1 and STOP1

The present study shows that ammonium-inducible STOP1 dominates ammonium

tolerance and target gene expression of STOP1 over NRT1.1. A possible scheme for

regulation of ammonium tolerance by NRT1.1 and STOP1 is presented (Figure 2e),

noting that STOP1 directly upregulates *NRT1.1* expression (Ye et al. 2021). The STOP1

upregulated by ammonium induces the expression of *NRT1.1* and other target gene(s).

The function(s) of some target gene product(s) (i.e., Target X in Figure 2e) could

enhance ammonium tolerance but be suppressed by NRT1.1, possibly via protein–

protein interaction(s).

In *A. thaliana*, STOP1 directly induces *CIPK23* expression (Sadhukhan et al.

2019), and CIPK23 phosphorylates NRT1.1 to alter its affinity with nitrate (Ho et al.

2009). Since CIPK23 also inhibits ammonium transport via phosphorylation of

high-affinity ammonium transporter AMT1, the loss-of-function of *CIPK23* confers ammonium-hypersensitivity (Hachiya et al. 2024; Straub et al. 2017). Meanwhile, a recent study reported that STOP1 indirectly activates the plasma membrane H^+-ATPase AHA2 by upregulating *SAUR55* whose translational product prevents PP2C.D2/5, negative regulators of AHA2 (Agrahari et al. 2023). The plant Cistrome database implies that STOP1 could directly bind to the promoter of *AHA2*. Moreover, NRT1.1 and receptor-kinase QSK1 inactivates the AHA2 via phosphorylation on S899 under low nitrate (Zhu et al. 2024). Hence, CIPK23 and AHA2 could be possible candidates for Target X in Figure 2e, although inhibition of CIPK23 function by NRT1.1 remains unknown. It would be valuable to explore the common target(s) of NRT1.1 and STOP1 for better understanding ammonium toxicity.

Acknowledgments

We would like to thank Dr. Thierry Desnos (Aix-Marseille University) for providing the *Arabidopsis* seeds, Dr. Alain Gojon for valuable discussion, and Mr. Shun Ohno for supporting experiments. The authors would like to thank Enago (www.enago.jp) for the English language review.

Disclosure statement

The authors report there are no competing interests to declare.

Funding

198   This study was supported by the Grants-in-Aid from the Ministry of Education, Culture,

199   Sports, Science and Technology, Japan under Grant

200   [number17K15237/20K05771/23K04978, the Inamori Foundation, the Agropolis

201   Foundation Grant [number1502-405], the Nagase Science and Technology Foundation,

202   and the Yanmar Joint Research to TH.

204   References

Agrahari, R. K., Y. Kobayashi, T. Enomoto, T. Miyachi, M. Sakuma, M. Fujita, T.

Ogata, et al. 2023. "STOP1-regulated *SMALL AUXIN UP RNA55* (*SAUR55*) is involved

in proton/malate co-secretion for Al tolerance in Arabidopsis." *Plant Direct* 8(1): e557.

doi: 10.1002/pld3.557.

Balzergue, C., T. Dartevelle, C. Godon, E. Laugier, C. Meisrimler, J. M. Teulon, A.

Creff, et al. 2017. "Low phosphate activates STOP1-ALMT1 to rapidly inhibit root cell

elongation." *Nature Communications* 8 (1): 15300. doi: 10.1038/ncomms15300.

Bouguyon, E., F. Brun, D. Meynard, M. Kubeš, M. Pervent, S. Leran, B. Lacombe, et al.

2015. "Multiple mechanisms of nitrate sensing by *Arabidopsis* nitrate transceptor

NRT1.1." *Nature Plants* 1 (3): 15015. doi:10.1038/nplants.2015.15.

Enomoto, T., M. Tokizawa, H. Ito, S. Iuchi, M. Kobayashi, Y. Y. Yamamoto, Y.

Kobayashi, et al. 2019. "STOP1 regulates the expression of *HsfA2* and *GDHs* that are

critical for low-oxygen tolerance in Arabidopsis." *Journal of Experimental Botany* 70
(12): 3297−3311. doi:10.1093/jxb/erz124.

Hachiya, T., Y. Mizokami, K. Mizokami, D. Tholen, C. K. Watanabe, and K. Noguchi.
2011. "Evidence for a nitrate-independent function of the nitrate sensor NRT1.1 in
Arabidopsis thaliana." *Journal of Plant Research* 124 (3): 425−430.
doi:10.1007/s10265-010-0385-7.

Hachiya, T. and K. Noguchi. 2011. "Mutation of NRT1.1 enhances ammonium/low
pH-tolerance in *Arabidopsis thaliana*." *Plant Signaling & Behavior* 6 (5): 706−708.
doi:10.4161/psb.6.5.15068.

Hachiya, T., J. Inaba, M. Wakazaki, M. Sato, K. Toyooka, A. Miyagi, M.
Kawai-Yamada et al. 2021. "Excessive ammonium assimilation by plastidic glutamine
synthetase causes ammonium toxicity in *Arabidopsis thaliana*." *Nature
Communications* 12 (1): 4944. doi:10.1038/s41467-021-25238-7.

Hachiya, T., N. Makita, L. Bach, A. Gojon, T. Nakagawa, and H. Sakakibara. 2024. "
Genetic and transcriptomic dissection of nitrate- independent function of Arabidopsis
NRT1.1/ NPF6.3/CHL1 under high ammonium condition." *Soil Science and Plant
Nutrition* in press. doi: 10.1080/00380768.2024.2326086.

Ho, C. H., S. H. Lin, H. C. Hu, and Y. F. Tsay. 2009. "CHL1 functions as a nitrate

sensor in plants." *Cell* 138 (6): 1184−1194. doi:10.1016/j.cell.2009.07.004.

Jian, S., Q. Liao, H. Song, Q. Liu, J. E. Lepo, C. Guan, J. Zhang, et al. 2018.

"NRT1.1-related NH_4^+ toxicity is associated with a disturbed balance between NH_4^+

uptake and assimilation." *Plant Physiology* 178 (4): 1473−1488.

doi:10.1104/pp.18.00410.

Liu, X. X., Y. X. Zhu, X. Z. Fang, J. Y. Ye, W. X. Du, Q. Y. Zhu, X. Y. Lin, et al., 2020.

"Ammonium aggravates salt stress in plants by entrapping them in a chloride

over-accumulation state in an NRT1.1-dependent manner." *Science of The Total*

Environment 746: 141244. doi:10.1016/j.scitotenv.2020.141244.

Sawaki, Y., S. Iuchi, Y. Kobayashi, Y. Kobayashi, T. Ikka, N. Sakurai, M. Fujita, K., et

al. 2009. "STOP1 regulates multiple genes that protect Arabidopsis from proton and

aluminum toxicities." *Plant Physiology* 150 (1): 281−294. doi:10.1104/pp.108.134700.

Sadhukhan, A., T. Enomoto, Y. Kobayashi, T. Watanabe, S. Iuchi, M. Kobayashi, L.

Sahoo, et al. 2019. "Sensitive to proton rhizotoxicity1 regulates salt and drought

tolerance of *Arabidopsis thaliana* through transcriptional regulation of *CIPK23*." *Plant*

and Cell Physiology 60 (9): 2113−2126. doi:10.1093/pcp/pcz120.

Straub, T., U. Ludewig, and B. Neuhäuser. 2017. "The kinase CIPK23 inhibits

ammonium transport in *Arabidopsis thaliana*." *The Plant Cell* 29 (2): 409−422. doi:

10.1105/tpc.16.00806.

Tian, W. H., J. Y. Ye, M. Q. Cui, J. B. Chang, Y. Liu, G. X. Li, Y. R. Wu, et al. 2021. "A

transcription factor STOP1-centered pathway coordinates ammonium and phosphate

acquisition in *Arabidopsis*." *Molecular Plant* 14 (9): 1554−1568.

doi:10.1016/j.molp.2021.06.024.

Tokizawa, M., T. Enomoto, H. Ito, L. Wu, Y. Kobayashi, J. Mora-Macías, D.

Armenta-Medina, et al. 2021. "High affinity promoter binding of STOP1 is essential for

early expression of novel aluminum-induced resistance genes *GDH1* and *GDH2* in

Arabidopsis." *Journal of Experimental Botany* 72 (7): 2769−2789.

doi:10.1093/jxb/erab031.

Tsay, Y. F., J. I. Schroeder, K. A. Feldmann, and N. M. Crawford. 1993. "The herbicide

sensitivity gene *CHL1* of Arabidopsis encodes a nitrate-inducible nitrate transporter."

Cell 72 (5): 705−713. doi:10.1016/0092-8674(93)90399-B.

Ye, J. Y., W. H. Tian, M. Zhou, Q. Y. Zhu, W. X. Du, Y. X. Zhu, X. X. Liu, et al. 2021.

"STOP1 activates NRT1.1-mediated nitrate uptake to create a favorable rhizospheric pH

for plant adaptation to acidity." *The Plant Cell* 33 (12): 3658−3674.

doi:10.1093/plcell/koab226.

Zhang, Y., J. Zhang, J. Guo, F. Zhou, S. Singh, X. Xu, Q. Xie, et al. "F-box protein

RAE1 regulates the stability of the aluminum-resistance transcription factor STOP1 in

Arabidopsis." *Proceedings of the National Academy of Sciences USA* 116 (1): 319−327.

doi: 10.1073/pnas.181442611.

Zhu, Z., L. Krall, Z. Li, L. Xi, H. Luo, S. Li, M. He, et al. "Transceptor NRT1.1 and

receptor-kinase OSK1 complex controls PM H^+-ATPase activity under low nitrate."

Current Biology 34(7): 1479−1491. doi: 10.1016/j.cub.2024.02.066.

Figure legends

Figure 1. *STOP1* deficiency suppresses enhanced ammonium tolerance in

NRT1.1-deficient mutants. (a) RT-PCR and agarose gel electrophoresis for *NRT1.1*,

STOP1, and *EF1α* using five-day-old seedlings of Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5*

stop1^{KO}, and *nrt1.1 stop1^{KO}* grown under 10 mM ammonium. (b,c) Representative

photographs (b) and fresh weights (c) of shoots from 11-day-old Col-0, *chl1-5*, *nrt1.1*,

stop1^{KO}, *chl1-5 stop1^{KO}*, and *nrt1.1 stop1^{KO}* seedlings grown under 10 mM ammonium

with 100 nM IAA (using 100 μM IAA (in dimethyl sulfoxide) stock solution). (d) Fresh

weights of shoots and roots from 11-day-old Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5*

stop1^{KO}, and *nrt1.1 stop1^{KO}* seedlings grown under 10 mM ammonium without IAA.

(b-d) In one dish, nine seedlings of each line were regarded as a single biological

replicate (Mean ± SD; n = 3–4). Different lowercase letters indicate significant

differences evaluated by the Tukey–Kramer multiple comparison test conducted at a

significance level of P <0.05.

Figure 2. NRT1.1 does not downregulate STOP1. (a) Relative transcript levels of

ALMT1, *CIPK23*, *GDH2*, *STOP1*, and *STOP2* in shoots and roots of five-day-old

seedlings from Col-0 and *chl1-5* grown under 10 mM ammonium or 10 mM nitrate

conditions without IAA. (b) Relative transcript levels of *ALMT1*, *CIPK23*, and *GDH2*

in five-day-old seedlings from Col-0, *chl1-5*, *nrt1.1*, *stop1^{KO}*, *chl1-5 stop1^{KO}*, and *nrt1.1*

stop1^{KO} grown under 10 mM ammonium without IAA. (a, b) Thirty-seven seedlings

from one plate were regarded as a single biological replicate (Mean ± SD; n = 3–5). (c)

Comparisons of the expression of STOP1-upregulated genes under acidic pH (Sawaki et

al. 2009) between Col-0 and *NRT1.1*-deficient mutants (*chl1-5*, *nrt1.1*) grown under 10

mM ammonium without IAA based on the microarray results (Hachiya et al. 2024).

Changes in gene expression levels were shown as logarithms to base 2 of fold change

(Mean ± SD). (d) GFP-STOP1 signals from the primary root tips of five-day-old

seedlings grown under 10 mM ammonium or 10 mM nitrate without IAA. The left four

images and lower right image were generated by the maximum intensity projection of

multiple slices of the range where fluorescence is clearly observed. The single cell

image in the upper right is a composite of bright and fluorescent images. Scale bars: 50

µm (left), 5 µm (upper right), and 20 µm (lower right). (e) A possible scheme for

ammonium tolerance regulation by NRT1.1 and STOP1.

Figure 1

Figure 2

a

ALMT1
fold change
a b c c d de ce ce

CIPK23
fold change
a b b b c de d ce

GDH2
fold change
a b c c d e f f

STOP1
fold change
a ab b ab c cd e de

STOP2
fold change
a ab b b c d e f

Col	chl1-5	Col	chl1-5	Col	chl1-5	Col	chl1-5
NH₄⁺		NO₃⁻		NH₄⁺		NO₃⁻	
Shoot				Root			

b

ALMT1
fold change
a b b c c c

CIPK23
fold change
a b b c c c

GDH2
fold change
a b b c c c

Col
chl1-5
nrt1.1
stop1^KO
chl1.5 stop1^KO
nrt1.1 stop1^KO

NH₄⁺ · Seedling

c STOP1-up-genes

Log₂(*NRT1.1*-deficient/Col)

1st 2nd 1st 2nd 1st 2nd 1st 2nd
chl1-5 *nrt1.1* *chl1-5* *nrt1.1*
3-day   5-day

d GFP-STOP1

stop1^KO	chl1-5 stop1^KO	stop1^KO	chl1-5 stop1^KO	stop1^KO
NH₄⁺		NO₃⁻		NH₄⁺

e

High NH₄⁺
↓
STOP1
↓
NRT1.1 —⊣ Target X Target Y
↓
NH₄⁺ tolerance