Induction of the aggresome and insoluble tau aggregation using an optogenetic tool

Shigeo Sakuragi[1], Tomoya Uchida[1#], Naoki Kato[1#], Boxiao Zhao[1], Akito Hattori[1$], Yoshihiro Sakata[1], Yoshiyuki Soeda[2], Akihiko Takashima[2], Hideaki Yoshimura[3], Gen Matsumoto[4], Hiroko Bannai[1*]

[1] *Department of Electrical Engineering and Biosciences, School of Advanced Science and Engineering, Waseda University, Shinjuku-Ku, Tokyo 162-0056, Japan.*

[2] *Laboratory for Alzheimer's Disease, Department of Life Science, Faculty of Science, Gakushuin University, Toshima-ku, Tokyo 171-8588, Japan*

[3] *Department of Chemistry, School of Science, The University of Tokyo, Tokyo 113-0033, Japan*

[4] *Department of Neurological Disease Control, Graduate School of Medicine, Osaka Metropolitan University, Osaka, 545-8585, Japan*

 # Equally contributed

$ Current affiliation：Department of Medical Laboratory Science, Kitasato University School of Health Sciences, Minami-Uonuma, 949-7241, Japan

Corresponding author: Hiroko Bannai, School of Advanced Science and Engineering, Department of Electrical Engineering and Biosciences, Waseda University, 2-2 Wakamatsu-cho, Shinjuku-Ku, Tokyo 162-0056, Japan. ORCID iD: https://orcid.org/0000-0002-0951-488X e-mail: hiroko.bannai@waseda.jp

ABSTRACT

Tauopathy is a group of diseases where fibrillary tau aggregates are formed in neurons and glial cells in

the brain. In Alzheimer's disease (AD), the most common form of tauopathy, tau aggregation begins in the

brainstem and entorhinal cortex and then spreads throughout the brain. Understanding the mechanism by

which locally formed tau pathology propagates throughout the brain is crucial for comprehending the

pathogenesis of AD. Therefore, a novel model of tau pathology that artificially induces tau aggregation in

targeted cells at specific times is essential. In this study, we report a novel optogenetic module, OptoTau, human

tau with the P301L mutation fused with a photosensitive protein CRY2Olig, which could induce various forms of

tau depending on the pattern of blue light illumination. Continuous blue light illumination for 12 h to Neuro2a

cells stably expressing OptoTau (OptoTauKI cells) resulted in cluster formation along microtubules, many of

which eventually accumulated in aggresomes.

On the other hand, when alternating light exposure and darkness in 30-minute cycles for 8 sets per day were

repeated over 8 days, methanol-insoluble tau aggregation was formed. Methanol-insoluble tau was induced more

rapidly by repeating cycles of 5-minute illumination followed by 25 minutes of darkness over 24 hours. These

findings suggest that OptoTau can induce various stages of tau aggregation depending on the pattern of blue

light exposure. Thus, this technique holds promise as a novel approach to creating specific tau aggregation in

targeted cells at desired time points.

Keywords: Tauopathy, Alzheimer's disease, CRY2Olig

◄ *Significance* ►

In this study, we established a technique to manipulate tau aggregation in a blue light-dependent manner using
21 cells stably expressing OptoTau, an optogenetic tool based on the CRY2Olig module. Tau accumulation in
aggresomes or insoluble tau aggregation were specifically induced in response to blue light illumination
22 conditions. These results are significant in that they provide a new technological basis for creating a singular
point of tau aggregation in specific targeted cells at a targeted time.

Introduction

Tauopathy is a group of diseases in which neurofibrillary tangles (NFTs), filamentous tau protein aggregates,

are formed in neurons and glial cells in the brain[1]. Despite being triggered by abnormalities in the same molecule tau,

tauopathy manifests as various disorders such as Alzheimer's disease (AD), Pick's disease, progressive supranuclear

palsy (PSP), corticobasal degeneration (CBD), chronic traumatic encephalopathy (CTE). In healthy neurons,

microtubule-binding protein tau plays physiological roles, such as neurite outgrowth and regulation of axonal transport

by motor proteins. However, tau detaches from microtubules and forms intracellular tau aggregates when tau is subject

to post-transcriptional modifications such as excessive phosphorylation [2]. In AD, the most common form of

tauopathy, the hyperphosphorylation of tau as a cause of NFT is initially formed in the neurons in locus coeruleus (LC)

and entorhinal cortex (EC) [3] [4]. This tau aggregation expands with age into the transentorhinal region, hippocampus,

and cerebral cortex (temporal lobe, prefrontal cortex, and higher sensory association neocortex) [4]. The progression of

tau aggregation is closely associated with the development of clinical pathology of AD [5] [6].

Several model experimental systems have contributed to understanding the relationship between the

accumulation of degenerative tau and higher brain functions in AD and the mechanisms underlying the expansion of tau

pathology. In mice overexpressing neurons with human tau with the mutation in 301st proline to serine or leucine

(P301S or P301L) found in familial frontotemporal dementia, tau aggregate formation, synapse loss, neurodegeneration,

and brain atrophy have been observed in a few months [7] [8] [9] [10]. Because these phenotypes have much in

common with the pathology in human patients, these mice have served as a model for AD, contributing to the

elucidation of the mechanism of toxicity from degenerated tau. Tau filaments polymerized in vitro and insoluble tau

fractions from AD patients injected into human tau-overexpressing mice form hyperphosphorylated tau and tau

aggregates in distant brain regions with neuronal connections [11] [12] [13]. Tau extracted from patient brains also

induces pathology, even in non-transgenic mice [14] [15]. The spread of degenerated tau to healthy neurons has been

shown to occur through synaptic contact or by uptake by other cells of degenerated tau released extracellularly in a

synaptic activity-dependent manner [16] [17]. These experiments have contributed to elucidating some aspects of the

pathogenesis of tauopathy as model systems that rationally reproduce some of the phenotypes seen in Alzheimer's

disease and other tauopathies in humans.

Recently, Nagai et al. defined a "singularity phenomenon" as a biological phenomenon in which a small

number of rare cells cause significant changes throughout a complex system, and these small number of rare cells are

named "singularity cells" [18] [19]. From this perspective, neurons that first accumulate hyperphosphorylated tau and NFTs in LCs and ECs can be regarded as "singularity cells" in AD [20]. How do these "singularity cells" that initially accumulate tau arise? How is the degeneration of tau transmitted from the singularity cells to the entire cortex? Addressing these questions is considered crucial for understanding the onset mechanisms of AD. However, to address this issue, the current experimental system of tauopathy models has several challenges to overcome. Although regions in the brain tend to aggregate tau, identified in tauopathy model mice with human tau overexpression, it remains unclear when the first "singularity cells" that accumulate tau are born and why they can produce tau aggregation. More seriously, the loss of endogenous genes and uncertain protein expression levels of transduced tau in human Tau overexpression model animals have been found, and whether the phenotypes obtained so far are due to abnormalities in tau requires reconsideration [10]. Another AD model, obtained by injecting exogenous tau seeds, bypasses the initial process of AD pathogenesis, tau aggregation. Tau seed uptake is entirely random, and it is impossible to identify the cell from which the pathology originates. A new model system is needed to create a singular point of tau aggregation at a targeted time in specifically targeted cells to elucidate the process of tau pathology from the singularity cells, neurons that first develop tau pathology.

In this study, we applied optogenetics to induce tau aggregation in targeted cells. CRY2, a photoreceptor in *Arabidopsis*, and its derivative CRY2Olig are optogenetic modules that form oligomers reversibly upon exposure to blue light [21]. Recently, researchers have introduced CRY2 to induce aggregation of proteins found in neurodegenerative diseases, including tau [22] [23] [24] [25]. CRY2 and tau fusion proteins accumulated on microtubules by liquid-liquid phase separation (LLPS) induced by blue light illumination [23]. Blue light illumination forms tau droplets and oligomers in primary cortical culture neurons of mice overexpressing tau fused with CRY2Olig, a higher capacity for homo-oligomer formation [24]. We have recently shown that 'OptoTau,' a fusion of human tau with the P301L mutation and CRY2Olig, transiently overexpressed in Neuo2a cells, could induce LLPS droplets and oligomers of tau by illuminating blue light for 24 hours [25]. In the present study, we manipulated intracellular tau by blue light illumination in Neuro2a cells stably expressing OptoTau. OptoTau, inserted into the safe-harbor region by CRISR/Cas9 technique, was subjected to various temporal patterns of blue light illumination, including eight-day illumination. Here, we report that different patterns of blue light illumination allow the same optogenetic tool, OptoTau, to induce opposite responses: induction of tau degradation by aggresome and formation of insoluble tau aggregates.

Materials and Methods

OptoTau knock-in into Neuro2a cells

OptoTau consists of the optogenetic module CRY2olig [21], a SNAP-tag, and a full-length human tau (2N4R) with a

P301L (Figure 1A). OptoTau-donor vector, OptoTau gene under the CMV promoter, and GFP and Puromycin resistant

genes under the EF1α promoter were constructed using ROSA26 MCS. The OptoTau-donor vector and

mROSA26sgRNA/Cas9 vector (GeneCopoeia), including the guide RNA and Cas9 sequences, were transfected into

mouse neuroblastoma-derived Neuro2a cells (JCRB Cell Bank) using Lipofectamine 3000 (Life Technologies). Drug

selection was performed by adding 400 µg/ml G418 (Nacalai tesque) on day one post-transfection and 1 µg/ml

puromycin (Nacalai tesque) 4 days later. GFP was used as a marker for gene insertion, and GFP-positive cells were

isolated into a 96-well plate. Single colonies of GFP-positive cells were replated into a 24-well plate and further

expanded on a 10 cm culture dish. After confirming the expression of OptoTau, one clone was used as OptoTau Knock-

in (OptoTauKI) cells in this study.

Cell Maintenance

OptoTauKI and Neuro2a wild type (WT) were maintained in Dulbecco's Modified Eagle's medium (4.5 g/l glucose;

Nacalai tesque) supplemented with 10% FBS, penicillin (100 units/ml) streptomycin (100 µg/ml) mixture, in 5% CO_2,

37°C environment. OptoTauKI cells were kept under light-shielded conditions to prevent irregular CRY2olig

conformational changes caused by room light. For the live imaging experiment, cells were plated on a 35 mm glass

bottom dish (MATSUNAMI) coated with 0.04% Polyethyleneimine (PEI, Sigma-Aldrich). Cells were plated on an 18

mm cover glass precoated with 0.04% PEI for fixation. The density of light-exposed cells was always maintained at 60-

80% confluence, except in the experiment where the cells were incubated for 48-72 hours after light illumination.

Blue light illumination

A blue light illumination device (TH2-211X200BL, CCS Inc.) was set in a 5% CO_2 incubator at 37°C. Plates of cells

were placed on a transparent plastic box with a height of 30 mm on the light-emitting device. The light intensity was

0.24 ± 0.04 mW/cm^2 throughout all experiments. The illumination protocol, one 12-hour illumination, 8 cycles of 30-

minute (min) illumination - 30-min light off per day for up to 8 days, and 48 cycles of 5-min illumination - 25-min light

off per day, was performed in this study. An automatic timer controlled the illumination timing.

Labeling of OptoTau using SNAP-tag

Fluorescent labeling of OptoTau molecules was performed by covalently binding a fluorescent-labeled SNAP ligand to a SNAP-tag inserted into the OptoTau. 0.1 µM SNAP-Cell 647-SiR (New England BioLabs) was administered into the culture medium, and OptoTau KI or WT was applied for 12 to 18 hours. Cells were washed with medium and then used for live cell imaging or immunocytochemistry.

Immunocytochemistry

Phosphate buffered saline (PBS) was used for dilution and washing throughout the immunocytochemistry. Cells were generally fixed with 4% fixation in 4% paraformaldehyde (PFA) for 15 min at room temperature (RT) and permeabilized with 0.1% Triton X-100 for 3 min at RT. For immunostaining of microtubule or aggresomal markers, cells were fixed with ice-cold Acetone/Methanol (Acetone: Methanol =1:1) for 3 minutes at -20°C, followed by post-fixation with 4% PFA (15 min, RT). To visualize insoluble tau, cells were fixed with ice-cold methanol for 20 min at -20°C. This treatment removes soluble tau in the cytosol [26]. After fixation and permeabilization, cells were incubated with 5% Bovine Serum Albumin (BSA) for 30 min RT for blocking, then with primary antibody treatment in 2.5% BSA for 1 hour at RT. Primary antibodies used were: anti-total tau antibody (RTM38, 1:1000, Fujifilm), antibody recognizing the initial structural change of pathological tau (MC1, 1:1000; gifted from Prof. Peter Davies [Albert Einstein College of Medicine Medicine], anti-αTubulin antibody (1:2000; Sigma-Aldrich), anti-γTubulin antibody (1:100; Santa Cruz), anti-HDAC6 antibody (1:100; Santa Cruz), anti-Dynein antibody (1: 100; Santa Cruz). After washing with PBS, cells were incubated with secondary antibodies (5 µg/ml in 2.5% BSA) for 40 min, RT. Secondary antibodies used were: Anti-rat IgG (H+L) Alexa Fluor® 555 conjugate, Anti-rat IgG (H+L) Alexa Fluor® 647 conjugate (Cell signaling) or Goat anti-Rat IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 594, Goat anti-Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 647, Goat anti- Mouse IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor™ 555 (Thermo Fisher Scientific).

Observation by epifluorescence microscopy and confocal laser scanning microscopy

Immunocytochemically or SNAP-tagged cells were observed using an inverted microscope (IX73; Evident) equipped with a high-intensity LED driver (DC4104, Thorlabs) and a digital CMOS camera (ORCA-Fish 4.0; HAMAMATSU). 60x oil immersion objective (NA 1.42; Evident), and the optical zoom was set to 5x. 5-10 fields of view were captured

from a single cover glass. Confocal laser scanning microscope (FV3000; Evident) images were taken with an oil

immersion objective (×60, NA1.42; Evident) at 5x digital zoom.

Image Analysis

Epifluorescence or confocal microscopy images were contrast-adjusted using MetaMorph (Molecular Devices) or

ImageJ (NIH) image analysis software. The percentage of cells with insoluble tau aggregates in the field of view was

calculated per images of PFA-fixed samples. The presence of the aggregates was defined when the maximum RTM38

intensity value in the cell was larger than "the mean of the RTM38 intensity + 3 standard deviation" of the dark

condition. The presence of aggregates around the nucleus was measured at the focal plane, where the nucleus was

maximally visible and calculated as a percentage per coverslip.

Western blotting

Immediately after completion of blue light illumination, 2.0×10^6 cells/mL of WT Neuro2a or OptoTau KI cells were

incubated in 10 mM HEPES buffer supplemented with 1% protease inhibitor (cOmplete, Roche), 1 µM okadaic acid

(Cayman Chemical), 1 mM sodium orthovanadate (MP Biomedicals), 1 mM sodium fluoride (Nacalai tesque), 1 mM β-

glycerophosphate disodium salt hydrate (Sigma-Aldrich), 250 mM Sucrose, 1 mM EDTA. Samples were subjected to

15 s- sonication followed by 15 s- ice-cooling 4 times and centrifuged at 3000 x g for 10 min at 4°C. The supernatant

was mixed with non-reducing Tris-Glycine SDS sample buffer (06-363No, TEFCO) and was loaded onto a 5-20%

gradient gel. Samples were electrophoresed at 40 mA and transferred to a PVDF membrane (170-4156, Bio-Rad). After

transfer, the membrane was blocked with 1% BSA with 0.05% Tween-20 and reacted with an anti-tau antibody

(RTM38, 1:2500) and anti-β-actin antibody (1:10,000; A5441, Sigma-Aldrich) at 1h RT (or 4°C O/N). After washing

three times with PBST (0.05% Tween20 in PBS), the membrane was incubated with the secondary antibodies

conjugated with peroxidase (1:10,000, Jackson Immuno Research) at RT for 1h. Chemiluminescence signals were

detected using a luminol-based chemiluminescence assay kit (Chemi-Lumi One L, Nacalai tesque) and

chemiluminescence detection system (ChemiDoc Touch MP imaging system, Bio-Rad).

Statistics

Numbers of data are listed in Figure legends. The Mann-Whitney U test was used to compare Western blots with and

without blue light illumination and the percentage of cells with insoluble tau aggregates. Student's t-test was used to

compare the percentage of tau aggregates in the perinuclear region for each light exposure pattern. The p-values in each

figure are as follows: NS: $p > 0.1$, *: $p < 0.05$, ***:$p < 0.001$, ****:$p < 0.0001$.

Results

Establishment of OptoTau knock-in (OptoTauKI) cells

OptoTau is a protein fused to CRY2Olig [21], which forms multimeric complexes when exposed to blue light, with

human 2N4R tau carrying the P301L mutation (Figure 1A). We have previously shown that transient overexpression of

OptoTau into Neuro2a and exposure to blue light can induce the clustering of tau [25]. In this study, we established

Neuro2a cells stably expressing OptoTau and investigated the effects of blue light illumination over a longer period

than in previous studies. OptoTau was introduced into the mouse Safe Harbor ROSA26 region using the CRISPR/Cas9

system. We designed the cells to express OptoTau under the CMV promoter and GFP for screening under the EF1α

promoter (Figure 1A). We isolated GFP-positive single colonies and defined them as OptoTau Knock-in (OptoTauKI)

cells. To confirm the expression and localization of OptoTau, we performed fluorescent labeling of SNAP-tag

incorporated into OptoTau. No signal was observed in non-genome-edited Neuro2a cells (WT) for both GFP and

SNAP647 labeling, whereas in OptoTauKI cells, GFP was observed throughout the cells, and SNAP-tag signal was

observed in the cytoplasm (Figure 1B). Live cell imaging with a confocal microscope revealed that OptoTau was

localized in the cytoplasm, avoiding the nucleus (Figure 1C). These results indicate that OptoTau is inserted in

OptoTauKI cells and that the expressed OptoTau resides mainly in the cytoplasm.

Formation of filament-like clusters of tau by 12-hour- blue light exposure

Next, we examined the behavior of OptoTau in OptoTauKI cells under blue light illumination. OptoTauKI cells

were continuously exposed to blue light for 12 hours using a blue LED in a CO_2 incubator, then tau localization by

SNAP staining or immunofluorescence staining. In OptoTauKI cells cultured in the dark, both the SNAP signal and the

signal of total tau visualized by the RTM38 antibody showed uniform distribution in the cytoplasm (Figure. 2A, Dark).

In contrast, in OptoTau cells exposed to blue light for 12 hours, OptoTau labeled with SNAP647 formed thick filament-

like structures, also recognized by RTM38 antibody (Figure. 2A, 12 h blue light).

Single-molecule localization microscopy has shown that tau forms oligomers on microtubules [27]. To determine

whether the filamentous tau formed by blue light illumination co-localizes with microtubules, we performed multicolor

immunofluorescence staining of tau and microtubules in the same cells. OptoTauKI cells illuminated with blue light for

12 hours were fixed with acetone methanol, and microtubules were labeled with α-tubulin antibody and tau with

RTM38 antibody (Figure. 2B). The signal of filamentous tau largely colocalized with that of α-tubulin (Figure. 2B,

Merge).

Furthermore, the expression level of tau in OptoTauKI cells was confirmed by Western blot using RTM38

antibody. Only a 115 kDa band was detected in Neuro2a cells (WT), while two bands of 115 kDa and 150 kDa were

detected in OptoTauKI cells (OptoTau) with or without blue light exposure. Based on the molecular weights, the 115

kDa band can be interpreted as the tau dimer and the 150 kDa band as the OptoTau, conjugation of CRY2olig, SNAP,

and tau. To examine whether the amount of OptoTau protein changes with or without blue light illumination, the

intensity of the 115 kDa and 150 kDa bands were quantified. Band intensities were normalized by β-actin, a loading

control, and calculated as "relative tau intensity" relative to the value of Neuro2a cells (WT). The relative tau intensity

of the 115 kDa band was not different between the cells with and without 12 hours of blue light exposure (Figure. 2D,

115 kDa), while the relative tau intensity of the 150 kDa band was higher in the blue light group in 5 out of the 6

samples. These results indicate that the amount of OptoTau protein in OptoTauKI cells was increased by 12-hour- blue

light illumination.

Tau dynamics in OptoTauKI cells after the termination of blue light illumination

Twelve hours of blue light irradiation induced filament-like structures in OptoTauKI cells (Fig. 2A, B). At the same

time, tau aggregates with large molecular weights, which could be oligomers, were not detected by Western Blot (Fig.

2C). To examine the stability of filamentous tau formed after 12 hours of blue light illumination, cells were fixed at 12,

24, 48, and 72 hours after termination of blue light. The subcellular localization of tau and microtubules was examined

by immunofluorescence staining (Fig. 3A). In OptoTauKI cells immediately after blue light termination, filament-like

tau co-localized with microtubules, and a strong signal was found in part of them (Fig. 3B, 0 h, arrowheads). Twenty-

four hours after the termination of blue light, a bright spot with dense tau was formed in the vicinity of the nucleus (Fig.

3B, 24 h, arrowheads). Bright spots were still present at 48 and 72 hours, but the tau bright spot was fragmented in most

cells. These results suggest that the filamentous tau formed in OptoTauKI cells after 12 h of blue light illumination is

not a stable structure but an unstable one that can dynamically change morphology over time.

Co-localization of tau with aggresome markers after blue light illumination

Proteasome inhibition and uptake of tau-paired helical protofilaments from AD patients have been shown to induce

aggresomes, inclusion bodies for misfolded proteins [28] [29]. Aggresomes are formed in the microtubule organizing

center (MTOC) located in the vicinity of the nucleus [30]. To investigate the possibility that the perinuclear tau

accumulation in OptoTauKI cells after 12 hours of exposure to blue light is an aggresome, we performed

immunofluorescence staining for γ-tubulin in the MTOC, together with tau. The tau accumulation in the vicinity of the

nucleus observed immediately and 24 hours after the cessation of blue light was co-localized with γ-tubulin in MTOCs

(Figure. 4A, arrowheads). This result indicates that the MTOC is the site of tau accumulation after blue light

termination.

The process of transporting degenerated proteins along the microtube to the aggresome at MTOCs involves the

microtubule motor protein dynein and histone deacetylase 6 (HDAC6), which recruits denatured proteins to dynein

[31]. To investigate the possibility that the solid bright spots of tau that appeared in OptoTauKI cells after blue light

arrest were contained in aggresome and its precursors, we examined the co-localization of these aggresome marker

proteins with tau. Tau signal co-localized with HDAC6 and dynein in the near-nuclear bright spots (arrowheads) and

thick filaments (Figure. 4B and C, 0 h). At 24 h after the cessation of blue light, tau showed stronger co-localization

with HDAC6, or dynein, near the nucleus and in the cytoplasm. These results suggest that tau proteins are recruited to

aggresome-associated proteins and ultimately accumulated to the aggresome in OptoTauKI cells with 12 h continuous

blue light exposure.

Western blot experiments showed an increase in OptoTau of 150 kDa after 12 hours of illumination (Figure. 2D).

Immunostaining with RTM38 antibody also showed cells with increased tau after 12 hours of light exposure compared

to dark conditions (Figure. 4D, top). Interestingly, cells with increased tau upon light illumination had increased levels

of HDAC6 protein (Figure. 4D, bottom). This result suggests that the increase in tau levels and filamentous tau induced

by light stimulation may activate the formation of aggresomes.

Insoluble tau aggregation induced by intermittent blue light illumination

Continuous blue light irradiation for 12 hours induced tau translocation to the aggresome, where denatured proteins

accumulated but did not form stable aggregates, as seen in AD and other tauopathy. We next searched for conditions

that would induce stable tau aggregation in OptoTauKI cells. To increase the probability of tau molecules interacting in

an optimal position, we expected repeated cluster formation and dissociation of OptoTau would be effective. Taking

advantage of the 23.1 min half-life for dissociation of the CRY2Olig cluster [21], we designed an illumination cycle in

which a 30-min blue light exposure is followed by a 30-min dark period during which the CRY2Olig interactions are

lost (Figure 5A). OptoTauKI cells express the OptoTau protein stably, allowing for prolonged blue light illumination,

which was impossible in transfected cells in previous studies. Therefore, we performed a 24-hour illumination sequence

consisting of a 16-hour dark period followed by eight 30-minute dark/30-minute blue light cycles (repeated

illumination), which was continued for 6 to 8 days (Figure. 5A). The stability of the tau aggregation-like structures

induced by blue light was evaluated based on whether they remained 16 hours after the illumination was terminated.

Total tau labeled with RTM38 antibody showed a thick linear filament structure immediately after repeated illumination

for 6 and 8 days (Figure. 5B, RTM38). This filament structure was not seen in OptoKI cells with a 6-day illumination

cycle plus 16 hours of darkness but remained in cells with an 8-day illumination cycle plus 16 hours of darkness. The

MC1 antibody is an antibody that recognizes the initial conformation of pathological tau [32] [33]. MC1 antibody also

recognized the filament-like conformation immediately after the termination of repeated illumination and after 16 hours

of illumination cycle for 8 days (Figure. 5B, MC1), as did the RTM38 antibody. The percentage of cells with bright

solid spots showing stable tau aggregation 16 hours after termination of repeated illumination was significantly higher

after 8 days of illumination than after 6 days (Figure. 5C).

Taking advantage of that methanol fixation had the effect of removing soluble tau in the cytoplasm [26], we

examined whether repeated illumination to OptoTauKI cells results in the formation the insoluble tau. Methanol-fixed

cells immediately after 6 days of repeated irradiation showed RTM38-positive aggregation in the cytoplasm, but this

aggregation disappeared 16 hours later (Fig. 5D, Top). Immediately after 8 days of repeated illumination, the amount of

insoluble tau remaining after methanol fixation was increased compared to that after 6 days of illumination and was

detected as thick filament structures (Fig. 5D, 8 days). This filament-like tau remained as a methanol-insoluble structure

after 16 hours (Fig. 5D, 8 days+16 h). The results indicate that repeated blue light exposure and dark incubation over 8

days can produce insoluble tau aggregation.

Acceleration of insoluble tau aggregation formation by short light-dark cycles

Finally, we examined whether the temporal pattern of blue light/dark cycle repetitions affected the time to the induction

of insoluble tau aggregation. Here, 25 min dark/5 min blue light illumination cycles were repeated 48 times (Figure

6A). The total required time was 24 hours, with a total illumination time of 4 hours. A 25 min dark/5 min blue light

cycle for 24 h formed thick tau filaments as detected by RTM38 and MC1 antibodies (Figure 6, top). MC1-positive tau

filaments remained after 16 hours of dark incubation after the illumination cycle (Figure 6B, bottom). The insolubility

of filamentous tau in OptoTauKI cells immediately after the end of the cycle and after 16 hours of dark incubation after

the cycle was also supported by immunofluorescent staining after methanol fixation (Figure 6C). These results indicate

that insoluble tau could be induced in only 24 hours by repeating the 25 min dark/5 min blue light exposure cycle. The

same 48 cycles of light illumination did not induce stable tau aggregation when 8 set cycles of 30 min dark / 30 min

blue light were repeated for 6 days; despite the longer duration of each illumination (30 min vs. 5 min), total

illumination time (24 h vs. 4 h), and total time required (6 days vs 24 hours) (Figure 5). Furthermore, fibrous tau

aggregation was hardly formed by 144 cycles of 1 min blue light/9 min dark illumination, despite the high number of

cycles (data not shown). The results indicate that the temporal pattern, including the duration of one illumination,

interval time, and number of repetitions, must be set appropriately to introduce insoluble aggregation with OptoTau.

Interestingly, insoluble tau aggregates induced by the 5 min light/25 min dark cycle were less frequently located

near the nucleus where aggresomes were formed (Figure 4). In OptoTauKI cells illuminated continuously for 12 h, by

which accumulation of tau in the aggresome was observed, an average of $69.64 \pm 2.90\%$ (n=9) of the tau aggregates

were located in the nuclear vicinity. On the other hand, in cells illuminated for 24 h with 5 min light/25 min dark cycles,

significantly less tau aggregation was found in the vicinity of the nucleus ($34.54 \pm 4.06\%$ (n=6)) (Figure 6D, $p<0.001$,

student's t-test). This result suggests that the insoluble tau induced by the 5 min blue light/25 min dark cycle escaped

translocation to aggresomes.

Discussion

In the present study, we generated cells knocking in OptoTau, an optogenetic tool based on the CRY2Olig module,

and established a technique to manipulate tau aggregation with blue light. We found light illumination conditions that

specifically induce tau accumulation in aggresomes or insoluble tau aggregation.

Recently, a series of studies have reported that transient overexpression of chimeric proteins combining tau with

CRY2 or its derivative CRY2Olig, i.e., families of OptoTau, results in the blue light-dependent formation of LLPS

droplets and tau oligomers [23] [24] [25]. In the present study, we manipulated tau by blue light in OptoTauKI cells, in

which OptoTau was inserted into the genome by genome editing. In dividing cells, the transfected plasmid is diluted by

cell division, and the expression level of the product is reduced. Here, using a cell line that stably expresses OptoTau

enabled us to validate the blue light illumination protocol over several days while maintaining OptoTau expression.

When SH SY5Y and HeLa cells overexpressing CRY2WT-mCherry-Tau, a protein fusing CRY2 and human wild-

type tau in a previous study, were exposed to blue light, tau condensate formed along microtubules within 50 seconds

[23]. Our result that tau formed filament structures colocalizing with microtubules in OptoTauKI cells exposed to blue

light for 12 hours is consistent with this report. On the other hand, exposure to blue light for 3 min to primary cortical

culture neurons overexpressing "Tau::CRY2", which is CRY2Olig fused to the C-terminus of human wild-type tau,

induced droplets and oligomers of tau [24]. In Neuro2a cells overexpressing our OptoTau, consisting of CRY2Olig-

Snap-human Tau with P301L, such a droplet of tau, which is considered to be an LLPS droplet, was induced by 24

hour- blue light illumination in the presence of microtubule depolymerizing agent nocodazole [25]. However, no

droplet-like structures were induced in OptoTauKI cells by any light illumination protocols in the present study. The

differences in the morphology of the OptoTau complex induced by blue light may depend on multiple parameters,

including the expression levels of OptoTau, endogenous tau in the cells, blue light exposure time, exposure intensity,

cell type, and microtubule integrity.

Some filamentous tau induced by 12 hours of blue light illumination on OptoTauKI accumulated in MTOCs

stained with γ-tubulin and co-localized with HDAC6 and dynein, both component proteins of aggresomes. OptoTau

accumulation in aggresomes was also observed in Neuro2a cells overexpressing OptoTau [25]. The result that HDAC6

amount is increased in cells with increased tau content implies that tau degradation is triggered by selective autophagy.

In this study, we did not confirm the up-regulation of tau ubiquitination, which is a signal for transport to aggresome. It

also remains to be verified whether aggresome activity is increased after light stimulation, leading to degradation of

OptoTau and endogenous Tau. Previous studies have shown that phosphorylated tau and tau oligomers are increased in

the hippocampus of mice in which the aggresomal protein p62 is knocked out [34]. Optogenetically accumulated

OptoTau co-localizes with endogenous p62 [25].　The role of p62 in light-induced aggresome formation and tau

degradation needs to be further verified.

In this study, we found two illumination conditions induce methanol-insoluble tau aggregation: one is to repeat a

24-hour illumination sequence consisting of a 16-hour dark period followed by eight 30 min dark/30 min blue light

cycles for 8 days, and another is repeating 25 min dark/5 min blue light cycles for 24 hours. This aggregation did not

disappear even 16 hours after the termination of light illumination. Therefore, these tau aggregates are interpreted as a

stable residual structure.　Initial conformational changes in pathological tau labeled by the MC1 antibody were detected

even before stable aggregation was generated but only remained when methanol-insoluble tau aggregation was present.

In a previous study, sarcosyl-insoluble tau aggregation could be induced optogenetically by OptoTau with truncated N-

terminal of tau. Still, OptoTau with full-length tau did not induce insoluble tau aggregation [25]. Since the N-terminal

of tau inhibits polymerization of full-length tau [35], it is likely that OptoTau lacking its N terminal is more likely to

produce aggregation seed. The present study indicates that even full-length tau could be transformed into stable

aggregation by light illumination with repeated On/Off cycles. Opto-α-synuclein, α-synuclein fused with CRY2Olig,

was also induced to form stable aggregates in human iPS cell-derived neurons by repeated stimulation at 0.17 Hz.

Repeated on/off stimulation and periods of no light exposure are effective means of avoiding phototoxicity. In addition,

it may be necessary to repeatedly stimulate the CRY2Olig module to induce stable aggregation. In particular, OptoTau

is a microtubule-binding protein that can be trapped in microtubules, making it difficult to interact with each other at

optimal positions. The frequent concentration of OptoTau by blue light followed by dissociation during the dark period

may increase the possibility of interaction between tau molecules at appropriate positions. Indeed, by accelerating the

cycles of light illumination, stable aggregation could be induced in a shorter period (8 days to 24 hours), even though

the total time of blue light illumination was shorter (24 hours to 4 hours). This result suggests that tau aggregation can

be induced more efficiently when CRY2Olig binding/dissociation is repeated rapidly. This study only demonstrated two

conditions under which stable induction of tau aggregation is possible, and the optimal aggregation induction system

should be further explored.

In this study, we showed that OptoTau could induce opposing behaviors such as stable tau aggregation and

aggresome formations, i.e., proteolysis and protein aggregation, depending on the pattern of light stimulation. OptoTau

presents a new experimental possibility for future introduction into the mouse brain, where the assembly state of tau can

be manipulated with light at the intended timing in specific cells at the targeted brain region. This optogenetic tool

could also help to validate the existence of "singularity cells," i.e., a small number of rare cells that cause significant

changes throughout the system [18]. In Alzheimer's disease, the neurons that first form tau aggregates in LCs and ECs

can be assumed to be "singularity cells" [20]. If OptoTau can specifically induce tau aggregation only in LCs and ECs,

it will be possible to verify whether they can be "singularity cells" that spread tau pathology to the cortex.

Some issues need to be resolved before OptoTau can be used as a model for tauopathies in the future. One is the

isoform of tau. The tau isoform with the 31 amino acid repeats encoded by exon 10 is called the 4R isoform, and the

isoform without the repeats is called the 3R. The tau aggregates of AD and CTE contain 3R and 4R tau, while Pick's

disease is composed of only 3R tau, and CBD and PSP are composed of only 4R tau. The molecular structure of the

filaments within tau aggregates differs in each tauopathic disease [10]. The molecular structure of the tau aggregation

core is the determinant of the tau aggregate properties (including the ratio of 3R tau to 4R tau) specific to tauopathies

[36]. Current OptoTau includes only 4R-type tau. In the future, it will be necessary to develop 3R-type OptoTau to

create optogenetic models specific to AD, CTE, and Pick's disease. The 3R to 4R tau expression ratio in OptoTau-

induced cells should be controlled. In addition, photogenetically inducing aggregation nuclei specific to each tauopathy

will be a significant challenge.

Conclusion

This study used the optogenetic tool OptoTau, based on the CRY2Olig module, to induce tau aggregation in

cells. Specific light exposure conditions introduced aggresome and insoluble aggregation in OptoTauKI cells. The

present results will provide a new technological basis for creating a singular point of tau aggregation at a targeted time

in targeted cells.

Conflict of Interest

The authors have no conflict of interest related to this publication.

Author Contributions

SS, TU, NK, BX, AH, and YS performed the experiment. SY, AT, HY, GM, and HB were involved in the design and

supervision of this work. SS, TU, NK, BX, AH, and HB processed the experimental data, performed the analysis,

drafted the manuscript, and designed the figures. All authors discussed the results and commented on the manuscript.

Data Availability

The evidence data generated and/or analyzed during the current study are available from the corresponding author upon

reasonable request.

Acknowledgments

The authors thank Prof. Peter Davies (Albert Einstein College of Medicine) for providing the MC1 antibody, Dr Sumiyo

Mimura (AS ONE Corporation) for technical support of single cell cloning, and Dr. Naruhiko Sahara (National

Institutes for Quantum Science and Technology) for insightful discussion. This research was funded by Grant-in-Aid for

Scientific Research on Innovative Areas "Singularity Biology" (JP18H05410 to HY, JP18H05414 to HB) from the

Ministry of Education, Culture, Sports, Science and Technology, Japan, JSPS Grants-in-Aid for Scientific Research

JP20K06896, JP23K05993 to YS, JP21H02450, 23K18116 to HB. and medical research continuous grants from Takeda

Science Foundation to HB. DeepL, ChatGPT 3.5, and Grammarly supported the English editing of this article.

References

[1] Gotz, J., Halliday, G., Nisbet, R.M. Molecular pathogenesis of the tauopathies. Annu Rev Pathol 14, 239-261 (2019). https://doi.org/10.1146/annurev-pathmechdis-012418-012936

[2] Alquezar, C., Arya, S., Kao, A.W. Tau post-translational modifications: Dynamic transformers of tau function, degradation, and aggregation. Front Neurol 11, 595532 (2020). https://doi.org/10.3389/fneur.2020.595532

[3] Braak, H., Del Tredici, K. The pathological process underlying alzheimer's disease in individuals under thirty. Acta Neuropathol 121, 171-181 (2011). https://doi.org/10.1007/s00401-010-0789-4

[4] Braak, H., Thal, D.R., Ghebremedhin, E., Del Tredici, K. Stages of the pathologic process in alzheimer disease: Age categories from 1 to 100 years. J Neuropathol Exp Neurol 70, 960-969 (2011). https://doi.org/10.1097/NEN.0b013e318232a379

[5] Ossenkoppele, R., Schonhaut, D.R., Schöll, M., Lockhart, S.N., Ayakta, N., Baker, S.L., et al. Tau pet patterns mirror clinical and neuroanatomical variability in alzheimer's disease. Brain 139, 1551-1567 (2016). https://doi.org/10.1093/brain/aww027

[6] Cho, H., Choi, J.Y., Hwang, M.S., Kim, Y.J., Lee, H.M., Lee, H.S., et al. In vivo cortical spreading pattern of tau and amyloid in the alzheimer disease spectrum. Annals of Neurology 80, 247-258 (2016). https://doi.org/https://doi.org/10.1002/ana.24711

[7] Allen, B., Ingram, E., Takao, M., Smith, M.J., Jakes, R., Virdee, K., et al. Abundant tau filaments and nonapoptotic neurodegeneration in transgenic mice expressing human p301s tau protein. J Neurosci 22, 9340-9351 (2002). https://doi.org/10.1523/jneurosci.22-21-09340.2002

[8] Santacruz, K., Lewis, J., Spires, T., Paulson, J., Kotilinek, L., Ingelsson, M., et al. Tau suppression in a neurodegenerative mouse model improves memory function. Science 309, 476-481 (2005). https://doi.org/10.1126/science.1113694

[9] Yoshiyama, Y., Higuchi, M., Zhang, B., Huang, S.M., Iwata, N., Saido, T.C., et al. Synapse loss and microglial activation precede tangles in a p301s tauopathy mouse model. Neuron 53, 337-351 (2007). https://doi.org/10.1016/j.neuron.2007.01.010

[10] Sahara, N., Yanai, R. Limitations of human tau-expressing mouse models and novel approaches of mouse modeling for tauopathy. Front Neurosci 17, 1149761 (2023). https://doi.org/10.3389/fnins.2023.1149761

[11] Iba, M., Guo, J.L., McBride, J.D., Zhang, B., Trojanowski, J.Q., Lee, V.M. Synthetic tau fibrils mediate transmission of neurofibrillary tangles in a transgenic mouse model of alzheimer's-like tauopathy. J Neurosci 33, 1024-1037 (2013). https://doi.org/10.1523/jneurosci.2642-12.2013

[12] Clavaguera, F., Akatsu, H., Fraser, G., Crowther, R.A., Frank, S., Hench, J., et al. Brain homogenates from human tauopathies induce tau inclusions in mouse brain. Proc Natl Acad Sci U S A 110, 9535-9540 (2013). https://doi.org/10.1073/pnas.1301175110

[13] Bell, B.J., Malvankar, M.M., Tallon, C., Slusher, B.S. Sowing the seeds of discovery: Tau-propagation models of alzheimer's disease. ACS Chem Neurosci 11, 3499-3509 (2020). https://doi.org/10.1021/acschemneuro.0c00531

[14] Narasimhan, S., Guo, J.L., Changolkar, L., Stieber, A., McBride, J.D., Silva, L.V., et al. Pathological tau strains from human brains recapitulate the diversity of tauopathies in nontransgenic mouse brain. J Neurosci 37, 11406-11423 (2017). https://doi.org/10.1523/JNEUROSCI.1230-17.2017

[15] Vogels, T., Leuzy, A., Cicognola, C., Ashton, N.J., Smolek, T., Novak, M., et al. Propagation of tau pathology: Integrating insights from postmortem and in vivo studies. Biol Psychiatry 87, 808-818 (2020). https://doi.org/10.1016/j.biopsych.2019.09.019

[16] Calafate, S., Buist, A., Miskiewicz, K., Vijayan, V., Daneels, G., de Strooper, B., et al. Synaptic contacts enhance cell-to-cell tau pathology propagation. Cell Reports 11, 1176-1183 (2015). https://doi.org/https://doi.org/10.1016/j.celrep.2015.04.043

[17] Pérez, M., Avila, J., Hernández, F. Propagation of tau via extracellular vesicles. Frontiers in Neuroscience 13 (2019). https://doi.org/10.3389/fnins.2019.00698

[18] Nagai, T. Editorial: Singularity biology and beyond. Biophysics and Physicobiology advpub (2024). https://doi.org/10.2142/biophysico.bppb-v21.s013

[19] Ichimura, T., Kakizuka, T., Sato, Y., Fujioka, Y., Ohba, Y., Horikawa, K., et al. Strength in numbers: Unleashing the potential of trans-scale scope amateras for massive cell quantification. Biophysics and Physicobiology advpub (2024). https://doi.org/10.2142/biophysico.bppb-v21.s017

[20] Bannai, H., Takashima, A., Soeda, Y., Yoshimura, H., Matsumoto, G., Sahara, N., et al. Research on the molecular mechanism of singularity phenomenon in neurological disorders. Biophysics and Physicobiology

advpub (2024). https://doi.org/10.2142/biophysico.bppb-v21.s008

[21] Taslimi, A., Vrana, J.D., Chen, D., Borinskaya, S., Mayer, B.J., Kennedy, M.J., et al. An optimized optogenetic
clustering tool for probing protein interaction and function. Nat Commun 5, 4925 (2014).
https://doi.org/10.1038/ncomms5925

[22] Tiwari, P., Tolwinski, N.S. Using optogenetics to model cellular effects of alzheimer's disease. Int J Mol Sci 24
(2023). https://doi.org/10.3390/ijms24054300

[23] Zhang, X., Vigers, M., McCarty, J., Rauch, J.N., Fredrickson, G.H., Wilson, M.Z., et al. The proline-rich domain
promotes tau liquid-liquid phase separation in cells. J Cell Biol 219 (2020).
https://doi.org/10.1083/jcb.202006054

[24] Jiang, L., Lin, W., Zhang, C., Ash, P.E.A., Verma, M., Kwan, J., et al. Interaction of tau with hnrnpa2b1 and
n(6)-methyladenosine rna mediates the progression of tauopathy. Mol Cell 81, 4209-4227 e4212 (2021).
https://doi.org/10.1016/j.molcel.2021.07.038

[25] Soeda, Y., Yoshimura, H., Bannai, H., Koike, R., Takashima, A. Intracellular tau fragment droplets serve as
seeds for tau fibrils. bioRxiv, 2023.2009.2010.557018 (2023). https://doi.org/10.1101/2023.09.10.557018

[26] Guo, J.L., Narasimhan, S., Changolkar, L., He, Z., Stieber, A., Zhang, B., et al. Unique pathological tau
conformers from alzheimer's brains transmit tau pathology in nontransgenic mice. J Exp Med 213, 2635-2654
(2016). https://doi.org/10.1084/jem.20160833

[27] Gyparaki, M.T., Arab, A., Sorokina, E.M., Santiago-Ruiz, A.N., Bohrer, C.H., Xiao, J., et al. Tau forms
oligomeric complexes on microtubules that are distinct from tau aggregates. Proc Natl Acad Sci U S A 118
(2021). https://doi.org/10.1073/pnas.2021461118

[28] Guthrie, C.R., Kraemer, B.C. Proteasome inhibition drives hdac6-dependent recruitment of tau to aggresomes.
J Mol Neurosci 45, 32-41 (2011). https://doi.org/10.1007/s12031-011-9502-x

[29] Santa-Maria, I., Varghese, M., Ksiezak-Reding, H., Dzhun, A., Wang, J., Pasinetti, G.M. Paired helical filaments
from alzheimer disease brain induce intracellular accumulation of tau protein in aggresomes. J Biol Chem 287,
20522-20533 (2012). https://doi.org/10.1074/jbc.M111.323279

[30] Kopito, R.R. Aggresomes, inclusion bodies and protein aggregation. Trends Cell Biol 10, 524-530 (2000).
https://doi.org/10.1016/s0962-8924(00)01852-3

[31] Kawaguchi, Y., Kovacs, J.J., McLaurin, A., Vance, J.M., Ito, A., Yao, T.-P. The deacetylase hdac6 regulates
aggresome formation and cell viability in response to misfolded protein stress. Cell 115, 727-738 (2003).
https://doi.org/https://doi.org/10.1016/S0092-8674(03)00939-5

[32] Jicha, G.A., Bowser, R., Kazam, I.G., Davies, P. Alz-50 and mc-1, a new monoclonal antibody raised to paired
helical filaments, recognize conformational epitopes on recombinant tau. J Neurosci Res 48, 128-132 (1997).
https://doi.org/10.1002/(sici)1097-4547(19970415)48:2<128::aid-jnr5>3.0.co;2-e

[33] Jicha, G.A., Lane, E., Vincent, I., Otvos, L., Jr., Hoffmann, R., Davies, P. A conformation- and phosphorylation-
dependent antibody recognizing the paired helical filaments of alzheimer's disease. J Neurochem 69, 2087-2095
(1997). https://doi.org/10.1046/j.1471-4159.1997.69052087.x

[34] Ono, M., Komatsu, M., Ji, B., Takado, Y., Shimojo, M., Minamihisamatsu, T., et al. Central role for p62/sqstm1
in the elimination of toxic tau species in a mouse model of tauopathy. Aging Cell 21, e13615 (2022).
https://doi.org/10.1111/acel.13615

[35] Horowitz, P.M., LaPointe, N., Guillozet-Bongaarts, A.L., Berry, R.W., Binder, L.I. N-terminal fragments of tau
inhibit full-length tau polymerization in vitro. Biochemistry 45, 12859-12866 (2006).
https://doi.org/10.1021/bi061325g

[36] Hosokawa, M., Masuda-Suzukake, M., Shitara, H., Shimozawa, A., Suzuki, G., Kondo, H., et al. Development
of a novel tau propagation mouse model endogenously expressing 3 and 4 repeat tau isoforms. Brain 145, 349-
361 (2021). https://doi.org/10.1093/brain/awab289

Figure Legends

Figure 1: Neuro2a knock-in Neuro2a cells

(A) Top: design of OptoTau, which consists of CRY2Olig, SNAP-tag, and human2N4R tau with P301L mutation/

Bottom: conceptual diagram of OptoTau knock-in to mouse ROSA26 locus using CRISPR/Cas9 system. The donor

vector contains OptoTau expressed under the CMV promoter and GFP and Puromycin (Puro) sequences for screening.

(B) Live cell imaging of GFP and SNAP647 fluorescence in Neuro2A cells (WT) and OptoTau knock-in Neuro2a cells

(OptoTauKI). The SNAP647 signal reflects the localization of OptoTau. Transmission images (Trans) were taken

from the same field of view. Bar: 10 μm.

(C) Confocal images of GFP and SNAP647 fluorescence in Neuro2A cells (WT) and Neuro2a cells knocking in OptoTau

knock-in (WT). Cells were fixed with 4% PFA. Bar: 10 μm.

Figure 2: Effect of 12 hours of blue light illumination on Tau in OptoTau knock-in cells.

(A) Subcellular localization of GFP (left), OptoTau (middle, SNAP647), and total Tau shown by immunofluorescence

staining with RTM38 antibody in OptoTau Knock-in cells. Top: cells kept in the dark condition (Dark, D); bottom:

cells fixed immediately after 12 h blue light (12 h blue light, L). Scale bar: 10 μm.

(B) Immunofluorescence staining images of microtubules (middle, α-tubulin) and total Tau (left, RTM38) in OptoTauKI

cells immediately after 12 hours of blue light illumination. The right panel shows the merge of these signals (a-

tubulin: green, total Tau: magenta). The scale bar represents 10 μm.

(C) Western blotting analysis of Tau using RTM38 antibody in Neuro2a cells (WT) and OptoTauKI cells (OptoTauKI).

D: cells under dark; L: cells illuminated with blue light for 12 h. 151 kDa and 115 kDa bands were detected. β-actin

was used as a loading control.

(D) Quantification of 115 kDa (left) and 151 kDa (right) tau. The ratio of β-actin-normalized tau band intensity to that

of Neuro2a cells (WT) was defined as relative Tau intensity. D: under dark, L: 12 h blue light exposure. NS: $p>0.1$,

*: $p<0.05$, Mann-Whitney U-test.

Figure 3: Localization change of Tau after cessation of blue light illumination

(A) Schematic illustration of the experimental design. Top: Localization of total Tau in OptoTauKI cells cultured in the

dark after 12 h of blue light illumination was verified by immunofluorescence staining with the RTM38 antibody;

the arrowhead indicates the timing of cell fixation. Dark indicates cells cultured in the dark for 12 h.

(B)   Immunofluorescence images of total tau and microtubules immediately (0 h), 24 h, 48 h, and 72 h after cessation of

blue light exposure. Top: RTM38 antibody, middle: α-tubulin antibody, bottom: Merge image. Yellow arrowheads

indicate large bright spots of Tau; arrows indicate fragmented bright spots of Tau. Scale bar: 10 μm.

Figure 4: Co-localization of tau with aggresome proteins after cessation of blue light illumination.

Multicolor immunofluorescence images of Tau and aggresomal proteins in OptoTauKI cells immediately (0 h) and 24 h

after stopping 12 h blue light illumination. Aggresomal proteins labeled were (A) γ-tubulin, (B) HDAC6, and (C)

Dynein. (D) Immunostaining of Tau (RTM38, top) and HDAC6 (bottom) under dark conditions (Dark) and

immediately after 12 h of blue light exposure (Blue light). rrowheads show strong bright spots of Tau. he scale Bar is 10

μm.

Figure 5: Formation of insoluble tau aggregates by repeated blue light illumination.

(A)   Protocol for repeated blue light stimulation. OptoTauKI cells were cultured in darkness for 16 hours, followed by 8

cycles of 30 min darkness/30 min blue light illumination, repeated for N days. Arrows indicate the timing of fixation.

(B)   Immunofluorescence staining using total tau antibody (RTM38) and tau conformation change recognition antibody

(MC1) in OptoTauKI cells after repeated blue light stimulation. Cells cultured in the dark for 6 days (6 days Dark),

cells fixed immediately after repeated blue light stimulation on day 6 (6 days), cells fixed at 16 hours after day 6

light stimulation (6 day+16 h), cells fixed immediately after repeated blue light stimulation on day 8 (8 days), cells

fixed at 16 hours after day 8 light stimulation (8 day+16 h). Bar: 10 μm.

(C)   Percentage of cells with insoluble tau aggregation 16 h after repeated blue light exposure. The red lines indicate

median. 6 days+16 h: N=46 cells; 8 days+16 h: N=31 cells. ****: $p < 0.0001$, Mann-Whitney U-Test.

(D)   Visualization of intracellular insoluble Tau by methanol fixation at 6 days and 6 days+16 h (top), 8 days and 8

days+16 h (bottom), using RTM38 antibody. Scale bar is 10 μm.

Figure 6: Acceleration of insoluble tau aggregation formation by repeated blue light illumination.

(A) Protocol for repeated blue light stimulation. The cycle was repeated for 48 consecutive cycles (24 h), each cycle

consisting of 25 min of darkness and 5 min of blue light illumination and. Arrows indicate the timing of fixation.

(B) Immunofluorescence staining using total tau antibody (RTM38) and tau conformation change recognition antibody

(MC1) in OptoTauKI cells after repeated blue light stimulation. PFA fixation was performed immediately after the

end of 48 cycles (48 cycles + 0 h) or 16 h after the end of 48 cycles (48 cycles + 16 h) in the blue light (Blue light)

and dark condition (Dark) groups, respectively. Bar: 10 μm.

(C) Visualization of intracellular insoluble Tau by methanol fixation. Methanol fixation was performed immediately after

the end of 48 cycles (48 cycles + 0 h) or 16 h after the end of 48 cycles (48 cycles + 16 h) in the blue light (Blue

light) and dark condition (Dark) groups, respectively. Scale bar is 10 μm.

(D) Comparison of the percentage of cells with perinuclear insoluble Tau accumulation of the single 12 h blue light

illumination (BL12 h x 1) and the 48 repetitions of 5 min off 25 min blue light illumination [(BL5m + DK25m) x

1]. Fixation was performed immediately after completion of the total blue light illumination program in both

groups. The red lines indicate median. BL12 h x 1: N=9 coverslips; (BL5m + DK25m) x 1: N=6 coverslips. ***:

$p < 0.001$, Student's t-test.

(A)

OptoTau

Cry2Olig — SNAP — Tau (human 2N4R) — P301L

Rosa26 HA left — OptoTau — GFP — Puro — Rosa26 HA right
CMV — EF1α — TA

Exon1 — ROSA26 gDNA — Cas9 — Exon2

(B)

	WT	OptoTauKI
GFP		
SNAP647		
Trans		

(C)

	WT	OptoTauKI
GFP		
SNAP647		
Merge		

Figure 1

(A)

GFP   SNAP647   Total Tau (RTM38)

Dark (D)

12 h blue light (L)

(B)

RTM38   α-Tubulin   RTM38 / α-Tubulin

(C)

WT   OptoTauKI

D   D   L

Total tau (RTM38)

(kDa)

◄ 150

◄ 115

β-actin

(D)

115 kDa

NS

150 kDa

*

Relative tau intensity

D   L   D   L

Figure 2

Figure 3

Figure 4

(A)

0 day   1 day   2 days   N days   +16 h

Repeated illumination

30 m | 30 m

8 cycles

(B)

| | 6 days Dark | 6 days | 6 days +16 h | 8 days | 8 days +16 h |

RTM38

MC1

(C)

Ratio of cell with aggregation

****

6 days +16 h   8 days +16 h

(D)

6 days | 6 days +16 h

RTM38

8 days | 8 days +16 h

RTM38

Figure 5

Figure 6