Title: Age-specific dynamics of neutralizing antibodies, cytokines, and chemokines in

response to La Crosse virus infection in mice

Authors: Reem Alatrash[1,2], Bobby Brooke Herrera[1,2*]

Affiliations:

[1]Rutgers Global Health Institute, Rutgers University, New Brunswick, NJ, USA

[2]Department of Medicine, Division of Allergy, Immunology, and Infectious Diseases and

Child Health Institute of New Jersey, Rutgers Robert Wood Johnson Medical School, New

Brunswick, NJ, USA

Corresponding Author: BBH, bherrera@globalhealth.rutgers.edu

Running Title: Age-specific cytokines and chemokines following LACV infection

Abstract

La Crosse virus (LACV) is a primary cause of pediatric arboviral encephalitis in the United States, with a notable impact on children aged 16 years or younger. This age-related susceptibility extends to murine models, where weanling mice (≤3 weeks old) succumb to LACV-induced disease, while adults (≥6 weeks old) demonstrate resistance. Despite its clinical significance, our understanding of host responses to LACV remains relatively unexplored. This study aims to elucidate the dynamics of neutralizing antibodies (nAbs), cytokines, and chemokines following LACV infection in both weanling and adult mice. Utilizing a highly infectious dose of LACV, our study reveals age-specific variations in viral titers, neutralizing antibody titers, and survivability. Weanling mice exhibit an early disease onset coupled with heightened peripheral viremia and nAbs compared to adults. Serum cytokine and chemokine profiling showed distinct kinetics and age-specific responses. Notably, adult mice exhibit significantly elevated levels of Th1 cytokines (IFN-γ, IL-18, GM-CSF, IL-1β and IL-12), juxtaposed with elevated levels of Th2 cytokines (IL-6 and IL-4) observed in weanling mice, often coinciding with the onset of symptoms. Furthermore, the heightened levels of Th9/Th17/Th22/Treg cytokines (IL-9, Il-10, IL-17A, IL-22, IL-23, and IL-27) and chemokines (CCL11, CCL3, and CCL5) in infected adult mice underscores their role in mounting protective immune responses against LACV. In conclusion, our study associates cytokines and chemokines with protective immunity in adult mice, contrasting with the pathogenesis observed in weanlings. This work emphasizes the need for further studies into mechanisms of innate and adaptive immune responses to LACV infection.

Introduction

La Crosse Virus (LACV) is a single-stranded, negative-sense RNA virus belonging to the *Peribunyaviridae* family in the order *Bunyavirales* (1). Its transmission to humans primarily occurs through the bite of infected Eastern Tree Hole mosquitoes (*Aedes triseriatus*), although other mosquito species may harbor LACV, potentially contributing to the emergence of disease (2). Upon entry into the host, LACV initially replicates in skin muscle cells, facilitating its spread into various tissues and organs (3, 4). While the majority of LACV-infected individuals manifest only mild, febrile symptoms, a subset experiences severe neurological complications, including altered behavior, seizures, coma, or fatality, as the virus invades the central nervous system (CNS) (5). Notably, severe disease and neurological manifestations of LACV are predominantly observed in children under the age of 16 years, indicating a higher susceptibility among this age group (6).

The clinical significance of LACV, coupled with its ability to also induce disease in murine models, has designated it as a prototype for studying viruses within the *Peribunyaviridae* family. Weanling mice (≤3 weeks old) exhibit susceptibility to LACV infection, whereas adult mice (≥6 weeks old) display resistance, mirroring the age-related susceptibility observed in humans (3, 7, 8). Unraveling the intricacies of host responses to LACV infection is important, particularly in delineating factors contributing to disease resistance or susceptibility. It is believed that the peripheral immune system's control of LACV access to the CNS plays a pivotal role in disease resistance in adult mice (9).

Prior investigations have shed light on the significance of both innate and adaptive immune responses following LACV infection (9, 10). Type I interferon responses have

garnered considerable attention, demonstrating their indispensable role in curbing LACV replication and mitigating disease severity (11). Early cellular innate immune responses, particularly those imparted by myeloid dendritic cells, proved critical for protection in wild type adult mice infected with LACV (10). The same study also showed that treatment of young wild type mice with type I interferon, particularly IFN-β resulted in protection mediated by myeloid dendritic cells (10). Moreover, the depletion of adaptive immune responses, including both B and T cells, resulted in neurological disease in adult mice infected with LACV, suggesting that these cell types are essential for protection against neuropathology and death (9). However, the broader immune response including cytokines and chemokines remains relatively unexplored. These signaling molecules, produced by various cell types in response to viral infections, orchestrate immune responses and exert influences on disease outcomes (12-15). Thus, deciphering the dynamics of cytokines and chemokines during LACV infection is important for unraveling their contributions to age-specific protection or pathogenesis.

Cytokines and chemokines undergo substantial modulation upon LACV infection across diverse cell types. Notably, human neuron-astrocyte co-cultures respond to LACV infection in vitro within 48 hours post infection by eliciting robust proinflammatory cytokines and chemokines including: IFN-γ, IL-6, TNF-α, IL-8, CCL4, CCL5, CXCL10 (16). In a separate study, glial cells derived from the brains of young LACV-infected mice produced significant levels of IFN-β (17). Furthermore, preliminary observations hint at a proinflammatory cytokine response in both the periphery and the brain following LACV infection, with elevated levels of IL-6 and IL-12p40, which were reported to be in comparable levels in young and adult mice (10, 18). Interestingly, administration of IL-12

or GM-CSF encoded by a plasmid prior to LACV infection in Ifnar1$^{-/-}$ mice was shown to

increase survival rates, suggesting a potential protective role of these cytokines (19). A

similar proinflammatory cytokine profile has also been reported with Oropouche virus, a

distantly related bunyavirus (20). Cytokines and chemokines seem to act as important

contributing factors in the immune response to LACV and other bunyavirus infections.

Given the important implications of cytokines and chemokines during LACV

infection, this study aims to address a knowledge gap by investigating the involvement of

these molecules in age-related susceptibility to LACV infection and their correlation with

other components of peripheral immune responses such as neutralizing antibodies

(nAbs). To address this knowledge gap, we present an in-depth analysis of cytokine and

chemokine levels in the sera of adult and weanling mice throughout the course of a highly

infectious dose of LACV infection. We identify age-specific differences in cytokine and

chemokine profiles between adult and weanling mice. Moreover, we establish correlations

between cytokine and chemokine levels and indicators of viral clearance and neutralizing

antibody responses.

Results

Age-related susceptibility is maintained despite a highly infectious dose of LACV

infection

An unpublished observation reported no significant difference in cytokine levels

between adult and weanling mice infected with a 1000 PFU dose of LACV (10). To

investigate the potential impact of infectious dose on immune system responsiveness to

LACV infection, we hypothesized that a higher dose might better reveal differences in

cytokine levels if they exist. Previous studies have shown that infection with 10^5 PFU of

LACV maintains age-related susceptibility in mice (10). Here, we aimed to assess the

effects of an infectious dose five-fold higher than the previously reported dose by

monitoring the development of neurological signs in infected mice and measuring viral

titers in peripheral blood.

117   Administration of a $5x10^5$ PFU dose of LACV demonstrated a similar difference in

survivability between adult and weanling mice, reinforcing the age-related susceptibility

to LACV. Adult mice exhibited markedly higher survival rates compared to weanling mice

(p < 0.001), with approximately 50% of adults surviving up to 12 days post-infection (dpi)

(Fig. 1A). Weanling mice began exhibiting signs of neurological disease starting at 4-5

dpi (Fig. 1B). Our findings show that the higher dose of LACV infection in adult and

weanling mice results in a timeline of symptom/disease onset like that of lower infectious

doses that have historically been used (Fig. 1B). Consistent with previous reports, viral

titers in sera were detectable between 1-3 dpi in both adult and weanling mice, with levels

significantly higher in weanling mice at 2 dpi (Fig. 1C) (3). This observation underscores

the retention of age-related susceptibility to LACV even at higher infectious doses,

reaffirming the correlation between viremia and survivability.

Neutralizing antibody titers do not correlate with viral clearance in weanling mice

131   nAb responses are pivotal for viral clearance in many infections, yet our findings

reveal an intriguing disparity between nAb titers and viral clearance in weanling mice.

Following infection with a highly infectious dose of LACV, nAbs peak in adult mice

between 8 and 12 dpi (Fig. 2A, 2C), whereas they peak earlier in weanling mice between

4 and 6 dpi (Fig. 2B, 2C), coinciding with the onset of neurological symptoms. Despite the

robust early nAbs responses in weanling mice, 100% succumbed to the infection by 7 dpi

(Fig. 1A). These results suggest that viral clearance and the onset of LACV-induced

neurological signs may not directly correlate with the nAb responses.

Cytokines and chemokines are differentially regulated in adult and weanling mice

To elucidate how serum cytokine and chemokine levels are modulated during

LACV infection in adult and weanling mice, we conducted a Luminex panel capable of

simultaneously detecting and quantifying Th1/Th2/Th9/Th17/Th22/Treg cytokines and

chemokines. Serum samples from adult mice and weanling mice were assessed,

revealing significant shifts in 21 cytokines in LACV-infected mice compared to the un-

infected phase (0 dpi) (Fig. 3). Notably, elevated levels of all cytokines and chemokines

were observed in response to LACV infection in adult and weanling animals, albeit with

varying levels, highlighting the importance of a chronological analysis to fully grasp the

dynamics of LACV-induced cytokine and chemokine production (Fig. 3).

To delineate the potential roles of serum cytokines and chemokines in protection

or pathogenesis at different infection stages, we classified the course of infection into

three phases: (1) peripheral viremia (1-3 dpi), (2) symptoms onset (4-6 dpi), and (3)

survival (>7 dpi). Additionally, we categorized the analyzed cytokines and chemokines

into three groups: Th1/Th2 cytokines, Th9/Th17/Th22/Treg cytokines, and chemokines.

Among the 21 modulated cytokines and chemokines, 9 belonged to the Th1/Th2 cytokine group (GM-CSF, IFN-γ, IL-1β, IL-4, IL-6, IL-12p70, IL-13, IL-18, TNF-α, 6 were classified under the Th9/Th17/Th22/Treg cytokine group (IL-9, IL-10, IL-17A/CTLA-8, IL-22, IL-23, IL-27), and the remaining 6 were identified as elevated chemokines in LACV-infected mice (Eotaxin/CCL11, GROα/CXCL1, IP-10/CXCL10, MCP-1/CCL2, MIP-1α/CCL3, RANTES/CCL5.

Dynamics of Th1/Th2 Cytokines during LACV Infection

IFN-γ, IL-18, TNF-α, and IL-13 emerged as the predominant Th1/Th2 cytokines detected in serum samples from both adult and weanling mice (Fig. 3-4). During the peripheral viremia phase, a significant elevation in IL-18 levels was observed as early as 1 dpi, followed by subsequent significant peaks in IL-12p70 and GM-CSF in adult mice (Fig. 4A, 4F, 4H). Notably, levels of IL-18 and IL-12p70 peak during this early phase before gradually declining as the infection progresses. Conversely, weanling mice display a notable significant increase in levels of IL-6 towards the latter part of the peripheral viremia phase, one day prior to symptoms onset (Fig. 4E).

At the symptoms onset phase, occurring between 4-6 dpi, several cytokines including IFN-γ, IL-1β, TNF-α, GM-CSF, IL-13, and IL-4 were significantly elevated in both adult and weanling mice (Fig. 4A-D, 4G, 4I). Most cytokines reach their peak levels during this phase, with adult mice exhibiting higher levels of IL-1β, TNF-α, GM-CSF, IL-13, and IFN-γ compared to weanlings. Conversely, IL-4 and IL-6 were significantly elevated in weanlings compared to adult mice (Fig. 4D-E). Interestingly, IL-6 levels in adult mice

remained comparable to those during the uninfected phase (Fig. 4E). Finally, during the

survival phase (≤>7 dpi) IFN-γ and IL-13 emerged as the primary cytokines specifically

elevated during this late phase of infection in adult mice.

Dynamics of Th9/Th17/Th22/Treg Cytokines during LACV Infection

A highly infectious dose of LACV revealed elevated levels across all

Th9/Th17/Th22/Treg cytokines analyzed in both adult and weanling (Fig. 5). Notably, IL-

9, IL-10, and IL-17A emerged as the most abundant cytokines within this category

throughout the infection course in both age groups (Fig. 3). During the early peripheral

viremia phase, peaks in IL-9 and IL-17A levels underscored their early involvement in the

immune response to LACV infection in adult and weanling mice (Fig. 5A, 5C). Additionally,

at 2 dpi, IL-23 levels were significantly higher in adult mice compared to weanlings,

mirroring the kinetics of IL-17A, which exhibited significantly elevated levels in adult mice

by 3 dpi (Fig. 5C, 5E).

During the symptoms onset phase, peaks in IL-22, IL-23, and IL-27 levels were

observed, with a notable increase observed in adult mice (Fig. 5D-F). In contrast, IL-10

levels exhibited a significant increase in weanling mice at 4 dpi, followed by a rapid decline

one day later, reaching levels significantly lower than those in adult mice. By 6 dpi,

Th9/Th17/Th22/Treg cytokine levels gradually decreased in adult mice, ultimately

returning to baseline levels.

Dynamics of Chemokines during LACV Infection

A highly infectious dose of LACV also triggers a significant increase in CXCL1 and

CCL11, followed by CXCL10, in both adult and weanling mice compared to uninfected

controls (Fig. 3, 6A-C). This increase manifests as early as 1 dpi for CCL11 and CXCL1.

Particularly, CCL11, CCL3, and CCL5 consistently exhibit higher levels in adult mice

compared to weanlings throughout the infection course (Figure 6A, 6E, 6F). Intriguingly,

CCL3 and CCL5 levels peak during the survival phase in adult mice, suggesting sustained

chemokine responses during later phases of infection.

Conversely, distinct chemokine patterns were observed in weanling mice, with

certain chemokines showing pronounced elevation whiles others exhibit minimal levels.

For example, CCL2 and CXCL10 levels were consistently higher in weanling mice across

all infection phases, whereas CXCL1 levels peak exclusively during the symptoms onset

phase (Fig. 6B, 6D). CCL5 and CCL3 levels in weanlings remain comparable to baseline

levels observed prior to infection (Figure 6E- F).

Discussion

Our study presents a comprehensive analysis of neutralizing antibody titers and

cytokine and chemokine dynamics during a highly infectious dose of LACV infection,

shedding light on their roles in protection and pathogenesis. Our findings demonstrate

distinct patterns of cytokines and chemokines specific to age and infection phase.

Notably, we observed significant increases in Th1/Th2/Th9/Th17/Th22/Treg cytokines

and chemokines in response to LACV infection across both adult and weanling mice.

Interestingly, we observed age-specific differences in many cytokines and chemokines

throughout the course of infection (Fig. 7).

Our data emphasize the consistent influence of age-related susceptibility,

regardless of the infecting dose. Despite prior indications of age-related vulnerability

using lower infectious doses of LACV, our findings expand on this concept by

demonstrating that age remains a pivotal determinant of susceptibility, even when using

highly infectious doses for infections (3). Specifically, we observed that weanling mice

infected with a highly infectious dose exhibited elevated peripheral viral load along with

neutralizing antibodies (nAbs) with an early onset of neurological disease. This

underscores the correlation between peripheral viremia and survivability, highlighting

the relevance of our model for investigating both immune response dynamics and

disease pathology.

Our investigation into viral neutralization yielded unexpected findings. Despite the

early and robust production of nAbs in weanling mice, these titers failed to correlate with

viral clearance, prompting speculation regarding the role of nAbs in early disease

progression, Notably, different neutralization capabilities have been associated with

isotype switching, a phenomenon documented in other viral infections such as HIV and

SARS-CoV-2 (21, 22). Interestingly, adult mice exhibited significantly lower nAb titers

during the symptoms onset phase, suggesting either rapid nAbs consumption in vivo or

the involvement of non-neutralizing antibodies in protection. However, given that weanling

nAbs were incapable of effectively eliminating the virus in vivo, the former explanation

seems less likely.

A possible explanation for these unexpected findings may stem from an imbalance

in the Th1/Th2 response among weanling mice, impacting viral clearance and disease

progression. Upon encountering pathogens, naive CD4$^+$ T cells differentiate into effector

Th1 and Th2 cells, with Th1 cells producing inflammatory cytokines crucial for cell-

mediated immune responses, while Th2 cells facilitate B cell activation and antibody

production (23). An imbalanced response can lead to ineffective immunity and contribute

to pathogenesis. Thus, a Th2-skewed response may account for disparities in the role of

nAbs. Further exploration of immune response dynamics, particularly the balance

between Th1 and Th2 responses, holds promise for elucidating the mechanisms

underlying our findings.

To advance this research direction, it is essential to conduct a thorough

characterization of not only Th1/Th2 cytokines but also Th9/Th17/Th22/Treg cytokines

and chemokines during LACV infection. The investigation of cytokine profiles becomes

particularly crucial in the context of age-dependent viral infections infiltrating the central

nervous system (CNS) (24). This study demonstrated that cytokine and chemokine

profiles in children and adolescents with tick-borne encephalitis vary depending on age

(24). Such variability in cytokine and chemokine responses may significantly influence the

outcome of viral infections and the subsequent risk of severe disease outcomes.

In the case of LACV infection, whether increased cytokines and chemokines are a

correlate of protection or pathogenesis remains to be fully investigated. However, in most

cases, an increase in specific cytokines at particular phases of infection in susceptible

hosts, such as weanling mice, may suggest a potential contribution of these pathways to

susceptibility and disease pathogenesis. Conversely, the increase in cytokines and

chemokines in disease-resistant hosts, such as adult mice, may indicate a protective role

facilitated by these molecules.

Our study reveals distinct patterns of Th1/Th2 cytokines around symptoms onset

during LACV infection in weanling mice. Particularly intriguing is the role of IL-6, a key

factor in determining the balance between Th1 and Th2 responses. Our data suggest that

IL-6, along with IL-4, promotes Th2-related responses during symptoms onset, correlating

with higher nAb levels in weanling mice, supporting the Th2 skey hypothesis (25).

Additionally, the substantial and sustained increase in Th1-related cytokines in adult mice

underscores their protective role against disease, consistent with previous observations

(10, 19).

Moreover, our findings in the context of Th1/Th2 cytokines are consistent with

previous studies on CNS viral infections. For instance, elevated IL-6 levels in the CNS in

infections with dengue and Japanese encephalitis have been associated with various

neurological manifestations and poor outcomes (26, 27). On the contrary, our observation

of the substantial and sustained upregulation of Th1-related cytokines, including IL-18,

IFN-γ, IL-12, and GM-CSF, throughout LACV infection supports previously reported data

of protective roles of GM-CSF and IL-12 in Ifnar1$^{-/-}$ mice and underscores their role in

eliciting protective responses in mice resistant to the disease (19).

Among these cytokines, the pronounced elevation of IL-18 in adult mice stands out

as the most significant modulation induced by LACV. IL-18 has been demonstrated to

stimulate the proliferation and differentiation of T cells toward a Th1 response and acts

as a crucial cofactor in IL-12-induced IFN-γ production by natural killer (NK) and T cells

(28). Moreover, IL-18 exerts its proinflammatory effects by upregulating cytokines such

as IL-13, IL-1β, and TNF-α, consistent with our findings (29). The protective role of IL-18

during viral infections is well-documented, as evidenced by studies showing that

administration of IL-18 induces antiviral activities in mice challenged with a lethal dose of

encephalomyocarditis virus (EMCV) (30). These findings highlight the importance of IL-

18 in mounting effective immune responses against LACV infections and suggest its

potential as a therapeutic target for enhancing antiviral immunity. Furthermore, IFN-γ

emerged as one of the most robust responses, particularly increased in adult mice,

especially during the survival phase. Our findings align with previous observations

indicating increased IFN-γ production from neuron/astrocyte co-cultures infected with

LACV in a dose-dependent manner (16). IFN-γ serves not only as a marker of Th1 CD4[+]

and CD8[+] T cells and NK cells but also as a crucial antiviral mediator central to eliminating

viruses from the CNS (31).

In our analysis of Th9/Th17/Th22/Treg cytokines, all exhibited increased levels in

adult mice in response to LACV infection. Among these, IL-23 and IL-27 were among the

most significantly increased. Although these cytokines have been predominantly studied

in the context of autoimmune diseases, emerging evidence suggests their crucial roles

during viral infections (32, 33). Specifically, IL-23 has been implicated in the control of

viral infections within the brain, indicating its potential importance in host defense

mechanisms against neurotropic viruses such as LACV (32). Similarly, IL-27 has been

found to mediate NK cell effector functions during viral infections, highlighting its

involvement in orchestrating innate immune responses against viral pathogens (33).

Chemokines, as a distinct group of immune mediators, demonstrated elevated

levels in response to LACV infection, with increases observed in CCL5, followed by CCL3,

in adult mice. In contrast, CCL2 increased in weanling mice throughout the course of

infection. The expression and release of CCL2 and CCL3 on cerebral endothelium

suggest an essential role for these chemokines in regulating the trafficking of

inflammatory cells across the blood-brain barrier (BBB) during CNS inflammation (34).

Furthermore, CCL3 has been shown to contribute to the differentiation and migration of

effector T cells in response to viral infection within the CNS (35). Additionally, CCL5 plays

a crucial role in the homing and migration of effector and memory T cells during acute

CNS infections (36). CCL2 emerges as a major player in mediating monocyte infiltration

and the trafficking of inflammatory monocytes to the brain during viral encephalitis (37).

Our findings, indicating increased levels of CCL2 in weanling mice, which by logic,

potentially contributes to pathogenesis possibly mediated through the recruitment of an

overwhelming inflammatory response in the infected brain. However, LACV-induced

lymphocyte infiltration of CNS itself is less likely to contribute to pathogenesis according

to a recent study (9). With this study being one of the few to focus specifically on LACV

infections, significant gaps persist in our understanding of immunity correlates and the

mechanisms underlying immune-mediated viral clearance.   More often than not,

successful clearance of virus infection from the CNS requires the elimination of virus from

the cells without causing damage to neurons. Ideally, immune-mediated virus clearance

from neurons occurs mainly through early cytolytic processes, followed by non-cytolytic

processes that are mediated by antibody specific for the virus and IFN-γ, which enable

neurons to survive (38). Building upon this observation, a better understanding of

adaptive immune responses is needed to address LACV-induced neurological disease.

It is important to acknowledge the limitations of our study. Characterization of

cytokines and chemokines secreted into the serum may not fully capture the immune

activity within the infected host, as the brain-resident immune response may differ from

the systemic immune response. Additionally, we cannot definitively conclude whether a

specific skew in Th1/Th2 responses occurred, as the cellular source of the cytokines

detected in the serum remains unknown. In conclusion, our study contributes to a deeper

understanding of immune responses to neurotropic bunyavirus infections, offering

insights into potential therapeutic targets and biomarkers for disease management.

Future research should focus on elucidating the underlying mechanisms driving age-

related susceptibility, further characterizing the immune response dynamics, and

exploring therapeutic interventions to mitigate the neurological complications associated

with LACV infection. By advancing our understanding of host-virus interactions, we can

develop more effective strategies for the prevention, diagnosis, and treatment of LACV-

induced disease, ultimately enhancing public health preparedness against emerging viral

threats.

Materials and Methods

Infection of mice with LACV and neurological disease progression criteria

All animal studies were approved by the Rutgers University IACUC (protocol

number 202300007). Wildtype (C57BL/6) mice were purchased from Jackson

Laboratories and maintained in a breeding colony at the Child Health Institute of New

Jersey. LACV (La Crosse Virus, NR-540), a human isolate was purchased obtained from

BEI Resources. Mice at 3 (weanling, n=13) or 6-8 (adult, n=13) weeks of age were

inoculated with 5×10^5 plaque forming units (PFU) of LACV in phosphate-buffered saline

(PBS) intraperitoneally in a volume of 200 μl/mouse. Mice were observed daily for signs

of neurological disease that included abnormal gait, seizures, and lethargy. Animals with

clear clinical signs of neurological disease were scored as clinical and euthanized

immediately and humanely.

Serum sample collection

Serum samples were collected from a total of 13 adult mice and 13 weanling mice

throughout the course of infection, including 0 (baseline uninfected phase) and 1, 2, 3, 4,

5, 6, 7, 8 and 12 dpi. Submandibular blood or cardiac blood (on days in which animals

were euthanized) were collected into heparin coated tubes. Blood samples were

centrifuged at 2000 x g for 10 minutes. After centrifugation, sera were aliquoted and

stored in -80°C until subsequent analysis.

Neutralizing antibody assay

A microneutralization assay was performed to detect neutralizing antibodies. 100

μl of the sera were diluted in phosphate buffer saline (PBS), and serial dilutions were

prepared in a 96-well plate. For each dilution, contact was performed with a 100 PFU

LACV dose for two hours at 37°C in a 5% CO_2 incubator. The complexes were then added

to Vero cells (ATCC) CCL-81, 4×10^4 cells/well. After 3-4 days of incubation, the

cytopathogenic effects were investigated with 0.1% crystal violet staining (SigmaAldrich,

St. Louis, MO) for 2-4 hours at room temperature prior to washing with water. The

neutralizing titer was defined as the inverse of the highest dilution resulting in a infectious

reduction of 50%. A titer lower than 1:20 was considered as negative if cytopathic effects

were not observed. Samples were considered as positive if the titer was >1:20.

qPCR

Viral titers were quantified by qPCR. Total viral RNA was extracted from pooled

serum samples from LACV-infected adult and weanling mice using Pure link viral RNA

mini kit (ThermoFisher, Waltham, MA) following manufacturer's protocol. Given that

submandibular blood collection results in small quantities of serum isolation necessitated

pooling of sera samples prior to viral RNA extraction. cDNA was prepared from RNA

samples using the SuperScript IV VILO reverse transcription kit (ThermoFisher, Waltham,

MA) following the manufacturer's instructions. cDNA samples were diluted fivefold in

Rnase-free water and those diluted samples were analyzed by qPCR using PowerTrack

SYBR Green kit (ThermoFisher, Waltham, MA) to detect LACV RNA. The following

primers were used: forward (ATTCTACCCGCTGACCATTG) and reverse

(GTGAGAGTGCCATAGCGTTG). CT values were measured using a QuantStudio 3Flex

instrument (Applied Biosysterms, Waltham, MA). QuantStudio Real-Time PCR software

was used to extract the data.

Bead-based multiplex immunoassay (Luminex)

Sera samples from adult male (n=3) and female (n=3) and weanling male (n=3) and female(n=3) mice at 0, 1, 3, 5, and 7 dpi were pooled for analysis. Additionally, sera samples from adult (n=7; 4 male, 3 female) and weanling (n=7; 4 male, 3 female) mice at 2, 4, 6, 8, and 12 dpi were individually analyzed. To quantify the levels of serum cytokines and chemokines in LACV-infected adult and weanling mice, a magnetic bead-based multiplex immunoassay was performed using a MAGPIX® (ThermoFisher, Waltham, MA) instrument. A mouse cytokine magnetic 26-plex panel (Invitrogen) was utilized to determine immune responses mediated through cytokine and chemokine such as (1) Th1/Th2: GM-CSF, IFN-γ, IL-1β , IL-2 , IL-4 , IL-5, IL-6, IL-12p70, IL-13, IL-18, TNF-α, and (2) Th9/Th17/Th22/Treg: IL-9, IL-10, IL-17A/CTLA-8, IL-22, IL-23, IL-27, (3) Chemokines: Eotaxiv/CCL11, GROα/CXCL1, IP-10/CXCL10, MCP-1/CCL2, MCP-3/CCL7, MIP-1α/CCL3, MIP-1β/CCL4, MIP-2α/CXCL2, RANTES/CCL5, MCP-3/CCL7 following the manufacturer's instructions. 5 (IL-2, IL-5, MIP-1β/CCL4, MIP-2α/CXCL2, and MCP-3/CCL7) of the analyzed cytokines were out of range and therefore were not included in the analysis for both adult and weanling mice.

Statistical analysis

All statistical analyses were performed using Prism software Version 7.01 (GraphPad) and are described in the figure legends.

Data Availability

The data that support the findings of this study are available within the manuscript or upon request from the corresponding author.

Acknowledgements

We would like to thank Rutgers Global Health Institute, Rutgers Robert Wood Johnson Medical School, and the Child Health Institute of New Jersey for their support.

Conflicts of Interest

BBH is a co-founder of Mir Biosciences, Inc., a biotechnology company focused on T cell-based diagnostics and vaccines for infectious diseases, cancer, and autoimmunity.

References

1. Alatrash R, Herrera BB. 2024. The Adaptive Immune Response against Bunyavirales. Viruses 16:483.

2. Westby KM, Fritzen C, Paulsen D, Poindexter S, Moncayo AC. 2015. La Crosse Encephalitis Virus Infection in Field-Collected Aedes albopictus, Aedes japonicus, and Aedes triseriatus in Tennessee. J Am Mosq Control Assoc 31:233-41.

3.   Bennett RS, Cress CM, Ward JM, Firestone CY, Murphy BR, Whitehead SS. 2008. La

Crosse virus infectivity, pathogenesis, and immunogenicity in mice and monkeys. Virol J

5:25.

4.   Schneider CA, Leung JM, Valenzuela-Leon PC, Golviznina NA, Toso EA, Bosnakovski

D, Kyba M, Calvo E, Peterson KE. 2024. Skin muscle is the initial site of viral replication

for arboviral bunyavirus infection. Nat Commun 15:1121.

5.   Haddow AD, Odoi A. 2009. The incidence risk, clustering, and clinical presentation of La

Crosse virus infections in the eastern United States, 2003-2007. PLoS One 4:e6145.

6.   Balfour HH, Jr., Siem RA, Bauer H, Quie PG. 1973. California arbovirus (La Crosse)

infections. I. Clinical and laboratory findings in 66 children with meningoencephalitis.

Pediatrics 52:680-91.

7.   Hartman AL, Myler PJ. 2023. Bunyavirales: Scientific Gaps and Prototype Pathogens for

a Large and Diverse Group of Zoonotic Viruses. The Journal of Infectious Diseases

228:S376-S389.

8.   López K, Wilson SN, Coutermash-Ott S, Tanelus M, Stone WB, Porier DL, Auguste DI,

Muller JA, Allicock OM, Paulson SL, Erasmus JH, Auguste AJ. 2021. Novel murine

models for studying Cache Valley virus pathogenesis and in utero transmission. Emerg

Microbes Infect 10:1649-1659.

9.   Winkler CW, Myers LM, Woods TA, Carmody AB, Taylor KG, Peterson KE. 2017.

Lymphocytes have a role in protection, but not in pathogenesis, during La Crosse Virus

infection in mice. J Neuroinflammation 14:62.

10.   Taylor KG, Woods TA, Winkler CW, Carmody AB, Peterson KE. 2014. Age-dependent

myeloid dendritic cell responses mediate resistance to la crosse virus-induced

neurological disease. J Virol 88:11070-9.

11.   Hefti HP, Frese M, Landis H, Di Paolo C, Aguzzi A, Haller O, Pavlovic J. 1999. Human

MxA protein protects mice lacking a functional alpha/beta interferon system against La

crosse virus and other lethal viral infections. J Virol 73:6984-91.

12.   Deng B, Zhang S, Geng Y, Zhang Y, Wang Y, Yao W, Wen Y, Cui W, Zhou Y, Gu Q,

Wang W, Wang Y, Shao Z, Wang Y, Li C, Wang D, Zhao Y, Liu P. 2012. Cytokine and

Chemokine Levels in Patients with Severe Fever with Thrombocytopenia Syndrome

Virus. PLOS ONE 7:e41365.

13.   McElroy AK, Nichol ST. 2012. Rift Valley fever virus inhibits a pro-inflammatory response

in experimentally infected human monocyte derived macrophages and a pro-

inflammatory cytokine response may be associated with patient survival during natural

infection. Virology 422:6-12.

14.   Borges AA, Campos GM, Moreli ML, Moro Souza RL, Saggioro FP, Figueiredo GG,

Livonesi MC, Moraes Figueiredo LT. 2008. Role of mixed Th1 and Th2 serum cytokines

on pathogenesis and prognosis of hantavirus pulmonary syndrome. Microbes Infect

10:1150-7.

15.   Salinas TP, Garrido JL, Salazar JR, Gonzalez P, Zambrano N, Fuentes-Villalobos F,

Bravo F, Fica-Leon V, Salas-Burgos A, Calvo M, Alvarez R, Armien B, Barria MI. 2021.

Cytokine Profiles and Antibody Response Associated to Choclo Orthohantavirus

Infection. Front Immunol 12:603228.

16.   Dawes BE, Gao J, Atkins C, Nelson JT, Johnson K, Wu P, Freiberg AN. 2018. Human

neural stem cell-derived neuron/astrocyte co-cultures respond to La Crosse virus

infection with proinflammatory cytokines and chemokines. Journal of Neuroinflammation

15:315.

17.   Kallfass C, Ackerman A, Lienenklaus S, Weiss S, Heimrich B, Staeheli P. 2012.

Visualizing production of beta interferon by astrocytes and microglia in brain of La

Crosse virus-infected mice. J Virol 86:11223-30.

18.   Taylor KG, Peterson KE. 2014. Innate immune response to La Crosse virus infection.

491   Journal of NeuroVirology 20:150-156.

492   19.   Operschall E, Schuh T, Heinzerling L, Pavlovic J, Moelling K. 1999. Enhanced protection

493   against viral infection by co-administration of plasmid DNA coding for viral antigen and

494   cytokines in mice. J Clin Virol 13:17-27.

495   20.   Proenca-Modena JL, Hyde JL, Sesti-Costa R, Lucas T, Pinto AK, Richner JM, Gorman

496   MJ, Lazear HM, Diamond MS. 2016. Interferon-Regulatory Factor 5-Dependent

497   Signaling Restricts Orthobunyavirus Dissemination to the Central Nervous System. J

498   Virol 90:189-205.

499   21.   Noval MG, Kaczmarek ME, Koide A, Rodriguez-Rodriguez BA, Louie P, Tada T, Hattori T,

500   Panchenko T, Romero LA, Teng KW, Bazley A, de Vries M, Samanovic MI, Weiser JN,

501   Aifantis I, Cangiarella J, Mulligan MJ, Desvignes L, Dittmann M, Landau NR, Aguero-

502   Rosenfeld M, Koide S, Stapleford KA. 2021. Antibody isotype diversity against SARS-

503   CoV-2 is associated with differential serum neutralization capacities. Sci Rep 11:5538.

504   22.   Scheepers C, Bekker V, Anthony C, Richardson SI, Oosthuysen B, Moyo T, Kgagudi P,

505   Kitchin D, Nonyane M, York T, Mielke D, Mabvakure BM, Sheng Z, Lambson BE, Ismail

506   A, Garrett NJ, Abdool Karim SS, Shapiro L, Williamson C, Morris L, Moore PL. 2020.

507   Antibody Isotype Switching as a Mechanism to Counter HIV Neutralization Escape. Cell

508   Reports 33.

509   23.   Paul WE, Seder RA. 1994. Lymphocyte responses and cytokines. Cell 76:241-251.

510   24.   Ygberg S, Fowler Å, Wickström R. 2022. Age-related changes in the inflammatory

511   responses to viral infections in the central nervous system during childhood. Pediatr Res

512   91:204-208.

513   25.   Rincón M, Anguita J, Nakamura T, Fikrig E, Flavell RA. 1997. Interleukin (IL)-6 directs

514   the differentiation of IL-4-producing CD4+ T cells. J Exp Med 185:461-9.

515   26.   Mehta VK, Verma R, Garg RK, Malhotra HS, Sharma PK, Jain A. 2017. Study of

516   interleukin-6 and interleukin-8 levels in patients with neurological manifestations of

517   dengue. J Postgrad Med 63:11-15.

518   27.   Winter PM, Dung NM, Loan HT, Kneen R, Wills B, Thu LT, House D, White NJ, Farrar JJ,

519   Hart CA, Solomon T. 2004. Proinflammatory Cytokines and Chemokines in Humans with

520   Japanese Encephalitis. The Journal of Infectious Diseases 190:1618-1626.

521   28.   Micallef MJ, Ohtsuki T, Kohno K, Tanabe F, Ushio S, Namba M, Tanimoto T, Torigoe K,

522   Fujii M, Ikeda M, Fukuda S, Kurimoto M. 1996. Interferon-gamma-inducing factor

523   enhances T helper 1 cytokine production by stimulated human T cells: synergism with

524   interleukin-12 for interferon-gamma production. Eur J Immunol 26:1647-51.

525   29.   Puren AJ, Fantuzzi G, Gu Y, Su MS, Dinarello CA. 1998. Interleukin-18 (IFNgamma-

526   inducing factor) induces IL-8 and IL-1beta via TNFalpha production from non-CD14+

527   human blood mononuclear cells. J Clin Invest 101:711-21.

528   30.   Tovey MG, Meritet JF, Guymarho J, Maury C. 1999. Mucosal cytokine therapy: marked

529   antiviral and antitumor activity. J Interferon Cytokine Res 19:911-21.

530   31.   Chesler DA, Reiss CS. 2002. The role of IFN-γ in immune responses to viral infections

531   of the central nervous system. Cytokine & Growth Factor Reviews 13:441-454.

532   32.   Yannam GR, Gutti T, Poluektova LY. 2012. IL-23 in infections, inflammation,

533   autoimmunity and cancer: possible role in HIV-1 and AIDS. J Neuroimmune Pharmacol

534   7:95-112.

535   33.   Amsden H, Kourko O, Roth M, Gee K. 2022. Antiviral Activities of Interleukin-27: A

536   Partner for Interferons? Front Immunol 13:902853.

537   34.   Chui R, Dorovini-Zis K. 2010. Regulation of CCL2 and CCL3 expression in human brain

538   endothelial cells by cytokines and lipopolysaccharide. Journal of Neuroinflammation 7:1.

539   35.   Trifilo MJ, Bergmann CC, Kuziel WA, Lane TE. 2003. CC chemokine ligand 3 (CCL3)

540   regulates CD8(+)-T-cell effector function and migration following viral infection. J Virol

541   77:4004-14.

542   36.   Glass WG, Lim JK, Cholera R, Pletnev AG, Gao J-L, Murphy PM. 2005. Chemokine

543   receptor CCR5 promotes leukocyte trafficking to the brain and survival in West Nile virus

544   infection. Journal of Experimental Medicine 202:1087-1098.

545   37.   Chen BP, Kuziel WA, Lane TE. 2001. Lack of CCR2 results in increased mortality and

546   impaired leukocyte activation and trafficking following infection of the central nervous

547   system with a neurotropic coronavirus. J Immunol 167:4585-92.

548   38.   Griffin DE. 2003. Immune responses to RNA-virus infections of the CNS. Nature

549   Reviews Immunology 3:493-502.

Figures

Figure 1. Impact of infectious dose on viremia and susceptibility to LACV infection. (A)

Percent survival of adult (black) and weanling (gray) LACV-infected mice with highly infectious

dose (P<0.001). **(B)** Percent mice showing LACV-induced outcomes in adult (top) and weanling

(bottom) mice. **(C)** Viral titer in sera of LACV-infected adult and weanling mice represented by

LACV RNA copies quantified by qPCR. *, p<0.05.

574 **Figure 2. Neutralizing antibody titers do not correlate with viral clearance in weanling mice.**

575 **(A, B)** NT50 titers in sera from LACV-infected adult (black) and weanling (gray) mice throughout

576 the course of infection. **(C)** Mean NT50 titers between adult and weanling mice. *, $p<0.05$.

Figure 3. Cytokines and chemokines are differentially regulated in adult and weanling mice. A heatmap for cytokine and chemokine levels analyzed in **(A)** adult and **(B)** weanling mice during the course of LACV infection (1-12 dpi) compared to uninfected phase (0 dpi). The rows represent cytokines and chemokines and the columns represent 1-12 dpi. The scale is represented in an arbitrary format, depicting low (0) to high (3) cytokines and/or chemokines detected in sera.

Figure 4. Dynamics of Th1/Th2 Cytokines During LACV Infection. (A-I) Bar charts representing levels of Th1/Th2 related cytokines in sera of adult (black) and weanling (gray) mice during LACV infection. *, p<0.05.

Figure 5. Dynamics of Th9/Th17/Th22/Treg Cytokines During LACV Infection. (A-F) Bar

charts representing levels of Th9/Th17/Th22/Treg related cytokines in sera of adult (black) and

weanling (gray) mice during LACV infection. Data represent average from 9 adult and 9 weanling

mice with SD. *, p<0.05.

634 **Figure 6. Dynamics of Chemokine Levels During LACV Infection. (A-F)** Bar charts

635 representing levels of chemokines in sera of adult (black) and weanling (gray) mice during LACV

636 infection. *, $p < 0.05$.

Cytokines modulated during LACV infection

654 **Figure 7. Summary of serum cytokine and chemokine responses during phases of LACV**

655 **infection.** Roadmap illustrating the relative changes in cytokine and chemokine levels across

656 different phases of LACV infection, categorized as peripheral viremia, symptom onset, and

657 survival stage. Blue arrows indicate trends observed in adult mice, while green arrows represent

658 trends observed in weanling mice.