Convergent evolution associated with the loss of developmental diapause may promote

extended lifespan in bees

Priscila K. F. Santos[1*], Karen M. Kapheim[1]

[1]Utah State University, Logan, UT, 84322, United States of America

*Author for Correspondence: Priscila K. F. Santos, Biology Department, Utah State University,

Logan, UT, USA, pkfsantos1@gmail.com

Abstract

Diapause has long been proposed to play a significant role in the evolution of eusociality in

Hymenoptera. Recent studies have shown that shifts in the diapause stage precede social

evolution in wasps and bees, however, the genomic basis remains unknown. Given the overlap

in molecular pathways that regulate diapause and lifespan, we hypothesized that the

evolutionary loss of developmental diapause may lead to extended lifespan among adults,

which is a prerequisite for the evolution of eusociality. To test this, we compared 27 bee

genomes with or without prepupal diapause. Our results point to several potential mechanisms

for lifespan extension in species lacking prepupal diapause, including the loss of the growth

hormone PTTH and its receptor TORSO, along with convergent selection in genes known to

regulates lifespan in animals. Specifically, we observed purifying selection of pro-longevity

genes and relaxed selection of anti-longevity genes within the IIS/TOR pathway in species that

have lost prepupal diapause. Changes in selection pressures on this pathway may lead to the

evolution of new phenotypes, such as lifespan extension and altered responses to nutritional

signals, that are crucial for social evolution.

Keywords:

Eusociality, comparative genomics, selection, dormancy, aging

Significance

Diapause has long been proposed to play a significant role in the evolution of eusociality in Hymenoptera. Recent studies have shown that the loss of diapause during the prepupal stage precedes social evolution in wasps and bees. However, the genomic mechanisms underlying this phenomenon remain unknown. Through comparative genomics, we showed that the convergent loss of prepupal diapause is associated with mechanisms that may promote lifespan extension, a prerequisite for social evolution. These mechanisms include genes losses and signals of selection on genes related to aging.

Introduction

The origin of eusociality is one of the major transitions in evolution, because it involves a shift from reproduction at the level of the individual to reproduction at the colony level (Szathmáry & Smith 1995). A typical eusocial colony of ants, bees, or wasps includes a long-lived queen who specializes on egg-laying living alongside her non-reproductive daughters who perform nest maintenance, foraging, and brood care (Wilson 1971). Understanding the selective pressures shaping this major transition requires investigating the origins of each of its defining features; (1) overlapping generations, (2) reproductive division of labor, and (3) cooperative brood care (Batra 1966; Michener 1969). For example, one of the preconditions for overlapping generations is that nest foundresses (queens) must live long enough to overlap with their adult daughters (workers). This suggests that the origins of eusociality were preceded by an extension of lifespan in the solitary ancestors (Michener 1974; Carey 2001; da Silva 2022).

A potential source of this lifespan extension could be rooted in the phenomenon of diapause.

Diapause is a period of dormancy in insects, in which development and reproduction are

suppressed, usually as a way to pass unfavorable periods such as extreme temperatures,

drought, or scarcity of resources (Denlinger 2022). In bees, diapause most commonly occurs in

larvae following the last molt, a stage referred to as the prepupa (Gauld & Bolton 1988; P.K.F.

Santos et al. 2019). However, there is a great deal of variation between bee species, with many

bees diapausing as adults, and many skipping diapause altogether (P.K.F. Santos et al. 2019).

Recent research indicates that prepupal diapause is the ancestral state in bees, but it has been

lost in the ancestor of every clade in which eusociality has evolved (P.K.F. Santos et al. 2019).

This pattern has been found in both bees and wasps (da Silva 2021), suggesting that while

diapause itself may not be essential for social evolution, its loss during the prepupal stage could

be pivotal if it leads to increased longevity, thus facilitating overlapping generations as required

for the evolution of eusociality.

A recent study of lifespan evolution in Hymenoptera gleaned longevity estimates from the

literature to show that primitively eusocial nest foundresses do indeed live significantly longer

than non-eusocial Hymenoptera foundresses (da Silva 2022). Although diapause was not

included in the final model of foundress lifespan (likely due to missing data in more than a

quarter of the species), foundresses from species included in the study that have lost prepupal

diapause had median lifespans 3.5 times as long as those with prepupal diapause (median = 87

d vs 25 d) (da Silva 2021). Here we attempt to explain these connections between lifespan,

eusociality, and the loss of prepupal diapause by identifying molecular pathways under

selection with the loss of prepupal diapause in bees.

Many of the molecular pathways that regulate diapause in insects also regulate aging and

longevity (Hutfilz 2022). Although diapause at different life stages shares common features –

such as reduced metabolism, increased lipid reserves, resistance to extreme environmental

conditions, and prolonged lifespan – its physiological regulation varies. In prepupal diapause,

development is halted due to a reduction in the production or release of the prothoracicotropic

hormone (PTTH) by the prothoracic gland (PG), leading to decreased ecdysone levels, which

delay metamorphosis (Denlinger 2022b; Rewitz et al. 2009). Conversely, adult diapause involves

the cessation of reproduction through the arrest of oocyte development, primarily regulated by

a shutdown in juvenile hormone (JH) production (Denlinger 2022b; C.G. Santos et al. 2019).

Despite these differences, certain pathways regulate diapause across life stages. The

insulin/insulin-like signaling and target of rapamycin (IIS/TOR) pathway is one of these

pathways, known for its role in regulating both diapause and aging. The pathway's specific

regulatory functions depend on the life stage, indicating that the same pathway may

experience different selective pressures depending on the stage: during developmental

diapause, the IIS/TOR pathway manages nutrient allocation for growth and metamorphosis,

while during adult diapause, it allocates nutrients for maintenance and reproduction (Denlinger

2022b). This suggests that shifts in the diapause stage could lead to changes in pathways

regulating aging, potentially facilitating lifespan extension.

One other important difference between prepupal and adult diapause is the opportunity for

cellular regeneration. Diapause during development is followed by the renewal of somatic

tissues through cell divisions and the process of metamorphosis itself (Tatar & Yin 2001).

Metamorphosis allows for a restructuring of the organism's body, which can rejuvenate tissues,

with potentially lifespan extending effects (Gilbert 2000; Seifert et al. 2012). However, in adults,

most of the tissues are post-mitotic, meaning they cannot undergo cell division to renew

themselves, limiting the extent to which lifespan can be extended by diapause alone. Adult

insects that go through diapause mitigate its potential negative effects by reducing age-specific

mortality through a reduced rate of senescence, an adaptation specific to the adult stage

(Tatar & Yin 2001; Easwaran & Montell 2023). The loss of prepupal diapause associated with

adaptations to reduce senescence rate may be key in promoting the overlap of generations and

the evolution of social behavior in bees.

If shifts in diapause stage result in prolonged lifespan, then the convergent loss of

developmental diapause in bees could be linked to genomic changes associated with

mechanisms that regulate adult lifespan extension. To test this hypothesis, we compared the

genomes of 27 bee species with and without prepupal diapause. We looked for signals of

positive, purifying, and relaxation of selection, as well as gene family gain and losses linked to

the absence of prepupal diapause. Next, we assessed whether genes under selection were

enriched for longevity or oxidative stress-related functions. We found that the loss of prepupal

diapause might be associated with mechanisms regulating adult lifespan extension through

processes including the loss of the growth prothoracicotropic hormone (PTTH) and its tyrosine-

protein kinase receptor torso (TORSO), purifying selection of pro-longevity genes, and relaxed

selection of anti-longevity genes in the IIS/TOR pathway.

Results

Genomic data selection and orthogroups identification

Out of the 74 bee genomes with available annotation files and information regarding the

diapause stage, 71 genomes exhibited over 80% complete BUSCO genes within their protein-

coding sequences. Following the exclusion of species from the same genus sharing identical life-

history traits, we retained 27 species for subsequent selection analysis (Figure 1; Table S1).

Orthofinder analysis yielded 2592 single-copy orthogroups across all 27 species. After filtering

to retain orthogroups present in all species, but being single-copy in at least 70% of them, we

identified and utilized 4173 orthogroups for further analysis.

Figure 1: The 27 species included in the molecular evolution analysis and their associated life-history traits. The prepupal diapause stage is depicted in dark blue, while the loss of developmental diapause is represented in lighter blue. Solitary bees are indicated by a stronger yellow color, and social species are denoted by a lighter yellow. Purple represents the presence of the prothoracicotropic hormone (PTTH) and its tyrosine-protein kinase receptor torso (TORSO), while the absence of color indicates orthologs could not be found. The species marked with "?" had conflicting results regarding the presence of PTTH, depending on whether the search was conducted using the genome or the annotated protein-coding genes set as a reference.

157 **Predominance of relaxation of selection linked to the loss of prepupal diapause**

158 The search for convergent evolutionary rate shifts linked to the loss of prepupal diapause using

159 RERconverge resulted in 914 significant orthogroups (Benjamini-Hochberg adjusted p-value <

160 0.05 after permulation). Among these orthogroups, 336 exhibited positive Rho values,

161 signifying genes undergoing accelerated evolution (attributed to the release of selective

162 constraint or adaptation), whereas 578 showed negative Rho values indicating a decelerated

163 evolutionary rate (increased selective constraint) in species without diapause during the

164 prepupal stage (Table S2).

165

166 aBSREL identified 98 orthogroups where positive selection occurred on a proportion of

167 branches in species lacking diapause in the prepupal stage. However, the significance in most

168 groups is associated with one species (at most three species), suggesting a weak convergent

169 signal of positive selection related to the loss of prepupal diapause (Table S3). Using Hyphy-

170 RELAX, 367 orthogroups were identified under intensification of selection (potentially positive

171 or purifying selection) and 1486 orthogroups under relaxation of selection (Table S3).

172

173 Combining results from RERconverge and Hyphy, we categorized orthogroups into groups of

174 interest (see Material and Methods for details): possible positive selection – 293 orthogroups;

175 possible purifying selection – 347 orthogroups; possible relaxed selection – 1217 orthogroups

176 (Figure 2A); and possible intensified selection – 260 orthogroups (Figure 2B). We also classified

177 these orthogroups as having the best evidence of positive selection (66 orthogroups) – those

178 that were significant in at least two of the following: aBSREL, RERconverge with Rho positive

values, and intensification of selection in RELAX (Figure 2B and Table S4); or having the best

evidence of purifying selection (45 orthogroups) – orthogroups significant in both RERconverge

with Rho negative values and intensification of selection from RELAX (Figure 2B; Table S5).

Figure 2: The Venn diagram depicts orthogroups categorized under relaxed (A) or intensified (B)

selection, which intersect with genes under positive selection (aBSREL), accelerated evolution

(Rho +), and/or decelerated evolution (Rho -). Different colors denote orthogroups potentially

experiencing positive (yellow), purifying (blue), relaxed (purple), or intensified (green) selection,

while dashed lines indicate orthogroups with the strongest evidence for positive (yellow) and

purifying (blue) selection. Refer to the Material and Methods section for comprehensive

classification details.

Gene ontology enrichment

Orthogroups exhibiting possible positive and intensified selection were significantly enriched

for terms associated with post-translational protein modifications, such as protein

dephosphorylation and ubiquitin protein ligase activity, respectively. Regulation of transcription

was also among the enriched terms for both positive and purifying selection (Table S6). There

were no significant terms among the orthogroups under relaxed selection at p < 0.01, with

double-strand break repair and RNA methylation being the top terms (p = 0.031) (Table S6).

Prepupal diapause loss is a strong predictor of *PTTH* and *torso* loss

An analysis of gene family gain and loss revealed that only two orthogroups were lost in the

majority of species that do not undergo diapause during development (Table S7). Specifically,

the prothoracicotropic hormone (*PTTH*) gene and its tyrosine-protein kinase receptor torso

(*torso*) were absent in all species lacking prepupae diapause, except for *Andrena dorsata*, which

has both orthogroups and two copies of the *torso* receptor (Figure 1, Table S7). Our results

show these losses are followed by positive selection in genes of the EGFR signaling pathway,

which is the main pathway producing ecdysone during metamorphosis (Cruz et al. 2020) (Table

S4). Notably, *epidermal growth factor receptor substrate 15* (*egfr15*) and *son of sevenless* (*SOS*)

genes show strong evidence of positive selection (Table S4 and Figure 3).

Figure 3: Simplified scheme illustrating two pathways regulating the production and release of

ecdysone by the prothoracic gland (PG). The pathway utilizing the tyrosine-protein kinase

receptor TORSO as a receptor for the prothoracicotropic hormone PTTH is absent in species

that do not undergo prepupal diapause. *epidermal growth factor receptor substrate 15*

(EGFR15), a substrate of the alternative pathway that uses EGFR as a receptor, is subject to

positive selection, along with the *son of sevenless* (*SOS*) gene, highlighting the significance of

this pathway in species without prepupal diapause.

PTTH and *torso* loss have been previously associated with social behavior in bees. In a previous

screening of 22 bee genomes, all 15 of the included social species were missing both *PTTH* and

torso (Costa et al. 2022). We aimed to discern whether the diapause stage or sociality exerted a

stronger influence on gene loss. To address this, we performed PGLS analyses, considering both

predictor variables independently and simultaneously (Table S8). When analyzed individually,

both diapause stage (Estimate: 0.81, Multiple R-squared: 0.87, p = 2e-13) and sociality

(Estimate: 0.48, Multiple R-squared: 0.44, p=8e-05) significantly predicted the

presence/absence of *PTTH* and *torso*. However, in the joint analysis, diapause stage is the only

significant predictor (Estimates: Diapause(yes) – 0.81, Sociality(solitary) – 0.10; p-values:

Diapause – 3e-10, Sociality – 0.1186). Furthermore, the model with both predictors had a

slightly higher Multiple R-squared = 0.88 (vs 0.87 including only diapause as predictor),

indicating diapause stage is a stronger predictor of *PTTH* and *torso* loss.

Orthogroups under selection are not enriched for oxidative stress and lifespan genes

Out of 198 genes related to aging from the GenAge database (Tacutu et al. 2018), we identified

101 among our 4172 orthogroups. We found aging genes among all genes we identified as

under possible positive, purifying, intensified, and relaxed selection associated with the loss of

prepupal diapause, with the orthogroups under relaxation of selection including the highest

number of aging genes (26 out of a total of 1217 genes). However, none of the groups exhibited

more aging genes than expected by chance (Hypergeometric test: p > 0.2). Many of the aging

genes under selection are part of the IIS/TOR pathway (Figure 4). One gene previously

described as pro-longevity, and at least five genes described as anti-longevity in *Drosophila*

(Tacutu et al. 2018) were identified as under purified and relaxed selection in the current study,

respectively (Figure 4 and Table 1).

Figure 4: Simplified scheme of the insulin/insulin-like signaling (IIS) and target of rapamycin

(TOR) pathways containing genes under selection in the current study. When the insulin

pathway is activated (high nutrition), Akt phosphorylates FoxO, preventing its entrance to the

nucleus of the cell, and activates Tor, leading to growth. When there is a reduction in the

IIS/TOR pathway activation (low-nutrition), FoxO enters in the nucleus of the cells, regulating

thousands of genes, leading to lifespan extension. Continuous arrows represent activation of

the pathway in an environment with high nutrition, for example, while dashed arrows

represent the alternative pathway in environments with low nutrition. Genes within green,

blue, or purple boxes represent those under intensified, purifying, and relaxed selection in the

current study, respectively. The box in the bottom right classifies the genes as pro- or anti-

longevity based on their known effects on *Drosophila* lifespan (Tacutu et al. 2018). Figure made

based on Kapahi and Zid 2004; Dutriaux et al. 2013; Texada et al. 2020; Denlinger 2022.

259 **Table 1:** Orthogroups under selection overlapping with the *Drosophila* aging genes list from

260 GenAge database (Tacutu et al. 2018). Longevity influence classifies the orthologs as pro- or

261 anti-longevity genes according to their effects on *Drosophila* lifespan. The asterisk (*) highlights

262 orthogroups that are also part of the best evidence of positive selection (Table 1).

Flybase ID	Orthogroup	Gene Symbol	Gene name	Longevity Influence	Selection
FBgn0014141	OG0000645	*cher*	cheerio	pro	intensification
FBgn0024248	OG0003925	*chico*	Insulin receptor substrate-1	anti	intensification
FBgn0000659	OG0002632	*fkh*	fork head	pro	intensification
FBgn0002566	OG0004524	*lt*	light	pro	intensification
FBgn0015269	OG0005277	*Nf1*	Neurofibromin 1	pro	intensification
FBgn0037802	OG0003529	*Sirt6*	CG6284-PB	pro	intensification
FBgn0000330	OG0005760	*cm*	carmine	pro	positive*
FBgn0037574	OG0005827	*Coq2*	Coenzyme Q biosynthesis protein 2	anti	positive
FBgn0001087	OG0003271	*g*	garnet	pro	positive
FBgn0260990	OG0005419	*yata*	CG1973 gene product from transcript CG1973-RA	pro	positive*
FBgn0031461	OG0004172	*daw*	dawdle	anti	purifying
FBgn0038197	OG0003188	*foxo*	Forkhead box, sub-group O	pro	purifying
FBgn0039044	OG0002793	*p53*	CG33336-PA	anti	purifying
FBgn0261549	OG0000874	*rdgA*	retinal degeneration A	anti	purifying
FBgn0044452	OG0005320	*Atg2*	Autophagy-specific gene 2	pro	relaxation
FBgn0034366	OG0003358	*Atg7*	Autophagy-specific gene 7	pro	relaxation
FBgn0030512	OG0004420	*CG9940*	CG9940	pro	relaxation
FBgn0262656	OG0001307	*dm*	diminutive	anti	relaxation
FBgn0000490	OG0004605	*dpp*	decapentaplegic	pro	relaxation
FBgn0026479	OG0002567	*Drp1*	Dynamin related protein 1	pro	relaxation
FBgn0004507	OG0005350	*GlyP*	Glycogen phosphorylase	anti	relaxation
FBgn0038074	OG0004077	*Gnmt*	Glycine N-methyltransferase	pro	relaxation
FBgn0028427	OG0002615	*Ilk*	Integrin linked kinase	anti	relaxation
FBgn0283499	OG0005890	*InR*	Insulin-like 1 receptor	anti	
FBgn0283499	OG0001807	*InR*	Insulin receptor-like	anti	relaxation
FBgn0011455	OG0002494	*l(3)neo18*	l(3)neo18	anti	relaxation
FBgn0002719	OG0002827	*Men*	Malic enzyme	pro	relaxation
FBgn0014863	OG0003311	*Mlp84B*	Muscle LIM protein at 84B	pro	relaxation
FBgn0023000	OG0004888	*mth*	methuselah	anti/pro	relaxation
FBgn0051216	OG0003476	*Naam*	Naam gene product from transcript Naam-RA	pro	relaxation
FBgn0017566	OG0002132	*ND75*	NADH:ubiquinone reductase 75kD subunit precursor	anti	relaxation
FBgn0030795	OG0001867	*ppk28*	pickpocket 28	anti	relaxation
FBgn0283472	OG0004887	*S6k*	RPS6-p70-protein kinase	anti	relaxation
FBgn0025391	OG0004306	*Scgdelta*	Sarcoglycan delta	pro	relaxation
FBgn0037248	OG0001135	*Spargel*	Spargel	pro	relaxation
FBgn0040268	OG0001708	*Top3alpha*	Topoisomerase 3alpha	pro	relaxation
FBgn0021796	OG0004069	*Tor*	Target of rapamycin	anti	relaxation
FBgn0004397	OG0005595	*Vinc*	Vinculin	pro	relaxation

264 Regarding oxidative stress genes (Kramer et al. 2021), 17 out of 34 had orthologs among the

265 bee orthogroups. Only the positive selection and the relaxation of selection groups included

266 oxidative stress orthologs in their lists, although not exceeding the amount of overlap expected

267 by chance (Hypergeometric test: p > 0.7).

Discussion

In this study, we tested the hypothesis that the evolutionary loss of prepupal diapause is

followed by genomic changes that might be associated with adult lifespan extension, a

prerequisite for the evolution of sociality in bees (Batra 1966; Carey 2001; Michener 1974). Our

findings revealed convergent signals of selection linked to the loss of prepupal diapause, with

the majority of genes experiencing relaxed selection. However, there is no significant overlap

between genes under selection and genes previously associated with aging or oxidative stress

in other groups. Below we discuss specific pathways that may be relevant to the adult lifespan

extension in the absence of prepupal diapause. These include endocrine pathways that regulate

metamorphosis, the IIS/TOR lifespan regulator pathway, and genes related to cell cycle and

DNA repair damage.

Loss of *PTTH/torso* promotes selection in the EGFR pathway

The loss of *PTTH and torso* along with positive selection in the *egfr15* and *SOS* genes (Figure 3)

suggest a change in the dynamics of the metamorphosis pathways when prepupal diapause is

lost. The onset of metamorphosis in holometabolous insects is regulated by the production and

release of ecdysone by the prothoracic gland (PG) in the brain (Henrich et al. 1999). When PTTH

binds to its tyrosine kinase receptor TORSO in the PG, the Ras/RAF/Erk pathway is activated,

which stimulates the release of ecdysone (Pan & O'Connor 2021). Although unknown for bees,

in other insects PTTH also plays a pivotal role in larval and pupal diapause. During diapause, the

289 brain-PG axis is shut down, inhibiting PTTH production and release, subsequently reducing

290 production of ecdysteroids, resulting in delayed metamorphosis (Figure 3) (Denlinger 2022).

292 For a considerable period, PTTH/TORSO was considered the primary pathway for ecdysone

293 production through the activation of the PG (Rewitz et al. 2009). However, there are at least

294 four other tyrosine kinase receptors expressed in the PG. A recent study demonstrated that

295 most of these are dispensable, because EGFR is the only receptor whose absence prevents

296 metamorphosis in flies (Cruz et al. 2020). This suggests EGFR is the ancestral ecdysone

297 biosynthesis pathway, with PTTH/TORSO likely co-opted in holometabolous insects to finely

298 adjust the timing of pupation in response to environmental cues (Shimell et al. 2018; Cruz et al.

299 2020). Recurrent *PTTH* and *torso* losses across species support the hypothesis that EGFR is the

300 foundational pathway regulating metamorphosis (Cruz et al. 2020). Moreover, it suggests that

301 PTTH/TORSO are dynamic elements susceptible to environmental influences (Skelly et al.

302 2019). Evidence of positive selection in two genes of the EGFR pathway indicates an amplified

303 role of this pathway in the absence of *PTTH* and *torso* in bees (Figure 3).

305 In Hymenoptera, the loss of *PTTH* and *torso* appears linked to the evolution of sociality in bees,

306 as all examined social bee genomes lack these genes (Costa et al. 2022) (Figure 1). Our findings

307 suggest that not only sociality, but also the loss of developmental diapause, predicts *PTTH* and

308 *torso* loss, with the latter being a stronger predictor. Since the loss of developmental diapause

309 is also a strong predictor of sociality in bees, future analysis of additional bee genomes will be

310 necessary to fully separate these effects.

The near-universal loss of *PTTH* and *torso* only in species lacking prepupal diapause suggests

strong selection against this pathway in the species that do not go through diapause during

development. In *Drosophila*, reducing expression of *PTTH* or *torso* delays pupariation (Shimell

et al. 2018; Kang et al. 2023; Rewitz et al. 2009), while constitutively knocking down *PTTH* or

torso in the PG results in prolonged adult lifespan, suggesting possible detrimental effects of

PTTH and *torso* in adulthood (Kang et al. 2023). This aligns with the antagonist pleiotropy

theory of aging (Williams 1957), where genes like *PTTH* and *torso* that confer benefits early in

life (e.g. promoting survival via diapause induction), but with detrimental effects later in life

(reduction of adult lifespan), are favored by natural selection and contribute to aging.

We propose that *PTTH* and *torso* loss is a consequence of the loss of prepupal diapause and

precedes the evolution of sociality. Selection may act against the PTTH/TORSO pathway once it

is no longer needed for diapause functions due to its lifespan reducing effects in adults. In this

way, the adult lifespan extension requisite for eusociality may be a result of reduced

antagonistic pleiotropy, freeing PTTH/TORSO from the selection shadow once their function in

early life was eliminated.

Loss of developmental diapause relaxes selection in the IIS/TOR pathways

Several genes in the insulin/insulin-like signaling (IIS) and target of rapamycin (TOR) pathways

are under relaxed selection in species that have lost prepupal diapause (Figure 4). These

pathways are highly conserved among metazoans and are key regulators of aging (Bitto et al.

2015; Guarente & Kenyon 2000). IIS and TOR work in parallel and interactively, such that when

either or both are downregulated, there is an increase in lifespan (Kapahi & Zid 2004;

McCormick et al. 2011).

In our study, *FoxO*, a pro-longevity gene in *Drosophila*, is under purifying selection in bee

species that lost prepupal diapause, reinforcing the importance of this gene, independent of

the diapause stage. All the anti-longevity genes in this pathway are under relaxed selection

(Figure 4). The exception is *mth*, which has both pro- and anti-longevity properties. Relaxed

selection in genes of the IIS/TOR pathway has been associated with both reduction of lifespan,

as observed in annual vs long-lived killifish (Cui et al. 2019), and increased lifespan as seen in

long vs short-lived mammals (Liu et al. 2023). Moreover, aging genes are under relaxed

selection in humans compared to chimpanzees, despite humans living approximately twice as

long (Cui et al. 2019). Similarly, relaxed selection on genes in the IIS/TOR pathway in bees that

have lost prepupal diapause could be associated with increased lifespan.

The hyperfunction theory of aging predicts that reduction in lifespan results from the

inappropriate activation of IIS/TOR pathway when growth is no longer beneficial (Blagosklonny

2006; Gems 2022). In accordance with this hypothesis, relaxed selection in IIS/TOR genes could

be associated with loss or shifts in function in some of the genes in this pathway, which could

result in reduced activation and increased lifespan. However, IIS/TOR is an essential pathway

for holometabolous development, so a complete loss of function is unlikely (Dutriaux et al.

2013; Fernandez et al. 1995; Hansen et al. 2004). Relaxed selection has also been associated

with the evolution of new phenotypes through the release of constraint in traits that were

under opposing selection pressures in different environments (Hunt et al. 2011). In

Hymenoptera, relaxed selection of caste-biased genes preceded the evolution of sociality and

could be the basis for phenotypic plasticity. Moreover, in honeybees, fast evolving genes were

more likely to be differentially expressed between castes (Hunt et al. 2011).

Loss of constraint in the IIS/TOR pathway suggests that different responses to nutritional cues

could evolve, a hallmark of caste differentiation in many social species (Kapheim 2017).

Differences in diet, either qualitative or quantitative, are associated at some level with caste

determination in all independent origins of castes in bees (Starr 2020; Carnell et al. 2020;

Kapheim et al. 2011; Richards & Packer 1994; Rehan & Richards 2010). A notable instance of a

unique nutritional response is observed in honeybee queens. Consumption of royal jelly, a

highly nutritious food, by the queens, leads to decreased activation of insulin pathway,

resulting in lifespan extension and increased fertility (Corona et al 2007). This is the opposite

effect of what is expected to other animals, in which a reduction of activity in the insulin

pathway leads to reduced lifespan and fertility (Ament et al. 2008). Relaxed selection on genes

in the IIS/TOR pathway could allow for such reversals to evolve.

Interestingly, a link between nutrition, caste differentiation, and diapause has long been

recognized (Hunt and Amdam 2005; Hunt 2006; Hunt et al. 2007). Our results suggest a possible

mechanism for this connection. Specifically, shifts in the timing of diapause may have allowed

molecular pathways important during development, such as IIS/TOR, to become co-opted for

social evolution (Hunt and Amdam 2005). We suggest that relaxation of the IIS/TOR pathway, as

consequence of the loss of prepupal diapause, precedes the evolution of sociality and could be

related to lifespan extension and/or the evolution of new phenotypes in response to

differences in nutrition.

Many cell cycle regulation and DNA damage repair genes are under positive selection in

species lacking prepupal diapause

Many genes involved in cell cycle regulation and DNA damage repair are subject to selection in

bee species that have lost prepupal diapause. These two biological processes are crucial for

diapause and aging mechanisms.

Diapause in the adult stage may require specific adaptations to maintain cell cycle arrest

without accumulating damage, given the lack of somatic tissue renewal in adults of

holometabolous insects (Tatar & Yin 2001). For instance, Fizzy (*fzr*), which is under positive

selection in bee species lacking prepupal diapause (Table S4), is involved in the natural

response to injuries in adult flies. Endocycles (consecutive cycles of cell growth and arrest in the

M phase), rather than mitosis, is the default response to injury in adult flies (Cohen et al. 2018).

However, the expression of *fzr* alone is sufficient to override the endocycle response, activating

mitosis after injury instead (Cohen et al. 2018). Positive selection of the *fzr* gene in bees that

have lost prepupal diapause suggests a potential mechanism to prevent organ injuries under

adverse environmental conditions, which could indirectly extend adult lifespan.

Other genes involved in cell cycle regulation and cell proliferation control, such as tumor

suppressor genes, are under positive selection in species that have lost prepupal diapause

(Table S4). Examples include *G1/S-specific cyclin-E1* and *cyclin-dependent kinase 14*, which

regulates the transition between phases G1 and S of the cell cycle (Chen et al. 2019; Fagundes

& Teixeira 2021), *ST7* and *TRIM33* that act as tumor suppressor genes (Charong et al. 2011; Xue

et al. 2015), and *Kruppel-like factor 6,* a transcription factor involved in the progression of

multiple malignant tumors (Sabatino et al. 2019). *p53* (Table 1), which also regulates cell cycle

arrest at both G1/S and G2/M checkpoints (Engeland 2018), is under purifying selection.

However, it remains unclear how selection on these genes could promote lifespan extension.

DNA repair mechanisms are crucial in preventing aging. Genes like *INO80 complex subunit D*,

which is involved in DNA recombination and DNA repair (Poli et al. 2017; Horigome et al. 2014),

and *ASH2*, which acts upstream of cellular response to DNA damage stimulus (Burgess et al.

2014), are under positive selection and are potentially adaptive for lifespan extension (Table

S4). Additionally, the gene ontology term related to double-strand break repair (p = 0.031,

Table S6) is enriched among the genes under relaxation of selection, suggesting loss of

constraint and possible evolution of new phenotypes associated with DNA repair in those bee

species.

Conclusions

Our study supports the hypothesis that convergent evolutionary changes associated with the

loss of prepupal diapause is related to molecular pathways that might regulate adult lifespan

extension. Our findings indicate that this extension appears to stem from a combination of

factors, such as the loss of *PTTH* and *torso*, coupled with a relaxation of selection in the IIS/TOR

pathway. This relaxation of selection, due to the loss of prepupal diapause, may facilitate the

emergence of new phenotypes, such as increased lifespan or altered responses to nutritional

signals, two important adaptations in the evolution of castes in social insects. Given the strong

correlation between loss of prepupal diapause and the evolution of social behavior in bees,

disentangling the effects of each phenotype on genome evolution proves challenging.

Therefore, advancing our understanding of the link between diapause and the evolution of

sociality demands expanding our understanding of the natural history of species presenting

different life history traits. Solitary bee species lacking prepupal diapause offer ideal

opportunities for investigating lifespan extension, examining signs of relaxed selection in *PTTH*

and *torso* (if these genes are still present) compared to solitary species retaining prepupal

diapause, and exploring potential innovations in how IIS/TOR pathway responds to diverse

stimuli.

Material and Methods

Genomes

For the comparative genomics analysis, we obtained bee genomes from the National Center for

Biotechnology Information (NCBI) (https://www.ncbi.nlm.nih.gov/datasets), the National

Genomics Data Center-China National Center for Bioinformation (NGDC-CNCB)

(https://ngdc.cncb.ac.cn/), and the Halictid Genome Browser

(https://beenomes.princeton.edu). Specifically, we selected genomes for which annotation files

were accessible and for which we possessed information regarding the diapause stage and the

social level of the bees (Table S1). Solitary and communal species were classified as 'solitary,'

while subsocial, primitively social, and advanced eusocial species were classified as 'social.'

Species that undergo diapause in the last larval instar before pupation were categorized as

having 'prepupal' diapause, whereas species with adult, reproductive, or no diapause were

classified as 'no prepupal'.

With the genome and annotation files, we used the agat-v0.9.2 software (Dainat 2023) to

retrieve the longest isoform for each gene. Subsequently, based on the longest isoform, we

extracted and translated the nucleotide coding sequences (CDS) for each gene. The protein

coding sequences were then used to run BUSCO-v5.3.2 (Simao et al. 2015) against the

Hymenoptera database, which encompasses 5991 genes. To ensure data quality, we retained

genomes in which > 80% of the proteins were classified as complete by BUSCO.

In the genome selection process, we retained one genome per genus, opting for the genome

with the highest percentage of complete proteins compared to the BUSCO database. More than

one genome per genus was kept in cases where distinct life history traits, such as solitary versus

social versus parasitic lifestyle, or variations in the diapause stage (prepupa versus adult) were

evident. Ultimately, a total of 27 genomes were included in the subsequent analysis (Figure 1).

Identification of Orthogroups

The protein coding sequences from the 27 different species served as input for Orthofinder-

v2.5.4 (Emms & Kelly 2015) to classify orthogroups. To proceed with further analysis, single-

copy orthologs were necessary. Orthofinder identified 2592 orthogroups with single-copy

orthologues. To augment the number of orthogroups for subsequent analysis, we filtered the

"Orthogroups.GeneCounts" table (output from Orthofinder) to retain orthogroups present in all

species, but single-copy in at least 70% of those species. This process yielded 4173 orthogroups.

In instances where species had paralogs, all copies were excluded from the orthogroups before

conducting the molecular evolution analysis, resulting in some orthogroups containing fewer

than 27 species.

Orthogroups nucleotide alignments and trees

We compiled the nucleotide coding sequences of all species into a single file to construct a

database. Utilizing the IDs of each sequence belonging to an orthogroup (generated using the

protein sequences), we retrieved the nucleotide sequences by comparing them with the

database. Species with more than one orthologous sequence (paralogs) in a given orthogroup

were excluded. Subsequently, we employed the orthogroup nucleotide sequences as input for

the MACSE_ALFIX_v01.sif pipeline, which encompasses codon alignment (Ranwez et al.

2021,https://hal.science/hal-03099847/document). The alignment pipeline included several

steps to ensure high-quality alignments. First, a pre-filtering step was performed to mask

potential untranslated regions (UTRs) and long non-homologous fragments. Next, HMMCleaner

was used to filter and mask amino acid residues that appeared to be misaligned. In the post-

processing stage, isolated codons were masked, and sequences that were more than 80%

masked were removed. Finally, the extremities of the alignments were trimmed to ensure that

at least 70% of the nucleotides (excluding gaps and ambiguous bases) were present at the first

and last sites. The orthogroup OG0003883 was excluded from the subsequent analysis due to

its alignment never concluding.

The resulting nucleotide alignment underwent filtering by removing positions with excessive

ambiguous data using the pxclsq function from the phyx-v1.3 software (Brown et al. 2017).

Specifically, alignment columns missing information for over 50% of the species were

eliminated. The filtered alignment was then used as input for iqtree2-v2.2.0 (Minh et al. 2020;

Kalyaanamoorthy et al. 2017) to generate phylogenetic trees for each orthogroup with the

options -st CODON and -m MFP, allowing for the automatic identification of the best codon

model for each alignment.

Molecular evolution analysis

RERconverge-v0.3.0 (Kowalczyk et al. 2019) was used to compute convergent evolutionary rate

shifts associated with the loss of developmental diapause. The input comprised nucleotide

alignments from MACSE and the species tree generated in Orthofinder. Some branches in the

species tree were adjusted using Mesquite-v3.81 (Maddison & Maddison 2023) to align with

the topology presented in (Almeida et al. 2023). The `estimatePhangornTreeAll` function with

the codon-based model YN98 (submodel = "YN98", type = "CODON") was employed to predict

the orthogroups tree while maintaining the same topology as the species tree. Four

orthogroups were excluded for the RERconverge analysis because they exceeded the cluster's

time limit for the branch length estimation step. Seventeen species that had lost prepupal

diapause (Figure 1) were used as foreground branches. Correlations were computed using the

terminal clades with unidirectional transitions.

To empirically calibrate the p-values, a permulation analysis was conducted (Saputra et al.

2021), as outlined in https://rdrr.io/github/nclark-

lab/RERconverge/f/vignettes/PermulationWalkthrough.Rmd. Briefly, 1000 permulated binary

trait trees were generated for each orthogroup using the Complete Case (CC) method with

Macropis europaea as the root species. The functions `getPermsBinary` and `permpvalcor` were

used to produce null p-values and correlation statistics. Empirical p-values were calculated as

the proportion of null statistics that were as extreme or more extreme than the observed

parametric statistics, followed by Benjamini-Hochberg correction. The analysis was performed

without specifying sister clades or including common ancestors. Orthogroups exhibiting

significant shifts in evolutionary rates were identified based on corrected permulation p-values

< 0.05. Orthogroups with Rho negative (Rho-) values indicate a slower evolutionary rate in the

foreground species, possibly reflecting increased selective constraint. Conversely, Rho positive

(Rho+) values denote genes undergoing faster evolution, which could result from a release of

selective constraint or adaptation. For additional details on the RERconverge pipeline, refer to

the GitHub repository.

To ascertain whether orthogroups exhibiting Rho+ values indicate an accelerated evolutionary

rate due to positive selection or relaxation of selection, we ran Hyphy-v2.5.55 (Kosakovsky

Pond et al. 2020) aBSREL (Smith et al. 2015) and RELAX (Wertheim et al. 2015). aBSREL

identifies signals of positive selection in single branches among the tested branches, whereas

RELAX discerns signals of convergent intensification of selection (K > 1, representing positive or

purifying selection) and relaxation of selection (K < 1) in tested (foreground) branches

compared to background branches. For both aBSREL and RELAX analyses, we used the

nucleotide alignment from MACSE and the corresponding trees from IQ-TREE as input files. Test

branches were specified as species that have lost developmental diapause. The parameters --

srv (yes) and --multiple-hits (Double+Triple) were employed to allow site-to-site synonymous

rate variation and multiple simultaneous hits. The threshold for positive selection in single

branches or convergent intensification or relaxation of selection in the tested branches was p =

0.05. aBSREL p-values was corrected by the Holm-Bonferroni method within Hyphy, and RELAX

p-values were corrected using the p.adjust function in the stats R package (R Core Team 2023)

using the Benjamini & Hochberg method.

Gene ontology enrichment

To identify gene ontology enrichment in orthogroups associated with the loss of developmental

diapause, we categorized the orthogroups of interest into four groups: possible positive

selection, possible purifying selection, possible relaxation of selection, and possible

intensification of selection. Possible positive selection included significant orthogroups with

Rho+ values in RERconverge and significant orthogroups from aBSREL that did not overlap with

relaxation of selection. Possible purifying selection included significant orthogroups with Rho-

values from RERconverge that did not overlap with aBSREL or relaxation of selection.

Orthogroups possibly under relaxation of selection were those significant in RELAX ($K<1$) that

did not overlap with aBSREL or RERconverge (Rho+ and Rho-) (Figure 2A). Lastly, orthogroups

under intensification of selection were those significant in RELAX ($K>1$) that did not overlap with

aBSREL or RERconverge (Rho+ and Rho-) (Figure 2B). The latter was considered a separate

category because it is not possible to differentiate between positive and purifying selection.

Gene ontology (GO) terms were annotated for the 4173 orthogroups using InterProScan-v5.63

(Jones et al. 2014) and the databases CDD, Pfam, SUPERFAMILY, and PANTHER. Out of these,

3267 orthogroups had at least one GO term annotated, forming the gene universe for the

enrichment analysis. We used topGO-v2.50.0 (Alexa & Rahnenfuhrer 2023) with the weight01

algorithm and Fisher's exact test statistics to identify enriched terms in each one of the

previously described categories. Terms were considered significant if they had a p-value < 0.01

and Annotated genes > 5.

Gene family evolution

To assess changes in gene families associated with the loss of developmental diapause, we used

the Orthogroup.GeneCount.tsv table generated in Orthofinder as input. The

`clade_and_size_filter.py` script from CAFE-v5.0 (Mendes et al. 2020) was used to filter out

gene families with 100 or more genes. The Base_count.tab output table was then used to

identify orthogroups that were gained or lost based on the diapause stage (prepupae diapause

or loss of developmental diapause). Filtering was performed for all species collectively and for

specific bee families with more than one species in each category (Apidae and Halictidae) (Table

S7) using a custom bash script.

Phylogenetic Generalized Least Squares (PGLS) analysis

We conducted a PGLS analysis using the R packages ape-v5.7.1 (Paradis & Schliep 2019) and

caper-v1.0.3 (Orme et al. 2023), to test whether the diapause stage and sociality are significant

predictors for the presence or absence of the prothoracicotropic hormone (*PTTH*) gene and its

receptor tyrosine-protein kinase receptor torso (*torso*). The dataset for this analysis included

the 27 species used in the molecular evolution analysis (Figure 1), along with *Exoneura robusta*

and *Exoneurella tridentata*. The phylogeny, generated by Orthofinder, encompasses all species,

and some branches were adjusted using Mesquite-v3.81 to align with the topology presented in

(Almeida et al. 2023). Additionally, a table of traits, including diapause stage, sociality, and

presence/absence of PTTH/torso, was utilized (Table S8).

The presence of *PTTH* and *torso* was first confirmed in the orthogroup files comprising the 29

species. Following this, a BLAST search was conducted using all protein coding sequence copies

of *PTTH* from the same orthogroup (11 sequences). This search was performed against the

genome of 118 species using tblastn and against the protein database of 48 species using blastp

(Table S9). Although additional species were included in this search, the discussion and PGLS

analysis were limited to the 29 species mentioned at the beginning of this section. This

limitation was due to a lack of information on diapause stages, lack of annotation for protein

retrieval and consequently the built of the phylogenetic tree, or because the species were from

already represented genus, leading to redundant information.

For both BLAST searches, the `max_target_seqs` was set to 100 and `evalue` to 1e-10. The

search for *torso* resulted in multiple hits across different genome locations, making it

challenging to identify the true ortholog accurately. Therefore, the presence of *torso* was only

verified in the orthogroup file. Since all species that lost *PTTH* also lost *torso*, we treated them

as a single trait in the PGLS analysis (Table S9). To model these traits, we used function pgls

from the R package caper-v1.0.3. Because diapause stage and sociality are correlated variables

(Fisher's exact test; p=0.0001), we employed the model with each variable one at a time

(PTTH.TORSO ~ Prepupae_Diapause; and PTTH.TORSO ~ Sociality) or both together

(PTTH.TORSO ~ Prepupae_Diapause + Sociality). The presence/absence of *PTTH* and *torso* was

set as a binary trait, with 0 indicating the absence and 1 indicating the presence of the

orthogroups. Diapause stage and sociality were set as categorical variables, yes/no for the

presence or absence or prepupal diapause, and solitary/social for social status.

Aging/longevity and oxidative genes

We compared the genes under selection with genes previously associated with aging/longevity

or oxidative stress in other species. For aging-related genes, we used the list of the *Drosophila*

aging-related genes obtained from the GenAge database (accessed in January 2024:

https://genomics.senescence.info/genes/index.html, (Tacutu et al. 2018)). We utilized Entrez

IDs to retrieve equivalent gene IDs and CDS nucleotide sequences from FlyBase

(https://flybase.org, Gramates et al. 2022). Concerning oxidative stress genes, we used as

reference the list of genes present in at least one of four Hymenoptera species from (Kramer et

al. 2021) and downloaded the sequences from Flybase based on the gene name.

For both sets of genes, we performed a reciprocal blast search against the translated nucleotide

coding sequences (CDS) of all 27 species combined in one file (322804 sequences). Initially, a

tblastn was conducted using either the aging or oxidative stress genes as the database and the

translated CDS sequences as query. Subsequently, using blastx, we used all translated CDS

sequences as the database and either the aging or oxidative stress genes as queries. In all four

searches `max_target_seqs` was set to 1 and `evalue` to 1e-10. The reciprocal best hits were

identified using the searchRBH.sh script adapted from

https://morphoscape.wordpress.com/2020/08/18/reciprocal-best-hits-blast-rbhb/. In the final

table, we included both orthogroups IDs and gene IDs based on the sequence IDs from the fasta

files using a custom script. The orthogroup OG0005890, which is the best orthologous match to

the *Drosophila insulin receptor* gene (InR), was initially excluded from the analysis due to its

absence in one species. However, since OG0005890 also corresponds to an InR, it has been

included in Table 1.

To assess whether any of our gene groups under selection (positive, purifying, relaxation, or

intensification) were enriched for aging or oxidative stress genes, we used the phyper function

from the hypeR-v2.0.0 R package (Federico & Monti 2020).

Acknowledgments

The support and resources from the Center for High Performance Computing at the University

of Utah are gratefully acknowledged. Special thanks to Tim Delory for support with snakemake

and discussions of the data analysis pipeline. We are grateful to the Kapheim Lab group for

their comments on earlier versions of the manuscript.

Funding

This work was supported by a grant from the National Science Foundation (NSF-IOS #2142778

to KMK).

Data availability

All scripts, input, and output files used in the analyses are available at:

https://github.com/kapheimlab/Comparative_genomics_loss_prepupal_diapause. Hyphy json

files will be available upon request.

Table and Figure Legends

Table 1: Orthogroups under selection overlapping with the *Drosophila* aging genes list from

GenAge database (Tacutu et al. 2018). Longevity influence classifies the orthologs as pro- or

anti-longevity genes according to their effects on *Drosophila* lifespan. The asterisk (*) highlights

orthogroups that are also part of the best evidence of positive selection (Table 1).

Figure 1: The 27 species included in the molecular evolution analysis and their associated life-

history traits. The prepupal diapause stage is depicted in dark blue, while the loss of

developmental diapause is represented in lighter blue. Solitary bees are indicated by a stronger

yellow color, and social species are denoted by a lighter yellow. Purple represents the presence

of the prothoracicotropic hormone (PTTH) and its tyrosine-protein kinase receptor torso

(TORSO), while the absence of color indicates orthologs could not be found. The two species

marked with "?" had conflicting results regarding the presence of the genes, depending on

whether the search was conducted using the genome or the annotated protein-coding genes

set as a reference.

Figure 2: The Venn diagram depicts orthogroups categorized under relaxed (A) or intensified (B)

selection, which intersect with genes under positive selection (aBSREL), accelerated evolution

(Rho +), and/or decelerated evolution (Rho -). Different colors denote orthogroups potentially

experiencing positive (yellow), purifying (blue), relaxed (purple), or intensified (green) selection,

while dashed lines indicate orthogroups with the strongest evidence for positive (yellow) and

purifying (blue) selection. Refer to the Material and Methods section for comprehensive

classification details.

Figure 3: Simplified scheme illustrating two pathways regulating the production and release of

ecdysone by the prothoracic gland (PG). The pathway utilizing the tyrosine-protein kinase

receptor TORSO as a receptor for the prothoracicotropic hormone PTTH is absent in species

that do not undergo prepupal diapause. egfr substrate 15 (EGFR15), a substrate of the

alternative pathway that uses EGFR as a receptor, is subject to positive selection, along with the

son of sevenless (*SOS*) gene, highlighting the significance of this pathway in species without

prepupal diapause.

Figure 4: Simplified scheme of the insulin/insulin-like signaling (IIS) and target of rapamycin

(TOR) pathways containing genes under selection in the current study. When the insulin

pathway is activated (high nutrition), Akt phosphorylates FoxO, preventing its entrance to the

nucleus of the cell, and activates Tor, leading to growth. When there is a reduction in the

IIS/TOR pathway activation (low-nutrition), FoxO enters in the nucleus of the cells, regulating

thousands of genes, leading to lifespan extension. Continuous arrows represent activation of

the pathway in an environment with high nutrition, for example, while dashed arrows

represent the alternative pathway in environments with low nutrition. Genes within green,

blue, or purple boxes represent those under intensified, purifying, and relaxed selection in the

current study, respectively. The box in the bottom right classifies the genes as pro- or anti-

longevity based on their known effects on *Drosophila* lifespan (Tacutu et al. 2018). Figure made

based on Kapahi and Zid 2004; Dutriaux et al. 2013; Texada et al. 2020; Denlinger 2022.

References

Alexa A, Rahnenfuhrer J. 2023. topGO: Enrichment Analysis for Gene Ontology. R package
version 2.50.0.
Almeida EAB et al. 2023. The evolutionary history of bees in time and space. Current Biology.
33:3409-3422.e6. doi: 10.1016/j.cub.2023.07.005.
Ament SA, Corona M, Pollock HS, Robinson GE. 2008. Insulin signaling is involved in the
regulation of worker division of labor in honey bee colonies. 105.
Batra SWT. 1966. Social behavior and nests of some Nomiine beed in India (Hymenoptera,
Halictidae). Insectes Soc. 8:145–154.

Bitto A, Wang AM, Bennett CF, Kaeberlein M. 2015. Biochemical genetic pathways that modulate aging in multiple species. Cold Spring Harb Perspect Med. 5. doi: 10.1101/cshperspect.a025114.

Blagosklonny M V. 2006. Aging and immortality: Quasi-programmed senescence and its pharmacologic inhibition. Cell Cycle. 5:2087–2102. doi: 10.4161/cc.5.18.3288.

Brown JW, Walker JF, Smith SA. 2017. Phyx: Phylogenetic tools for unix. Bioinformatics. 33:1886–1888. doi: 10.1093/bioinformatics/btx063.

Burgess RC, Burman B, Kruhlak MJ, Misteli T. 2014. Activation of DNA Damage Response Signaling by Condensed Chromatin. Cell Rep. 9:1703–1717. doi: 10.1016/j.celrep.2014.10.060.

Carey JR. 2001. Demographic mechanisms for the evolution of long life in social insects. Exp Gerontol. 36:713–722. www.elsevier.nl/locate/expgero.

Carnell JD, Hulse RA, Hughes WOH. 2020. A review of nutrition in bumblebees: The effect of caste, life-stage and life history traits. In: Advances in Insect Physiology.Vol. 59 Academic Press Inc. pp. 71–129. doi: 10.1016/bs.aiip.2020.09.003.

Charong N, Patmasiriwat P, Zenklusen JC. 2011. Localization and characterization of ST7 in cancer. J Cancer Res Clin Oncol. 137:89–97. doi: 10.1007/s00432-010-0863-2.

Chen L et al. 2019. CDK14 involvement in proliferation migration and invasion of esophageal cancer. Ann Transl Med. 7.

Cohen E, Allen SR, Sawyer JK, Fox DT. 2018. Fizzy-Related dictates A cell cycle switch during organ repair and tissue growth responses in the Drosophila hindgut. Elife. 7:1–22. doi: 10.7554/eLife.38327.001.

Costa CP, Okamoto N, Orr M, Yamanaka N, Woodard SH. 2022. Convergent Loss of Prothoracicotropic Hormone , A Canonical Regulator of Development , in Social Bee Evolution. Front Physiol. 13:1–6. doi: 10.3389/fphys.2022.831928.

Cruz J, Martín D, Franch-Marro X. 2020. Egfr Signaling Is a Major Regulator of Ecdysone Biosynthesis in the Drosophila Prothoracic Gland. Current Biology. 30:1547-1554.e4. doi: 10.1016/j.cub.2020.01.092.

Cui R et al. 2019. Relaxed Selection Limits Lifespan by Increasing Mutation Load. Cell. 178:385-399.e20. doi: 10.1016/j.cell.2019.06.004.

Dainat J. 2023. AGAT: Another Gff Analysis Toolkit to handle annotations in any GTF/GFF format. https://www.doi.org/10.5281/zenodo.3552717.

Denlinger DL. 2022a. *Insect Diapause*. Cambridge University Press doi: 10.1017/9781108609364.

Denlinger DL. 2022b. Molecular Signaling Pathways that Regulate Diapause. In: Insect Diapause. Cambridge University Press pp. 240–292.

Dutriaux A, Godart A, Brachet A, Silber J. 2013. The Insulin Receptor Is Required for the Development of the Drosophila Peripheral Nervous System. PLoS One. 8:e71857. doi: 10.1371/journal.pone.0071857.

Easwaran S, Montell DJ. 2023. The molecular mechanisms of diapause and diapause-like reversible arrest. Biochem Soc Trans. 51:1847–1856. doi: 10.1042/BST20221431.

Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. Genome Biol. 16:157. doi: 10.1186/s13059-015-0721-2.

Engeland K. 2018. Cell cycle arrest through indirect transcriptional repression by p53: I have a
DREAM. Cell Death Differ. 25:114–132. doi: 10.1038/cdd.2017.172.
Fagundes R, Teixeira LK. 2021. Cyclin E/CDK2: DNA Replication, Replication Stress and Genomic
Instability. Front Cell Dev Biol. 9. doi: 10.3389/fcell.2021.774845.
Federico A, Monti S. 2020. hypeR: an R package for geneset enrichment workflows.
Bioinformatics. 36:1307–1308. doi: 10.1093/bioinformatics/btz700.
Fernandez R, Tabarini D, Azpiazu N, Frasch M, Schlessinger J. 1995. The Drosophila insulin
receptor homolog: A gene essential for embryonic development encodes two receptor isoforms
with different signaling potential. EMBO Journal. 14:3373–3384. doi: 10.1002/j.1460-
2075.1995.tb07343.x.
Gauld I, Bolton B. 1988. *The Hymenoptera*. Oxford University Press: Oxford.
Gems D. 2022. The hyperfunction theory: An emerging paradigm for the biology of aging.
Ageing Res Rev. 74. doi: 10.1016/j.arr.2021.101557.
Gilbert SF. 2000. The Hormonal Reactivation of Development. In: Developmental Biology.
Sinauer Associates: Sunderland, MA.
Gramates LS et al. 2022. FlyBase: a guided tour of highlighted features. Genetics. 220. doi:
10.1093/genetics/iyac035.
Guarente L, Kenyon C. 2000. Genetic pathways that regulate ageing in model organisms.
Nature. 408:255–262. www.nature.com.
Hansen IA, Attardo GM, Park JH, Peng Q, Raikhel AS. 2004. Target of rapamycin-mediated
amino acid signaling in mosquito anautogeny. Proc Natl Acad Sci U S A. 101:10626–10631. doi:
10.1073/pnas.0403460101.
Henrich VC, Rybczynski R, Gilbert LI. 1999. Peptide Hormones, Steroid Hormones, and Puffs:
Mechanisms and Models in Insect Development. Vitam Horm. 55:73–125. doi:
https://doi.org/10.1016/S0083-6729(08)60934-6.
Horigome C et al. 2014. SWR1 and INO80 chromatin remodelers contribute to dna double-
strand break perinuclear anchorage site choice. Mol Cell. 55:626–639. doi:
10.1016/j.molcel.2014.06.027.
Hunt BG et al. 2011. Relaxed selection is a precursor to the evolution of phenotypic plasticity.
Proc Natl Acad Sci U S A. 108:15936–15941. doi: 10.1073/pnas.1104825108.
Hunt JH et al. 2007. A diapause pathway underlies the gyne phenotype in *Polistes* wasps,
revealing an evolutionary route to caste-containing insect societies. Proc Natl Acad Sci U S A.
104:14020–14025. doi: 10.1073/pnas.0705660104.
Hunt JH. 2006. Evolution of castes in *Polistes*. Ann Zool Fennici. 43:407–422. doi:
10.2307/23736751.
Hunt JH, Amdam GV. 2005. Bivoltinism as an antecedent to eusociality in the paper wasp genus
Polistes. Science (1979). 308:264–267. doi: 10.1126/science.1109724.
Hutfilz C. 2022. Endocrine Regulation of Lifespan in Insect Diapause. Front Physiol. 13:1–18. doi:
10.3389/fphys.2022.825057.
Jones P et al. 2014. InterProScan 5: Genome-scale protein function classification.
Bioinformatics. 30:1236–1240. doi: 10.1093/bioinformatics/btu031.
Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. 2017. ModelFinder: fast
model selection for accurate phylogenetic estimates. Nat Methods. 14:587–589. doi:
10.1038/nmeth.4285.

Kang P et al. 2023. Ptth regulates lifespan through innate immunity pathway in Drosophila.
bioRxiv. doi: 10.1101/2023.09.30.560323.
Kapahi P, Zid B. 2004. TOR pathway: linking nutrient sensing to life span. Sci Aging Knowledge
Environ. 2004. doi: 10.1126/sageke.2004.36.pe34.
Kapheim KM. 2017. Nutritional, endocrine, and social influences on reproductive physiology at
the origins of social behavior. Curr Opin Insect Sci. 22:62–70. doi: 10.1016/j.cois.2017.05.018.
Kapheim KM, Bernal SP, Smith AR, Nonacs P, Wcislo WT. 2011. Support for maternal
manipulation of developmental nutrition in a facultatively eusocial bee, Megalopta genalis
(Halictidae). Behav Ecol Sociobiol. 65:1179–1190. doi: 10.1007/s00265-010-1131-9.
Kosakovsky Pond SL et al. 2020. HyPhy 2.5 - A Customizable Platform for Evolutionary
Hypothesis Testing Using Phylogenies. Mol Biol Evol. 37:295–299. doi:
10.1093/molbev/msz197.
Kowalczyk A et al. 2019. RERconverge: An R package for associating evolutionary rates with
convergent traits. Bioinformatics. 35:4815–4817. doi: 10.1093/bioinformatics/btz468.
Kramer BH et al. 2021. Oxidative stress and senescence in social insects: A significant but
inconsistent link? Philosophical Transactions of the Royal Society B: Biological Sciences. 376.
doi: 10.1098/rstb.2019.0732.
Liu W et al. 2023. Large-scale across species transcriptomic analysis identifies genetic selection
signatures associated with longevity in mammals. EMBO J. 42. doi: 10.15252/embj.2022112740.
Maddison WP, Maddison DR. 2023. Mesquite: a modular system for evolutionary analysis.
Version 3.81. http://www.mesquiteproject.org. http://www.mesquiteproject.org.
McCormick MA, Tsai SY, Kennedy BK. 2011. TOR and ageing: A complex pathway for a complex
process. Philosophical Transactions of the Royal Society B: Biological Sciences. 366:17–27. doi:
10.1098/rstb.2010.0198.
Mendes FK, Vanderpool D, Fulton B, Hahn MW. 2020. CAFE 5 models variation in evolutionary
rates among gene families. Bioinformatics. 36:5516–5518. doi:
10.1093/bioinformatics/btaa1022.
Michener CD. 1969. Comparative Social Behavior of Bees. Annu Rev Entomol. 14:299–342. doi:
10.1146/annurev.en.14.010169.001503.
Michener CD. 1974. *The social behavior of the bees : a comparative study*. Harvard University
Press: Cambridge.
Minh BQ et al. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. Mol Biol Evol. 37:1530–1534. doi: 10.1093/molbev/msaa015.
O'Leary NA et al. 2016. Reference sequence (RefSeq) database at NCBI: Current status,
taxonomic expansion, and functional annotation. Nucleic Acids Res. 44:D733–D745. doi:
10.1093/nar/gkv1189.
Orme D et al. 2023. caper: Comparative Analyses of Phylogenetics and Evolution in R.
Pan X, O'Connor MB. 2021. Coordination among multiple receptor tyrosine kinase signals
controls Drosophila developmental timing and body size. Cell Rep. 36. doi:
10.1016/j.celrep.2021.109644.
Paradis E, Schliep K. 2019. Ape 5.0: An environment for modern phylogenetics and evolutionary
analyses in R. Bioinformatics. 35:526–528. doi: 10.1093/bioinformatics/bty633.

Poli J, Gasser SM, Papamichos-Chronakis M. 2017. The INO80 remodeller in transcription,
replication and repair. Philosophical Transactions of the Royal Society B: Biological Sciences.
372. doi: 10.1098/rstb.2016.0290.
R Core Team. 2023. R: A Language and Environment for Statistical Computing.
Ranwez V, Chantret N, Delsuc F. 2021. Aligning Protein-Coding Nucleotide Sequences with
MACSE. In: pp. 51–70. doi: 10.1007/978-1-0716-1036-7_4.
Rehan SM, Richards MH. 2010. The influence of maternal quality on brood sex allocation in the
small carpenter bee, Ceratina calcarata. Ethology. 116:876–887. doi: 10.1111/j.1439-
0310.2010.01804.x.
Rewitz KF, Yamanaka N, Gilbert LI, O'Connor MB. 2009. The insect neuropeptide PTTH activates
receptor tyrosine kinase torso to initiate metamorphosis. Science (1979). 326:1403–1405. doi:
10.1126/science.1176450.
Richards MH, Packer L. 1994. Trophic aspects of caste determination in *Halictus ligatus*, a
primitively eusocial sweat bee. Behav Ecol Sociobiol. 34:385–391. doi: 10.1007/BF00167329.
Sabatino ME et al. 2019. Krüppel-Like Factor 6 Is Required for Oxidative and Oncogene-Induced
Cellular Senescence. Front Cell Dev Biol. 7. doi: 10.3389/fcell.2019.00297.
Santos C.G., Humann FC, Hartfelder K. 2019. Juvenile hormone signaling in insect oogenesis.
Curr Opin Insect Sci. 31:43–48. doi: 10.1016/j.cois.2018.07.010.
Santos P.K.F., Arias MC, Kapheim KM. 2019. Loss of developmental diapause as a prerequisite
for social evolution in bees. Biol Lett. 15:1–7. doi: 10.1101/649897.
Saputra E, Kowalczyk A, Cusick L, Clark N, Chikina M. 2021. Phylogenetic Permulations: A
Statistically Rigorous Approach to Measure Confidence in Associations in a Phylogenetic
Context. Mol Biol Evol. 38:3004–3021. doi: 10.1093/molbev/msab068.
Seifert AW et al. 2012. The influence of fundamental traits on mechanisms controlling
appendage regeneration. Biological Reviews. 87:330–345. doi: 10.1111/j.1469-
185X.2011.00199.x.
Shimell MJ et al. 2018. Prothoracicotropic hormone modulates environmental adaptive
plasticity through the control of developmental timing. Development (Cambridge). 145. doi:
10.1242/dev.159699.
da Silva J. 2021. Data from: The extension of foundress lifespan and the evolution of eusociality
in the Hymenoptera [Dataset]. Dryad. https://doi.org/10.5061/dryad.6wwpzgn0p.
da Silva J. 2021. Life History and the Transitions to Eusociality in the Hymenoptera. Front Ecol
Evol. 9:1–20. doi: 10.3389/fevo.2021.727124.
da Silva J. 2022. The Extension of Foundress Life Span and the Evolution of Eusociality in the
Hymenoptera. Am Nat. 199. doi: 10.1086/718594.
Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinformatics.
31:3210–3212. doi: 10.1093/bioinformatics/btv351.
Skelly J, Pushparajan C, Duncan EJ, Dearden PK. 2019. Evolution of the Torso activation
cassette, a pathway required for terminal patterning and moulting. Insect Mol Biol. 28:392–
408. doi: 10.1111/imb.12560.
Smith MD et al. 2015. Less is more: An adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. Mol Biol Evol. 32:1342–1353. doi:
10.1093/molbev/msv022.

Starr CK. 2020. *Encyclopedia of Social Insects*. Springer International Publishing.
Szathmáry E, Smith JM. 1995. The major evolutionary transitions. Nature. 374:227–232. doi:
doi.org/10.1038/374227a0.
Tacutu R et al. 2018. Human Ageing Genomic Resources: new and updated databases. Nucleic
Acids Res. 46:D1083–D1090. doi: 10.1093/nar/gkx1042.
Tatar M, Yin C. 2001. Slow aging during insect reproductive diapause : why butterflies ,
grasshoppers and flies are like worms. Exp Gerontol. 36:723–738.
Texada MJ, Koyama T, Rewitz K. 2020. Regulation of body size and growth control. Genetics.
216:269–313. doi: 10.1534/genetics.120.303095.
Wertheim JO, Murrell B, Smith MD, Pond SLK, Scheffler K. 2015. RELAX: Detecting relaxed
selection in a phylogenetic framework. Mol Biol Evol. 32:820–832. doi:
10.1093/molbev/msu400.
Williams GC. 1957. Pleiotropy, Natural Selection, and the Evolution of Senescence. Evolution (N
Y). 11:398–411.
Wilson EO. 1971. *The Insect Societies*. Belknap Press of Harvard University Press.
Xue J et al. 2015. Tumour suppressor TRIM33 targets nuclear β-catenin degradation. Nat
Commun. 6. doi: 10.1038/ncomms7156.
897