Artesunate, EDTA and colistin work synergistically against MCR-negative and

-positive colistin-resistant *Salmonella*

Yajun Zhai[1]†, Peiyi Liu[1]†, Xueqin Hu[1]†, Changjian Fan[1], Xiaodie Cui[1], Qibiao He[1], Dandan He[1], Xiaoyuan

Ma[1], Gongzheng Hu[1*], Yajun Zhai[1*]

[1]Department of Pharmacology and Toxicology, College of Veterinary Medicine, Henan Agricultural University,

Zhengzhou, Henan 450046, China

† The authors contributed equally as the first author.

*Corresponding authors Address: College of Veterinary Medicine, Henan Agricultural University, Zhengzhou

450002, People's Republic of China;

E-mail address: zyj90518@126.com (Yajun Zhai), yaolilab@126.com (Gongzheng Hu)

Running title: Synergy of artesunate, EDTA and colistin

Abstract

Discovering new strategies to combat the multi-drug resistance bacteria constitutes a major medical challenge of our time. Previously, artesunate (AS) has been reported to exert antibacterial enhancement activity in combination with β-lactam antibiotics, via inhibition of the efflux pump AcrB. However, combination of AS and colistin (COL) revealed weak synergistic effect against a limited number of strains, and few studies have further explored its possible mechanism of synergistic action. In this paper, we found that AS and EDTA could strikingly enhance the antibacterial effects of COL against mcr-1^- and mcr-1^+ $Salmonella$ strains either in $vitro$ or in $vivo$, when used in triple combination. The excellent bacteriostatic effect was primarily related to the increased cell membrane damage, accumulation of toxic compounds and inhibition of MCR-1. The potential binding sites of AS to MCR-1 (THR283, SER284, and TYR287) were critical for its inhibition of MCR-1 activity. Additionally, we also demonstrated that the CheA of chemosensory system and virulence-related protein SpvD were critical for the bacteriostatic synergistic effects of the triple combination. Selectively targeting CheA, SpvD or MCR using the natural compound artesunate could be further investigated as an attractive strategy for treatment of $Salmonella$ infection. Collectively, our work opens up avenues towards the potentiation of colistin and revealed an alternative drug combination strategy to overcome COL resistant bacterial infections.

Keywords: artesunate/EDTA/colistin/triple combination/colistin resistant/$Salmonella$

Introduction

Infections of antibiotic-resistant pathogens, especially the carbapenemase-producing *Enterobacteriaceae*, pose an impending catastrophe of a return to the pre-antibiotic era. The drying up of the antibiotic discovery pipeline has resulted in increased use of polymyxins as the "last resort" antimicrobial drug with the inevitable risk of emerging resistance (Falagas & Kasiakou, 2005). Colistin (polymyxin E, COL), a fatty acyl oligopeptide antibiotic, is an active agent against multi-drug resistant (MDR) Gram-negative (G⁻) pathogens. Generally, COL kills bacteria through a detergent-like effect, that the polycationic ring of COL electrostatically interacted with the cell envelope components, causing the competitive displacement of divalent cations calcium (Ca^{2+}) and magnesium (Mg^{2+}), destabilizing the membrane, thus killing the bacterium via the "self-promoted uptake" pathway (Kaye *et al*, 2016). Beyond that, other models for the antibacterial activity have been reported, including vesicle-vesicle contact, hydroxyl radical death, inhibition of respiratory enzymes and anti-endotoxin COL activity pathways (El-Sayed Ahmed *et al*, 2020). Until now, numerous chromosomally- or plasmid-mediated mechanisms underlying polymyxins resistance in G⁻ bacteria have been identified, including intrinsic, mutation (eg. PmrAB, PhoPQ or AcrAB-TolC mutants), adaptation mechanisms or horizontally acquired resistance via the phosphoethanolamine (pEtN) transferase genes *mcr-1* to *9* (Carroll *et al*, 2019; Lima *et al*, 2018; Poirel *et al*, 2017).

To tackle the increasing emergence of MDR pathogens, many alternative therapies, less costly and time-consuming, become the areas of current research interest, involving the combination therapy of existing agents, the drug discovery from natural products, and the evaluation of drug resistance reversers (Rosenthal, 2003). A variety of antibiotic adjuvants that may or may not have direct antibacterial effects have been widely investigated to increase the effectiveness of current antibiotics or delay the emergence of drug resistance, such

as β-lactamase inhibitors, aminoglycoside-modifying enzyme inhibitors, membrane permeabilisers, and efflux

pump inhibitors (Laws *et al*, 2019). Several promising inhibitors, for example zidebactam and

pyrazolopyrimidine compounds have been described as the β-lactam and aminoglycoside enhancers against G⁻

bacteria (Moya *et al*, 2017; Stogios *et al*, 2013). Alternatively, numerous synthetic antimicrobial peptides and

plant-derived natural products have been shown to possess membrane permeabilising or efflux pump inhibitory

activity, with the combination of azithromycin, ciprofloxacin, imipenem *et al* (Aron & Opperman, 2016; Lin *et*

al, 2015; Su & Wang, 2018).

Artesunate (AS) is a semi-synthetic derivative of anti-malarial compound artemisinin that extracted from

the traditional Chinese herb Artemisia annua. Beyond remarkable antimalarial action, AS and other artemisinin

derivatives, eg. dihydroartemisinin (DHA), have been proven to restored the antibacterial effect of COL against

Escherichia coli (*E. coli*), while themselves didn't exhibited intrinsic antimicrobial activity against clinical *E.*

coli isolates as well as ATCC 25922 (Wei *et al*, 2020; Zhou *et al*, 2022). In addition, AS has also been proved to

enhance the effectiveness of various β-lactam and fluoroquinolones antibiotics against MDR *E. coli* via

inhibiting the efflux pump AcrAB-TolC (Pan *et al*, 2020; Wei *et al*, 2020). Nonetheless, the synergistic effects

between AS and COL were only observed in a limited number of strains, with a modest reversal effect.

Therefore, few studies have been undertaken to evaluate its underlying mechanism. Under this circumstance, it

is meaningful to explore new medication regimens between AS and COL against MDR bacteria. Encouragingly,

in this study, we confirmed the prominent synergistic effects of AS and EDTA to restore the antimicrobial

activity of COL and its possible molecular mechanism.

Results

88 **Artesunate and EDTA could enhance the effects of colistin against *Salmonella* strains**

89 The antimicrobial activities of AS, EDTA or COL alone were initially investigated to the COL-sensitive strains

90 of *Salmonella* (JS, S34), COL-resistant clinical strains of *Salmonella* (S16, S20, S13, and S30), *E. coli* (E16),

91 and intrinsically COL-resistant species (*Morganella morganii* strain M15, *Proteus mirabilis* strain P01). Results

92 showed that AS or EDTA alone had no direct antibacterial activity against these strains, with the minimum

93 inhibitory concentrations (MICs) 1250 or > 125 mg/L (Table supplement 1). Excepting for the two intrinsically

94 COL-resistant strains M15 and P01, there were only slight decrease of COL MICs for other strains (fold changes

95 ranging from 0 to 133), when the sub-inhibitory concentrations (1/4, 1/8, 1/16 MIC) of AS or EDTA were

96 combined with COL (namely AC or EC) (Table supplement 1). Whease, we found marked decrease of COL

97 MICs (up to 60000 fold), after three drug combinations (namely AEC) (Table 1). These results indicated that

98 when used simultaneously with COL, AS and EDTA exerted antibacterial enhancement activity for *Salmonella*

99 and *E. coli*, but not the intrinsically COL-resistant species. Thus, AS and EDTA could be considered as

100 adjuvants to reverse COL resistance in *Salmonella*.

101 To verify the antibacterial enhancement activity of AS and EDTA, the growth curves of *Salmonella* JS, S16

102 (*mcr-1⁻*), and S30 (*mcr-1⁺*) for the combined treatments were generated within 12 h (Figure 1). Generally, the

103 higher concentration of COL (2 mg/L), either alone or in combinations, was found more effective against these

104 strains than that of the lower concentration (0.1 mg/L). Compared with the control groups, when these bacteria

105 were grown in the presence of COL alone (0.1 mg/L) or two drug combinations (AC and EC), the antimicrobial

106 activity were not significant, and only the three drug combination AEC exhibited a certain degree of growth

107 inhibition after incubation for 8-12 h (Figure 1a, b, c). By contrast, higher concentration of COL (2mg/L), alone

108 or in combinations, showed better antibacterial activity, and a gradual increase in antibacterial activity was

observed: AEC > AC > EC > C (Figure 1d, e, f). It's worth noting that the effect of different combinations

against *mcr-1*[+] strain S30 was weaker than that of *mcr-1*[-] S16 and standard sensitive strain JS.

Artesunate and EDTA enhanced the membrane-damaging effect of colistin on *Salmonella*

In order to gain insight into the membrane-damaging bactericidal mechanism of COL alone or combinations,

the damages to the bacterial outer and inner membrane (OM and IM) were severally monitored by measuring

the fluorescent intensity of *Salmonella* strains S16 and S30 mixed with NPN and PI. Overall, the fluorescence

signals of NPN and PI increased progressively with increases in the concentration of COL. After the treatment

of AEC, S16 and S30 both showed strongest fluorescence signals of NPN, compared to those of other groups

(Figure 2a, b). Nevertheless, AS treatment group also exerted a significant increase of fluorescent signal,

although it is not as strong as that of AEC treated group. Meanwhile, there were rapid and significant increase in

fluorescence of PI, when AEC or EC were added to bacterial cultures, and the two regimens played dominant

roles in low or high concentrations of COL, respectively (Figure 2c, d). Therefore, these results indicated that

the bacterial cell surfaces were severely damaged after the AEC treatment, via the rapid perturbations of OM

and IM. Subsequently, the morphological changes of *Salmonella* strain S16 treated with different regimens were

further investigated using Scanning electron microscope (SME) analysis to confirm the above

membrane-damaging effect. As shown in Figure 3, SEM micro-graphs of control and solvent treated groups

revealed that cells were short and rod-shaped, with rounded ends and intact cell membranes. As expected, after

exposure to COL alone and different regimens, noticeable damage to the OM were observed, especially in AC

and AEC treated group. Exposure to AC and AEC leading to cell damages characterized by folds, crevices, and

depressions, which further suggested that AS and EDTA combined with COL resulted in remarkable bacterial

membrane injury for antibacterial activity. Collectively, these data confirmed the membrane-damaging effects

of AS, EDTA and COL combination against *Salmonella*.

AEC combination could collapse the Δψ component of Proton Motive Force (PMF) in *Salmonella*

In bacteria, the PMF, alternatively known as electrochemical proton gradient, is result from the extrusion of

protons by the electron transport chain, and made up of the sum of two parameters: an electric potential ($\Delta\psi$)

and an osmotic component (ΔpH) (Farha *et al*, 2013). It has been reported to drives vital cellular processes in

bacteria, including ATP synthesis, antibiotic transport, and cell division (Le *et al*, 2021). Therefore, PMF

dissipation was considered as a promising strategy for combating microbial pathogens. In this work, we

explored whether the antibacterial synergism activities of different combinations were accompanied by

dissipation of PMF in cells. We uncovered that when compared to that of other groups, AEC treatment caused a

rapid collapse of $\Delta\psi$ component, as shown by the increase in fluorescence values of $DISC_3(5)$ (Figure 2e, f). But

none of these combinations observed significant dissipation of ΔPH, when compared to that of control group

(Figure 2g, h). The above results indicated that AEC treatment was able to dissipate selectively the $\Delta\psi$

component of PMF.

Other reactive oxygen species (ROS), not H_2O_2, contributed to the AS and EDTA-mediated efficacy

enhancement of COL

ROS including superoxide (O^{2-}), hydrogen peroxide (H_2O_2), and hydroxyl radical ($\cdot OH$), are commonly

generated during the electron transfer process, which has been considered to associated with the lethal action of

diverse antimicrobials. Subsequently, we investigated whether the addition of AS and EDTA could facilitate the

intracellular ROS generation and stimulate the ROS-mediated killing. As shown in Figure 2i, j, compared to

control group, total ROS increases in S16 and S30 strains were observed in AC and AEC groups either after 0.1

or 2 mg/L COL was added, but for COL alone and EC groups, the increase were only occured after the addition

of a relatively high COL concentration (2 mg/L). In addition, the ROS accumulation level was further increased

in AEC group, when extending the incubation time period to 6 h (Figure supplement 1). Moreover, we found

that H_2O_2 did not contribute to the increase of total ROS, as there was no significant difference in intracellular

level of H_2O_2 among all the groups Figure 2k, l. Collectively, these observations support a role for ROS in AS

and EDTA-mediated efficacy enhancement of COL.

The transcriptome data exhibited more robust changes than that of metabolome among different

comparison groups.

A total of 6944 differentially expressed genes (DEGs) were performed KEGG pathway enrichment analysis, of

which 1832 and 5112 transcripts were included in S16 and S30 strains, respectively (Figure supplement 2).

Several canonical pathways, including two-component system (TCS), flagellar assembly and ABC transporters

pathways, indicating similar directional changes in both strains were selected for further analysis, as shown in

Figure 4. Since AEC incubation has displayed excellent antibacterial effects, we expected to screen the

significantly differentially expressed genes (SDEGs) with similar variations in AEC vs. C, AEC vs. AC, and

AEC vs. EC groups, and the SDEGs involved in TCS (*pagC*, *cheA*, *ompF*, *et al.*), flagellar assembly (*flgK*, *flgL*,

fliD, *et al*) and ABC transporters (*oppuBB*, *osmX*, *gltI*, *dppA*, *et al.*) were selected (mostly down-regulated) and

summarized in Figure 5.

Unlike transcriptome changes, the metabolite alterations were much less abundant among AEC vs. C, AEC

vs. AC, and AEC vs. EC groups, either in S16 or S30 strain. Unfortunatly, we demonstrated that there were low

correlation between the metabolome and transcriptome data. According to the enrichment analysis, archidonic

acid metabolism (down-regulated), degradation of aromatic compounds (up-regulated), taurine and hypotaurine

metabolism (up-regulated) were the most prominent pathways showing differences primarily in AEC vs. C

group (Figure 6).

AS+EDTA+COL combination therapy is a promising therapeutic against *Salmonella* infection *in vivo*

The excellent bacteriostatic synergistic effect against *Salmonella in vitro* of AEC combination further prompted

us to confirm the effect *in vivo* for *Salmonella* S30 (*mcr-1*$^+$) infected mouse models. Consistent with the

synergistic bactericidal activity of AS, EDTA, and COL, the combination of AC (7.13 $\log_{10}$CFU/g Liver, 6.91

$\log_{10}$CFU/g spleen), EC (7.33 $\log_{10}$CFU/g Liver, 6.88 $\log_{10}$CFU/g spleen), and AEC (6.51 $\log_{10}$CFU/g Liver,

6.52 $\log_{10}$CFU/g spleen) outperformed single-drug treatments of COL (7.44 $\log_{10}$CFU/g Liver, 7.05 $\log_{10}$CFU/g

spleen) and AS (7.54 $\log_{10}$CFU/g Liver, 7.02 $\log_{10}$CFU/g spleen). In particular, in the AEC combination-treated

samples, there were far fewer bacteria burden in spleen and liver when compared to other groups (Figure 7).

These disparity *in vivo* further illustrate the synergy of AS and EDTA in combination with COL.

Discussion

Membrane-damaging played a crucial role in the bacteriostatic synergistic effects of AS+EDTA+COL

combination

COL is an increasingly important antibiotic against serious infections caused by G$^-$ bacteria. It damages both the

OM and IM layers of the cell surface by targeting LPS, displacing cations that form bridges between LPS

molecules, and thereby leading to disruption of the cell envelope and bacterial lysis (Sabnis *et al*, 2021). Our

data showed that the AEC combination could permit ingress of NPN fluorophore into the OM, as well as the

membrane impermeant dye PI into the IM, which fluoresces upon contact with DNA in the bacterial cytoplasm

(Figure 2a-d). Thus, it indicated AEC combination has punched holes in both the OM and IM of whole bacterial

cells. The considerably deformed of cell membranes observed by SEM further supported the above results that

the membrane integrity, and permeability were damaged after AEC incubation. Considering that EDTA is used

as a complexing agent, we tried to explore if it could assist COL to destroy the bacterial membrane and reduce

their survival by chelating cations that stabilize LPS and the outer membrane. Unfortunately, we found minimal

change in MICs of different combinations to S16 and S30 strains, after different cations (Na^+, K^+, Ca^{2+}, Mg^{2+},

Mn^{2+}, Zn^{2+}) were supplemented (Table supplement 2). In contrast, LPS treatment resulted in dose-dependent

changes in MICs of different combinations to S16 and S30 strains (Table supplement 3), which indicated that

LPS on cell membrane was a crucial target for AEC to injure cell membrane and exert prominent antibacterial

effects.

Whilst it is commonly believed that COL acts against G$^-$ bacteria by cell membrane lysis, we hypothesised

that alterations in bacterial membrane lipid composition may also be a possible Achilles' heel to increase

efficacy of COL after the AEC incubation in this paper. The metabolomic results of AEC vs. C group in this

study showed a significant decrease of prostaglandins (PGs) in arachdonic acid metabolism pathway (Figure 6).

Arachidonic acid is a highly abundant long chain polyunsaturated fatty acid in vertebrates, which have been

proposed to have antibacterial roles. Exogenous Arachidonic acid has been reported to readily incorporated into

the synthesis pathways of membrane phospholipids, and exert detrimental effects on membrane integrity by

perturbing membrane ordering, altering membrane composition and increasing fluidity (Eijkelkamp *et al*, 2018;

MacDermott-Opeskin *et al*, 2022). PGs are lipid compounds derived from arachidonic acid, which has been

demonstrate to enhance biofilm development and fungal load in the murine vaginae of *Candida albicans* (Ells *et*

al, 2011). Therefore, we speculated that the down-regulated lipid compounds PGs may lead to perturbtion of

membrane phospholipids in cell membranes and reduce microbial viability.

Accumulation of toxic compounds may also be held responsible for the membrane damage. We noted that

there were significant accumulation of styrene in both S16 and S30 strains (Figure 6). Styrene is naturally

present as a minor metabolite, that can be synthesized at low levels by several microorganisms, like *Pencillium*

camemberti and members of the *Styracaceae* family (Yeh *et al*, 2022). However, styrene itself is toxic to most

cell types, and its hydrophobic molecules could readily partition into bacterial membrane, resulting in

membrane disruption and cell death (Lian *et al*, 2016). Additionally, we found that the taurocholic acid and

protocatechuic acid were up-regulated respectively in S16 and S30 strains (Figure 6). Taurocholic acid is

usually a major component of the selective culture medium, for example MacConkey agar, for G$^-$ bacteria,

which has also been confirmed as a secondary metabolite of marine isolates and the soil bacterium

Streptococcus faecium. Sannasiddappa *et al* have proved that taurocholic acid were able to inhibit growth of

Staphylococcus aureus through increasing membrane permeability and disruption of the PMF (Sannasiddappa

et al, 2017). Protocatechuic acid has been demonstrated to exert antimicrobial effects by disrupting the cell

membranes, preventing bacterial adhesion and biofilm formation (Bernal-Mercado *et al*, 2018; Stojković *et al*,

2013). Consequently, these toxic compounds were anticipated to accelerate the destruction of cell membrane

and thus enhance the antibacterial activity.

The CheA of chemosensory system and virulence-related protein SpvD were critical for the

bacteriostatic synergistic effect of AEC combination

Flagellar motility is intimately connected to chemotaxis, biofilm formation, colonisation and virulence of many bacterial pathogens. It is generally regulated by a chemotactic signaling system, which enables their movement toward favorable conditions and invade their hosts (Bolton, 2015). The chemotaxis proteins CheA, CheW, CheY, methyl-accepting chemotaxis proteins (MCPs) have already been identified as core components and present in all chemotaxis systems (Minamino *et al*, 2022). The *Salmonella* flagellum is composed of about 30 different proteins, such as FliD (the filament cap), FlgK and FlgL (the hook-filament junction) *et al* (Minamino *et al*, 2022). Tellingly, we have found an extensive down-regulation of chemotaxis and flagellar assembly related genes in S16 and S30 strains after AEC incubation (Figure 5a-d). The overexpression of *cheA* in S16 strain, but not *cheY*, *STMDT2-34621*, *aer*, *fliD*, *fliT*, could lead to noticeable increases of MICs by 4 to 32 fold, after EC or AEC incubation (Table supplement 4). These results suggested that the down regulation of the central component of chemosensory system CheA may affect both chemotactic motility and general structure of flagellum, thus attenuating *Salmonella* survival after AEC treatment.

In addition to the above findings, we also noted that there were relatively large number of SDEGs, most of them were down-regulated, enriched in the ABC transports pathway in S16 and S30 strains after AEC incubation (Figure 5e, f). ABC transporters are a class of transmembrane transporters, which mediate uptake of micronutrients, including saccharides, amino acids, metal ions *et al* and have also been shown to protect bacteria from hazardous compounds (Nguyen & Götz, 2016; Wang *et al*, 2023). The *opuBB* and *opuBA* encode components of the ABC-type proline/glycine betaine transport system, and their up-regulation were proved to promote accumulation of proline that acts as an osmoprotectant (Dupre *et al*, 2019). Similarly, the OsmU osmoprotectant systems, consisting of OsmV, OsmW, OsmY, and OsmX, were identified to enable bacterial

survival at high-osmolarity through the accumulation of glycine betaine (Frossard *et al*, 2012). The *gltI* gene

encodes glutamate/aspartate transport protein, and its deletion in *E. coli* were shown to result in attenuated

survival under antibiotics, acid, and hyperosmotic stressors (Niu *et al*, 2023). Furthermore, the dipeptide

permease operon (*dpp*), especially *dppA*, has been reported as an essential enzyme for survival of *Mycobacteria*

tuberculosis under nutrient starvation conditions, and was associated with reduced bacterial burden in

chronically infected mice in knockout studies (Fernando *et al*, 2022). Unfortunately, the overexpression of

opuBB, *gltI*, *dppB* and *dppC* genes caused no detectable change in MICs for S16 strain after AEC incubation

(Table supplement 4). Thus, these observations suggest that overexpression of these SDEGs in ABC

transporters were not sufficient enough for causing changes in COL susceptibility, and additional factors may

be required for the excellent bactericidal activity of AEC.

Besides that, we also noticed that the expression levels of *Salmonella* infection related genes *sseJ*, *spvD*

and ribosomal protein genes *rpmJ*, *rpmE* were all strikingly changed in S16 strain (Figure supplement 4a). SseL

and SpvD are effectors of *Salmonella* pathogenicity islands 1 and 2 (SPI1 and SPI2), which are required for full

virulence during animal infections (Coombes *et al*, 2007; Grabe *et al*, 2016). The *rpmJ* gene codified a part of

50S ribosomal subunit, and its up-regulation has been related to size increase and slow growth of *E. coli* cells

under starvation conditions (Peredo-Lovillo *et al*, 2019). The overexpression of *spvD* gene in S16 strain could

attenuate the antibacterial activity of AC and AEC regimen with 4 fold increase in MICs (Table supplement 4),

which indicated that the virulence-related protein SpvD contributed to the increase of COL susceptibility after

AEC incubation.

Artesunate can be considered as a potential MCR-1 inhibitor that enhances the efficacy of colistin

During our research process, several phenomenon have caught our attention, that the synergistic effect of AEC

combination was irrelevant to whether the *mcr-1* gene exists or not, and the *mcr-1⁻* strain S16 exhibited more

robust changes than that of *mcr-1⁺* strain S30 in different KEGG pathways. These results indicated that AS and

EDTA were possible to exert synergistic effects by blocking the broad-spectrum resistance mechanisms (eg.

efflux pumps, membrane damage), and coupling with the drug-specific resistance mechanisms (eg. MCR-1,

β-lactamase). Previously, AS was capable of significantly enhancing the antibacterial activity of β-lactam

antibiotics against *E. coli*, via inhibition of the efflux pumps such as AcrB, NorA, NorB, and NorC (Jiang *et al*,

2013; Li *et al*, 2011). Molecular docking experiments showed that AS could dock into AcrB very well forming

five hydrogen bonds with Ser46, Gln89 and Gln176 (Wu *et al*, 2013). Whereas, in this paper, the inhibitory

actions of different medication regimens on efflux pump were only observed in *mcr-1⁻* strain S16, regardless of

whether AS was added (Figure supplement 4c). Therefore, we hypothesised that AS may exert synergistic

effects with COL against *mcr-1⁺* S30 strain by targeting MCR-1 rather than efflux pump.

MCR-1 comprises two distinct domains, an N-terminal transmembrane domain and a soluble C-terminal

α/β/α sandwich domain where the active site located. The active site contains a concentration of metal-binding

residues to accommodate between one and four zinc ions (Wei *et al*, 2018). Several crystal structures of the

soluble domain have been well studied, and six residues GLU246, THR285, HIS395, ASP465, HIS466, and

HIS478 were found conserved among pEtN transferases (Ma *et al*, 2016). Especially the THR285 residue,

which was highly conserved and providing a distinct electronegative potential to attract and bind the substrate

pEtN (Son *et al*, 2019). Mutations of these residues and stripping the metals by EDTA could re-establish

polymyxin B antibacterial action (Hinchliffe *et al*, 2017; Hu *et al*, 2016; Stojanoski *et al*, 2016). In this paper,

we firstly analyzed the relative expression of *mcr-1* in S30 strain after the incubation of different medication

regimens, and found a striking down-regulation of *mcr-1* gene whether after AC, EC incubation or AEC

incubation, when compered to that of A and E treatment (Figure supplement 4b). Additionally, we performed

the molecular docking between AS and MCR-1 to predict if there were possible interactions, and found that AS

could bind to the residues arrounding the reported key residues within MCR-1 of *E. coli*, forming seven

hydrogen bonds with THR283, SER284, TYR287, PRO481, and ASN482 residues (Figure supplement 4e).

The interaction between AS and MCR-1 was further proved by competitive inhibitory assays, that the binding

of AS with MCR-1 was blocked after the addition of polypeptide P_u (containing unmutated THR283, SER284,

and TYR287 sites) and lead to significant increases of MICs by 8 fold after AC or AEC treatment (Table

supplement 5). Nevertheless, we also noticed that the blocking effect could not be removed by peptide P_m

(containing mutated THR283, SER284, and TYR287 sites) when EDTA exists. We supposed that EDTA may

chelate zinc ions that required for MCR-1 activity. Thus, we suggested that AS could be developed as the a

MCR-1 inhibitor, and coupled with the chelating agent EDTA may favor to additionally magnify its inhibitory

effect.

In summary, our results established that the combination of COL with AS and EDTA was a promising

candidate for combating infections caused by MCR-negative and -positive COL-resistant *Salmonella*. The

membrane-damaging effect, accumulation of toxic compounds, inhibition of MCR-1 were supposed to play

synergistic roles in reversing COL resistance of *Salmonella* (Figure 8). The CheA of chemosensory system and

virulence-related protein SpvD were critical for the bacteriostatic synergistic effects of AEC combination.

Selectively targeting CheA, SpvD or MCR using the natural compound AS could be further investigated as an

attractive strategy for treatment of *Salmonella* infection.

Materials and Methods

Bacterial strains and agents

A total of 9 bacteria strains were used in this study (Table 1), including a multidrug-susceptible standard strain

of *Salmonella* Typhimurium CVCC541 (named as JS), a COL-susceptible clinical strain of *Salmonella* (named

as S34), four COL-resistant clinical strains of *Salmonella* (named as S16, S20, S13, and S30), a COL-resistant

clinical strain of *Escherichia coli* (named as E16), and two strains of intrinsically COL-resistant species

(*Morganella morganii* strain M15 and *Proteus mirabilis* strain P01). COL was purchased from Shengxue

Dacheng Pharmaceutical, China. COL and EDTA-2Na were both dispersed into water at final concentration of

640 mg/L and 10000 mg/L, respectively. AS was purchased from Meilunbio (Dalian, China) and dissolved in

water with 10% (V/V) N, N-dimethylformamide at a final concentration of 5000 mg/L. The

1-N-phenylnaphthylamine (NPN) was purchased from Sigma-Aldrich, USA. Propidium iodide (PI) was

purchased from Thermo Fisher Scientific, USA. The 3,3-dipropylthiadicarbocyanine iodide $DiSC_3(5)$ and

ethidium bromide (EtBr) were purchased from Aladdin, China. BCECF-AM, Reactive Oxygen Species Assay

Kit, and Hydrogen Peroxide Assay Kit were purchased from Beyotime, China.

Antibacterial activity *in vitro*

1. Antimicrobial susceptibility testing. The MICs of COL, AS, and EDTA against all strains were determined

by the 2-fold serial broth microdilution method according to CLSI guidelines (Wayne, 2021) The double and

triple combination strategies were carried out as follow: AS (1/4, 1/8, or 1/16 MIC of AS)+ COL, EDTA (1/4,

1/8, or 1/16 MIC of EDTA) + COL, AS (1/4 MIC of AS)+ EDTA (1/4, 1/8, 1/16 or 1/32 MIC of EDTA) + COL,

AS (1/8 MIC of AS)+ EDTA (1/4, 1/8, 1/16 or 1/32 MIC of EDTA) + COL. These medication strategies,

including COL alone, AS + COL, EDTA + COL, AS + EDTA + COL, were abbreviated as C, AC, EC, and AEC,

respectively. When determining the MICs of COL after drug combinations, AS or/and EDTA were pre-added

into MHB broth with different final concentrations, namely 1/4, 1/8, 1/16 or 1/32 MIC of AS or EDTA, then

COL was added and diluted to make a 2-fold dilution series. The lowest concentrations with no visible growth

of bacteria were defined as MIC values of COL after drug combinations.

2. Time-kill assays. Time-kill assays were performed against the *Salmonella* strains JS, S16 (*mcr-1*⁻), and S30

(*mcr-1*⁺) with COL alone as well as in combinations (AC, EC, AEC). When combined with COL (0.1 or 2 mg/L),

AS and EDTA were added at final concentrations equivalent to their 1/8 MICs. Overnight cultures were diluted

1:100 in fresh LB medium and grown to an OD_{600} of 0.5, then treated with different combinations for 12 h. The

cultures were serially diluted 10-fold and spreaded over sterile nutrient agar at 0.5, 4, 8 and 12 h. Bacterial

colonies on individual plates were counted after overnight incubationat 37 ℃ and expressed as the $\log_{10}$ of

colony forming units/mL (CFU/mL).

Fluorescent probe-permeability assays

Overnight cultures of *Salmonella* strains S16 (*mcr-1*⁻) and S30 (*mcr-1*⁺) were diluted 1:100 in fresh LB medium

and grown to an OD_{600} of 0.7. Cells were harvested and washed twice with PBS or HEPES, then re-suspended in

the same buffer to $OD_{600} \approx 0.5$ for further analysis. different drug combinations or Fluorescent probes were

added and incubated when necessary. The medication strategies used in the fluorescence probe assays were as

follow: C, AC, EC, and AEC. The final concentration of COL was 0.1 or 2 mg/L, when used alone or in drug

combinations. AS and EDTA were added at final concentrations equivalent to their 1/8 MICs, when used in drug

combinations. Fluorescence intensity were measured with Spark 10M microplate spectrophotometer (Tecan,

Switzerland).

1. Cell membrane integrity assay. Bacterial suspensions in HEPES were mixed with either fluorescent probe

1-N-phenylnaphthylamine (NPN) or propidium iodide (PI) to a final probe concentrations of 10 μM for NPN or

15 μM for PI. After incubation at 37°C for 0.5 h, bacterial suspensions were then mixed with different drug

combinations and incubated for another 1 h. Fluorescence measurements were then taken with the excitation

wavelength at 350 nm (or 535 nm) and emission wavelength at 420 nm (or 615 nm) for NPN (or PI).

2. Proton motive force assay. Bacterial suspensions in PBS were incubated with either

3,3-dipropylthiadicarbocyanine iodide (DiSC$_3$(5), 0.5 μM) or pH-sensitive fluorescent probe BCECF-AM (20

μM) for 0.5 h to determine the membrane potential ($\Delta\psi$) and pH gradient (ΔpH). Then these suspensions were

mixed with different drug combinations and incubated for another 1 h. Finally, the fluorescence were measured

with the excitation wavelength of 622 nm (or 488 nm) and emission wavelength of 670 nm (or 535 nm) for

DiSC$_3$(5) (or BCECF-AM).

3. Total ROS measurement

The ROS-sensitive fluorescence indicator 2', 7'-dichlorodihydro-fluorescein diacetate (DCFH-DA, 10 μM) was

used to assess the ROS levels in bacterial cells. Bacterial suspensions in PBS were mixed with DCFH-DA and

incubated for 0.5 h, then different drug combinations were added and incubated for another 1 h and 6 h. Finally,

fluorescence intensity were measured at an excitation wavele of 488 nm and an emission wavelength of 525 nm.

4. Efflux pump assay. Bacterial suspensions in PBS were incubated with ethidium bromide (EtBr) for 0.5 h, in

a final concentration of 5 μM. Then different medication regimens were added and incubated for another 1 h,

then the accumulation of EtBr in the cells were evaluated with excitation wavelength of 530 nm and barrier filter

of 600 nm.

H₂O₂ assay

The bacterial culture conditions and sample preparation were same as the fluorescent probe-permeability assays

section. The luminescence absorbancy were measured by a Spapk 10 M Microplate reader (Tecan, Switzerland).

The cellular H_2O_2 levels were assessed according to the kit procedure by using an Hydrogen Peroxide Assay Kit.

Cells samples were prepared as described above and incubated with different drug combinations for 1.5 h. Cell

precipitates were collected by centrifugation (10000 rpm), and 200 μL lysis solution were added under gentle

shaking. Supernatants were then taken for luminescence (Abosorbance) detection at 560 nm.

Scanning electron microscope (SME)

Overnight S16 (*mcr-1⁻*) culture was diluted 1:100 in fresh LB medium and grown to an OD_{600} of 0.7. Cells were

then divided equally and different drug combinations were added, same as that in the fluorescent

probe-permeability assays section. After 6 h incubation at 37 ℃, cells were washed three times with PBS and

fixed with 2.5% glutaraldehyde at 4 ℃ for 24 h. Samples were then stained in 1% osmium tetroxide and

dehydrated in a series of increasing ethanol concentrations (30% – 100%). The processed samples were dried in

Critical Point Dryer (Quorum K850) and sputter-coated with gold. Finally, samples were observed and images

were taken with SEM (HITACHI, SU8100).

Omics analysis

Overnight cultures of S16 (*mcr-1⁻*) and S30 (*mcr-1⁺*) were diluted 1:100 in fresh LB medium and grown to an

OD_{600} of 0.5. Cells were divided equally and different medication regimens were used, including C, AC, EC,

and AEC. When combined with COL (2 mg/L), AS and EDTA were added at final concentrations equivalent to

their 1/8 MICs. Samples were continually incubated 6 h, then washed three times with PBS and fast-frozen in

liquid nitrogen for further use. Transcriptome and metabolome analysis were performed among different

comparison groups, including AC vs. C, EC vs. C, AEC vs. AC, and AEC vs. EC, and carried out by Novogene

Co. Ltd (Beijing, China).

1. Transcriptome analysis

The clean reads were mapped to the *Salmonella* Typhimurium DT2 genome (HG326213.1) from NCBI using

Bowtie2. The SDEGs were screened with a p value ≤ 0.05 and |log$_2$Fold Change| ≥ 1. ClusterProfiler software

were used to analysis the Gene Ontology functional enrichment (GO, https://geneontology.org/) or Kyoto

Encyclopedia of Genes and Genomes pathway enrichment (KEGG, https://www.kegg.jp/kegg/pathway.html) of

SDEGs.

2. Metabolome analysis

In this study, Samples were analyzed by the non-targeted metabolomics with Liquid Chromatography-tandem

Mass Spectrometry (LC-MS/MS) in either positive ion or negative ion mode. Compound Discoverer 3.1

software (Thermo Scientific, Waltham, MA, USA) was used for identification of metabolites based on the exact

masses and fragmentation spectra. The significant differential metabolites (SDMs) were identified with VIP ≥

1.0, Fold Change ≥ 1.2 or Fold Change ≤ 0.833, p value < 0.05. SDMs were then annotated and classified by

KEGG Database (https://www.genome.jp/kegg/pathway.html), Human Metabolome Database (HMDB,

https://hmdb.ca/metabolites), and Lipidmaps Database (https://www.lipidmaps.org/).

Antibacterial activity *in vivo*

Bacterial preparation: Overnight culture of S30 (*mcr-1*[+]) was diluted 1:100 in fresh LB medium and grown to

an OD_{600} of 0.7. Cells were harvested and washed twice with PBS, then adjusting the concentration of bacterial

suspensions to 1.31×10^6 CFU/mL for further use.

Animals and treatments: A total of 36 SPF Kunming mice (Six to eight-week-old, 18-22 g, half male and half

female) were purchased from the Huaxing Experimental Animal Center of Zhengzhou (Zhengzhou, China) and

divided into six groups (n = 6 per group): (1) PBS control group; (2) COL (10 mg/kg) group; (3) AS (15 mg/kg)

group; (4) AS (15 mg/kg) + COL (10 mg/kg) group; (5) EDTA (50 mg/kg) + COL (10 mg/kg) group; (6) AS (15

mg/kg) + EDTA (50 mg/kg)+ COL (10 mg/kg) group. Each mice was intraperitoneally injected with 100 μL

bacterial solution (1.31×10^5 CFU). Treatment was initiated at 2 h post infection and continued for 3 days.

Treatment was administered once per day by intraperitoneal injection according to the therapeutic dose

mentioned above. Mices were all euthanized and spleen, liver of aseptic were collected, weighed, homogenized

with PBS. Then the tissue homogenates were serially diluted with PBS in an appropriate amount and 100 μL of

each dilution was withdrawn and uniformly spread on SS agar plates. Bacterial counts were computed and

presented as the mean ± SD $\log_{10}$ CFU/mL, after incubated at 37°C for 16 to 18 h. Mice were maintained in a

barrier facility and guaranteed strict compliance with the regulations for the Administration of Affairs

Concerning Experimental Animals approved by the State Council of People's Republic of China (11–14–

1988).The mouse experiments were approved by the Henan Science and Technology Department (protocol

number SCXK 2019-0002).

Data available

Transcriptome data have been submitted to the Sequence Read Archive database (SRA,

https://www.ncbi.nlm.nih.gov/sra) under the BioProject accession number PRJNA1036120 (S16 strain) and

PRJNA1036408 (S30 strain). Metabolome data have been submitted to the MetaboLights database

(https://www.ebi.ac.uk/metabolights) under accession number MTBLS8875. It is anticipated that this accession

number will be released as soon as the manuscript is received.

Statistical analysis

Statistical analysis was conducted using Graphpad prism 9 and SPSS software. All data were derived from $n \geq 3$

biological replicates and presented as mean ± SD. Without specific indication, differences between the

independent groups (* $p < 0.001$) were assessed with student's t-test or one-way ANOVA.

Supplement materials

Figure supplement 1. Intracellular accumulation of ROS in S16 and S30 strains after 6 h treatment.

Figure supplement 2. The number of DEGs are identified in S16 and S30 strains among different

comparison groups.

Figure supplement 3. KEGG pathway analysis of SDEGs in S16 (a, b) and S30 (c, d) strains within the

AC .vs. C, and EC .vs. C groups.

Figure supplement 4. (a) The SDEGs detected in *Salmonella* infection and ribosome pathways; (b)

Expression level of *mcr-1*; (c, d) efflux pump activity in S16 or S30 strain; (e) Putative pattern of interaction

between AS and MCR-1 protein.

Table supplement 1. The antibacterial activities of COL, AS and EDTA against the tested strains after

single and double combinations

Table supplement 2. MIC values of COL against S16 and S30 strains after addition of exogenous cations

Table supplement 3. MIC values of COL against S16 and S30 strains after addition of exogenous LPS

Table supplement 4. MIC values of COL against S16 strain after the overexpression of different genes

Table supplement 5. MIC values of COL against S30 strain after the incubation of polypeptides

Table supplement 6. Sequences of primers used in this study

Supplement materials and methods

Acknowledgements

474 This work was supported by the National Natural Science Foundation of China (No. 32102716 and 32373069).

475   Ethical statement: This study was carried out in accordance with the guidelines of Henan Agricultural

476 University Animal Ethics Committee.

Author contributions

479 YZ and GH conceived the study. PL, XH, and CF performed experiments, results analysis and wrote the original

480 draft. XC and QH analyzed the results and wrote the original draft. DH and XM analyzed the results. YZ, PL,

481 and GH approved the final version of the manuscript.

Disclosure and competing interests statement

484 The authors declare that they have no conflict of interest.

Data Availability

Data will be made available freely online.

References

Aron Z and Opperman TJ (2016) Optimization of a novel series of pyranopyridine RND efflux pump inhibitors.

Curr Opin Microbiol 33: 1-6.

Bernal-Mercado AT, Vazquez-Armenta FJ, Tapia-Rodriguez MR, Islas-Osuna MA and Mata-Haro V (2018).

Comparison of single and combined use of catechin, protocatechuic, and vanillic acids as antioxidant and

antibacterial agents against uropathogenic *Escherichia Coli* at planktonic and biofilm levels. *Molecules*

23:2813.

Bolton DJ (2015). *Campylobacter* virulence and survival factors. *Food Microbiol* 48: 99-108.

Carroll LM, Gaballa A, Guldimann C, Sullivan G, Henderson LO and Wiedmann M (2019) Identification of

novel mobilized colistin resistance gene *mcr-9* in a multidrug-resistant, colistin-susceptible *Salmonella*

enterica Serotype Typhimurium isolate. *mBio* 10: e00853-19.

Coombes BK, Lowden MJ, Bishop JL, Wickham ME, Brown NF, Duong N, Osborne S, Gal-Mor O and Finlay

BB (2007) SseL is a *Salmonella*-specific translocated effector integrated into the SsrB-controlled

Salmonella pathogenicity island 2 type III secretion system. *Infect Immun* 75: 574-580.

Dupre JM, Johnson WL, Ulanov AV, Li Z, Wilkinson BJ and Gustafson JE (2019) Transcriptional profiling and

metabolomic analysis of *Staphylococcus aureus* grown on autoclaved chicken breast. *Food Microbiol* 82:

46-52.

Eijkelkamp BA, Begg SL, Pederick VG, Trapetti C, Gregory MK, Whittall JJ, Paton JC and McDevitt CA (2018)

Arachidonic acid stress impacts pneumococcal fatty acid homeostasis. *Front Microbiol* 9: 813

El-Sayed Ahmed MAE, Zhong LL and Shen C (2020) Colistin and its role in the Era of antibiotic resistance: an

extended review (2000-2019). *Emerg Microbes Infect* 9: 868-885.

Ells R, Kock JL, Albertyn J, Kemp G and Pohl CH (2011) Effect of inhibitors of arachidonic acid metabolism on

prostaglandin E_2 production by *Candida albicans* and *Candida dubliniensis* biofilms. *Med Microbiol*

Immunol 200: 23-28.

Falagas ME and Kasiakou SK (2005) Colistin: the revival of polymyxins for the management of

multidrug-resistant gram-negative bacterial infections. *Clin Infect Dis* 40: 1333-1341.

Farha MA, Verschoor CP, Bowdish D and Brown ED (2013) Collapsing the proton motive force to identify

synergistic combinations against *Staphylococcus aureus*. *Chem Biol* 20: 1168-1178.

Fernando DM, Gee CT, Griffith EC, Meyer CJ, Wilt LA, Tangallapally R, Wallace MJ, Miller DJ and Lee RE

(2022) Biophysical analysis of the *Mycobacteria tuberculosis* peptide binding protein DppA reveals a

stringent peptide binding pocket. *Tuberculosis (Edinb)* 132: 102157.

Frossard SM, Khan AA, Warrick EC, Gately JM, Hanson AD, Oldham ML, Sanders DA and Csonka LN (2012)

Identification of a third osmoprotectant transport system, the OsmU system, in *Salmonella enterica*. *J*

Bacteriol 194: 3861-3871.

Grabe GJ, Zhang Y, Przydacz M, Rolhion N, Yang Y, Pruneda JN, Komander D, Holden DW and Hare SA (2016)

The *Salmonella* effector SpvD is a cysteine hydrolase with a serovar-specific polymorphism influencing

catalytic activity, suppression of immune responses, and bacterial virulence. *J Biol Chem* 291: 25853-25863.

Hinchliffe P, Yang QE, Portal E, Young T, Li H, Tooke CL, Carvalho MJ, Paterson NG, Brem J, Niumsup PR *et*

al (2017) Insights into the mechanistic basis of plasmid-mediated colistin resistance from crystal structures

of the catalytic domain of MCR-1. *Sci Rep* 7: 39392.

Hu M, Guo J, Cheng Q, Yang Z, Chan EWC, Chen S and Hao Q (2016) Crystal structure of *Escherichia coli*

originated MCR-1, a phosphoethanolamine transferase for colistin resistance. *Sci Rep* 6: 38793.

Jiang W, Li B, Zheng X, Liu X, Pan X, Qing R, Cen Y, Zheng J and Zhou H (2013) Artesunate has its

enhancement on antibacterial activity of β-lactams via increasing the antibiotic accumulation within

methicillin-resistant *Staphylococcus aureus* (MRSA). *J Antibiot (Tokyo)* 66: 339-345.

Kaye KS, Pogue JM, Tran TB, Nation RL and Li J (2016) Agents of last resort: polymyxin resistance. *Infect Dis*

Clin North Am 30: 391-414.

Laws M, Shaaban A and Rahman KM (2019) Antibiotic resistance breakers: current approaches and future

directions. *FEMS Microbiol Rev* 43: 490-516.

Le D, Krasnopeeva E, Sinjab F, Pilizota T and Kim M (2021) Active efflux leads to heterogeneous dissipation of

proton motive force by protonophores in bacteria. *mBio* 12: e0067621.

Li B, Yao Q, Pan XC, Wang N, Zhang R, Li J, Ding G, Liu X, Wu C, Ran D *et al* (2011) Artesunate enhances the

antibacterial effect of β-lactam antibiotics against *Escherichia coli* by increasing antibiotic accumulation via

inhibition of the multidrug efflux pump system AcrAB-TolC. *J Antimicrob Chemother* 66: 769-777.

Lian J, McKenna R, Rover MR, Nielsen DR, Wen Z and Jarboe LR (2016) Production of biorenewable styrene:

utilization of biomass-derived sugars and insights into toxicity. *J Ind Microbiol Biotechnol* 43: 595-604.

Lima WG, Alves MC, Cruz WS and Paiva MC (2018) Chromosomally encoded and plasmid-mediated

polymyxins resistance in *Acinetobacter baumannii*: a huge public health threat. *Eur J Clin Microbiol Infect*

Dis 37: 1009-1019.

Lin L, Nonejuie P, Munguia J, Hollands A, Olson J, Dam Q, Kumaraswamy M, Rivera H, Jr., Corriden R, Rohde

M *et al* (2015) Azithromycin synergizes with cationic antimicrobial peptides to exert bactericidal and

therapeutic activity against highly multidrug-resistant Gram-Negative bacterial pathogens. *EBioMedicine* 2:

690-698.

Ma G, Zhu Y, Yu Z, Ahmad A and Zhang H (2016) High resolution crystal structure of the catalytic domain of

MCR-1. *Sci Rep* 6: 39540.

MacDermott-Opeskin HI, Panizza A, Eijkelkamp BA and O'Mara ML (2022) Dynamics of the *Acinetobacter*

baumannii inner membrane under exogenous polyunsaturated fatty acid stress. *Biochim Biophys Acta*

Biomembr 1864: 183908.

Minamino T, Kinoshita M and Namba K (2022) Insight into distinct functional roles of the flagellar ATPase

complex for flagellar assembly in *Salmonella*. *Front Microbiol* 13: 864178.

Moya B, Barcelo IM, Bhagwat S, Patel M, Bou G, Papp-Wallace KM, Bonomo RA and Oliver A (2017) WCK

5107 (Zidebactam) and WCK 5153 are novel inhibitors of PBP2 showing potent "β-Lactam Enhancer"

activity against *Pseudomonas aeruginosa*, including multidrug-resistant metallo-β-lactamase-producing

high-risk clones. *Antimicrob Agents Chemother* 61: e02529-16.

Nguyen MT and Götz F (2016) Lipoproteins of Gram-Positive bacteria: key players in the immune response and

virulence. *Microbiol Mol Biol Rev* 80: 891-903.

Niu H, Li T, Du Y, Lv Z, Cao Q and Zhang Y (2023) Glutamate transporters GltS, GltP and GltI are involved in

Escherichia coli tolerance in vitro and pathogenicity in mouse urinary tract infections. *Microorganisms* 11:

1173.

Pan X, Cen Y, Kuang M, Li B, Qin R and Zhou H (2020) Artesunate interrupts the self-transcriptional activation

of MarA to inhibit RND family pumps of *Escherichia coli*. *Int J Med Microbiol* 310: 151465.

Peredo-Lovillo A, Dorantes-Alvarez L, Hernández-Sánchez H, Ribas-Aparicio R, Cauich-Sánchez P and

Aparicio-Ozores G (2019) Functional properties and microstructure of *Leuconostoc citreum* and

Lactobacillus casei shirota exposed to habanero pepper extract and inhibition of *Staphylococcus aureus*.

Revista Mexicana de Ingeniería Química 18: 115-129.

Poirel L, Jayol A and Nordmann P (2017) Polymyxins: Antibacterial activity, susceptibility testing, and

resistance mechanisms encoded by plasmids or chromosomes. *Clin Microbiol Rev* 30: 557-596.

Rosenthal PJ (2003) Antimalarial drug discovery: old and new approaches. *J Exp Biol* 206: 3735-3744

Sabnis A, Hagart KL, Klöckner A, Becce M, Evans LE and Furniss RCD (2021) Colistin kills bacteria by

targeting lipopolysaccharide in the cytoplasmic membrane. *eLife* 10: e65836.

Sannasiddappa TH, Lund PA and Clarke SR (2017) *In vitro* antibacterial activity of unconjugated and

conjugated bile salts on *Staphylococcus aureus*. *Front Microbiol* 8: 1581.

Son SJ, Huang R, Squire CJ and Leung IKH (2019) MCR-1: a promising target for structure-based design of

inhibitors to tackle polymyxin resistance. *Drug Discov Today* 24: 206-216.

Stogios PJ, Spanogiannopoulos P, Evdokimova E, Egorova O, Shakya T, Todorovic N, Capretta A, Wright GD

and Savchenko A (2013) Structure-guided optimization of protein kinase inhibitors reverses aminoglycoside

antibiotic resistance. *Biochem J* 454: 191-200.

Stojanoski V, Sankaran B, Prasad BV, Poirel L, Nordmann P and Palzkill T (2016) Structure of the catalytic

domain of the colistin resistance enzyme MCR-1. *BMC biology* 14: 81.

Stojković DS, Zivković J, Soković M, Glamočlija J, Ferreira IC, Janković T and Maksimović Z (2013)

Antibacterial activity of *Veronica montana* L. extract and of protocatechuic acid incorporated in a food

system. *Food Chem Toxicol* 55: 209-213.

Su F and Wang J (2018) Berberine inhibits the MexXY-OprM efflux pump to reverse imipenem resistance in a

clinical carbapenem-resistant *Pseudomonas aeruginosa* isolate in a planktonic state. *Exp Ther Med* 15:

467-472.

Wang J, Sheng Z, Liu Y, Chen X, Wang S and Yang H (2023) Combined proteomic and transcriptomic analysis

of the antimicrobial mechanism of tannic acid against *Staphylococcus aureus*. *Front Pharmacol* 14:

1178177.

Wayne, PA (2021) Clinical and Laboratory Standards Institute. Performance standards for antimicrobial

susceptibility testing: 31st ed. CLSI Supplement M100.

Wei P, Song G, Shi M, Zhou Y, Liu Y, Lei J, Chen P and Yin L (2018) Substrate analog interaction with MCR-1

offers insight into the rising threat of the plasmid-mediated transferable colistin resistance. *Faseb j* 32:

1085-1098.

Wei S, Yang Y, Tian W, Liu M, Yin S and Li J (2020) Synergistic activity of fluoroquinolones combining with

artesunate against multidrug-resistant *Escherichia coli*. *Microb Drug Resist* 26: 81-88.

Wu C, Liu J, Pan X, Xian W, Li B, Peng W, Wang J, Yang D and Zhou H (2013) Design, synthesis and

evaluation of the antibacterial enhancement activities of amino dihydroartemisinin derivatives. *Molecules*

18: 6866-6882.

Yeh V, Goode A, Johnson D, Cowieson N and Bonev BB (2022) The role of lipid chains as determinants of

membrane stability in the presence of styrene. *Langmuir* 38: 1348-1359.

Yi K, Liu S, Liu P, Luo X, Zhao J, Yan F, Pan Y, Liu J, Zhai Y, Hu G (2022) Synergistic antibacterial activity

of tetrandrine combined with colistin against MCR-mediated colistin-resistant *Salmonella*. Biomed

Pharmacother 149: 112873.

Zhou Y, Liu B, Chu X, Su J, Xu L, Li L, Deng X and Li D (2022) Commercialized artemisinin derivatives

combined with colistin protect against critical Gram-negative bacterial infection. *Commun Biol* 5: 931.

Tables

Table 1. The antibacterial activities of COL against the tested strains after single and triple combinations

Strains	COL alone	COL + 1/4 AS +EDTA				COL + 1/8 AS +EDTA				Fold change
		1/4 EDTA	1/8 EDTA	1/16 EDTA	1/32 EDTA	1/4 EDTA	1/8 EDTA	1/16 EDTA	1/23 EDTA	
JS	0.25	0.015	0.125	0.125	0.125	0.03	0.25	0.25	0.25	**0-17**
S34	0.125	0.008	0.015	0.015	0.015	0.02	0.0625	0.125	0.125	**0-16**
S16	4	0.015	0.03	0.0625	0.0625	0.0625	0.125	0.125	0.5	**8-267**
S20	2	0.0625	0.125	0.125	0.125	0.25	0.5	0.5	0.5	**4-32**
S13	2	0.015	0.125	0.25	0.25	0.0625	0.25	0.5	0.5	**4-133**
S30	4	0.015	0.0625	0.0625	0.125	0.125	0.125	0.25	0.5	**8-266.6**
E16	2	0.00003	0.002	0.25	0.5	0.004	0.015	0.25	1	**2-66667**
M15	**> 64**	24	24	**> 64**	**> 64**	24	24	**> 64**	**> 64**	**0-2**
P01	**> 64**	**> 64**	**> 64**	**> 64**	**> 64**	**> 64**	**> 64**	**> 64**	**> 64**	**0**

Note: COL: colistin; AS: artesunate

Figure Legends

Figure 1. Time-kill curves of *Salmonella* strains JS, S16, and S30 with COL alone and in combinations.

a-f   Samples were treated with different concentration of COL (0.1 or 2 mg/L), alone or in drug combinations,

for 12 h. When used in combination, 1/8 MIC of AS or EDTA was added to a final concentration of 156.3

or 15.6 mg/L, respectively. Counts of CFU/mL were performed on all cultures at each time point, and data

are mean ± SD from representative of three independent experiments. CON indicates the negative control

group.

Figure 2. AS, EDTA, and COL affected membrane integrity, PMF, ROS, and H_2O_2 levels in S16 and S30

strains..

Different concentration of COL (0.1 or 2 mg/L) were used alone or in combination with AS and EDTA. When

used in combination, 1/8 MIC of AS or EDTA was added to a final concentration of 156.3 or 15.6 mg/L,

respectively.

a-d   AS, EDTA, and COL affected membrane integrity as measured by fluorescence probes NPN and PI. Error

bars indicated standard deviations for 3 replicas (* $p < 0.001$, ns not significant). CON indicates the

negative control group, and SOV indicates the solvent-exposed group.

e-h   Disruption of PMF is shown by measuring the dissipation of electric potential (Δψ) (a, b) and osmotic

component (ΔpH) (c, d). Error bars indicate standard deviations for 3 replicas (* $p < 0.001$, ns not

significant). CON indicates the negative control group, and SOV indicates the solvent-exposed group.

i-l   Intracellular accumulation of ROS (i, j) and H_2O_2 (k, l) in S16 and S30 strains after 1 h treatment. Data were

shown in the mean of triplicates $\pm$ SD (* $p < 0.001$, ns not significant). CON indicates the negative control

group, SOV indicates the solvent-exposed group, and POS indicates the positive control group.

Figure 3. Morphological changes of S16 (*mcr-1*⁻) strain.

The images were obtained after the treatment with COL (2 mg/L) alone or in combination with 1/8 MIC of AS

(156.3 mg/L) or EDTA (15.6 mg/L). CON indicates the negative control group, and SOV indicates the

solvent-exposed group. Red arrows indicate the cell damages characterized by folds, crevices, or depressions.

Figure 4. KEGG pathway analysis of SDEGs in S16 (a, b, c) and S30 (d, e, f) strains within the AEC .vs. C,

AEC .vs. AC, and AEC .vs. EC groups.

Samples were harvested after the trestment of COL (2 mg/L) alone or in combination with 1/8 MIC of AS

(156.3 mg/L) or EDTA (15.6 mg/L) for 6 h. Pathway name and number of down-regulated (blue),

up-regulated (red) genes in each pathway are indicated in parentheses on the left (down/up). Highlighted

with red circles are the pathways that SDEGs mainly enriched and appeared simultaneously in different

comparison groups.

Figure 5. The SDEGs detected in two component system (a, b), flagellar assembly (c, d) and ABC

transporters (e, f) pathways among different comparison groups, within S16 and S30 strains.

Samples were harvested after the treatment of COL (2 mg/L) alone or in combination with 1/8 MIC of AS

(156.3 mg/L) or EDTA (15.6 mg/L) for 6 h. Labels in each square indicate the $\log_2$ (fold change) of

corresponding genes. Squares without label and gray background indicate the data are not credible ($p >$

0.05, |$\log_2$Fold Change| < 1.0). Background colors indicate the expression levels of the respective genes,

red = up-regulated, blue = down-regulated. $\log_2$FC: $\log_2$Fold Change.

Figure 6. The SDMs detected in archidonic acid metabolism, degradation of aromatic compounds ,

taurine and hypotaurine metabolism pathways among different comparison groups, within S16 (a) and

S30 (b) strains.

Samples were harvested after the treatment of COL (2 mg/L) alone or in combination with 1/8 MIC of AS

(156.3 mg/L) or EDTA (15.6 mg/L) for 6 h. Labels in each square indicate the fold changes of

corresponding metabolites. Squares without label and gray background indicate the data are not credible

(VIP < 1.0, 0.833 > Fold Change < 1.2 or Fold Change ≤ 0.833, $p \geq 0.05$). Background colors indicate the

fold changes of the respective metabolites, red = increased, blue = decreased.

Figure 7. AS and EDTA potentiate colistin activity against *Salmonella* S30 (*mcr-1*[+]) *in vivo*.

Kunming mice (n=6 per group) were intraperitoneally given a non-lethal dose of *Salmonella* S30 (1.31 $\times$

10^5 CFU), then treated with PBS, COL (10 mg/kg), AS (15 mg/kg), AS (15 mg/kg) + COL (10 mg/kg),

EDTA (50 mg/kg) + COL (10 mg/kg), and AS (15 mg/kg) + EDTA (50 mg/kg)+ COL (10 mg/kg) by

intraperitoneal injection. Bacterial loads were determined in spleen and liver and bacterial counts were

computed and presented as the mean ± SD $\log_{10}$ CFU/mL. The p values were determined by one-way

ANOVA (* $p < 0.001$, ns not significant).

Figure 8. Scheme summarizing the proposed mechanisms that AS and EDTA enhancing the antibacterial

effect of COL.

①COL and EDTA cause the membrane damages by targeting LPS and chelating cations, which punch

holes in both the outer membrane (OM) and inner membrane (IM). ②AEC combination collapses the Δψ

component of Proton Motive Force (PMF) in *Salmonella*. ③The down-regulated lipid compounds

prostaglandins (PGs) lead to perturbtion of membrane phospholipids in cell membranes. ④Accumulation

of toxic compounds (eg. styrene, ROS *et al*) could accelerate the destruction of cell membrane. ⑤The

down-regulation of chemotaxis, flagellar assembly, *Salmonella* infection related genes indicate impaired

virulence of *Salmonella*. ⑥The down-regulation of genes in ABC transporters indicate impaired stress

tolerance of *Salmonella*. ⑦AEC treatment results in the down-rugulation of flagellar assembly related

genes and defective of flagellum. ⑧AS was capable of significantly enhancing the antibacterial activities

of antibiotics against *E. coli,* via inhibition of the efflux pumps. ⑨AS could be developed as the a MCR-1

inhibitor, which may work synergistically with the EDTA chelation to inhibit MCR-1 and contribute to

reverse the COL resistance of *mcr-1*-harboring *Salmonella* strains. FC: Fold Change.

Outer membrane

a S16 (*mcr-1⁻*)

b S30 (*mcr-1⁺*)

Inner permeability

c S16 (*mcr-1⁻*)

d S30 (*mcr-1⁺*)

ΔΨ

e S16 (*mcr-1⁻*)

f S30 (*mcr-1⁺*)

ΔPH

g S16 (*mcr-1⁻*)

h S30 (*mcr-1⁺*)

ROS

i S16 (*mcr-1⁻*)

j S30 (*mcr-1⁺*)

H₂O₂

k S16 (*mcr-1⁻*)

l S30 (*mcr-1⁺*)

S16 (*mcr-1⁻*)

S30 (*mcr-1⁺*)

Two-component system
Flagellar assembly
ABC transporters

a S16 (*mcr-1⁻*)

	AC.vs.C	EC.vs.C	AEC.vs.C	AEC.vs.AC	AEC.vs.EC
Arachidonic acid metabolism — Prostaglandin G$_2$ (PGG$_2$)	-0.98	-1.92	-1.71		
Prostaglandin H$_2$ (PGH$_2$)	-1.01	-2.57	-2.61		
Prostaglandin D$_2$ (PGD$_2$)	-1.22	-2.69	-2.46		
Prostaglandin J$_2$ (PGJ$_2$)	-1.15	-3.25	-3.03		
Prostaglandin E$_2$ (PGE$_2$)		-1.84	-1.92		
Prostaglandin B$_2$ (PGB$_2$)		-1.49	-1.51		
Degradation of aromatic compounds — Styrene	2.86		3.08		2.98
o-Cresol	3.34		1.92	-1.41	1.75
Taurine and hypotaurine metabolism — Taurocholic acid	5.06		3.36	-1.71	3.52

b S30 (*mcr-1⁺*)

	AC.vs.C	EC.vs.C	AEC.vs.C	AEC.vs.AC	AEC.vs.EC
Degradation of aromatic compounds — Styrene	2.56		2.74		2.74
Protocatechuic acid		1.63	1.50	1.81	

Liver

a

Spleen

b

ROS

a S16 (*mcr-1⁻*)

b S30 (*mcr-1⁺*)

Legend: CON, SOV, POS, C, AC, EC, AEC

S30 (*mcr-1*⁺)

a AC.vs.C

- Two-component system (8/43)
- ABC transporters (8/43)
- Flagellar assembly (7/16)
Fructose and mannose metabolism (3/14)
Pentose and glucuronate interconversions (2/15)
Porphyrin metabolism (0/14)
Glyoxylate and dicarboxylate metabolism (3/10)
Citrate cycle (TCA cycle) (12/1)
Oxidative phosphorylation (12/1)
Sulfur metabolism (1/12)
Propanoate metabolism (1/11)
Pyruvate metabolism (1/11)
Glycolysis / Gluconeogenesis (5/6)
Bacterial secretion system (1/9)
Nitrogen metabolism (1/9)
Pentose phosphate pathway (1/9)
Purine metabolism (3/7)
Ascorbate and aldarate metabolism (0/8)
Starch and sucrose metabolism (2/6)
Glycerophospholipid metabolism (0/7)

Down | Up

b EC.vs.C

- ABC transporters (5/48)
- Two-component system (2/41)
- Flagellar assembly (1/15)
Pentose and glucuronate interconversions (1/14)
Porphyrin metabolism (0/15)
Sulfur metabolism (0/15)
Pyruvate metabolism (1/13)
Fructose and mannose metabolism (0/12)
Propanoate metabolism (0/11)
Glyoxylate and dicarboxylate metabolism (0/9)
Nitrogen metabolism (0/9)
Pentose phosphate pathway (0/9)
Pyrimidine metabolism (3/6)
Ascorbate and aldarate metabolism (0/8)
Bacterial secretion system (0/8)
Glycerophospholipid metabolism (0/8)
Purine metabolism (1/7)
Thiamine metabolism (0/7)
Starch and sucrose metabolism (0/7)
Glycolysis / Gluconeogenesis (1/6)

Down | Up

S16 (*mcr-1*⁻)

c AC.vs.C

- Two-component system (13/16)
- ABC transporters (11/7)
- Flagellar assembly (6/5)
Oxidative phosphorylation (9/0)
Citrate cycle (TCA cycle) (8/0)
- *Salmonella* infection (6/1)
Ascorbate and aldarate metabolism (0/6)
Propanoate metabolism (1/5)
Fructose and mannose metabolism (0/6)
Pentose and glucuronate interconversions (1/5)
Bacterial secretion system (2/3)
Nitrogen metabolism (2/3)
Porphyrin metabolism (0/5)
Glyoxylate and dicarboxylate metabolism (0/4)

Down | Up

d EC.vs.C

- ABC transporters (3/3)
Pentose and glucuronate interconversions (2/2)
- Two-component system (3/1)
Fructose and mannose metabolism (3/0)
Amino sugar and nucleotide sugar metabolism (3/0)
- *Salmonella* infection (0/3)
Bacterial secretion system (1/1)
- Flagellar assembly (1/1)
Ribosome (0/2)
Phosphotransferase system (PTS) (0/1)
Fatty acid biosynthesis (0/1)
Tyrosine metabolism (1/0)
Galactose metabolism (0/1)
Glycine, serine and threonine metabolism (1/0)
Propanoate metabolism (1/0)

Down | Up