Full title:

Sex differences in change-of-mind neuroeconomic decision-making is modulated by LINC00473 in medial

prefrontal cortex

Short title:

Neuroeconomic decision-making in males and females

Authors:

Romain Durand-de Cuttoli[1], Orna Issler[2], Benjamin Yakubov[3], Nusrat Jahan[3], Aisha Abid[3], Susanna

Kasparov[3], Kastalia Granizo[3], Sanjana Ahmed[3], Scott J. Russo[1], Eric J. Nestler[1,4], Brian M. Sweis*[1,4]

*Corresponding: brian.sweis@mountsinai.org

Affiliations:

[1] Nash Family Department of Neuroscience, Friedman Brain Institute, Icahn School of Medicine at Mount

Sinai, New York, NY 10029

[2] Neuroscience Institute, Department of Anesthesiology, Perioperative Care and Pain Medicine, and

Department of Neuroscience and Physiology, NYU Langone, New York, NY 10016

[3] CUNY Hunter, New York, NY 10023

[4] Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY 10029

Abstract:

Changing one's mind is a complex cognitive phenomenon involving a continuous re-appraisal of the trade-off between past costs and future value. Recent work modeling this behavior across species has established associations between aspects of this choice process and their contributions to altered decision-making in psychopathology. Here, we investigated the actions in medial prefrontal cortex (mPFC) neurons of long intergenic non-coding RNA, LINC00473, known to induce stress resilience in a striking sex-dependent manner, but whose role in cognitive function is unknown. We characterized complex decision-making behavior in male and female mice longitudinally in our neuroeconomic foraging paradigm, Restaurant Row, following virus-mediated LINC00473 expression in mPFC neurons. On this task, mice foraged for their primary source of food among varying costs (delays) and subjective value (flavors) while on a limited time-budget during which decisions to accept and wait for rewards were separated into discrete stages of primary commitments and secondary re-evaluations. We discovered important differences in decision-making behavior between female and male mice. LINC00473 expression selectively influenced multiple features of re-evaluative choices, without affecting primary decisions, in female mice only. These behavioral effects included changing how mice (i) cached the value of the passage of time and (ii) weighed their history of economically disadvantageous choices. Both processes were uniquely linked to change-of-mind decisions and underlie the computational bases of distinct aspects of counterfactual thinking. These findings reveal a key bridge between a molecular driver of stress resilience and psychological mechanisms underlying sex-specific decision-making proclivities.

Introduction:

How an individual changes one's mind when making choices is a challenging yet important phenomenon that has been gaining more traction recently in the field of decision science [1]. This mental process involves complex interactions between cognitive and affective systems and presents an opportunity to peer into how the brain represents internal conflict [2, 3]. This captures conflict at multiple levels, including the economic level: between competing interests of managing ongoing expenditures while minimizing potential losses vs. maximizing future gains [4], as well as the psychological level: re-evaluating one's own recent actions, which could be perceived as egodystonic [5-7]. Aspects of these processes are likely dysfunctional in psychiatric illnesses and may manifest as the cognitive changes observed in mood disorders like depression, considering these mental operations could contribute to affective burden [8, 9]. While the neural mechanisms underlying change-of-mind decision-making remain underexplored, expanding this line of investigation holds promise to enrich our understanding of the repertoire of behavioral processes at play in depression.

Depression is a debilitating mental illness characterized by pervasive mood disturbances, altered cognitive capabilities, and maladaptations in motivated behavior [9]. The incidence of depression is two times higher in women than in men and can manifest different symptomatology, including negative rumination [10]; however, the molecular mechanisms mediating these sex differences are unclear [11-15]. Progress is being made toward appreciating neuroanatomical, genetic, epigenetic, and pathophysiological differences between depressed males and females based on advances in clinical and preclinical research, with the latter enabling mechanistic studies of sex as a biological factor [16, 17]. Such advances are also due to adopting novel approaches that leverage unsupervised analyses of large datasets in emerging fields of systems biology leading to breakthroughs, for example, in identifying key regulatory roles of non-coding elements of the genome [18].

Recent transcriptomics work carried out in human postmortem brain tissue samples obtained from subjects carrying a diagnosis of major depressive disorder discovered sex-dependent functions of a

primate-specific long intergenic non-coding RNA: LINC00473 [19]. Overall, long non-coding RNAs

represent a significant percentage of all of the regulated transcripts in human depression and are regulated

in a sex- and brain region-specific manner. [19]. Lower LINC00473 expression levels were found in the

medial prefrontal cortex (mPFC) of depressed females only [19]. Cross-species validation studies in rodents

demonstrated expressing LINC00473 in mPFC neurons promotes resilience to stress in females only [19].

This included ameliorating depressive- and anxiety-like behaviors while blunting hypothalamic-pituitary-

adrenal-axis activity as well as the induction of gene expression changes induced by chronic stress [19].

Other studies have implicated LINC00473 in molecular scaffolding, inter-organelle communication, and

transcriptional regulation in non-neuronal tissue, while exhibiting immediate early gene-like properties in

cultured neurons [20-25]. Beyond this, LINC00473 has not been studied in neural tissue in any other

capacity, with no work performed to date on the role of long non-coding RNAs in regulating decision-making

information processing.

Strides made in the field of neuroeconomics, an area focused on understanding how the brain is

comprised of multiple valuation algorithms, have leveraged complex approaches in decision neuroscience

to quantify how circuits constrain behavior [26-29]. This encompasses characterizing multifactorial aspects

of motivation that operationalize reward value along several dimensions, integrate choice processes with

environmental circumstances, and segregate stages of a decision stream into its component parts [30-33].

This work includes isolating properties of change-of-mind re-evaluative decisions, and points to the mPFC

as a critical hub for regulating reward value, self-control, and mood [4, 34-38]. Insights from neuroeconomic

principles offer novel approaches to investigate decision-making capable of resolving behavior into

discretely measurable computational units in a manner that is biologically tractable and readily translatable

across species [30, 39, 40]. This approach has yielded efforts from our team and others developing

innovative tasks adapted for use in mice, rats, monkeys, and humans [5, 41-44]. The use of these

translational tasks have led to cross-species discoveries uncovering the behavioral underpinnings of

signatures of higher-order cognitive processes, previously thought to be unique to humans, supported by

physically separable circuits in the brain [45-49].

Here, we characterized complex decision-making behavior in mice tested on our longitudinal

neuroeconomic foraging paradigm, Restaurant Row. This task has been utilized previously to gain access

to behavioral and neurophysiological correlates of multiple decision-making systems in the brain, including

fundamentally distinct valuation algorithms engaged within the same trial [5, 47, 50-55]. On this task,

rodents forage for their primary source of food among varying costs (delays cued by tone pitch) and

subjective value (flavors tied to spatially fixed and contextualized locations, or "restaurants") while on a

limited time-budget. Decisions to accept and wait for rewards are separated into discrete stages of primary

commitment (accept vs. reject in an offer zone) and secondary re-evaluation (stay vs. quit in a wait zone).

We tested both male and female mice – the first ever analysis of females in this task – transfected with a

newly packaged adeno-associated virus expressing LINC00473 or only a green fluorescent protein targeted

to mPFC neurons (**Fig. 1a**, **Supplementary Fig. 1**). Mice were tested daily over two months across a

changing economic landscape during which complex foraging strategies developed. We report sex- and

LINC00473-dependent differences in neuroeconomic decision-making behavior specific to alterations in

multiple features of change-of-mind re-evaluative choices without affecting primary decisions. Differences

were rooted in dissociable forms of counterfactual thinking, or how sensitive individuals were to missed

opportunities for alternative actions that could have led to hypothetically better outcomes – the

computational basis of regret processing previously captured by leveraging this task across species [5, 45,

46, 48, 56, 57].

Results:

<u>*Male and female mice similarly acquire the Restaurant Row task.*</u>

All mice underwent virus transfection surgeries and were allowed 2-3 weeks to recover before

beginning the longitudinal Restaurant Row paradigm (**Fig. 1a**). During the first week of Restaurant Row

testing, all reward offers were fixed at 1 s only (cued by a 4 KHz tone) as mice acquired the basic structure

of the task (**Fig. 1a-b**), learning to run in the correct direction to forage for low-cost food rewards. All mice

were food restricted to the same percentage (~80-85%) of baseline free-feeding weight (**Supplementary**

Fig. 2). Both males and females, regardless of GFP or LINC00473 treatment, grossly traveled similar total

distances across the first week (**Fig. 1c**, sex*virus: $F=0.971$, $p=0.331$). Behavior shaped equally among all

groups as the number of laps run in the correct direction (**Fig. 1d**, day: $F=334.219$, $p<0.0001$; sex*virus:

$F=2.780$, $p=0.104$) and total pellets earned (**Fig. 1e**, day: $F=393.991$, $p<0.0001$; sex*virus: $F=3.062$,

$p=0.089$) increased across the first week. All mice displayed subjective flavor preferences, revealed by

splitting earnings among flavors ranked from least to most preferred by summing end-of-session totals in

each restaurant (**Fig. 1f**, rank: $F=255.250$, $p<0.0001$). There was a modest upward shift in earns split by

flavor ranking in LINC00473-treated females across the first week (**Fig. 1f**, sex*virus: $F=3.334$, $p=0.076$).

Overall, these data indicate all groups were able to demonstrate the ability to acquire the basic structure of

the task and forage for food in a self-paced manner without overt locomotor or gross feeding differences.

<u>*LINC00473 alters value-dependent, task-driven conditioned place preference behavior in females.*</u>

Before and after the first week of behavioral testing, we characterized exploratory behavior in the

maze arena (**Fig. 2a-b**). That is, without any active task, mice were allowed to roam the maze freely on day

0 and again on day 8 of the experimental timeline (**Fig. 2b**). This allowed us to probe a conditioned place

preference (CPP) response for reward sites driven by experience that naturally developed throughout the

first week of self-paced task performance. We measured the number of entries into and time spent near

the feeder site in each restaurant's wait zone on these two probe sessions (**Fig. 2c-e**). All mice explored

the arena, including each restaurant, to similar degrees at baseline (day 0) and did not differ between

groups (sex*virus [entries]: $F=0.259$, $p=0.614$; sex*virus [time]: $F=1.155$, $p=0.290$, **Fig. 2c**). Overall,

compared to the day 0 pre-test baseline, we found that during the day 8 post-test all mice robustly increased

total time spent (F=75.818, p<0.0001) and to a lesser degree entries into the reward sites F=8.501, p<0.01,

Fig. 2c). We then split these metrics by restaurant ranked according to day 7's task earnings and found

that CPP scores scaled with flavor preferences (rank [entries]: F=16.307, p<0.0001; rank [time]: F=9.829,

p<0.0001, **Fig. 2f-g**). We found significantly lower CPP scores (post-pre delta score of 0 on average) with

respect to time spent in the wait zone of the most preferred flavor in LINC00473-treated females only,

despite being matched to males for equivalent number of rewards earned on day 7 of the task

(sex*virus*rank [time]: F=2.844, p<0.05, **Fig. 2f-g**). This sex-dependent effect was replicated in an

independent cohort of mice separately transfected with a lower titer of the AAV-expressing LINC00473

(**Supplementary Fig. 3**). Conversely, LINC00473-treated females displayed a higher number of post-pre

test entries into less preferred restaurants compared to males or GFP controls (sex*virus*rank [entries]:

F=5.584, p<0.001, **Fig. 2f-g**, **Supplementary Fig. 3**). These findings highlight that mPFC LINC00473

expression can have sex-specific effects on select actions during value-based reward-seeking behavior.

More specifically, LINC00473 treatment in females altered preference-dependent entering vs. exiting

choices that may access dissociable computations involved in checking-waiting-leaving behavioral

sequences [51, 58, 59], shedding light on more discrete interpretations of CPP metrics than are traditionally

reported in simpler assays and explored in detail below.

Escalating reward scarcity unmasks sex-dependent effects of LINC00473 on complex foraging profiles.

We next examined task foraging profiles across the subsequent weeks as we increased the

distribution of offer costs (the presented delays) available in the task environment without adjusting the

limited time budget mice had to forage for food (**Fig. 3a**). Thus, in this closed-economy system, tightening

budget constraints pressured animals to adapt to the changing landscape of the environment as reward

scarcity increased – a model of economic stress we recently characterized [33]. Costs increased in a

stepwise manner across blocks of days (i.e., offer delays ranged from 1-5 s on days 8-12, 1-15 s on days

13-17, and finally 1-30 s on days 18-45, as previously published [5, 33, 60]) and were randomly sampled

from a uniform distribution (the starting offer of each longer delay is associated with a discretely higher tone

pitch that is presented in the offer zone and counts down in the wait zone). As expected, we found a

decrease in pellets earned in response to each stepwise economic challenge (sign test [-] on earns: day
after minus day before transition, collapsed across epochs: $t=-12.655$, $p<0.0001$, **Fig. 3b**). These deficits
in earnings scaled by flavor and were matched between all groups (rank: $F=6.938$, $p<0.001$, timepoint*rank:
$F=3.651$, $p<0.01$; sex*virus: $F=0.016$, $p=0.899$, **Fig. 3c-d**, **Supplementary Fig. 4**). The most striking
change in foraging behavior that accounted for environmental scarcity-induced reward deficits across the
experiment was the proportion of session time re-allocated to quitting behavior in the wait zone, which
significantly increased at each economic challenge step and more prominently in females
(timepoint*choice: $F=226.523$, $p<0.0001$; timepoint*sex [quits]: $F=11.821$, $p<0.0001$, **Fig. 3e**,
Supplementary Fig. 4). Across the final epoch of testing (the 1-30 s offer range period: days 18-45), mice
displayed a complex behavioral response profile, including running more laps that developed over multiple
weeks in an effort to recover and increase total food earnings (laps [days 18-45]: $F=355.340$, $p<0.0001$;
earns [days 18-14]: $F=252.218$, $p<0.0001$, **Fig. 3b-d**). All groups were matched in their ability to recover
earnings (sex*virus: $F=0.234$, $p=0.631$, **Fig. 3d**, **Supplementary Fig. 4**). However, we found that females
displayed a greater increase in the number of laps run compared to males throughout this process (sex*day:
$F=7.848$, $p<0.01$, **Fig. 3b,f**, **Supplementary Fig. 4**). Interestingly, LINC00473-expressing females
displayed a significantly higher propensity to engage in wait zone quitting behavior compared to males
(sex*virus: $F=8.201$, $p<0.01$, **Fig. 3g**, **Supplementary Fig. 5**). Furthermore, LINC00473-expressing
females displayed shorter latencies to quit upon arrival once in the wait zone compared to males particularly
for more preferred flavors (sex*virus*day*rank: $F=2.805$, $p<0.05$), despite no changes in offer zone reaction
times ($F=1.713$, $p=0.162$, **Supplementary Fig. 5**). Taken together, these data reflect rearrangements in
complex foraging patterns that emerged across longitudinal time scales in response to worsening economic
demand. These analyses reveal sex-dependent effects of mPFC LINC00473 expression that alter quit
decisions and are not unmasked until mice encounter a reward-scarce environment. This effect of
LINC00473 in the wait zone is reminiscent of the altered check-then-leave behaviors in LINC00473-
expressing females during the CPP probe session (see **Fig. 2**)
196
**Females demonstrate unique neuroeconomic decision-making policies during change-of-mind choices.**

To characterize the economic nature of the decisions made during distinct action-selection

processes in the offer zone and wait zone, we quantified choice outcomes to enter vs. skip (offer zone) or

earn vs. quit (wait zone) separated by both the cued offer cost and flavor identity (**Fig. 4a-c**). All mice

acquired the economic structure of the task, making cost-informed enter vs. skip choices in the offer zone

or earn vs. quit choices in the wait zone, discriminating tones as well as demonstrating different degrees of

willingness to wait in each restaurant (offer*flavor [offer zone]: $F=13.129$, $p<0.001$; offer*flavor [wait zone]:

$F=18.006$, $p<0.001$, **Fig. 4a-c**). Thresholds of willingness to accept and separately wait for offers can be

calculated independently in the offer zone and wait zone by fitting a Heaviside step regression to the choice

outcome as a function of cost and extracting the inflection (indifference) point in each restaurant (**Fig. 4d**).

Overall, in all mice, we found that offer zone and wait zone thresholds significantly differed from each other

in magnitude of response to the changing economic landscape (timepoint*zone: $F=296.648$, $p<0.0001$, **Fig.**

4d). Upon transitioning to the 1-30 s epoch of testing (day 18), offer zone thresholds drastically increased

in all groups to similar degrees while retaining the ordinal ranking of flavor preferences (rank: $F=17.247$,

$p<0.0001$; sex: $F=0.170$, $p=0.681$; sex*virus: $F=0.031$, $p=0.861$, **Fig. 4d-f**). Wait zone thresholds, on the

other hand, increased only in male mice following the transition to the 1-30 s epoch (rank: $F=58.574$,

$p<0.0001$; sex: $F=17.776$, $p<0.0001$; sex*virus: $F=1.238$, $p=0.268$, **Fig. 4d-f**). Over the subsequent week

(approximately days 18-24), wait zone thresholds in males decreased before stabilizing across the

remainder of the experiment (approximately days 25-45, **Fig. 4d-f**). In females, wait zone thresholds

remained relatively unchanged upon transition to day 18 and were significantly lower than males across

the entire 1-30 s epoch (sex*rank [offer zone]: $F=0.484$, $p=0.694$; sex*rank [wait zone]: $F=3.689$, $p<0.05$,

Fig. 4d-f). Offer zone thresholds on the other hand gradually decreased, equally in all groups, but across

a much slower timescale over several weeks (**Fig. 4d-f**). Leveraging an economic analysis we recently

developed (taking individual differences in flavor-specific earnings on day 7 and projecting profiles to the

decision threshold learning that takes place during the 1-30 s epoch [33]), we found that female decision-

making policies were more optimized for both recovering total food intake as well as yielding maximal

subjective value once in a reward-scarce environment (**Fig. 4g-i**, **Supplementary Fig. 6**). LINC00473

expression in mPFC did not significantly alter any decision-making thresholds (sex*virus [offer zone]:

$F=1.781$, $p=0.185$; sex*virus [wait zone]: $F=187$, $p=0.175$, **Fig. 4d-f**). These data highlight a robust sex

difference between distinct offer zone and wait zone decision-making processes that we previously

demonstrated derive from separable circuits [48, 50].

LINC00473 in mPFC enhances change-of-mind contrast via sensitivity to sunk costs and action-specific

regret.

Wait zone thresholds capture economic indifference points of re-evaluation decisions defined by

choice outcome as a function of the starting offer. Next, we quantified more dynamically at what timepoint

quit decisions were made. Thus, we calculated the value remaining in the countdown at the moment of

quitting by subtracting the within-trial countdown time remaining from an individual mouse's wait zone

threshold (**Fig. 5a**). We found all mice generally made quit decisions (i) for offers that were typically above

one's wait zone threshold (i.e., negatively valued offers) and (ii) with ample time remaining in the countdown

such that the remaining time investment required to obtain the reward was still above one's threshold (i.e.,

negative value left), and therefore would be economically advantageous or efficient and timesaving to quit

(**Fig. 5a**). Thus, overall, quit decisions reflected corrective re-evaluations of sub-optimal offer zone

decisions relative to one's own policy, consistent with previous reports [5, 33, 48, 49, 51]. We found that

females made quit decisions with significantly lower and more negative value remaining in the countdown

compared to males (sex*virus: F=195.823, p<0.0001, **Fig. 5b**). This sex difference was more pronounced

in LINC00473-expressing animals and thus made females quit decisions even more efficient compared to

males (sex*virus: F=12.957, p<0.001, **Fig. 5b**). Furthermore, only LINC0047-expressing mice revealed a

sex difference in the number of quit decisions that could be characterized as economically advantageous

(as described above, sex*virus: F=153.236, p<0.0001), with a greater proportion of these events in females

compared to other quit events that were less common and more disadvantageous (i.e., quitting negatively

valued offers [above one's threshold] with positive value left [below one's threshold]; or, quitting positively

valued offers [below one's threshold] – both reflecting quits passing the timepoint after which mice should

have chosen to stay instead (quit type*sex*virus: F=13.293, p<0.0001, **Fig. 5c-d**).

We then characterized the temporal dynamics of how mice made continuous re-evaluation

decisions in the wait zone as a function of the passage of time, which wait zone thresholds do not capture.

We calculated the instantaneous probability of staying and earning a reward as a continuous function of

not only (i) the time remaining in the countdown at each snapshot in time but also (ii) the time already spent

waiting. We computed a survival analysis using a sliding 1 s window after binning all trials of varying starting

offers into [time spent, time left] pairs (**Fig. 5f**). In all groups, we found that the passage of time drove an

increased likelihood that mice would continue to wait, the effect of which grew additively stronger with

escalating time spent independent of and orthogonal to temporal distance to the goal measured at any

given moment in time (time spent*time left: F=891.924, p<0.0001, **Fig. 5f-h**). This captures a well-studied

cognitive phenomenon known as the sunk cost bias whereby irrecoverable past expenditures can enhance

the value of continuing to invest in and perseverate on an ongoing endeavor, as we have reported

previously captured across species [46, 49]. We found a significant sex difference in sensitivity to sunk

costs only in LINC00473-expressing mice. LINC00473-expressing females demonstrated increased

sensitivity to sunk costs compared to LINC00473-expressing males, which instead demonstrated relatively

diminished sensitivity to sunk costs during quit decisions (sex*virus: F=15.944, p<0.0001, **Fig. 5i**). This

effect only emerged after mice transitioned to the 1-30 s reward-scarce environment (time spent in wait

zone [1-15 s offers]: F=0.174, p=0.677, **Fig. 5i**, **Supplementary Fig. 7**) and uniquely affected wait zone

decisions compared to other forms of time spent on this task, including in the offer zone (time spent in offer

zone: F=0.558, p=0.455; time spent*sex*virus: F-0.418, p-0.518, **Supplementary Fig. 7**). These data

highlight sex- and LINC00473-dependent alterations in how the passage of time can differentially enhance

reward value and escalate the commitment to an ongoing endeavor specifically during re-evaluation

decisions.

Next, we aimed to characterize how events related to quit decisions on a given trial influenced

subsequent behavior. Because trials are interdependent, a strength of the Restaurant Row task lies in being

able to subdivide trial sequences into unique economic scenarios in which animals naturalistically find

themselves. Analysis of mice after they make suboptimal choices and potentially realize when gaining

access to information that an alternative, unselected action could have led to a better outcome has led to

breakthrough discoveries modeling across species aspects of the behavioral and neurophysiological

signatures of counterfactual thinking – the computational basis of regret processing [5, 47, 48, 61-63]. We

previously linked such processes to quit decisions as well as other types of economic choices on this task

[5, 47, 48]. In past reports, we and others behaviorally operationalized counterfactual sensitivity on

Restaurant Row by identifying trial pair sequences [trial t-1 → trial t] in which animals made economically

suboptimal choices [on trial t-1] and by measuring the change in economic choice behavior on the next trial

[on trial t: the read-out trial] [5, 47, 48]. Importantly, these events are compared to specific control sequences

that remove the error of one's own agency by examining non-violation events [on trial t-1], matched for all

other economic factors (e.g., including the offer values and flavor identities on both trials t-1 and trial t) in

order to behaviorally measure changes in decision-making counterfactual based on one's choice history:

our operational behavioral definition of counterfactual sensitivity. Recent work demonstrated the importance

of the mPFC in these behavioral sensitivity measures on our neuroeconomic task [48]. Sequences were

extracted post hoc from natural encounters while foraging amid randomly presented offers, not constructed

a priori nor built into the task design.

295   We previously defined two distinct regret-related sequences of events on this task [5, 47, 48].

Scenario 1: situations in which animals inappropriately forgo a high-value offer [on trial t-1] only to

subsequently encounter a worse low-value offer on the next trial [trial t] (**Fig. 5j**). This violation sequence

can be compared to the non-violation control sequence in which mice instead appropriately accepted the

high-value offer on trial t-1, holding all other variables constant (**Fig. 5j**). Here, we found that all groups

were insensitive to scenario 1, with no effect of sex or LINC00473 expression (+ sign test: [male GFP]: t=-

0.282, p=0.608; [female GFP]: t=0.101, p=0.461; [male LINC]: t=0.521, p=0.308; [female LINC]: t=-0.144,

p=0.556, **Fig. 5j**). That is, there were no changes in economic choice behavior on the read-out trial when

comparing violation sequences to non-violation sequences [thus yielding a delta score of zero in change of

choice probabilities on trial t] (**Fig. 5j**). **Scenario 2**: situations in which animals inappropriately accepted a

low-value offer [on trial t-1] that typically precipitates a resultant change-of-mind decision (**Fig. 5k**). This

violation sequence can be compared to the non-violation control sequence in which mice instead

appropriately skipped the low-value offer on trial t-1, holding all other variables constant and matched for

the same low-value offer on trial t as in scenario 1 (**Fig. 5j-k**). Overall, we found that mice demonstrated

sensitivity to scenario 2, whereby individuals were more likely to compensate for and accept low-value

offers that they typically would not on the read-out trial following violation sequences compared to non-

violation sequences (**Fig. 5k**). Interestingly, only GFP-treated females did not display scenario 2 sensitivity,

while all other groups including LINC00473-expressing females displayed delta scores significantly greater

than zero (+ sign test: [male GFP]: t=2.931, p<0.01; [female GFP]: t=1.123, p=0.145; [male LINC]: t=3.056,

p<0.01; [female LINC]: t=2.196, p<0.05, **Fig. 5k**). Taken together, these data reveal how mice can

differentially carry the weight of distinct types of past mistakes into future choices, with a unique association

between change-of-mind-related counterfactual sensitivity (scenario 2, not scenario 1) and mPFC

LINC00473 expression in females. These findings provide an example whereby LINC00473 expression in

this brain region overcomes a baseline sex difference that potentially normally disadvantages females.

Lastly, we characterized how access to task-relevant information may be critically important for

sex- and LINC00473-dependent differences during change-of-mind decisions. After day 45, we tested mice

for an additional 2 days during which tone playback on the task was silenced only in the wait zone but

remained audible during offer zone presentations (**Fig. 6a**). That is, mice had access to cost information

signaled by tone pitch when making initial enter vs. skip decisions in the offer zone, but received no

countdown information once in the wait zone while making secondary re-evaluative decisions to quit or stay

before earning rewards. Overall, we found that males were more sensitive to this informational manipulation

compared to females. Both GFP- and LINC00473-expressing males displayed a modest decreased number

of laps run and total number of pellets earned, while female mice remained relatively unchanged between

audible and silent conditions (laps [timepoint*sex]: F=10.678, p<0.01; laps [timepoint*virus]: F=0.449,

p=0.507; earns [timepoint*sex]: F=10.124, p<0.01; earns [timepoint*virus]: F=0.002, p=0.969, **Fig. 6b-c**).

This task manipulation did not alter any overt changes in offer zone choices in any group as animals

continued to discriminate the audible tones during cost-informed enter vs. skip decisions (offer:

F=1186.818, p<0.0001; offer*timepoint: F=1.365, p=0.243; **Fig. 6d**). Interestingly, when examining changes

in wait zone thresholds, we found a significant sex difference when wait zone countdown information was

withheld that was compounded by an additional LINC00473 effect in females only (**Fig. 6e**). In males,

compared to the audible condition, the lack of auditory information cuing mice on progress during the

countdown in the silent wait zone condition drove an increase in wait zone thresholds for most preferred

flavors (timepoint: F=22.976, p<0.0001; timepoint*virus: F=0.181, p=0.671 **Fig. 6e**). Males also displayed a concomitant decrease in wait zone thresholds for least preferred flavors (timepoint: F=19.561, p<0.0001; timepoint*virus: F=0.526, p=0.470, **Fig. 6e**). These data indicate that, without cued auditory guidance in the wait zone, males were more patient waiting for favorite flavors and less patient waiting for least favorite flavors (defined by a change in the indifference point of earn outcomes as a function of the trial's starting offer). Conversely, GFP-expressing females only displayed a decrease in wait zone thresholds for less preferred flavors, with no change in thresholds for most preferred flavors (**Fig. 6e**). LINC00473-expressing females however did not display any changes in wait zone thresholds in any restaurant compared between audible and silent wait zone conditions (timepoint [most preferred]: F=2.569, p=0.112; timepoint*virus [most preferred]: F=0.137, p=0.712; timepoint [least preferred]: F=12.185, p<0.0001; timepoint*virus [least preferred]: F=8.901, p<0.01). These data indicate LINC00473-expressing females valued offers in a consistent manner with or without externally guided information, suggesting that their internal measures of time or how they hold and maintain information from the offer zone in working memory may be more stable. When examining quit times in the wait zone, all mice displayed a modest increased latency to make quit decisions in the silent condition compared to the audible condition (timepoint: F=11.764, p<0.001; sex: F=2.358, p=0.126; virus: F=0.836, p=0.361, **Fig. 6f**). Finally, when examining sensitivity to sunk costs, we found that the silent condition unmasked a sex difference in GFP-treated mice whereby females show increased sunk cost sensitivity as a function of the passage of time compared to males – a sex difference that was previously only apparent during the audible condition in LINC00473-expressing groups (time spent: F=97.196, p<0.001; sex: F=36.989, p<0.0001; virus: F=1.899, p=0.168; **Fig. 6g**, see **Supplementary Fig. 8** for alternate task information manipulations). Taken together, these findings highlight the complexity of change-of-mind decisions, how mice internally generate value representations with vs. without access to explicit, exogenous task-relevant information, and the multifactorial ways in which this re-evaluative choice process can be uniquely modulated in a sex-dependent manner.

Discussion:

In this study, we characterized highly complex behaviors in mice tested on the neuroeconomic foraging paradigm, Restaurant Row. We explored the role of a non-coding RNA, a relatively poorly understood class of molecules, on motivated behavior. We discovered sex-dependent effects of LINC00473 expression in mPFC neurons on specific aspects of value-based decision-making information processing. This was typified by selectively altering the way in which an individual changes one's mind during an ongoing decision. Importantly, we discovered a role for mPFC LINC00473 expression – which we recently found increases stress resilience in simple behavioral screening tests in female mice only [19] – in sharpening the contrast of re-evaluation processes in females. Our findings extend from simple conditioned place preference behaviors like those commonly assayed using simpler goal-oriented paradigms [40, 64, 65] to more complex, higher order neuroeconomic phenomena. These latter effects included enhancing specific aspects of sensitivity to sunk costs and counterfactual thinking. These data provide a deeper understanding of how multiple stages of a continuous decision stream derive from dissociable psychological mechanisms supported by physically separable circuits in the brain. Our findings have important implications for how molecular determinants of stress responses and non-coding regions of the genome can influence changes in motivated behavior via unique region-specific decision-making computations differentially leveraged by males and females.

The distinction between offer zone choices and wait zone choices on the Restaurant Row task is critically important. They reflect a separation of fundamentally distinct decision-making algorithms that occur within the same trial but are segregated across space and time [51]. Numerous reports have tested subjects on this paradigm translated for use across species in task variants adapted for mice, rats, monkeys, and humans [41, 42, 46]. These studies have shed light on the biological and psychological implications of resolving dissociable, evolutionarily conserved decision-making processes. In the offer zone, individuals engage deliberative systems that represent alternating options of competing actions between which one must choose [54, 55]. This has been shown in rodents to derive in part from hippocampal (HPC)-dependent processes [50, 51]. Population level activity of neurons in HPC is capable of decoding multiple, alternating

future paths through the offer zone to potential goals in the current vs. next restaurant [58, 64, 66, 67].

Similar findings have been observed from fMRI studies in humans tested on a translated version of the

Restaurant Row task [45]. Default mode network activation patterns can decode competing future options

during offer zone deliberation, but seldom do such representations include past events or previous

restaurants [2, 68]. Decisions in the wait zone on the other hand are much less understood. The HPC has

been implicated in wait zone decisions as well: it switches oscillatory activity from a predominately theta-

driven state sensitive to the cued offer costs in the offer zone [54] instead to an increase in sharp wave

ripple activity sensitive to time spent waiting in the wait zone [55]. Further, wait zone decision periods also

depend on HPC synchrony with the mPFC [54, 55]. Recordings in rodents from the mPFC on Restaurant

Row surprisingly found no evidence of encoding of the cued offer costs when animals were in the offer zone

[52]. However, once in the wait zone, mPFC neurons strongly represented offer value independent of time

spent waiting and could predict whether or not the trial would terminate in a quit [52]. Within the mPFC,

pairs of neurons also drastically changed excitation-inhibition balance specifically when animals were in the

wait zone [69]. Taken together, these findings indicate a qualitative shift in mental state upon transition from

the offer zone to the wait zone that recruits the mPFC exclusively during change-of-mind decisions.

This study represents the first analysis of females in the Restaurant Row task. Nearly all of the sex

differences captured here, including those altered by mPFC LINC00473 expression, are restricted to wait

zone and not offer zone behaviors. Furthermore, sex differences not only persist but are even exaggerated

when access to task-relevant cost information is restricted in the wait zone during change-of-mind

decisions. We showed recently that LINC00473 expression diminishes spontaneous excitatory post-

synaptic current frequency and amplitude in excitatory mPFC pyramidal neurons of female mice, with no

effect seen in males [19]. Chemogenetic inactivation of the mPFC during Restaurant Row diminishes wait

zone quit decisions [53-55]. Optogenetic induction of long-term depotentiation in excitatory mPFC neurons

that project to the nucleus accumbens (NAc) [50], where negative deflections in dopamine signals have

been found to causally precede quit choices on Restaurant Row [70], selectively decreases quitting

behavior for several days without altering offer zone behavior [50]. Outside of Restaurant Row, intracranial

optogenetic excitatory self-stimulation of mPFC terminals in the NAc can be reinforcing while also promoting

enter-then-quitting behavior [71]. These converging findings suggest that LINC00473 expression is likely

mediating sex-dependent effects on change-of-mind decisions by at least in part dampening and

restructuring mPFC activity. How these alterations in change-of-mind decisions could relate to mPFC

LINC00473's known role in enhancing resilience to stress in a sex-dependent manner will help bridge

interpretations between molecular drivers of stress responses and the behavioral computations associated

with susceptibility vs. resilience to stress in males and females.

Recent transcriptomic analyses of human postmortem mPFC obtained from individuals diagnosed

with major depressive disorder revealed a female-specific decrease in LINC00473 expression in the mPFC

[19]. While LINC00473 is not normally expressed in rodents – it arose in the primate lineage – expressing

LINC00473 in mouse mPFC neurons demonstrated pro-resilient effects in females, but not males,

protecting against several depressive-like phenotypes in simple behavioral tests after exposure to chronic

social defeat or variable stress [19]. LINC00473 expression, which predominates in pyramidal neurons in

human mPFC [19], has been shown to be regulated by conserved Ca^{2+}/calmodulin signaling pathways and

is likely induced by cAMP response element-binding protein (CREB) [19-21, 23]. CREB has long been

studied based on more traditional candidate gene approaches and shown to regulate numerous other

genes involved in neuronal excitability and synaptic plasticity as well as stress sensitivity [72-77]. Like

LINC00473, CREB function in the mPFC has been implicated in promoting resilience to stress, however,

unlike LINC00473, this has been shown to be true for both males and females [19, 78]. These reports

suggest either that there must be some divergence of intracellular CREB-LINC00473-interacting molecular

pathways between sexes conferring differential sensitivities to stress, or that prior studies were limited in

their ability to resolve more nuanced sex differences at the behavioral level.

We recently characterized the impact of chronic social defeat stress on Restaurant Row behavior

in male mice, in tandem with disruptions of CREB function in mPFC [33, 48, 79]. First, we found that male

mice were required to progress through worsening economic challenges in order to extract any group

differences in foraging profiles [33]. This is consistent with results presented here that emphasize the

importance of considering the environmental demand and stressors of a task necessary to unmask disease-

relevant phenotypes. Second, we found that stress-resilient mice – defined by a simple social interaction

screening test measured outside of the Restaurant Row task – mounted the most robust behavioral

activation in response to the transition into the 1-30 s reward-scarce environment in efforts to recover and

maintain food security despite budget constraints, similar to LINC00473-expressing female mice as shown

here [33]. Third, without grossly altering decision-making thresholds, we found that stress-susceptible male

mice demonstrated increased regret-related sensitivity to type 1 scenarios (economically risky decisions to

skip, followed by poor outcomes), a phenotype no group demonstrated in the present study [48].

Additionally, stress-susceptible mice demonstrated diminished regret-related sensitivity to type 2 scenarios

(change-of-mind decisions) compared to stress-resilient individuals who instead displayed enhanced

change-of-mind sensitivity to both regret and sunk costs related to quitting behavior [48]. Interestingly, in

the present study, this constellation of differences was only apparent among the GFP-expressing and

LINC00473-expressing females, respectively, suggesting that females at baseline have a decision-making

vulnerability specifically in change-of-mind choice processes that is restored to male levels by LINC00473

expression in mPFC. In our earlier study, knocking down CREB function had multifactorial effects

depending on the neuroeconomic valuation algorithm in question, the brain region targeted (mPFC vs.

NAc), and whether or not animals were stress-naïve at baseline vs. exposed to subsequent stress [48].

CREB function in the mPFC was required to maintain suppression of regret-related sensitivity to type 1

scenarios and, when knocked down, rendered only type 2 scenarios sensitive to subsequent stress – a

response profile distinct from CREB function in the NAc where it is known to drive stress susceptibility [48,

72, 80, 81]. These data shed light on a divergence between CREB and LINC00473 in mediating dissociable

aspects of counterfactual thinking that could independently contribute to the affective toll of past mistakes

and one's overall mood burden. These data also point to which consequence of changes in counterfactual

sensitivity (i.e., resisting an increase in type 1 or promoting type 2) could be uniquely ascribed to the pro-

resilient features of mPFC LINC00473 expression. While these prior studies were conducted in males [33,

48, 79], they highlight multiple dimensions with which to consider how information integrates across

fundamentally distinct choice processes in order to ultimately precipitate sex-dependent behavioral

phenotypes related to one's stress vulnerability.

LINC00473's sex-dependent effects on behavior have only been causally tested in the mPFC to

date [19]. Although postmortem human tissue studies implicate mPFC as a top region to investigate [19],

understanding LINC00473's role in other regions, especially in how it may be interacting with other key

molecular players such as CREB in the NAc [23, 72, 73], will be critically important in future studies to

elucidate how information processing integrates across neural networks. For instance, LINC00473 has

been shown to increase cellular excitability in cultured HPC neurons [23], contrary to its effects in mPFC

[19]. Future decision-making studies should combine experiments of LINC00473 manipulations with

exposure to other stressors, beyond the economic challenges tested in the present study, such as chronic

social defeat or variable stress as well as metabolic effects of these stressors to explore compounding

effects on decision-making [19]. Further, more work needs to be done toward investigating the role of

gonadal steroid hormones, sex chromosomes, and sex-dependent noncoding RNAs in mediating aspects

of the reported sex differences. As just one example, CREB function has been shown to fluctuate with

estrus cycle [82-86]. Additionally, given previous reports that LINC00473 expression in mPFC of stress-

naïve male mice can induce behavioral changes in certain simpler assays (i.e., marble burying) [19], it

would be fruitful to explore in future work how Restaurant Row decision-making profiles could potentially

relate to individual differences in other traits outside of the task itself (e.g., anxiety-related behavior

measured on independent tests). Lastly, it is critically important to consider how the decision-making

processes as measured here could relate directly to and drive information processing during the expression

of one's symptomology itself (e.g., underlying the choices leading to anhedonia or social avoidance related

behaviors themselves) as opposed to occupying a more general role as a trait of the individual.

Taken together, we discovered robust sex differences in the overall contrast of how an individual

decides to change one's mind. This is conceptualized as how dynamic a value representation can be when

deciding to quit an ongoing endeavor and the strength with which quitting behavior can influence the

psychology of multiple cognitive processes. We characterized this as both the steepness of a tipping point

during change-of-mind decisions to either stay or quit and the weight with which such events can be carried

into future choices. We found that this lability depends on individual differences in one's subjective value,

on the state of the environment in the form of how scarce reward availability and access to information

might be, and is uniquely enhanced by LINC00473 expression in the mPFC of females. Counterfactual

thinking during change-of-mind decisions can elicit "should not have done that" mental representations that

map hypothetical outcomes onto alternative actions that could have been selected instead. Even if change-

of-mind decisions are an unpleasant experience or evoke to some degree a form of cognitive dissonance,

enhancing the sharpness and salience of this process could increase attention paid toward realized losses,

can possess utility for fictive learning, and collectively may be a sex-dependent feature of stress resilience

altering the lens through which individuals process their prior mistakes.

Figure Captions:

Fig. 1 | Male and female mice acquire the Restaurant Row task. a Experimental timeline. 40 C57BL/6J

mice (20 male, 20 female) underwent stereotaxic surgery, half of each sex receiving AAV-LINC00473 or

AAV-GFP viruses injected bilaterally in the mPFC. Mice were allowed to recover before beginning

longitudinal testing on the Restaurant Row task for 45 consecutive days. Dashed box indicates time period

relevant for this figure: week 1 of behavioral testing. **b** Task schematic. Mice were allotted 30 min daily to

invest time foraging for their primary source of food in a self-paced manner. Costs to obtain rewards were

in the form of delays mice would have to wait near feeder sites. Mice were required to run in a

counterclockwise direction encountering offers for different flavors at each "restaurant" in serial order. Each

restaurant, separated by hallways, was divided into a T-shaped "offer zone" choice point and a separate

"wait zone" that housed the pellet dispenser. Upon offer zone entry from the correct heading direction, a

tone sounded whose pitch indicated the delay mice would have to wait if accepting the offer by entering the

wait zone. If entered, tone pitch descended in the wait zone, cuing the indicated delay. Each trial terminated

if mice skipped in the offer zone, quit during the countdown in the wait zone, or earned a reward, after which

animals were required to proceed to the next restaurant. **c-f** Simple behavioral metrics across the first week

of testing during which all offers were 1 s only (lowest pitch, 4 kHz): (c) total distance traveled, (d) laps run

in the correct direction, (e) total rewards earned, (f) earnings split by flavors ranked from least to most

preferred by summing each day's end-of-session totals in each restaurant. No effect of sex or LINC00473

expression was grossly apparent in these initial metrics. Error bars represent ±1 SEM.

Fig. 2 | LINC00473 expression in mPFC alters value-dependent, task-driven conditioned place

preference behavior in females. a Experimental timeline. Dashed box indicates time period relevant for

this figure: two special testing days marked with cyan arrows. **b** Experimental design. Animals were placed

in the arena free to roam for 20 min on day 0 and day 8 of the experimental timeline but with no active task

to obtain pre-task baseline and post-task experience-related exploratory behavior. **c** Total number of entries

and total time spent in the wait zone, aggregated in all restaurants. **d-e** Representative video tracking data

from an example mouse [GFP-treated female] recorded during the pre (day 0) and post (day 8) tests. (d)

Track plots displaying body position when traveling between restaurants (gray) and when in the wait zone

of each restaurant (colored). (e) Occupancy heat maps of the same example mouse in (d). The change in

entries into and time spent in each restaurant's wait zone can be calculated as a delta score (post minus

pre). **f-g** Delta scores in change of (f) total entries and (g) total time comparing post (day 8) minus pre (day

0) time points split by restaurants ranked according to rewards earned on the active task on day 7. Only

the most preferred (1st) and least preferred (4th, faint) restaurants are depicted here (see **Supplementary**

Fig. 3 for full dataset and replication cohort). Horizontal dashed gray line indicates delta score of 0.

LINC00473 expression in mPFC abolished CPP behavior in most preferred restaurants only in females and

only in the time but not entry domain. *Represent significant sex differences. Error bars represent ±1 SEM.

Fig. 3 | Escalating reward scarcity unmasks sex-dependent effects of LINC00473 expression in

mPFC on complex foraging profiles. a Experimental timeline. Dashed box indicates time period relevant

for this figure: the entire 45-day Restaurant Row paradigm. **b** Total rewards earned (black) and laps run in

the correct direction (gray). All 40 mice collapsed. Visual guidance color bars along the x-axis reflect the

experimental schedules of the change the distribution of offer costs illustrated in the timeline in (a). Vertical

dashed green-yellow-red lines indicate the transition points of the stepwise schedule of and are re-used

throughout all other figures as a visual aid. Inset depicts "hysteresis" plot of laps against earns (epochs of

days are color-coded). **c** Rewards earned split by flavors ranked from most preferred (1st) to least preferred

(4th, faint). All 40 mice collapsed. **d** Change in rewards earned in each restaurant comparing two days at

various color-coded transition points (delta scores calculated by subtracting the following: i, day 8-7

(yellow); ii, day 13-12 (orange); iii, day 18-17 (red); iv, day 45-18 (purple). All 40 mice collapsed (see

Supplementary Fig. 4 for comparison between groups). **e** The portion of total session time engaged in one

of the four major choice behaviors: offer zone – enter (green) or skip (red); wait zone – quit (magenta) or

earn (blue). Rings split restaurants ranked from most preferred (1st, inner ring) to least preferred (4th, outer

ring). Each of the 5 pie charts represent data from each stage of testing, with the early and late epochs

during the 1-30 s offer range extracted from the first 5 and last 5 days of that period. **f** Laps run in the correct

direction split by sex and treatment groups. **g** Proportion of enter decisions that terminated in a quit outcome

once in the wait zone. LINC00473 expression in mPFC drove females to run more laps in response to a

worsening economic landscape and quit more frequently. ***Represent significant sex differences. Shading

/ error bars represent ±1 SEM.

Fig. 4 | Females demonstrate unique neuroeconomic decision-making policies during change-of-

mind choices. a Experimental timeline. Dashed box indicates time period relevant for this figure: the entire

45-day Restaurant Row paradigm. **b** Example behavior from a single mouse early or late during the final 1-

30 s testing epoch (red portion of the experimental timeline). Data depict offer zone choice outcomes on

the y-axis (enter vs. skip) as a function of the randomly presented cued offer cost (tone pitch) on the x-axis

in each restaurant (dots represent individual trial choice outcomes). Quit outcomes in the wait zone are

indicated in black dots. **c** Aggregate example summary choice data from all 40 mice early or late during the

final 1-30 s testing epoch. Data depict separate offer zone or wait zone summary probability of choice

outcome. Horizontal dashed gray line indicates chance at 0.5. Mirror curves for skip and quit (1-

probabilities) are intended to highlight cross over points that approximate threshold fits. **d** Thresholds or

indifference points of willingness to accept offers in the offer zone or earn offers in the wait zone. Inset

depicts an example session from a single mouse in one restaurant where vertical solid green or blue lines

illustrate the inflection of Heaviside-step regression fits to choice outcomes in each zone as a function of

starting offer costs. All 40 mice collapsed. Inset depicts "hysteresis" plot of offer zone against wait zone

thresholds (epochs of days are color-coded). **e** Offer zone thresholds (top) and wait zone thresholds

(bottom) split by sex and treatment groups. GFP groups in faint dashed lines. **f** Thresholds from (e)

extracted from early and late 1-30 s epoch (red). GFP groups in dashed lines. **g-i** Economic analysis of

how well offer zone and wait zone decision policies during the 1-30 s epoch approximate subjective value

yield as determined from individual differences in earnings by flavor acquired during the low-cost

environment earlier in testing (day 7) when all rewards cost only 1 s. This is compared against theoretical

decision policies that maximize total food intake during the 1-30 s epoch irrespective of flavor. See **Methods**

and **Supplementary Fig. 6** for a full explanation of this analysis. (g) Individual differences in (top) the

standard deviation of rewards earned across flavors on day 7, (middle) total rewards earned on day 7, and

(bottom) earns split by restaurant coded in color (flavor: chocolate-brown, banana-yellow, grape-purple,

plain-cyan) and size (reward count) on day 7. All data in (g) is sorted by stdev data in (top). Small colored

boxes along the x-axis indicate sex (male-blue, female-red) and treatment groups (LINC-dark, GFP-light).

h Example data from a single mouse on day 7 showing value ratios among earns between restaurants

relative to the least preferred flavor (faint). **i** Data from (g-h) used in a computational modeling analysis to

calculate the Euclidean distance between observed decision policies each day from days 18-45 (cool to

warm colors, each dot indicates group average per day) to theoretical policies that could achieve either

maximal food (x-axis) or maximal subjective value (y-axis) for offer zone and wait zone thresholds. Inset

diagonal lines indicate reference point for the 1:1 unity line. Prominent sex difference, without apparent

effect of LINC00473 expression in mPFC, in present in wait zone but not offer zone economic decision

policies. *Represent significant sex differences. Shading / error bars represent ±1 SEM.

Fig. 5 | LINC00473 expression in mPFC enhances change-of-mind contrast via sensitivity to sunk

costs and action-specific regret. a Distribution of quit trials as a function of offer value (x-axis) or value

left at the moment of quitting (y-axis) defined relative to one's wait zone threshold (origin vertical and

horizontal dashed blue lines). Data split by restaurants ranked from least to most preferred. All 40 mice

collapsed. Data extracted from the entire 1-30 s epoch. **b** Average value left at the moment of quitting split

by sex and treatment groups. Horizontal dashed gray line indicates 0 value left (or wait zone threshold). **c**

Three categories of quit events based on offer value and value left terms. The green quadrant (Vo<0 &

V_L<0) represent the most efficient quit decisions from economically disadvantageous offers. **d** Counts of

total number of quit events aggregated across all restaurants from each economic quit category described

in (c). **e** Counts of total number of quit events from each economic quit category in (d) split by restaurants

ranked from least (4th) to most preferred (1st) extracted from the entire 1-30 s epoch (red). **f-h** Sunk cost

analysis of staying behavior in the wait zone, demonstrated using all 40 mice. (f) The likelihood of staying

in the wait zone and earning a reward (e.g., not quit) is plotted as a function of time left in the countdown

along the x-axis and time already spent waiting orthogonally in color. Note the black 0 s time spent curve

represents animals having just entered the wait zone from the offer zone. Inset vertical dashed gray line

illustrates an example analysis point comparing three sunk cost conditions originating from different starting

offers but matched at 10 s left. Data from (f) dimensioned reduced in (g) collapsing across time left,

highlighting the grand mean of each time spent sunk cost condition (color and x-axis). Insets in (g) depict

data from curves in (f) are collapsed into the observed (sunk condition) and control (0 s condition) lines.

Difference between curves in (g) are plotted in (h) in order to summarize the envelope of the overall effect

of time already spent on escalating the commitment of staying in the wait zone. Horizontal dashed line

represents delta score of 0. **i** Sensitivity to sunk costs plotted across the entire Restaurant Row paradigm

where the delta curve in (h) is plotted as vertical slices in (i) with time already spent along the y-axis and

heatmap representing delta score magnitude. **j-k** Regret-related sequence analyses comparing trial pairs

between (j) type 1 scenarios and (k) type 2 scenarios each. Read out trials captures a change in behavior

depending on choice history of distinct economic violations (black) compared to non-violations (gray) on

trial t-1. Read out trials report delta scores in likelihood change of entering by subtracting choice outcomes

between violation-nonviolation sequences. Horizontal dashed gray line indicates delta score of 0. See

Methods for a full explanation of these analyses. Prominent sex-dependent effect revealed by LINC00473

expression in mPFC on change-of-mind related sunk costs (enhanced in females) and regret sensitivity

(type 2: enhanced in females). * in (b-i) represent significant sex differences. * in (j-k) represent delta score

significantly different from 0. Shading / error bars represent ±1 SEM.

Fig. 6 | Sex- and LINC00473-dependent wait zone behaviors are differentially modulated by access

to task information. a Informational manipulation schematic: (left) standard Restaurant Row task followed

by (right) two days of tones silenced only in the wait zone, but not offer zone (remains audible as before).

b Total number of laps run in the correct direction (top) and total pellets earned aggregated among

restaurants (bottom) across the audible and silent wait zone task conditions (x-axis). **c** Rewards earned in

each restaurant ranked from least (4th) to most preferred (1st) across the audible (black) and silent wait zone

(gray) task conditions. **d** Overall offer zone choice probabilities to make enter decisions as a function of

audible cued offer costs collapsed across all flavors to illustrate no effect of task condition on auditory

discriminatory behaviors. Horizontal dashed line indicates change at 0.5. **e** Wait zone thresholds in each

restaurant across task conditions. **f** Latency to quit reaction times once in the wait zone in each restaurant

across task conditions. **g** Sensitivity to sunk costs in the wait zone depicted here only in the silent condition

as a function of time already spent (x-axis). Y-axis reflects delta earn probabilities as depicted in Fig. 5h.

Prominent sex-dependent effects of withholding wait zone information on overall task performance

(observed in males), wait zone economic decision policies (bidirectional in males, unidirectional in GFP-

females, and no effect in LINC00473-females), and sunk cost sensitivity (sex difference revealed for the

first time GFP groups unlike in Fig. 5). Horizontal dashed line indicates delta score of 0. * in (b-f) represent

significant effects of task condition within group. * in (g) represent significant sex differences. Shading /

error bars represent ±1 SEM.

Acknowledgments:

We thank members of the labs of Eric Nestler and Scott Russo for helpful discussion and technical

assistance. Open-source illustrations obtained from SciDraw (www.scidraw.io), credit Federico

Claudi.

Funding:

National Institute of Mental Health grant R01MH051399 (EJN)

National Institute of Mental Health grant R01MH129306 (EJN)

National Institute of Mental Health grant R01MH114882 (SJR)

National Institute of Mental Health grant R01MH127820 (SJR)

National Institute of Mental Health grant R01MH104559 (SJR)

National Institute of Mental Health grant L40MH127601 (BMS)

National Institute of Mental Health training grant R25MH129256 (BMS)

National Institute of Mental Health supplement grant R01MH051399-31S1 (BMS)

Leon Levy Scholarship in Neuroscience, New York Academy of Sciences (BMS)

Burroughs Wellcome Fund Career Award for Medical Scientists (BMS)

Animal Models for the Social Dimensions of Health and Aging Research Network via NIH/NIA R24

673   AG065172 (BMS)

Brain & Behavior Research Foundation Young Investigator Award 31140 (RDC)

Author contributions:

Conceptualization: RDC, OI, EJN, BMS

Methodology: RDC, OI, BMS

Investigation: RDC, BY, NJ, AA, SK, KG, SA, SJR, EJN, OI, BMS

Data curation: RDC, OI, BMS

682  Formal analysis: BMS

683  Visualization: RDC, OI, BMS

684  Funding acquisition: OI, BMS

685  Supervision: RDC, BMS

686  Writing – original draft: RDC, OI, BMS

687  Writing – review & editing: RDC, BY, NJ, AA, SK, KG, SA, SJR, EJN, OI, BMS

689  **Competing interests:**

691  Authors declare that they have no competing interests.

693  **Data and materials availability:**

695  All data and materials used in the analysis are available in the manuscript, materials and methods

696  section, or supplementary information.

Supplementary Figure Captions

Supplementary Fig. 1 | Validation of newly packaged adeno-associated virus expression of

LINC00473. Raw qPCR cycle threshold count (ct) obtain from mPFC tissue punches extracted from mice

separately transfected with either AAV2-hSyn-GFP (control) or AAV2-hSyn-LINC00473-GFP (treatment).

qPCR readouts obtained for expression levels of GFP, HPRT1 (housekeeping), ZFP189 (another known

regulator of stress-resilience), and LINC00473. Note similar levels of expression of GFP (t=2.03, p=0.089),

HPRT1 (t=0.41, p=0.693), and ZFP189 (t=0.91, p=0.396) between control and treatment viruses. Also note

no detectable levels of LINC00473 in mice transfected with the control virus, as this is not endogenously

expressed in mice. Dots represent individual samples. Error bars represent ±1 SEM.

Supplementary Fig. 2 | Bodyweight measurements. Percentage of a 2-day average (circled in gray,

horizontal dashed line for reference) of baseline body weight obtained 4 days prior (black epoch) to the

start of the longitudinal Restaurant Row paradigm. During this period (black epoch), mice were food

restricted and fed a limited ration to intentionally decrease weights to approximately 80-85% of baseline

that was steadily maintained throughout the remainder of the experiment (days 1-45, see **Fig. 1** for an

explanation of the color-coded epochs of testing across the paradigm). These body weights reflect the

measurements immediately before task performance on each day of testing. There were no differences in

percentage baseline bodyweight between groups (virus: F=0.292, p=0.593; sex: F=0.118, p=0.733).

Shaded area represents ±1 SEM.

717
Supplementary Fig. 3 | Replication cohort of two different virus titers on sex-specific Restaurant

Row conditioned place preference (CPP) findings. Animals were placed in the arena free to roam for

20 min on day 0 and day 8 of the experimental timeline but with no active task to obtain pre-task baseline

and post-task experience-related exploratory behavior (see **Fig. 2**). **a** Total number of entries and total time

spent in the wait zone, aggregated in all restaurants. Depicting findings from independent cohorts: surgeries

and behavioral experiments were carried out across separate calendar months and using different virus

titers. **b** Delta scores in change of entries and time comparing post (day 8) minus pre (day 0) time points

split by restaurants ranked according to rewards earned on the active task on day 7. **c** Data from (b, only

1st and 4th ranking restaurants) replotted with number of rewards earned on day 7 on the x-axis. Horizontal

dashed gray line indicates delta score of 0. LINC00473 expression in mPFC abolished CPP behavior in

most preferred restaurants only in females and only in the time but not entry domain, replicated in both

cohorts. +Sign-test on delta score of time: low virus cohort: male GFP: ($t=2.142$, $p<0.05$); female GFP:

($t=4.424$, $p<0.01$); female LNC: ($t=0.04$, $p=0.516$); high virus cohort: male GFP: ($t=8.681$, $p<0.0001$); male

LINC: ($t=5.087$, $p<0.001$); female GFP: ($t=7.157$, $p<0.0001$); female LNC: ($t=0.962$, $p=0.181$). +Sign-test

on delta score of entries: low virus cohort: male GFP: ($t=0.955$, $p=0.197$); female GFP: ($t=3.155$, $p<0.05$);

female LNC: ($t=3.766$, $p<0.01$); high virus cohort: male GFP: ($t=3.520$, $p<0.01$); male LINC: ($t=2.493$,

$p<0.05$); female GFP: ($t=3.236$, $p<0.01$); female LNC: ($t=2.355$, $p<0.05$). Error bars represent ±1 SEM.

Supplementary Fig. 4 | Basic Restaurant Row laps and earnings metrics across testing. a Overall

sex difference comparisons between males and females (collapsing across GFP and LINC00473 virus

treatment within each sex). Average number of laps run in the correct direction (sex*day: $F=78.040$,

$p<0.0001$), total rewards earned (sex*day: $F=1.331$, $p=0.249$), ratio of earns / laps (sex*day: $F=28.155$,

$p<0.0001$), and standard deviation of earns among flavors (sex*day: $F=13.014$, $p<0.001$). **b** Data from (a)

split by GFP and LINC groups: laps (sex*virus*day: $F=0.008$, $p=0.927$), total earns (sex*virus*day: $F=7.470$,

$p<0.01$), ratio of earns / laps (sex*virus*day: $F=1.130$, $p=0.288$), standard deviation of earns by flavor

(sex*virus*day: $F=15.486$, $p<0.0001$). Statistical tests from (a) and (b) were performed on the entire

experimental timeline (days 1-45). **c** Data from main Fig. 3d split by sex and virus groups depicting change

in rewards earned in each restaurant comparing two days at each color-coded transition point. Delta scores

calculated by subtracting the following: i, day 8-7 (yellow, rank: $F=1.361$, $p=0.259$; sex*virus: $F=0.017$,

$p=0.897$); ii, day 13-12 (orange, rank: $F=1.123$, $p=0.343$; sex*virus: $F=0.086$, $p=0.769$); iii, day 18-17 (red,

rank: $F=18.116$, $p<0.0001$; sex*virus: $F=0.076$, $p=0.783$); iv, day 45-18 (purple, rank: $F=0.603$, $p=0.614$;

sex*virus: $F=0.375$, $p=0.542$). Shaded area and error bars represent ±1 SEM.

Supplementary Fig. 5 | Restaurant Row choice metrics split by flavor ranking across testing. a

Average number of enter decisions made in the offer zone split by sex and virus treatment groups as well

as the 4th and 1st ranking restaurant (sex*rank*day: $F=16.043$, $p<0.0001$; sex*virus*rank*day: $F=3.967$,

$p<0.01$). **b** Number of skip decisions made in the offer zone (sex*rank*day: $F=9.894$, $p<0.0001$;

sex*virus*rank*day: $F=2.617$, $p<0.05$). **c** Offer zone reaction time before making an enter or skip decision

calculated from offer zone entry (tone onset) to either a right turn into the wait zone (enter) or a left turn into

the hallway (skip) (sex*rank*day: $F=0.325$, $p=0.808$; sex*virus*rank*day: $F=2.186$, $p=0.087$). **d** Number of

quit decisions made in the wait zone during a countdown (sex*rank*day: $F=11.725$, $p<0.0001$;

sex*virus*rank*day: $F=1.216$, $p=0.302$). **e** The probability of quitting in the wait zone given the animal

entered (sex*rank*day: $F=4.145$, $p<0.01$; sex*virus*rank*day: $F=0.463$, $p=0.708$). This normalizes the

likelihood of quitting to enter frequency. **f** Wait zone latency to quit reaction time from countdown onset

(wait zone entry) until mice leave the wait zone prematurely (sex*rank*day: $F=1.427$, $p=0.233$;

sex*virus*rank*day: $F=3.160$, $p<0.05$). Statistical tests here were performed on the entire experimental

timeline (days 1-45). Shaded area represents ±1 SEM.

Supplementary Fig. 6 | Longitudinal economic analysis of decision policy optimization for food

security vs. subjective value. Analysis underlying data from main Fig. 4g-i derived from Durand-de Cuttoli

et al 2023 *Biological Psychiatry* [33]. **a** Computer Restaurant Row simulation of total number of pellets

earned. The ideal threshold required to obtain the theoretical maximum number of pellets when ignoring

flavor preferences was empirically determined to be 10 s. **b** Relative ratio of earns among restaurants for

an example mouse on day 7 between the flavor rankings (i.e., this mouse had a 3.7 : 2.1 : 1.4 : 1 ratio of

earns across flavors capturing a summary of idealized relative subjective value for this mouse when all

offers were 1 s only). **c** Two intersecting planes of decision policies that yield varying amounts of theoretical

value either for maximal food intake (as determined by computer simulations, brown plane) or subjective

value (as determined by multiplying day 7 preference ratios on a mouse-by-mouse basis, pink plane) when

in a reward-scarce environment (1 to 30 s offers). Here, only two decision policy dimensions (least preferred

and most preferred restaurants) of the four dimensions (all four ranked restaurants) are displayed. Both

planes are normalized to each's min and max values for the purpose of plotting both in the same space

while preserving the threshold coordinate locations that achieve either theoretical maximum (brown

[location always fixed at threshold coordinates of 10 s] or pink beacons [location of discovered threshold

coordinates that vary from mouse-to-mouse]). Actual observed daily decision policies represented by the

cyan beacon wander throughout this space from day 18 to 45 in a reward-scarce environment. Trajectories

projected to the floor of this display trace out individual mouse decision policy paths. Example coordinates

of brown, pink, and cyan beacons illustrated as dots on the x and y axes. Euclidean distance from cyan

coordinates to either brown or pink coordinates were calculated (question mark symbols). **d** Euclidean

distances from observed decision policies in the offer zone (left) or wait zone (right) to either food (brown)

or preference (magenta) theoretical maximum across testing in a reward-scarce environment split by GFP

(top) and LINC (bottom) groups and sex. Offer zone: food distance: day: $F=49.362$, $p<0.0001$; sex: $F=1.115$,

$p=0.298$; virus: $F=0.008$, $p=0.929$; value distance: day: $F=42.042$, $p<0.0001$, sex: $F=1.546$, $p=0.221$; virus:

$F=0.001$, $p=0.989$; Wait zone: food distance: day: $F=16.774$, $p<0.001$, sex: $F=13.796$, $p<0.001$; virus:

$F=0.598$, $p=0.445$; value distance: day: $F=38.046$, $p<0.0001$, sex: $F=18.034$, $p<0.0001$; virus: $F=0.261$,

$p=0.613$. Shaded area represents ±1 SEM.

Supplementary Fig. 7 | Sensitivity to sunk costs analysis during the 1-15 s testing epoch. Mice do

not display sensitivity to sunk costs during days 13-17 in a relatively reward rich environment compared to

days 18+ (when offers range 1-30 s, as in main Fig. 5). **a-c** Sunk cost analysis of staying behavior in the

wait zone, demonstrated using all 40 mice. (f) The likelihood of staying in the wait zone and earning a

reward (e.g., not quit) is plotted as a function of time left in the countdown along the x-axis and time already

spent waiting orthogonally in color. Note the black 0 s time spent curve represents animals having just

entered the wait zone from the offer zone. Inset vertical dashed gray line illustrates an example analysis

point comparing three sunk cost conditions originating from different starting offers but matched at 10 s left.

Data from (a) dimensioned reduced in (b) collapsing across time left, highlighting the grand mean of each

time spent sunk cost condition (color and x-axis). Difference between curves in (b) are plotted in (c) in order

to summarize the envelope of the overall effect of time already spent on escalating the commitment of

staying in the wait zone. Horizontal dashed line represents delta score of 0. **d** Data in (c) split by sex and

virus treatment groups depicting no effect of time spent during the countdown in the wait zone on changing

the probability of quitting vs. staying to earn a reward (time spent: $F=0.522$, $p=0.470$; sex*virus*time spent:

$F=0.809$, $p=0.369$). **e** To compare to time spent waiting in the wait zone, data here depict a different type

of time spent on the task: time spent in the offer zone prior to accepting an offer and entering the wait zone

does not influence the probability of quitting vs. staying to earn a reward once in the wait zone (time spent

in offer zone: $F=1.574$, $p=0.210$; sex*virus*time spent: $F=0.916$, $p=0.339$). These data are from the 1-30 s

reward-scarce epoch (days 18-45) to demonstrate the unique value of time spent waiting in the wait zone

uniquely conferring sunk cost sensitivity. Error bars represent ±1 SEM.

Supplementary Fig. 8 | Full set of task information manipulations on Restaurant Row. a After day 45,

mice were challenged with special probe days on three different task informational manipulations each for

2 days. Manipulations took the form of silencing and withholding tone presentation either in the offer zone,

wait zone, or both: (i) silent offer zone but audible wait zone [days 46-47], (ii) audible offer zone but silent

wait zone [days 48-49], and (iii) silent offer zone and silent wait zone [days 50-51]. **b** Offer zone choice

probability to enter as a function of offer cost split by restaurants ranked from least (4th) to most (1st)

preferred separating sex and virus treatment groups (rows) across the 4 different task information

conditions (columns). **c** Data in (b) collapsed across all four restaurants showing summary enter

probabilities as a function of offer cost. Note that enter probabilities are largely unaffected when the offer

zone tones are audible comparing baseline (black, regular task) to silent wait zone (blue) conditions,

however, as intended, offer zone choice probabilities change into a flat horizontal line in both the silent offer

zone (green) and silent task (gray) conditions, as animals do not have any tone information in the offer zone

during these conditions (audible task: offer: $F=2017.115$, $p<0.0001$; sex*virus*offer: $F=1.912$, $p=0.167$;

silent offer zone task: offer: $F=1.783$, $p=0.182$; sex*virus*offer: $F=0.227$, $p=0.634$; silent wait zone task:

offer: $F=1004.552$, $p<0.0001$; sex*virus*offer: $F=0.147$, $p=0.702$; fully silent task: offer: $F=0.723$, $p=0.395$;

sex*virus*offer: $F=1.723$, $p=0.190$). **d** Wait zone thresholds plotted as a relative fold change comparing

each condition to the regular task audible baseline separated across the three information manipulations

(silent offer zone, left, green; silent wait zone, middle, blue; silent task, right, gray) split by restaurants

ranked from least (4th) to most (1st) preferred, sex, and virus treatment groups. Note the difference in

patterns in changes in wait zone thresholds across restaurants between sex/virus groups and across task

information manipulation conditions. Sign tests: silent offer zone: male GFP: least preferred: $t=+1.365$,

$p=0.103$; most preferred: $t=+2.639$, $p<0.05$; male LNC: least preferred: $t=+2.699$, $p<0.05$; most preferred:

$t=+1.538$, $p=0.079$; female GFP: least preferred: $t=+3.638$, $p<0.01$; most preferred: $t=+0.360$, $p=0.363$;

female LNC: least preferred: t=+2.125, p=<0.05; most preferred: t=+0.188, p=0.427; silent wait zone: male

GFP: least preferred: t=-2.684, p<0.05; most preferred: t=+3.191, p<0.01; male LNC: least preferred: t=-

2.624, p<0.05; most preferred: t=+2.685, p<0.05; female GFP: least preferred: t=-4.733, p<0.001; most

preferred: t=+1.325, p=0.109; female LNC: least preferred: t=-0.411, p=0.345; most preferred: t=+0.886,

p=0.199; silent task: male GFP: least preferred: t=-0.358, p=0.364; most preferred: t=+5.554, p<0.001; male

LNC: least preferred: t=+0.794, p=0.225; most preferred: t=+3.049, p<0.001; female GFP: least preferred:

t=-3.584, p<0.01; most preferred: t=+2.345, p<0.05; female LNC: least preferred: t=-1.009, p=0.170; most

preferred: t=+1.671, p=0.064. Shaded area and error bars represent ±1 SEM.

Methods

Subjects: Adult male and female C57BL/6J mice (Jackson, 10wks, n=40) were used. Mice

were individually housed and maintained on a 12-hr light/dark cycle with ad libitum water, only

food-restricted during Restaurant Row testing, conducted during their light phase. Experiments

were approved by the Mount Sinai Institutional Animal Care and Use Committee (IACUC; protocol

number LA12-00051) and adhered to the National Institutes of Health (NIH) guidelines.

Viral constructs: To achieve long-term neuronal expression of LINC00473, it was

subcloned into an AAV plasmid with human synapsin promoter (hSyn) driving gene expression.

LINC00473 transcript variant 1, (NR_026860.1 /AK289375) of 1821bp length was subcloned from

p1005+ HSV vector (PMID: 32304628) into pAAV-hSyn-eGFP-hSny plasmid (PMID: 36449610)

using blunt ligation into a PMEI and EcoRV restriction sites following the second hSyn promoter.

The orientation and sequence were validated using Sanger sequencing. Empty plasmid

expressing eGFP only was used as control. These vectors were used to generate AAV serotype

2 at the University of Maryland virus vector core. Viral vector placement and expression were

confirmed in vivo by a GFP fluorescent flashlight and qPCR.

RNA extraction, RT, and qPCR: Total RNA was isolated and purified with RNeasy Micro

Kit (Qiagen), using QIAzol as a lysis reagent, including DNase treatment. All samples were tested

for concentration and purity using a NanoDrop (Thermo Fisher). RNA amount was normalized

across samples (400ng), and cDNA was created using iScript (Bio-Rad). Real-time PCR reactions

were run in triplicate using PowerUp SYBR Master Mix (Applied Biosystems) in a QuantStudio 7

(Thermo   Fisher)   qPCR   machine.   Primer   sequences: LINC00473:   forward:

GCATACTTTGGCGGACCTTTT,   reverse   TGTGCCTCCCTGTGAATTCTC;   HPRT1   forward:

GCAGTACAGCCCCAAAATGG,   reverse: GGTCCTTTTCACCAGCAAGCT;   GFP   forward:

CATGCCCGAAGGCTACGT,   reverse   CGATGCCCTTCAGCTCGAT.   Raw   Ct   values   are

presented and not normalized amounts, as LINC00473 is not endogenously expressed in mice.

Virus surgery: Mice were anesthetized by intraperitoneal injection with a mixture of

Ketamine HCl (100 mg/kg) and Xylazine (10 mg/kg) and positioned on a stereotaxic instrument

(David Kopf Instruments). In the medial prefrontal cortex (mPFC, from bregma with an angle of

15 degrees: AP +1.8 mm; ML ±0.75 mm; DV −2.7 mm), 0.7 µL of virus (AAV2-hSyn-GFP or AAV2-

hSyn-LINC00473-GFP) was bilaterally infused using 33-Gauge Hamilton needles over 5 min at a

rate of 0.1 µL/min, and the needle was left in place for 10-15 min after the injection before

retraction. Mice were allotted 2 weeks to recover before beginning food restriction across 3-4 days

in preparation for testing on Restaurant Row. An initial virus validation cohort (independent cohort

of n=10 mice) was injected with GFP or LINC00473 viruses bilaterally in mPFC at a titer of 1×10^{12}

at. These animals were tested on Restaurant Row for only 1 week (offers remained at 1 s only,

first epoch, green) after which they were sacrificed for PCR validation on tissue punches obtained

from mPFC. Behavioral data from these animals on the conditioned-place preference probe (days

0 and 8) appear in Supplementary Fig. 3. A second replication cohort (n=40 mice) was injected

with GFP or LINC00473 viruses bilaterally in mPFC at a titer of 5×10^{12} and served as the primary

cohort of animals for the main figures that underwent the full 45-day longitudinal behavioral

testing.

Neuroeconomic Task: Following surgery, we characterized mice longitudinally in

Restaurant Row [5]. On this task, mice had a limited time period each day to forage for food by

navigating a maze with four uniquely flavored (chocolate, banana, grape, and plain) and

contextualized (horizontal stripes, dots, triangles, vertical stripes) feeding sites, or "restaurants"

(Fig. 1b). Animals were allotted 30 min to forage for their primary source of food for the day.

Rewards used on this task were obtained from BioServe (20 mg pellets #F007120). These pellets

are full nutrition and not sucrose-only pellets that only differed in flavoring but not nutritional profile

(3.6 kcal/gm calories, 59.1% carbohydrate, 18.7% protein, 6.5% ash, 5.6% fat, 4.7% fiber).

Choices on this task were interdependent across trials and across days. This makes the task

economic in nature, requiring animals to budget their limited time effectively in a self-paced

manner in order to earn a sufficient amount of food in a closed-economy system. The cost of

earning a reward took place in the form of a delay animals would have to wait cued by the pitch

of a tone that sounded upon entry into each restaurant. Each restaurant had a separate offer zone

(OZ) and wait zone (WZ). Upon entry into the OZ, a tone (500 ms) sounded whose pitch indicated

how long of a delay mice would have to wait in a cued countdown should they enter the WZ. Mice

could decide to skip an offer presented in the OZ and proceed down the hallway to the next

restaurant. In the OZ, tones repeated every second held at the same pitch to repeatedly cue the

offer of the current trial until mice either made a skip or enter decision. If mice decide instead to

enter the WZ, this triggered a countdown with descending tone pitch stepping every second until

mice either waited out the full delay and earned a food pellet or quit prematurely during the

countdown. If mice made a skip or quit decision, the tone terminated, the offer was rescinded,

and mice must proceed forward to the next restaurant. Mice quickly learned to only travel in one

direction as animals must encounter offers serially navigating the maze in a counterclockwise

direction (i.e., nothing happens if mice traverse in the wrong direction other than wasting time).

Mice were tested across 45 consecutive days for the main experiment where strategies developed

longitudinally as animals learned the structure of the task and the changing economic landscape.

During the first week of testing (block 1, green epoch), all trials consisted of 1 s offers only.

1 s offers were indicated by tones played at the 4,000 Hz pitch. A special probe session was

inserted into this timeline immediately before (day 0) and after (day 8) the first week of testing

during which mice essentially were allowed to roam the empty maze arena freely without any task

components (no tones, no rewards) to examine conditioned place preference behaviors learned

through the natural experience of the first week of the task itself. The remainder of the

experimental timeline continued thereafter. During the next 5 days of testing (block 2, yellow

epoch), offers ranged from 1 to 5 s (4,000 Hz to 5,548 Hz in 387 Hz steps) randomly selected

from a uniform distribution. For instance, if a 4 s offer was selected on a given trial, the tone

presented in the OZ would be 5,161 Hz for 500 ms, and this would repeat every second indefinitely

until a skip or enter decision was made. If the WZ was entered, using this example trial, this 5,161

Hz tone would step down by 387 Hz every second descending in pitch until earning (with the final

tone of 4,000 Hz) or quitting prematurely. Block 3 (5 days, orange epoch) consisted of a 1 to 15

s (4,000–9,418 Hz) range. The fourth and final block (red epoch) consisted of offers ranging from

1 to 30 s (4,000–15,223 Hz) and continued for the remainder of the experiment. This block-wise

protocol allowed us to examine how mice initially shaped their behavior as they learned the basic

structure of the task as well as how they adjusted their foraging strategies across long timescales

(days, weeks, and months) as they transitioned from reward-rich to reward-scarce environments

in a stepwise manner. That is, blocks 1-3 are relatively reward rich as costs are generally low

delays and rewards are easier to earn compared to block 4 which is relatively reward scarce as

there are far more expensive reward offers distributed in the environment for which animals must

learn to forage efficiently. The rewards used in this task only differ in flavor for which animals have

subjective differences in preferences quantified below. All flavor options become increasingly

scarce, equally across the longitudinal paradigm.

Rewards were delivered using a custom-built 3D-printed automated pellet dispenser that

was triggered by a computer running the task programmed in ANY-Maze (Stoelting), run under

dim lighting. The task was run by video-tracking animal position in real-time where all choice

contingencies are determined by the centroid of the animal's body crossing territories in the maze

in combination with being required to run in a counterclockwise direction. The key zones in the

maze include each restaurant's separate OZ and WZ as well as the hallways in between

restaurants (thus, 12 total main zones). 6 key behavioral epochs make up all possible, mutually

exclusive behaviors animals could be engaged in while spending their limited time budget. 1)

Enter time: defined from OZ entry in the correct heading direction (counterclockwise around the

maze) triggering tone onset until the animal physically enters right into the WZ triggering tone

countdown. 2) **Skip time**: defined from OZ entry in the correct heading direction triggering tone

onset until the animal physically exits left into the hallway. 3) **Quit time**: defined from WZ entry

and tone countdown onset until the animal physically exits the WZ prematurely before the

countdown completes, rescinding the offer and terminating the countdown. 4) **Earn time**: defined

from WZ entry and tone countdown onset until the countdown completes and a pellet is delivered.

5) **Consumption time**: defined from reward delivery onset until the animal physically leaves the

reward site and exits the WZ. 6) **Travel time**: defined from previous trial termination (either from

a skip, quit, or consumption event as defined above) and hallway entry until the next trial begins,

triggered by OZ entry into the next restaurant.

Body weight was measured twice daily, immediately before and after testing on the

Restaurant Row task.

Data analysis: In addition to simple behavioral metrics like rewards earned, laps run,

reaction times as described above, a key neuroeconomic behavioral metric is decision thresholds.

This metric approximates indifference points in choices made either in the OZ or WZ as a function

of cost and flavor. We measured the inflection point of fitting heaviside step regressions to enter

versus skip choices in the OZ or earn versus quit choices in the WZ as a function of cued offer

cost in each restaurant. Thresholds determined this way were used for several subsequent

analyses, including calculating reward value. The value of an offer can be calculated by

subtracting a trial's offer from one's WZ threshold as this reflects the relative cost normalized to

each animals' willingness to wait and earn a reward (i.e., offer value <0 indicate offers typically

not earned and thus should be skipped in the OZ or quit in the WZ as determined by each animal's

subjective flavor-specific decision policy which remained relatively stable across block 4 of

testing). Similarly, when quitting in the WZ, the value left at the moment of quitting can be

calculated by subtracting the time remaining in that trial's countdown from the WZ threshold (i.e.,

value left <0 indicates an amount of time left that is not worth waiting for and should have been

quit whereas value left >0 would have been more advantageous to finish waiting at that point

instead of quitting). The amount of time spent remaining in the countdown at the moment of

quitting relative to what the starting offer was and how much time was spent also served as the

basis of our analysis of sensitivity to sunk costs, as previously published in mice, rats, and humas

whereby time spent waiting can decrease the probability of quitting independent of time left in the

countdown, see Sweis et al *Science* for the development of this analysis [44, 46, 48, 49, 53].

Thresholds as well as economic choice history serve as the basis and criteria used in regret-

related analysis whereby distinct violation vs. non-violation sequences are compared head-to-

head in order to quantify the effects of mistake history on influencing choice behavior on the next

trial. See Steiner and Redish 2014 *Nature Neuroscience* [47], Sweis et al 2018 *PloS Biology* [5],

and Durand-de Cuttoli 2022 *Science Advances* [48] for the development of these analyses.

Thresholds were also used for an analysis we recently developed to capture how mice develop

strategies that optimize food yield while also maximizing subjective value, see Durand-de Cuttoli

et al 2023 *Biological Psychiatry* [33] for the development of this analysis.

Data were processed in Matlab with statistical analyses in JMP Pro 17. All data are

expressed as mean ± 1 standard error. Statistical significance was assessed using student's t

tests, sign tests, and one-way, two-way, three-way, and repeated measures ANOVAs. No data

were excluded as outliers.

References

1.   Stone, C., J.B. Mattingley, and D. Rangelov, *On second thoughts: changes of mind in*
1001   *decision-making.* Trends Cogn Sci, 2022. **26**(5): p. 419-431.

1002   2.   Liu, X., et al., *Common and distinct networks underlying reward valence and processing*
1003   *stages: a meta-analysis of functional neuroimaging studies.* Neurosci Biobehav Rev,
1004   2011. **35**(5): p. 1219-36.

1005   3.   Redish, A.D., *The mind within the brain: How we make decisions and how those decisions*
1006   *go wrong.* 2013: Oxford University Press.

1007   4.   McGuire, J.T. and J.W. Kable, *Medial prefrontal cortical activity reflects dynamic re-*
1008   *evaluation during voluntary persistence.* Nat Neurosci, 2015. **18**(5): p. 760-6.

1009   5.   Sweis, B.M., M.J. Thomas, and A.D. Redish, *Mice learn to avoid regret.* PLoS Biol, 2018.
1010   **16**(6): p. e2005853.

1011   6.   Kirkebøen, G., E. Vasaasen, and K. Halvor Teigen, *Revisions and Regret: The Cost of*
1012   *Changing your Mind.* Journal of Behavioral Decision Making, 2011. **26**(1): p. 1-12.

1013   7.   Brenner, C., *The mind in conflict.* 1982, New York: International Universities Press. 266 p.

1014   8.   Baumeister, R.F., K.D. Vohs, and D.M. Tice, *The Strength Model of Self-Control.* Current
1015   Directions in Psychological Science, 2016. **16**(6): p. 351-355.

1016   9.   American Psychiatric Association. and American Psychiatric Association. DSM-5 Task
1017   Force., *Diagnostic and statistical manual of mental disorders : DSM-5*. 5th ed. 2013,
1018   Washington, D.C.: American Psychiatric Association. xliv, 947 p.

1019   10.   Mezo, P.G. and R.M. Baker, *The moderating effects of stress and rumination on*
1020   *depressive symptoms in women and men.* Stress Health, 2012. **28**(4): p. 333-9.

1021   11.   Seedat, S., et al., *Cross-National Associations Between Gender and Mental Disorders in*
1022   *the World Health Organization World Mental Health Surveys.* Archives of General
1023   Psychiatry, 2009. **66**(7).

12.   Ferrari, A.J., et al., *Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010.* PLoS Med, 2013. **10**(11): p. e1001547.

13.   Marcus, S.M., et al., *Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study.* Compr Psychiatry, 2008. **49**(3): p. 238-46.

14.   Silverstein, B., *Gender differences in the prevalence of somatic versus pure depression: a replication.* Am J Psychiatry, 2002. **159**(6): p. 1051-2.

15.   Martin, L.A., H.W. Neighbors, and D.M. Griffith, *The experience of symptoms of depression in men vs women: analysis of the National Comorbidity Survey Replication.* JAMA Psychiatry, 2013. **70**(10): p. 1100-6.

16.   Eid, R.S., A.R. Gobinath, and L.A.M. Galea, *Sex differences in depression: Insights from clinical and preclinical studies.* Prog Neurobiol, 2019. **176**: p. 86-102.

17.   Akil, H., et al., *Treatment resistant depression: A multi-scale, systems biology approach.* Neurosci Biobehav Rev, 2018. **84**: p. 272-288.

18.   Ulitsky, I. and D.P. Bartel, *lincRNAs: genomics, evolution, and mechanisms.* Cell, 2013. **154**(1): p. 26-46.

19.   Issler, O., et al., *Sex-Specific Role for the Long Non-coding RNA LINC00473 in Depression.* Neuron, 2020. **106**(6): p. 912-926 e5.

20.   Chen, Z., et al., *cAMP/CREB-regulated LINC00473 marks LKB1-inactivated lung cancer and mediates tumor growth.* J Clin Invest, 2016. **126**(6): p. 2267-79.

21.   Jiang, L.Y., et al., *CREB-induced LINC00473 promotes chemoresistance to TMZ in glioblastoma by regulating O6-methylguanine-DNA-methyltransferase expression via CEBPalpha binding.* Neuropharmacology, 2024. **243**: p. 109790.

22.   Ataman, B., et al., *Evolution of Osteocrin as an activity-regulated factor in the primate brain.* Nature, 2016. **539**(7628): p. 242-247.

23.   Pruunsild, P., et al., *Expression of the primate-specific LINC00473 RNA in mouse neurons*

promotes excitability and CREB-regulated transcription. J Biol Chem, 2023. **299**(5): p.

104671.

24.   Pruunsild, P., C.P. Bengtson, and H. Bading, *Networks of Cultured iPSC-Derived Neurons*

Reveal the Human Synaptic Activity-Regulated Adaptive Gene Program. Cell Rep, 2017.

18(1): p. 122-135.

25.   Tran, K.V., et al., *Human thermogenic adipocyte regulation by the long noncoding RNA*

LINC00473. Nat Metab, 2020. **2**(5): p. 397-412.

26.   Glimcher, P.W. and A. Rustichini, *Neuroeconomics: the consilience of brain and decision.*

Science, 2004. **306**(5695): p. 447-52.

27.   Kenning, P. and H. Plassmann, *Neuroeconomics: an overview from an economic*

perspective. Brain Res Bull, 2005. **67**(5): p. 343-54.

28.   Loewenstein, G., S. Rick, and J.D. Cohen, *Neuroeconomics.* Annu Rev Psychol, 2008.

59: p. 647-72.

29.   Kalenscher, T. and C.M. Pennartz, *Is a bird in the hand worth two in the future? The*

neuroeconomics of intertemporal decision-making. Prog Neurobiol, 2008. **84**(3): p. 284-

315.

30.   Clithero, J.A. and A. Rangel, *Informatic parcellation of the network involved in the*

computation of subjective value. Soc Cogn Affect Neurosci, 2014. **9**(9): p. 1289-302.

31.   Levy, D.J. and P.W. Glimcher, *The root of all value: a neural common currency for choice.*

Curr Opin Neurobiol, 2012. **22**(6): p. 1027-38.

32.   Louie, K. and P.W. Glimcher, *Efficient coding and the neural representation of value.* Ann

N Y Acad Sci, 2012. **1251**: p. 13-32.

33.   Durand-de Cuttoli, R., et al., *A double-hit of social and economic stress in mice precipitates*

changes in decision-making strategies. Biol Psychiatry, 2023.

34. Resulaj, A., et al., *Changes of mind in decision-making.* Nature, 2009. **461**(7261): p. 263-6.

35. Hare, T.A., C.F. Camerer, and A. Rangel, *Self-control in decision-making involves modulation of the vmPFC valuation system.* Science, 2009. **324**(5927): p. 646-8.

36. Alagapan, S., et al., *Cingulate dynamics track depression recovery with deep brain stimulation.* Nature, 2023. **622**(7981): p. 130-138.

37. Knutson, B., et al., *Neural responses to monetary incentives in major depression.* Biol Psychiatry, 2008. **63**(7): p. 686-92.

38. Holton, E., et al., *Goal commitment is supported by vmPFC through selective attention.* Nature Human Behaviour, 2024.

39. Sweis, B.M. and E.J. Nestler, *Pushing the boundaries of behavioral analysis could aid psychiatric drug discovery.* PLoS Biol, 2022. **20**(12): p. e3001904.

40. Sweis, B.M., M.J. Thomas, and A.D. Redish, *Beyond simple tests of value: measuring addiction as a heterogeneous disease of computation-specific valuation processes.* Learn Mem, 2018. **25**(9): p. 501-512.

41. Abram, S.V., et al., *The Web-Surf Task: A translational model of human decision-making.* Cogn Affect Behav Neurosci, 2016. **16**(1): p. 37-50.

42. Maisson, D., et al., *Widespread coding of navigational variables in prefrontal cortex* biorxiv, 2022.

43. McInnes, A.N., et al., *Psychometric validation and clinical correlates of an experiential foraging task.* BioRxiv, 2023. **2023.12.28.573439**.

44. Kazinka, R., A.W. MacDonald, 3rd, and A.D. Redish, *Sensitivity to Sunk Costs Depends on Attention to the Delay.* Front Psychol, 2021. **12**: p. 604843.

45. Abram, S.V., et al., *Neural signatures underlying deliberation in human foraging decisions.* Cogn Affect Behav Neurosci, 2019. **19**(6): p. 1492-1508.

1099   46.   Sweis, B.M., et al., *Sensitivity to "sunk costs" in mice, rats, and humans.* Science, 2018.

1100   **361**(6398): p. 178-181.

1101   47.   Steiner, A.P. and A.D. Redish, *Behavioral and neurophysiological correlates of regret in*

1102   *rat decision-making on a neuroeconomic task.* Nat Neurosci, 2014. **17**(7): p. 995-1002.

1103   48.   Durand-de Cuttoli, R., et al., *Distinct forms of regret linked to resilience versus*

1104   *susceptibility to stress are regulated by region-specific CREB function in mice.* Science

1105   Advances, 2022. **8**(42): p. eadd5579.

1106   49.   Redish, A.D., et al., *Sunk cost sensitivity during change-of-mind decisions is informed by*

1107   *both the spent and remaining costs.* Commun Biol, 2022. **5**(1): p. 1337.

1108   50.   Sweis, B.M., et al., *Altering gain of the infralimbic-to-accumbens shell circuit alters*

1109   *economically dissociable decision-making algorithms.* Proc Natl Acad Sci U S A, 2018.

1110   **115**(27): p. E6347-E6355.

1111   51.   Sweis, B.M., A.D. Redish, and M.J. Thomas, *Prolonged abstinence from cocaine or*

1112   *morphine disrupts separable valuations during decision conflict.* Nat Commun, 2018. **9**(1):

1113   p. 2521.

1114   52.   Diehl, G. and A.D. Redish, *Differential processing of decision information in subregions of*

1115   *rodent medial prefrontal cortex.* biorxiv, 2022.

1116   53.   Duin, A.A., et al., *Certainty and uncertainty of the future changes planning and sunk costs.*

1117   Behav Neurosci, 2021.

1118   54.   Schmidt, B., A.A. Duin, and A.D. Redish, *Disrupting the medial prefrontal cortex alters*

1119   *hippocampal sequences during deliberative decision making.* J Neurophysiol, 2019.

1120   **121**(6): p. 1981-2000.

1121   55.   Schmidt, B. and A.D. Redish, *Disrupting the medial prefrontal cortex with designer*

1122   *receptors exclusively activated by designer drug alters hippocampal sharp-wave ripples*

1123   *and their associated cognitive processes.* Hippocampus, 2021.

56.   Abram, S.V., A.D. Redish, and A.W. MacDonald, 3rd, *Learning From Loss After Risk:*
Dissociating Reward Pursuit and Reward Valuation in a Naturalistic Foraging Task. Front
Psychiatry, 2019. **10**: p. 359.

57.   Vialou, V., et al., *DeltaFosB in brain reward circuits mediates resilience to stress and*
antidepressant responses. Nat Neurosci, 2010. **13**(6): p. 745-52.

58.   German, P.W. and H.L. Fields, *Rat nucleus accumbens neurons persistently encode*
locations associated with morphine reward. J Neurophysiol, 2007. **97**(3): p. 2094-106.

59.   Sjulson, L., et al., *Cocaine Place Conditioning Strengthens Location-Specific*
Hippocampal Coupling to the Nucleus Accumbens. Neuron, 2018. **98**(5): p. 926-934 e5.

60.   Nwakama, C.A., et al., *Diabetes alters neuroeconomically dissociable forms of mental*
accounting. bioRxiv, 2024.

61.   Camille, N., et al., *The involvement of the orbitofrontal cortex in the experience of regret.*
Science, 2004. **304**(5674): p. 1167-70.

62.   Coricelli, G., et al., *Regret and its avoidance: a neuroimaging study of choice behavior.*
Nat Neurosci, 2005. **8**(9): p. 1255-62.

63.   Coricelli, G., R.J. Dolan, and A. Sirigu, *Brain, emotion and decision making: the*
paradigmatic example of regret. Trends Cogn Sci, 2007. **11**(6): p. 258-65.

64.   Bardo, M.T. and R.A. Bevins, *Conditioned place preference: what does it add to our*
preclinical understanding of drug reward? Psychopharmacology, 2000. **153**(1): p. 31-43.

65.   Tzschentke, T.M., *Measuring reward with the conditioned place preference (CPP)*
paradigm: update of the last decade. Addict Biol, 2007. **12**(3-4): p. 227-462.

66.   Johnson, A. and A.D. Redish, *Neural ensembles in CA3 transiently encode paths forward*
of the animal at a decision point. J Neurosci, 2007. **27**(45): p. 12176-89.

67.   Kay, K., et al., *Constant Sub-second Cycling between Representations of Possible*
Futures in the Hippocampus. Cell, 2020. **180**(3): p. 552-567 e25.

68.   Huynh, T., et al., *Vicarious Trial-and-Error Is Enhanced During Deliberation in Human*
1150   *Virtual Navigation in a Translational Foraging Task.* Front Behav Neurosci, 2021. **15**: p.
1151   586159.

1152   69.   Diehl, G. and A.D. Redish, *Measuring excitation-inhibition balance through spectral*
1153   *components of local field potentials.* bioRxiv, 2024.

1154   70.   Kocharian, A., *Dopaminergic control of neuroeconomic decision-making.* ProQuest,
1155   University of Minnesota, 2023.

1156   71.   Lind, E.B., et al., *A quadruple dissociation of reward-related behaviour in mice across*
1157   *excitatory inputs to the nucleus accumbens shell.* Commun Biol, 2023. **6**(1): p. 119.

1158   72.   Barrot, M., et al., *CREB activity in the nucleus accumbens shell controls gating of*
1159   *behavioral responses to emotional stimuli.* Proc Natl Acad Sci U S A, 2002. **99**(17): p.
1160   11435-40.

1161   73.   Carlezon, W.A., Jr., R.S. Duman, and E.J. Nestler, *The many faces of CREB.* Trends
1162   Neurosci, 2005. **28**(8): p. 436-45.

1163   74.   Kim, J., et al., *Memory recall and modifications by activating neurons with elevated CREB.*
1164   Nat Neurosci, 2014. **17**(1): p. 65-72.

1165   75.   Lonze, B.E. and D.D. Ginty, *Function and regulation of CREB family transcription factors*
1166   *in the nervous system.* Neuron, 2002. **35**(4): p. 605-23.

1167   76.   McClung, C.A. and E.J. Nestler, *Regulation of gene expression and cocaine reward by*
1168   *CREB and DeltaFosB.* Nat Neurosci, 2003. **6**(11): p. 1208-15.

1169   77.   Sakai, N., et al., *Inducible and brain region-specific CREB transgenic mice.* Mol
1170   Pharmacol, 2002. **61**(6): p. 1453-64.

1171   78.   Lorsch, Z.S., et al., *Stress resilience is promoted by a Zfp189-driven transcriptional*
1172   *network in prefrontal cortex.* Nat Neurosci, 2019. **22**(9): p. 1413-1423.

1173   79.   Durand-de Cuttoli, R., et al., *Chronic social stress induces isolated deficits in reward*
1174   *anticipation on a neuroeconomic foraging task.* biorxiv, 2022.

80.   Barrot, M., et al., *Regulation of anxiety and initiation of sexual behavior by CREB in the*
nucleus accumbens. Proc Natl Acad Sci U S A, 2005. **102**(23): p. 8357-62.

81.   Dong, Y., et al., *CREB modulates excitability of nucleus accumbens neurons.* Nat
Neurosci, 2006. **9**(4): p. 475-7.

82.   Zhou, J., et al., *Effects of Estrogen Treatment on Expression of Brain-Derived*
Neurotrophic Factor and cAMP Response Element-Binding Protein Expression and
Phosphorylation in Rat Amygdaloid and Hippocampal Structures. Neuroendocrinology,
2005. **81**(5): p. 294-310.

83.   Sharma, K., et al., *Chronic exposure to estrogen and tamoxifen regulates synaptophysin*
and phosphorylated cAMP response element-binding (CREB) protein expression in CA1
of ovariectomized rat hippocampus. Brain Res, 2007. **1132**(1): p. 10-9.

84.   Lorsch, Z.S., et al., *Estrogen receptor alpha drives pro-resilient transcription in mouse*
models of depression. Nat Commun, 2018. **9**(1): p. 1116.

85.   Shansky, R.M., et al., *Estrogen mediates sex differences in stress-induced prefrontal*
cortex dysfunction. Mol Psychiatry, 2004. **9**(5): p. 531-8.

86.   Issler, O., et al., *The long noncoding RNA FEDORA is a cell type– and sex-specific*
regulator of depression. Science Advances, 2022. **8**(48).

1192

Figure 1

a surgery & recovery | day: | longitudinal neuroeconomic paradigm

-17 | 1 | 8 | 13 | 18 | 45

AAV-GFP vs. AAV-LINC00473 | 7 days | 5 days | 5 days | 28 days

target: medial prefrontal cortex

male C57BL/6J N=20 | female C57BL/6J N=20

1s only | 1-5s | 1-15s | 1-30s offer range

distribution of wait time (costs)

b Restaurant Row Task

plain

offer zone | wait zone

3 3 3 | 3 2 1

quit
earn
grape

skip | enter

must forage for offers in serial order

chocolate

➤ 30 min daily budget
➤ primary source of food
➤ cost indicated by tone pitch
➤ subjective flavor preferences

banana

c GFP | LINC
distance traveled (m) | 250 ... 50 | ♂ ♀

d laps correct direction | 40 ... 0

e total rewards earned | 90 ... 0

day 1 ... 7 | day 1 ... 7

f GFP
4th rank | 3rd rank | 2nd rank | 1st rank
earns by flavor | 40 ... 0 | ♂ ♀
day 1 ... 7

LINC
4th rank | 3rd rank | 2nd rank | 1st rank
day 1 ... 7

Figure 2

longitudinal neuroeconomic paradigm

a day: 1   7   45

relevant experimental timeline   0   7 days   8

1s only

RRow CPP
pre   post

b — **RRow Conditioned Place Preference** —

pre test → *post test*

NO TASK   7 days of full task   NO TASK

1s only

day 0   days 1 - 7   day 8

c GFP   LINC

♂ ♀

entries in zone — entries: 135 / 75

time in zone — time (s): 1000 / 550

pre   post   pre   post

d plain   grape

chocolate   *pre*   example mouse GFP ♀   *post*   banana

distance traveled: 135.182 m
average speed: 14.2 cm/s
max speed: 78.5 cm/s

distance traveled: 177.657 m
average speed: 17.9 cm/s
max speed: 107.3 cm/s

e plain   grape

chocolate   *pre*   example mouse GFP ♀   *post*

banana   Δ time (s) →   time (s): 130 / 0

f GFP   LINC

♂ ♀

Δ entries [post - pre]: 17 / -7

1st rank

4th rank

0   day 7 task earns   35   0   day 7 task earns   35

g GFP   LINC

♂ ♀

Δ time (s) [post - pre]: 160 / -30

*

0   day 7 task earns   35   0   day 7 task earns   35

Figure 3

a longitudinal neuroeconomic paradigm

day: 1   8   13   18   45

relevant experimental timeline | 7 days | 5 days | 5 days | 28 days

distribution of wait time (costs) | 1s only | 1-5s | 1-15s | 1-30s offer range

b all 40 mice

c all 40 mice — earns by flavor — 1st, 2nd, 3rd, 4th — i ii iii iv

d all 40 mice — Δ earns — transition — iv recovery, i 1-5 s, ii 1-15 s, iii 1-30 s — 4th rank 1st

e session time — quit, earn, enter, skip — rank — 1s only, 1-5s, 1-15s, 1-30s offer range — early, late — all 40 mice

f GFP   LINC — laps — ♂ ♀ — day

g GFP   LINC — p(quit | enter) — ♂ ♀ — day

Figure 4

Figure 5

Figure 6

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4

Supplementary Fig. 8