Full title:

Sex differences in change-of-mind neuroeconomic decision-making is modulated by LINC00473 in medial

prefrontal cortex

Short title:

Neuroeconomic decision-making in males and females

Authors:

Romain Durand-de Cuttoli[1], Orna Issler[2], Benjamin Yakubov[3], Nusrat Jahan[3], Aisha Abid[3], Susanna

Kasparov[3], Kastalia Granizo[3], Sanjana Ahmed[3], Scott J. Russo[1], Eric J. Nestler[1,4], Brian M. Sweis*[1,4]

*Corresponding: brian.sweis@mountsinai.org

Affiliations:

[1] Nash Family Department of Neuroscience, Friedman Brain Institute, Icahn School of Medicine at Mount

Sinai, New York, NY 10029

[2] Neuroscience Institute, Department of Anesthesiology, Perioperative Care and Pain Medicine, and

Department of Neuroscience and Physiology, NYU Langone, New York, NY 10016

[3] CUNY Hunter, New York, NY 10023

[4] Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY 10029

Abstract:

Changing one's mind is a complex cognitive phenomenon involving a continuous re-appraisal of the trade-off between past costs and future value. Recent work modeling this behavior across species has established associations between aspects of this choice process and their contributions to altered decision-making in psychopathology. Here, we investigated the actions in medial prefrontal cortex (mPFC) neurons of long intergenic non-coding RNA, LINC00473, known to induce stress resilience in a striking sex-dependent manner, but whose role in cognitive function is unknown. We characterized complex decision-making behavior in male and female mice longitudinally in our neuroeconomic foraging paradigm, Restaurant Row, following virus-mediated LINC00473 expression in mPFC neurons. On this task, mice foraged for their primary source of food among varying costs (delays) and subjective value (flavors) while on a limited time-budget during which decisions to accept and wait for rewards were separated into discrete stages of primary commitments and secondary re-evaluations. We discovered important differences in decision-making behavior between female and male mice. LINC00473 expression selectively influenced multiple features of re-evaluative choices, without affecting primary decisions, in female mice only. These behavioral effects included changing how mice (i) cached the value of the passage of time and (ii) weighed their history of economically disadvantageous choices. Both processes were uniquely linked to change-of-mind decisions and underlie the computational bases of distinct aspects of counterfactual thinking. These findings reveal a key bridge between a molecular driver of stress resilience and psychological mechanisms underlying sex-specific decision-making capabilities.

Introduction:

How an individual changes one's mind when making choices is a challenging yet important phenomenon that has been gaining more traction recently in the field of decision science [1]. This mental process involves complex interactions between cognitive and affective systems and presents an opportunity to peer into how the brain represents internal conflict [2, 3]. This captures conflict at multiple levels, including the economic level: between competing interests of managing ongoing expenditures while minimizing potential losses vs. maximizing future gains [4], as well as the psychological level: re-evaluating one's own recent actions, which could be perceived as egodystonic [5-7]. Aspects of these processes are likely dysfunctional in neuropsychiatric illnesses and may manifest as the cognitive changes observed in mood disorders like depression, considering these mental operations could contribute to affective burden [8, 9]. While the neural mechanisms underlying change-of-mind decision-making remain underexplored, expanding this line of investigation holds promise to enrich our understanding of the repertoire of behavioral processes at play in depression.

Depression is a debilitating mental illness characterized by pervasive mood disturbances, altered cognitive capabilities, and maladaptations in motivated behavior [9]. The incidence of depression is two times higher in women than in men and can manifest different symptomatology, including negative rumination [10]; however, the molecular mechanisms mediating these sex differences are unclear [11-15]. Progress is being made toward appreciating neuroanatomical and pathophysiological differences between depressed males and females based on advances in clinical and preclinical research, with the latter enabling mechanistic studies of sex as a biological factor [16, 17]. Such advances are also due to adopting novel approaches that leverage unsupervised analyses of large datasets in emerging fields of systems biology leading to breakthroughs, for example, in identifying key regulatory roles of non-coding elements of the genome [18].

Recent transcriptomics work carried out in human postmortem brain tissue samples obtained from subjects carrying a diagnosis of major depressive disorder discovered sex-dependent functions of a

primate-specific long intergenic non-coding RNA: LINC00473 [19]. Overall, long non-coding RNAs

represent a significant percentage of all of the regulated transcripts in human depression and are regulated

in a sex- and brain region-specific manner. [19]. Lower LINC00473 expression levels were found in the

medial prefrontal cortex (mPFC) of depressed females only [19]. Cross-species validation studies in rodents

demonstrated expressing LINC00473 in mPFC neurons promotes resilience to stress in females only [19].

This included ameliorating depressive- and anxiety-like behaviors while blunting hypothalamic-pituitary-

adrenal-axis activity as well as the induction of gene expression changes induced by chronic stress [19].

Other studies have implicated LINC00473 in molecular scaffolding, inter-organelle communication, and

transcriptional regulation in non-neuronal tissue, while exhibiting immediate early gene-like properties in

cultured neurons [20-25]. Beyond this, LINC00473 has not been studied in neural tissue in any other

capacity, with no work performed to date on the role of long non-coding RNAs in regulating decision-making

information processing.

Strides made in the field of neuroeconomics, an area focused on understanding how the brain is

comprised of multiple valuation algorithms, have leveraged complex approaches in decision neuroscience

to quantify how circuits constrain behavior [26-29]. This encompasses characterizing multifactorial aspects

of motivation that operationalize reward value along several dimensions, integrate choice processes with

environmental circumstances, and segregate stages of a decision into its component parts [30-33]. This

work includes isolating properties of change-of-mind re-evaluative decisions, and points to the mPFC as a

critical hub for regulating reward value, self-control, and mood [4, 34-38]. Insights from neuroeconomic

principles offer novel approaches to investigate decision-making capable of resolving behavior into

discretely measurable computational units in a manner that is biologically tractable and readily translatable

across species [30, 39, 40]. This approach has yielded efforts from our team and others developing

innovative tasks adapted for use in mice, rats, monkeys, and humans [5, 41-44]. The use of these

translational tasks have led to cross-species discoveries uncovering the behavioral underpinnings of

signatures of higher-order cognitive processes, previously thought to be unique to humans, supported by

physically separable circuits in the brain [45-49].

Here, we characterized complex decision-making behavior in mice tested on our longitudinal

neuroeconomic foraging paradigm, Restaurant Row. This task has been utilized previously to gain access

to behavioral and neurophysiological correlates of multiple decision-making systems in the brain, including

fundamentally distinct valuation algorithms engaged within the same trial [5, 47, 50-55]. On this task,

rodents forage for their primary source of food among varying costs (delays cued by tone pitch) and

subjective value (flavors tied to spatially fixed and contextualized locations, or "restaurants") while on a

limited time-budget. Decisions to accept and wait for rewards are separated into discrete stages of primary

commitment (accept vs. reject in an offer zone) and secondary re-evaluation (stay vs. quit in a wait zone).

We tested both male and female mice – the first ever analysis of females in this task – transfected with a

newly packaged adeno-associated virus expressing LINC00473 or only a green fluorescent protein targeted

mPFC neurons (**Supplementary Fig. 1**). Mice were tested daily over two months across a changing

economic landscape during which complex foraging strategies developed. We report sex- and LINC00473-

dependent differences in neuroeconomic decision-making behavior specific to alterations in multiple

features of change-of-mind re-evaluative choices without affecting primary decisions. Differences were

rooted in dissociable forms of counterfactual thinking, or how sensitive individuals were to missed

opportunities for alternative actions that could have led to hypothetically better outcomes – the

computational basis of regret processing previously captured by leveraging this task across species [5, 45,

46, 48, 56, 57].

Results:

<u>*Male and female mice similarly acquire the Restaurant Row task.*</u>

During the first week of Restaurant Row testing, all reward offers were fixed at 1 s only as mice

acquired the basic structure of the task (**Fig. 1a-b**), learning to run in the correct direction to forage for low-

cost food rewards. All mice were food restricted to the same percentage (~80-85%) of baseline free-feeding

weight (**Supplementary Fig. 2**). Both males and females, regardless of GFP or LINC00473 treatment,

grossly traveled similar total distances across the first week (**Fig. 1c**, sex*virus: F=0.971, p=0.331).

Behavior shaped equally among all groups as the number of laps run in the correct direction (**Fig. 1d**, day:

F=334.219, p<0.0001; sex*virus: F=2.780, p=0.104) and total pellets earned (**Fig. 1e**, day: F=393.991,

p<0.0001; sex*virus: F=3.062, p=0.089) increased across the first week. All mice displayed subjective flavor

preferences, revealed by splitting earnings by flavors ranked from least to most preferred by summing end-

of-session totals in each restaurant (**Fig. 1f**, rank: F=255.250, p<0.0001). There was a modest upward shift

in earns split by flavor ranking in LINC00473-treated females across the first week (**Fig. 1f**, sex*virus:

F=3.334, p=0.076). Overall, these data indicate all groups were able to demonstrate the ability to forage for

food in a self-paced manner without overt locomotor or feeding changes.

<u>*LINC00473 alters value-dependent, task-driven conditioned place preference behavior in females.*</u>

Before and after the first week of behavioral testing, we characterized exploratory behavior in the

maze arena (**Fig. 2a**). That is, without any task, mice were allowed to roam the maze freely on day 0 and

again on day 8 of the experimental timeline (**Fig. 2b**). This allowed us to probe a conditioned place

preference (CPP) response for reward sites driven by experience that naturally developed throughout the

first week of self-paced task performance. We measured the number of entries into and time spent near

the feeder site in each restaurant's wait zone on the probe sessions (**Fig. 2c**). All mice explored the arena,

including each restaurant, to similar degrees at baseline (day 0) and did not differ between groups

(sex*virus [entries]: F=0.259, p=0.614; sex*virus [time]: F=1.155, p=0.290, **Fig. 2c**). Overall, compared to

the day 0 pre-test baseline, we found that during the day 8 post-test all mice robustly increased total time

spent (F=75.818, p<0.0001) and to a lesser degree entries into the reward sites F=8.501, p<0.01, **Fig. 2c**).

We then split these metrics by restaurant ranked according to day 7's task earnings and found that CPP

scores scaled with flavor preferences (rank [entries]: F=16.307, p<0.0001; rank [time]: F=9.829, p<0.0001,

Fig. 2d-e). We found a significant decrease in CPP scores with respect to time spent in the wait zone of

the most preferred flavor in LINC00473-treated females only, despite being matched to males for equivalent

number of rewards earned on day 7 (sex*virus*rank [time]: F=2.844, p<0.05, **Fig. 2d-e**). This sex-

dependent effect was replicated in an independent cohort of mice separately transfected with a lower titer

of the AAV-expressing LINC00473 (**Supplementary Fig. 3**). Conversely, LINC00473-treated females

displayed an increased number of entries into less preferred restaurants compared to males or GFP

controls (sex*virus*rank [entries]: F=5.584, p<0.001, **Fig. 2d-e**, **Supplementary Fig. 3**). These findings

highlight that mPFC LINC00473 expression can have sex-specific effects on select actions during value-

based reward-seeking behavior. More specifically, LINC00473 treatment in females altered preferences-

dependent entering vs. exiting choices that may access dissociable computations involved in checking-

waiting-leaving behavioral sequences [51, 58, 59], shedding light on more precise interpretations of CPP

metrics than are traditionally reported and explored in detail below.

Escalating reward scarcity unmasks sex-dependent effects of LINC00473 on complex foraging profiles.

We next examined task foraging profiles across the subsequent weeks as we increased the

distribution of offer costs available in the task environment without adjusting the limited time budget mice

had to forage for food (**Fig. 3a**). Thus, in this closed-economy system, tightening budget constraints

pressured animals to adapt to the changing landscape of the environment as reward scarcity increased –

a model of economic stress we characterized recently [33]. Costs increased in a stepwise manner across

blocks of days (i.e., offers ranged from 1-5 s on days 8-12, 1-15 s on days 13-17, and finally 1-30 s on days

18-45, as previously published [5, 33, 60]) and were randomly sampled from a uniform distribution. As

expected, we found a decrease in pellets earned in response to each stepwise economic challenge (sign

test [-]: t=-12.655, p<0.0001, **Fig. 3b**). These deficits scaled by flavor and were matched between all groups

(rank: F=6.938, p<0.001, timepoint*rank: F=3.651, p<0.01; sex*virus: F=0.016, p=0.899, **Fig. 3c-d**,

Supplementary Fig. 4). The most striking change in behavior that accounted for environmental scarcity-

induced reward deficits across the experiment was the proportion of session time re-allocated to quitting

behavior in the wait zone, which significantly increased at each economic challenge step and more

prominently in females (timepoint*choice: F=226.523, p<0.0001; timepoint*sex [quits]: F=11.821, p<0.0001,

Fig. 3e, **Supplementary Fig. 4**). Across the final epoch of testing (the 1-30 s offer range period: days 18-

45), mice displayed a complex response profile, including running more laps that developed over multiple

weeks in an effort to recover and increase total food earnings (laps [days 18-45]: F=355.340, p<0.0001;

earns [days 18-14]: F=252.218, p<0.0001, **Fig. 3b-d**). All groups were matched in their ability to recover

earnings (sex*virus: F=0.234, p=0.631, **Fig. 3d**, **Supplementary Fig. 4**). However, we found that females

displayed a greater increase in the number of laps run compared to males throughout this process (sex*day:

F=7.848, p<0.01, **Fig. 3b,f**, **Supplementary Fig. 4**). Interestingly, LINC00473-expressing females

displayed a significantly higher propensity to engage in wait zone quitting behavior compared to males that

persisted across the entire experiment (sex*virus: F=8.201, p<0.01, **Fig. 3g**, **Supplementary Fig. 5**).

Furthermore, LINC00473-expressing females displayed shorter latencies to quit upon arrival once in the

wait zone compared to males particularly for more preferred flavors (sex*virus*day*rank: F=2.805, p<0.05),

despite no changes in offer zone reaction times (F=1.713, p=0.162, **Supplementary Fig. 5**). Taken

together, these data reflect rearrangements in complex foraging patterns that emerged across longitudinal

time scales in response to worsening economic demand. These analyses reveal sex-dependent effects of

mPFC LINC00473 expression that alter quit decisions unmasked in a reward-scarce environment. This

effect of LINC00473 is reminiscent of the altered check-then-leave behaviors in LINC00473-expressing

females during the CPP probe session (see Fig. 2)

Females demonstrate unique neuroeconomic decision-making policies during change-of-mind choices.

To characterize the economic nature of the decisions made during distinct action-selection

processes in the offer zone and wait zone, we quantified choice outcomes to enter vs. skip (offer zone) or

earn vs. quit (wait zone) separated by both the cued offer cost and flavor identity (**Fig. 4a-c**). All mice

acquired the economic structure of the task, making cost-informed enter vs. skip choices in the offer zone

or earn vs. quit choices in the wait zone, discriminating tones as well as demonstrating different degrees of

willingness to wait in each restaurant (offer*flavor [offer zone]: F=13.129, p<0.001; offer*flavor [wait zone]:

F=18.006, p<0.001, **Fig. 4a-c**). Thresholds of willingness to accept and separately wait for offers can be

calculated independently in the offer zone and wait zone by fitting a Heaviside step regression to the choice

outcome as a function of cost and extracting the inflection (indifference) point in each restaurant (**Fig. 4d**).

Overall, in all mice, we found that offer zone and wait zone thresholds significantly differed from each other

in magnitude of response to the changing economic landscape (timepoint*zone: $F=296.648$, $p<0.0001$, **Fig.**

4d). Upon transitioning to the 1-30 s epoch of testing (day 18), offer zone thresholds drastically increased

in all groups to similar degrees while retaining the ordinal ranking of flavor preferences (rank: $F=17.247$,

$p<0.0001$; sex: $F=0.170$, $p=0.681$; sex*virus: $F=0.031$, $p=0.861$, **Fig. 4d-f**). Wait zone thresholds, on the

other hand, increased only in male mice following the transition to the 1-30 s epoch (rank: $F=58.574$,

$p<0.0001$; sex: $F=17.776$, $p<0.0001$; sex*virus: $F=1.238$, $p=0.268$, **Fig. 4d-f**). Over the subsequent week

(days 18-24), wait zone thresholds in males decreased before stabilizing across the remainder of the

experiment (days 25-45, **Fig. 4d-f**). In females, wait zone thresholds remained relatively unchanged upon

transition to day 18 and were significantly lower than males across the entire 1-30 s epoch (sex*rank [offer

zone]: $F=0.484$, $p=0.694$; sex*rank [wait zone]: $F=3.689$, $p<0.05$, **Fig. 4d-f**). Offer zone thresholds on the

other hand gradually decreased, equally in all groups, but across a much slower timescale over several

weeks (**Fig. 4d-f**). Leveraging an economic analysis we recently developed [33], we found that female

decision-making policies were more optimized for both recovering food as well as yielding subjective value

in a reward-scarce environment (**Fig. 4g-i**, **Supplementary Fig. 6**). LINC00473 expression in mPFC did

not significantly alter any decision-making thresholds (sex*virus [offer zone]: $F=1.781$, $p=0.185$; sex*virus

[wait zone]: $F=187$, $p=0.175$, **Fig. 4d-f**). These data highlight a robust sex difference between distinct offer

zone and wait zone decision-making processes that we previously demonstrated derive from separable

circuits [48, 50].

LINC00473 in mPFC enhances change-of-mind contrast via sensitivity to sunk costs and action-specific

regret.

Wait zone thresholds capture economic indifference points of re-evaluation decisions defined by

choice outcome as a function of the starting offer. Next, we quantified more dynamically at what timepoint

quit decisions were made. Thus, we calculated the value remaining in the countdown at the moment of

quitting by subtracting the within-trial time remaining from an individual mouse's wait zone threshold (**Fig.**

5a). We found all mice generally made quit decisions: (i) for offers that were typically above one's wait zone

threshold (i.e., negatively valued offers) and (ii) with ample time remaining in the countdown such that the

time investment required to obtain the reward was still above one's threshold (i.e., negative value left), and

therefore would be advantageous to quit (**Fig. 5a**). Thus, overall, quit decisions reflected economically

advantageous re-evaluations of sub-optimal offer zone decisions relative to one's own policy, consistent

with previous reports [5, 33, 48, 49, 51]. We found that females made quit decisions with significantly lower

value remaining in the countdown compared to males (sex*virus: $F=195.823$, $p<0.0001$, **Fig. 5b**). This sex

difference was more pronounced in LINC00473-expressing animals and thus made females even more

efficient compared to males (sex*virus: $F=12.957$, $p<0.001$, **Fig. 5b**). Furthermore, only LINC0047-

expressing mice revealed a sex difference in the number of quit decisions that could be characterized as

economically advantageous (as described above, sex*virus: $F=153.236$, $p<0.0001$), with a greater

proportion in females compared to other quit events that were less common and more disadvantageous

(i.e., quitting negatively valued offers [above one's threshold] with positive value left [below one's threshold];

or, quitting positively valued offers [below one's threshold] – both reflecting quits passing the timepoint after

which mice should have chosen to stay instead (quit type*sex*virus: $F=13.293$, $p<0.0001$, **Fig. 5c-d**).

We then characterized the temporally dynamics of how mice made continuous re-evaluation

decisions in the wait zone as a function of the passage of time. We calculated the probability of staying and

earning a reward as a continuous function of not only (i) the time remaining in the countdown but also (ii)

the time already spent waiting. We computed a survival analysis using a sliding 1 s window after binning

all trials of varying starting offers into [time spent, time left] pairs (**Fig. 5f**). In all groups, we found that the

passage of time drove an increased likelihood that mice would continue to wait, the effect of which grew

additively stronger with escalating time spent (time spent*time left: $F=891.924$, $p<0.0001$, **Fig. 5f-h**). This

captures a well-studied cognitive phenomenon known as the sunk cost bias whereby irrecoverable past

expenditures can enhance the value of continuing to invest in and perseverate on an ongoing endeavor, as

we have reported previously across species [46, 49]. We found a significant sex difference in sensitivity to

sunk costs only in LINC00473-expressing mice. LINC00473-expressing females demonstrated increased

sensitivity to sunk costs compared to LINC00473-expressing males which instead demonstrated relatively

diminished sensitivity to sunk costs during quit decisions (sex*virus: $F=15.944$, $p<0.0001$, **Fig. 5i**). This

effect was independent of and orthogonal to temporal distance to the goal measured at any given moment

in time (**Fig. 5f**), only emerged after mice transitioned to the 1-30 s reward-scarce environment (time spent

in wait zone [1-15 s offers]: $F=0.174$, $p=0.677$, **Fig. 5i**, **Supplementary Fig. 7**), and uniquely affected wait

zone decisions compared to other forms of time spent on this task (time spent in offer zone: $F=0.558$,

$p=0.455$; time spent*sex*virus: F-0.418, p-0.518, **Supplementary Fig. 7**). These data highlight sex- and

LINC00473-dependent alterations in how the passage of time can differentially enhance reward value and

escalate the commitment to an ongoing endeavor specifically during re-evaluation decisions.

Next, we aimed to characterize how events related to quit decisions on a given trial influenced

subsequent behavior. Because trials are interdependent, a strength of the Restaurant Row task lies in being

able to subdivide trial sequences into unique economic scenarios in which animals naturalistically find

themselves. Analysis of mice after they make suboptimal choices and realize or gain access to information

that an alternative, unselected action could have led to a better outcome has led to breakthrough

discoveries modeling aspects of the behavioral and neurophysiological signatures of counterfactual thinking

– the computational basis of regret processing [5, 47, 48, 61-63]. We previously linked such processes to

quit decisions as well as other types of choices on this task [5, 47, 48]. In past reports, we and others

behaviorally operationalized counterfactual sensitivity on Restaurant Row by identifying trial pair sequences

[trial t-1 → trial t] in which animals made suboptimal choices [on trial t-1] and by measuring the change in

behavior on the next trial [on trial t: the read-out trial] [5, 47, 48]. Importantly, these events are compared

to specific control sequences that remove the error of one's own agency [on trial t-1], matched for all other

economic factors (e.g., including the offer values and flavor identities on both trials t-1 and trial t) in order

to behaviorally measure counterfactual sensitivity based on one's choice history. Recent work

demonstrated the importance of the mPFC in these behavioral responses [48]. Sequences were extracted

post hoc from natural encounters while foraging amid randomly presented offers, not constructed a priori

nor built into the task design.

We previously defined two distinct regret-related sequences of events on this task [5, 47, 48].

Scenario 1: situations in which animals inappropriately forgo a high-value offer [on trial t-1] only to

subsequently encounter a worse low-value offer on the next trial [trial t] (**Fig. 5j**). This violation sequence

can be compared to the non-violation control sequence in which mice instead appropriately accepted the

high-value offer on trial t-1, holding all other variables constant (**Fig. 5j**). Here, we found that all groups

were insensitive to scenario 1, with no effect of sex or LINC00473 expression (+ sign test: [male GFP]: t=-

0.282, p=0.608; [female GFP]: t=0.101, p=0.461; [male LINC]: t=0.521, p=0.308; [female LINC]: t=-0.144,

p=0.556, **Fig. 5j**). That is, there were no changes in behavior on the read-out trial when comparing violation

sequences to non-violation sequences [thus yielding a delta score of zero in change of choice probabilities

on trial t] (**Fig. 5j**). **Scenario 2**: situations in which animals inappropriately accepted a low-value offer [on

trial t-1] that typically precipitates a resultant change-of-mind decision (**Fig. 5k**). This violation sequence

can be compared to the non-violation control sequence in which mice instead appropriately skipped the

low-value offer on trial t-1, holding all other variables constant and matched for the same low-value offer on

trial t as in scenario 1 (**Fig. 5j-k**). Overall, we found that mice demonstrated sensitivity to scenario 2,

whereby individuals were more likely to compensate for and accept low-value offers that they typically would

not on the read-out trial following violation sequences compared to non-violation sequences (**Fig. 5k**).

Interestingly, only GFP-treated females did not display scenario 2 sensitivity, while all other groups including

LINC00473-expressing females displayed delta scores significantly greater than zero (+ sign test: [male

GFP]: t=2.931, p<0.01; [female GFP]: t=1.123, p=0.145; [male LINC]: t=3.056, p<0.01; [female LINC]:

t=2.196, p<0.05, **Fig. 5k**). Taken together, these data reveal how mice can differentially carry the weight of

different types of past mistakes into future choices, with a unique association between change-of-mind-

related counterfactual sensitivity and mPFC LINC00473 expression in females. These findings provide an

example whereby LINC00473 expression in this brain region overcomes a baseline sex difference that

normally disadvantages females.

Lastly, we characterized how access to task-relevant information may be critically important for

sex- and LINC00473-dependent differences during change-of-mind decisions. After day 45, we tested mice

for an additional 2 days during which tone playback was silenced only in the wait zone but remained audible

during offer zone presentations (**Fig. 6a**). That is, mice had access to cost information signaled by tone

pitch when making initial enter vs. skip decisions in the offer zone, but received no countdown information

once in the wait zone while making secondary re-evaluative decisions to quit or stay before earning

rewards. Overall, we found that males were more sensitive to this informational manipulation compared to

females. Both GFP- and LINC00473-expressing males displayed a modest decreased number of laps run

and total number of pellets earned, while female mice remained relatively unchanged between audible and

silent conditions (laps [timepoint*sex]: $F=10.678$, $p<0.01$; laps [timepoint*virus]: $F=0.449$, $p=0.507$; earns

[timepoint*sex]: $F=10.124$, $p<0.01$; earns [timepoint*virus]: $F=0.002$, $p=0.969$, **Fig. 6b-c**). This task

manipulation did not alter any overt changes in offer zone choices in any group as animals continued to

discriminate the audible tones during cost-informed enter vs. skip decisions (offer: $F=1186.818$, $p<0.0001$;

offer*timepoint: $F=1.365$, $p=0.243$; **Fig. 6d**). Interestingly, when examining changes in wait zone thresholds,

we found a significant sex difference that was compounded by an additional LINC00473 effect in females

only (**Fig. 6e**). In males, compared to the audible condition, the lack of auditory information cuing mice on

progress during the countdown in the silent wait zone condition drove an increase in wait zone thresholds

for most preferred flavors (timepoint: $F=22.976$, $p<0.0001$; timepoint*virus: $F=0.181$, $p=0.671$ **Fig. 6e**).

Males also displayed a concomitant decrease in wait zone thresholds for least preferred flavors (timepoint:

$F=19.561$, $p<0.0001$; timepoint*virus: $F=0.526$, $p=0.470$, **Fig. 6e**). These data indicate that, without cued

auditory guidance in the wait zone, males were more patient waiting for favorite flavors and less patient

waiting for least favorite flavors. Conversely, GFP-expressing females only displayed a decrease in wait

zone thresholds for less preferred flavors, with no change in thresholds for most preferred flavors (**Fig. 6e**).

LINC00473-expressing females however did not display any changes in wait zone thresholds in any

restaurant between audible and silent wait zone conditions (timepoint [most preferred]: $F=2.569$, $p=0.112$;

timepoint*virus [most preferred]: $F=0.137$, $p=0.712$; timepoint [least preferred]: $F=12.185$, $p<0.0001$;

timepoint*virus [least preferred]: $F=8.901$, $p<0.01$). These data indicate LINC00473-expressing females

valued offers in a consistent manner with or without externally guided information, suggesting that their

internal measures of time or holding information from the offer zone in working memory may be more stable.

When examining quit times in the wait zone, all mice displayed a modest increased latency to quit in the

silent condition compared to the audible condition (timepoint: $F=11.764$, $p<0.001$; sex: $F=2.358$, $p=0.126$;

virus: F=0.836, p=0.361, **Fig. 6f**). Finally, when examining sensitivity to sunk costs, we found that the silent

condition unmasked a sex difference in GFP-treated mice whereby females show increased sensitivity as

a function of the passage of time compared to males that was previously only apparent in LINC00473-

expressing groups during the audible condition (time spent: F=97.196, p<0.001; sex: F=36.989, p<0.0001;

virus: F=1.899, p=0.168; **Fig. 6g**, see **Supplementary Fig. 8** for alternate task information manipulations).

Taken together, these findings highlight the complexity of change-of-mind decisions, how mice internally

generate value representations with vs. without access to explicit task-relevant information, and the

multifactorial ways in which this re-evaluative choice process can be uniquely modulated in a sex-

dependent manner.

Discussion:

In this study, we characterized highly complex behavior in mice tested on the neuroeconomic foraging paradigm, Restaurant Row. We explored the role of a non-coding RNA, a relatively poorly understood class of molecules, on motivated behavior. We discovered sex-dependent effects of LINC00473 expression in mPFC neurons on specific aspects of value-based decision-making information processing. This was typified by selectively altering the way in which an individual changes one's mind during an ongoing decision. Importantly, we discovered a role for mPFC LINC00473 expression – which increases stress resilience in simple behavioral screening tests in female mice only [19] – in sharpening the contrast of re-evaluation processes in females. Our findings extend from simple conditioned place preference behaviors like those commonly assayed using simpler paradigms [40, 64, 65] to more complex, higher order neuroeconomic phenomena. These latter effects included enhancing specific aspects of sensitivity to sunk costs and counterfactual thinking. These data provide a better understanding of how multiple stages of a continuous decision stream derive from dissociable psychological mechanisms supported by physically separable circuits in the brain. Our findings have important implications for how molecular determinants of stress responses influence changes in motivated behavior via unique decision-making computations differentially leveraged by males and females.

The distinction between offer zone choices and wait zone choices on the Restaurant Row task is critically important. They reflect a separation of fundamentally distinct decision-making algorithms that occur within the same trial but are segregated across space and time [51]. Numerous reports have tested subjects on this paradigm translated for use across species in task variants adapted for mice, rats, monkeys, and humans [41, 42, 46]. These studies have shed light on the biological and psychological implications of resolving dissociable, evolutionarily conserved decision-making processes. In the offer zone, individuals engage deliberative systems that represent alternating options of competing actions between which one must choose [54, 55]. This has been shown in rodents to derive in part from hippocampal (HPC)-dependent processes [50, 51]. Population level activity of neurons in HPC is capable of decoding multiple, alternating future paths through the offer zone to potential goals in the current vs. next restaurant [58, 64, 66, 67].

Similar findings have been observed from fMRI studies in humans tested on a translated version of the

Restaurant Row task [45]. Default mode network activation patterns can decode competing future options

during offer zone deliberation, but seldom do such representations include past events or previous

restaurants [2, 68]. Decisions in the wait zone on the other hand are much less understood. The HPC has

been implicated in wait zone decisions as well: it switches oscillatory activity from a predominately theta-

driven state sensitive to the cued offer costs in the offer zone [54] instead to an increase in sharp wave

ripple activity sensitive to time spent waiting in the wait zone [55]. Further, wait zone decision periods also

depend on HPC synchrony with the mPFC [54, 55]. Recordings in rodents from the mPFC surprisingly

found no evidence of encoding of the cued offer costs when animals were in the offer zone [52]. However,

once in the wait zone, mPFC neurons strongly represented offer value independent of time spent waiting

and could predict whether or not the trial would terminate in a quit [52]. Within the mPFC, pairs of neurons

also drastically changed excitation-inhibition balance specifically when animals were in the wait zone [69].

Taken together, these findings indicate a qualitative shift in mental state upon transition from the offer zone

to the wait zone that recruits the mPFC exclusively during change-of-mind decisions.

This study represents the first analysis of females in the Restaurant Row task. Nearly all of the sex

differences captured here, including those altered by mPFC LINC00473 expression, are restricted to wait

zone and not offer zone behaviors. Furthermore, sex differences not only persist but are even exaggerated

when access to task-relevant cost information is restricted in the wait zone during change-of-mind

decisions. We showed recently that LINC00473 expression diminishes spontaneous excitatory post-

synaptic current frequency and amplitude in excitatory mPFC pyramidal neurons of female mice, with no

effect seen in males [19]. Chemogenetic inactivation of the mPFC diminishes wait zone quit decisions [53-

55]. Optogenetic induction of long-term depotentiation in excitatory mPFC neurons that project to the

nucleus accumbens (NAc) [50], where negative deflections in dopamine signals have been found to

causally precede quit choices [70], selectively decreases quitting behavior for several days without altering

offer zone behavior [50]. Outside of Restaurant Row, intracranial optogenetic excitatory self-stimulation of

mPFC terminals in the NAc can be reinforcing while also promoting enter-then-quitting behavior [71]. These

converging findings suggest that LINC00473 expression is likely mediating sex-dependent effects on

change-of-mind decisions by at least in part dampening and restructuring mPFC activity. How these

alterations in change-of-mind decisions relate to mPFC LINC00473's known role in enhancing resilience to

stress in a sex-dependent manner will help bridge interpretations between molecular drivers of stress

responses and the behavioral computations associated with susceptibility vs. resilience to stress in males

and females.

Recent transcriptomic analyses of human postmortem mPFC obtained from individuals diagnosed

with major depressive disorder revealed a female-specific decrease in LINC00473 expression in the mPFC

[19]. While LINC00473 is not normally expressed in rodents – it arose in the primate lineage – expressing

LINC00473 in mouse mPFC neurons demonstrated pro-resilient effects in females, but not males,

protecting against several depressive-like phenotypes in simple behavioral tests after exposure to chronic

social defeat or variable stress [19]. LINC00473 expression, which predominates in pyramidal neurons in

human mPFC [19], has been shown to be regulated by conserved Ca^{2+}/calmodulin signaling pathways and

is likely induced by cAMP response element-binding protein (CREB) [19-21, 23]. CREB has long been

studied based on more traditional candidate gene approaches and shown to regulate numerous other

genes involved in neuronal excitability and synaptic plasticity as well as stress sensitivity [72-77]. Like

LINC00473, CREB function in the mPFC has been implicated in promoting resilience to stress, however,

unlike LINC00473, this has been shown to be true for both males and females [19, 78]. These reports

suggest either that there must be some divergence of intracellular CREB-LINC00473-interacting molecular

pathways between sexes conferring differential sensitivities to stress, or that prior studies were limited in

their ability to resolve more nuanced sex differences at the behavioral level.

We recently characterized the impact of chronic social defeat stress on Restaurant Row behavior

in male mice, in tandem with disruptions of CREB function in mPFC [33, 48, 79]. First, we found that male

mice were required to progress through worsening economic challenges in order to extract any group

differences in foraging profiles [33]. This is consistent with results presented here that emphasize the

importance of considering the environmental demand and stressors of a task. Second, we found that stress-

resilient mice – defined by a simple social interaction screening test – mounted the most robust behavioral

activation in response to the transition into the 1-30 s reward-scarce environment in efforts to recover and

maintain food security despite budget constraints, similar to LINC00473-expressing female mice as shown

here [33]. Third, without grossly altering decision-making thresholds, we found that stress-susceptible male

mice demonstrated increased regret-related sensitivity to type 1 scenarios (economically risky decisions to

skip, followed by poor outcomes), a phenotype no group demonstrated in the present study [48].

Additionally, stress-susceptible mice demonstrated diminished regret-related sensitivity to type 2 scenarios

(change-of-mind decisions) compared to stress-resilient individuals who instead displayed enhanced

change-of-mind sensitivity to both regret and sunk costs related to quitting behavior [48]. Interestingly, in

the present study, this constellation of differences was only apparent among the GFP-expressing and

LINC00473-expressing females, respectively, suggesting that females at baseline have a decision-making

vulnerability specifically in change-of-mind choice processes that is restored to male levels by LINC00473

expression in mPFC. In our earlier study, knocking down CREB function had multifactorial effects

depending on the neuroeconomic valuation algorithm in question, the brain region targeted (mPFC vs.

NAc), and whether or not animals were stress-naïve at baseline vs. exposed to subsequent stress [48].

CREB function in the mPFC was required to maintain suppression of regret-related sensitivity to type 1

scenarios and, when knocked down, rendered only type 2 scenarios sensitive to subsequent stress – a

response profile distinct from CREB function in the NAc where it is known to drive stress susceptibility [48,

72, 80, 81]. These data shed light on a divergence between CREB and LINC00473 in mediating dissociable

aspects of counterfactual thinking that could independently contribute to the affective toll of past mistakes

and one's overall mood burden. While these prior studies were conducted in males [33, 48, 79], they

highlight multiple dimensions with which to consider how information integrates across fundamentally

distinct choice processes in order to ultimately precipitate sex-dependent behavioral phenotypes related to

one's stress vulnerability.

LINC00473's sex-dependent effects on behavior have only been causally tested in the mPFC to

date [19]. Although postmortem human tissue studies implicate mPFC as a top region to investigate [19],

understanding LINC00473's role in other regions, especially in how it may be interacting with other key

molecular players such as CREB in the NAc [23, 72, 73], will be critically important in future studies to

elucidate how information processing integrates across neural networks. For instance, LINC00473 has

been shown to increase cellular excitability in cultured HPC neurons [23], contrary to its effects in mPFC

[19]. Future decision-making studies should combine experiments of LINC00473 manipulations with

exposure to other stressors, beyond the economic challenges tested in the present study, such as chronic

social defeat or variable stress to explore compounding effects [19]. Further, more work needs to be done

toward investigating the role of gonadal steroid hormones, sex chromosomes, and sex-dependent

noncoding RNAs in mediating aspects of the reported sex differences. As just one example, CREB function

has been shown to fluctuate with estrus cycle [82-86]. Additionally, given previous reports that LINC00473

expression in mPFC of stress-naïve male mice does induce behavioral changes in certain simpler assays

(i.e., marble burying) [19], it would be fruitful to explore in future work how Restaurant Row decision-making

profiles could potentially relate to individual differences in task-unrelated traits (e.g., anxiety-related

behavior measured on independent tests). Lastly, it is critically important to consider how the decision-

making processes as measured here could relate directly to and drive information processing during the

expression of one's symptomology itself (e.g., underlying the choices leading to anhedonia or social

avoidance related behaviors themselves) as opposed to occupying a more general role as a trait of the

individual.

Taken together, we discovered robust sex differences in the overall contrast of how an individual

decides to change one's mind. This is conceptualized as how dynamic a value representation can be when

deciding to quit an ongoing endeavor and the strength with which quitting behavior can influence the

psychology of multiple cognitive processes. We characterized this as both the steepness of a tipping point

during change-of-mind decisions to either stay or quit and the weight such events can carry into future

choices. We found that this lability depends on individual differences in one's subjective value, on the state

of the environment in the form of how scarce reward availability and access to information might be, and is

uniquely enhanced by LINC00473 expression in mPFC of females. Counterfactual thinking during change-

of-mind decisions can elicit "should not have done that" mental representations that map hypothetical

outcomes onto alternative actions that could have been selected instead. Even if change-of-mind decisions

are an unpleasant experience or evoke to some degree a form of cognitive dissonance, enhancing the

sharpness of this process could increase attention paid toward realized losses, can possess utility for fictive

learning, and collectively may be a sex-dependent feature of stress resilience altering the lens through

which individuals process their prior mistakes.

Figure Captions:

Fig. 1 | Male and female mice acquire the Restaurant Row task. a Experimental timeline. 40 C57BL/6J mice (20 male, 20 female) underwent stereotaxic surgery, half of each sex receiving AAV-LINC00473 or AAV-GFP viruses injected bilaterally in the mPFC. Mice were allowed to recover before beginning longitudinal testing on the Restaurant Row task for 45 consecutive days. Dashed box indicates time period relevant for this figure: week 1 of behavioral testing. **b** Task schematic. Mice were allotted 30 min daily to invest time foraging for their primary source of food in a self-paced manner. Costs to obtain rewards were in the form of delays mice would have to wait near feeder sites. Mice were required to run in a counterclockwise direction encountering offers for different flavors at each "restaurant" in serial order. Each restaurant, separated by hallways, was divided into a T-shaped "offer zone" choice point and a separate "wait zone" that housed the pellet dispenser. Upon offer zone entry from the correct heading direction, a tone sounded whose pitch indicated the delay mice would have to wait if accepting the offer by entering the wait zone. If entered, tone pitch descended in the wait zone, cuing the indicated delay. Each trial terminated if mice skipped in the offer zone, quit during the countdown in the wait zone, or earned a reward, after which animals were required to proceed to the next restaurant. **c-f** Simple behavioral metrics across the first week of testing during which all offers were 1 s only (lowest pitch, 4 kHz): (c) total distance traveled, (d) laps run in the correct direction, (e) total rewards earned, (f) earnings split by flavors ranked from least to most preferred by summing each day's end-of-session totals in each restaurant. No effect of sex or LINC00473 expression was grossly apparent in these initial metrics. Error bars represent ±1 SEM.

Fig. 2 | LINC00473 expression in mPFC alters value-dependent, task-driven conditioned place preference behavior in females. a Experimental timeline. Dashed box indicates time period relevant for this figure: two special testing days marked with cyan arrows. **b** Experimental design. Animals were placed in the arena free to roam for 20 min on day 0 and day 8 of the experimental timeline but with no active task to obtain pre-task baseline and post-task experience-related exploratory behavior. **c** Total number of entries and total time spent in the wait zone, aggregated in all restaurants. **d-e** Delta scores in change of (d) total entries and (e) total time comparing post (day 8) minus pre (day 0) time points split by restaurants ranked

according to rewards earned on the active task on day 7. Only the most preferred (1st) and least preferred

(4th, faint) restaurants are depicted here (see Supplementary Fig. 3 for full dataset and replication cohort).

Horizontal dashed gray line indicates delta score of 0. LINC00473 expression in mPFC abolished CPP

behavior in most preferred restaurants only in females and only in the time but not entry domain. *Represent

significant sex differences. Error bars represent ±1 SEM.

Fig. 3 | Escalating reward scarcity unmasks sex-dependent effects of LINC00473 expression in

mPFC on complex foraging profiles. a Experimental timeline. Dashed box indicates time period relevant

for this figure: the entire 45-day Restaurant Row paradigm. **b** Total rewards earned (black) and laps run in

the correct direction (gray). All 40 mice collapsed. Visual guidance color bars along the x-axis reflect the

experimental schedules of the change the distribution of offer costs illustrated in the timeline in (a). Vertical

dashed green-yellow-red lines indicate the transition points of the stepwise schedule of and are re-used

throughout all other figures as a visual aid. Inset depicts "hysteresis" plot of laps against earns (epochs of

days are color-coded). **c** Rewards earned split by flavors ranked from most preferred (1st) to least preferred

(4th, faint). All 40 mice collapsed. **d** Change in rewards earned in each restaurant comparing two days at

various color-coded transition points (delta scores calculated by subtracting the following: i, day 8-7

(yellow); ii, day 13-12 (orange); iii, day 18-17 (red); iv, day 45-18 (purple). All 40 mice collapsed (see

Supplementary Fig. 4 for comparison between groups). **e** The portion of total session time engaged in one

of the four major choice behaviors: offer zone – enter (green) or skip (red); wait zone – quit (magenta) or

earn (blue). Rings split restaurants ranked from most preferred (1st, inner ring) to least preferred (4th, outer

ring). Each of the 5 pie charts represent data from each stage of testing, with the early and late epochs

during the 1-30 s offer range extracted from the first 5 and last 5 days of that period. **f** Laps run in the correct

direction split by sex and treatment groups. **g** Proportion of enter decisions that terminated in a quit outcome

once in the wait zone. LINC00473 expression in mPFC drove females to run more laps in response to a

worsening economic landscape and quit more frequently. *Represent significant sex differences. Shading

/ error bars represent ±1 SEM.

Fig. 4 | Females demonstrate unique neuroeconomic decision-making policies during change-of-mind choices. a Experimental timeline. Dashed box indicates time period relevant for this figure: the entire 45-day Restaurant Row paradigm. **b** Example behavior from a single mouse early or late during the final 1-30 s testing epoch (red portion of the experimental timeline). Data depict offer zone choice outcomes on the y-axis (enter vs. skip) as a function of the randomly presented cued offer cost (tone pitch) on the x-axis in each restaurant (dots represent individual trial choice outcomes). Quit outcomes in the wait zone are indicated in black dots. **c** Aggregate example summary choice data from all 40 mice early or late during the final 1-30 s testing epoch. Data depict separate offer zone or wait zone summary probability of choice outcome. Horizontal dashed gray line indicates chance at 0.5. Mirror curves for skip and quit (1-probabilities) are intended to highlight cross over points that approximate threshold fits. **d** Thresholds or indifference points of willingness to accept offers in the offer zone or earn offers in the wait zone. Inset depicts an example session from a single mouse in one restaurant where vertical solid green or blue lines illustrate the inflection of Heaviside-step regression fits to choice outcomes in each zone as a function of starting offer costs. All 40 mice collapsed. Inset depicts "hysteresis" plot of offer zone against wait zone thresholds (epochs of days are color-coded). **e** Offer zone thresholds (top) and wait zone thresholds (bottom) split by sex and treatment groups. GFP groups in faint dashed lines. **f** Thresholds from (e) extracted from early and late 1-30 s epoch (red). GFP groups in dashed lines. **g-i** Economic analysis of how well offer zone and wait zone decision policies during the 1-30 s epoch approximate subjective value yield as determined from individual differences in earnings by flavor acquired during the low-cost environment earlier in testing (day 7) when all rewards cost only 1 s. This is compared against theoretical decision policies that maximize total food intake during the 1-30 s epoch irrespective of flavor. See Methods and Supplementary Fig. 6 for a full explanation of this analysis. (g) Individual differences in (top) the standard deviation of rewards earned across flavors on day 7, (middle) total rewards earned on day 7, and (bottom) earns split by restaurant coded in color (flavor: chocolate-brown, banana-yellow, grape-purple, plain-cyan) and size (reward count) on day 7. All data in (g) is sorted by stdev data in (top). Small colored boxes along the x-axis indicate sex (male-blue, female-red) and treatment groups (LINC-dark, GFP-light). **h** Example data from a single mouse on day 7 showing value ratios among earns between restaurants relative to the least preferred flavor (faint). **i** Data from (g-h) used in a computational modeling analysis to

calculate the Euclidean distance between observed decision policies each day from days 18-45 (cool to

warm colors, each dot indicates group average per day) to theoretical policies that could achieve either

maximal food (x-axis) or maximal subjective value (y-axis) for offer zone and wait zone thresholds. Inset

diagonal lines indicate reference point for the 1:1 unity line. Prominent sex difference, without apparent

effect of LINC00473 expression in mPFC, in present in wait zone but not offer zone economic decision

policies. ***Represent significant sex differences. Shading / error bars represent ±1 SEM.**

Fig. 5 | LINC00473 expression in mPFC enhances change-of-mind contrast via sensitivity to sunk

costs and action-specific regret. a Distribution of quit trials as a function of offer value (x-axis) or value

left at the moment of quitting (y-axis) defined relative to one's wait zone threshold (origin vertical and

horizontal dashed blue lines). Data split by restaurants ranked from least to most preferred. All 40 mice

collapsed. Data extracted from the entire 1-30 s epoch. **b** Average value left at the moment of quitting split

by sex and treatment groups. Horizontal dashed gray line indicates 0 value left (or wait zone threshold). **c**

Three categories of quit events based on offer value and value left terms. The green quadrant (V_o<0 &

V_L<0) represent the most efficient quit decisions from economically disadvantageous offers. **d** Counts of

total number of quit events aggregated across all restaurants from each economic quit category described

in (c). **e** Counts of total number of quit events from each economic quit category in (d) split by restaurants

ranked from least (4th) to most preferred (1st) extracted from the entire 1-30 s epoch (red). **f-h** Sunk cost

analysis of staying behavior in the wait zone, demonstrated using all 40 mice. (f) The likelihood of staying

in the wait zone and earning a reward (e.g., not quit) is plotted as a function of time left in the countdown

along the x-axis and time already spent waiting orthogonally in color. Note the black 0 s time spent curve

represents animals having just entered the wait zone from the offer zone. Inset vertical dashed gray line

illustrates an example analysis point comparing three sunk cost conditions originating from different starting

offers but matched at 10 s left. Data from (f) dimensioned reduced in (g) collapsing across time left,

highlighting the grand mean of each time spent sunk cost condition (color and x-axis). Insets in (g) depict

data from curves in (f) are collapsed into the observed (sunk condition) and control (0 s condition) lines.

Difference between curves in (g) are plotted in (h) in order to summarize the envelope of the overall effect

of time already spent on escalating the commitment of staying in the wait zone. Horizontal dashed line

represents delta score of 0. **i** Sensitivity to sunk costs plotted across the entire Restaurant Row paradigm

where the delta curve in (h) is plotted as vertical slices in (i) with time already spent along the y-axis and

heatmap representing delta score magnitude. **j-k** Regret-related sequence analyses comparing trial pairs

between (j) type 1 scenarios and (k) type 2 scenarios each. Read out trials captures a change in behavior

depending on choice history of distinct economic violations (black) compared to non-violations (gray) on

trial t-1. Read out trials report delta scores in likelihood change of entering by subtracting choice outcomes

between violation-nonviolation sequences. Horizontal dashed gray line indicates delta score of 0. See

Methods for a full explanation of these analyses. Prominent sex-dependent effect revealed by LINC00473

expression in mPFC on change-of-mind related sunk costs (enhanced in females) and regret sensitivity

(type 2: enhanced in females). * in (b-i) represent significant sex differences. * in (j-k) represent delta score

significantly different from 0. Shading / error bars represent ±1 SEM.

Fig. 6 | Sex- and LINC00473-dependent wait zone behaviors are differentially modulated by access

to task information. a Informational manipulation schematic: (left) standard Restaurant Row task followed

by (right) two days of tones silenced only in the wait zone, but not offer zone (remains audible as before).

b Total number of laps run in the correct direction (top) and total pellets earned aggregated among

restaurants (bottom) across the audible and silent wait zone task conditions (x-axis). **c** Rewards earned in

each restaurant ranked from least (4th) to most preferred (1st) across the audible (black) and silent wait zone

(gray) task conditions. **d** Overall offer zone choice probabilities to make enter decisions as a function of

audible cued offer costs collapsed across all flavors to illustrate no effect of task condition on auditory

discriminatory behaviors. Horizontal dashed line indicates change at 0.5. **e** Wait zone thresholds in each

restaurant across task conditions. **f** Latency to quit reaction times once in the wait zone in each restaurant

across task conditions. **g** Sensitivity to sunk costs in the wait zone depicted here only in the silent condition

as a function of time already spent (x-axis). Y-axis reflects delta earn probabilities as depicted in Fig. 5h.

Prominent sex-dependent effects of withholding wait zone information on overall task performance

(observed in males), wait zone economic decision policies (bidirectional in males, unidirectional in GFP-

females, and no effect in LINC00473-females), and sunk cost sensitivity (sex difference revealed for the

first time GFP groups unlike in Fig. 5). Horizontal dashed line indicates delta score of 0. * in (b-f) represent

significant effects of task condition within group. * in (g) represent significant sex differences. Shading /

error bars represent ±1 SEM.

Acknowledgments:

We thank members of the labs of Eric Nestler and Scott Russo for helpful discussion and technical

assistance. Open-source illustrations obtained from SciDraw (www.scidraw.io), credit Federico

Claudi.

Funding:

National Institute of Mental Health grant R01MH051399 (EJN)

National Institute of Mental Health grant R01MH129306 (EJN)

National Institute of Mental Health grant R01MH114882 (SJR)

National Institute of Mental Health grant R01MH127820 (SJR)

National Institute of Mental Health grant R01MH104559 (SJR)

National Institute of Mental Health grant L40MH127601 (BMS)

National Institute of Mental Health supplement grant R01MH051399-31S1 (BMS)

Leon Levy Scholarship in Neuroscience, New York Academy of Sciences (BMS)

Burroughs Wellcome Fund Career Award for Medical Scientists (BMS)

Animal Models for the Social Dimensions of Health and Aging Research Network via NIH/NIA R24

AG065172 (BMS)

Brain & Behavior Research Foundation Young Investigator Award 31140 (RDC)

Author contributions:

Conceptualization: RDC, OI, EJN, BMS

Methodology: RDC, OI, BMS

Investigation: RDC, BY, NJ, AA, SK, KG, SA, SJR, EJN, OI, BMS

Data curation: RDC, OI, BMS

Formal analysis: BMS

Visualization: RDC, OI, BMS

Funding acquisition: OI, BMS

Supervision: RDC, BMS

Writing – original draft: RDC, OI, BMS

Writing – review & editing: RDC, BY, NJ, AA, SK, KG, SA, SJR, EJN, OI, BMS

Competing interests:

Authors declare that they have no competing interests.

Data and materials availability:

All data and materials used in the analysis are available in the manuscript, materials and methods

section, or supplementary information.

Supplementary Figure Captions

Supplementary Fig. 1 | Validation of newly packaged adeno-associated virus expression of LINC00473. Raw qPCR cycle threshold count (ct) obtain from mPFC tissue punches extracted from mice separately transfected with either AAV2-hSyn-GFP (control) or AAV2-hSyn-LINC00473-GFP (treatment). qPCR readouts obtained for expression levels of GFP, HPRT1 (housekeeping), ZFP189 (another known regulator of stress-resilience), and LINC00473. Note similar levels of expression of GFP, HPRT1, and ZFP189 between control and treatment viruses. Also note no detectable levels of LINC00473 in mice transfected with the control virus, as this is not endogenously expressed in mice. Dots represent individual samples. Error bars represent ±1 SEM.

Supplementary Fig. 2 | Bodyweight measurements. Percentage of a 2-day average (circled in gray, horizontal dashed line for reference) of baseline body weight obtained 4 days prior (black epoch) to the start of the longitudinal Restaurant Row paradigm. During this period (black epoch), mice were food restricted and fed a limited ration to intentionally decrease weights to approximately 80-85% of baseline that was steadily maintained throughout the remainder of the experiment (days 1-45, see main Fig. 1 for an explanation of the color-coded epochs of testing across the paradigm). These body weights reflect the measurements immediately before task performance on each day of testing. Shaded area represents ±1 SEM.

Supplementary Fig. 3 | Replication cohort of two different virus titers on sex-specific Restaurant Row conditioned place preference (CPP) findings. Animals were placed in the arena free to roam for 20 min on day 0 and day 8 of the experimental timeline but with no active task to obtain pre-task baseline and post-task experience-related exploratory behavior (see main Fig. 2). **a** Total number of entries and total time spent in the wait zone, aggregated in all restaurants. Depicting findings from independent cohorts. **b** Delta scores in change of entries and time comparing post (day 8) minus pre (day 0) time points split by restaurants ranked according to rewards earned on the active task on day 7. **c** Data from (b, only 1st and 4th ranking restaurants) replotted with number of rewards earned on day 7 on the x-axis. Horizontal dashed gray line indicates delta score of 0. LINC00473 expression in mPFC abolished CPP behavior in most

preferred restaurants only in females and only in the time but not entry domain, replicated in both cohorts.

Error bars represent ±1 SEM.

Supplementary Fig. 4 | Basic Restaurant Row laps and earnings metrics across testing. a Overall

sex difference comparisons between males and females (collapsing across GFP and LINC00473 virus

treatment within each sex). Average number of laps run in the correct direction, total rewards earned, ratio

of earns / laps, and standard deviation of earns among flavors. **b** Data from (a) split by GFP and LINC

groups. **c** Data from main Fig. 3d split by sex and virus groups depicting change in rewards earned in each

restaurant comparing two days at each color-coded transition point. Delta scores calculated by subtracting

the following: i, day 8-7 (yellow); ii, day 13-12 (orange); iii, day 18-17 (red); iv, day 45-18 (purple). Shaded

area and error bars represent ±1 SEM.

Supplementary Fig. 5 | Restaurant Row choice metrics split by flavor ranking across testing. a

Average number of enter decisions made in the offer zone split by sex and virus treatment groups as well

as the 4th and 1st ranking restaurant. **b** Number of skip decisions made in the offer zone. **c** Offer zone

reaction time before making an enter or skip decision calculated from offer zone entry (tone onset) to either

a right turn into the wait zone (enter) or a left turn into the hallway (skip). **d** Number of quit decisions made

in the wait zone during a countdown. **e** The probability of quitting in the wait zone given the animal entered.

This normalizes the likelihood of quitting to enter frequency. **f** Wait zone latency to quit reaction time from

countdown onset (wait zone entry) until mice leave the wait zone prematurely. Shaded area represents ±1

SEM.

Supplementary Fig. 6 | Longitudinal economic analysis of decision policy optimization for food

security vs. subjective value. Analysis underlying data from main Fig. 4g-i derived from Durand-de Cuttoli

et al 2023 *Biological Psychiatry* [33]. **a** Computer Restaurant Row simulation of total number of pellets

earned. The ideal threshold required to obtain the theoretical maximum number of pellets when ignoring

flavor preferences was empirically determined to be 10 s. **b** Relative ratio of earns among restaurants for

an example mouse on day 7 between the flavor rankings (i.e., this mouse had a 3.7 : 2.1 : 1.4 : 1 ratio of

earns across flavors capturing a summary of idealized relative subjective value for this mouse when all

offers were 1 s only). **c** Two intersecting planes of decision policies that yield varying amounts of theoretical

value either for maximal food intake (as determined by computer simulations, brown plane) or subjective

value (as determined by multiplying day 7 preference ratios on a mouse-by-mouse basis, pink plane) when

in a reward-scarce environment (1 to 30 s offers). Here, only two decision policy dimensions (least preferred

and most preferred restaurants) of the four dimensions (all four ranked restaurants) are displayed. Both

planes are normalized to each's min and max values for the purpose of plotting both in the same space

while preserving the threshold coordinate locations that achieve either theoretical maximum (brown

[location always fixed at threshold coordinates of 10 s] or pink beacons [location of discovered threshold

coordinates that vary from mouse-to-mouse]). Actual observed daily decision policies represented by the

cyan beacon wander throughout this space from day 18 to 45 in a reward-scarce environment. Trajectories

projected to the floor of this display trace out individual mouse decision policy paths. Example coordinates

of brown, pink, and cyan beacons illustrated as dots on the x and y axes. Euclidean distance from cyan

coordinates to either brown or pink coordinates were calculated (question mark symbols). **d** Euclidean

distances from observed decision policies in the offer zone (left) or wait zone (right) to either food (brown)

or preference (magenta) theoretical maximum across testing in a reward-scarce environment split by GFP

(top) and LINC (bottom) groups and sex. Shaded area represents ±1 SEM.

Supplementary Fig. 7 | Sensitivity to sunk costs analysis during the 1-15 s testing epoch. Mice do

not display sensitivity to sunk costs during days 13-17 in a relatively reward rich environment compared to

days 18+ (when offers range 1-30 s, as in main Fig. 5). **a-c** Sunk cost analysis of staying behavior in the

wait zone, demonstrated using all 40 mice. (f) The likelihood of staying in the wait zone and earning a

reward (e.g., not quit) is plotted as a function of time left in the countdown along the x-axis and time already

spent waiting orthogonally in color. Note the black 0 s time spent curve represents animals having just

entered the wait zone from the offer zone. Inset vertical dashed gray line illustrates an example analysis

point comparing three sunk cost conditions originating from different starting offers but matched at 10 s left.

Data from (a) dimensioned reduced in (b) collapsing across time left, highlighting the grand mean of each

time spent sunk cost condition (color and x-axis). Difference between curves in (b) are plotted in (c) in order

to summarize the envelope of the overall effect of time already spent on escalating the commitment of

staying in the wait zone. Horizontal dashed line represents delta score of 0. **d** Data in (c) split by sex and

virus treatment groups depicting no effect of time spent during the countdown in the wait zone on changing

the probability of quitting vs. staying to earn a reward. **e** To compare to time spent waiting in the wait zone,

data here depict a different type of time spent on the task: time spent in the offer zone prior to accepting an

offer and entering the wait zone does not influence the probability of quitting vs. staying to earn a reward

once in the wait zone. These data are from the 1-30 s reward-scarce epoch (days 18-45) to demonstrate

the unique value of time spent waiting in the wait zone uniquely conferring sunk cost sensitivity. Error bars

represent ±1 SEM.

Supplementary Fig. 8 | Full set of task information manipulations on Restaurant Row. A After day

45, mice were challenged with special probe days on three different task informational manipulations each

for 2 days. Manipulations took the form of silencing and withholding tone presentation either in the offer

zone, wait zone, or both: (i) silent offer zone but audible wait zone [days 46-47], (ii) audible offer zone but

silent wait zone [days 48-49], and (iii) silent offer zone and silent wait zone [days 50-51]. **b** Offer zone

choice probability to enter as a function of offer cost split by restaurants ranked from least (4th) to most (1st)

preferred separating sex and virus treatment groups (rows) across the 4 different task information

conditions (columns). **c** Data in (b) collapsed across all four restaurants showing summary enter

probabilities as a function of offer cost. Note that enter probabilities are largely unaffected when the offer

zone tones are audible comparing baseline (black, regular task) to silent wait zone (blue) conditions,

however, as intended, offer zone choice probabilities change into a flat horizontal line in both the silent offer

zone (green) and silent task (gray) conditions, as animals do not have any tone information in the offer zone

during these conditions. **d** Wait zone thresholds plotted as a relative fold change comparing each condition

to the regular task audible baseline separated across the three information manipulations (silent offer zone,

left, green; silent wait zone, middle, blue; silent task, right, gray) split by restaurants ranked from least (4th)

to most (1st) preferred, sex, and virus treatment groups. Note the difference in patterns in changes in wait

zone thresholds across restaurants between sex/virus groups and across task information manipulation

conditions. Shaded area and error bars represent ±1 SEM.

Methods

Subjects: Adult male and female C57BL/6J mice (Jackson, 10wks, n=40) were used. Mice

were individually housed and maintained on a 12-hr light/dark cycle with ad libitum water, only

food-restricted during Restaurant Row testing, conducted during their light phase. Experiments

were approved by the Mount Sinai Institutional Animal Care and Use Committee (IACUC; protocol

number LA12-00051) and adhered to the National Institutes of Health (NIH) guidelines.

Viral constructs: To achieve long-term neuronal expression of LINC00473, it was

subcloned into an AAV plasmid with human synapsin promoter (hSyn) driving gene expression.

LINC00473 transcript variant 1, (NR_026860.1 /AK289375) of 1821bp length was subcloned from

p1005+ HSV vector (PMID: 32304628) into pAAV-hSyn-eGFP-hSny plasmid (PMID: 36449610)

using blunt ligation into a PMEI and EcoRV restriction sites following the second hSyn promoter.

The orientation and sequence were validated using Sanger sequencing. Empty plasmid

expressing eGFP only was used as control. These vectors were used to generate AAV serotype

2 at the University of Maryland virus vector core. Viral vector placement and expression were

confirmed in vivo by a GFP fluorescent flashlight and qPCR.

RNA extraction, RT, and qPCR: Total RNA was isolated and purified with RNeasy Micro

Kit (Qiagen), using QIAzol as a lysis reagent, including DNase treatment. All samples were tested

for concentration and purity using a NanoDrop (Thermo Fisher). RNA amount was normalized

across samples (400ng), and cDNA was created using iScript (Bio-Rad). Real-time PCR reactions

were run in triplicate using PowerUp SYBR Master Mix (Applied Biosystems) in a QuantStudio 7

(Thermo   Fisher)   qPCR   machine.   Primer   sequences: LINC00473:   forward:

GCATACTTTGGCGGACCTTTT,   reverse   TGTGCCTCCCTGTGAATTCTC;   HPRT1   forward:

GCAGTACAGCCCCAAAATGG,   reverse: GGTCCTTTTCACCAGCAAGCT;   GFP   forward:

CATGCCCGAAGGCTACGT, reverse CGATGCCCTTCAGCTCGAT. Raw Ct values are

presented and not normalized amounts, as LINC00473 is not endogenously expressed in mice.

Virus surgery: Mice were anesthetized by intraperitoneal injection with a mixture of

Ketamine HCl (100 mg/kg) and Xylazine (10 mg/kg) and positioned on a stereotaxic instrument

(David Kopf Instruments). In the medial prefrontal cortex (mPFC, from bregma with an angle of

15 degrees: AP +1.8 mm; ML ±0.75 mm; DV −2.7 mm), 0.7 μL of virus (AAV2-hSyn-GFP or AAV2-

hSyn-LINC00473-GFP) was bilaterally infused using 33-Gauge Hamilton needles over 5 min at a

rate of 0.1 μL/min, and the needle was left in place for 10-15 min after the injection before

retraction. Mice were allotted 2 weeks to recover before beginning food restriction across 3-4 days

in preparation for testing on Restaurant Row. An initial virus validation cohort (independent cohort

of n=10 mice) was injected with GFP or LINC00473 viruses bilaterally in mPFC at a titer of 1×10^{12}

at. These animals were tested on Restaurant Row for only 1 week (offers remained at 1 s only,

first epoch, green) after which they were sacrificed for PCR validation on tissue punches obtained

from mPFC. Behavioral data from these animals on the conditioned-place preference probe (days

0 and 8) appear in Supplementary Fig. 3. A second replication cohort (n=40 mice) was injected

with GFP or LINC00473 viruses bilaterally in mPFC at a titer of 5×10^{12} and served as the primary

cohort of animals for the main figures that underwent the full 45-day longitudinal behavioral

testing.

Neuroeconomic Task: Following surgery, we characterized mice longitudinally in

Restaurant Row [5]. On this task, mice had a limited time period each day to forage for food by

navigating a maze with four uniquely flavored (chocolate, banana, grape, and plain) and

contextualized (horizontal stripes, dots, triangles, vertical stripes) feeding sites, or "restaurants"

(Fig. 1b). Animals were allotted 30 min to forage for their primary source of food for the day.

Rewards used on this task were obtained from BioServe (20 mg pellets #F007120). These pellets

are full nutrition and not sucrose-only pellets that only differed in flavoring but not nutritional profile

(3.6 kcal/gm calories, 59.1% carbohydrate, 18.7% protein, 6.5% ash, 5.6% fat, 4.7% fiber).

Choices on this task were interdependent across trials and across days. This makes the task

economic in nature, requiring animals to budget their limited time effectively in a self-paced

manner in order to earn a sufficient amount of food in a closed-economy system. The cost of

earning a reward took place in the form of a delay animals would have to wait cued by the pitch

of a tone that sounded upon entry into each restaurant. Each restaurant had a separate offer zone

(OZ) and wait zone (WZ). Upon entry into the OZ, a tone (500 ms) sounded whose pitch indicated

how long of a delay mice would have to wait in a cued countdown should they enter the WZ. Mice

could decide to skip an offer presented in the OZ and proceed down the hallway to the next

restaurant. In the OZ, tones repeated every second held at the same pitch to repeatedly cue the

offer of the current trial until mice either made a skip or enter decision. If mice decide instead to

enter the WZ, this triggered a countdown with descending tone pitch stepping every second until

mice either waited out the full delay and earned a food pellet or quit prematurely during the

countdown. If mice made a skip or quit decision, the tone terminated, the offer was rescinded,

and mice must proceed forward to the next restaurant. Mice quickly learned to only travel in one

direction as animals must encounter offers serially navigating the maze in a counterclockwise

direction (i.e., nothing happens if mice traverse in the wrong direction other than wasting time).

Mice were tested across 45 consecutive days for the main experiment where strategies developed

longitudinally as animals learned the structure of the task and the changing economic landscape.

During the first week of testing (block 1, green epoch), all trials consisted of 1 s offers only.

1 s offers were indicated by tones played at the 4,000 Hz pitch. A special probe session was

inserted into this timeline immediately before (day 0) and after (day 8) the first week of testing

during which mice essentially were allowed to roam the empty maze arena freely without any task

components (no tones, no rewards) to examine conditioned place preference behaviors learned

through the natural experience of the first week of the task itself. The remainder of the

experimental timeline continued thereafter. During the next 5 days of testing (block 2, yellow

epoch), offers ranged from 1 to 5 s (4,000 Hz to 5,548 Hz in 387 Hz steps) randomly selected

from a uniform distribution. For instance, if a 4 s offer was selected on a given trial, the tone

presented in the OZ would be 5,161 Hz for 500 ms, and this would repeat every second indefinitely

until a skip or enter decision was made. If the WZ was entered, using this example trial, this 5,161

Hz tone would step down by 387 Hz every second descending in pitch until earning (with the final

tone of 4,000 Hz) or quitting prematurely. Block 3 (5 days, orange epoch) consisted of a 1 to 15

s (4,000–9,418 Hz) range. The fourth and final block (red epoch) consisted of offers ranging from

1 to 30 s (4,000–15,223 Hz) and continued for the remainder of the experiment. This block-wise

protocol allowed us to examine how mice initially shaped their behavior as they learned the basic

structure of the task as well as how they adjusted their foraging strategies across long timescales

(days, weeks, and months) as they transitioned from reward-rich to reward-scarce environments

in a stepwise manner. That is, blocks 1-3 are relatively reward rich as costs are generally low

delays and rewards are easier to earn compared to block 4 which is relatively reward scarce as

there are far more expensive reward offers distributed in the environment for which animals must

learn to forage efficiently. The rewards used in this task only differ in flavor for which animals have

subjective differences in preferences quantified below. All flavor options become increasingly

scarce, equally across the longitudinal paradigm.

Rewards were delivered using a custom-built 3D-printed automated pellet dispenser that

was triggered by a computer running the task programmed in ANY-Maze (Stoelting), run under

dim lighting. The task was run by video-tracking animal position in real-time where all choice

contingencies are determined by the centroid of the animal's body crossing territories in the maze

in combination with being required to run in a counterclockwise direction. The key zones in the

maze include each restaurant's separate OZ and WZ as well as the hallways in between

restaurants (thus, 12 total main zones). 6 key behavioral epochs make up all possible, mutually

exclusive behaviors animals could be engaged in while spending their limited time budget. 1)

Enter time: defined from OZ entry in the correct heading direction (counterclockwise around the

maze) triggering tone onset until the animal physically enters right into the WZ triggering tone

countdown. 2) **Skip time**: defined from OZ entry in the correct heading direction triggering tone

onset until the animal physically exits left into the hallway. 3) **Quit time**: defined from WZ entry

and tone countdown onset until the animal physically exits the WZ prematurely before the

countdown completes, rescinding the offer and terminating the countdown. 4) **Earn time**: defined

from WZ entry and tone countdown onset until the countdown completes and a pellet is delivered.

5) **Consumption time**: defined from reward delivery onset until the animal physically leaves the

reward site and exits the WZ. 6) **Travel time**: defined from previous trial termination (either from

a skip, quit, or consumption event as defined above) and hallway entry until the next trial begins,

triggered by OZ entry into the next restaurant.

Body weight was measured twice daily, immediately before and after testing on the

Restaurant Row task.

Data analysis: In addition to simple behavioral metrics like rewards earned, laps run,

reaction times as described above, a key neuroeconomic behavioral metric is decision thresholds.

This metric approximates indifference points in choices made either in the OZ or WZ as a function

of cost and flavor. We measured the inflection point of fitting heaviside step regressions to enter

versus skip choices in the OZ or earn versus quit choices in the WZ as a function of cued offer

cost in each restaurant. Thresholds determined this way were used for several subsequent

analyses, including calculating reward value. The value of an offer can be calculated by

subtracting a trial's offer from one's WZ threshold as this reflects the relative cost normalized to

each animals' willingness to wait and earn a reward (i.e., offer value <0 indicate offers typically

not earned and thus should be skipped in the OZ or quit in the WZ as determined by each animal's

subjective flavor-specific decision policy which remained relatively stable across block 4 of

testing). Similarly, when quitting in the WZ, the value left at the moment of quitting can be

calculated by subtracting the time remaining in that trial's countdown from the WZ threshold (i.e.,

value left <0 indicates an amount of time left that is not worth waiting for and should have been

quit whereas value left >0 would have been more advantageous to finish waiting at that point

instead of quitting). The amount of time spent remaining in the countdown at the moment of

quitting relative to what the starting offer was and how much time was spent also served as the

basis of our analysis of sensitivity to sunk costs, as previously published in mice, rats, and humas

whereby time spent waiting can decrease the probability of quitting independent of time left in the

countdown, see Sweis et al *Science* for the development of this analysis [44, 46, 48, 49, 53].

Thresholds as well as economic choice history serve as the basis and criteria used in regret-

related analysis whereby distinct violation vs. non-violation sequences are compared head-to-

head in order to quantify the effects of mistake history on influencing choice behavior on the next

trial. See Steiner and Redish 2014 *Nature Neuroscience* [47], Sweis et al 2018 *PloS Biology* [5],

and Durand-de Cuttoli 2022 *Science Advances* [48] for the development of these analyses.

Thresholds were also used for an analysis we recently developed to capture how mice develop

strategies that optimize food yield while also maximizing subjective value, see Durand-de Cuttoli

et al 2023 *Biological Psychiatry* [33] for the development of this analysis.

Data were processed in Matlab with statistical analyses in JMP Pro 17. All data are

expressed as mean ± 1 standard error. Statistical significance was assessed using student's t

tests, sign tests, and one-way, two-way, three-way, and repeated measures ANOVAs. No data

were excluded as outliers.

References

1. Stone, C., J.B. Mattingley, and D. Rangelov, *On second thoughts: changes of mind in decision-making.* Trends Cogn Sci, 2022. **26**(5): p. 419-431.

2. Liu, X., et al., *Common and distinct networks underlying reward valence and processing stages: a meta-analysis of functional neuroimaging studies.* Neurosci Biobehav Rev, 2011. **35**(5): p. 1219-36.

3. Redish, A.D., *The mind within the brain: How we make decisions and how those decisions go wrong.* 2013: Oxford University Press.

4. McGuire, J.T. and J.W. Kable, *Medial prefrontal cortical activity reflects dynamic re-evaluation during voluntary persistence.* Nat Neurosci, 2015. **18**(5): p. 760-6.

5. Sweis, B.M., M.J. Thomas, and A.D. Redish, *Mice learn to avoid regret.* PLoS Biol, 2018. **16**(6): p. e2005853.

6. Kirkebøen, G., E. Vasaasen, and K. Halvor Teigen, *Revisions and Regret: The Cost of Changing your Mind.* Journal of Behavioral Decision Making, 2011. **26**(1): p. 1-12.

7. Brenner, C., *The mind in conflict*. 1982, New York: International Universities Press. 266 p.

8. Baumeister, R.F., K.D. Vohs, and D.M. Tice, *The Strength Model of Self-Control.* Current Directions in Psychological Science, 2016. **16**(6): p. 351-355.

9. American Psychiatric Association. and American Psychiatric Association. DSM-5 Task Force., *Diagnostic and statistical manual of mental disorders : DSM-5*. 5th ed. 2013, Washington, D.C.: American Psychiatric Association. xliv, 947 p.

10. Mezo, P.G. and R.M. Baker, *The moderating effects of stress and rumination on depressive symptoms in women and men.* Stress Health, 2012. **28**(4): p. 333-9.

11. Seedat, S., et al., *Cross-National Associations Between Gender and Mental Disorders in the World Health Organization World Mental Health Surveys.* Archives of General Psychiatry, 2009. **66**(7).

12. Ferrari, A.J., et al., *Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010.* PLoS Med, 2013. **10**(11): p. e1001547.

13. Marcus, S.M., et al., *Sex differences in depression symptoms in treatment-seeking adults: confirmatory analyses from the Sequenced Treatment Alternatives to Relieve Depression study.* Compr Psychiatry, 2008. **49**(3): p. 238-46.

14. Silverstein, B., *Gender differences in the prevalence of somatic versus pure depression: a replication.* Am J Psychiatry, 2002. **159**(6): p. 1051-2.

15. Martin, L.A., H.W. Neighbors, and D.M. Griffith, *The experience of symptoms of depression in men vs women: analysis of the National Comorbidity Survey Replication.* JAMA Psychiatry, 2013. **70**(10): p. 1100-6.

16. Eid, R.S., A.R. Gobinath, and L.A.M. Galea, *Sex differences in depression: Insights from clinical and preclinical studies.* Prog Neurobiol, 2019. **176**: p. 86-102.

17. Akil, H., et al., *Treatment resistant depression: A multi-scale, systems biology approach.* Neurosci Biobehav Rev, 2018. **84**: p. 272-288.

18. Ulitsky, I. and D.P. Bartel, *lincRNAs: genomics, evolution, and mechanisms.* Cell, 2013. **154**(1): p. 26-46.

19. Issler, O., et al., *Sex-Specific Role for the Long Non-coding RNA LINC00473 in Depression.* Neuron, 2020. **106**(6): p. 912-926 e5.

20. Chen, Z., et al., *cAMP/CREB-regulated LINC00473 marks LKB1-inactivated lung cancer and mediates tumor growth.* J Clin Invest, 2016. **126**(6): p. 2267-79.

21. Jiang, L.Y., et al., *CREB-induced LINC00473 promotes chemoresistance to TMZ in glioblastoma by regulating O6-methylguanine-DNA-methyltransferase expression via CEBPalpha binding.* Neuropharmacology, 2024. **243**: p. 109790.

22. Ataman, B., et al., *Evolution of Osteocrin as an activity-regulated factor in the primate brain.* Nature, 2016. **539**(7628): p. 242-247.

23.   Pruunsild, P., et al., *Expression of the primate-specific LINC00473 RNA in mouse neurons*
990   *promotes excitability and CREB-regulated transcription.* J Biol Chem, 2023. **299**(5): p.
991   104671.

992   24.   Pruunsild, P., C.P. Bengtson, and H. Bading, *Networks of Cultured iPSC-Derived Neurons*
993   *Reveal the Human Synaptic Activity-Regulated Adaptive Gene Program.* Cell Rep, 2017.
994   **18**(1): p. 122-135.

995   25.   Tran, K.V., et al., *Human thermogenic adipocyte regulation by the long noncoding RNA*
996   *LINC00473.* Nat Metab, 2020. **2**(5): p. 397-412.

997   26.   Glimcher, P.W. and A. Rustichini, *Neuroeconomics: the consilience of brain and decision.*
998   Science, 2004. **306**(5695): p. 447-52.

999   27.   Kenning, P. and H. Plassmann, *Neuroeconomics: an overview from an economic*
1000   *perspective.* Brain Res Bull, 2005. **67**(5): p. 343-54.

1001   28.   Loewenstein, G., S. Rick, and J.D. Cohen, *Neuroeconomics.* Annu Rev Psychol, 2008.
1002   **59**: p. 647-72.

1003   29.   Kalenscher, T. and C.M. Pennartz, *Is a bird in the hand worth two in the future? The*
1004   *neuroeconomics of intertemporal decision-making.* Prog Neurobiol, 2008. **84**(3): p. 284-
1005   315.

1006   30.   Clithero, J.A. and A. Rangel, *Informatic parcellation of the network involved in the*
1007   *computation of subjective value.* Soc Cogn Affect Neurosci, 2014. **9**(9): p. 1289-302.

1008   31.   Levy, D.J. and P.W. Glimcher, *The root of all value: a neural common currency for choice.*
1009   Curr Opin Neurobiol, 2012. **22**(6): p. 1027-38.

1010   32.   Louie, K. and P.W. Glimcher, *Efficient coding and the neural representation of value.* Ann
1011   N Y Acad Sci, 2012. **1251**: p. 13-32.

1012   33.   Durand-de Cuttoli, R., et al., *A double-hit of social and economic stress in mice precipitates*
1013   *changes in decision-making strategies.* Biol Psychiatry, 2023.

1014   34.   Resulaj, A., et al., *Changes of mind in decision-making.* Nature, 2009. **461**(7261): p. 263-

1015   6.

1016   35.   Hare, T.A., C.F. Camerer, and A. Rangel, *Self-control in decision-making involves*

1017   *modulation of the vmPFC valuation system.* Science, 2009. **324**(5927): p. 646-8.

1018   36.   Alagapan, S., et al., *Cingulate dynamics track depression recovery with deep brain*

1019   *stimulation.* Nature, 2023. **622**(7981): p. 130-138.

1020   37.   Knutson, B., et al., *Neural responses to monetary incentives in major depression.* Biol

1021   Psychiatry, 2008. **63**(7): p. 686-92.

1022   38.   Holton, E., et al., *Goal commitment is supported by vmPFC through selective attention.*

1023   Nature Human Behaviour, 2024.

1024   39.   Sweis, B.M. and E.J. Nestler, *Pushing the boundaries of behavioral analysis could aid*

1025   *psychiatric drug discovery.* PLoS Biol, 2022. **20**(12): p. e3001904.

1026   40.   Sweis, B.M., M.J. Thomas, and A.D. Redish, *Beyond simple tests of value: measuring*

1027   *addiction as a heterogeneous disease of computation-specific valuation processes.* Learn

1028   Mem, 2018. **25**(9): p. 501-512.

1029   41.   Abram, S.V., et al., *The Web-Surf Task: A translational model of human decision-making.*

1030   Cogn Affect Behav Neurosci, 2016. **16**(1): p. 37-50.

1031   42.   Maisson, D., et al., *Widespread coding of navigational variables in prefrontal cortex*

1032   biorxiv, 2022.

1033   43.   McInnes, A.N., et al., *Psychometric validation and clinical correlates of an experiential*

1034   *foraging task.* BioRxiv, 2023. **2023.12.28.573439**.

1035   44.   Kazinka, R., A.W. MacDonald, 3rd, and A.D. Redish, *Sensitivity to Sunk Costs Depends*

1036   *on Attention to the Delay.* Front Psychol, 2021. **12**: p. 604843.

1037   45.   Abram, S.V., et al., *Neural signatures underlying deliberation in human foraging decisions.*

1038   Cogn Affect Behav Neurosci, 2019. **19**(6): p. 1492-1508.

46. Sweis, B.M., et al., *Sensitivity to "sunk costs" in mice, rats, and humans.* Science, 2018.
361(6398): p. 178-181.

47. Steiner, A.P. and A.D. Redish, *Behavioral and neurophysiological correlates of regret in*
rat decision-making on a neuroeconomic task. Nat Neurosci, 2014. **17**(7): p. 995-1002.

48. Durand-de Cuttoli, R., et al., *Distinct forms of regret linked to resilience versus*
susceptibility to stress are regulated by region-specific CREB function in mice. Science
Advances, 2022. **8**(42): p. eadd5579.

49. Redish, A.D., et al., *Sunk cost sensitivity during change-of-mind decisions is informed by*
both the spent and remaining costs. Commun Biol, 2022. **5**(1): p. 1337.

50. Sweis, B.M., et al., *Altering gain of the infralimbic-to-accumbens shell circuit alters*
economically dissociable decision-making algorithms. Proc Natl Acad Sci U S A, 2018.
115(27): p. E6347-E6355.

51. Sweis, B.M., A.D. Redish, and M.J. Thomas, *Prolonged abstinence from cocaine or*
morphine disrupts separable valuations during decision conflict. Nat Commun, 2018. **9**(1):
p. 2521.

52. Diehl, G. and A.D. Redish, *Differential processing of decision information in subregions of*
rodent medial prefrontal cortex. biorxiv, 2022.

53. Duin, A.A., et al., *Certainty and uncertainty of the future changes planning and sunk costs.*
Behav Neurosci, 2021.

54. Schmidt, B., A.A. Duin, and A.D. Redish, *Disrupting the medial prefrontal cortex alters*
hippocampal sequences during deliberative decision making. J Neurophysiol, 2019.
121(6): p. 1981-2000.

55. Schmidt, B. and A.D. Redish, *Disrupting the medial prefrontal cortex with designer*
receptors exclusively activated by designer drug alters hippocampal sharp-wave ripples
and their associated cognitive processes. Hippocampus, 2021.

56.   Abram, S.V., A.D. Redish, and A.W. MacDonald, 3rd, *Learning From Loss After Risk: Dissociating Reward Pursuit and Reward Valuation in a Naturalistic Foraging Task.* Front Psychiatry, 2019. **10**: p. 359.

57.   Vialou, V., et al., *DeltaFosB in brain reward circuits mediates resilience to stress and antidepressant responses.* Nat Neurosci, 2010. **13**(6): p. 745-52.

58.   German, P.W. and H.L. Fields, *Rat nucleus accumbens neurons persistently encode locations associated with morphine reward.* J Neurophysiol, 2007. **97**(3): p. 2094-106.

59.   Sjulson, L., et al., *Cocaine Place Conditioning Strengthens Location-Specific Hippocampal Coupling to the Nucleus Accumbens.* Neuron, 2018. **98**(5): p. 926-934 e5.

60.   Nwakama, C.A., et al., *Diabetes alters neuroeconomically dissociable forms of mental accounting.* bioRxiv, 2024.

61.   Camille, N., et al., *The involvement of the orbitofrontal cortex in the experience of regret.* Science, 2004. **304**(5674): p. 1167-70.

62.   Coricelli, G., et al., *Regret and its avoidance: a neuroimaging study of choice behavior.* Nat Neurosci, 2005. **8**(9): p. 1255-62.

63.   Coricelli, G., R.J. Dolan, and A. Sirigu, *Brain, emotion and decision making: the paradigmatic example of regret.* Trends Cogn Sci, 2007. **11**(6): p. 258-65.

64.   Bardo, M.T. and R.A. Bevins, *Conditioned place preference: what does it add to our preclinical understanding of drug reward?* Psychopharmacology, 2000. **153**(1): p. 31-43.

65.   Tzschentke, T.M., *Measuring reward with the conditioned place preference (CPP) paradigm: update of the last decade.* Addict Biol, 2007. **12**(3-4): p. 227-462.

66.   Johnson, A. and A.D. Redish, *Neural ensembles in CA3 transiently encode paths forward of the animal at a decision point.* J Neurosci, 2007. **27**(45): p. 12176-89.

67.   Kay, K., et al., *Constant Sub-second Cycling between Representations of Possible Futures in the Hippocampus.* Cell, 2020. **180**(3): p. 552-567 e25.

68.   Huynh, T., et al., *Vicarious Trial-and-Error Is Enhanced During Deliberation in Human Virtual Navigation in a Translational Foraging Task.* Front Behav Neurosci, 2021. **15**: p. 586159.

69.   Diehl, G. and A.D. Redish, *Measuring excitation-inhibition balance through spectral components of local field potentials.* bioRxiv, 2024.

70.   Kocharian, A., *Dopaminergic control of neuroeconomic decision-making.* ProQuest, University of Minnesota, 2023.

71.   Lind, E.B., et al., *A quadruple dissociation of reward-related behaviour in mice across excitatory inputs to the nucleus accumbens shell.* Commun Biol, 2023. **6**(1): p. 119.

72.   Barrot, M., et al., *CREB activity in the nucleus accumbens shell controls gating of behavioral responses to emotional stimuli.* Proc Natl Acad Sci U S A, 2002. **99**(17): p. 11435-40.

73.   Carlezon, W.A., Jr., R.S. Duman, and E.J. Nestler, *The many faces of CREB.* Trends Neurosci, 2005. **28**(8): p. 436-45.

74.   Kim, J., et al., *Memory recall and modifications by activating neurons with elevated CREB.* Nat Neurosci, 2014. **17**(1): p. 65-72.

75.   Lonze, B.E. and D.D. Ginty, *Function and regulation of CREB family transcription factors in the nervous system.* Neuron, 2002. **35**(4): p. 605-23.

76.   McClung, C.A. and E.J. Nestler, *Regulation of gene expression and cocaine reward by CREB and DeltaFosB.* Nat Neurosci, 2003. **6**(11): p. 1208-15.

77.   Sakai, N., et al., *Inducible and brain region-specific CREB transgenic mice.* Mol Pharmacol, 2002. **61**(6): p. 1453-64.

78.   Lorsch, Z.S., et al., *Stress resilience is promoted by a Zfp189-driven transcriptional network in prefrontal cortex.* Nat Neurosci, 2019. **22**(9): p. 1413-1423.

79.   Durand-de Cuttoli, R., et al., *Chronic social stress induces isolated deficits in reward anticipation on a neuroeconomic foraging task.* biorxiv, 2022.

80.   Barrot, M., et al., *Regulation of anxiety and initiation of sexual behavior by CREB in the*
1116   *nucleus accumbens.* Proc Natl Acad Sci U S A, 2005. **102**(23): p. 8357-62.

1117   81.   Dong, Y., et al., *CREB modulates excitability of nucleus accumbens neurons.* Nat
1118   Neurosci, 2006. **9**(4): p. 475-7.

1119   82.   Zhou, J., et al., *Effects of Estrogen Treatment on Expression of Brain-Derived*
1120   *Neurotrophic Factor and cAMP Response Element-Binding Protein Expression and*
1121   *Phosphorylation in Rat Amygdaloid and Hippocampal Structures.* Neuroendocrinology,
1122   2005. **81**(5): p. 294-310.

1123   83.   Sharma, K., et al., *Chronic exposure to estrogen and tamoxifen regulates synaptophysin*
1124   *and phosphorylated cAMP response element-binding (CREB) protein expression in CA1*
1125   *of ovariectomized rat hippocampus.* Brain Res, 2007. **1132**(1): p. 10-9.

1126   84.   Lorsch, Z.S., et al., *Estrogen receptor alpha drives pro-resilient transcription in mouse*
1127   *models of depression.* Nat Commun, 2018. **9**(1): p. 1116.

1128   85.   Shansky, R.M., et al., *Estrogen mediates sex differences in stress-induced prefrontal*
1129   *cortex dysfunction.* Mol Psychiatry, 2004. **9**(5): p. 531-8.

1130   86.   Issler, O., et al., *The long noncoding RNA FEDORA is a cell type– and sex-specific*
1131   *regulator of depression.* Science Advances, 2022. **8**(48).

1132

Figure 1

a surgery & recovery | day: 1 8 13 18 45 | longitudinal neuroeconomic paradigm

AAV-GFP vs. AAV-LINC00473 | 7 days | 5 days | 5 days | 28 days

1s only *1-5s* *1-15s* *1-30s offer range*

target: medial prefrontal cortex

male C57BL/6J N=20 female C57BL/6J N=20

distribution of wait time (costs)

b Restaurant Row Task

plain

offer zone wait zone

skip enter quit earn grape

must forage for offers in serial order

chocolate

➤ 30 min daily budget
➤ primary source of food
➤ cost indicated by tone pitch
➤ subjective flavor preferences

banana

c GFP LINC — distance traveled (m)

d laps correct direction

e total rewards earned

day

f GFP — 4th rank 3rd rank 2nd rank 1st rank — earns by flavor — day

LINC — 4th rank 3rd rank 2nd rank 1st rank — day

Figure 2

a longitudinal neuroeconomic paradigm

day: 1 7 45

relevant experimental timeline 0 7 days 8

1s only

RRow CPP
pre *post*

b — RRow Conditioned Place Preference —

pre test → *post test*

NO TASK — 7 days of full task — NO TASK

1s only

day 0 days 1 - 7 day 8

c

entries in zone

time in zone

GFP LINC

entries 135 / 75
♂ ♀

pre post *pre post*

time (s) 1000 / 550

d

GFP LINC

Δ entries [post - pre] 17 / -7
♂ ♀

1st rank

4th rank

0 day 7 task earns 35   0 day 7 task earns 35

e

GFP LINC

Δ time (s) [post - pre] 160 / -30
♂ ♀

*

0 day 7 task earns 35   0 day 7 task earns 35

Figure 3

a longitudinal neuroeconomic paradigm

day: 1   8   13   18   45

relevant experimental timeline | 7 days | 5 days | 5 days | 28 days

distribution of wait time (costs) — 1s only | 1-5s | 1-15s | 1-30s offer range

b all 40 mice — earns / laps / day

c all 40 mice — earns by flavor — 1st, 2nd, 3rd, 4th — i ii iii iv

d all 40 mice — Δ earns — transition: iv recovery, i 1-5 s, ii 1-15 s, iii 1-30 s — 4th rank 1st

e session time — quit, earn, enter, skip, rank — 1s only | 1-5s | 1-15s | 1-30s offer range — early late — all 40 mice

f GFP — laps — ♂ ♀ — day 1–45 | LINC — laps — day 1–45 *

g GFP — p(quit | enter) — ♂ ♀ — day 1–45 | LINC — p(quit | enter) — day 1–45 *

Figure 4

a longitudinal neuroeconomic paradigm

day: 1 — 8 — 13 — 18 — 45

relevant experimental timeline | 7 days | 5 days | 5 days | *early* | 28 days | *late*

distribution of wait time (costs): 1s only | 1-5s | 1-15s | 1-30s offer range

b offer zone choice — early 1-30 s / late 1-30 s — earn, quit
banana, plain, chocolate, grape — enter / skip — offer (s) 1–30

c oz early WZ / oz late WZ — p(choice) — enter, earn, skip, quit — 1st rank, 2nd rank, 3rd rank, 4th rank — all 40 mice — offer (s) 1–30

d all 40 mice — threshold (s) — offer zone / wait zone — oz, wz — skip enter, earn, quit — offer (s) 1–30 — day 1–45

e oz threshold (s) — GFP / LINC — ♂ ♀ — wz threshold (s) — day 1–45

f oz threshold (s) early / late — wz threshold (s) — 4th rank – 1st rank

g day 7 — stdev of flavor earnings (sorted) — total earns — earns by flavor — mouse (sorted) 1–40

h value ratio — 1x 1.4x 2.1x 3.7x — example mouse — earns by ranking 5–45 — day 7

i GFP ♂ ♀ / LINC ♂ ♀ — day 18–45 — value Euclidean distance — offer zone policy / wait zone policy — food Euclidean distance

Figure 5

Figure 6

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Fig. 8