<u>Repeated loss of floral scent in independent transitions from bee to</u>

<u>hummingbird pollination in spiral gingers</u>

Kathy Darragh[1,2,+], Kathleen M. Kay[3], Santiago R. Ramírez[1]

[1]Department of Evolution and Ecology, University of California, Davis, CA, USA

[2]Department of Biology, Indiana University, Bloomington, IN, USA

[3]Department of Ecology and Evolutionary Biology, University of California, Santa Cruz,

CA, USA

[+]Corresponding author (Email: kdarrag@iu.edu)

ORCID Kathy Darragh 0000-0002-9215-014X

ORCID Kathleen M. Kay 0000-0001-8858-110X

ORCID Santiago R. Ramírez 0000-0003-1306-1315

Keywords

Abstract

The repeated evolution of the same trait in different lineages provides powerful natural experiments to study the phenotypic and genotypic predictability of how traits are gained and lost. A fascinating example of this is the repeated evolution of hummingbird pollination in plant lineages in the Americas, a widespread and often unidirectional phenomenon. The spiral gingers in the genus *Costus* are ancestrally bee-pollinated and hummingbird pollination has evolved multiple times independently in the tropical Americas. These pollinator transitions are accompanied by predictable morphological and color changes but the changes in floral scent have not been described. In this study, we describe the floral scent composition of 30 species of *Costus* sampled across the phylogeny to understand how floral scent has evolved across the genus with respect to pollinator transitions. We then combine transcriptomics and genomics in one clade of *Costus* to identify genetic expression differences and gene family evolution associated with pollinator transitions. We show that hummingbird-pollinated species have mostly lost their floral scent, whereas bee-pollinated species exhibit either floral scent maintenance or in some cases, gains of more diverse scent profiles. The remarkable consistency of scent loss in hummingbird-pollinated species highlights the shared strong selection pressures experienced by these lineages. Even species with more recent transitions from bee to hummingbird pollination exhibit scent loss, highlighting the rapid breakdown of scent production following pollinator transitions. We also find the floral scent loss appears to be due to gene downregulation rather than pseudogenization. This research highlights the capacity for rapid changes when selection pressures are strong through downregulation of floral scent genes.

Resumen

La evolución independiente de un rasgo en diferentes linajes ofrece la oportunidad de

estudiar la predictibilidad de fenotipos y sus bases genéticas. Un ejemplo fascinante es la

evolución repetida de polinización por colibríes en varios linajes de plantas en las Américas, un

fenómeno generalizado y típicamente unidireccional. Las especies de caña agria del género

Costus son polinizadas ancestralmente por abejas y la polinización por colibríes ha evolucionado

varias veces de forma independiente en América tropical. Cada transición de polinizador ha sido

acompañada de varios cambios morfológicos y colores predecibles, pero los cambios en el aroma

floral no han sido descritos. En este estudio describimos la composición del aroma floral de 30

especies de *Costus* para comprender cómo el aroma floral ha evolucionado con respecto a las

transiciones de polinizadores. Combinamos transcriptómica y genómica en un clado de *Costus*

para identificar diferencias en la expresión genética y la evolución de la familia de genes

asociadas con las transiciones de polinizadores. Nuestros resultados muestran que las especies

polinizadas por colibríes han perdido en su mayoría su aroma floral, mientras que las especies

polinizadas por abejas ya sea mantienen el aroma floral y, en algunos casos, muestran ganancias

de perfiles de aroma más diversos. La consistencia en la pérdida de olor en especies polinizadas

por colibríes resalta las fuertes presiones de selección natural que experimentan estos linajes.

Incluso las especies con transiciones más recientes de polinización de abejas a colibríes exhiben

pérdida de olor, lo que destaca la rápida interrupción de la producción de olores después de las

transiciones de los polinizadores. También encontramos que la pérdida de aroma floral parece

estar controlada por regulación génica negativa y no por la pérdida de genes de producción de

aromas (*e.g.* pseudogenización). Esta investigación destaca la rapidez de cambios a través de la

regulación negativa de genes que controlan aromas florales cuando las presiones de selección

son fuertes.

Introduction

The repeated evolution of the same traits in different lineages is thought to reflect shared selection pressures. These cases of repeated evolution apply not only to the gain of traits, such as the evolution of antifreeze proteins in fish (Chen et al. 1997) or carnivory in plants (Albert et al. 1992), but also the loss of traits, as seen in the eye regression of cavefish (Sifuentes-Romero et al. 2023), or reduction of attractive traits such as flower size or floral scent in self-fertilizing plants (Tsuchimatsu and Fujii 2022). Repeated evolution provides us with systems where we can study the process of adaptation and ask questions about the predictability of evolution, and the extent of constraint, or flexibility, of the underlying molecular mechanisms (Orr 2005; Losos 2011; Sackton and Clark 2019).

One such example of repeated evolution is pollinator transitions in flowering plants. For example, the evolution of hummingbird pollination has occurred many times across flowering plants, and often involves convergence in a suite of traits. Flowers pollinated by hummingbirds are characterized by traits thought to attract and fit hummingbirds (such as a narrow shape and large amounts of nectar) and in addition, traits thought to deter bees (red coloration, lack of nectar guides, lack of scent) (Schemske and Bradshaw 1999; Castellanos et al. 2004; Bergamo et al. 2016). Interestingly, the evolution of hummingbird pollination has primarily been documented in one direction, from ancestral bee pollination to derived hummingbird pollination (Rosas-Guerrero et al. 2014; Abrahamczyk and Renner 2015; Kay and Grossenbacher 2022). However, there is clade-specific variation in this directionality, with the reasons for pollinator transitions remaining unclear (Rosas-Guerrero et al. 2014; Abrahamczyk and Renner 2015; Kay and Grossenbacher 2022; Barreto et al. 2024).

Hummingbird-pollinated flowers lack floral scent (Knudsen et al. 2004; Wessinger 2024). Genomic studies have shown that, even for birds, hummingbirds have a low number of olfactory receptors (Driver and Balakrishnan 2021; Driver 2022), suggesting at least a reduced investment in this sensory modality. Behavioral experiments have also suggested that hummingbirds rely more on visual than olfactory cues when foraging (Goldsmith and Goldsmith 1982). In fact, floral scent analyses of hummingbird pollinated flowers have shown either complete lack of floral scent or low emission rates of floral scent compounds. (Knudsen et al. 2004; Byers et al. 2014; Amrad et al. 2016).

A fascinating group in which to study the evolution of hummingbird pollination is the

genus *Costus* (Costaceae). The genus contains over 80 species of perennial understory

herbaceous monocots, the majority of which have diverged rapidly within the last 3 million years

in Central and South America (Maas 1972, 1977; Vargas et al. 2020). Since establishing in the

American tropics, hummingbird pollination has evolved repeatedly and independently in *Costus*

from an ancestral state of bee pollination (Vargas et al. 2020). Specifically, bee-pollinated

species are pollinated primarily by orchid bees (Apidae: Euglossini), with females visiting

flowers for nectar and pollen, and males visiting for nectar (Kay and Schemske 2003).

Hummingbird-pollinated species are often pollinated by hermit hummingbirds (Kay and

Schemske 2003). While different species were initially classified as hummingbird or bee

pollinated (Maas 1972, 1977), many species do not fit the classic paradigm of long tubular red

flowers associated with hummingbird pollination (Grant and Grant 1968). For example, many

bee-pollinated flowers have some red coloration (Kay and Schemske 2003), and hummingbird-

pollinated flowers are short (Yost and Kay 2009). Despite this variation, a single phenotypic

optimum is shared across most of the independent transitions to hummingbird pollination in the

clade (Kay and Grossenbacher 2022). Traits including small flower size, the absence of nectar

guides, and brightly colored floral bracts are all predictive of hummingbird pollination (Kay and

Grossenbacher 2022). The combination of recent genomic resources, a well-resolved phylogeny,

and the possibility of collecting samples both in the field and in greenhouses, makes *Costus* a

powerful system to study the evolution of convergence on hummingbird pollination (Vargas et

al. 2020; Kay and Grossenbacher 2022; Harenčár et al. 2023).

While the morphological and color changes involved in the evolution of hummingbird

pollination have been well-studied in *Costus*, the expected scent changes have not been

characterized. In this study, we describe floral scent in 30 species of *Costus* sampled across the

phylogeny and including eight independent origins of hummingbird pollination. We test for

differences in floral scent between bee-pollinated and hummingbird-pollinated species and

investigate how floral scent has evolved across the genus. We then focus on one clade of *Costus*

with variation in floral scent production and generate floral transcriptomic data. We use this data

to identify gene expression differences associated with floral scent production, helping us to

better understand the genetic changes underlying floral scent evolution in this group. Finally, we

annotated a family of genes important for the production of terpenoid floral scent compounds

(the terpene synthase genes (TPS)) in three previously available *Costus* genomes, and a high-

quality genome assembly for *Costus allenii* generated in this study, encompassing two

independent transitions to hummingbird pollination. High-quality annotations allow us to

analyze the evidence of gene loss or pseudogenization following transition to hummingbird

pollination in this gene family that plays an important role in floral scent evolution. Our study,

therefore, combines multiple approaches to characterize floral scent changes, gene expression

changes, and gene family evolution associated with shifts to hummingbird pollination in *Costus*.

Results

<u>Bee-pollinated species produce more compounds and more diverse floral scents</u>

We collected 100 samples from 30 *Costus* species including species previously described

as both bee- and hummingbird-pollinated (Fig 1A, Table S1). Using multivariate techniques, we

visualized the floral scent profiles to identify the broad patterns in our dataset. Before filtering

our data, we found that samples from hummingbird-pollinated species did not differ from

ambient control samples (Fig 1B). In contrast, samples from bee-pollinated species spread across

a wider section of chemospace, with many samples visually separated from both ambient and

hummingbird-pollinated samples (Fig. 1B). We found that both pollination group, and species

(nested within pollination group) explained variation in floral scent (PERMANOVA, pollination

group: $F_{2,167}=8.0$, p<0.001; species: $F_{28,167}=2.5$, p<0.001). Interestingly, bee-pollinated species

were significantly different from both hummingbird-pollinated species and ambient control

samples (Pairwise PERMANOVA, p=0.003, whereas hummingbird-pollinated species did not

differ from ambient control samples (Pairwise PERMANOVA, p=0.07) (Figure S1).

After filtering likely contaminants from the ambient air, we obtained a final dataset with

60 floral scent compounds (Table S2). The majority of these compounds were terpenoid (87%,

52/60), with a few aromatics and compounds of unknown class. The dataset included both

monoterpenes and sesquiterpenes. No one compound class was restricted to a specific pollinator

group. After removing the ambient compounds, we found that one bee-pollinated sample and

eighteen hummingbird-pollinated samples had no detectable floral scent, confirming the

similarity of the hummingbird-pollinated samples to the ambient air samples. The unscented

hummingbird-pollinated samples were from six species. For two species, *C. montanus* and *C.*

stenophyllus, all individuals produced no detectable floral scent (4/4 individuals). Furthermore,

most C. chartaceus (4/5), *C. lasius* (4/7), and *C. wilsonii* (5/8), and a single *C. productus*

individual (1/5) lacked floral scent.

We found that bee-pollinated flowers produced more floral scent compounds, averaging 9

(s.d. 10) compounds in comparison to only 2 (s.d. 4) for hummingbird-pollinated flowers (Figure

1C). Bee-pollinated flowers contained more compounds than hummingbird-pollinated flowers

(Fig 2A). Similar to the number of compounds, we used a measure of biochemical and structural

diversity of the compounds present in a sample to calculate the overall chemical diversity. The

Functional Hill Diversity index considers not only the total number of compounds but also their

biosynthetic complexity. We found that the bee-pollinated floral scents had a higher chemical

diversity (Fig 2B). Bee-pollinated flowers also produced a higher abundance of floral scent, as

measured by total ion abundance in each sample, a proxy for compound amount (Fig. 2C).

To see how individual compounds varied between bee- and hummingbird-pollinated

species we calculated the percentage of individuals in each pollination group that produced the

most common floral scent compounds in the dataset. We found that these compounds are present

in a higher proportion of bee-pollinated flowers than hummingbird-pollinated flowers (Fig. 2D).

In particular, (*E*)-beta-ocimene and beta-caryophyllene show the largest percentage differences

between the two pollination groups (Fig. 2D) and are the only two compounds to exhibit a

significant difference in frequency between pollinator groups (Table S2).

To facilitate visualization of our final filtered dataset using NMDS we removed the

nineteen samples with no detectable floral scent. When this filtered dataset was plotted, we

observed more overlap between the hummingbird- and bee-pollinated samples, but the two sets

of samples significantly different (PERMANOVA, pollination group: $F_{1,80}=4.2$, $p<0.001$;

species: $F_{26,80}=2.4$, $p<0.001$; Pairwise PERMANOVA bee- vs hummingbird-pollinated species

$p=0.001$) (Figure S1).

Time

Figure 1. A) Representative images of two pairs of closely related bee- and
hummingbird-pollinated species: *Costus allenii* **and** *C. lasius* **and** *C. erythrophyllus* **and** *C.*
*spiralis***. B) NMDS (nonmetric multidimensional scaling) plot illustrating in the variation in**
the unfiltered dataset (Stress=0.12). Points are labelled as bee-pollinated, hummingbird-
pollinated, or ambient samples. Ambient samples are control samples of the air taken in the

same area as the focal plant individual. C) Total ion chromatogram of floral scent from

***Costus allenii* and *C. lasius*. We chose the individuals with the second-highest amount of**

scent from each species. The *C. lasius* sample was shifted on the x-axis by 1.5 minutes to

match the *C. allenii* sample due to changes in the GC/MS column over time. Abundance is

scaled to the highest peak present in *C. allenii*. 1, alpha-pinene; 2, beta-pinene; 3, beta-

myrcene; 4, cineole; 5, alpha-copaene; 6, (*E*)-beta-caryophyllene; 7, gamma-elemene; 8,

alpha-humulene; 9, (*Z*)-beta-caryophyllene; 10, germacrene D; 11, alpha-elemene; 12,

gamma-cadinene; 13, delta-cadinene; 14, germacrene-d-4-ol; 15, unidentified sesquiterpene

(RI=1616). Contaminants are identified with an asterisk (*).

Figure 2. Differences in bee-pollinated and hummingbird-pollinated *Costus* species
in terms of A) Number of scent compounds produced (ANOVA, pollination group:
$F_{1,70}=60.4$, p<0.0001, species: $F_{28,70}=9.4$, p<0.0001), B) The diversity of scent compounds
produced (ANOVA, pollination group: $F_{1,51}=5.3$, p=0.03, species: $F_{27,51}=1.4$, p=0.1), C)
Total compound amount produced (measured by proxy of total ion abundance) (ANOVA,
pollination group: $F_{1,70}=109$, p=<0.0001, species: $F_{28,70}=2.8$, p<0.001) and D) the percentage
of individuals that produced the ten most common floral scent compounds in the dataset.

205 **Significant differences in frequency between bee- and hummingbird-pollinated flowers as**

206 **determined by Chi-squared testing are highlighted with an asterisk. Fifteen individuals**

207 **with the highest compound production were removed from C for visualization.**

208 <u>Ancestral state reconstructions show that hummingbird-pollinated species have lost floral scent</u>

209 To understand the evolution of floral scent in *Costus*, we carried out phylogenetic

210 analyses using a previously published phylogeny (Vargas et al. 2020). In our dataset, we found a

211 strong correlation between pollination group and the phylogenetic covariance matrix using two-

212 block partial least squares analysis (rPLS = 0.56; P=0.01). This reduces the statistical power of

213 testing group differences using phylogenetic ANOVA due to the high aggregation of pollinator

214 groups on the phylogeny (Adams and Collyer 2018). We tested for evidence of phylogenetic

215 signal in compound number, compound diversity, compound amount (total area of floral scent

216 peaks), (*E*)-beta-ocimene production, and (*E*)-beta-caryophyllene production, detecting weak and

217 non-significant phylogenetic signal overall (λ= 4.14795e-05, p=1). We also tested for evidence

218 of phylogenetic signal in the multivariate floral scent dataset, detecting non-significant almost

219 zero phylogenetic signal (Kmult=0.43, p=0.68). We therefore decided against using phylogenetic

220 ANOVA with simulations based on Brownian motion and instead used a method of randomizing

221 residuals in a permutation procedure (Adams and Collyer 2018). We found that compound

222 number, compound diversity, compound amount, and (*E*)-beta-ocimene production were all

223 significantly different between bee-pollinated and hummingbird-pollinated species (compound

224 number, $F_{1,26}$=7.2, p=0.008; compound diversity, $F_{1,26}$=5.8, p=0.01; compound amount,

225 $F_{1,26}$=627.5, p=0.008; (*E*)-beta-ocimene production, $F_{1,26}$=4.7, p=0.03). (*E*)-beta-caryophyllene

226 production did not differ between pollinator groups ($F_{1,26}$=1.0, p=0.4). Our comparative analyses

227 complement our previous analyses, again showing that bee-pollinated species produce higher

228 amounts of more diverse and complex floral scents and have higher (*E*)-beta-ocimene

229 production.

230 To further investigate how floral scent has evolved across the *Costus* phylogeny, we

231 carried out maximum likelihood ancestral state reconstruction of mean compound production.

232 We overlaid the eight independent origins of hummingbird pollination represented by our dataset

233 as determined in (Vargas et al. 2020). The ancestral state of all *Costus* species found in tropical

234 America is to produce more compounds than average hummingbird-pollinated species (4.5 vs 2).

In this reconstruction, the ancestral *Costus* population produced an average floral scent – more
compounds than hummingbird-pollinated species but less than the average bee-pollinated species
(Figure 3). Many hummingbird-pollinated species have since completely lost floral scent, as
shown by the white circles (Figure 3). We see evidence for scent reduction or loss in all eight
origins of hummingbird pollination in our dataset. In contrast, some bee-pollinated species, such
as *Costus allenii*, have evolved much stronger floral scents than the ancestral state.

**Figure 3. Ancestral state reconstruction of floral scent in the genus *Costus*. Pollinator for
each species is illustrated by the right-hand dots. Mean compound number produced by
each species is shown by the left-hand dots with each shade of gray representing a range of
means. Darker dots indicate more compounds are produced. Predicted compound
production at each node is shown following the same color scale. Origins of hummingbird
pollination (as noted by hummingbird symbols on branches) and tip labels following
(Vargas et al. 2020).**

Loss of floral scent is correlated with downregulation of floral terpene synthase genes.

To analyze the genetic basis of floral scent evolution, we first chose a clade of closely
related species with variation in floral scent production and pollination systems. We chose the

clade containing *Costus lasius* due to the variation in floral scent production and the availability

of a high-quality genome. In this clade, *Costus lasius* (hummingbird-pollinated) produces on

average three floral scent compounds, whereas *Costus allenii* (bee-pollinated) produces 24 (one

of the highest in our dataset), and *Costus villosissimus* (bee-pollinated) produces seven

compounds (Figure 4A).

We first produced floral transcriptomes of all three species. After clustering transcripts

with over 95% similarity, we found 112998 transcripts present in *C. allenii*, 80208 in *C. lasius*,

and 96886 in *C. villosissimus*. Re-mapping rates were high in all three species: 99.2% in *C.*

allenii, 99% in *C. lasius*, and 99.1% in *C. villosissimus*. All three also had BUSCO

(Benchmarking Universal Single-Copy Orthologs) scores of more than 80% (Table S3),

suggesting high-quality transcriptomes with most single-copy orthologs from the Embryophyta

database present in all species.

Due to the dominance of terpenoid compounds in the floral scent, we focused on the

expression of terpene synthase genes (TPS). TPS genes are responsible for producing terpenoid

compounds. We found that 20 TPS genes were expressed in the floral transcriptome of *C. allenii*,

compared to five in *C. villosissimus*, and only one in *C. lasius*, when expression was defined as

one transcript per million (TPM) in at least one sample of the species. Furthermore, we found

that counts of TPS expression per TPS gene expressed were much higher in both *C. allenii* and

C. villosissimus when compared to *C. lasius* (Fig. 4B). In fact, in *C. allenii,* a TPS is the highest

expressed gene in the floral transcriptome, and TPSs are also the 4th and 6th most expressed genes

in the transcriptome, highlighting the extent of expression. On average, 12% of total expression

can be attributed to TPSs in *C. allenii*, 0.17% in *C. villosissimus* and 0.00014% in *C. lasius*,

highlighting the orders of magnitude differences between species.

Figure 4. Differences in floral scent and gene expression between *C. allenii*, *C. lasius*, and *C. villosissimus*. Species differ significantly in A) number of floral scent components (ANOVA, $F_{2,19}=20.9$, $p<0.001$), and (B) expression of terpene synthase transcripts (ANOVA, $F_{2,16}=508.4$, $p<0.001$).

<u>Floral scent loss is not driven by gene loss in *Costus*</u>

While the transcriptome assemblies can give us insight into levels of TPS expression, it is hard to determine gene copy number due to the redundancies in the assemblies. For this, we need genome data to allow us to determine the number of TPS genes in each species. There are currently three publicly available *Costus* genomes: *C. bracteatus* (bee), *C. lasius (hummingbird)*, and *C. pulverulentus* (hummingbird) (Valderrama et al. 2022; Harenčár et al. 2023). *Costus lasius* and *C. pulverulentus* represent two independent transitions to hummingbird pollination (Kay and Grossenbacher 2022). To generate a dataset of two hummingbird- and two bee-pollinated species, we assembled an additional high-quality reference genome for bee-pollinated *C. allenii* (Table S1 for assembly statistics, Figure S2 to see alignment with *C. lasius* genome). We generated high-quality whole genome annotations for each species with high completeness and consistency (Table 2).

We annotated between 63-75% of each genome as repeat elements (Tables S4, S5, S6,
S7). We found that, of the elements that could be classified, retrotransposable elements
represented the largest percentage of the genome. These are the most abundant form of repeat
elements found in angiosperm genomes. Many repeats were not classified into previously known
families, perhaps suggesting the existence of new repeat elements.

Table 1. *Costus allenii* genome assembly statistics

N50 (Mb)	120
L50	5
Total length (Gbp)	1.137
GC%	40.07
Number of Ns per 100 kbp	0
Longest contig/scaffold (Mbp)	143
Number of contigs/scaffolds	423
BUSCO complete	98.8% (1594/1614)
BUSCO single copy	90.8% (1465/1614)

302
303

304 Table 2. BUSCO scores for high quality annotations of each *Costus* genome searched against the
305 Embryophyta database.

	C. allenii	*C. bracteatus*	*C. lasius*	*C. pulverulentus*
Complete	1493 (92.5%)	1457 (90.2%)	1480 (91.7%)	1466 (90.9%)
Complete and single copy	1139 (70.6%)	1116 (69.1%)	1121 (69.5%)	1152 (71.4%%)
Complete and duplicated	354 (21.9%)	341 (21.1%)	359 (22.2%)	314 (19.5%)
Fragmented	12 (0.7%)	25 (1.5%)	19 (1.2%)	21 (1.3%)
Missing	109 (6.8%)	132 (8.3%)	115 (7.1%)	127 (7.8%)
Total searched	1614	1614	1614	1614
Consistent lineage placement (OMArk)	86%	87%	86%	87%
Total gene number	28,442	27,794	28,409	26,696

We annotated a total of 66 TPSs in *C. allenii*, 44 in *C. bracteatus*, 40 in *C. lasius*, and 44
in *C. pulverulentus*. All TPSs belonged to previously described TPS families TPS-a, TPS-b,
TPS-c, TPS-e/f, and TPS-g (Fig. 5). Those TPSs responsible for floral scent biosynthesis are
primarily found in the TPS-a, TPS-b, and TPS-g families. TPS-a members tend to produce
monoterpenes, TPS-b members sesquiterpenes, and TPS-g members either monoterpenes or
sesquiterpenes. We identified lineage-specific duplications in both TPS-a and TPS-b, as well as
TPS-e/f in the bee-pollinated *C. allenii*. We did not find any evidence for pseudogenization or
gene loss in the two hummingbird-pollinated species included in the gene family analyses, *C.*
lasius and *C. pulverulentus*, with TPS genes containing complete protein domains and start
codons.

We mapped the RNAseq reads both to the *C. allenii* genome and the *C. lasius* genome. In
C. allenii, we found a consistent percentage of successfully mapped reads attributed to TPSs as
with the transcriptome (12% transcriptome, 13% mapped to either genome). We found some
differences for both *C. lasius* and *C. villosissimus*. For the *C. villosissimus* transcriptome
mapping, we found 0.17% of reads were attributed to TPSs, in comparison to 0.009% when
mapped to either genome. Due to a lack of genome for *C. villosissimus*, it is unclear whether this
could be due to genetic changes or rearrangements not captured by the available genomes. For *C.*
lasius, we found only 0.00014% of reads mapped to the transcriptome were due to TPSs, in
contrast to 0.011% of the genome-mapped reads.

When *C. allenii* reads were mapped to the *C. allenii* genome, we found high levels of

multi-mapping. This is due to recent TPS duplications in *C. allenii* which are highly expressed.

To illustrate this, the highest gene expressed in *C. allenii* is *TPS49*, and the third most expressed

"gene" consists of reads that multimap to *TPS49* and *TPS52*. This clade is represented by only

one gene in each of the other species. Other recent lineage-specific duplications include *TPS2*,

TPS3, *TPS4*, and *TPS5* where the majority of reads multimap due to sequence similarity between

duplicates. Both of these clades have only one member in each of the other species with an

available genome sequence (Fig S3). This highlights the complexity of mapping to gene families

with recent duplications, especially when they are highly expressed, meaning that the typical

choice of ignoring multi-mapped reads will lose large percentages of mapped reads from these

duplications. We, therefore, decided to carry out our differential expression analysis using the *C.*

lasius genome.

By mapping all reads to the *C. lasius* genome, we identified 4790 genes differentially

expressed between *C. allenii* and *C. lasius*, 7678 between *C. allenii* and *C. villosissimus*, and

6191 between *C. lasius* and *C. villosissimus*. Of the 40 TPS genes found in the *C. lasius* genome,

15 were differentially expressed between *C. allenii* and *C. lasius*, with 13 of these upregulated in

C. allenii. This includes *TPS33*, orthologous to the *TPS49/52* clade, the most highly expressed

genes in *C. allenii*. We found 19 TPSs that were differentially expressed between *C. allenii* and

C. villosissimus, with 14 upregulated in *C. allenii* and 5 in *C. villosissimus*. Ten TPS genes were

differentially expressed between *C. lasius* and C. *villosissimus*, with 7 upregulated in *C.*

villosissimus and 3 in *C. lasius*. Overall, more TPSs were upregulated in *C. allenii* when

compared to either *C. lasius* or *C. villosissimus*. Our gene ontology analysis found little evidence

for functional enrichment in differentially expressed genes between species (Table S8).

Figure 5. Phylogenetic analysis of terpene synthase gene families. The phylogeny was constructed in IQ-TREE with 1000 bootstraps and model LG. TPS family classifications based on *Arabidopsis thaliana* are illustrated.

Discussion

Repeated evolutionary transitions provide unique opportunities to study the extent of convergence both at the phenotypic and molecular level, shedding light on the predictability of evolution. Transitions to hummingbird pollination are widespread in the Americas and are predicted to be associated with loss of floral scent, however, this has not been studied in a comparative framework across multiple independent transitions within the same genus. In this

study, we combine comparative chemical ecology with genomics and transcriptomics and

demonstrate that floral scent evolution in the genus *Costus* is highly associated with pollinator

type. We found that hummingbird-pollinated *Costus* species have lost or reduced their floral

scent across eight independent evolutionary origins, whereas bee-pollinated species in the

Americas exhibit either maintenance of floral scent or gain of a more diverse scent profile. We

find that this is related to downregulation of genes associated with floral scent production (in this

case terpene synthases) in floral tissues of hummingbird-pollinated species but is not

accompanied by loss of terpene synthase genes.

Bee-pollinated *Costus* species showed much greater variation in their floral scents than

hummingbird-pollinated species, with some having extremely diverse floral scents, yet others

producing far fewer compounds. Bee-pollinated species also show greater morphological

variation, possibly reflecting variation in orchid bee diversity (Kay and Grossenbacher 2022).

Furthermore, not all species in our dataset follow the expected patterns. For example, *Costus*

arabicus and *Costus vinosus* (bee-pollinated) produce little scent. It is unclear whether the

variation in floral scent between different bee-pollinated species is related to pollinator

preference, other abiotic or biotic factors, or historical contingency.

There are multiple hypotheses for why floral scent would be lost in hummingbird-

pollinated species. Firstly, it could be because there is simply no selection to maintain the

production due to poor sense of smell, or lack of preference shown for floral scent, by

hummingbirds (Goldsmith and Goldsmith 1982; Knudsen et al. 2004; Driver and Balakrishnan

2021). If this were the case, we might expect to find a correlation between length of time

following pollinator transition and extent of floral scent loss. However, our data shows that

downregulation of floral scent can occur on relatively quick evolutionary timescales. *Costus*

wilsonii, *C. lasius*, and *C. spiralis* are all hummingbird-pollinated species that are most closely

related to bee-pollinated species, representing more recent evolutionary shifts to hummingbird

pollination. Yet all three species produce little to no floral scent, suggesting there can be quick

breakdown of floral scent production following pollinator transitions. This observation is

consistent with previous studies that have shown that floral scent often evolves rapidly in

response to selection (Ramos and Schiestl 2019; Zu et al. 2020; Liu et al. 2024).

The rapid and repeated loss of scent in hummingbird-pollinated species suggests a shared
strong selection pressure against floral scent production. One potential mechanism is the
energetic cost of terpene production. A similar argument has been made in the case of eye
regression in cavefish. If vision is energetically costly to maintain, then it should be selected
against when no longer useful (Moran et al. 2015; Sifuentes-Romero et al. 2023). This is
supported by evidence of the energetic costs of maintaining a visual system (Moran et al. 2015).
The metabolic cost of terpene production is unknown, but is thought to be minimal (Raguso
2016; Pichersky and Raguso 2018). In fact, it has been argued that the complex network of
terpene metabolism, in which individual enzymes produce multiple products, makes terpene
pathways more efficient, compared with a linear pathway with separate enzymes and regulators
for each step (Lanier et al. 2023).

An alternative scenario to the energetic cost is that terpene production has ecological
costs in hummingbird-pollinated species. For example, there may be strong selection against bee
attraction (Castellanos et al. 2004; Bergamo et al. 2016). Many benefits to hummingbird,
compared with bee, pollination have been described. Birds do not consume pollen leading to less
pollen loss, and birds can travel longer distances between plants and promote higher gene flow,
which is beneficial to combat inbreeding depression (Thomson and Wilson 2008; Krauss et al.
2017; Dellinger et al. 2022). One caveat is that comparisons in pollination effectiveness between
bees and hummingbirds have not been carried out in orchid bees. Both male and female orchid
bees tend to forage over long distances, perhaps resulting in a more "bird-like" pollen transport
(Janzen 1971, 1981; Wikelski et al. 2010; Opedal et al. 2017; Kay and Grossenbacher 2022;
Gamba and Muchhala 2023). While pollinator efficiency is unknown in *Costus*, rapid floral scent
loss is congruent with the hypothesis of "bee-avoidance" following establishment of
hummingbird pollination.

A final piece of evidence to support bee-repellence is the large reduction in beta-ocimene
production in hummingbird-pollinated flowers relative to bee-pollinated flowers, as this
compound is a common insect pollinator attractant (Farré-Armengol et al. 2017). In *Mimulus*,
loss of monoterpenes, including beta-ocimene, maintains reproductive isolation between bee-
and hummingbird-pollinated species (Byers et al. 2014). Interestingly, we found that α- and β-
pinene were produced by both bee- and hummingbird-pollinated species. One potential reason is
that pinenes do not repel orchid bees but do repel other types of bees. Terpenoid-dominated

scents have been found to repel honey bees (Larue et al. 2016), and in particular α-pinene has a

repellent effect on honey bees (Fernandes et al. 2019). We speculate that bee-pollinated species

emit a mix of compounds, some of which potentially attract orchid bees, and some of which

potentially repel other pollinators to narrow attraction to orchid bees, specifically. Then when

floral scent is reduced in hummingbird-pollinated flowers, the compounds which are attractive to

orchid bees are lost but not those which are general bee repellants. Further experimental work is

needed to test these hypotheses.

We found that the floral scents of *Costus* are dominated by terpene compounds. Terpene

diversity in a given organism is mainly determined by the terpene synthase (TPS) gene family,

which carry out crucial steps in terpene formation (Tholl 2006). Neither hummingbird-pollinated

species with an available genome sequence (*C. lasius* and C. *pulverulentus*) exhibited loss of

TPS genes. There are multiple reasons why trait loss would be associated with changes in gene

regulation rather than gene loss. Firstly, these cases may represent recent transitions to

hummingbird pollination, and at least some of these TPSs that were responsible for floral scent

production will eventually be lost. This has been proposed as a "step-wise" model to

pseudogenization, whereby reduction in expression is followed by relaxed selection and then

loss-of-function mutations, as seen in an enzyme lost in marine mammals (Graham et al. 2023).

Alternatively, we might not expect any TPS genes to be lost even in the long term.

Terpenes, and therefore TPSs, are important for many other roles in plants. Plants use terpenes to

deter herbivores, to attract parasitoids or predators of herbivores, and as anti-fungal and anti-

bacterial compounds to prevent disease (Gershenzon and Dudareva 2007). These pleiotropic

functions of TPSs could explain why we observe downregulation in floral tissues specifically

rather than any pseudogenization. One argument against this is that in other systems TPS

pseudogenization does occur following pollinator transitions. In *Mimulus*, loss of monoterpene

production (specifically beta-ocimene), is not due to expression differences but loss-of-function

mutations (Peng et al. 2017). It is unclear whether pseudogenization will occur at some point in

Costus.

Although we did not observe gene loss in hummingbird-pollinated lineages, we did

observe extensive lineage-specific duplications in the bee-pollinated species *C. allenii*. *Costus*

allenii was one of the species with highest scent diversity in our dataset, and we found multiple

TPS genes to exhibit several recent lineage-specific duplication events. We also found extremely

high TPS expression, accounting for approximately 12% of expression in floral tissue. The gene

with the highest expression in floral tissue was found to be a TPS-a family member, with high

expression also found also for several TPS-b members. These two families produce

sesquiterpenes and monoterpenes, respectively, both of which are found in the floral scent of *C.*

allenii. In *C. lasius* and *C. villosissimus*, we found reduced TPS expression compared to *C.*

allenii, as expected based off their reduced floral scent production. We also saw high levels of

phenotypic variation in *C. villosissimus*, and there is no publicly available genome, making

interpretation of the results more difficult. We speculate that *C. allenii* has higher floral scent

production than closely related *C. villosissimus* because *C. allenii* grows as isolated individuals

in deeply shaded, perennially wet forest, whereas *C. villosissimus* grows at forest edges and large

treefall gaps, and often has conspecifics nearby (Chen and Schemske 2015). Thus *C. allenii* may

need more scent to attract pollinators. While the overall trends are clear, we hesitate to over-

interpret the gene expression data as we have only sampled at one timepoint and it is possible

that there is floral scent production happening earlier that is not captured in our data.

We observe more consistency and predictability in the phenotypic patterns than the

underlying molecular mechanisms. In many systems, parallel changes in floral scent have

accompanied independent pollinator shifts, including the current study (Knudsen et al. 2004;

Byers et al. 2014). This can also be seen in other reproductive shifts, such as transitions to self-

fertilization (Wozniak et al. 2022). In some systems, the underlying molecular mechanisms of

floral scent loss are also convergent, having been shown to involve the same loci in multiple

independent evolutionary events. For example, loss of floral scent in different *Mimulus* lineages

is due to changes at the same loci, including frameshifts, deletions, and potentially post-

transcriptional regulation (Peng et al. 2017). A similar pattern was found in two different systems

that have lost benzaldehyde floral scent production due to either evolution of self-fertilization

(Sas et al. 2016), or transition to hummingbird pollination (Amrad et al. 2016). The same locus

contains loss-of-function mutations in both cases (Raguso 2016). However, this is not universally

true. In *Capsella*, the loss of beta-ocimene following transition to self-fertilization in one lineage

is due to changes in subcellular enzyme localization, but QTL results suggests another locus is

responsible for scent loss in a related species (Wozniak et al. 2022). In this study, we saw no

evidence for pseudogenization. Instead, we found evidence that gene expression differences are

responsible for scent differences between *Costus* species, with scent loss likely due to changes
in, potentially tissue-specific, gene expression.

In summary, we have shown that *Costus* species exhibit repeated loss of floral scent
following transitions from bee to hummingbird pollination, suggesting a shared strong selection
pressure favoring scent loss. We also find bee species that have recently gained incredibly
diverse scents, which, at least in *C. allenii*, is associated with lineage-specific terpene synthase
duplications. We show that scent loss is not associated with loss of terpene synthase genes but
rather appears to be due to gene downregulation. This demonstrates the capacity for rapid
metabolic changes in response to selection following pollinator transitions. This system provides
us with the exciting opportunity to identify the cis- or trans-regulatory changes driving gene
expression differences in multiple independent transitions to hummingbird pollination, allowing
us to study the extent of genetic convergence.

Materials and Methods

<u>Floral scent sample collection</u>

We collected 100 samples of 30 species of *Costus* between 2019 and 2023 (Table S1).
All samples for *Costus glaucus*, *Costus laevis*, *Costus montanus*, *Costus stenophyllus*, and
Costus wilsonii were collected in Costa Rica (La Gamba, Las Cruces, Las Alturas, and
Monteverde). All other species were sampled in the UCSC greenhouses in Santa Cruz,
California, USA from plants composing the UCSC living *Costus* collection. For each day of
sampling, both floral scent samples and ambient samples were collected (with one exception: a
Costus scaber sample collected on April 8[th] 2019 was missing a control and so we used a control
from the nearest date available, March 7th). Scent was sampled using dynamic headspace
sampling from approximately 9 am until 3 pm. Flowers from living plants were placed inside
oven bags (Reynold's). A Spectrex PAS-500 air pump with Tygon tubing was used to pull
scented-air through a volatile trap for 6 hours at a flow rate of 200 ml air / min. The volatile trap
contained a filter of glass wool and 50 mg Porapak. We conditioned scent traps by passing
5 mL of hexane before usage. Volatiles were eluted from the trap using 400μl of hexane and
stored at -20°C.

<u>Chemical analysis</u>

Samples were analyzed using Agilent model 5977A mass-selective detector with an

Agilent GC model 7890B. Samples were run on a HP-5 Ultra Inert column (Agilent, 30m x

0.25mm, 0.25 μm). For analysis, one microliter of each sample was injected in splitless mode

using the ALS 7694 autosampler with helium as the carrier gas (250°C injector temperature).

The program started at 40 °C for 3 minutes and then rose at 5 °C/min to 210°C. The temperature

was held at 210°C for 1 minute, before rising at 20°C/min to 300°C where it was held for 2.5

minutes, and 315°C for a final minute.

Compounds were identified by comparing mass spectra and retention indices with

reference libraries. Compounds that were not found in at least two samples were removed. A

two-step process was used to filter datasets for floral scent components. First, compounds were

filtered to include only those which are found in five times higher amounts in floral than ambient

samples using the "filter_ambient_ratio" function from the *bouquet* package (Powers et al.

2023). This resulted in a list of ambient compounds which were not considered floral scent

compounds. Second, using the raw data matrices, we subtracted the ambient compound amounts

from the corresponding floral scent amount collected on the same date. We then removed the

ambient compounds identified by the package *bouquet* from this subtracted matrix to create the

final data matrix of floral scent.

<u>Statistical analyses of floral scent</u>

To visualize divergence in floral scent among samples we used nonmetric

multidimensional scaling (NMDS) using the "metaMDS" function in *vegan* (Oksanen et al.

2020). To test for differences both between pollination groups, and between floral and ambient

samples, we carried out a PERMANOVA (permutational multivariate analysis of variance) using

the "adonis2" function in *vegan* (Bray–Curtis distance matrix, 1000 permutations). To identify

which groups were significantly different, we carried out post hoc pairwise testing using the

"pairwise.perm.MANOVA" function in the *RVAideMemoire* package with a Bonferroni

correction (Hervé 2021). We did this for both filtered and unfiltered datasets.

We tested for differences in compound number produced by bee-pollinated and

hummingbird-pollinated species using ANOVA with a nested model including species within

pollinator group. To test for differences in the diversity of compounds produced by bee-

pollinated and hummingbird-pollinated species, we used the package *chemodiv* (Petrén et al.

2023). This package considers the biochemical and structural diversity of the compounds present

in a sample to calculate the overall chemodiversity per sample. First, we used the "calcDiv"

function to calculate the functional Hill diversity index for each sample. Then we carried out an

ANOVA test for the diversity indices, using a nested model of species nested within pollinator

group to test for differences in chemodiversity. We also tested for differences in compound

production using ANOVA. In this case a proxy for compound amount was used: the sum of the

total area of floral scent peaks in the sample, or the total ion abundance in the sample. To test for

difference in compound frequency between bee- and hummingbird-pollinated samples, we

carried out chi-squared tests and adjusted the p-values using the "p.adjust" function in R.

<u>Phylogenetic analyses of floral scent</u>

To investigate the evolution of floral scent in a phylogenetic context we used a previously

published phylogeny (Vargas et al. 2020). First, we trimmed the phylogeny to include only those

species for which we have scent data using the "drop.tip" function in *ape* (Paradis and Schliep

2018). To test for correlation between pollination group and the phylogenetic covariance matrix,

we used a two-block partial least squares analysis using the "two.b.pls" function in the *geomorph*

package (Adams and Otárola-Castillo 2013; Adams and Collyer 2018). A significant result

means that our power to test for differences between groups using phylogenetic simulation-based

is weakened due to group aggregation on the phylogeny. We then used the "phylosig" function in

the package *phytools* to test for phylogenetic signal in multiple traits (compound number,

compound diversity, compound amount, (*E*)-beta-ocimene production, and (E)-beta-

caryophyllene production) (Revell 2012). We also tested if floral scent data overall shows

phylogenetic signal using the multivariate generalization of Bloomberg's K (K_{mult}) (Adams

2014). Under Brownian motion, K_{mult} is expected to be 1. If $K_{mult}>1$, higher phylogenetic signal

is detected than the null expectation under Brownian motion. If $K_{mult}<1$, lower phylogenetic

signal is detected than the null expectation under Brownian motion. To calculate the K_{mult} of

Costus floral scent, we used the filtered dataset with 1000 iterations to test significance. We

carried out the analysis using the "physignal" function in the *geomorph* package (Adams and

Otárola-Castillo 2013). Due to the weak phylogenetic signal detected in both multivariate and

univariate analyses, we used the "procD.pgls" function in the *RRPP* package to test for

differences in traits between pollinator groups rather than a simulation-based ANOVA (Adams

and Collyer 2018; Collyer and Adams 2018). This function uses a method of randomizing

residuals in a permutation procedure to evaluate the significance of phenotypic differences

between groups considering the phylogeny and has higher statistical power than methods using

phylogenetic simulations (Adams and Collyer 2018). We set lambda at 4.14795e-05 as

calculated for the univariate variables. We corrected for multiple testing using the "p.adjust"

function in R.

To better understand how floral scent has evolved in the genus *Costus*, we carried out an

ancestral state reconstruction of the number of compounds produced. We used the function "ace"

in the *ape* package for ancestral character estimation (Paradis and Schliep 2018). We divided the

compound number trait into twelve categories, each of which were assigned a different shade of

blue. We plotted both the compound number for each phylogenetic tip label and internal node to

illustrate how floral scent has evolved in *Costus*. We overlaid the origins of hummingbird

pollination as described in (Vargas et al. 2020).

<u>Sample collection for RNAseq</u>

Live individuals of *Costus allenii* and *Costus villosissimus* were collected in Soberanía

National Park, Colón Province, Panama. Vouchers are deposited at University of Panama

(*Costus allenii*: 0132264, 0132265; *Costus villosissimus*: 0132270). Live individuals of *Costus*

*lasiu*s were collected in El Valle de Antón, Cocle Province, Panama. Voucher is deposited in

Michigan State University (MSC) herbarium (Kay 0321). For all three species, the individuals

used for RNAseq analyses were grown in greenhouses at UCSC. Freshly opened flowers of each

species were removed at approximately 9 am and flash frozen immediately in liquid nitrogen

before being stored at -80°C until extraction. Flowers of *Costus* open early in the morning and

fall off by mid-afternoon.

<u>RNA extraction and sequencing</u>

Frozen flowers were removed and dissected on dry ice. Tissue from the petals and labella

was broken down using a TissueLyser (Qiagen). RNA was then extracted using an RNeasy Mini

Kit (Qiagen) with Qiashredder spin columns (Qiagen) following standard protocols. Samples

were sent to Novogene for quality control, library preparation, and sequencing. Our final dataset

consisted of 19 samples: 8 *Costus allenii* (from 4 individual plants), 6 *Costus lasius* (from 2

individual plants), and 5 *Costus villosissimus* (from 1 individual plant). Samples were sequenced

on a NovaSeq 6000 using 150 bp paired end reads. This generated approximately 53 million

reads per library (mean=52.74 million, SD=12.01 million, N=19). The raw sequence data were

deposited in the SRA under the BioProject accession PRJNA1010186.

<u>Transcriptome assembly</u>

Before assembling transcriptomes, we trimmed the raw reads using TrimGalore! (Martin

2011). We then assembled transcriptomes for each species using Trinity version 2.5.1 with

default settings (Grabherr et al. 2011; Haas et al. 2013). To reduce redundancy, we clustered

contigs with a minimum sequence identity of 95% using CD-HIT version 4.6.8 (Fu et al. 2012).

We assessed transcriptome completeness and quality using BUSCO (v5.6.1)

(embryophyta_odb10 dataset) (Simão et al. 2015; Manni et al. 2021; Zdobnov et al. 2021). We

also checked assembly quality by mapping RNAseq reads back to the assembly using Bowtie

(v2.4.1) to calculate mapping rates (Langmead and Salzberg 2012).

<u>Terpene synthase gene expression analysis</u>

We extracted the longest isoforms for each Trinity "gene" using the Trinity script

"get_longest_isoform_seq_per_trinity_gene.pl" and translated these sequences. We then

searched for potential TPSs using hmmsearch two HMM profiles: Terpene_syth_C (PF03936)

and Terpene Synthase N-terminal domain (PF01397). We used the R package rhmmer to parse

the output from hmmsearch resulting in a list of putative TPS sequences for each species. To

reduce the TPSs to only those expressed in the dataset, we used a threshold of TPM>1

(transcripts per million) in at least one individual of a species (Eddy 2011; Arendsee 2017). To

calculate expression of each putative TPS we normalized counts using Gene length corrected

trimmed mean of M-values (GeTMM), an approach that improves inter- and intra-sample

comparisons (Smid et al. 2018). To do this, we first calculated FPKM (fragments per kilobase of

transcript per million fragments mapped) for each transcript and carried out TMM-normalization

using the package *edgeR* (Robinson et al. 2010). We then combined GeTMM values for

transcripts which were identified by Trinity to be the same gene. This does not normalize counts

between species but is used to determine whether genes are lowly or highly expressed within a

species.

<u>*Costus allenii* genome assembly</u>

We selected a wild-collected individual *C. allenii* from Pipeline Road, Soberanía

National Park, Colón Province, Panama for *de novo* genome assembly. A voucher for this

population of *C. allenii* is deposited at the University of Panama Herbarium (PMA:0132264).

This individual was grown to flowering in the UCSC greenhouses from a cutting, and we

harvested 1.5 g fresh tissue from leaf meristems and flower buds. We isolated cell nuclei

following Workman (Workman et al. 2019) and then used the Circulomics Nanobind Plant

Nuclei Big DNA kit to extract HMW DNA. We assessed DNA concentration with a Qubit

Fluorometer (Thermo Fisher Scientific), purity with 260/280 and 260/230 absorbance ratios from

a NanoDrop Spectrophotometer (Thermo Fisher Scientific) and ran a 7% agarose gel at 70 V for

over 8 h to assess DNA fragment integrity and size. HMW DNA was sent to the UC Davis

Genomics Center for additional cleaning, Pacific Biosciences HiFi library preparation, and

sequencing on one PacBio Revio SMRT cell. This produced 74.8 Gb of HiFi yield and 7.9 M

HiFi reads. We assembled the PacBio data into a primary and alternative assembly using

HiFiasm (v0.19.8)(Cheng et al. 2021). We then used RagTag (v2.1.0)(Alonge et al. 2022) to

scaffold *C. allenii* contigs using the closely related *C. lasius* as a reference. We used D-GENIES

(Cabanettes and Klopp 2018) with Minimap2 (Li 2018) to generate a dotplot comparing the two

assemblies both before and after scaffolding. The genome assembly quality was assessed using

BUSCO v 5.6.1 (Simão et al. 2015). Raw sequence data and the final genome assembly were

submitted to NCBI under BioProject PRJNA1091680.

<u>Genome annotations</u>

Of the three species for which we generated RNAseq data, an unannotated genome was

previously available for *C. lasius* (Harenčár et al. 2023). In this study we also generated a

genome assembly for *C.allenii*. In addition, previously published genomes are available for both

C. bracteatus and the more distantly-related *C. pulverulentus* (Valderrama et al. 2022; Harenčár

et al. 2023). We started by processing RNAseq data as evidence for annotations. Trimmed reads

from *C. lasius* were mapped using 2pass mapping in STAR to the *C. lasius* genome, and the *C.*

pulverulentus genomes (Martin 2011; Dobin et al. 2013). Bam files were concatenated for use as

described below. Trimmed reads from *C. allenii* were mapped using 2pass mapping in STAR to

both the *C. allenii* and the closely related *C. bracteatus* genomes, and bam files concatenated per

species (Martin 2011; Dobin et al. 2013).

We annotated the genomes using BRAKER3 (v3.0.7) which combines RNA-seq and

protein data in an automated pipeline (Lomsadze et al. 2005, 2014; Stanke et al. 2006, 2008;

Gotoh 2008; Iwata and Gotoh 2012; Buchfink et al. 2015; Hoff et al. 2016, 2019; Kovaka et al.

2019; Brůna et al. 2020, 2021; Pertea and Pertea 2020; Gabriel et al. 2023). This pipeline

requires a masked genome. We identified and annotated tandem repeats and transposable

elements using RepeatModeler (Flynn et al. 2020) (v.2.0.4 for *C. lasius* and v.2.0.5 for the other

genomes) and RepeatMasker v.4.1.5 (Tarailo-Graovac and Chen 2009). We then used

RepeatMasker with the outputs of RepeatModeler to generate softmasked genomes. The masked

genomes were then provided to BRAKER along with plant proteins downloaded from OrthoDB

(odb10_plants) (Zdobnov et al. 2021). For all species, BRAKER was run in ETP mode which

combines evidence from both proteins and RNAseq (Gabriel et al. 2021). BUSCO v 5.6.1

(Simão et al. 2015) and OMArk (Nevers et al. 2022) were used to evaluate the completeness and

consistency of the sets of gene models. During the project, the *C. lasius* genome was updated

(direction of arms on chromosome 1 were altered) and so we re-scaffolded chromosome 1 of *C.*

allenii and transferred our annotation using liftoff (Shumate and Salzberg 2021).

<u>Terpene synthase gene family annotation</u>

Following whole-genome annotation, we manually curated the gene models for terpene

synthase (TPS) genes in the annotation. To do this, we used bitacora (Vizueta et al. 2020), a

pipeline which curates an existing annotation and finds additional gene family members. In this

pipeline, BLASTP and hmmer are used to search for gene family members of interest within the

existing annotation (Altschul et al. 1990; Eddy 2011). Additional regions of the genome are then

searched using TBLASTN and are annotated using GeMoMa (Keilwagen et al. 2019). Hmmer is

used to validate the gene models. As input, we used two HMM profiles: Terpene_syth_C

(PF03936) and Terpene Synthase N-terminal domain (PF01397) downloaded from Pfam. In

addition, we provided previously annotated genes from *Arabidopsis thaliana* and *Oryza sativa*

(Chen et al. 2011; Yu et al. 2020; Jia et al. 2022). Gene models from bitacora and the genome-

wide annotation from BRAKER were compared and TPS genes were manually curated using

IGV (Robinson et al. 2011). We also searched for additional TPSs by carrying out exonerate

searches using the curated *Costus* gene models (Slater and Birney 2005). We checked all

annotated TPSs for complete protein domains using the NCBI conserved domain search

(Marchler-Bauer et al. 2015). We excluded proteins with less than 250 amino acids.

<u>Gene family analysis</u>

To classify these TPSs in previously described TPS families, we constructed a

phylogenetic tree based on amino acid sequences with TPSs from *Arabidopsis thaliana*. The

Costus TPS protein sequences, combined with those from *A. thaliana*, were aligned with

MAFFT v7.508 (-maxiterate 1000, using L-INS-I algorithm) (Katoh and Standley 2013). We

then trimmed the alignment using trimAl (-gt0.6, sites only included when present in 60% of

sequences) (Capella-Gutierrez et al. 2009). We used this alignment to construct a gene tree with

IQ-TREE and the ModelFinder function to determine the best-fit model (Nguyen et al. 2015;

Kalyaanamoorthy et al. 2017; Hoang et al. 2018). We rooted the tree using the *phytools* package

in R ("midpoint.root" function)(Revell 2012; R Core Team 2023). We plotted the resulting gene

trees using the following packages in R: *ape* (Paradis and Schliep 2018), *evobiR* (Blackmon and

Adams 2015), *ggnewscale* (Campitelli 2023), *ggstar* (Xu 2022), *ggtree* (Yu et al. 2017),

ggtreeExtra (Xu et al. 2021), and *tidytree* (Yu 2022).

<u>Differential expression analyses</u>

First, we trimmed the reads using TrimGalore! (Martin 2011). We then mapped the reads

from all three species to both the *Costus allenii* and *Costus lasius* genome (Harenčár et al. 2023).

We ran 2pass mapping using STAR (Dobin et al. 2013). In general, we found higher levels of

multi-mapped *C. allenii* reads when mapped to *C. allenii*. To investigate this we used mmquant

(with merge, -m, set to 10) to count the reads, a tool which can account for multi-mapping reads

(Zytnicki 2017). The output from mmquant was read into R for further analysis. To calculate

percentage of gene expression due to TPS expression, we normalized reads by library size using

TMM (trimmed mean of M values) (Robinson and Oshlack 2010). Due to the consistency in TPS

expression as calculated by mapping to both genomes, we decided to use the *C. lasius* genome

assembly to identify genes differentially expressed between the three species to remove the

complication of multi-mapping. To do this, we first counted reads using featureCounts (Liao et

al. 2014) and again read the output into R. We filtered genes to include those with at least one

count per million (CPM) in at least one library and normalized reads by library size using TMM

(trimmed mean of M values). Differential gene expression was evaluated using the non-

parametric "noiseqbio" function from the *NOISeq* package (Tarazona et al. 2011, 2015). We

used the default cut-off of q=0.95 in the "degenes" function which is equivalent to an adjusted p-
value of 0.05.

<u>GO Analysis</u>

We functionally annotated *C. alennii* genes using Blast2GO (Götz et al. 2008; Burge et
al. 2012). We imported these functional annotations into R to carry out enrichment analysis with
the package *topGO* (Alexa and Rahnenfuhrer 2022). We considered the background to be all
genes included in that specific differential expression analysis after filtering.

<u>Plotting and data manipulation</u>

Additional packages used for general plotting: *cowplot* (Wilke 2020), *ggplot2* (Wickham
2009), and *scico* (Pedersen and Crameri 2023). Packages used for data transformation and
manipulation include: *data.table* (Barrett et al. 2024), *dplyr* (Wickham et al. 2021), *knitr* (Xie
2023), *magrittr* (Bache and Wickham 2023), *readr* (Wickham et al. 2023), *reshape2* (Wickham
2007), and *tibble* (Müller and Wickham 2022). Analyses were carried out in R version 4.3.2 (R
Core Team 2023).

Data availability

Data and R scripts used for analysis are available from Open Science Framework:
https://osf.io/2ap4k/?view_only=6b8f5ff7d2ae499b871e1b1fb6fe6219. The raw sequence reads
and final genome assembly for *Costus allenii* are deposited in the SRA (BioProject
PRJNA1091680)
https://dataview.ncbi.nlm.nih.gov/object/PRJNA1091680?reviewer=p4jo1ioj3vqualqk1kvege7aq
l. The raw RNAseq data is deposited in the SRA (BioProject PRJNA1010186)
https://dataview.ncbi.nlm.nih.gov/object/PRJNA1010186?reviewer=fil9vpq6up5g3pj7e654rtokje

Acknowledgements

This work was funded by the National Science Foundation Dimensions of Biodiversity

grant awarded to K.M.K. (DEB-1737889) and S.R.R. (DEB-1737771), and the Jean H.

Langenheim Chair in Plant Ecology and Evolution held by K.M.K. We thank the Organization

for Tropical Studies, Estación Biológica Monteverde, and Estación Tropical La Gamba for

facilitating field work in Costa Rica and the Smithsonian Tropical Research Institute for

facilitating field work in Panama. All field research was conducted with appropriate research

permits in Costa Rica (M-P-SINAC-PNI-ACAT-026-2018, ACC-PI-0272018, MPC-SINAC-

PNI-ACLAP-020-2018, INV-ACOSA-076-18, MPC-SINAC-PNI-ACTo-020-18, CONAGEBIO

R-056-2019OT and R-058-2019-OT) and Panama (SE/AP-13-19, 09-99) We thank Julia

Harenčár for assistance with lab work, Pedro Juarez and Cecilia Girvin for assistance with field

sampling, and Alyssa Kaatmann, Aubrie Tait, Kate Uckele, Melina Sauerman Carrizosa, and

Selena Vengco for assistance with greenhouse sampling. We thank J. Velzy and S. Childress for

greenhouse care of the UCSC living *Costus* collection. We thank the Ramírez lab for helpful

discussions.

Author Contributions

758   Conceptualization, K.D., K.M.K., and S.R.R.; Investigation, K.D. and K.M.K.; Formal

759   Analysis, K.D.; Visualization, K.D.; Resources, K.M.K. and S.R.R.; Writing – Original Draft,

760   K.D.; Writing – Review and Editing, K.D., K.M.K., and S.R.R.; Supervision, K.M.K. and

761   S.R.R.; Project Administration, K.M.K. and S.R.R.; Funding Acquisition, K.M.K. and S.R.R.

Declaration of Interests

764   The authors declare no competing interests.

References

References

Abrahamczyk, S., and S. S. Renner. 2015. The temporal build-up of hummingbird/plant

mutualisms in North America and temperate South America. BMC Evol. Biol. 15:104.

Adams, D. C. 2014. A generalized K statistic for estimating phylogenetic signal from shape and

other high-dimensional multivariate data. Syst. Biol. 63:685–697.

Adams, D. C., and M. L. Collyer. 2018. Phylogenetic ANOVA: Group-clade aggregation,

biological challenges, and a refined permutation procedure. Evolution 72:1204–1215.

Adams, D. C., and E. Otárola-Castillo. 2013. geomorph: an r package for the collection and

analysis of geometric morphometric shape data. Methods Ecol. Evol. 4:393–399.

Albert, V. A., S. E. Williams, and M. W. Chase. 1992. Carnivorous Plants: Phylogeny and

Structural Evolution. Science 257:1491–1495. American Association for the

Advancement of Science.

Alexa, A., and J. Rahnenfuhrer. 2022. topGO: Enrichment Analysis for Gene Ontology.

Alonge, M., L. Lebeigle, M. Kirsche, K. Jenike, S. Ou, S. Aganezov, X. Wang, Z. B. Lippman,

M. C. Schatz, and S. Soyk. 2022. Automated assembly scaffolding using RagTag elevates

a new tomato system for high-throughput genome editing. Genome Biol. 23:258.

Amrad, A., M. Moser, T. Mandel, M. de Vries, R. C. Schuurink, L. Freitas, and C. Kuhlemeier.

2016. Gain and Loss of Floral Scent Production through Changes in Structural Genes

during Pollinator-Mediated Speciation. Curr. Biol. 26:3303–3312.

Arendsee, Z. 2017. rhmmer: utilities parsing "HMMER" results.

Bache, S. M., and H. Wickham. 2023. magrittr: A Forward-Pipe Operator for R.

Barreto, E., M. M. A. Boehm, E. Ogutcen, S. Abrahamczyk, M. Kessler, J. Bascompte, A. S.

Dellinger, C. Bello, D. M. Dehling, F. Duchenne, M. Kaehler, L. P. Lagomarsino, L. G.

Lohmann, M. A. Maglianesi, H. Morlon, N. Muchhala, J. F. Ornelas, M. Perret, N. R.

Salinas, S. D. Smith, J. C. Vamosi, I. G. Varassin, and C. H. Graham. 2024.

Macroevolution of the plant–hummingbird pollination system. Biol. Rev. 000.

Barrett, T., M. Dowle, A. Srinivasan, J. Gorecki, M. Chirico, and T. Hocking. 2024. data.table:

Extension of `data.frame`.

Bergamo, P. J., A. R. Rech, V. L. G. Brito, and M. Sazima. 2016. Flower colour and visitation

rates of *Costus arabicus* support the 'bee avoidance' hypothesis for red-reflecting

hummingbird-pollinated flowers. Funct. Ecol. 30:710–720.

Blackmon, H., and R. A. Adams. 2015. EvobiR: Tools for comparative analyses and teaching

evolutionary biology. , doi: 10.5281/zenodo.30938.

Burge, S., E. Kelly, D. Lonsdale, P. Mutowo-Muellenet, C. McAnulla, A. Mitchell, A.

Sangrador-Vegas, S.-Y. Yong, N. Mulder, and S. Hunter. 2012. Manual GO annotation

of predictive protein signatures: the InterPro approach to GO curation. Database J. Biol.

Databases Curation 2012:bar068.

Byers, K. J. R. P., H. D. Bradshaw, and J. A. Riffell. 2014. Three floral volatiles contribute to

differential pollinator attraction in monkeyflowers (*Mimulus*). J. Exp. Biol. 217:614–623.

Cabanettes, F., and C. Klopp. 2018. D-GENIES: dot plot large genomes in an interactive,

efficient and simple way. PeerJ 6:e4958. PeerJ Inc.

Campitelli, E. 2023. ggnewscale: Multiple Fill and Colour Scales in "ggplot2."

Capella-Gutierrez, S., J. M. Silla-Martinez, and T. Gabaldon. 2009. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

Castellanos, M. C., P. Wilson, and J. D. Thomson. 2004. 'Anti-bee' and 'pro-bird' changes

during the evolution of hummingbird pollination in *Penstemon* flowers. J. Evol. Biol.

17:876–885.

Chen, G. F., and D. W. Schemske. 2015. Ecological differentiation and local adaptation in two

sister species of Neotropical *Costus* (Costaceae). Ecology 96:440–449.

Chen, L., A. L. DeVries, and C.-H. C. Cheng. 1997. Convergent evolution of antifreeze

glycoproteins in Antarctic notothenioid fish and Arctic☐cod. Proc. Natl. Acad. Sci.

94:3817–3822. Proceedings of the National Academy of Sciences.

Cheng, H., G. T. Concepcion, X. Feng, H. Zhang, and H. Li. 2021. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. Nat. Methods 18:170–175. Nature

Publishing Group.

Collyer, M. L., and D. C. Adams. 2018. RRPP: An r package for fitting linear models to high-

dimensional data using residual randomization. Methods Ecol. Evol. 9:1772–1779.

Dellinger, A. S., O. Paun, J. Baar, E. M. Temsch, D. Fernández-Fernández, and J.

Schönenberger. 2022. Population structure in Neotropical plants: Integrating pollination

biology, topography and climatic niches. Mol. Ecol. 31:2264–2280.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson,

and T. R. Gingeras. 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics

29:15–21.

Driver, R. J. 2022. Evolution of olfactory receptors in birds. East Carolina University.

Driver, R. J., and C. N. Balakrishnan. 2021. Highly Contiguous Genomes Improve the

Understanding of Avian Olfactory Receptor Repertoires. Integr. Comp. Biol. 61:1281–

1290.

Eddy, S. R. 2011. Accelerated Profile HMM Searches. PLOS Comput. Biol. 7:e1002195.

Farré-Armengol, G., I. Filella, J. Llusià, and J. Peñuelas. 2017. β-Ocimene, a Key Floral and

Foliar Volatile Involved in Multiple Interactions between Plants and Other Organisms.

Molecules 22:1148.

Fernandes, N. de S., F. A. N. Silva, F. A. S. de Aragão, G. J. Zocolo, and B. M. Freitas. 2019.

Volatile Organic Compounds Role in Selective Pollinator Visits to Commercial Melon

Types. J. Agric. Sci. 11:p93.

Fu, L., B. Niu, Z. Zhu, S. Wu, and W. Li. 2012. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinformatics 28:3150–3152.

Gamba, D., and N. Muchhala. 2023. Pollinator type strongly impacts gene flow within and

among plant populations for six Neotropical species. Ecology 104:e3845.

Gershenzon, J., and N. Dudareva. 2007. The function of terpene natural products in the natural

world. Nat. Chem. Biol. 3:408–414.

Goldsmith, K. M., and T. H. Goldsmith. 1982. Sense of Smell in the Black-Chinned

Hummingbird. The Condor 84:237.

Götz, S., J. M. García-Gómez, J. Terol, T. D. Williams, S. H. Nagaraj, M. J. Nueda, M. Robles,

M. Talón, J. Dopazo, and A. Conesa. 2008. High-throughput functional annotation and

data mining with the Blast2GO suite. Nucleic Acids Res. 36:3420–3435.

Grabherr, M. G., B. J. Haas, M. Yassour, J. Z. Levin, D. A. Thompson, I. Amit, X. Adiconis, L.

Fan, R. Raychowdhury, Q. Zeng, Z. Chen, E. Mauceli, N. Hacohen, A. Gnirke, N. Rhind,

F. di Palma, B. W. Birren, C. Nusbaum, K. Lindblad-Toh, N. Friedman, and A. Regev.

2011. Full-length transcriptome assembly from RNA-Seq data without a reference

genome. Nat. Biotechnol. 29:644–652. Nature Publishing Group.

Graham, A. M., J. M. Jamison, M. Bustos, C. Cournoyer, A. Michaels, J. S. Presnell, R. Richter,

D. E. Crocker, A. Fustukjian, M. E. Hunter, L. D. Rea, J. Marsillach, C. E. Furlong, W.

K. Meyer, and N. L. Clark. 2023. Reduction of Paraoxonase Expression Followed by

Inactivation across Independent Semiaquatic Mammals Suggests Stepwise Path to

Pseudogenization. Mol. Biol. Evol. 40:msad104.

Grant, K. A., and V. Grant. 1968. Hummingbirds and their flowers. Columbia University Press,

New York.

Haas, B. J., A. Papanicolaou, M. Yassour, M. Grabherr, P. D. Blood, J. Bowden, M. B. Couger,

D. Eccles, B. Li, M. Lieber, M. D. MacManes, M. Ott, J. Orvis, N. Pochet, F. Strozzi, N.

Weeks, R. Westerman, T. William, C. N. Dewey, R. Henschel, R. D. LeDuc, N.

Friedman, and A. Regev. 2013. De novo transcript sequence reconstruction from RNA-

seq using the Trinity platform for reference generation and analysis. Nat. Protoc. 8:1494–

1512. Nature Publishing Group.

Harenčár, J., O. M. Vargas, M. Escalona, D. W. Schemske, and K. M. Kay. 2023. Genome

assemblies and comparison of two neotropical spiral gingers: *Costus pulverulentus* and *C.*

lasius. J. Hered. esad018.

Hervé, M. 2021. RVAideMemoire: Testing and Plotting Procedures for Biostatistics. R package

version 0.9-81.

Hoang, D. T., O. Chernomor, A. von Haeseler, B. Q. Minh, and L. S. Vinh. 2018. UFBoot2:

Improving the Ultrafast Bootstrap Approximation. Mol. Biol. Evol. 35:518–522. Oxford

Academic.

Janzen, D. H. 1981. Bee Arrival at Two Costa Rican Female Catasetum Orchid Inflorescences,

and a Hypothesis on Euglossine Population Structure. Oikos 36:177–183.

Janzen, D. H. 1971. Euglossine Bees as Long-Distance Pollinators of Tropical Plants. Science

171:203–205.

Kalyaanamoorthy, S., B. Q. Minh, T. K. F. Wong, A. von Haeseler, and L. S. Jermiin. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods

14:587–589.

Katoh, K., and D. M. Standley. 2013. MAFFT Multiple Sequence Alignment Software Version

7: Improvements in Performance and Usability. Mol. Biol. Evol. 30:772–780. Oxford

Academic.

Kay, K. M., and D. L. Grossenbacher. 2022. Evolutionary convergence on hummingbird

pollination in Neotropical *Costus* provides insight into the causes of pollinator shifts.

New Phytol. 236:1572–1583.

Kay, K. M., and D. W. Schemske. 2003. Pollinator Assemblages and Visitation Rates for 11

Species of Neotropical *Costus* (Costaceae). Biotropica 35:198–207.

Knudsen, J. T., L. Tollsten, I. Groth, G. Bergstöm, and R. A. Raguso. 2004. Trends in floral

scent chemistry in pollination syndromes: floral scent composition in hummingbird-

pollinated taxa. Bot. J. Linn. Soc. 146:191–199.

Krauss, S. L., R. D. Phillips, J. D. Karron, S. D. Johnson, D. G. Roberts, and S. D. Hopper. 2017.

Novel Consequences of Bird Pollination for Plant Mating. Trends Plant Sci. 22:395–410.

Langmead, B., and S. L. Salzberg. 2012. Fast gapped-read alignment with Bowtie 2. Nat.

Methods 9:357–359. Nature Publishing Group.

Lanier, E. R., T. B. Andersen, and B. Hamberger. 2023. Plant terpene specialized metabolism:

complex networks or simple linear pathways? Plant J. 114:1178–1201.

Larue, A.-A. C., R. A. Raguso, and R. R. Junker. 2016. Experimental manipulation of floral

scent bouquets restructures flower–visitor interactions in the field. J. Anim. Ecol.

85:396–408.

Li, H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34:3094–

3100.

Liao, Y., G. K. Smyth, and W. Shi. 2014. featureCounts: an efficient general purpose program

for assigning sequence reads to genomic features. Bioinforma. Oxf. Engl. 30:923–930.

Liu, J. W., P. Milet-Pinheiro, G. Gerlach, M. Ayasse, C. E. P. Nunes, I. Alves-Dos-Santos, and

S. R. Ramírez. 2024. Macroevolution of floral scent chemistry across radiations of male

euglossine bee-pollinated plants. Evolution 78:98–110.

Losos, J. B. 2011. CONVERGENCE, ADAPTATION, AND CONSTRAINT. Evolution

65:1827–1840.

Maas, P. J. M. 1972. Costoideae (Zingiberaceae). Flora Neotropica 8:1–139. New York

Botanical Garden Press.

Maas, P. J. M. 1977. Renealmia (Zingiberaceae - Zingiberoideae) Costoideae (Additions)

(Zingiberaceae). Flora Neotropica 18:1–218. New York Botanical Garden Press.

Manni, M., M. R. Berkeley, M. Seppey, F. A. Simão, and E. M. Zdobnov. 2021. BUSCO

Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic

Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol. Biol. Evol.

38:4647–4654.

Martin, M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet.journal 17:10–12.

Moran, D., R. Softley, and E. J. Warrant. 2015. The energetic cost of vision and the evolution of

eyeless Mexican cavefish. Sci. Adv. 1:e1500363. American Association for the

Advancement of Science.

Müller, K., and H. Wickham. 2022. tibble: Simple Data Frames.

Nguyen, L.-T., H. A. Schmidt, A. von Haeseler, and B. Q. Minh. 2015. IQ-TREE: A Fast and

Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. Mol.

Biol. Evol. 32:268–274.

Oksanen, J., F. Guillaume Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P.

Minchin, R. O'Hara, G. Simpson, P. Solymos, H. Stevens, E. Szoecs, and H. Wagner.

2020. vegan: Community Ecology Package. R package version 2.5-7.

Opedal, Ø. H., M. Falahati-Anbaran, E. Albertsen, W. S. Armbruster, R. Pérez-Barrales, H. K.

Stenøien, and C. Pélabon. 2017. Euglossine bees mediate only limited long-distance gene

flow in a tropical vine. New Phytol. 213:1898–1908.

Orr, H. A. 2005. The Probability of Parallel Evolution. Evolution 59:216–220.

Paradis, E., and K. Schliep. 2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics, doi: 10.1093/bioinformatics/bty633.

Pedersen, T. L., and F. Crameri. 2023. scico: Colour Palettes Based on the Scientific Colour-

Maps.

Peng, F., K. J. R. P. Byers, and H. D. Bradshaw Jr. 2017. Less is more: Independent loss-of-

function OCIMENE SYNTHASE alleles parallel pollination syndrome diversification in

monkeyflowers (*Mimulus*). Am. J. Bot. 104:1055–1059.

Petrén, H., T. G. Köllner, and R. R. Junker. 2023. Quantifying chemodiversity considering

945   biochemical and structural properties of compounds with the R package chemodiv. New

946   Phytol. 237:2478–2492.

947   Pichersky, E., and R. A. Raguso. 2018. Why do plants produce so many terpenoid compounds?

948   New Phytol. 220:692–702.

949   Powers, J., K. Eisen, D. Campbell, and R. Raguso. 2023. bouquet: Filter and Augment Biological

950   GC-MS Data.

951   R Core Team. 2023. R: A language and environment for statistical computing. R Foundation for

952   Statistical Computing, Vienna, Austria.

953   Raguso, R. A. 2016. Plant Evolution: Repeated Loss of Floral Scent — A Path of Least

954   Resistance? Curr. Biol. 26:R1282–R1285.

955   Ramos, S. E., and F. P. Schiestl. 2019. Rapid plant evolution driven by the interaction of

956   pollination and herbivory. Science 364:193–196. American Association for the

957   Advancement of Science.

958   Revell, L. J. 2012. phytools: an R package for phylogenetic comparative biology (and other

959   things). Methods Ecol. Evol. 3:217–223.

960   Robinson, M. D., D. J. McCarthy, and G. K. Smyth. 2010. edgeR: a Bioconductor package for

961   differential expression analysis of digital gene expression data. Bioinformatics 26:139–

962   140.

963   Robinson, M. D., and A. Oshlack. 2010. A scaling normalization method for differential

964   expression analysis of RNA-seq data. Genome Biol. 11:R25.

Rosas-Guerrero, V., R. Aguilar, S. Martén-Rodríguez, L. Ashworth, M. Lopezaraiza-Mikel, J.

M. Bastida, and M. Quesada. 2014. A quantitative review of pollination syndromes: do

floral traits predict effective pollinators? Ecol. Lett. 17:388–400.

Sackton, T. B., and N. Clark. 2019. Convergent evolution in the genomics era: new insights and

directions. Philos. Trans. R. Soc. B Biol. Sci. 374:20190102. Royal Society.

Sas, C., F. Müller, C. Kappel, T. V. Kent, S. I. Wright, M. Hilker, and M. Lenhard. 2016.

Repeated Inactivation of the First Committed Enzyme Underlies the Loss of

Benzaldehyde Emission after the Selfing Transition in *Capsella*. Curr. Biol. 26:3313–

3319.

Schemske, D. W., and H. D. Bradshaw. 1999. Pollinator preference and the evolution of floral

traits in monkeyflowers (*Mimulus*). Proc. Natl. Acad. Sci. 96:11910–11915. Proceedings

of the National Academy of Sciences.

Shumate, A., and S. L. Salzberg. 2021. Liftoff: accurate mapping of gene annotations.

Bioinformatics 37:1639–1643.

Sifuentes-Romero, I., A. M. Aviles, J. L. Carter, A. Chan-Pong, A. Clarke, P. Crotty, D.

Engstrom, P. Meka, A. Perez, R. Perez, C. Phelan, T. Sharrard, M. I. Smirnova, A. J.

Wade, and J. E. Kowalko. 2023. Trait Loss in Evolution: What Cavefish Have Taught Us

about Mechanisms Underlying Eye Regression. Integr. Comp. Biol. 63:393–406.

Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov. 2015.

BUSCO: assessing genome assembly and annotation completeness with single-copy

orthologs. Bioinformatics 31:3210–3212.

Slater, G. S. C., and E. Birney. 2005. Automated generation of heuristics for biological sequence

comparison. BMC Bioinformatics 6:31.

Smid, M., R. R. J. Coebergh van den Braak, H. J. G. van de Werken, J. van Riet, A. van Galen,

V. de Weerd, M. van der Vlugt-Daane, S. I. Bril, Z. S. Lalmahomed, W. P. Kloosterman,

S. M. Wilting, J. A. Foekens, J. N. M. IJzermans, P. P. L. O. Coene, J. W. T. Dekker, D.

D. E. Zimmerman, G. W. M. Tetteroo, W. J. Vles, W. W. Vrijland, R. Torenbeek, M.

Kliffen, J. H. Carel Meijer, A. A. vd Wurff, J. W. M. Martens, A. M. Sieuwerts, and on

behalf of the MATCH study group. 2018. Gene length corrected trimmed mean of M-

values (GeTMM) processing of RNA-seq data performs similarly in intersample analyses

while improving intrasample comparisons. BMC Bioinformatics 19:236.

Tarazona, S., P. Furió-Tarí, D. Turrà, A. D. Pietro, M. J. Nueda, A. Ferrer, and A. Conesa. 2015.

Data quality aware analysis of differential expression in RNA-seq with NOISeq R/Bioc

package. Nucleic Acids Res. 43:e140.

Tarazona, S., F. García-Alcalde, J. Dopazo, A. Ferrer, and A. Conesa. 2011. Differential

expression in RNA-seq: a matter of depth. Genome Res. 21:2213–2223.

Tholl, D. 2006. Terpene synthases and the regulation, diversity and biological roles of terpene

metabolism. Curr. Opin. Plant Biol. 9:297–304.

Thomson, J. D., and P. Wilson. 2008. Explaining Evolutionary Shifts between Bee and

Hummingbird Pollination: Convergence, Divergence, and Directionality. Int. J. Plant Sci.

169:23–38. The University of Chicago Press.

Tsuchimatsu, T., and S. Fujii. 2022. The selfing syndrome and beyond: diverse evolutionary

consequences of mating system transitions in plants. Philos. Trans. R. Soc. B Biol. Sci.

377:20200510.

Valderrama, E., J. B. Landis, D. Skinner, P. J. M. Maas, H. Maas-van de Kramer, T. André, N.

Grunder, C. Sass, M. Pinilla-Vargas, C. J. Guan, H. R. Phillips, A. M. R. de Almeida, and

1011   C. D. Specht. 2022. The genetic mechanisms underlying the convergent evolution of

1012   pollination syndromes in the Neotropical radiation of *Costus* L. Front. Plant Sci.

1013   13:874322.

1014  Vargas, O. M., B. Goldston, D. L. Grossenbacher, and K. M. Kay. 2020. Patterns of speciation

1015   are similar across mountainous and lowland regions for a Neotropical plant radiation

1016   (Costaceae: *Costus*). Evolution 74:2644–2661.

1017  Wessinger, C. A. 2024. How the switch to hummingbird pollination has greatly contributed to

1018   our understanding of evolutionary processes. New Phytol. 241:59–64.

1019  Wickham, H. 2009. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.

1020  Wickham, H. 2007. Reshaping Data with the reshape Package. J. Stat. Softw. 21:1–20.

1021  Wickham, H., R. François, L. Henry, and K. Müller. 2021. dplyr: A Grammar of Data

1022   Manipulation. R package version 1.0.7.

1023  Wickham, H., J. Hester, and J. Bryan. 2023. readr: Read Rectangular Text Data.

1024  Wikelski, M., J. Moxley, A. Eaton-Mordas, M. M. López-Uribe, R. Holland, D. Moskowitz, D.

1025   W. Roubik, and R. Kays. 2010. Large-Range Movements of Neotropical Orchid Bees

1026   Observed via Radio Telemetry. PLOS ONE 5:e10738.

1027  Wilke, C. O. 2020. cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2."

1028  Workman, R., R. Fedak, D. Kilburn, S. Hao, K. Liu, and W. Timp. 2019. High Molecular

1029   Weight DNA Extraction from Recalcitrant Plant Species for Third Generation

1030   Sequencing.

1031  Wozniak, N., K. Sartori, C. Kappel, L. Zhao, A. Erban, I. Fehrle, F. Jantzen, M. Orsucci, S.

1032   Rosa, M. Lenhard, J. Kopka, and A. Sicard. 2022. Convergence and molecular evolution

1033   of floral fragrance after independent transitions to self–fertilization. bioRxiv.

Xie, Y. 2023. knitr: A General-Purpose Package for Dynamic Report Generation in R.

Xu, S. 2022. ggstar: Multiple Geometric Shape Point Layer for "ggplot2."

Xu, S., Z. Dai, P. Guo, X. Fu, S. Liu, L. Zhou, W. Tang, T. Feng, M. Chen, L. Zhan, T. Wu, E.

Hu, Y. Jiang, X. Bo, and G. Yu. 2021. ggtreeExtra: Compact Visualization of Richly

Annotated Phylogenetic Data. Mol. Biol. Evol. 38:4039–4042.

Yost, J. M., and K. M. Kay. 2009. The evolution of postpollination reproductive isolation in

Costus. Sex. Plant Reprod. 22:247–255.

Yu, G. 2022. Data Integration, Manipulation and Visualization of Phylogenetic Trees. Chapman

& Hall.

Yu, G., D. K. Smith, H. Zhu, Y. Guan, and T. T.-Y. Lam. 2017. ggtree: an r package for

visualization and annotation of phylogenetic trees with their covariates and other

associated data. Methods Ecol. Evol. 8:28–36.

Zdobnov, E. M., D. Kuznetsov, F. Tegenfeldt, M. Manni, M. Berkeley, and E. V. Kriventseva.

2021. OrthoDB in 2020: evolutionary and functional annotations of orthologs. Nucleic

Acids Res. 49:D389–D393.

Zu, P., F. P. Schiestl, D. Gervasi, X. Li, D. Runcie, and F. Guillaume. 2020. Floral signals evolve

in a predictable way under artificial and pollinator selection in *Brassica rapa*. BMC Evol.

Biol. 20:127.

Zytnicki, M. 2017. mmquant: how to count multi-mapping reads? BMC Bioinformatics 18:411.

1053