H₂ consumption by acetogenic bacteria follows first-order kinetics

Laura Muñoz-Duarte[1] Laura.munoz@bce.au.dk https://orcid.org/0000-0003-0705-9999

Susmit Chakraborty[1], such@bce.au.dk

Louise Vinther Grøn[1] louise.groen@bce.au.dk

Maria Florencia Bambace[1] mfbambace@bce.au.dk

Jacopo Catalano[2] jcatalano@au.dk   https://orcid.org/0000-0003-4819-5436

Jo Philips[1]* jo.philips@bce.au.dk https://orcid.org/0000-0003-0471-7802

Department of Biological and Chemical Engineering

Aarhus University

[1]Gustav Wieds vej 10C,

[2] Åbogade 40A,

Aarhus, 8200 Denmark

*Corresponding author (jo.philips@bce.au.dk), Phone: +4593521569

Abstract

Acetogenic bacteria play an important role in various environmental and biotechnological

processes, because of their autotrophic metabolism converting carbon dioxide with

molecular hydrogen (H_2) as electron donor into acetate. The main factor limiting acetogenesis

is often H_2. Assessing the potential of acetogens in environmental and biotechnological

processes thus requires insights into their H_2 consumption kinetics. In this study, initial H_2

consumption rates at a range of different initial H_2 concentrations were measured for three

different acetogens. Interesting, for all three strains, H_2 consumption was found to follow

first-order kinetics, i.e. the H_2 consumption rate increased linearly with the dissolved H_2

concentration up to almost saturated H_2 levels. This contradicts Monod kinetics, which is

commonly assumed for acetogens. The obtained first-order rate coefficients (k_1) were further

validated by fitting first-order kinetics on previous time-course experimental results. The

latter method was also used to determine the k_1 value of five additional acetogens strains.

Biomass specific first-order rate coefficients were found to vary up to six-fold, with the highest

k_1 for *Acetobacterium wieringae* and the lowest for *Sporomusa sphaeroides*. Overall, our

results demonstrate the importance of the dissolved H_2 concentration to understand the rate

of acetogenesis in environmental and biotechnological settings.

Keywords

Acetogenic bacteria, H_2 consumption characteristics, first-order kinetics, CO_2 reduction, strain

selection, gas fermentation, microbial electrosynthesis.

Synopsis

Acetogenic bacteria are diverse, although they catalyze the same overall reaction. This study

provides insights into their different H_2 consumption rates, which has implications for their

role in the environment and in biotechnological applications.

Introduction

Acetogens are an interesting group of strictly anaerobic bacteria with remarkable metabolic capabilities. They are phylogenetically diverse, but have the common characteristic of utilizing the Wood–Ljungdahl pathway (WLP) for CO_2 fixation and acetate production while conserving energy, using H_2 as electron donor [1], according to the reaction:

$$4\ H_2 + 2\ CO_2 \rightarrow CH_3COOH + 2\ H_2O \quad \Delta G^{0'} = -95\ kJ/mol \quad (1)$$

Acetogens are widely distributed across diverse environments, ranging from soils to the human gut [2,3]. In addition, the metabolic abilities of acetogens are attractive for the development of novel biotechnological processes to convert carbon dioxide into valuable products. In gas fermentation, CO_2 is metabolized in a bioreactor fed with H_2 gas, while an electrode is used to produce H_2 *in situ* in the case of microbial electrosynthesis [4]. In these biotechnological settings, as well as in natural environments, the supply of H_2 often limits acetogenesis [5]. Understanding the relationship between the H_2 level and the kinetics of H_2 consumption and acetogenesis is thus essential for assessing the potential of acetogens in biotechnological and environmental processes.

Most often, H_2 consumption and growth by acetogens is assumed to follow Monod kinetics (similar to Michaelis Menten kinetics for enzymes)[5], in which the rate (μ) increases with the substrate concentration *[S]*, until a maximum rate (μ_{max}) is reached [6]:

$$\mu = \mu_{\max} \frac{[S]}{K_s + [S]}$$

$$(2)$$

The half-saturation coefficient K_s gives the substrate concentration at which half of the maximum rate is reached. So far, (maximum) growth rates or doubling times have frequently

81   been reported for acetogens[7,8], but only few studies have reported half-saturation

82   coefficients for H_2 consumption by acetogens (**Table 1**). Moreover, not all of these studies

83   carefully characterized whether the relationship between the H_2 consumption rate and the

84   H_2 concentration is well described by Monod kinetics (Eq. 2).

86   **Table 1.** Reported values of half-saturation coefficients (K_s) for H_2 consumption by acetogens.

Acetogen	K_s (µM)	Method	Reference
Acetobacterium woodii	0.73	Chemostat; based on half of the max growth rate	[9]
Acetobacterium woodii	2500	Enzyme assay on purified hydrogenases from cells grown on fructose	[15]
Acetobacterium bakii	4	Batch; H_2 depletion curves over time, constant biomass	[10]
Sporomusa termitida	6	Batch without headspace; H_2 depletion in the liquid	[11]
Eubacterium limosum	340	Batch; H_2 depletion curves over time- initial rates (similar as our procedure)	[12]
Clostridium strain P11	233	Hydrogenase assay on permeabilized whole cells from on syngas	[13]
Clostridium ljungdahlii	327	Hydrogenase assay on permeabilized whole cells grown on syngas	[13]
Sporomusa sphaeroides	341	Enzyme assay on purified hydrogenase from betaine grown cells	[14]

Diverse acetogens are available as pure culture. Several reviews enlisted available acetogenic

strains, their ability to consume H_2 and CO_2, and their main products [7,16,13]. Although

acetogens perform the same overall reaction (Eq. 1), their metabolic diversity is reflected by

differences in their energy conservation strategies [17–22], leading to different H_2 thresholds, i.e.

the H_2 partial pressure at which acetogenesis halts [23].

H_2 consumption kinetics likely also differ significantly among acetogenic strains, as different

doubling times or growth rates have been reported [5,8]. However, the few studies that have

published half-saturation coefficients for H_2 consumption by acetogens (**Table 1**), tested each

just one acetogenic strain. Moreover, these studies used various approaches to quantify the

half-saturation coefficient (**Table 1**), resulting in parameter values that are hard to compare

among the different studies.

With this work, we aim to investigate how the H_2 consumption rate of acetogens depends on

the H_2 concentration, as well as to compare the H_2 consumption kinetics of different

acetogens. We first measured initial H_2 consumption rates at different initial H_2 levels (ranging

from ~0.5 to 600 µM) for three different strains using the same pH, temperature, medium

and agitation. These analyses showed that H_2 consumption follows first-order kinetics, rather

than Monod kinetics, as no saturation was observed over the whole range of tested H_2 levels.

The obtained first-order rate coefficients were further validated by fitting first-order kinetics

on previous time-course experimental results [23]. Moreover, first-order kinetic coefficients of

five additional acetogens strains were also obtained by fitting time-course experimental

results.

Materials and methods

Strains and culture conditions

The acetogenic strains *Sporomusa ovata* (DSM 2662), *Acetobacterium woodii* (DSM 1030) and *Clostridium ljungdahlii* (DSM 13528) were revived from frozen stocks in the recommended medium (DSMZ 311, 135 and 879, respectively) using an organic substrate (betaine or fructose). After one transfer in the same medium, when an optical density (OD_{600nm}) of approximately 0.6 was reached, the cultures were filtered through a 0.2 µm polyethersulfone membrane. The cells were washed three times with autotrophic growth medium inside an anaerobic chamber with an N_2/CO_2 atmosphere (Jacomex, France) to remove acetate and organic substrates resulting from the heterotrophic growth medium. The washed cells were resuspended in autotrophic growth medium, with a composition of 1 g NH_4Cl, 0.1 g KCl, 0.8 g NaCl, 0.1 g KH_2PO_4, 0.077 g $MgCl_2$, 0.02 g $CaCl_2.2H_2O$, 0.3 g L-Cysteine- $HCl.H_2O$, 20 mM $NaHCO_3$, 50 mM 3-(N-morpholino) propane sulfonic acid (MOPS) brought to pH 7.0 with NaOH, 0.1 g yeast extract, 10 mL vitamin solution DSMZ 141, 1 mL trace element solution and 0.1 mL tungstate-selenite solution per liter [24]. The strains were not grown on H_2 before the start of the experiment, to avoid possible adaptations to high H_2 levels [25].

Initial consumption rate experiment

The H_2 consumption of the three acetogenic strains was characterized at a range of different initial H_2 concentrations, in the same conditions of temperature, medium composition, electron acceptor concentration, pH and agitation. Serum bottles (120 mL) were filled with 40 mL of autotrophic growth medium, while the headspace was flushed with gas mixes with different H_2 concentrations (ranging from ~0.1 to 60%) and with 20% CO_2 in N_2. These gas mixes were prepared using Brooks SLA5800 mass flow controllers. The serum bottles were

pressurized to 1.2 bar absolute pressure. At least five different H_2 initial concentrations were

tested in triplicate for each strain. The gas phase was allowed to equilibrate with the liquid

phase by agitation on an orbital shaker (Gerhardt, DE) at 100 rpm overnight at 30°C. After

equilibration, the bottles were inoculated with 5 mL of washed-cell suspension to reach a final

starting OD_{600} of 0.2. The bottles were incubated at 30°C and shaken at 100 rpm, while kept

upside down to avoid gas loss through the rubber septa.

After inoculation, the cells were given at least 1 hour of equilibration time. After that period

(set as time zero), samples (1.5 mL) were taken from the headspace for GC analysis until the

H_2 concentration had decreased by approx. 10% of the initial H_2 concentration. This allowed

us to measure the reaction rate at almost constant H_2 concentrations. The total pressure was

measured using a digital manometer (Omega DPG108, UK) before each headspace sampling.

The partial pressure of H_2 was calculated from the headspace H_2 concentration (%) measured

by GC and the total pressure at each sampling point. The concentration of dissolved H_2 in the

liquid phase was calculated using the Henry constant for H_2 ($7.8 \cdot 10^{-4}$ M bar^{-1}, at 30 °C) [26].

Abiotic controls with different initial H_2 concentrations were sampled using an identical

protocol to confirm that the H_2 depletion rates resulted only from the metabolic activities of

the acetogens. Liquid samples were taken for measurement of the OD_{600} and acetate

concentration in the initial and final sampling time.

Analytical methods

The analytical methods to measure the H_2 headspace concentration by GC, acetate

concentration by GC-VFA and the OD_{600} by spectrophotometry were previously described [23].

In addition, correlation curves of the cell density and dry weight with the optical density

(OD_{600}) were obtained for each acetogenic strain (**Table S1**). The cell density (cells mL^{-1}) was

quantified by Impedance Flow Cytometry (IFC) (Bactobox, SBT, Denmark). A heterotrophically

grown culture was washed and resuspended in autotrophic growth medium as described

above, and thoroughly homogenized using a vortex. Afterwards, the culture was diluted in

1:9 diluted Phosphate Buffer Saline (PBS) and measured in triplicates with IFC. Total cell

numbers were used to calculate the correlation factors. The IFC method was validated using

qPCR on the bacterial 16S rRNA gene, which gave results in the same order of magnitude.

Likewise, the dry weight was correlated to OD_{600} for each acetogenic strain in quintuplicates.

Cell suspensions were dried in an oven at 80°C until constant weight was reached and the

average dry weight was used to calculate the correlation factors.

Determination of kinetic order and kinetic coefficients from the initial rate experiment

The rate of H_2 consumption was correlated to the initial H_2 concentration to determine the

kinetic order. The rate of H_2 consumption was derived from the decrease of the H_2 partial

pressure over time. For each sampling event, the headspace pressure was measured before

and after sampling, to estimate the small loss of H_2 related to the withdrawal of the gaseous

sample. This loss of H_2 due to sampling was subtracted from the overall decrease of the H_2

total mole [27] to avoid overestimating the microbial consumption. The volumes of gas and

liquid phases were considered constant throughout the experiment. Our calculations assume

that H_2 was the only limiting substrate and that the agitation resulted in instantaneous

equilibrium between the gas and liquid phases in the bottles. The latter assumption is further

evaluated and discussed below.

H_2 consumption followed first-order kinetics over the range of H_2 concentrations evaluated

in this study, as described below. Since the H_2 consumption by microbial metabolism only

occurs in the liquid phase, but dissolved H_2 is replenished by the dissolution of gaseous H_2,

the first-order kinetic coefficients were calculated based on the total mole change over time,

similar as described previously [28]. The total moles ($n_{T,H2}$) of H_2 at any time point is described

by Eq. 3:

186
$$n_{T,H_2} = n_{l,H_2} + n_{g,H_2} = C_{H_2} \cdot V_l + \frac{p_{H_2} \cdot V_g}{RT} = C_{H_2} \cdot \left(V_l + \frac{V_g}{HRT}\right) \tag{3}$$

With n_l moles in the liquid phase, n_g moles in the gas phase, C_{H_2} the H_2 concentration in the

liquid phase, V_l the volume of the liquid phase, V_g the volume in the gas phase, p_{H_2} the partial

pressure of H_2 in the headspace, R the ideal gas constant, T the temperature in Kelvin, H

Henry's constant of H_2 at 30 °C.

The ordinary differential equation describing the time dependence of the total H_2 moles in

the system and considering first-order kinetics reads as:

194
$$\frac{dn_T}{dt} = \frac{dC_{H_2}}{dt} \cdot \left(V_l + \frac{V_g}{HRT}\right) = -k_1 \left[C_{H_2} - C_{H_2}^*\right] \cdot V_l \tag{4}$$

Where k_1 is the first-order rate coefficient and has units of h^{-1} and $C_{H_2}^*$ is the H_2 threshold as

previously experimentally determined [23]. The H_2 threshold is defined as the minimum H_2

dissolved concentration at which acetogenesis is thermodynamically feasible, and below

which acetogenesis becomes unfavorable and halts. The expression of the H_2 threshold in Eq.

4 is as previously proposed [10,29].

The first-order rate coefficients and their confidence intervals were obtained from the slope

of the total mol of H_2 change per time per volume of liquid culture vs the initial H_2

concentration (rearranging Eq. 4), using the linear regression function of GraphPad 9, which

also calculated the 95% confidence interval. The cell-specific first-order rate coefficients were

obtained by dividing k_1 by the cell density or dry weight, which was constant over the

experimental time.

Determination of first-order kinetic coefficients from time course experiment

To validate the first-order rate coefficients obtained from the initial rate experiment, the

kinetic coefficients were also calculated by numerical fitting of our previous experimental H_2

consumption data [23]. This study included the three strains used in the initial rate experiment,

as well as five additional strains: *S. ovata* (DSM 2663), *S. sphaeroides* (DSM 2875), *A. malicum*

(DSM 4132, *A. wieringae* (DSM 1911), and *C. autoethanogenum* (DSM 10061), for which also

first-order rate coefficients were fitted.

This earlier study [23] tested all strains at OD_{600} of 0.2 (similarly as in our initial rate experiment)

and used about 25 µM as starting H_2 concentration, while H_2 consumption was monitored

until the specific threshold was reached. In addition to these earlier experimental data, the

same experiment was repeated for the two *Clostridium* strains at pH 5.9 (instead of pH 7), as

this is closer to their optimal pH [30].

The first-order H_2 consumption coefficient of those eight acetogenic strains was determined

by fitting the above ordinary differential equation Eq. (4), on the experimental data, using the

starting H_2 concentration as initial condition. Eq. (4) was numerically integrated by using the

ode15s algorithm in Matlab. In addition, a non-linear Nelder-Mead simplex algorithm[31]

(*fminsearch*) was used to obtain the fitting curves by minimizing the sum of squares errors

(SSE) between the model prediction and the experimental data. Equal weight was applied to

all the data points for calculating SSE. The input parameters, V_l, V_g, H, R, T, and $C^*_{H_2}$ were taken

as experimentally determined [23]. The 95% confidence interval (CI) for each predicted k_1 value

was further calculated as described in the **Supplementary Method 1**.

Evaluation of mass transfer limitations

The calculations described above assume instant equilibrium between the gas and liquid

phase. To evaluate the possible influence of mass transfer limitations on our results, we

calculated the overall mass transfer coefficient $k_L a$ from our orbital shaker settings, based on

previously reported mass transfer correlations for shakers [32] (**Supplemental Method 2**). This

parameter was used to evaluate whether H_2 mass transfer limitations could have impacted

the type of the kinetic order, observed by the initial rate experiment. Hereto, the outcome of

the initial rate experiment was simulated using a system of ODE's, including a term for the

mass transport of H_2 from the headspace to the liquid phase (**Supplemental Method 3**). The

outcome of an initial rate experiment was simulated assuming first-order or Monod kinetics.

Furthermore, the possible effect of mass transfer limitations on the fitted k_1 values was

assessed, by repeating the fitting of the first-order kinetic coefficients (as described above),

using the system of ODE's including mass transfer.

Results

Initial consumption rates experiment (IR)

H_2 consumption rates for at least five different initial H_2 concentrations were determined in

comparable conditions for *S. ovata*, *A. woodii* and *C. ljungdahlii*. Biological consumption was

confirmed by the faster depletion of H_2 than in abiotic controls under the same conditions

(**Figure S1**). Headspace sampling ended when the headspace H_2 concentration had decreased

by about 10%. In all tested conditions, the OD_{600} did not change during the short experimental

timeframe, as the amount of consumed H_2 was insufficient to lead to observable growth. In

addition, changes in the acetate concentration were insufficient to be detected.

Initially, H_2 consumption rates were measured up to a dissolved H_2 concentration of 180 µM

(approx. 20% in headspace), since the H_2 consumption rate was expected to reach a maximum

already in the lower µM range, based on previously reported Monod half-saturation

coefficients for *A. woodii* and *S. termitida* [9,11]. Plotting the experimentally determined H_2

consumption rates versus the dissolved H_2 concentration, however, showed a linear

relationship and no indication of a deflection or a maximum rate at higher H_2 concentrations

(**Figure 1**). For this reason, the range of initial concentrations was extended to 450 µM for *A.*

woodii and *C. ljungdahlii*, and even to 600 µM for *S. ovata*. Note that the latter concentration

is close to the dissolved H_2 concentration 780 µM at saturation (1 atm of pure H_2 in headspace

at 30 °C). Even with this extended range of initial H_2 concentration, the relationship between

the initial H_2 concentration and the H_2 consumption rate remained linear, for all three tested

strains, without any observable deflection towards a maximum (**Figure 1**). This linear

relationship indicates that H_2 consumption by acetogens follows first-order reaction kinetics,

at least at H_2 concentrations up to saturation (not considering overpressure). The absence of

a maximum rate at the highest H_2 concentrations suggests that H_2 consumption remains

limited by the H_2 concentration and not by the uptake or conversion capacity of the cells.

It can also be observed from **Figure 1** that *S. ovata* consumed H_2 faster at the same initial H_2

concentration than *A. woodii* and *C. ljungdahlii.* The slopes of the plots of **Figure 1** indeed

relate to the first-order rate coefficient k_1 (Eq. 4). The first-order rate coefficient is more than

six times higher for *S. ovata* than for *C. ljungdahlii* and almost double that for *A. woodii* (**Table**

2). *C. ljungdahlii* has the lowest *k_1* among the three strains evaluated in the initial rates

experiment, which is most likely due to the experimental pH 7, which deviated from its

optimum pH 6 [30]. Also, when expressed relative to the cell density or dry weight, the cell-

specific first-order coefficient was highest for *S. ovata* (**Table 2**).

Evaluation of the impact of mass transfer resistance on the kinetic order type

We evaluated whether H_2 mass transfer limitations could impact the type of kinetics revealed

by **Figure 1.** We first estimated the overall mass transfer coefficient k_La from the

characteristics from our specific setup and orbital shaker settings, which was 22.7 per hour

(**Supplementary Methods S2**). This value was used to simulate the outcome of an initial rate

experiment, under the assumption of first-order or Monod kinetics. Our simulations showed

that our initial rate experiment revealed the true kinetic order type, as the shape of the curve

was not impacted by the actual mass transfer limitations (**Figure S2**). Only at a 100 times lower

k_La (which would correspond to static incubation), mass transfer limitations would be so

strong that they would lead to a linear relationship between the consumption rate and the H_2

concentration, also in case of Monod kinetics.

Figure 1. The observed H_2 consumption rates in function of the initial H_2 concentration for *S.*

ovata (top), *A. woodii* (middle) and *C. ljungdahlii* (bottom). The observed H_2 consumption

rates (*y*-axis) were expressed as total mmol H_2 per hour per L of liquid phase (Eq. 4) to account

for the replenishment of H_2 from the headspace. The H_2 concentrations (*x*-axis) are corrected

by the H_2 threshold [23], thus the origin of the axis corresponds with the respective H_2 threshold

of each strain. The continuous line reflects the linear relationship as a first-order reaction. The

dotted line around the linear regression represents the 95% confidence interval. The liquid

medium is saturated with H_2 at 780 μM in our conditions. Each replicate is shown as a

separate dot in the plot.

Fitting time course experiment (TC)

To validate the obtained first-order rate coefficients for *S. ovata* 2662, *A. woodii* and *C.*

ljungdahlii, the kinetic coefficients were also calculated by fitting Eq. 4 on previous

experimental H_2 consumption time course data [23]. These independent experiments were

performed at the same pH, temperature, medium and agitation, but used only one initial H_2

concentration (~25 μM) and followed the consumption of H_2 until the H_2 threshold was

reached. The fitted curves described the H_2 depletion reasonably well (**Figure 2**). The k_1 values

obtained using the two experimental approaches are highly comparable (**Figure 3, Table 2**),

which validates the method used for the initial rate experiment.

In addition, the same fitting method was used to obtain first-order rate coefficients from

previous experimental results for five additional acetogenic strains, as well as from new

experimental H_2 depletion data for the *Clostridium* strains at pH 5.9 (**Figure 2**).

Among the tested strains, *A. wieringae* exhibited the highest k_1 and dry weight specific-k_1

(**Table 2, Figure 4, Figure S4**). Interestingly, both *S. ovata* 2662 and 2663 had significantly

higher k_1 values (based on both DW and cell density), compared to *A. woodii*, indicating its

higher overall H_2 consumption capacity under the tested conditions (**Table 2, Figure 4, Figure**

S4).

C. ljungdahlii displayed a pH-dependent k_1, with a higher value observed at its more optimal

pH 5.9 compared to pH 7 (**Table 2, Figure 4, Figure S4**), which aligns with the expected

influence of optimal growth conditions on metabolic activity. Notably, *C. ljungdahlii* also

demonstrated the highest cell density specific-k_1 at pH 5.9 (**Table 2, Figure S4**) of all the strains

tested. On the contrary, *C. autoethanogenum* did not exhibit a higher k_1 at pH 5.9 than at pH

7.0, even though is optimal pH is described to be 5.9 [33]. This discrepancy could relate to the

narrow concentration difference between the initial H_2 concentration and the H_2 threshold

for *C. autoethanogenum* at pH 7 (**Figure 2**), which resulted in a wide confidence interval for

the fitted k_1 (**Table 2**). Nevertheless, the H_2 threshold for *C. autoethanogenum* did reflect its

more optimal pH at 5.9 (**Figure 2**). Also *C. ljungdahlii* had a lower H_2 threshold at pH 5.9 than

at pH 7.0 (**Figure 2**), while in all conditions the two *Clostridium* strains still had a higher H_2

threshold, than all other tested strains.

Finally, *S. sphaeroides* displayed the lowest k_1 and biomass-specific k_1 values (DW and cell

density) (**Table 2**, **Figure S4**), but when analyzing the k_1 values expressed by dry weight, no

significant differences between *S. sphaeroides* and *A. malicum*, *A. woodii*, *C.*

autoethanogenum at pH 5.9, *Clostridium ljungdahlii* were observed (**Table 2**, **Figure 4**).

Evaluation of the impact of mass transfer on the fitted k_1 values

Our simulations (**Figure S2**) showed that the assumption of instant equilibrium between the

gas and liquid phase could have underestimated the actual k_1 values, which we evaluated

further. The overall mass transfer coefficient k_La (22.7 h^{-1}) is about four times higher than the

highest k_1 obtained in this study. We evaluated its influence on the fitted value of the first-

order rate coefficient obtained for *A. wieringae*, which is the strain with the highest

consumption rate and thus potentially most susceptible to mass transfer limitations. Our

analysis revealed that the kinetic coefficient obtained incorporating mass transfer resistance

is at the most 32% higher than when assuming gas-liquid equilibrium (**Figure S3**, **Table S2**).

For all the other strains, this difference is even lower, due to their lower k_1 (**Table S2**). As this

difference is much less than the difference observed between the strains (6-fold difference in

k_1 between highest and lowest), as well as the differences that result from other conditions

(for instance pH), we conclude that the results of **Table 2** (assuming instant equilibrium) are

sufficiently accurate and can be considered as lower limits for the H_2 consumption rates.

Figure 2. Best-fitting curves on the change of the dissolved H_2 concentration over time for

eight acetogenic strains[23] (pH 7), as well as on similar experimental data for *C.*

autoethanogenum and *C. ljungdahlii* at pH 5.9. Data points (symbols) represent the mean H_2

concentration (3 to 5 replicates) with the standard deviation given by the error bars. The best

fit is given by the full black lines, while the confidence interval (95%) is aligned by pink dotted

lines. The H_2 threshold is marked by a horizontal orange dotted line.

Table 2. First-order rate coefficients (k_1) of H_2 consumption obtained by the initial rates

experiment (IR) (grey) or by fitting the time course experiment (FT) for different acetogenic

bacteria. First-order rate coefficients are also given normalized by the biomass concentration,

expressed as cell density or as dry weight per L. All parameters are at pH 7 unless stated

otherwise. The 95% confidence intervals are presented in brackets next to the values. These

parameters were obtained under the assumption of instant equilibrium between the gas and

liquid phase.

Organism	Experiment	k_1 (h^{-1})	DW Specific k_1 $(h^{-1} (g\ DW\ L^{-1})^{-1})$	Cell density specific k_1 $x10^{-10} (h^{-1} (cell\ L^{-1})^{-1})$
S. sphaeroides	TC	0.88 (0.79 to 0.97)	10.63	0.06
C. ljungdahlii	TC	0.94 (0.79 to 1.08)	13.28	0.64
C. ljungdahlii	IR	0.58 (0.53 to 0.62)	8.19	0.25
C. autoethanogenum pH 5.9	TC	1.13 (0.87 to 1.37)	12.28	0.14
A. woodii	TC	1.74 (1.51 to 1.97)	15.47	0.39
A. woodii	IR	2.18 (2.03 to 2.33)	19.36	0.44
C. ljungdahlii pH 5.9	TC	2.76 (2.65 to 2.87)	39.02	1.11
S. ovata 2662	TC	3.27 (2.82 to 3.72)	29.81	1.09
S. ovata 2662	IR	3.84 (3.62 to 4.06)	34.97	1.21
C. autoethanogenum	TC	3.49 (3.03 to 3.95)	38.19	0.44
S. ovata 2663	TC	3.83 (3.60 to 4.06)	53.98	0.88
A. malicum	TC	2.38 (2.07 to 2.68)	13.39	0.15
A. wieringae	TC	5.79 (5.61 to 5.96)	60.93	0.67

Figure 3. Comparative graph of the numerical values of k_1 obtained from the two

experimental approaches: Initial rates experiment (IR) and fitting on the time course

experimental data (TC) for *S. ovata* 2662, *A. woodii* and *C. ljungdahlii.* The error bar

corresponds to the 95% confidence interval.

Figure 4. Comparison of the dry weight specific first-order coefficient k_1 obtained by fitting

the time course experiment for the acetogenic strains tested. The symbol represents the

mean value, and the bars represent the 95% confidence interval.

Discussion

H$_2$ consumption by acetogens follows first-order kinetics and not Monod kinetics up to saturated H$_2$ levels

In this work, we found that the H$_2$ consumption rates of three different acetogenic strains linearly increased with the dissolved H$_2$ concentration, up to almost saturated H$_2$ concentrations (maximal dissolved H$_2$ concentration was 450 to 600 μM) (**Figure 1**). This linear trend shows that H$_2$ consumption by the different acetogenic strains followed first-order kinetics. For practical reasons, we did not further increase the H$_2$ partial pressure, even though acetogens possibly exhibit a maximum H$_2$ consumption rate (Monod kinetics) at oversaturated H$_2$ concentrations. Overpressures of H$_2$ (>1 atm partial pressure) are relevant for gas fermentation applications, but in most environmental conditions, as well as during microbial electrosynthesis, only undersaturated H$_2$ levels (<1 atm total pressure) are expected.

Our finding of first-order H$_2$ consumption kinetics contradicts with previous studies, which assumed Monod kinetics (Eq. 2) and reported half-saturation coefficients for H$_2$ consumption spanning four orders of magnitude (**Table 1**). Previously, Egli (2009)[6] pointed out that determining K_s is overall difficult, even for soluble substrates, e.g. also for glucose uptake by the model organism *E. coli*, reported K_s values range over three orders of magnitude. Egli (2009)[6] related this to differences in the experimental approach. In addition, the use of growth rates is problematic, as changes in biomass are hard to measure at low substrate concentrations [6]. Also Robinson and Tiedje (1982)[34] pointed out that underestimated H$_2$ half-saturation coefficients can result from relating H$_2$ consumption to growth. Consequently, the direct determination of the consumption rate in function of the substrate concentration, as in our study, reduces the associated error.

The previous studies reporting half-saturation coefficients for acetogenic H_2 consumption

(**Table 1**) did not report the reliability of their K_s value. Smith *et al*. (2020) [29] clearly illustrated

the importance of reporting the confidence interval of parameter estimates. They found that

both Monod kinetics and first-order kinetics fitted batch growth data for the acetogen *Blautia*

hydrogenotrophica well, but described that in the case of Monod kinetics, the values of K_s and

μ_{max} were strongly correlated, which was reflected by an extremely wide confidence interval

for K_s. In contrast, they reported a narrow confidence interval for their first-order rate

coefficient. In our work, the confidence intervals associated with the first-order kinetic

coefficient are also narrow (**Table 2**). In addition, we applied two different experimental

approaches (initial rate experiment and fitting time course data), which resulted in highly

comparable k_1 values (**Figure 3**). Our k_1 values can thus be considered as highly reliable.

Possibly, H_2 mass transfer limitations affect the observed H_2 consumption kinetic order type

[35,36]. Through simulations (**Figure S2**), we showed that the actual mass transfer resistance in

our system could not have masked the true kinetic order type. The mass transfer limitations

did affect the estimated k_1 values, but this was maximally 32% for the strain with the highest

k_1 (**Figure S3**, **Table S2**).

Besides Smith *et al*. (2020) [29], few other studies previously found indications for first-order H_2

consumption by acetogens. For instance, a study of *A. woodii* in chemostats to determine its

maximal H_2 uptake rate concluded that a physiological limitation was likely not reached in

their experiments [37]. Also the relatively high K_s found using several hydrogenase assays (**Table**

1), show that the maximum H_2 conversion rate was not reached below saturation.

First-order H_2 consumption rate coefficients differ up to six times among acetogenic strains

Our results demonstrate that acetogenic strains strongly differ in their first-order H_2

consumption rate coefficient (**Table 2**, **Figure 4**). The highest k_1 coefficient was obtained for

A. wieringae, which had a six-fold higher consumption rate than *S. sphaeroides,* for which the

lowest k_1 value was found. The same strains have respectively the highest and lowest k_1 value

when k_1 is expressed per dry weight. *C. ljungdahlii* at pH 5.9 had one of the highest k_1

expressed per cell density, but an average k_1 expressed per dry weight. This discrepancy likely

stems from cell size differences, as *C. ljungdahlii* possibly has larger cells, corresponding to

fewer cells per unit of dry weight.

Contrary to what was previously observed for H_2 thresholds [23], k_1 values did not correlate with

the genus, indicating that k_1 values might be strain-specific. Likewise, Groher and Weuster-

Botz [8] described variations in the maximal H_2 uptake rate and maximal growth rate among

three *Acetobacterium* and two *Sporomusa* species. Differences in growth rates at comparable

conditions within the same genus were also observed by Bengelsdorf *et al*. 2016 [38] for four

different *Clostridium* strains.

In part, these variations in k_1 are likely due to differences in optimal conditions, as illustrated

by the different k_1 values at different pH for the *Clostridium* strains (**Table 2**, **Figure 4**). Besides

pH, the H_2 consumption rate of a single acetogenic strain varies with the temperature [10] and

the medium composition. For example, differences in the concentrations of yeast extract,

reductive agents or trace elements (selenium and tungsten) are known to affect acetogenesis

rates [39–41].

In addition to varying optimal conditions, differences in H_2 consumption rates between

acetogenic strains could be due to different kinetics of the involved hydrogenases, which are

the metalloenzymes catalyzing the redox conversion between H_2 and protons. The

hydrogenases type [NiFe] and the [FeFe] hydrogenases are commonly found in acetogens [42–

[44]. These enzymes exhibit complex structural features, including hydrophobic channels for gas

diffusion to the active site [45]. This structural conformation can influence the affinity and the

sensitivity of the enzyme [46]. Indeed, some studies already demonstrated differences in the

hydrogenase kinetics of different acetogens [13,15,47].

In our study, *A. wieringae* exhibited the highest H_2 consumption rate under our experimental

conditions, followed by *S. ovata* and the two *Clostridium* strains (**Table 2**, **Figure 4**). Numerous

studies already explored the capabilities of *C. ljungdahlii* and *C. autoethanogenum* as catalysts

in gas conversion technologies [48–52], because their high conversion rates. Maybe less known,

but also *A. wieringae* has already been recognized for its high conversion rates on C_1 carbon

sources [53,54]. In addition, *A. wieringae* was the keystone member in an enriched microbial

community leading to high rates of microbial electrosynthesis [55,56], making it a highly

promising strain for applications.

Implications

Our finding that H_2 consumption by acetogens follows first-order kinetics up to almost

saturated H_2 levels, means that the dissolved H_2 concentration is a major factor limiting

acetogenesis in both environmental and biotechnological systems. First-order kinetics entails

that the rate of H_2 conversion per unit biomass is higher at an increased dissolved H_2

concentration (Eq. 4). This insight is highly relevant to optimize acetogenic biotechnologies.

In gas fermentation, increasing the H_2 partial pressure is often used to increase the H_2

solubility and thus to overcome mass transfer resistance[57]. Our results suggest that this also

leads to higher consumption rates by acetogens. In contrast, low dissolved H_2 levels often

prevail in microbial electrosynthesis systems, which likely contributes to their limited acetate

production rates [58].

In addition, the understanding that H_2 consumption by acetogens follows first-order kinetics

is important to understand their relevance in environmental settings. For instance,

competition between acetogenic bacteria and methanogenic archaea has mostly been

studied by comparing their Monod kinetic parameters for H_2 consumption [10,59], but the use

of the right kinetic order could lead to more accurate interpretations. Furthermore, to

understand acetogenesis in both natural and biotechnological settings, numerical simulations

are highly interesting, but the value of models entirely depends on the use of the correct

expressions and parameter values governing acetogenic kinetics. Consequently, the insights

generated by our study are of high relevance for any research related to acetogenesis.

This work also showed that first-order rate coefficients between acetogens differ up to six

times (**Table 2**, **Figure 4**). The k_1 values obtained by this work are of high value for optimal

strain selection for the different biotechnological applications. Nowadays, acetogenic strain

selection is mostly based on the desired end-product. However, genetic engineering can alter

the formed end-product [54,60,61], while H_2 consumption and product formation rates are much

harder to engineer. Consequently, it remains important to understand the intrinsic

differences in kinetics between the available acetogenic strains to allow optimal strain

selection.

Finally, it should be noted that this work only characterized H_2 consumption rates. This work

did not analyze growth nor acetate production rates. Moreover, the H_2 consumption rate

does not only depend on the (biomass specific) k_1, but definitely also on the biomass density

in the system, which in turn depends on growth. Acetogens likely differ in their growth yields,

as they have different ATP gains [23]. Furthermore, product versus biomass yields might vary

among acetogens. Insights into all these aspects are required, before a comprehensive

understanding of the product and growth kinetics of acetogens becomes possible.

Author contributions

LM and JP designed the experiments; LM performed the experimental work; SC fitted the k_1 values on our experimental results; MFB performed qPCR; JC performed the simulations investigating mass transfer limitations; LM and LVG analyzed the experimental data; LM and SC constructed the figures; LM and JP wrote the manuscript; SC, FMB, LVG and JC edited the manuscript; JC and JP acquired funding and performed supervision. All authors contributed to the article and approved the submitted version.

Acknowledgements

This work was financed by a starting grant of Aarhus Universitets Forskningsfond (AUFF), as well as by a Green Transition Project I grant (MES-MODEL, 1127-00048B) from the Independent Research Fund Denmark (DFF).

We thank Lars-Erik Meyer for his advice on how to determine kinetics. We also thank Thomas Thomas Dyekjær, Cecilie Nielsen, Gitte Hastrup and Heidi Skov Johansen for their technical and practical support of this project. Miguel Oliveira is acknowledged for his contribution to the Bactobox setup.

References

(1) Basen, M.; Müller, V. Editorial: Acetogens - from the Origin of Life to Biotechnological Applications. *Frontiers in Microbiology*. Frontiers Media S.A. 2023. https://doi.org/10.3389/fmicb.2023.1186930.

(2) Smith, N. W.; Shorten, P. R.; Altermann, E.; Roy, N. C.; McNabb, W. C. Competition for Hydrogen Prevents Coexistence of Human Gastrointestinal Hydrogenotrophs in Continuous Culture. *Front Microbiol* **2020**, *11*. https://doi.org/10.3389/fmicb.2020.01073.

(3) Drake, H. L.; Küsel, K.; Matthies, C. Ecological Consequences of the Phylogenetic and Physiological Diversities of Acetogens. *Antonie van Leeuwenhoek, International Journal of General and Molecular Microbiology* **2002**, *81* (1–4), 203–213. https://doi.org/10.1023/A:1020514617738.

(4) Harnisch, F.; Deutzmann, J. S.; Boto, S. T.; Rosenbaum, M. A. Microbial Electrosynthesis: Opportunities for Microbial Pure Cultures. *Trends Biotechnol* **2024**. https://doi.org/10.1016/j.tibtech.2024.02.004.

(5) Philips, J. Extracellular Electron Uptake by Acetogenic Bacteria: Does H_2 Consumption Favor the H_2 Evolution Reaction on a Cathode or Metallic Iron ? *Front Microbiol* **2020**, *10*, 1–13. https://doi.org/10.3389/fmicb.2019.02997.

(6) Egli, T. Growth Kinetics, Bacterial. In *Encyclopedia of Microbiology*; Elsevier Inc., 2009; pp 180–193. https://doi.org/10.1016/B978-012373944-5.00260-1.

(7) Bengelsdorf, F. R.; Beck, M. H.; Erz, C.; Hoffmeister, S.; Karl, M. M.; Riegler, P.; Wirth, S.; Poehlein, A.; Weuster-Botz, D.; Dürre, P. Bacterial Anaerobic Synthesis Gas (Syngas) and CO_2 + H_2 Fermentation. *Adv Appl Microbiol* **2018**, *103*, 143–221. https://doi.org/10.1016/BS.AAMBS.2018.01.002.

(8) Groher, A.; Weuster-Botz, D. Comparative Reaction Engineering Analysis of Different Acetogenic Bacteria for Gas Fermentation. *J Biotechnol* **2016**, *228*, 82–94. https://doi.org/10.1016/j.jbiotec.2016.04.032.

(9) Peters, V.; Janssen, P. H.; Conrad, R. Efficiency of Hydrogen Utilization during Unitrophic and Mixotrophic Growth of *Acetobacterium Woodii* on Hydrogen and Lactate in the Chemostat. *FEMS Microbiol Ecol* **1998**, *26*, 317–324. https://doi.org/10.1111/j.1574-6941.1998.tb00516.x.

(10) Kotsyurbenko, O. R.; Glagolev, M. V.; Nozhevnikova, A. N.; Conrad, R. Competition
between Homoacetogenic Bacteria and Methanogenic Archaea for Hydrogen at Low
Temperature. *FEMS Microbiol Ecol* **2001**, *38* (2–3), 153–159.
https://doi.org/10.1016/S0168-6496(01)00179-9.

(11) Breznak, J. A.; Switzer, J. M.; Seitz, H.-J. Sporomusa Termitida Sp. Nov., an H2/CO2-
Utilizing Acetogen Isolated from Termites. *Arch Microbiol* **1988**, *50*, 282–288.

(12) Genthner, B. R.; Bryant, M. P. *Additional Characteristics of One-Carbon-Compound
Utilization by Eubacterium Limosum and Acetobacterium Woodii*; 1987.

(13) Skidmore, B. E.; Baker, R. A.; Banjade, D. R.; Bray, J. M.; Tree, D. R.; Lewis, R. S. Syngas
Fermentation to Biofuels: Effects of Hydrogen Partial Pressure on Hydrogenase
Efficiency. **2013**. https://doi.org/10.1016/j.biombioe.2013.01.034.

(14) Dobrindt, U.; Ulrich, M. Purification and Characterization of a Membrane-Bound
hydrogenase from Sporomusa Sphaeroides involved in Energy-Transducing Electron
Transport. *Arch Microbiol* **1996**, *165*, 141–147.

(15) Ragsdale, S. W.; Ljungdahl, L. G. Hydrogenase from Acetobacterium Woodii*. *Arch
Microbiol* **1984**, *139*, 361–365.

(16) Schiel-Bengelsdorf, B.; Dürre, P. Pathway Engineering and Synthetic Biology Using
Acetogens. *FEBS Lett* **2012**, *586*, 2191–2198.
https://doi.org/10.1016/j.febslet.2012.04.043.

(17) Wiechmann, A.; Müller, V. Energy Conservation in the Acetogenic Bacterium
Clostridium Aceticum. *Microorganisms* **2021**, *9* (2), 1–12.
https://doi.org/10.3390/microorganisms9020258.

(18) Katsyv, A.; Müller, V. Overcoming Energetic Barriers in Acetogenic C1 Conversion.
Frontiers in Bioengineering and Biotechnology | www.frontiersin.org **2020**, *8*, 621166.
https://doi.org/10.3389/fbioe.2020.621166.

(19) Rosenbaum, F. P.; Müller, V. Energy Conservation under Extreme Energy Limitation:
The Role of Cytochromes and Quinones in Acetogenic Bacteria. *Extremophiles* **2021**,
25, 413–424. https://doi.org/10.1007/s00792-021-01241-0.

(20) Kremp, F.; Roth, J.; Müller, V. A Third Way of Energy Conservation in Acetogenic
Bacteria. *Microbiol Spectr* **2022**.

(21)   Kremp, F.; Roth, J.; Müller, V. The *Sporomusa* Type Nfn Is a Novel Type of Electron-
581   Bifurcating Transhydrogenase That Links the Redox Pools in Acetogenic Bacteria. *Sci*
582   *Rep* **2020**, *10* (1). https://doi.org/10.1038/S41598-020-71038-2.

583   (22)   Wiechmann, A.; Ciurus, S.; Oswald, F.; Seiler, V. N.; Volker Müller, •. It Does Not Always
584   Take Two to Tango: "Syntrophy" via Hydrogen Cycling in One Bacterial Cell. *ISME J*
585   **2020**, *14*, 1561–1570. https://doi.org/10.1038/s41396-020-0627-1.

586   (23)   Munoz, L.; Philips, J. No Acetogen Is Equal: Strongly Different H2 Thresholds Reflect
587   Diverse Bioenergetics in Acetogenic Bacteria. *Environ Microbiol* **2023**, *25* (10), 2032–
588   2040. https://doi.org/10.1111/1462-2920.16429.

589   (24)   Patil, S. A.; Arends, J. B. A.; Vanwonterghem, I.; Van Meerbergen, J.; Guo, K.; Tyson, G.
590   W.; Rabaey, K. Selective Enrichment Establishes a Stable Performing Community for
591   Microbial Electrosynthesis of Acetate from CO_2. *Environ Sci Technol* **2015**, *49* (14),
592   8833–8843. https://doi.org/10.1021/es506149d.

593   (25)   Kato, S.; Yumoto, I.; Kamagata, Y. Isolation of Acetogenic Bacteria That Induce
594   Biocorrosion by Utilizing Metallic Iron as the Sole Electron Donor. *Appl Environ*
595   *Microbiol* **2015**, *81* (1), 67–73. https://doi.org/10.1128/AEM.02767-14.

596   (26)   Sander, R. Compilation of Henry's Law Constants (Version 4.0) for Water as Solvent.
597   *Atmos. Chem. Phys* **2015**, *15*, 4399–4981. https://doi.org/10.5194/acp-15-4399-2015.

598   (27)   Dopffel, N.; Mayers, K.; Kedir, A.; Alagic, E.; Annie An-Stepec, B.; Djurhuus, K.; Boldt, D.;
599   Beeder, J.; Hoth, S. Microbial Hydrogen Consumption Leads to a Significant PH Increase
600   under High-Saline-Conditions: Implications for Hydrogen Storage in Salt Caverns.
601   *Scientific Reports* **2023**, *13*. https://doi.org/10.1038/s41598-023-37630-y.

602   (28)   Karadagli, F.; Rittmann, B. E. Thermodynamic and Kinetic Analysis of the H_2 Threshold
603   for *Methanobacterium Bryantii* M.o.H. *Biodegradation* **2007**, *18* (4), 439–452.
604   https://doi.org/10.1007/s10532-006-9073-7.

605   (29)   Smith, N. W.; Shorten, P. R.; Altermann, E.; Roy, N. C.; Mcnabb, W. C. Mathematical
606   Modelling Supports the Existence of a Threshold Hydrogen Concentration and Media-
607   Dependent Yields in the Growth of a Reductive Acetogen. *Bioprocess Biosyst Eng* **2020**,
608   *43*, 885–894. https://doi.org/10.1007/s00449-020-02285-w.

609   (30)   Tanner, R. S.; Miller, L. M.; Decheng, A. *Clostridium Ljungdahlii Sp. Nov., an Acetogenic*
610   *Species in Clostridial RRNA Homology Group I*; 1993; Vol. 43.

(31) Lagarias, J. C.; Reeds, J. A.; Wright, M. H.; Wright, P. E.; Optim, S. J. *CONVERGENCE PROPERTIES OF THE NELDER-MEAD SIMPLEX METHOD IN LOW DIMENSIONS *; 1998; Vol. 9, pp 112–147. http://www.siam.org/journals/siopt/9-1/30347.html.

(32) Klöckner, W.; Büchs, J. Advances in Shaking Technologies. *Trends in Biotechnology*. June 2012, pp 307–314. https://doi.org/10.1016/j.tibtech.2012.03.001.

(33) Abrini, J.; Naveau, H.; Nyns, E.-J. *Clostridium Autoethanogenum, Sp. Nov., an Anaerobic Bacterium That Produces Ethanol from Carbon Monoxide*; 1994; Vol. 161.

(34) Robinson, J. A.; Tiedje, J. M. Kinetics of Hydrogen Consumption by Rumen Fluid, Anaerobic Digestor Sludge, and Sedimentt. *Appl Environ Microbiol* **1982**, *44* (6), 1374–1384.

(35) Karadagli Andrew Marcus Bruce E Rittmann, F. K. Role of Hydrogen (H 2) Mass Transfer in Microbiological H 2-Threshold Studies. *Biodegradation* **2019**, *30*, 113–125. https://doi.org/10.1007/s10532-019-09870-1.

(36) Robinson, J. A.; Tiedje, J. M. *Nonlinear Estimation of Monod Growth Kinetic Parameters from a Single Substrate Depletion Curve*; 1983; Vol. 45. https://journals.asm.org/journal/aem.

(37) Novak, K.; Neuendorf, C. S.; Kofler, I.; Kieberger, N.; Klamt, S.; Pflügl, S. Blending Industrial Blast Furnace Gas with H2 Enables Acetobacterium Woodii to Efficiently Co-Utilize CO, CO2 and H2. *Bioresour Technol* **2021**, *323*. https://doi.org/10.1016/j.biortech.2020.124573.

(38) Bengelsdorf, F. R.; Poehlein, A.; Linder, S.; Erz, C.; Hummel, T.; Hoffmeister, S.; Daniel, R.; Dürre, P. Industrial Acetogenic Biocatalysts: A Comparative Metabolic and Genomic Analysis. *Front Microbiol* **2016**, *7* (JUL). https://doi.org/10.3389/fmicb.2016.01036.

(39) An, T.; Kim, Y. K. Effect of Selenium and Tungsten on Cell Growth and Metabolite Production in Syngas Fermentation Using "Clostridium Autoethanogenum." *J Biotechnol* **2022**, *356*, 60–64. https://doi.org/10.1016/j.jbiotec.2022.07.004.

(40) Lanzillo, F.; Pisacane, S.; Raganati, F.; Russo, M. E.; Salatino, P.; Marzocchella, A. Optimization of CO Fermentation by Clostridium Carboxidivorans in Batch Reactors: Effects of the Medium Composition. *Anaerobe* **2024**, *87*. https://doi.org/10.1016/j.anaerobe.2024.102855.

(41) Groher, A.; Weuster-Botz, D. General Medium for the Autotrophic Cultivation of
Acetogens. *Bioprocess Biosyst Eng* **2016**, *39* (10), 1645–1650.
https://doi.org/10.1007/s00449-016-1634-5.

(42) Schuchmann, K.; Chowdhury, N. P.; Müller, V. Complex Multimeric [FeFe]
Hydrogenases: Biochemistry, Physiology and New Opportunities for the Hydrogen
Economy. *Front Microbiol* **2018**, *9*. https://doi.org/10.3389/fmicb.2018.02911.

(43) Madjarov, J.; Soares, R.; Paquete, C. M.; Louro, R. O. *Sporomusa Ovata* as Catalyst for
Bioelectrochemical Carbon Dioxide Reduction : A Review across Disciplines from
Microbiology to Process Engineering. *Front Microbiol* **2022**, *13*, 1–13.
https://doi.org/10.3389/fmicb.2022.913311.

(44) Peters, J. W.; Schut, G. J.; Boyd, E. S.; Mulder, D. W.; Shepard, E. M.; Broderick, J. B.;
King, P. W.; Adams, M. W. W. [FeFe]- and [NiFe]-Hydrogenase Diversity, Mechanism,
and Maturation. *Biochimica et Biophysica Acta - Molecular Cell Research*. Elsevier June
1, 2015, pp 1350–1369. https://doi.org/10.1016/j.bbamcr.2014.11.021.

(45) Zacarias, S.; Temporão, A.; Barrio, M. Del; Fourmond, V.; Léger, C.; Matias, P. M.;
Pereira, I. A. C. A Hydrophilic Channel Is Involved in Oxidative Inactivation of a [NiFeSe]
Hydrogenase. *ACS Catal* **2019**, *9* (9), 8509–8519.
https://doi.org/10.1021/acscatal.9b02347.

(46) Leroux, F.; Bastien Dementin, S.; Né Dicte Burlat, B.; Cournac, L.; Volbeda, A.; Phanie
Champ, S.; Martin, L.; Guigliarelli, B.; Bertrand, P.; Fontecilla-Camps, J.; Rousset, M.; Lé,
C. *Experimental Approaches to Kinetics of Gas Diffusion in Hydrogenase*; 2008.
www.pnas.orgcgidoi10.1073pnas.0803689105.

(47) Daniel, S. L.; Hsu, T.; Dean, S. I.; Drake, H. L. *Characterization of the H2-and CO-*
Dependent Chemolithotrophic Potentials of the Acetogens Clostridium
Thermoaceticum and Acetogenium Kivuit; 1990; Vol. 172.
https://journals.asm.org/journal/jb.

(48) Zhu, H.-F.; Liu, Z.-Y.; Zhou, X.; Yi, J.-H.; Lun, Z.-M.; Wang, S.-N.; Tang, W.-Z.; Li, F.-L.
Energy Conservation and Carbon Flux Distribution During Fermentation of CO or
H2/CO2 by Clostridium Ljungdahlii. *Front Microbiol* **2020**, *11*.
https://doi.org/10.3389/fmicb.2020.00416.

(49) Bardone, E.; Marzocchella, A.; Keshavarz, T.; Stoll, K. I.; Herbig, S.; Zwick, M.; Boukis,
N.; Sauer, J.; Neumann, A.; Oswald, F. Fermentation of H2 and CO2 with Clostridium

673   Ljungdahlii at Elevated Process Pressure-First Experimental Results. *Chem Eng Trans*

674   **2018**, *64*. https://doi.org/10.3303/CET1864026.

675 (50) Boto, S. T.; Bardl, B.; Harnisch, F.; Rosenbaum, M. A. Microbial Electrosynthesis with

676   Clostridium Ljungdahlii Benefits from Hydrogen Electron Mediation and Permits a

677   Greater Variety of Products †. **2023**. https://doi.org/10.1039/d3gc00471f.

678 (51) Martínez-Ruano, J.; Suazo, A.; Véliz, F.; Otálora, F.; Conejeros, R.; González, E.; Aroca,

679   G. Electro-Fermentation with Clostridium Autoethanogenum: Effect of PH and Neutral

680   Red Addition. *Environ Technol Innov* **2023**, *31*.

681   https://doi.org/10.1016/j.eti.2023.103183.

682 (52) Valgepea, K.; Talbo, G.; Takemori, N.; Takemori, A.; Ludwig, C.; Mahamkali, V.; Mueller,

683   A. P.; Tappel, R.; Köpke, M.; Simpson, S. D.; Nielsen, L. K.; Marcellin, E. Absolute

684   Proteome Quantification in the Gas-Fermenting Acetogen Clostridium

685   Autoethanogenum . *mSystems* **2022**, *7* (2). https://doi.org/10.1128/msystems.00026-

686   22.

687 (53) Arantes, A. L.; Moreira, J. P. C.; Diender, M.; Parshina, S. N.; Stams, A. J. M.; Alves, M.

688   M.; Alves, J. I.; Sousa, D. Z. Enrichment of Anaerobic Syngas-Converting Communities

689   and Isolation of a Novel Carboxydotrophic Acetobacterium Wieringae Strain JM. *Front*

690   *Microbiol* **2020**, *11* (January). https://doi.org/10.3389/fmicb.2020.00058.

691 (54) Moreira, J. P. C.; Heap, J. T.; Alves, J. I.; Domingues, L. Developing a Genetic Engineering

692   Method for Acetobacterium Wieringae to Expand One-Carbon Valorization Pathways.

693   *Biotechnology for Biofuels and Bioproducts* **2023**, *16* (1).

694   https://doi.org/10.1186/s13068-023-02259-6.

695 (55) Marshall, C. W.; Ross, D. E.; Handley, K. M.; Weisenhorn, P. B.; Edirisinghe, J. N.; Henry,

696   C. S.; Gilbert, J. A.; May, H. D.; Norman, & R. S. Metabolic Reconstruction and Modeling

697   Microbial Electrosynthesis. **2017**, *7* (839). https://doi.org/10.1038/s41598-017-08877-

698   z.

699 (56) Mills, S.; Dessì, P.; Pant, D.; Farràs, P.; Sloan, W. T.; Collins, G.; Zeeshan Ijaz, U. A Meta-

700   Analysis of Acetogenic and Methanogenic Microbiomes in Microbial Electrosynthesis.

701   *NPJ Biofilms Microbiomes* **2022**, *8* (73). https://doi.org/10.1038/s41522-022-00337-5.

702 (57) Van Hecke, W.; Bockrath, R.; De Wever, H. Effects of Moderately Elevated Pressure on

703   Gas Fermentation Processes. *Bioresource Technology*. Elsevier Ltd December 1, 2019.

704   https://doi.org/10.1016/j.biortech.2019.122129.

705   (58)   Prévoteau, A.; Carvajal-Arroyo, J. M.; Ganigué, R.; Rabaey, K. Microbial Electrosynthesis
706   from CO2: Forever a Promise? *Curr Opin Biotechnol* **2020**, *62*, 48–57.
707   https://doi.org/10.1016/j.copbio.2019.08.014.

708   (59)   Tsapekos, P. ; Alvarado-Morales, M. ;; Angelidaki, I. H2 Competition between
709   Homoacetogenic Bacteria and Methanogenic Archaea during Biomethanation from a
710   Combined Experimental-Modelling Approach. *J Environ Chem Eng* **2022**, *10* (2).
711   https://doi.org/10.1016/j.jece.2022.107281.

712   (60)   Tremblay, P.-L.; Zhang, T. Genetic Tools for the Electrotroph Sporomusa Ovata and
713   Autotrophic Biosynthesis. *Biotechnology | Full-Length Text* **2023**.
714   https://doi.org/10.1128/aem.01757-23.

715   (61)   Bourgade, B.; Minton, N. P.; Islam, M. A. Genetic and Metabolic Engineering Challenges
716   of C1-Gas Fermenting Acetogenic Chassis Organisms. *FEMS Microbiol Rev* **2021**, *008*,
717   1–20. https://doi.org/10.1093/femsre/fuab008.
718