Concurrent Joint Contact in Anterior Cruciate Ligament Injury induces cartilage

micro-injury and subchondral bone sclerosis, resulting in knee osteoarthritis

Authors

Kei Takahata1,2, Yu-Yang Lin3, Benjamin Osipov3, Kohei Arakawa2,4, Saaya Enomoto5,

Blaine A. Christiansen3*, Takanori Kokubun1,6*

Affiliations

1 Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University,

Koshigaya, Saitama, Japan

2 Japan Society for the Promotion of Science, Tokyo, Japan

3 Department of Orthopaedic Surgery, University of California Davis Health, Sacramento,

CA, United States

4 Laboratory of Animal Regeneration Systemology, Department of Life Sciences, School of

Agriculture, Meiji University, Kanagawa, Japan

5 Saitama Prefecture Rehabilitation Center, Saitama, Japan

6 Department of Physical Therapy, School of Health and Social Services, Saitama Prefectural

University, Koshigaya, Saitama, Japan

Co-Correspondence

[3]Blaine A. Christiansen

University of California Davis Health, Department of Orthopaedic Surgery, Lawrence J.

Ellison Musculoskeletal Research Center, 2700 Stockton Ave, Suite 2301 Sacramento, CA

95817, USA. Tel.: 916-734-3974

bchristiansen@ucdavis.edu

[1,6]Takanori Kokubun, Ph.D.

820 San-nomiya, Koshigaya, Saitama, 343-8540. Japan

Tel: +81 48-973-417617

kokubun-takanori@spu.ac.jp

ABSTRACT

Objective. Anterior Cruciate Ligament (ACL) injury develops the Osteoarthritis (OA) via

two distinct processes: initial direct micro-injury of the cartilage surface by compressive force

during ACL injury, and secondary joint instability due to the deficiency of the ACL. Using

the conventional Compression-induced ACL-R and novel Non-Compression ACL-R models,

we aimed to reveal the individual effects on OA progression after ACL injury.

Methods. Twelve-week-old C57BL/6 male were randomly divided to three experimental

groups: Compression ACL-R, Non-Compression ACL-R, and Intact. We performed the joint

laxity test and microscope analysis at 0 days, in vivo imaging with matrix-metalloproteinases

(MMPs) at 3 and 7 days, histological and micro-CT analysis at 0, 7, 14, and 28 days.

Results. Although no differences in the joint laxity were observed between both ACL-R

groups, the Compression ACL-R group exhibited a significant increase of cartilage roughness

immediately after injury compared with the Non-Compression group. At 7 days, Compression

group increased MMPs-induced fluorescence intensity slightly and MMP-13 positive cell

ratio of chondrocytes significantly than that in the Non-Compression group. Moreover,

histological cartilage degeneration initiated in the whole joint level of the Compression group

at the same time point. Micro-CT analysis revealed that sclerosis of tibial Subchondral bone

in the Compression group developed significantly more than in the Non-Compression group

at 28 days, especially in the medial tibial compartment.

51   **Conclusions.** Concurrent joint contact during ACL rupture caused initial micro-damage

52   on the cartilage surface and early cartilage degeneration with MMP-13 production, leading to

53   later bone formation in the subchondral bone.

Key Words:

56   Osteoarthritis, Animal model, ACL injury, Mechanical stress, Subchondral bone

INTRODUCTION

Osteoarthritis (OA) is a major chronic musculoskeletal disease that progresses

irreversibly and negatively influences the quality of life of OA patients. The number of OA

patients is rapidly increasing, with a 30.8% increase in OA prevalence between 2007 and

2017 [1]. OA can be divided into two types depending on the mechanisms of initiation:

primary OA, which is idiopathic and generally occurs during aging, and secondary post-

traumatic OA (PTOA), which is initiated by a traumatic joint injury such as a meniscus tear or

ligament injury. PTOA accounts for approximately 12% of symptomatic OA patients [2] and

has an increasing incidence rate due to the increasing popularity of high-impact sports [3].

Therefore, elucidating the mechanisms of PTOA initiation and progression is necessary to

find therapeutic strategies to prevent or slow the progression of OA.

Anterior Cruciate Ligament (ACL) injury is a known initiator of PTOA

development. The annual incidence of ACL injury in the general population is 68.6 per

100,000 people [4], and the incidence of PTOA following ACL injury is as high as 87% [5].

In the etiology of OA following ACL injury, concomitant injuries of other joint structures

along with ACL rupture and increased mechanical stress caused by joint instability are likely

contributing factors. Superphysiological compressive force to the joint surfaces are common

during ACL injury. Almost half of ACL injury patients suffer from initial damage of articular

cartilage [6], and concomitant cartilage injury may increase OA risk up to 2.4 times at 19

years [7]. In addition, 80-90% of patients with an acute ACL injury also show signs of

subchondral bone lesions measured using magnetic resonance imaging (MRI). [5]

Subchondral bone changes can affect cartilage degeneration via crosstalk in molecular and

biomechanical interactions [8], suggesting that acute subchondral bone injuries may also

initiate OA progression.

Because the ACL restricts the anterior-posterior (AP) translation of the tibia relative

to the femur [9], joint instability after ACL injury can change the articulation of the knee joint

and induce abnormal mechanical stress, which may stimulate knee joint tissues to produce

catabolic enzymes. Since knee PTOA develops several decades after ACL injury, it was

assumed that secondary joint instability is the most important factor, and surgical

reconstruction of the ACL has been widely used to restore joint stability and reduce the risk

of PTOA development [10]. However, a recent systematic review reported that reconstruction

surgery may not affect PTOA progression [11], which suggests that the initial response to

compressive forces during injury may also plays a significant role in the pathogenesis of

PTOA. Since cartilage is an avascular tissue with poor self-healing capability, acute damage

to the articular chondral surface and subchondral bone may be involved with the initiation of

PTOA.

ACL transection (ACL-T) is a common model for inducing knee OA in small

animals, but this method also induces unnecessary acute inflammation in articular tissues due

to the invasive surgical procedures [12-15]. In contrast, the Non-Invasive Compression ACL

Rupture (ACL-R) model was developed and used to explore mechanisms of PTOA initiation

and progression following ACL injury without the intra-articular inflammation associated

with surgical procedures [16-20]. ACL-R is induced by applying compressive force to the

joint, which causes anterior tibial dislocation and ACL injury. This method largely imitates

acute ACL injury and knee PTOA progression after injury. However, it is difficult to separate

the effects of initial overload of the articular cartilage and the secondary effects of joint

instability. We recently established a novel Non-Compression ACL-R model, which is made

without compression force on the cartilage surface and therefore induces no articular injuries

other than ACL rupture [21]. Based on the findings of this model, we hypothesized that

comparing the Compression ACL-R model and the novel Non-Compression ACL-R model

will allow us to investigate the individual effects of initial tissue damage due to compression

vs. secondary instability on PTOA progression.

MATERIALS AND METHODS

Animals and Experimental Design

Twelve-week-old C57BL/6J male mice were randomly divided into three

experimental groups: Compression ACL-R (n = 49), Non-Compression ACL-R (n = 48), and

Intact (n=21). Mice were euthanized at 0, 3, 7, 14, and 28 days after injury, and knee joints

were dissected for analysis (Fig. 1A). In the joint laxity test, microscopic, morphological, and

histological analysis at day 0 after injury, contralateral knee joints from the Non-Compression

group were used as the Intact group. Mice were allowed to acclimate for 1-2 weeks in the

vivarium before the start of the experiment, and were cared for in accordance with the

guidelines set by the National Institutes of Health (NIH) on the care and use of laboratory

animals. All procedures were conducted under approved Institutional Animal Care and Use

Committee protocols at Saitama Prefectural University and the University of California

Davis.

Creating Compression ACL-R and Non-Compression ACL-R Models

All procedures were performed on the right knee joint of each anesthetized mouse

under 1-4% inhaled isoflurane. The Compression ACL-R model was created via tibial

compression overload as previously described [18, 22]. Right knee joints were positioned in

an electromagnetic materials machine (ElectroForce 3200, TA Instruments, New Castle, DE),

and a single tibial compressive overload was applied at 1 mm/s to induce ACL rupture. The

Non-Compression ACL-R model was created based on our previous study [23]. The knee

joint was fixed at 90 degrees using surgical tape on a stand, and a force was slowly applied by

the thumb tip of the operator along the long axis of the femur. The applied force was stopped

quickly after hearing a distinct popping sound, which indicated ACL rupture. Anterior tibial

translation was manually confirmed to verify ACL injury. For both models, buprenorphine

analgesia (0.1 mg/kg) was injected immediately following injury for pain relief.

Anterior-Posterior Joint Laxity Testing

Knee joints were collected immediately post-injury, and femurs and tibias were

embedded in brass tubes using polymethylmethacrylate as previously described [22, 24].

Tibias were fixed to the load cell; then femurs were secured so that the angle of the knee joint

was either 60 □ or 90 □. Tibias were able to freely translate and rotate about the superior-

inferior axis. Five anterior-posterior (AP) loading cycles were applied perpendicular to the

longitudinal axis of the tibia to a target force of ±1.5 N at a loading rate of 0.5 mm/s. The

degree of AP joint laxity was quantified based on the difference between displacement at +0.8

N and -0.8 N.

Microscopic Analysis of Cartilage Surface Roughness

For samples collected 0 days following injury, soft tissue was removed carefully to

expose the tibial plateau, then 3D optical profilograph images of the whole tibial plateau were

taken (VR-6000; Keyence, JPN). The region of interest (ROI) was set as an ellipse with 0.6-

1.2 mm diameters in the center area of the medial and lateral tibial plateau (Figure 2A). The

arithmetical mean height and maximum height of the cartilage surface were measured by

normalizing to the contralateral intact knee of each mouse. Methods are further described in

the Supplementary Materials.

Fluorescent Reflectance Imaging (FRI)

157   At 3 and 7 days post-injury, mice were imaged in vivo using an optical imaging

system (IVIS Spectrum, PerkinElmer, Waltham, MA), after IV administration of a near

infrared probe that is activated by matrix-metalloproteinases (MMPSense 680, PerkinElmer,

Waltham, MA). The ROI was set as a circle of 0.7 mm^2 that surrounded the tibial tuberosity

to the superior border of the patella, and image processing and quantification were performed

via IVIS Living Image software as previously described [25]. To unify the mouse-to-mouse

variation in the delivery of the fluorescent probe, the radiant efficiency of the ACL-R knee

was normalized to the contralateral intact knee of each mouse. Methods are further described

in the Supplementary Materials.

Micro-Computed Tomography Analysis of Osteophyte Formation and Epiphyseal Bone

Microstructure

169   Bilateral knees were scanned using micro-computed tomography (μCT 35,

SCANCO, Brüttisellen, Switzerland) with the following parameters: nominal voxel size = 10

μm, energy = 55 kVp, intensity = 72μA, integration time = 800 ms. For analysis of

osteophyte formation at 7, 14, and 28 days after injury, VOI contouring included all

heterotopic mineralized tissue around the joint, as well as the patella, fabellae, and menisci.

The average bone volume of the patella, fabellae, and menisci in the Intact group was used as

the "baseline" volume to calculate osteophyte formation in the Compression and Non-

Compression groups. The difference in total bone volume between the injured mouse knees

and the baseline volume was calculated to determine the total osteophyte volume for each

injured joint.

We also assessed the morphological changes in epiphyseal trabecular bone at 0, 7,

14, and 28 days after injury. Morphological analysis of trabecular bone in the tibia and femur

was performed by manually drawing contours on 2D transverse slices; the ROI was designed

as the trabecular bone enclosed by the growth plate and subchondral cortical bone plate

(Figure 3B). Additionally, whole subchondral bone including the cortical bone plate and

trabecular bone in the medial and lateral tibial compartments were also measured (Figure 4A).

Using the manufacturer's analysis software, we quantified apparent bone mineral density

(BMD, g/cm^3), bone volume per total volume (BV/TV, %), trabecular number (Tb.N, 1/mm),

trabecular thickness (Tb.Th, mm), and trabecular separation (Tb.Sp, mm).

Histological Analysis for Cartilage Degeneration

After performing µCT analysis, knee joints were decalcified in 10%

ethylenediaminetetraacetic acid (EDTA) for 2 weeks, dehydrated in 70% and 100% ethanol

and xylene, and embedded in paraffin blocks. The samples were cut in the sagittal plane (7

µm thickness) using a microtome (ROM-360; Yamato Kohki Industrial Co., Ltd., Saitama,

JPN). Then, safranin-O/fast green staining was performed to evaluate articular cartilage

degeneration in the medial and lateral tibial compartments using the Osteoarthritis Research

Society International (OARSI) histopathological grading system [26]. These assessments

were conducted by two independent observers blinded to all other sample information. We

initially assessed the whole joint pathology in the knee joint, then assessed the medial and

lateral tibial plateau separately. The mean of the observer's scores was used as a

representative value.

Immunohistochemical Analysis

Immunohistochemical (IHC) staining was performed using anti-MMP-13 (1:200, bs-

0575R, Bioss). Detailed protocols are described in the Supplementary Materials. We

calculated the ratio between the number of MMP-13 positive cells and the number of

chondrocytes in the anterior and posterior area of the articular cartilage with regions of

interest of 40,000 µm^2 (200 mm × 200 mm). As with the OARSI score, we initially evaluated

the whole joint and then subsequently assessed the medial and lateral tibial plateau separately.

Statistical Analysis

Statistical analysis was performed using RStudio. We conducted the Shapiro-Wilk test

to verify the normality of all data. Student's t-test was used for the osteophyte formation and

Wilcoxon rank sum was used for the microscope and FRI data. One-way analysis of variance

was performed for the joint laxity test and MMP-13 IHC analysis initially, and then the

Tukey-Kramer test was used for post-hoc analysis. The Kruskal-Wallis test was used to

compare subchondral bone μCT data and OARSI scores, and the Steel-Dwass method was

used for the subsequent multiple comparisons. Parametric data are expressed as the mean ±

95% confidence intervals (95% CI); non-parametric data are expressed as the median ±

interquartile ranges. Statistical significance was set at $p < 0.05$.

RESULTS

AP joint Laxity Test of the Knee

Compared with the Intact group, both the Compression and Non-Compression ACL-

R groups had significantly increased AP joint laxity at both 60 ☐ and 90 ☐ (Figure 1B).

However, no significant differences were observed between the Compression and Non-

Compression groups at either angle. This suggests that any differences in OA progression

between the ACL-R groups can be attributed to the initial effect of compressive force at the

time of ACL injury.

[Insert Figure 1 here]

Figure 1. (A) Experimental design. We made the Compression ACL-R, Non-

Compression ACL-R, and Intact groups. Mice were euthanized at 0, 3, 7, 14, and 28 days

after injury and following analyses were performed; joint laxity test, microscopic, microCT,

and histological analysis. At day 0 after injury, contralateral knee joints from the Non-

Compression group were used as the Intact group. (B) A result of joint laxity test. The

Compression and Non-Compression ACL-R groups significantly increased Anterior-Posterior

joint laxity at both 60 ▢ and 90 ▢. However, no significant differences were observed

between both injury groups. Data are presented as the mean ± 95% CI. *P< 0.05; ***P<

0.001.

Microscope Analysis of Cartilage Roughness in the Tibial Plateau

Surface roughness in the region not covered by menisci was increased in the

Compression ACL-R group, especially on the medial tibial plateau. The normalized

arithmetical mean height of the medial compartment in the Compression group was

significantly higher than that in the Non-Compression group, but no difference was observed

in the lateral compartment (Figure 2A). In addition, there were no significant differences in

the maximum height of the medial and lateral compartments between groups. These results

indicate that compressive force during compression ACL injury caused initial micro-damage

in the surface layer of articular cartilage.

Quantification of MMP Fluorescence Intensity

At 3 days following injury, MMP-induced fluorescence intensity around ACL-R

knees increased slightly in the Compression (Normalized data: 0.31 ± 0.659) and Non-

Compression groups (0.512 ± 0.472) compared with each contralateral intact knee, and no

significant difference was observed between groups (Figure 2B). Whereas, at 7 days,

fluorescence intensity increased moderately in the Compression (0.684 ± 0.597) and slightly

in the Non-Compression (0.17 ± 0.368) groups. However, there were no significant

differences between these groups.

Figure 2. (A) Microscope Analysis of Cartilage Roughness in the Tibial Plateau at

day 0 after injury. The ROI was set as an ellipse with 0.6-1.2 mm diameters in the center area

of the medial and lateral tibial plateau. Compression group significantly increased the

normalized arithmetical mean height in the medial compartment compared with the Non-

Compression group. (B) Quantification of MMP Fluorescence Intensity. Normalized MMP-

induced fluorescence intensity around ACL-R knees increased in the Compression, especially

at 7 days following injury. However, there were no significant differences between these

groups. Data are presented as the median ± interquartile range. *$P< 0.05$.

MicroCT Measurement of Osteophyte Formation

Mineralized osteophyte volume was quantified at 14 and 28 days because no

osteophytes were observed at 7 days following injury. Similarly, only a small amount of

mineralized osteophyte formation was observed in both ACL-R groups at 14 days (Figure

3A). At 28 days following injury, a greater volume of ectopic mineralized tissue was

observed in the medial femoral condyle and tibial plateau of injured joints. However, there

were no significant differences between ACL-R groups.

MicroCT Analysis of Epiphysis Trabecular Bone of the Distal Femur and Proximal

Tibia

At 0 days post-injury, no micro injuries or fractures were observed in the femoral and

tibial epiphysis trabecular bone in either ACL-R group, and there were no significant

differences in the microstructure between any experimental groups (Supplementary Figure

2A-B). This indicates that compressive force while inducing ACL rupture didn't cause

concomitant structural failures in the subchondral trabecular bone. Femoral epiphysis

trabecular bone showed a significant decrease of BMD, BV/TV, and Tb.Th in the

Compression and Non-Compression groups compared with the Intact group at 7 days, which

suggests acute bone absorption at an early time point in both ACL-R groups (Figure 3C).

These parameters were still significantly lower in the Compression and Non-Compression

groups at 14 days. However, the Compression group recovered bone volume moderately at 28

days, and no significant difference in the BMD was observed between the Intact group and

the Compression group at this time point, although the Non-Compression group was still

significantly lower than the Intact group. Tb.N and Tb.Sp results are described in the

Supplementary Material (Supplementary Figure 3A).

Similarly, tibial epiphysis trabecular bone also exhibited a significant decrease in

BMD and BV/TV in the Compression and Non-Compression groups at 7 days following

injury (Figure 3D). At 14 days, the significantly decreased BMD and BV/TV was still present

in the Non-Compression group, whereas the Compression group slightly increased bone

volume and there was no significant difference in BV/TV between the Intact and the

Compression groups. Furthermore, in the Compression group at 28 days, the bone volume

increased to the same level as in the Intact group. The Non-Compression group still showed

deficits in bone microstructure at this time point, therefore the BMD and BV/TV in the Intact

and Compression group were significantly greater than in the Non-Compression group. Tb.N

and Tb.Sp results are described in the Supplementary Material (Supplementary Figure 3B).

Figure 3. (A) Reconstructed images obtained by microCT and measurement of

osteophyte formation. Only a small amount of mineralized osteophyte at 14 days and a greater

volume of ectopic mineralized tissue in the medial femoral condyle and tibial plateau of

injured joints at 28 days were observed in the Compression and Non-Compression groups.

However, there were no significant differences. Black arrows show the osteophyte. (B) The

ROI to measure the tibial and femoral epiphysis trabecular bone and (C) the result of

microCT Analysis. Femoral epiphysis trabecular bone significantly decreased BMD, BV/TV,

and Tb.Th in the Compression and Non-Compression groups compared with the Intact group

at 7 days. At 28 days, no significant difference in the BMD was observed between the Intact

and the Compression groups. However, the Non-Compression group was still significantly

lower than the Intact group at the same timepoint. Tibial epiphysis trabecular bone also

showed a significant decrease in BMD and BV/TV in both injury groups at 7 days. However,

at 28 days, the BMD and BV/TV in the Intact and Compression groups were significantly

greater than in the Non-Compression group. Data are presented as the median ± interquartile

range. *P< 0.05. **P< 0.01.

MicroCT Analysis of Subchondral Bone in the Medial and Lateral Tibial Compartment

Analysis of subchondral bone microstructure in the medial and lateral tibial

compartments at 7, 14, and 28 days (Figure 4A) showed that in the medial compartment, the

Compression and the Non-Compression groups caused bone loss with a significant decrease

in BV/TV and Tb.Th compared to the Intact group at 7 days (Figure 4B). Although BV/TV

and Tb.Th in the Non-Compression group were still significantly lower than in the Intact

group at 14 days, the Compression group increased BV/TV and Tb.Th and no differences

were observed compared to other groups. Additionally, the Compression group exhibited

significantly increased BV/TV compared to the Non-Compression group at 28 days.

In the lateral compartment, both ACL-R groups exhibited significant bone loss

compared with the Intact group at 7 days similar to what was observed in the medial

compartment (Figure 4C). At 14 days, the Compression and Non-Compression groups

increased BV/TV and Tb.Th, and no differences were observed between any experimental

groups. Unlike the medial compartment, the Compression group exhibited a significant

increase of BV/TV and Tb.Th compared to the Intact group at 28 days. Tb.N and Tb.Sp

results for both compartments are described in the Supplementary Materials (Supplementary

Figure 4A-B).

Figure 4. (A) The ROI of medial and lateral tibial trabecular bone and microCT

images. (B) Analysis of subchondral bone microstructure in the medial and lateral tibial

compartments. In the medial compartment, the Compression and the Non-Compression

groups caused bone loss with a significant decrease in BV/TV and Tb.Th compared to the

Intact group at 7 days. At 28 days, the Compression group showed significantly increased

BV/TV compared to the Non-Compression group. In the lateral compartment, both ACL-R

groups exhibited significant bone loss compared with the Intact group at 7 days. Unlike the

medial compartment, the Compression group exhibited a significant increase of BV/TV and

Tb.Th compared to the Intact group at 28 days. Data are presented as the median ±

interquartile range. *P< 0.05. **P< 0.01.

Histological Analysis of Cartilage Degeneration in the Medial and Lateral Tibial

Compartment

Histological analysis did not detect any acute injuries of articular tissues other than

the ACL (cartilage, menisci, subchondral bone) in the knee joint at 0 days following injury

(Supplementary Figure 5). In the medial tibial plateau at 7 days, over half of the mice in the

Compression group and some mice in the Non-Compression group developed a small loss of

cartilage around the posterior edge, and cell proliferation and staining of Safranin-O were

observed in the posterior region (Figure 5A). At 14 days, fibrillation of the cartilage surfaces,

mild to moderate cartilage erosion, and osteophytes on the posterior tibia were observed in

both ACL-R groups, especially in the posterior region. At 28 days, erosion extending to the

center and posterior region of the subchondral bone and growth plate were observed in both

ACL-R groups. These histological changes were observed in the lateral tibial plateau as well;

however, the degree of OA progression was milder compared to the medial tibial plateau at all

time points (Figure 5B).

The OARSI score for the whole joint pathology in the Compression group increased

significantly compared with the Intact group at 7 days (Figure 5C). Moreover, OARSI score

in both ACL-R groups increased significantly compared with the Intact group at 14 days and

28 days. For the medial tibial plateau, OARSI score in both ACL-R groups were significantly

higher than that in the Intact group at all time points (Figure 5D). For the lateral tibial plateau,

OARSI score in both ACL-R groups were significantly higher than that in the Intact group at

14 days and 28 days (Figure 5E).

[Insert Figure 5 here] scale bar 500μm

Figure 5. (A) Histological images of cartilage degeneration in the medial and lateral

tibial compartment. (B) The OARSI score for the whole joint pathology in the Compression

group was significantly higher than that in the Intact group at 7 days. At 14 and 28 days, both

ACL-R groups significantly increased the OARSI score compared with the Intact group. (C)

In the medial compartments, OARSI score in both ACL-R groups were significantly higher

than that in the Intact group at all time points. (D) In the lateral compartment, both ACL-R

groups caused a significant increase of OARSI score compared with the Intact group at 14

and 28 days. Data are presented as the median ± interquartile range. *P< 0.05. **P< 0.01.

Black scale bar, 500 µm.

Immunohistochemical Analysis of the Articular Cartilage

IHC analysis of MMP-13 was performed at 7 and 14 days following injury because

cartilage was highly degenerated by 28 days (Figure 6A-B). At 7 days, the number of MMP-

13-positive cells of the whole joint in the Compression group increased significantly

compared with the Intact and Non-Compression groups (Figure 6C). Moreover, the Non-

Compression group also significantly increased the positive cell ratio compared to the Intact

group. Analysis of the medial tibial plateau showed similar results to the whole joint (Figure

6D), whereas for the lateral tibial plateau, the number of positive cells in the Compression

group increased significantly compared with the other groups (Figure. 6E). At 14 days, the

positive cells ratio of the whole joint in both ACL-R groups increased significantly compared

with the Intact group. Similar results were observed in the medial and lateral tibial plateau.

Figure 6. (A-B) Immunohistochemical images of MMP-13 in cartilage in the medial

and lateral tibial plateau. (C-D) In the whole joint level and medial tibial plateau,

Compression group increased the MMP-13 positive cell rate significantly compared with the

Intact and Non-Compression groups at 7 days. Furthermore, the Non-Compression group also

significantly increased the positive cell ratio compared to the Intact group. At 14 days, the

positive cells ratio in both ACL-R groups increased significantly compared with the Intact

group. (E) In the lateral tibial plateau at 7 days, the number of MMP-13 positive cells in the

Compression group increased significantly compared with the other groups. At 14 days, both

ACL-R groups increased the MMP-13 positive cell rate significantly compared with the Intact

group. Data are presented as the mean $\pm$ 95% CI. **$P< 0.01$; ***$P< 0.001$. Black scale bar,

100 μm. (F) Schematic summarizing the mechanism of OA development following ACL

injury with or without compression concomitant compression force. Compression-ACL

rupture in mice caused initial micro-damage on the cartilage surface and early cartilage

degeneration with MMP-13 production, leading to later bone formation in the subchondral

bone.

DISCUSSION

The mechanism of OA development following ACL injury involves two distinct

processes: initial direct micro-injury of the cartilage surface as a result of the compressive

force experienced during ACL injury, and secondary joint instability due to the deficiency of

the ACL. In this study, we compared the conventional Compression-induced ACL-R and

Non-Compression ACL-R models, with the aim of clarifying the individual effects of the

initial injury response and secondary joint instability on OA progression after ACL injury.

Although no significant differences in the joint laxity were observed between ACL-R groups,

the roughness of the cartilage immediately after injury was increased significantly in the

Compression ACL-R group but not the Non-Compression group. Similarly, MMP-induced

fluorescence intensity at 7 days was higher in the Compression group than in the Non-

Compression group. Moreover, the MMP-13 positive cell ratio in the Compression group was

also increased significantly compared to the Non-Compression group at 7 days and micro-CT

analysis revealed that osteophyte volume in the Compression group was slightly higher than

in the Non-Compression group. Furthermore, sclerosis of tibial SCB in the Compression

group developed significantly more than in the Non-Compression group at 28 days, especially

in the medial tibial compartment.

The ACL contributes to joint stability and ACL deficiency results in an increase in

mechanical stress [27]. Our assessment of AP joint laxity immediately after injury revealed

that the Compression and Non-Compression groups had significantly increased joint laxity

compared to the Intact group. However, no differences were observed between the ACL-R

groups (Figure 1B). This result indicates that any differences in OA progression between the

ACL-R groups can be attributed to the direct effect of compressive stress experienced while

inducing ACL rupture.

Interestingly, microscopic analysis demonstrated that roughness on the surface of the

medial tibial plateau in the Compression group was significantly increased compared to the

Non-Compression group. However, no injuries such as cartilage erosion, meniscus injury, or

subchondral bone loss were observed histologically and morphologically immediately

following injury. A previous study using an intra-articular tibial plateau fracture mouse model

reported that compression force to the tibial plateau to a target load of 55 N at a rate of 60 N/s

caused massive soft tissue injuries, including to the synovium, meniscus, tendon, and

cartilage [28]. The mean compressive force applied with the Compression ACL-R model was

approximately 10 N at a loading rate of 1 mm/s [18]. Thus, the compressive force in the joint

during Compression ACL-R was remarkably lower than in the previous study using the tibial

plateau fracture model, therefore roughness on the cartilage surface indicated mild articular

damage in the Compression group rather than severe damage.

In addition to the initial micro-injury on the cartilage surface induced by compressive

loading, we also measured the biological response in the early stage using FRI with an MMP-

activatable probe. MMP-3, MMP-9, and MMP-13 levels increased in animal models of

osteoarthritis [29-33], and MMP-2, MMP-3, MMP-9, and MMP-13 increased in in vitro

models of osteoarthritis [34], which can degrade collagens and proteoglycans of the articular

cartilage. Compressive loading during ACL injury increases the production of matrix-

degrading enzymes and inflammatory cytokines in chondrocytes, which results in

chondrocyte apoptosis and cartilage degeneration [6, 35, 36]. A previous animal study

reported that even a single compressive load at 6 N without ACL rupture caused cartilage

degeneration with chondrocyte apoptosis [20]. However, in the current study no significant

differences were observed between ACL-R groups; the Compression group showed a high

MMP-induced fluorescence intensity using FRI at 7 days following injury. Furthermore, IHC

analysis demonstrated that MMP-13 expression in chondrocytes of the Compression group

increased significantly compared to the Non-Compression group. These molecular changes

may be due to the initial micro-injury detected as surface roughness, which is consistent with

early stage pathology of ACL injury in human patients. Interestingly, histological data also

showed that the OARSI score in the whole joint of the Compression group significantly

increased compared to the Intact group at 7 days. These results suggested that the

compressive force on the cartilage surface while inducing ACL rupture may have accelerated

cartilage degeneration in the early stage through the activation of MMPs. Alternatively,

considering that there was no difference in OARSI score at 14 and 28 days, it is also possible

that secondary joint instability is a more important factor in the progression of cartilage

degeneration after the initial stage.

Subchondral bone provides mechanical and nutritional support for cartilage, and

microenvironmental changes in subchondral bone might affect cartilage metabolism directly

or indirectly [37]. Generally, subchondral bone reacts to mechanical stress and is faster to

remodel than articular cartilage due to its robust innervation and blood supply that provide it

with a high capacity of turnover [38, 39]. We hypothesized that subchondral bone would react

to compressive force and induced structural changes during the early stage following ACL

injury. Indeed, we observed loss of subchondral bone volume in both ACL-R groups

following injury. However, μCT analysis showed that the Compression group induced

subchondral bone sclerosis in the medial tibial plateau compared to the Non-Compression

group at 28 days, and the lesion area was consistent with the cartilage degeneration. Some

reports have shown that cartilage degeneration precedes subchondral bone changes [40-42]

and osteoarthritic chondrocytes enhanced osteoblast differentiation in subchondral bone via

the ERK1/2 pathway [43]. Our results revealed that ACL injury with concomitant

compressive force initially caused micro-injury on the cartilage surface, then subchondral

bone remodeling and sclerosis in the medial compartment at an earlier time point than ACL

injury without compressive loading (Figure 6F). This earlier switch from bone loss to bone

sclerosis is consistent with an accelerated PTOA progression in the Compression ACL-R

group compared to the Non-Compression group.

There are two main limitations to this study. First, articular surface roughness

analysis was only performed immediately after the induction of each model. Microscopic

evaluation can determine the degree of cartilage degeneration as a surface rather than a line in

more detail. Roughness analysis was able to detect micro-injury on the cartilage surface at 0

days (Figure 2A), which was not possible to see with histological observation (Supplementary

Figure 5). Further roughness analysis at additional time points may enable us to reveal the

reaction of cartilage degeneration to compression force, especially in the early phase. Second,

we didn't investigate the molecular biological mechanism of subchondral bone remodeling

and sclerosis during PTOA progression. It has been shown that the proliferation and

differentiation of osteoprogenitors are stimulated by platelet-derived growth factor-A,

transforming gene factor-β1, and fibroblast growth factor-1, resulting in subchondral bone

formation [44]. We revealed that chondrocytes reacted to initial compression force and

increased MMP-13; therefore, additional experiments about the biological mechanism should

be performed to develop further understanding of these mechanisms.

In conclusion, concomitant joint contact while non-invasively inducing ACL rupture

in mice caused initial micro-damage on the cartilage surface and early cartilage degeneration

with MMP-13 production, leading to later bone formation in the subchondral bone.

Understanding the initial pathology of the ACL injury may be an important indicator of

disease etiology and represent a potential preventative approach for mitigating secondary OA

development.

Acknowledgments

The author(s) received no financial support for the research, authorship, and/or publication of

this article. We would like to thank Editage (www.editage.com) for English language editing.

The study design and summary scheme were created with BioRender.com. Research reported

in this publication was partially supported by the National Institute of Arthritis and

Musculoskeletal and Skin Diseases, part of the National Institutes of Health, under Award

Number R01 AR075013.

Author contributions

All authors approved the final submitted manuscript.

Study design: TK and BC

Making model and Data collection: KT and YL

Mechanical analysis: KT and BO

Fluorescent reflectance imaging: KT and YL

Morphological analysis: KT, YL, and KA

Histological analysis: KT, KA, and SE

Manuscript composition: KT, TK, and BC

Role of funding source

Competing interest statement

533   All authors have no conflicts of interest related to the manuscript.

References

536   1.   Disease, G.B.D., I. Injury, and C. Prevalence, *Global, regional, and national incidence, prevalence, and*
537   *years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990-2017: a*

systematic analysis for the Global Burden of Disease Study 2017. Lancet, 2018. **392**(10159): p. 1789-
1858.

2. Thomas, A.C., et al., *Epidemiology of Posttraumatic Osteoarthritis.* J Athl Train, 2017. **52**(6): p. 491-
496.

3. Lohmander, L.S., et al., *High prevalence of knee osteoarthritis, pain, and functional limitations in*
female soccer players twelve years after anterior cruciate ligament injury. Arthritis Rheum, 2004.
50(10): p. 3145-52.

4. Sanders, T.L., et al., *Incidence of Anterior Cruciate Ligament Tears and Reconstruction: A 21-Year*
Population-Based Study. Am J Sports Med, 2016. **44**(6): p. 1502-7.

5. Friel, N.A. and C.R. Chu, *The role of ACL injury in the development of posttraumatic knee*
osteoarthritis. Clin Sports Med, 2013. **32**(1): p. 1-12.

6. Riordan, E.A., C. Little, and D. Hunter, *Pathogenesis of post-traumatic OA with a view to intervention.*
Best Pract Res Clin Rheumatol, 2014. **28**(1): p. 17-30.

7. Whittaker, J.L., et al., *Risk factors for knee osteoarthritis after traumatic knee injury: a systematic*
review and meta-analysis of randomised controlled trials and cohort studies for the OPTIKNEE
Consensus. Br J Sports Med, 2022. **56**(24): p. 1406-1421.

8. Yuan, X.L., et al., *Bone-cartilage interface crosstalk in osteoarthritis: potential pathways and future*
therapeutic strategies. Osteoarthritis Cartilage, 2014. **22**(8): p. 1077-89.

9. Ajuied, A., et al., *Anterior cruciate ligament injury and radiologic progression of knee osteoarthritis: a*
systematic review and meta-analysis. Am J Sports Med, 2014. **42**(9): p. 2242-52.

10. Thorstensson, C.A., et al., *Choosing surgery: patients' preferences within a trial of treatments for*
anterior cruciate ligament injury. A qualitative study. BMC Musculoskelet Disord, 2009. **10**: p. 100.

11. Ferrero, S., et al., *Impact of anterior cruciate ligament surgery on the development of knee*
osteoarthritis: A systematic literature review and meta-analysis comparing non-surgical and surgical
treatments. Osteoarthr Cartil Open, 2023. **5**(3): p. 100366.

12. Goldring, M.B. and M. Otero, *Inflammation in osteoarthritis.* Current Opinion in Rheumatology, 2011.
23(5): p. 471-478.

13. Liu-Bryan, R., *Synovium and the Innate Inflammatory Network in Osteoarthritis Progression.* Current
Rheumatology Reports, 2013. **15**(5).

14. Mathiessen, A. and P.G. Conaghan, *Synovitis in osteoarthritis: current understanding with therapeutic*
implications. Arthritis Research & Therapy, 2017. **19**(1).

15. Scanzello, C.R. and S.R. Goldring, *The role of synovitis in osteoarthritis pathogenesis.* Bone, 2012.
51(2): p. 249-257.

16. Ramme, A.J., et al., *A novel rat model for subchondral microdamage in acute knee injury: a potential*
mechanism in post-traumatic osteoarthritis. Osteoarthritis Cartilage, 2016. **24**(10): p. 1776-1785.

17. Brown, S.B., et al., *Characterization of Post-Traumatic Osteoarthritis in Rats Following Anterior*
Cruciate Ligament Rupture by Non-Invasive Knee Injury (NIKI). J Orthop Res, 2020. **38**(2): p. 356-367.

18. Christiansen, B.A., et al., *Musculoskeletal changes following non-invasive knee injury using a novel*
mouse model of post-traumatic osteoarthritis. Osteoarthritis Cartilage, 2012. **20**(7): p. 773-82.

19.   Maerz, T., et al., *Articular cartilage degeneration following anterior cruciate ligament injury: a*
578   *comparison of surgical transection and noninvasive rupture as preclinical models of post-traumatic*
579   *osteoarthritis.* Osteoarthritis Cartilage, 2016. **24**(11): p. 1918-1927.

580   20.   Wu, P., et al., *Early response of mouse joint tissue to noninvasive knee injury suggests treatment*
581   *targets.* Arthritis Rheumatol, 2014. **66**(5): p. 1256-65.

582   21.   Takahata, K., et al., *Joint instability causes catabolic enzyme production in chondrocytes prior to*
583   *synovial cells in novel non-invasive ACL ruptured mouse model.* Osteoarthritis Cartilage, 2023. **31**(5):
584   p. 576-587.

585   22.   Hsia, A.W., et al., *Osteophyte formation after ACL rupture in mice is associated with joint*
586   *restabilization and loss of range of motion.* J Orthop Res, 2017. **35**(3): p. 466-473.

587   23.   Takahata, K., et al., *Joint instability causes catabolic enzyme production in chondrocytes prior to*
588   *synovial cells in novel non-invasive ACL ruptured mouse model.* Osteoarthritis Cartilage, 2022.

589   24.   Lockwood, K.A., et al., *Comparison of loading rate-dependent injury modes in a murine model of post-*
590   *traumatic osteoarthritis.* J Orthop Res, 2014. **32**(1): p. 79-88.

591   25.   Satkunananthan, P.B., et al., *In vivo fluorescence reflectance imaging of protease activity in a mouse*
592   *model of post-traumatic osteoarthritis.* Osteoarthritis Cartilage, 2014. **22**(10): p. 1461-9.

593   26.   Glasson, S.S., et al., *The OARSI histopathology initiative - recommendations for histological*
594   *assessments of osteoarthritis in the mouse.* Osteoarthritis Cartilage, 2010. **18 Suppl 3**: p. S17-23.

595   27.   Kraeutler, M.J., et al., *Anatomy and Biomechanics of the Native and Reconstructed Anterior Cruciate*
596   *Ligament: Surgical Implications.* Journal of Bone and Joint Surgery, 2017. **99**(5): p. 438-445.

597   28.   Stiffel, V., et al., *A Mouse Noninvasive Intraarticular Tibial Plateau Compression Loading-Induced*
598   *Injury Model of Posttraumatic Osteoarthritis.* Calcif Tissue Int, 2020. **106**(2): p. 158-171.

599   29.   Hu, H., et al., *Blocking of the P2X7 receptor inhibits the activation of the MMP-13 and NF-kappaB*
600   *pathways in the cartilage tissue of rats with osteoarthritis.* Int J Mol Med, 2016. **38**(6): p. 1922-1932.

601   30.   Raghu, H., et al., *CCL2/CCR2, but not CCL5/CCR5, mediates monocyte recruitment, inflammation and*
602   *cartilage destruction in osteoarthritis.* Ann Rheum Dis, 2017. **76**(5): p. 914-922.

603   31.   Qu, R., et al., *Ghrelin protects against osteoarthritis through interplay with Akt and NF-kappaB*
604   *signaling pathways.* FASEB J, 2018. **32**(2): p. 1044-1058.

605   32.   Schmidli, M.R., et al., *Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate*
606   *ligament disease.* BMC Vet Res, 2018. **14**(1): p. 161.

607   33.   Li, H., et al., *TNF-alpha increases the expression of inflammatory factors in synovial fibroblasts by*
608   *inhibiting the PI3K/AKT pathway in a rat model of monosodium iodoacetate-induced osteoarthritis.*
609   Exp Ther Med, 2018. **16**(6): p. 4737-4744.

610   34.   Jiang, Y., et al., *BRD4 has dual effects on the HMGB1 and NF-kappaB signalling pathways and is a*
611   *potential therapeutic target for osteoarthritis.* Biochim Biophys Acta Mol Basis Dis, 2017. **1863**(12): p.
612   3001-3015.

613   35.   Carbone, A. and S. Rodeo, *Review of current understanding of post-traumatic osteoarthritis resulting*
614   *from sports injuries.* J Orthop Res, 2017. **35**(3): p. 397-405.

36.   Wang, L.J., et al., *Post-traumatic osteoarthritis following ACL injury.* Arthritis Res Ther, 2020. **22**(1):
p. 57.

37.   Hu, Y., et al., *Subchondral bone microenvironment in osteoarthritis and pain.* Bone Res, 2021. **9**(1): p.
20.

38.   Sniekers, Y.H., et al., *A role for subchondral bone changes in the process of osteoarthritis; a micro-CT*
study of two canine models. BMC Musculoskelet Disord, 2008. **9**: p. 20.

39.   Botter, S.M., et al., *Cartilage damage pattern in relation to subchondral plate thickness in a*
collagenase-induced model of osteoarthritis. Osteoarthritis Cartilage, 2008. **16**(4): p. 506-14.

40.   Panula, H.E., et al., *Subchondral bone remodeling increases in early experimental osteoarthrosis in*
young beagle dogs. Acta Orthop Scand, 1998. **69**(6): p. 627-32.

41.   Kwon, M., D. Nam, and J. Kim, *Pathological Characteristics of Monosodium Iodoacetate-Induced*
Osteoarthritis in Rats. Tissue Eng Regen Med, 2023. **20**(3): p. 435-446.

42.   Intema, F., et al., *In early OA, thinning of the subchondral plate is directly related to cartilage damage:*
results from a canine ACLT-meniscectomy model. Osteoarthritis Cartilage, 2010. **18**(5): p. 691-8.

43.   Prasadam, I., et al., *Osteoarthritic cartilage chondrocytes alter subchondral bone osteoblast*
differentiation via MAPK signalling pathway involving ERK1/2. Bone, 2010. **46**(1): p. 226-35.

44.   Hu, W., et al., *Microenvironment in subchondral bone: predominant regulator for the treatment of*
osteoarthritis. Ann Rheum Dis, 2021. **80**(4): p. 413-422.