Concurrent Joint Contact in Anterior Cruciate Ligament Injury induces cartilage

micro-injury and subchondral bone sclerosis, resulting in knee osteoarthritis

Authors

Kei Takahata1,2, Yu-Yang Lin3, Benjamin Osipov3, Kohei Arakawa2,4, Saaya Enomoto5,

Blaine A. Christiansen3*, Takanori Kokubun1,6*

Affiliations

1 Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University,

Koshigaya, Saitama, Japan

2 Japan Society for the Promotion of Science, Tokyo, Japan

3 Department of Orthopaedic Surgery, University of California Davis Health, Sacramento,

CA, United States

4 Laboratory of Animal Regeneration Systemology, Department of Life Sciences, School of

Agriculture, Meiji University, Kanagawa, Japan

5 Saitama Prefecture Rehabilitation Center, Saitama, Japan

6 Department of Physical Therapy, School of Health and Social Services, Saitama Prefectural

University, Koshigaya, Saitama, Japan

Co-Correspondence

[3]Blaine A. Christiansen

University of California Davis Health, Department of Orthopaedic Surgery, Lawrence J.

Ellison Musculoskeletal Research Center, 2700 Stockton Ave, Suite 2301 Sacramento, CA

95817, USA. Tel.: 916-734-3974

bchristiansen@ucdavis.edu

[1,6]Takanori Kokubun, Ph.D.

820 San-nomiya, Koshigaya, Saitama, 343-8540. Japan

Tel: +81 48-973-417617

kokubun-takanori@spu.ac.jp

ABSTRACT

Objective:

Anterior Cruciate Ligament (ACL) injury initiates post-traumatic osteoarthritis (PTOA)

via two distinct processes: initial direct contact injury of the cartilage surface during ACL

injury, and secondary joint instability due to the ACL deficiency. Using the well-established

Compression-induced ACL rupture method (ACL-R) and a novel Non-Compression ACL-R

model, we aimed to reveal the individual effects of cartilage compression and joint instability

on PTOA progression after ACL injury in mice.

Design:

Twelve-week-old C57BL/6J male were randomly divided to three experimental groups:

Compression ACL-R, Non-Compression ACL-R, and Intact. Following ACL injury, we

performed joint laxity testing and microscopic analysis of the articular cartilage surface at 0

days, in vivo optical imaging of matrix-metalloproteinase (MMP) activity at 3 and 7 days, and

histological and microCT analysis at 0, 7, 14, and 28 days.

Results:

The Compression ACL-R group exhibited a significant increase of cartilage roughness

immediately after injury compared with the Non-Compression group. At 7 days, the

Compression group exhibited increased MMP-induced fluorescence intensity and MMP-13

positive cell ratio of chondrocytes. Moreover, histological cartilage degeneration was

observable in the Compression group at the same time point. Sclerosis of tibial subchondral

bone in the Compression group was more significantly developed than in the Non-

Compression group at 28 days.

Conclusions:

Both Compression and Non-Compression ACL injury initiated PTOA progression due to

joint instability. However, joint contact during ACL rupture also caused initial micro-damage

on the cartilage surface and initiated early MMP activity, which could accelerate PTOA

progression compared to ACL injury without concurrent joint contact.

Key Words:

Osteoarthritis, Animal model, ACL injury, Cartilage compression, Subchondral bone, Joint

instability

INTRODUCTION

Osteoarthritis (OA) is a major chronic musculoskeletal disease that progresses irreversibly and negatively influences the quality of life of OA patients. OA can be divided into two types depending on the mechanisms of initiation: primary OA, which is idiopathic and generally occurs during aging, and secondary post-traumatic OA (PTOA), which is initiated by a traumatic joint injury such as a meniscus tear or ligament injury. PTOA accounts for approximately 12% of symptomatic OA patients [1] and has an increasing incidence rate due to the increasing popularity of high-impact sports [2]. Therefore, elucidating the mechanisms of PTOA initiation and progression is necessary to find therapeutic strategies to prevent or slow the progression of OA.

Anterior Cruciate Ligament (ACL) injury is a known initiator of PTOA development. The annual incidence of ACL injury in the general population is 68.6 per 100,000 people [3], and the incidence of PTOA following ACL injury is as high as 87% [4]. In the etiology of OA following ACL injury, concomitant injuries of other joint structures along with ACL rupture and increased mechanical stress caused by joint instability are likely contributing factors. Almost half of ACL injury patients suffer from initial damage of articular cartilage [5], and concomitant cartilage injury may increase OA risk up to 2.4 times at 19 years [6]. In addition, 80-90% of patients with an acute ACL injury also show signs of subchondral bone lesions measured using magnetic resonance imaging. [4] Subchondral bone

changes can affect cartilage degeneration via crosstalk in molecular and biomechanical

interactions. [7].

Because the ACL restricts the anterior-posterior (AP) translation of the tibia relative

to the femur [8], joint instability after ACL injury can induce abnormal mechanical stress,

which may stimulate knee joint tissues to produce catabolic enzymes. Since knee PTOA

develops several decades after ACL injury, it was assumed that secondary joint instability is

the most important factor and surgical reconstruction of the ACL has been widely used to

reduce the risk of PTOA development [9]. However, a recent systematic review reported that

reconstruction surgery may not affect PTOA progression [10]. This indicates that concurrent

joint contact when the ACL is injured may also play a significant role in the pathogenesis of

PTOA. Acute damage to the articular chondral surface and subchondral bone could be the

initiation of PTOA, however, it remains unclear which tissue would be more involved in the

OA onset.

ACL transection (ACL-T) is a common model for inducing knee OA in small

animals, but this method also induces unnecessary acute inflammation in articular tissues due

to invasive surgical procedures [11-14]. In contrast, the Non-Invasive Compression ACL Rupture

(ACL-R) model was developed and used to explore mechanisms of PTOA progression

following ACL injury [15-19]. ACL-R is induced by applying compressive force to the joint,

which causes anterior tibial dislocation and ACL injury. This method largely imitates acute

ACL injury and knee PTOA progression after injury. However, it is difficult to separate the

effects of initial overload of the articular cartilage and the secondary effects of joint

instability. We recently established a novel Non-Compression ACL-R model, which is made

without compression force on the cartilage surface and induces non-articular injuries other

than ACL rupture [20]. Based on the findings of this model, we hypothesized that comparing the

Compression ACL-R model and the novel Non-Compression ACL-R model will allow us to

investigate the individual effects of initial cartilage damage due to compression vs. secondary

instability on PTOA progression.

MATERIALS AND METHODS

Animals and Experimental Design

Twelve-week-old C57BL/6J male mice were randomly divided into three

experimental groups: Compression ACL-R (n = 49), Non-Compression ACL-R (n = 48), and

Intact (n=21). Mice were euthanized at 0, 3, 7, 14, and 28 days after injury, and knee joints

were dissected for analysis (Fig. 1A). In the joint laxity test, microscopic, morphological, and

histological analysis at day 0 after injury, contralateral knee joints from the Non-Compression

group were used as the Intact group. Mice were allowed to acclimate for 1-2 weeks in the

vivarium before the start of the experiment and were cared for in accordance with the

guidelines set by the National Institutes of Health (NIH) on the care and use of laboratory

animals. All procedures were conducted under approved Institutional Animal Care and Use

Committee protocols at Saitama Prefectural University and the University of California

Davis.

Creating Compression ACL-R and Non-Compression ACL-R Models

All procedures were performed on the right knee joint of each anesthetized mouse

under 1-4% inhaled isoflurane. The Compression ACL-R model was created via tibial

compression overload as previously described [17, 21]. Right knee joints were positioned in an

electromagnetic materials machine (ElectroForce 3200, TA Instruments, New Castle, DE),

and a single tibial compressive overload was applied at 1 mm/s to induce ACL rupture. The

Non-Compression ACL-R model was created based on our previous study [22]. The knee joint

was fixed at 90 degrees using surgical tape on a stand, and a force was slowly applied by the

thumb tip of the operator along the long axis of the femur. The applied force was stopped

quickly after hearing a distinct popping sound, which indicated ACL rupture. For both

models, buprenorphine analgesia (0.1 mg/kg) was injected immediately following injury for

pain relief.

Anterior-Posterior Joint Laxity Testing

Knee joints were collected immediately post-injury, and femurs and tibias were

embedded in brass tubes using polymethylmethacrylate as previously described [21, 23]. Tibias

were fixed to the load cell; then femurs were secured so that the angle of the knee joint was

either 60˚ or 90˚. Five anterior-posterior (AP) loading cycles were applied perpendicular to the

longitudinal axis of the tibia to a target force of ±1.5 N at a loading rate of 0.5 mm/s. The

degree of AP joint laxity was quantified based on the difference between displacement at +0.8

N and -0.8 N.

Microscopic Analysis of Cartilage Surface Roughness

For samples collected 0 days following injury, soft tissue was removed carefully to

expose the tibial plateau, then 3D optical profilograph images of the whole tibial plateau were

taken (VR-6000; Keyence, JPN). The region of interest (ROI) was set as an ellipse with 0.6-

1.2 mm diameters in the center area of the medial and lateral tibial plateau (Figure 2A). The

arithmetical mean height and maximum height of the cartilage surface were measured by

normalizing to the contralateral intact knee of each mouse. Methods are further described in

the Supplementary Materials.

Fluorescent Reflectance Imaging (FRI)

At 3 and 7 days post-injury, mice were imaged in vivo using an optical imaging

system (IVIS Spectrum, PerkinElmer, Waltham, MA), after IV administration of a near-

infrared probe that is activated by matrix-metalloproteinases (MMPSense 680, PerkinElmer,

Waltham, MA). The ROI was set as a circle of 0.7 mm^2 that surrounded the tibial tuberosity

to the superior border of the patella, and image processing and quantification were performed

via IVIS Living Image software as previously described [24]. To unify the mouse-to-mouse

variation in the delivery of the fluorescent probe, the radiant efficiency of the ACL-R knee

was normalized to the contralateral intact knee of each mouse. Methods are further described

in the Supplementary Materials.

Micro-Computed Tomography Analysis of Osteophyte Formation and Epiphyseal Bone

Microstructure

For analysis of osteophyte formation at 7, 14, and 28 days after injury, ROI

contouring included all heterotopic mineralized tissue around the joint, as well as the patella,

fabellae, and menisci. The average bone volume in the Intact group was used as the

"baseline" volume to calculate osteophyte formation in the Compression and Non-

Compression groups.

We also assessed the morphological changes in epiphyseal trabecular bone at 0, 7,

14, and 28 days after injury. The ROI was designed as the femoral and tibial trabecular bone

enclosed by the growth plate and subchondral cortical bone plate (Figure 3B). Additionally,

whole subchondral bone including the cortical bone plate and trabecular bone in the medial

and lateral tibial compartments were also measured (Figure 4A). We quantified apparent bone

mineral density (BMD, g/cm^3), bone volume per total volume (BV/TV, %), trabecular

number (Tb.N, 1/mm), trabecular thickness (Tb.Th, mm), and trabecular separation (Tb.Sp,

mm).

Histological Analysis for Cartilage Degeneration

Safranin-O/fast green staining was performed to evaluate articular cartilage

degeneration in the medial and lateral tibial compartments using the Osteoarthritis Research

Society International (OARSI) histopathological grading system [25]. We initially assessed the

whole joint pathology in the knee joint, then assessed the medial and lateral tibial plateau

separately.

Immunohistochemical Analysis

Immunohistochemical (IHC) staining was performed using anti-MMP-13 (1:200, bs-

0575R, Bioss). Detailed protocols are described in the Supplementary Materials. We

calculated the ratio between the number of MMP-13 positive cells and the number of

chondrocytes in the anterior and posterior area of the articular cartilage with regions of

interest of 40,000 μm^2 (200 $\mu m \times$ 200 μm). As with the OARSI score, we initially evaluated

the whole joint and then subsequently assessed the medial and lateral tibial plateau separately.

Statistical Analysis

Statistical analysis was performed using RStudio. Student's t-test was used for the

osteophyte formation and Wilcoxon rank sum was used for the microscope and FRI data.

One-way analysis of variance was performed for the joint laxity test and MMP-13 IHC

analysis initially, and then the Tukey-Kramer test was used for post-hoc analysis. The

Kruskal-Wallis test was used to compare subchondral bone μCT data and OARSI scores, and

the Steel-Dwass method was used for the subsequent multiple comparisons. Parametric data

are expressed as the mean ± 95% confidence intervals (95% CI); non-parametric data are

expressed as the median ± interquartile ranges. Statistical significance was set at $p < 0.05$.

RESULTS

RESULTS

AP joint Laxity Test of the Knee

Compared with the Intact group, both the Compression and Non-Compression ACL-

R groups had significantly increased AP joint laxity at both 60°and 90°(Figure 1B).

However, no significant differences were observed between the Compression and Non-

Compression groups at either angle. This suggests that any differences in OA progression

between the ACL-R groups can be attributed to the initial effect of compressive force at the

time of ACL injury.

Figure 1. (A) Experimental design. We made the Compression ACL-R, Non-

Compression ACL-R, and Intact groups. Mice were euthanized at 0, 3, 7, 14, and 28 days

after injury and following analyses were performed; joint laxity test, microscopic, microCT,

and histological analysis. At day 0 after injury, contralateral knee joints from the Non-

Compression group were used as the Intact group. (B) A result of joint laxity test. The

Compression and Non-Compression ACL-R groups significantly increased Anterior-Posterior

joint laxity at both 60° and 90°. However, no significant differences were observed between

both injury groups. Data are presented as the mean ± 95% CI. *P< 0.05; ***P< 0.001.

Microscope Analysis of Cartilage Roughness in the Tibial Plateau

228   Surface roughness in the region not covered by menisci was increased in the

Compression ACL-R group, especially on the medial tibial plateau. The normalized

arithmetical mean height of the medial compartment in the Compression group was

significantly higher than that in the Non-Compression group, but no difference was observed

in the lateral compartment (Figure 2A). In addition, there were no significant differences in

the maximum height of the medial and lateral compartments between groups. These results

indicate that compressive force during compression ACL injury caused initial micro-damage

in the surface layer of articular cartilage.

Quantification of MMP Fluorescence Intensity

238   At 3 days following injury, MMP-induced fluorescence intensity around ACL-R

239   knees increased slightly in the Compression (Normalized data: 0.31 ± 0.659) and Non-

240   Compression groups (0.512 ± 0.472) compared with each contralateral intact knee, and no

significant difference was observed between groups (Figure 2B). Whereas, at 7 days,

fluorescence intensity increased moderately in the Compression (0.684 ± 0.597) and slightly

in the Non-Compression (0.17 ± 0.368) groups. However, there were no significant

differences between these groups.

Figure 2. (A) Microscope Analysis of Cartilage Roughness in the Tibial Plateau at

day 0 after injury. The ROI was set as an ellipse with 0.6-1.2 mm diameters in the center area

of the medial and lateral tibial plateau. Compression group significantly increased the

normalized arithmetical mean height in the medial compartment compared with the Non-

Compression group. (B) Quantification of MMP Fluorescence Intensity. Normalized MMP-

induced fluorescence intensity around ACL-R knees increased in the Compression, especially

at 7 days following injury. However, there were no significant differences between these

groups. Data are presented as the median ± interquartile range. *P< 0.05.

MicroCT Measurement of Osteophyte Formation

Mineralized osteophyte volume was quantified at 14 and 28 days because no

osteophytes were observed at 7 days following injury. Similarly, only a small amount of

mineralized osteophyte formation was observed in both ACL-R groups at 14 days (Figure

3A). At 28 days following injury, a greater volume of ectopic mineralized tissue was

observed in the medial femoral condyle and tibial plateau of injured joints. However, there

were no significant differences between ACL-R groups.

MicroCT Analysis of Epiphysis Trabecular Bone of the Distal Femur and Proximal

Tibia

At 0 days post-injury, no micro injuries or fractures were observed in the femoral and

tibial epiphysis trabecular bone in either ACL-R group, and there were no significant

differences in the microstructure between any experimental groups (Supplementary Figure

2A-B). This indicates that compressive force while inducing ACL rupture didn't cause

concomitant structural failures in the subchondral trabecular bone. Femoral epiphysis

trabecular bone showed a significant decrease of BMD, BV/TV, and Tb.Th in the

Compression and Non-Compression groups compared with the Intact group at 7 days, which

suggests acute bone absorption at an early time point in both ACL-R groups (Figure 3C).

These parameters were still significantly lower in the Compression and Non-Compression

groups at 14 days. However, the Compression group recovered bone volume moderately at 28

days, and no significant difference in the BMD was observed between the Intact group and

the Compression group at this time point, although the Non-Compression group was still

significantly lower than the Intact group. Tb.N and Tb.Sp results are described in the

Supplementary Material (Supplementary Figure 3A).

280   Similarly, tibial epiphysis trabecular bone also exhibited a significant decrease in

BMD and BV/TV in the Compression and Non-Compression groups at 7 days following

injury (Figure 3D). At 14 days, the significantly decreased BMD and BV/TV was still present

in the Non-Compression group, whereas the Compression group slightly increased bone

volume and there was no significant difference in BV/TV between the Intact and the

Compression groups. Furthermore, in the Compression group at 28 days, the bone volume

increased to the same level as in the Intact group. The Non-Compression group still showed

deficits in bone microstructure at this time point, therefore the BMD and BV/TV in the Intact

and Compression group were significantly greater than in the Non-Compression group. Tb.N

and Tb.Sp results are described in the Supplementary Material (Supplementary Figure 3B).

Figure 3. (A) Reconstructed images obtained by microCT and measurement of

osteophyte formation. Only a small amount of mineralized osteophyte at 14 days and a greater

volume of ectopic mineralized tissue in the medial femoral condyle and tibial plateau of

injured joints at 28 days were observed in the Compression and Non-Compression groups.

However, there were no significant differences. Black arrows show the osteophyte. (B) The

ROI to measure the tibial and femoral epiphysis trabecular bone and (C) the result of

microCT Analysis. Femoral epiphysis trabecular bone significantly decreased BMD, BV/TV,

and Tb.Th in the Compression and Non-Compression groups compared with the Intact group

at 7 days. At 28 days, no significant difference in the BMD was observed between the Intact

and the Compression groups. However, the Non-Compression group was still significantly

lower than the Intact group at the same timepoint. Tibial epiphysis trabecular bone also

showed a significant decrease in BMD and BV/TV in both injury groups at 7 days. However,

at 28 days, the BMD and BV/TV in the Intact and Compression groups were significantly

greater than in the Non-Compression group. Data are presented as the median ± interquartile

range. *$P < 0.05$. **$P < 0.01$.

MicroCT Analysis of Subchondral Bone in the Medial and Lateral Tibial Compartment

Analysis of subchondral bone microstructure in the medial and lateral tibial

compartments at 7, 14, and 28 days (Figure 4A) showed that in the medial compartment, the

Compression and the Non-Compression groups caused bone loss with a significant decrease

in BV/TV and Tb.Th compared to the Intact group at 7 days (Figure 4B). Although BV/TV

and Tb.Th in the Non-Compression group were still significantly lower than in the Intact

group at 14 days, the Compression group increased BV/TV and Tb.Th and no differences

were observed compared to other groups. Additionally, the Compression group exhibited

significantly increased BV/TV compared to the Non-Compression group at 28 days.

In the lateral compartment, both ACL-R groups exhibited significant bone loss

compared with the Intact group at 7 days similar to what was observed in the medial

compartment (Figure 4C). At 14 days, the Compression and Non-Compression groups

increased BV/TV and Tb.Th, and no differences were observed between any experimental

groups. Unlike the medial compartment, the Compression group exhibited a significant

increase of BV/TV and Tb.Th compared to the Intact group at 28 days. Tb.N and Tb.Sp

results for both compartments are described in the Supplementary Materials (Supplementary

Figure 4A-B).

Figure 4. (A) The ROI of medial and lateral tibial trabecular bone and microCT

images. (B) Analysis of subchondral bone microstructure in the medial and lateral tibial

compartments. In the medial compartment, the Compression and the Non-Compression

groups caused bone loss with a significant decrease in BV/TV and Tb.Th compared to the

Intact group at 7 days. At 28 days, the Compression group showed significantly increased

BV/TV compared to the Non-Compression group. In the lateral compartment, both ACL-R

groups exhibited significant bone loss compared with the Intact group at 7 days. Unlike the

medial compartment, the Compression group exhibited a significant increase of BV/TV and

Tb.Th compared to the Intact group at 28 days. Data are presented as the median ±

interquartile range. *P< 0.05. **P< 0.01.

Histological Analysis of Cartilage Degeneration in the Medial and Lateral Tibial

Compartment

Histological analysis did not detect any acute injuries of articular tissues other than

the ACL (cartilage, menisci, subchondral bone) in the knee joint at 0 days following injury

(Supplementary Figure 5). In the medial tibial plateau at 7 days, over half of the mice in the

Compression group and some mice in the Non-Compression group developed a small loss of

cartilage around the posterior edge, and cell proliferation and staining of Safranin-O were

observed in the posterior region (Figure 5A). At 14 days, fibrillation of the cartilage surfaces,

mild to moderate cartilage erosion, and osteophytes on the posterior tibia were observed in

both ACL-R groups, especially in the posterior region. At 28 days, erosion extending to the

center and posterior region of the subchondral bone and growth plate were observed in both

ACL-R groups. These histological changes were observed in the lateral tibial plateau as well;

however, the degree of OA progression was milder compared to the medial tibial plateau at all

time points (Figure 5B).

The OARSI score for the whole joint pathology in the Compression group increased

significantly compared with the Intact group at 7 days (Figure 5C). Moreover, OARSI score

in both ACL-R groups increased significantly compared with the Intact group at 14 days and

28 days. For the medial tibial plateau, OARSI score in both ACL-R groups were significantly

higher than that in the Intact group at all time points (Figure 5D). For the lateral tibial plateau,

OARSI score in both ACL-R groups were significantly higher than that in the Intact group at

14 days and 28 days (Figure 5E).

[Insert Figure 5 here] scale bar 500μm

Figure 5. (A) Histological images of cartilage degeneration in the medial and lateral

tibial compartment. (B) The OARSI score for the whole joint pathology in the Compression

group was significantly higher than that in the Intact group at 7 days. At 14 and 28 days, both

ACL-R groups significantly increased the OARSI score compared with the Intact group. (C)

In the medial compartments, OARSI score in both ACL-R groups were significantly higher

than that in the Intact group at all time points. (D) In the lateral compartment, both ACL-R

groups caused a significant increase of OARSI score compared with the Intact group at 14

and 28 days. Data are presented as the median ± interquartile range. *$P< 0.05$. **$P< 0.01$.

Black scale bar, 500 µm.

Immunohistochemical Analysis of the Articular Cartilage

IHC analysis of MMP-13 was performed at 7 and 14 days following injury because

cartilage was highly degenerated by 28 days (Figure 6A-B). At 7 days, the number of MMP-

13-positive cells of the whole joint in the Compression group increased significantly

compared with the Intact and Non-Compression groups (Figure 6C). Moreover, the Non-

Compression group also significantly increased the positive cell ratio compared to the Intact

group. Analysis of the medial tibial plateau showed similar results to the whole joint (Figure

6D), whereas for the lateral tibial plateau, the number of positive cells in the Compression

group increased significantly compared with the other groups (Figure. 6E). At 14 days, the

positive cells ratio of the whole joint in both ACL-R groups increased significantly compared

with the Intact group. Similar results were observed in the medial and lateral tibial plateau.

Figure 6. (A-B) Immunohistochemical images of MMP-13 in cartilage in the medial

and lateral tibial plateau. (C-D) In the whole joint level and medial tibial plateau,

Compression group increased the MMP-13 positive cell rate significantly compared with the

Intact and Non-Compression groups at 7 days. Furthermore, the Non-Compression group also

significantly increased the positive cell ratio compared to the Intact group. At 14 days, the

positive cells ratio in both ACL-R groups increased significantly compared with the Intact

group. (E) In the lateral tibial plateau at 7 days, the number of MMP-13 positive cells in the

Compression group increased significantly compared with the other groups. At 14 days, both

ACL-R groups increased the MMP-13 positive cell rate significantly compared with the Intact

group. Data are presented as the mean ± 95% CI. **$P< 0.01$; ***$P< 0.001$. Black scale bar,

100 μm. (F) Schematic summarizing the mechanism of OA development following ACL

injury with or without compression concomitant compression force. Compression-ACL

rupture in mice caused initial micro-damage on the cartilage surface and early cartilage

degeneration with MMP-13 production, leading to later bone formation in the subchondral

bone.

DISCUSSION

The mechanism of OA development following ACL injury involves two distinct

processes: initial direct injury of the cartilage surface when the ACL is injured, and secondary

joint instability due to the deficiency of the ACL. In this study, we compared the conventional

Compression-induced ACL-R and Non-Compression ACL-R models, to clarify the individual

effects of the initial injury response and secondary joint instability on OA progression after

ACL injury. Although no significant differences in joint laxity were observed between ACL-

R groups, the roughness of the cartilage after injury was increased significantly in the

Compression ACL-R group but not the Non-Compression group. Similarly, MMP-induced

fluorescence intensity at 7 days was higher in the Compression group than in the Non-

Compression group. Moreover, the MMP-13 positive cell ratio in the Compression group was

also increased significantly compared to the Non-Compression group at 7 days. Furthermore,

sclerosis of tibial SCB in the Compression group developed significantly more than in the

Non-Compression group at 28 days.

The ACL contributes to joint stability and ACL deficiency results in an increase in

mechanical stress [26]. Our assessment of AP joint laxity immediately after injury revealed that

the Compression and Non-Compression groups had significantly increased joint laxity

compared to the Intact group. However, no differences were observed between the ACL-R

groups (Figure 1B). This result indicates that any differences in OA progression between the

ACL-R groups can be attributed to the direct effect of compressive stress experienced while

inducing ACL rupture.

Interestingly, microscopic analysis demonstrated that roughness on the surface of the

medial tibial plateau in the Compression group was significantly increased compared to the

Non-Compression group. However, no injuries such as cartilage erosion, meniscus injury, or

subchondral bone loss were observed histologically and morphologically immediately

following injury. A previous study using an intra-articular tibial plateau fracture mouse model

reported that compression force to the tibial plateau to a target load of 55 N at a rate of 60 N/s

caused massive soft tissue injuries, including to the meniscus, tendon, and cartilage [27]. The

mean compressive force applied with the Compression ACL-R model was approximately 10

N at a loading rate of 1 mm/s [17]. The compressive force in the joint during Compression

ACL-R was remarkably lower than in the previous study using the tibial plateau fracture

model, therefore roughness on the cartilage surface indicated mild articular damage in the

Compression group rather than severe damage.

In addition to the initial micro-injury, we also measured the biological response in

the early stage using FRI with an MMP-activatable probe. MMP-3, MMP-9, and MMP-13

levels increased in animal models of osteoarthritis [28-32], and MMP-2, MMP-3, MMP-9, and

MMP-13 increased in in vitro models of osteoarthritis [33], which can degrade collagens and

proteoglycans of the articular cartilage. Compressive loading during ACL injury increases the

production of matrix-degrading enzymes and inflammatory cytokines in chondrocytes, which

results in chondrocyte apoptosis and cartilage degeneration [5, 34, 35]. A previous animal study

reported that even a single compressive load at 6 N without ACL rupture caused cartilage

degeneration with chondrocyte apoptosis [19]. However, in the current study no significant

differences were observed between ACL-R groups; the Compression group showed a high

MMP-induced fluorescence intensity using FRI at 7 days following injury. Furthermore, IHC

analysis demonstrated that MMP-13 expression in chondrocytes of the Compression group

increased significantly compared to the Non-Compression group. These molecular changes

may be due to the initial micro-injury detected as surface roughness, which is consistent with

early stage pathology of ACL injury in human patients. Interestingly, histological data also

showed that the OARSI score in the whole joint of the Compression group significantly

increased compared to the Intact group at 7 days. These results suggested that the

compressive force on the cartilage surface while inducing ACL rupture may have accelerated

cartilage degeneration in the early stage through the activation of MMPs. Alternatively,

considering that there was no difference in OARSI score at 14 and 28 days, it is also possible

that secondary joint instability is a more important factor in the progression of cartilage

degeneration after the initial stage.

Subchondral bone provides mechanical and nutritional support for cartilage, and

microenvironmental changes in subchondral bone might affect cartilage metabolism directly

or indirectly [36]. Generally, subchondral bone reacts to mechanical stress and is faster to

remodel than articular cartilage due to its robust innervation and blood supply that provide it

with a high capacity of turnover [37, 38]. We hypothesized that subchondral bone would react to

compressive force and induced structural changes during the early stage. Indeed, we observed

loss of subchondral bone volume in both ACL-R groups following injury. However, the

Compression group induced subchondral bone sclerosis in the medial tibial plateau compared

to the Non-Compression group at 28 days, and the lesion area was consistent with the

cartilage degeneration. Some reports have shown that cartilage degeneration precedes

subchondral bone changes [39-41] and osteoarthritic chondrocytes enhanced osteoblast

differentiation in subchondral bone via the ERK1/2 pathway [42]. Our results revealed that ACL

injury with concomitant compressive force initially caused micro-injury on the cartilage

surface, then subchondral bone remodeling and sclerosis in the medial compartment at an

earlier time point than ACL injury without compressive loading (Figure 6F). This earlier

switch from bone loss to bone sclerosis is consistent with an accelerated PTOA progression in

the Compression ACL-R group compared to the Non-Compression group.

472   There are two main limitations to this study. First, articular surface roughness

analysis was only performed immediately after the induction of each model. Microscopic

evaluation can determine the degree of cartilage degeneration as a surface rather than a line in

more detail. Roughness analysis was able to detect micro-injury on the cartilage surface at 0

days (Figure 2A), which was not possible to see with histological observation (Supplementary

Figure 5). Further roughness analysis at additional time points may enable us to reveal the

reaction of cartilage degeneration to compression force, especially in the early phase. Second,

we didn't investigate the molecular biological mechanism of subchondral bone remodeling

and sclerosis during PTOA progression. It has been shown that the proliferation and

differentiation of osteoprogenitors are stimulated by platelet-derived growth factor-A,

transforming gene factor-β1, and fibroblast growth factor-1, resulting in subchondral bone

formation [43]. We revealed that chondrocytes reacted to initial compression force and

increased MMP-13; therefore, additional experiments about the biological mechanism should

be performed to develop further understanding of these mechanisms.

In conclusion, concomitant joint contact while non-invasively inducing ACL rupture

in mice caused initial micro-damage on the cartilage surface and early cartilage degeneration

with MMP-13 production, leading to later bone formation in the subchondral bone.

Understanding the initial pathology of the ACL injury may be an important indicator of

disease etiology and represent a potential preventative approach for mitigating secondary OA

development.

493   Acknowledgments

The author(s) received no financial support for the research, authorship, and/or publication of

this article. We would like to thank Editage (www.editage.com) for English language editing.

The study design and summary scheme were created with BioRender.com. Research reported

in this publication was partially supported by the National Institute of Arthritis and

Musculoskeletal and Skin Diseases, part of the National Institutes of Health, under Award

Number R01 AR075013.

501   Author contributions

All authors approved the final submitted manuscript.

Study design: TK and BC

Making model and Data collection: KT and YL

Mechanical analysis: KT and BO

Fluorescent reflectance imaging: KT and YL

Morphological analysis: KT, YL, and KA

Histological analysis: KT, KA, and SE

Manuscript composition: KT, TK, and BC

Role of funding source

Competing interest statement

514   All authors have no conflicts of interest related to the manuscript.

References

1. Thomas AC, Hubbard-Turner T, Wikstrom EA, Palmieri-Smith RM. Epidemiology of
Posttraumatic Osteoarthritis. J Athl Train 2017;52:491-6.
2. Lohmander LS, Ostenberg A, Englund M, Roos H. High prevalence of knee osteoarthritis,
pain, and functional limitations in female soccer players twelve years after anterior cruciate
ligament injury. Arthritis Rheum 2004;50:3145-52.
3. Sanders TL, Maradit Kremers H, Bryan AJ, Larson DR, Dahm DL, Levy BA, *et al.* Incidence of
Anterior Cruciate Ligament Tears and Reconstruction: A 21-Year Population-Based Study.
Am J Sports Med 2016;44:1502-7.
4. Friel NA, Chu CR. The role of ACL injury in the development of posttraumatic knee
osteoarthritis. Clin Sports Med 2013;32:1-12.

5. Riordan EA, Little C, Hunter D. Pathogenesis of post-traumatic OA with a view to intervention.
Best Pract Res Clin Rheumatol 2014;28:17-30.

6. Whittaker JL, Losciale JM, Juhl CB, Thorlund JB, Lundberg M, Truong LK, *et al.* Risk factors
for knee osteoarthritis after traumatic knee injury: a systematic review and meta-analysis of
randomised controlled trials and cohort studies for the OPTIKNEE Consensus. Br J Sports
Med 2022;56:1406-21.

7. Yuan XL, Meng HY, Wang YC, Peng J, Guo QY, Wang AY, *et al.* Bone-cartilage interface
crosstalk in osteoarthritis: potential pathways and future therapeutic strategies. Osteoarthritis
Cartilage 2014;22:1077-89.

8. Ajuied A, Wong F, Smith C, Norris M, Earnshaw P, Back D, *et al.* Anterior cruciate ligament
injury and radiologic progression of knee osteoarthritis: a systematic review and meta-
analysis. Am J Sports Med 2014;42:2242-52.

9. Thorstensson CA, Lohmander LS, Frobell RB, Roos EM, Gooberman-Hill R. Choosing
surgery: patients' preferences within a trial of treatments for anterior cruciate ligament injury. A
qualitative study. BMC Musculoskelet Disord 2009;10:100.

10. Ferrero S, Louvois M, Barnetche T, Breuil V, Roux C. Impact of anterior cruciate ligament
surgery on the development of knee osteoarthritis: A systematic literature review and meta-
analysis comparing non-surgical and surgical treatments. Osteoarthr Cartil Open
2023;5:100366.

11. Goldring MB, Otero M. Inflammation in osteoarthritis. Current Opinion in Rheumatology
2011;23:471-8.

12. Liu-Bryan R. Synovium and the Innate Inflammatory Network in Osteoarthritis Progression.
Current Rheumatology Reports 2013;15.

13. Mathiessen A, Conaghan PG. Synovitis in osteoarthritis: current understanding with
therapeutic implications. Arthritis Research & Therapy 2017;19.

14. Scanzello CR, Goldring SR. The role of synovitis in osteoarthritis pathogenesis. Bone
2012;51:249-57.

15. Ramme AJ, Lendhey M, Raya JG, Kirsch T, Kennedy OD. A novel rat model for subchondral
microdamage in acute knee injury: a potential mechanism in post-traumatic osteoarthritis.
Osteoarthritis Cartilage 2016;24:1776-85.

16. Brown SB, Hornyak JA, Jungels RR, Shah YY, Yarmola EG, Allen KD, *et al.* Characterization
of Post-Traumatic Osteoarthritis in Rats Following Anterior Cruciate Ligament Rupture by
Non-Invasive Knee Injury (NIKI). J Orthop Res 2020;38:356-67.

17. Christiansen BA, Anderson MJ, Lee CA, Williams JC, Yik JH, Haudenschild DR.
Musculoskeletal changes following non-invasive knee injury using a novel mouse model of
post-traumatic osteoarthritis. Osteoarthritis Cartilage 2012;20:773-82.

18. Maerz T, Newton MD, Kurdziel MD, Altman P, Anderson K, Matthew HW, *et al.* Articular
cartilage degeneration following anterior cruciate ligament injury: a comparison of surgical

transection and noninvasive rupture as preclinical models of post-traumatic osteoarthritis.

Osteoarthritis Cartilage 2016;24:1918-27.

19.   Wu P, Holguin N, Silva MJ, Fu M, Liao W, Sandell LJ. Early response of mouse joint tissue to

noninvasive knee injury suggests treatment targets. Arthritis Rheumatol 2014;66:1256-65.

20.   Takahata K, Arakawa K, Enomoto S, Usami Y, Nogi K, Saitou R*, et al.* Joint instability causes

catabolic enzyme production in chondrocytes prior to synovial cells in novel non-invasive ACL

ruptured mouse model. Osteoarthritis Cartilage 2023;31:576-87.

21.   Hsia AW, Anderson MJ, Heffner MA, Lagmay EP, Zavodovskaya R, Christiansen BA.

Osteophyte formation after ACL rupture in mice is associated with joint restabilization and loss

of range of motion. J Orthop Res 2017;35:466-73.

22.   Takahata K, Arakawa K, Enomoto S, Usami Y, Nogi K, Saitou R*, et al.* Joint instability causes

catabolic enzyme production in chondrocytes prior to synovial cells in novel non-invasive ACL

ruptured mouse model. Osteoarthritis Cartilage 2022.

23.   Lockwood KA, Chu BT, Anderson MJ, Haudenschild DR, Christiansen BA. Comparison of

loading rate-dependent injury modes in a murine model of post-traumatic osteoarthritis. J

Orthop Res 2014;32:79-88.

24.   Satkunananthan PB, Anderson MJ, De Jesus NM, Haudenschild DR, Ripplinger CM,

Christiansen BA. In vivo fluorescence reflectance imaging of protease activity in a mouse

model of post-traumatic osteoarthritis. Osteoarthritis Cartilage 2014;22:1461-9.

25.   Glasson SS, Chambers MG, Van Den Berg WB, Little CB. The OARSI histopathology initiative

- recommendations for histological assessments of osteoarthritis in the mouse. Osteoarthritis

Cartilage 2010;18 Suppl 3:S17-23.

26.   Kraeutler MJ, Wolsky RM, Vidal AF, Bravman JT. Anatomy and Biomechanics of the Native

and Reconstructed Anterior Cruciate Ligament: Surgical Implications. Journal of Bone and

Joint Surgery 2017;99:438-45.

27.   Stiffel V, Rundle CH, Sheng MH, Das S, Lau KW. A Mouse Noninvasive Intraarticular Tibial

Plateau Compression Loading-Induced Injury Model of Posttraumatic Osteoarthritis. Calcif

Tissue Int 2020;106:158-71.

28.   Hu H, Yang B, Li Y, Zhang S, Li Z. Blocking of the P2X7 receptor inhibits the activation of the

MMP-13 and NF-kappaB pathways in the cartilage tissue of rats with osteoarthritis. Int J Mol

Med 2016;38:1922-32.

29.   Raghu H, Lepus CM, Wang Q, Wong HH, Lingampalli N, Oliviero F*, et al.* CCL2/CCR2, but

not CCL5/CCR5, mediates monocyte recruitment, inflammation and cartilage destruction in

osteoarthritis. Ann Rheum Dis 2017;76:914-22.

30.   Qu R, Chen X, Wang W, Qiu C, Ban M, Guo L*, et al.* Ghrelin protects against osteoarthritis

through interplay with Akt and NF-kappaB signaling pathways. FASEB J 2018;32:1044-58.

31.   Schmidli MR, Fuhrer B, Kurt N, Senn D, Drogemuller M, Rytz U*, et al.* Inflammatory pattern of

the infrapatellar fat pad in dogs with canine cruciate ligament disease. BMC Vet Res

2018;14:161.

32.   Li H, Xie S, Qi Y, Li H, Zhang R, Lian Y. TNF-alpha increases the expression of inflammatory
factors in synovial fibroblasts by inhibiting the PI3K/AKT pathway in a rat model of
monosodium iodoacetate-induced osteoarthritis. Exp Ther Med 2018;16:4737-44.
33.   Jiang Y, Zhu L, Zhang T, Lu H, Wang C, Xue B, *et al.* BRD4 has dual effects on the HMGB1
and NF-kappaB signalling pathways and is a potential therapeutic target for osteoarthritis.
Biochim Biophys Acta Mol Basis Dis 2017;1863:3001-15.
34.   Carbone A, Rodeo S. Review of current understanding of post-traumatic osteoarthritis
resulting from sports injuries. J Orthop Res 2017;35:397-405.
35.   Wang LJ, Zeng N, Yan ZP, Li JT, Ni GX. Post-traumatic osteoarthritis following ACL injury.
Arthritis Res Ther 2020;22:57.
36.   Hu Y, Chen X, Wang S, Jing Y, Su J. Subchondral bone microenvironment in osteoarthritis
and pain. Bone Res 2021;9:20.
37.   Sniekers YH, Intema F, Lafeber FP, van Osch GJ, van Leeuwen JP, Weinans H, *et al.* A role
for subchondral bone changes in the process of osteoarthritis; a micro-CT study of two canine
models. BMC Musculoskelet Disord 2008;9:20.
38.   Botter SM, van Osch GJ, Waarsing JH, van der Linden JC, Verhaar JA, Pols HA, *et al.*
Cartilage damage pattern in relation to subchondral plate thickness in a collagenase-induced
model of osteoarthritis. Osteoarthritis Cartilage 2008;16:506-14.
39.   Panula HE, Nieminen J, Parkkinen JJ, Arnala I, Kroger H, Alhava E. Subchondral bone
remodeling increases in early experimental osteoarthrosis in young beagle dogs. Acta Orthop
Scand 1998;69:627-32.
40.   Kwon M, Nam D, Kim J. Pathological Characteristics of Monosodium Iodoacetate-Induced
Osteoarthritis in Rats. Tissue Eng Regen Med 2023;20:435-46.
41.   Intema F, Hazewinkel HA, Gouwens D, Bijlsma JW, Weinans H, Lafeber FP, *et al.* In early
OA, thinning of the subchondral plate is directly related to cartilage damage: results from a
canine ACLT-meniscectomy model. Osteoarthritis Cartilage 2010;18:691-8.
42.   Prasadam I, Friis T, Shi W, van Gennip S, Crawford R, Xiao Y. Osteoarthritic cartilage
chondrocytes alter subchondral bone osteoblast differentiation via MAPK signalling pathway
involving ERK1/2. Bone 2010;46:226-35.
43.   Hu W, Chen Y, Dou C, Dong S. Microenvironment in subchondral bone: predominant
regulator for the treatment of osteoarthritis. Ann Rheum Dis 2021;80:413-22.
635