1 **Plasmodesmal closure elicits stress responses**
2

3 Author names and affiliations: Estee E. Tee[1,†], Andrew Breakspear[1,†], Diana Papp[1,2,†],
4 Hannah R. Thomas[1,3], Catherine Walker[1], Annalisa Bellandi[1,4], Christine Faulkner[1]
5
[1]Cell and Developmental Biology, John Innes Centre, Norwich Research Park,
Norwich, NR4 7UH, UK
[2]Present address: Department of Metabolism, Digestion and Reproduction, Imperial
College London, London, W12 0NN, UK
[3]Present address: Department of Horticulture, Zijingang Campus, Zhejiang University,
Hangzhou, China
[4]Present address: Laboratoire de Reproduction et Développement des Plantes, École
Normale Supérieure de Lyon, 69342 Lyon, France
[†]These authors contributed equally to this work
Corresponding author: Christine Faulkner

16
17 **Abstract**
18

Plant cells are connected to their neighbors via plasmodesmata facilitating the
exchange of nutrients and signaling molecules. During immune responses,
plasmodesmata close, but how this contributes towards a full immune response is
unknown. To investigate this, we developed two transgenic lines with which we could
induce plasmodesmal closure independently of immune elicitors, using the over-active
CALLOSE SYNTHASE3 allele *icals3m* and the C-terminus of PDLP1 to drive callose
deposition at plasmodesmata. Induction of plasmodesmal closure increased the
expression of stress responsive genes, salicylic acid accumulation and resistance to
Pseudomonas syringae DC3000. More homogeneous plasmodesmal closure using
icals3m also led to the accumulation of starch and sugars, decreased leaf growth, as
well as hypersusceptibility to *Botrytis cinerea*. Based on the profile of responses, we
conclude that plasmodesmal closure itself activates stress signaling, raising questions
of what signals mediate this response and whether these responses occur in all
circumstances when plasmodesmata close.

33
34 Keywords: plasmodesmata / defense / callose / stress signaling
35

Introduction

Plants are multicellular organisms, with almost every plant cell cytoplasmically connected to its neighbors. While most plant cells are equipped with machinery to respond autonomously to a range of stress signals, optimal responses involve the regulation of this cytoplasmic connectivity between cells. This is particularly evident in immune signaling; most cells produce the receptors that perceive microbial threats and can activate appropriate responses. This response can include intracellular, as well as non-cell autonomous regulation, where the degree of connectivity to neighboring cells is a critical part of the full immune response. If plants cannot regulate the cytoplasmic connectivity between cells, they are more susceptible to infection by a range of pathogenic microbes. However, how cell-to-cell connectivity contributes to an overall immune response is an outstanding question.

Cytoplasmic connectivity between plant cells is established via membrane-lined bridges called plasmodesmata. Plasmodesmata are dynamic structures and the oscillation in their aperture, i.e., closing and opening, allows control over the flux of soluble molecules between cells. We broadly assume that many molecules that move through plasmodesmata, including plant hormones, mRNAs, and proteins, are information carriers and that aperture changes thus impact cell-to-cell communication. Indeed, if an immune response involves the production of soluble defense-associated molecules, their passage through plasmodesmata could transmit critical information to neighboring (or distant) naïve cells.

Plant immune responses include an upregulation of callose synthesis at plasmodesmata leading to a decrease in plasmodesmal aperture, which reduces cell-to-cell movement of molecules (as recently reviewed by Wang et al. 2021b; German et al. 2023; Alazem and Burch-Smith 2024). The plasmodesmal aperture itself is regulated via callose deposition and degradation at the plasmodesmal neck by callose synthases (CalS) and β-glucanases respectively (Levy et al. 2007; Vatén et al. 2011). The decrease in plasmodesmal aperture in response to stress is a process activated by signaling cascades mediated by specific machinery. For example, LYSM-CONTAINING GPI-ANCHORED PROTEIN 2 (LYM2; Faulkner et al. 2013) and CALMODULIN-LIKE 41 (CML41; Xu et al. 2017) specifically mediate plasmodesmal responses to the microbial elicitors chitin and flg22 respectively, and

PLASMODESMATA LOCATED PROTEIN 5 (PDLP5) integrates these responses, as well as plasmodesmal responses triggered by the defense hormone salicylic acid (SA; Wang et al. 2013; Tee et al. 2023b). The observation that *lym2* mutants cannot close their plasmodesmata in response to chitin but execute normal chitin-triggered mitogen-activated protein kinase (MAPK) activation and apoplastic ROS production (Faulkner et al. 2013) indicates that plasmodesmal signaling cascades can act independently of other immune responses. This independence of response suggests that whether plasmodesmata are open or closed might be informed by another layer of cellular regulation.

It has been observed that plasmodesmal permeability decreases upon PAMP perception within 30 min (Xu et al. 2017) and is still detected after 24 hours (Lim et al. 2016; Li et al. 2021). Both PAMP perception and pathogen infection initiate a wide array of defense response making it challenging to identify the specific contribution of plasmodesmal closure to immunity. Further, in the context of an infection, many pathogens deploy effectors targeting plasmodesmata that suppress plasmodesmal closure (e.g. Aung et al, 2020; Tomczynska et al. 2020; Li et al. 2021; Ohtsu et al. 2024), further complicating the analysis of the role of plasmodesmata in host immune execution in an infection context.

Whether plasmodesmata can close or not during immune responses determines whether a full defense response can be executed, but untangling how plasmodesmal closure contributes to overall immunity is a complex problem. To begin to address this we have simplified the question and asked what cellular responses are triggered by plasmodesmal closure, and how closing plasmodesmata affects elements of an immune response. We generated two genetic tools with which we can induce plasmodesmal closure independently of an immune or stress elicitor. By identifying what responses are trigged by plasmodesmal closure in both tools, we found that plasmodesmal closure itself instigates a subset of stress responses including transcriptional reprogramming and salicylic acid (SA) production. This raises further questions regarding how this contributes to the orchestration of immune responses, how plasmodesmal closure triggers stress responses and whether these stress responses occur during the broad range of physiological processes during which plasmodesmata close.

Results

Genetic tools induce plasmodesmal closure independent of a physiological elicitor

To manipulate plasmodesmal closure independently of external signals we generated two transgenic lines in which estradiol application can induce callose deposition at plasmodesmata. First, we exploited the *icals3m* overactive *CalS3* allele (Vatén et al. 2011) that enhances callose deposition at plasmodesmata; *icals3m* has been used extensively as a tool to understand symplastic connection in development (e.g., Paterlini et al. 2021; Ross-Elliott et al. 2017; Sevilem et al. 2012; Wu et al. 2016; Yadav et al. 2014) but less is known about its affect on immunity. As a callose synthase, this enzyme acts terminally in the plasmodesmal-associated callose deposition pathway. Secondly, we utilized the observation that a synthetic peptide that fuses the PDLP1 signal peptide to the 5' end and the PDLP1 transmembrane domain and cytoplasmic tail to the 3' end of mCherry also promotes callose deposition at plasmodesmata (Caillaud et al. 2014). We fused these two transgenes independently to the *XVE* chimeric transcription activator and *LexA* operator (Zuo et al. 2000) and transformed them into *Arabidopsis thaliana* Col-0 to create two different inducible tools, named LexA::icals3m and LexA::PD-Plug respectively. To identify functional transgenic lines, we performed qPCR, copy number analysis and/or live imaging (Supplemental Table 1; Supplemental Fig. S1).

To functionally characterize the selected LexA::PD-Plug and LexA::icals3m lines, we first measured callose deposition by quantitative live cell imaging of aniline blue stained plasmodesmal callose. We found that there was an increase in callose deposition at plasmodesmata 24 hours (h) post estradiol treatment in both LexA::PD-Plug and LexA::icals3m when compared to the DMSO treatment, although there was a greater response in LexA::icals3m (Fig. 1a; Supplemental Fig. S2) suggesting that this line deposits more callose at plasmodesmata than LexA::PD-Plug. To determine whether increased callose deposition perturbed plasmodesmal permeability, we used microprojectile bombardment assays of *35S::GFP* to measure GFP flux between cells in DMSO or estradiol treated leaves. While estradiol treatment did not impact GFP flux in comparison to DMSO in wild-type Col-0 plants, we found that estradiol induction of either transgene reduced GFP flux between cells, indicating plasmodesmal aperture

was reduced and cell-to-cell connectivity decreased (Fig. 1b-c; Supplemental Data Set

1). Correlated with the greater callose deposition at plasmodesmata in LexA::icals3m,

the extent of GFP flux was significantly decreased in induced LexA::icals3m compared

to induced LexA::PD-Plug (Fig. 1c). From these results, we conclude that estradiol

application to both LexA::PD-Plug and LexA::icals3m, but not Col-0, induces

plasmodesmal callose deposition, and plasmodesmal closure, which reduced

symplastic flux.

We noted that the data obtained from GFP mobility assays indicates that induced

LexA::PD-Plug displays a heterogeneous response in cell-to-cell connectivity; while

there is an increase in the population of cells that exhibit plasmodesmal closure in

induced tissues, a subset of the population of cells remain connected to their

neighbors (Fig. 1c; Supplemental Fig. S3). This is reminiscent of the data obtained

from GFP mobility assays of leaves treated with elicitors such as chitin and flg22

(Cheval et al. 2020; Tee et al. 2023). By comparison, induced LexA::icals3m displays

a more homogenous response in which low cell-to-cell movement with low variance is

observed (Fig. 1c; Supplemental Fig. S3), similar to transgenic lines such as the

PDLP1-overexpressor (Tee et al. 2023). Despite differences in the degree of

plasmodesmal closure, these transgenic lines allow us to manipulate plasmodesmata

independently of both endogenous and non-endogenous signals and investigate plant

responses to plasmodesmal closure.

Plasmodesmal closure triggers stress transcriptional responses and salicylic acid

production

As pathogens are capable of manipulating plasmodesmal aperture during infection, it

is difficult to determine whether plants sustain immune-associated plasmodesmal

closure when challenged with a pathogen over time. Thus, we utilized the secretion-

defective *Pseudomonas syringae* DC3000 (*Pst* DC3000) mutant strain *hrcC* that would

not secrete effectors that manipulate plasmodesmal aperture, and assayed for

plasmodesmal callose deposition at 24 h, 48 h and 72 h. We found that callose was

deposited throughout the leaf at all time points, and that plasmodesmal callose

deposition significantly increased at 48 h and 72 h post treatment (Supplemental Fig.

S4). This indicates that plasmodesmal closure can be sustained throughout an
infection context.

To investigate the impact of plasmodesmal closure on leaf tissue, we next
characterized our plasmodesmal closure inducible tools using RNAseq. Given our
findings that plasmodesmal closure is sustained up to 72 h after initial infection
(Supplemental Fig. S4), we profiled transcriptional changes at 12 h, 24 h, 48 h and 72
h following DMSO or estradiol treatment.

Preliminary analysis of the data first revealed that a critical driving factor of variance
was time (Supplemental Fig. S5). Therefore, all subsequent analyses were performed
within single time points. We assessed the effect of estradiol relative to DMSO
treatment within a genotype, identifying differentially expressed genes (DEGs). The
greatest response was in Col-0 at 12 h, where estradiol led to the downregulation of
over a thousand genes (Supplemental Fig. S6). The presence of shared estradiol-
responsive DEGs in Col-0 and the transgenic genotypes support an early estradiol
specific response not correlated to transgene induction. The number of these shared
estradiol-induced DEGs detected was dramatically reduced after 24 h (Supplemental
Data Set 2; Supplemental Fig. S6), suggesting secondary effects of estradiol are
negligible after this point.

Assessing transgene induction across the sampling period, both the PD-Plug
transcripts (as indicated by read counts of mCherry) and icals3m transcripts (as
indicated by read counts of CalS3) were upregulated compared to controls by 12 h
post estradiol treatment and remained significantly elevated at 72 h (Supplemental
Fig. S7). Specifically, the PD-Plug transgene (measured by mCherry counts) was
induced similarly to other highly expressed genes such as photosynthetic gene
RBCS1A (AT1G67090; Supplemental Dataset 3). To determine that the upregulation
of the transgenes did not inconsequentially result in stress, particularly in the ER from
high levels of ectopic protein production, we examined the expression of genes known
to be ER stress markers, defined as either ER stress sensors or unfolded protein
response (UPR) effectors (Beaugelin et al. 2020). Of the 13 genes examined, only
BIP3 (AT1G09080) was significantly upregulated in estradiol treated LexA::PD-Plug
and LexA::icals3m at 72 h (Supplemental Fig. S8), suggesting that transgene
expression in these lines does not induce ER-stress.

Next we investigated the gene expression changes specific to the induction of each
transgene. Thus, we used a likelihood-ratio test of all genotypes and treatments to
determine differentially expressed genes specific to estradiol treated LexA::PD-Plug
and estradiol treated LexA::icals3m (Supplementary Dataset 4; Supplemental Fig.
S9). At both 12 h and 24 h, we found no genes uniquely upregulated in the estradiol
treated samples from either line. At 48 h, 28 and five genes were uniquely upregulated
in estradiol treated LexA::PD-Plug and estradiol treated LexA::icals3m respectively. At
72h, 33 genes were differentially upregulated in estradiol treated LexA::icals3m only;
there were no genes differentially upregulated exclusively in estradiol treated
LexA::PD-Plug at 72 h. Based on the low rate of transgene-specific upregulation, we
next investigated genes shared between the two tools, as these would be most
informative for identifying core plasmodesmal closure related changes.

While expression of both the *icals3m* and *PD-Plug* transgenes have specific
transcriptional dynamics (Supplemental Fig. S7), and different modes of action as
proteins, we reasoned that the genes that are commonly up- or down-regulated
following induction are associated with their common effect on plasmodesmata.
Therefore, to identify genes which are uniquely upregulated when plasmodesmata are
closed, we defined two different groups for comparison: plasmodesmal state closed
(estradiol treated LexA::PD-Plug and LexA::icals3m) and plasmodesmal state open
(DMSO treated LexA::PD-Plug and LexA::icals3m, and all Col-0 samples). A
likelihood-ratio test was performed on these two groupings at each time point to
identify genes that were significantly differentially expressed only in the closed
plasmodesmal state (Fig. 2a; Supplemental Fig. S10; Supplemental Data Set 5-8) and
observed that the number of upregulated DEGs increased over time, suggesting that
the longer plasmodesmata remain closed, the greater the cellular response. Notably,
the number of differentially expressed genes that was shared between LexA::icals3m
and LexA::PD-Plug in the grouping of plasmodesmal state closed was greater than
the number of up-regulated DEGs specific to each line (Fig. 2a). A GO term analysis
identified that genes upregulated when plasmodesmata are closed at 48 and 72 h are
enriched in defensive processes (Fig. 2b; Supplemental Fig. S11). These include
response to bacterium, oomycetes, chitin and fungus, regulation of systemic acquired
resistance, and response to salicylic acid (SA).

As response to SA at 72 h was the most significantly enriched GO term in the entire
analysis, we further explored this by analyzing DEGs related to SA to determine what
aspects of SA signaling were impacted when plasmodesmata were closed (Fig. 2c;
see Supplemental Data Set 9 for associated references; Vlot et al. 2008; Seyfferth et
al. 2014; Yang et al. 2015; van Butselaar and Van den Ackerveken 2020) and found
both SA biosynthesis/catabolism and SA response pathways were significantly
upregulated at 48 and 72 h. To confirm that the transcriptional signature of SA is a
consequence of the activation of SA synthesis when plasmodesmata are closed, we
measured SA content in Col-0, LexA::PD-Plug and LexA::icalsm3 72 h post DMSO
and estradiol treatment. We found induced plasmodesmal closure increased SA
content in both LexA::PD-Plug and LexA::icals3m, with induced LexA::icals3m having
a greatly increased SA content, being three-fold higher when compared to DMSO
treatment (Fig. 2d). Thus, plasmodesmal closure leads to the upregulation of
transcriptional defense responses, and SA synthesis and accumulation, independent
of an immune stimulus.

Plasmodesmal closure differentially impacts pathogen resistance and immune
responses

Plants that lack plasmodesmata-specific immune signaling machinery and cannot
close their plasmodesmata in response to pathogen infection, show increased
susceptibility to different pathogenic microbes (Lee et al. 2011; Faulkner et al. 2013;
Xu et al. 2017). To determine whether an inverse correlation exists for induced
plasmodesmal closure we examined the susceptibility of induced, transgenic lines to
the bacterial, biotrophic pathogen *Pst* DC3000. We inoculated mature leaves of Col-
0, LexA::PD-Plug and LexA::icals3m plants 72 h post DMSO or estradiol treatment
(Fig. 3a) and found that induction of both transgenes reduced bacterial growth 3 days
post infection (dpi). Thus, plasmodesmal closure is correlated with enhanced
resistance to *Pst* DC3000. Next, we assayed for resistance to the fungal necrotroph,
Botrytis cinerea. In contrast with *Pst* DC3000 infection, we found induction of the PD-
Plug transgene had no impact on *B. cinerea* infection when compared to the disease
caused in Col-0 plants, but induction of *icals3m* increased susceptibility to *B. cinerea*
(Fig. 3b; Supplemental Fig. S12). This indicates that plasmodesmal closure
specifically does not enhance resistance in all infection contexts.

Overall susceptibility is driven by multiple components of host defense responses, and plasmodesmal closure is a key immune response to the pathogen associated molecular patterns (PAMPs), flg22 and chitin. Previously, we have observed that PAMP-triggered plasmodesmal closure occurs independently of other PAMP responses (Faulkner et al. 2013; Xu et al. 2017). To confirm that plasmodesmal closure in our transgenic lines is likewise independent of PAMP responses and not the cause of the increased resistance we observed to *Pst* DC3000, we examined the effect of plasmodesmal closure on the activation of MAPK signaling and ROS production by flg22. When we assayed for flg22-triggered MAPK phosphorylation in LexA::PD-Plug and LexA::icals3m lines following estradiol treatment, we saw no difference in the timing or magnitude of the response in either line when compared to Col-0 in mature leaves and seedlings (Fig. 3c; Supplemental Fig. S13). However, while induction of *LexA::PD-Plug* did not perturb the flg22-triggered ROS burst in either seedlings or mature leaves (Fig. 3d,e), when we assayed for the same response in LexA::icals3m, induction of the transgene reduced the flg22-triggered ROS burst in mature leaves (Fig. 3d) and increased the response in seedlings (Fig. 3e). To determine whether there were differences in general basal cellular ROS levels that might explain this observation, we stained estradiol treated Col-0 and LexA::icals3m leaves with H_2DCFDA but observed no quantitative differences between the genotypes or treatments (Supplemental Fig. S14), indicating plasmodesmal closure itself does not alter basal ROS cellular levels.

These results indicate that plasmodesmal closure differentially influences the outcomes of host interactions with different pathogenic microbes, possibly correlated with differences in pathogen lifestyle. Increased resistance to *Pst* DC3000 is not associated with enhanced MAPK activation or, given that our pathoassays were performed on mature leaves, changes in ROS burst responses. Further, our ROS data raises the possibility that plasmodesmal closure differentially interacts with immune responses in different developmental stages.

Plasmodesmal closure induces sugar accumulation and growth defects

Given the increased defense related transcripts, we next explored how these modulated differences in plasmodesmal permeability could translate phenotypically.

As plasmodesmata connect the phloem to cells in young sink tissues and allow sugar translocation, we expect that plasmodesmal closure might reduce the flux of sugars to growing tissues and limit growth. Alternatively, activation of defense processes negatively regulate growth and therefore it is possible that plasmodesmal closure impairs growth by one or both pathways. Indeed, several transgenic lines known for constitutive plasmodesmal closure exhibit a significant retardation in growth such as *PDLP1, PDLP5* and *PDLP6* overexpressors (Thomas et al. 2008; Lee et al. 2011; Li et al. 2024). For the *PDLP1* overexpressor, the reduced cell-to-cell connectivity profile is more homogenous than observed with physiological responses, similar to the LexA::icals3m induced tool (Fig. 1c, Supplemental Fig. S3). Thus, we explored whether sustained plasmodesmal closure reduced growth in induced LexA::PD-Plug and LexA::icals3m. In 9-day old seedlings, we applied either DMSO or estradiol treatment and measured the first true leaves (Fig. 4a) six days post treatment. We found that estradiol reduced leaf growth in both Col-0 and LexA::PD-Plug in comparison to the DMSO treated seedlings. Further, the estradiol-induced growth reduction was greater in LexA::icals3m relative to LexA::PD-Plug and Col-0. Alongside reduced leaf size, we observed yellowing and chlorosis on the true leaves of estradiol treated LexA::icals3m (Supplemental Fig. S15), a finding similar to increased senescence observed in the PDLP5 overexpressor (Lee et al. 2011).

Plasmodesmal closure might be expected to restrict photoassimilate distribution between source and sink tissue, and indeed, a reduction in plasmodesmal permeability and rosette size has been documented in *Arabidopsis* alongside a significant accumulation of soluble sugars and starch (e.g., in the MOVEMENT PROTEIN 17 *MP17* overexpressor, *CHOLINE TRASPORTER-LIKE 1 cher1* mutant, *PDLP5* and *PDLP6* overexpressors; Kronberg et al. 2007; Kraner et al. 2017; Li et al. 2024). Therefore, we quantified sugar accumulation in source tissues following plasmodesmal closure, measuring the sucrose, fructose, glucose, and starch content of source leaves of plants 72 h post treatment. These data reveal a significant increase in all four sugars in estradiol treated LexA::icals3m samples (Fig. 4b). Reasoning that an increase in sugar content might down-regulate photosynthesis, we used chlorophyll fluorescence imaging to infer photosynthetic yield and found there was no difference in photosynthetic yield of PSII (as indicated by Y(II)) after 72 h in Col-0, LexA::PD-Plug and LexA::icals3m between the DMSO or estradiol treatments (Table 1). As there were

no changes in the photosynthetic yield that correlated with the increase in sugar content in induced LexA::icals3m, we next hypothesized that plasmodesmal closure and sugar accumulation might activate the expression of sugar transporters and starch biosynthesis genes to reduce the concentration of soluble sugars in the cytosol. However, out of 125 genes known to be sugar and starch related (Supplemental Data Set 10; Büttner and Sauer 2000; Williams et al. 2000; Chen et al. 2010; Monroe and Storm 2018; Abt and Zeeman 2020; Preiser et al. 2020; Smith and Zeeman 2020; David et al. 2022; Bavnhøj et al. 2023), only the transporter *SENESCENCE-ASSOCIATED GENE 29 (SAG29)/SUGARS WILL EVENTUALLY BE EXPORTED TRANSPORTERS 15 (SWEET15)* was significantly upregulated when plasmodesmata are closed, in LexA::icals3m at 48h and 72 h (Supplemental Fig. S16, S17). AT3G20460 was differentially expressed in LexA::icals3m estradiol treatment at 72 h and is proposed to be a monosaccharide transporter (Johnson et al. 2006). However, this gene was downregulated in response to plasmodesmal closure and as the function of this gene is not yet well characterized it is difficult to speculate on any possible role in the sugar/starch phenotype (Johnson et al. 2006). These results indicate that while plasmodesmal closure induces starch and soluble sugar accumulation, it does not perturb photosynthesis or the expression of sugar transporters.

Discussion

Plants initiate a suite of defense responses when challenged by a microbial threat. Plasmodesmal closure is an early reaction, initiated within 30 minutes upon perceiving pathogen-derived molecules and stress-associated signals including SA and hydrogen peroxide (Xu et al. 2017; Cheval et al. 2020; Tee et al. 2023b), and is critical for the execution of a full response. Thus, we reason that plasmodesmal closure regulates a range of downstream processes. However, considering the plethora of responses elicited during immunity, and the array of elicitors that trigger plasmodesmal closure via converging signaling pathways (Tee et al. 2023b), it is challenging to identify what specific contribution plasmodesmal closure makes to the overall execution of immunity. Using lines with which we can induce plasmodesmal closure independently of elicitors, we explored what occurs when plasmodesmata close

without other co-incident processes and determined that plasmodesmal closure itself
induces stress.

Transcriptional profiling of responses to plasmodesmal closure identified
transcriptional signatures of defense responses associated with SA production and
signaling. It has long been known that constitutive over-production of the
plasmodesmal protein PDLP5 closes plasmodesmata and leads to high levels of SA
(Wang et al. 2013). However, as PDLP5 is transcriptionally responsive to SA, it is
viewed as a stress response element (Wang et al. 2013). We have previously shown
that microbial elicitors, flg22 and chitin, also require PDLP5 to elicit plasmodesmal
closure, but these elicitors interestingly do not require ISOCHORISMATE SYNTHASE
1 which is involved in the synthesis of SA (Wildermuth et al. 2001), for the
plasmodesmal response (Tee et al. 2023b). This poses a potential model in which
plasmodesmal closure is first initiated upon pathogen perception, leading to the
accumulation of SA at plasmodesmata to create a feedback loop to sustain closure.
Our finding that plasmodesmal closure leads to SA accumulation also raises the
question of whether plasmodesmal closure in all contexts induces SA synthesis and
accumulation, and thus if SA plays a role in developmental contexts which require
plasmodesmal closure.

The accumulation of SA when plasmodesmata are closed might be elicited directly in
response to plasmodesmal signaling, or indirectly as a secondary consequence of
another process triggered by a change in cellular status induced by isolation. The
greater accumulation of SA observed in induced LexA::icals3m occurs in parallel with
an observed increase in sugar levels (Fig. 4c) and is correlated with the greater degree
of plasmodesmal closure observed in this line relative to LexA::PD-Plug. It is possible
that the increase in sugar content induces osmotic stress or another process that feeds
into SA signaling, and therefore that these stress responses are secondary to
plasmodesmal closure. Further, the combined increase in sugar content and SA might
explain our pathoassay data; the necrotrophic pathogen *B. cinerea* might be able to
access increased intracellular sugar stores and be unimpeded by SA defense
pathways (Ferrari et al. 2003; El-Oirdi et al. 2011), while *Pst* DC3000 might be unable
to access intracellular sugars and is impeded by SA (Velásquez et al. 2017; Wilson et
al. 2017; Howlader et al. 2020).

Sugars can induce specific signal transduction pathways, with some studies finding
that increasing source leaf carbohydrate content leads to a feedback loop of
decreasing photosynthetic yield (Araya et al. 2006). Additionally, an inverse
relationship between plasmodesmal aperture and the expression/activity of sugar
transporters (Ruan et al. 2001; Zhang et al. 2017; Liu et al. 2022) has been reported
suggesting that cells maintain the capability to transport sugars between cells by
balancing plasmodesmal aperture with transporter activity (Zhang et al. 2017). On this
basis, we might expect that induction of LexA::icals3m would activate a compensatory
mechanism for sugar distribution. However, we only observed that one sugar
transporter was upregulated in induced LexA::icals3m, SAG29/SWEET15, which is
also activated by SA during the process of leaf senescence (Seo et al. 2011; Wang et
al. 2021a). Therefore, while this may be a component of the senescing phenotype
observed in induced LexA::icals3m, the increased expression of this transporter does
not counteract the intracellular sugar accumulation observed (Chen et al. 2015).
Irrespective, the question arises as to whether there is a cellular mechanism that
perceives an increase in soluble sugars and whether this is incapacitated when
plasmodesmata are closed independently of other signal inputs.

The variance in multiple sets of experimental data suggest that physiological
plasmodesmal responses are heterogeneous within a tissue, i.e., not all cells close
their plasmodesmata in response to an elicitor such as chitin or flg22 (Faulkner et al.
2013; Xu et al. 2017; Cheval et al. 2020; Tee et al. 2023b). Our transgenic lines show
two different patterns of plasmodesmal response: we see variability in GFP mobility in
the induced LexA::PD-Plug line that resembles physiological data sets despite
homogeneous induction of the transgene (Supplemental Fig. S1), while data from
induced LexA::icals3m (Fig. 1b,c) exhibits much lower variance (Supplemental Fig.
S3). This suggests that more cells are isolated from their neighbors than normally
occurs and LexA::icals3m induces plasmodesmal closure more homogeneously than
in LexA::PD-Plug. This increased homogeneity of plasmodesmal closure correlates
with the magnitude of the responses measured in most of the assays performed in this
study, with effects on disease resistance, growth, ROS burst, sugar and SA
accumulation all greater in magnitude in, or only detected in, LexA::icals3m relative to
LexA::PD-Plug. Thus, our data suggests that we have detected some of these effects
only when they have been elicited in more cells. However, an alternative possibility is

that some responses to plasmodesmal closure occur only when a threshold of isolated
cells is reached, which would imply that responses to plasmodesmal closure is a tissue
level response, and that connectivity is a population-level parameter. Indeed, the
resistance of both LexA::PD-Plug and LexA::icals3m to *Pst* DC3000, and the increase
in salicylic acid and defensive transcripts, indicates that heterogenous plasmodesmal
closure can intersect with immune signaling. Further, that homogeneous cellular
isolation results in undesired outcomes suggests a degree of heterogeneity is optimal
for overall physiology.

In this study we have induced plasmodesmal closure in a non-physiological manner
to isolate it from other immune responses and study its effects. This raises the
consideration of how responses to induced plasmodesmal closure correspond to the
effects of plasmodesmal closure that occurs as a physiological response to stress. For
example, how long plasmodesmal closure is sustained in different response contexts
is not well described, and whether SA synthesis and accumulation occur within these
timeframes is an important question. However, during an infection, the exposure of
cells to PAMPs is likely to be a sustained process suggesting that prolonged
plasmodesmal closure occurs during infection which is supported by our findings of
increased plasmodesmal callose deposits 72 h post *hrcC* infection (Supplemental Fig.
S4). Further, given we have identified plasmodesmal closure without immune
activation triggers stress-associated responses, whether these responses occur when
cells are isolated during developmental transitions must also be considered. Likewise,
in many investigations regarding developmental transitions, the icals3m tool is used
to induce ectopic plasmodesmal closure at different cell boundaries and our data
indicates that this is also likely to block the distribution of sugars to, or from, targeted
cells and activate defense. Thus, phenotypes that arise from blocking plasmodesmata
at a critical developmental transition must be interpreted with these factors in mind.

We have found that plasmodesmal closure triggers stress responses. This suggests
that cellular isolation is a stress-inducing state and, conversely, that connection to
neighboring cells is essential for optimal growth-associated cellular functions. Thus,
understanding how a cell responds when it becomes isolated from its neighbors, and
why cell-to-cell connectivity regulates homeostatic cellular processes, is key to
understanding how plants function as multicellular organisms.

Methods

Plant material

All *Arabidopsis thaliana* used in this study is in the Columbia-0 ecotype background. For assays on mature plants, *Arabidopsis* plants were grown at 10 h light at 22 °C, on soil or for the RNA-seq experiment, on Murashige and Skoog (MS) 1% sucrose 0.8% agar in 50 mm diameter petri dishes for 4–5 weeks. For MAPK and ROS measurement assays on seedlings, seeds were germinated on MS 1% sucrose 0.8% agar at 16 h light at 22 °C and grown for 8 days, then 10 seedlings were transferred per well on a 6-well plate together with 8 mL of liquid MS and grown for an additional 7 days.

Confocal microscopy

Microprojectile bombardment, transgene induction visualization and aniline blue plasmodesmal callose measurements utilized confocal microscopy (LSM Zeiss 800). RFP/mCherry was excited with a 561-nm DPSS laser and collected at 600 to 640 nm, GFP/H_2DCFDA was excited with a 488-nm argon laser and collected at 505 to 530 nm, and aniline blue was excited with a 405-nm UV laser and collected at 430 to 550 nm.

Microprojectile bombardment

Microprojectile bombardment of *35S::GFP* was performed as described in (Tee et al. 2022). 24 h after DMSO or estradiol treatment, 5-week-old leaves were co-bombarded with pB7WG2.0.RFP$_{ER}$ and pB7WG2.0.GFP. Bombardment sites were then imaged 16 h after using confocal microscopy with a 20× water dipping objective (W N-Achroplan 20×/0.5).

Generation of transgenic lines and induction of transgene

The *LexA::icals3m* construct is as described (Bellandi et al. 2022). The *LexA::PD-Plug* construct was assembled into a binary vector via Golden Gate cloning; first the 'PD-Plug' sequence was created by fusing the PDLP1 signal peptide and the PDLP1 transmembrane domain and cytoplasmic tail to the N- and C-termini of mCherry respectively (Caillaud et al. 2014). This synthetic peptide was inserted downstream of the *LexA* promoter sequence and upstream of a NOS terminator and assembled into

the backbone of pICSL0022013 containing the *Nos::BAR::Nos* selection cassette and
Act2::XVE::Act2 cassette.

LexA::PD-Plug and *LexA::icals3m* constructs were transformed into the Col-0
Arabidopsis background and screened with phosphinothricin (10 µg/mL concentration)
or kanamycin selection (50 µg/mL concentration) respectively. For induction of the
transgene in all experiments (with the exception of size phenotyping), leaves were
painted on both the abaxial and adaxial side with either 0.1% DMSO (mock treatment)
or 20 µM β-Estradiol 17-acetate (estradiol treatment) containing 0.01% Silwet L-77 in
ddH$_2$O, using a fine paint brush. For size phenotyping, seedlings were sprayed with
the above specified DMSO or estradiol treatment.

Plasmodesmal callose measurements

24 h post treatment with either DMSO or estradiol, expanded leaves of 5-week-old
plants were infiltrated with 1% aniline blue in PBS buffer (pH 7.4) then imaged using
confocal microscopy from the abaxial side using a 63× water immersion objective
(Plan-Apochromat 63×/1.20). Z-stacks from multiple areas from a given treated leaf,
with a minimum of three biological replicates per genotype and treatment were
collected. An automated image analysis pipeline to quantify aniline blue-stained
plasmodesmata is available at (Johnston et al. 2022).

Quantitative PCR

To determine transgene expression, *Arabidopsis* T2 seedlings were grown on MS 1%
sucrose 0.8% agar with phosphinothricin (10 µg/mL concentration) or kanamycin (50
µg/mL concentration), for 21 days in 16 h light at 22 °C. Leaves from each line was
treated with DMSO or estradiol as above and harvested 48 h later in liquid nitrogen.
Samples were homogenized in a GenoGrinder 2100, with RNA extracted, DNase
treatment, first strand cDNA synthesis and qPCR performed as previously described
(Bellandi et al. 2022). Genes tested are *icals3m*/*CalS3* (using primers
AGGACGTTACTTCATGGCGG and GGCTCCGGGAATACAGTCTG) and the
housekeeper *AtUBQ10* (using primers AGTCTACTCTTCACTTGGTCCTGC and
GCCCCAAAACACAAACCACCAAAG). Normalized relative quantities (NRQs) were
generated using the qBase model (Hellemans et al. 2008).

Pseudomonas syringae infection assays

Three leaves from 4–5-week-old plants were treated with DMSO or estradiol. After 72 h, leaves were infiltrated with 5×10^4 cfu/mL *Pseudomonas syringae* DC3000 and harvested 72 h later. Each leaf was harvested with a 5 mm corer and the three leaves from a single plant combined as a sample. Leaf tissue was homogenized with 1 mL ddH$_2$O for 2 mins at 1200 rpm in a GenoGrinder 2100. 5 µL of multiple 10-fold dilutions were plated out with three technical replicates per plant sample, then incubated for 48 h at 28 °C with colony counts determined from countable dilutions. Four independent experiments with five plants per treatment/genotype per experiment was performed.

To determine plasmodesmal associated callose deposits after *Pseudomonas syringae* DC3000 *hrcC* (*hrcC*) infection, a culture at an OD of 0.2 was prepared. Leaves from 5-week-old Col-0 plants were infiltrated either with H$_2$O (mock) or *hrcC*, and then imaged 24 h, 48 h or 72 h post treatment. Aniline blue staining was performed as per the above plasmodesmal callose measurements with the exception that 0.1% aniline blue was used. 3-4 z-stacks from multiple areas per leaf was performed, with eight biological replicates per treatment/timepoint collected. Only callose deposits at plasmodesmata were analyzed.

Botrytis cinerea infection assays

B. cinerea spores were applied to leaves from 4–5-week-old plants 72 h post DMSO or estradiol treatment. *B. cinerea* spores were harvested and adjusted to a concentration of 2.5×10^5 spores/mL in 0.25× potato dextrose broth and incubated at room temperature for 2 h with continual shaking for spore germination. Leaves were adhered to 1.5% agar H$_2$O plates, and droplets of 2 µL spore suspensions were placed in between the mid vein and leaf edge on each side. Plates were sealed with parafilm and incubated in a cabinet set at 22 °C with 10 h light. Developing disease lesions were photographed 2 dpi and measured using FIJI. Each prepared plate had four leaves per genotype and treatment, and the average lesion area was calculated per genotype/treatment in a given plate, defined as a single biological replicate. Three independent experiments were conducted with a minimum of five biological replicates per genotype/treatment.

MAPK activation assays

MAPK activation assays were performed on leaf discs from expanded leaves of 5-week-old plants or on 15-day-old seedlings 72 h post treatment with DMSO or

estradiol. Samples were harvested 0, 5 mins, 15 mins, 30 mins and 45 mins post 100

nM flg22 treatment, and flash frozen in liquid nitrogen. Frozen samples were

homogenized via 90 seconds of shaking (1100 rpm) in a GenoGrinder 2100, then 500

μL protein extraction buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA pH

8.0, 10% glycerol, 0.5% NP40 IGEPAL® CA-630, protease inhibitor cocktail (Sigma)

1:100, phosphatase inhibitor (Sigma) 1:200, 1 mM $Na_2MoO_4 \times 2H_2O$, 1mM NaF, 1.5

mM activated Na_3VO_4, 5 mM DTT, 1mM PMSF) was added and the sample left to mix

for 30 mins at 4 °C. Samples were centrifuged at max speed for 10 mins at 4 °C,

supernatant transferred to a new tube and re-centrifuged at max speed for another 10

mins at 4 °C. The resulting supernatant was incubated at 95 °C for 15 mins with

Laemmli buffer, then proteins were separated by SDS-PAGE and transferred to

Immuno-blot® PVDF membrane. Proteins were detected using p44/42 MAPK (1:2000;

Cell Signaling Technologies, #9102) and α-Rabbit-HRP (1:10000, Sigma, A0545).

ROS burst measurements

ROS assays were performed on leaf discs taken from 5-week-old plants or 15-day-old

seedlings 72 h post DMSO or estradiol treatment. Leaf discs were first floated on H_2O

overnight; the following day water was removed, then 20 μg/mL HRP and 6 μM L-012

was added, with or without 100 nm flg22. The same was performed for seedlings

except liquid media was removed before adding the HRP and L-012.

Chemiluminescence was recorded using a Varioskan Flash (Thermo Fisher) over 45

mins, with the total luminescence emitted used for analysis.

General ROS quantification

20 μM H_2DCFDA (Thermo Fisher) was syringe infiltrated into expanded leaves of 5-

week-old plants 72 h post DMSO or estradiol treatment, and confocal microscopy was

performed on the abaxial side of the leaf. Z-stacks were taken from four areas of a

given leaf from four plants per genotype/treatment. Quantification of fluorescence was

performed in FIJI using a sum projection.

RNA-seq

Two mature leaves per 5-week-old plant were DMSO or estradiol treated, and

harvested 12 h, 24 h, 48 h and 72 h post treatment. Each biological replicate contained

four plants (i.e., 8 leaves), with three biological replicates for each

genotype/treatment/time point. Leaves were flash frozen in liquid nitrogen, then

homogenized via 90 seconds of shaking (1100 rpm) in a 2100 Geno/Grinder (Spex

SamplePrep, USA). Total RNA was extracted using the RNEasy Plant Mini Kit

(QIAGEN, Germany) and eluted into 60 µl of water. Purified RNA was treated with the

TURBO DNA-free kit (Thermo Fisher, USA). RNA quantification and library

construction was conducted by Novogene Co., Ltd (Beijing). RNA quality and quantity

were access via 1% agarose gel, NanoPhotometer ® spectrophotometer (IMPLEN,

USA), and the RNA Nano 6000 Assay Kit for the Bioanalyzer 2100 System (Agilent

Technologies, USA). Libraries were constructed using 0.4 µg of RNA and the

NEBNext ® UltraTM RNA Library Prep Kity for Illumina (NEB, USA). mRNA was

purified via poly-T magnetic beads. Fragmentation, cDNA synthesis, and NEBNext

Adaptor hybridization were performed according to manufacturer's instructions and

purified with the AMPure XP system (Beckman Coulter, USA). Index-coded samples

were clustered on a cBot Cluster Generation System using TruSeq PE Cluster Kit v3-

cBot-HS (Illumina, USA) and sequenced on an Illumina Novaseq platform to generate

150bp paired-end reads with a sequencing depth of 10 million reads per sample. All

sequencing data are available on GEO at GSE248301.

Bioinformatic analyses

Adaptors and low-quality (phred33<20) reads were trimmed from the data using

TrimGalore v0.5.0 (Krueger et al. 2021) and Cutadapt v1.7 (Martin 2011). RNA quality

was accessed using FastQC v0.11.8 (Babraham Bioinformatics - FastQC A Quality

Control tool for High Throughput Sequence Data) and MultiQC v.1.7 (Ewels et al.

2016). The data was mapped against the *Arabidopsis* reference genome (TAIR10;

NCBI Refseq GCA_000001735.1) using HISAT2 v2.1.0 (Kim et al. 2019). Mapped

reads were sorted using Samtools v1.4.1 Sort (Danecek et al. 2021). Transcript

assembly was completed using Stringtie v1.3.3 (Pertea et al. 2015) and gene counts

(see Supplemental Data Set 3 and 11 for gene count matrix and metadata

respectively) were extracted using the built-in python wrapper, prepDE.py available at

http://ccb.jhu.edu/software/stringtie/dl/prepDE.py. Principle component analysis was

conducted using DEseq2, ggplot2 (Wickham 2016), ggConvexHull, and TidyVerse

(Wickham et al. 2019) in R (R Core Team 2022). Differentially expressed genes were

determined using DEseq2 (Love et al. 2014) in R. Genes with less than 10 counts

across all samples were excluded from the analysis. The data was subset by timepoints, and Wald Tests or Likelihood Ratio Tests (LRT) were performed based on comparisons of interest. Wald Tests were used to extract DEGs based on differences between treatment or genotype. LRT were used to extract genes differentially expressed based on difference between closed plasmodesmata (induced/estradiol treated LexA::PD-Plug and LexA::icals3m combined) and open plasmodesmata (uninduced/DMSO treated LexA::PD-Plug, uninduced/DMSO treated LexA::icals3m, and DMSO and estradiol treated Col-0). LRT was used in this way to determine genes uniquely up- or down-regulated in the inducible lines treated with estradiol. Differentially expressed genes from the Wald Test were designated if the log2 fold change (lfc) was greater than |1.5| and adjusted p-value<0.05. Differentially expressed genes from the LRT were considered significant if the adjusted p-value was less than 0.05. LRT significantly differentially expressed genes were clustered using Lasso2 (Lokhorst et al. 2021) and DEGreports v1.36.0 (Pantano 2023) and normalized read counts plotted using ggplot2 and pheatmaps v1.0.12 (Kolde 2019). Genes involved in Salicylic Acid biosynthesis/catabolism and response were determined based on a Wald Test of open plasmodesmata versus closed plasmodesmata with a lfc cut-off of +/-0.5 and adjusted p-value of 0.05 (Supplemental Data Set 12). Gene ontology enrichment analysis was conducted based on differentially expressed genes. Enrichment was conducted in TopGo v2.52.0 (Alexa and Rahnenfuhrer 2023) based on TAIR10 GO terms, with a topgoFisher cut-off of p<0.05. GO terms were plotted in ggplot2. GO parent terms were determined using the Revigo's (Supek et al. 2011) redundancy feature (cut-off 0.7) and utilized to extract defense-specific enriched GO terms.

Transgene Mapping

Transgenes were mapped by generating a pseudo-genome consisting of the *Arabidopsis* TAIR genome as well as the transgene of interest (mCherry) using SeqKit v0.9.1 (Shen et al. 2016) and STAR v2.5a (Dobin et al. 2013). The reads were then mapped and sorted by coordinates to the pseudo-genome using STAR v2.5a. The data was processed the same as previously described using Samtools and Stringtie. Gene counts of mCherry (Supplemental Data Set 13) and CalS3 (Supplemental Data Set 14) were extracted using DESeq2 and plotted in ggplot2.

 Size phenotyping

9-day-old seedlings of Col-0, LexA::PD-Plug and LexA:: icals3m were finely sprayed
with treatments DMSO or estradiol and 6 days later, the first two true leaves were
adhered to a microscope slide and images taken on a Leica M205 FA stereo
microscope. Leaf area was measured using FIJI.

Sugar and starch measurements

Leaf tissue was harvested in liquid nitrogen from plants 72 h post treatment of either
DMSO or estradiol (3 leaves from 5-week-old plants), with sugars and starch
separately extracted from approximately 100 mg total leaf tissue as described by
(Smith and Zeeman 2006). Frozen tissue was homogenized for 2 mins at 1200 rpm in
a GenoGrinder 2100. Samples were further homogenized in 1 mL of 0.7 M cold
perchloric acid and then centrifuge at max speed 4 °C for 5 min. The supernatant was
transferred to process for sugar, while the pellet was kept on ice for starch
quantification. For sugar, 600 µL supernatant was transferred to a new 1.5 mL tube,
and neutralization buffer (2 M KOH, 400 mM MES) was added to achieve a pH of 6-7.
The sample was then spun at max speed for 4 min and supernatant recovered. Soluble
sugars were quantified by enzymatic assay in technical triplicates by monitoring NADH
production absorbance at 340 nm in a 96-well format using an Omega FLUOstar
microplate reader. Each sample was added to 50 µL of buffer (50 mM HEPES NaOH
pH 7.4-7.6, 1 mM $MgCl_2$, 1mM ATP, 1 mM NAD, 1.4 U Hexokinase), topped with H_2O
to a total volume of 198 µL. An initial A_{340} reading was taken, before adding glucose-
6-phosphate dehydrogenase for a kinetic reading to determine glucose content,
followed by adding phosphoglucose isomerase to determine fructose content, followed
by invertase to determine sucrose content. To determine starch content, the sample
pellet was first washed by adding 1 mL H_2O and vortexing, followed by centrifugation
for 3 mins at 3,000 ×g. Supernatant was removed, then pellet washed again by adding
1 mL 80% ethanol and vortexing, followed by centrifugation for 3 mins at 3,000 ×g;
this ethanol wash was repeated for a total of three washes. Excess ethanol was
evaporated, then resuspended with 750 µL of H_2O. 2 × 200 µL of each sample was
transferred to a new screw cap tube and incubated at 95 °C for 12 minutes. The
samples were left to briefly cool to room temperature (~5 min), then one aliquot was
digested by incubation with α-amylase (10 U) and amyloglucosidase (1.26 U) in 0.1045

M sodium acetate buffer (pH 4.8) for two hours at 37 °C, alongside the other aliquot
serving as the non-digested control (i.e., containing only the 0.1045 M sodium acetate
buffer and two hour 37 °C incubation). Starch content was quantified through
enzymatic assay of the digested aliquot sample by monitoring NADH production as
per the above glucose measurement. Technical triplicates were performed for each
digested sample, alongside a single technical replicate for the non-digested sample.
Starch content was calculated from the glucose assay values in mg per gram of fresh
weight of sample.

Salicylic acid measurements

Leaves from 5-week-old plants were treated with DMSO or estradiol and harvested 72
h post treatment and flash frozen in liquid nitrogen; per biological replicate, a minimum
of six leaves from a minimum of three plants were pooled together to obtain between
150-220mg of fresh tissue weight. Frozen tissue was homogenized for 2×2 mins at
1200 rpm in a GenoGrinder 2100. 300 µL of buffer (10% methanol, 1% acetic acid in
H_2O) containing an internal standard of 1 µM salicylic acid-d_4 (SA-d_4, Merck) was
added, mixed well with a vortex then left to mix at 4 °C for 20 mins. Samples were
centrifuged for 25 mins at 4°C at 15,000 RCF, and the supernatant recovered. The
pellet was re-extracted with another 300 µL buffer without the internal standard, mixed
and centrifuged as before, with the resulting supernatant combined with the first
recovered supernatant. The total supernatant was centrifuged at 4 °C for 10 mins at
max speed, and each sample was then run for analysis. Quantification was performed
on an Acquity UPLC attached to a Xevo TQS tandem mass spectrometer (Waters).
Separation was on a 50×2.1mm 2.6µM Kinetex EVO C18 column (Phenomenex) using
the following gradient of acetonitrile (solvent B) versus 0.1% formic acid in water
(solvent A), run at 600 µL.min-1 and 35°C: 0 min, 10% B; 2 min, 80% B; 2.15 min,
80% B; 2.2 min, 10% B; 3.4 min, 10% B. The injection volume was 5 µL. Salicylate
was quantified by negative mode electrospray, monitoring the transition *m/z* 137→93
(and for the internal standard the corresponding transition 141→97) at 16V collision
energy. Spray chamber conditions were 1.5kV spray voltage, 500°C desolvation
temperature, 900 L.hr^{-1} desolvation gas, 150 L.hr^{-1} cone gas, and 7 bar nebulizer
pressure. Results were processed using TargetLynx software V4.1 SCN876 (2012).

Chlorophyll fluorescence measurements

Chlorophyll fluorescence measurements were performed using an Imaging-PAM
chlorophyll fluorometer and ImagingWin software application (Walz). The effective
PSII quantum yield as indicated by Y(II) was taken following a 30 min dark adaptation
of leaves from 5-week-old plants 72 h post DMSO or estradiol treatment. Four leaves
per treatment/genotype were measured in a given run, with 8 runs in total over two
independent experiments.

Statistical analyses

Statistical analyses were performed using RStudio 2021.09.01 Build 351/R version
4.0.3. Data presented as boxplots have the middle line indicating the median, the box
representing the upper and lower quartiles, and the whiskers indicating the minimum
and maximum values with 1.5× interquartile range. For bootstrap analyses, data were
analyzed using *medianBootstrap* (Johnston and Faulkner 2021) or *medianBootstrap2*
(Tee et al. 2023a). Pathoassays, basal ROS quantification, and chlorophyll
fluorescence measurements were analyzed using a linear-mixed effects model using
the R package, *lmerTest*. For pathoassays the random effect was 'independent
experiment repeat'; for the basal ROS quantification the random effect was 'biological
replicate'; for chlorophyll fluorescence measurements the random effect was
'experimental run'. Size phenotyping, starch, sucrose, glucose and fructose
measurements were analyzed using a linear model. All ANOVAs specified are ANOVA
Satterthwaite's Method, and significant differences between factors were determined
by post hoc Tukey HSD using the R package, *emmeans*. Salicylic acid quantification
was analyzed using a Mann-Whitney test.

Data availability

The original images for callose and GFP mobility quantification for Fig. 1 will be
deposited on the Bioimage Archive (https://www.ebi.ac.uk/bioimage-archive/). Raw
counts for bombardment data generated for this study (Fig. 1C and Supplemental Fig.
S3) is available in Supplemental Data Set 1. Additional raw counts for bombardment
data for Supplemental Fig. S3 are sourced from Cheval et al. 2020 and Tee et al.
2023b. RNAseq data has been deposited in the Gene Expression Omnibus under the
accession code GSE248301, with further analysis included in Supplemental Data Sets
as specified in-text.

Supplemental Data Sets

Supplemental Data Set 1: Data for number of cells showing GFP within a genotype/treatment

Supplemental Data Set 2: Background response to estradiol: differentially regulated genes in response to estradiol treatment shared across all genotypes

Supplemental Data Set 3: Gene count matrix from the RNAseq analysis

Supplemental Data Set 4: Genes differentially expressed in category specified, identified by a likelihood ratio test analysis

Supplemental Data Set 5: Genes differentially expressed in response to 12h plasmodesmal closure identified by a likelihood ratio test analysis

Supplemental Data Set 6: Genes differentially expressed in response to 24h plasmodesmal closure identified by a likelihood ratio test analysis

Supplemental Data Set 7: Genes differentially expressed in response to 48 h plasmodesmal closure identified by a likelihood ratio test analysis

Supplemental Data Set 8: Genes differentially expressed in response to 72h plasmodesmal closure identified by a likelihood ratio test analysis

Supplemental Data Set 9: Genes identified from the literature as associated with salicylic acid biosynthesis, catabolism and response

Supplemental Data Set 10: Genes identified from the literature as associated with the synthesis and transport of soluble sugars and starch

Supplemental Data Set 11: Metadata related to the gene count matrix

Supplemental Data Set 12: Log2 fold change of genes associated with SA that are differentially expressed at 12h, 24h, 48h and 72h

Supplemental Data Set 13: Gene counts of mCherry from pseudo-genome GEO GSE248301

Supplemental Data Set 14: Gene counts of CalS3 from pseudo-genome GEO GSE248301

Author Contributions

CREDIT authorship breakdown:

Conceptualization – all authors

Data curation – EET, ABr, DP, HRT

Formal analysis – EET, ABr, DP, HRT

Funding acquisition – CF

Investigation – EET, ABr, DP, HRT, CW, ABe

Methodology – EET, ABr, DP, HRT, CW, ABe, CF

Project administration – CF

Resources – DP, ABr, CF

Supervision – CF

Validation – EET, ABr, DP, HRT

Visualization – EET, ABr, DP, HRT

Writing – original draft – EET, HRT, CF

Writing – review & editing – EET, ABr, DP, HRT, CW, ABe, CF

Disclosure and competing interests statement

The authors declare no competing interests.

Acknowledgements

We acknowledge access to the John Innes Centre Biomolecular Analysis Facility and thank Dr. Lionel Hill for their assistance and training with mass spectrometry, as well as access to the John Innes Centre Bioimaging Facility. We also acknowledge the Seung lab for technical assistance in the sugar and starch measurements.

Work in the Faulkner lab is funded by the Biotechnology and Biological Science Research Council (Grants: BB/X010996/1 and BB/X007685/1) and the European Research Council (Grant: 725459 - "INTERCELLAR).

References

Abt MR and Zeeman SC. Evolutionary innovations in starch metabolism. Curr Opin Plant Biol. 2020:55:109–117. https://doi.org/10.1016/J.PBI.2020.03.001

Alazem M and Burch-Smith TM. Roles of ROS and redox in regulating cell-to-cell communication: Spotlight on viral modulation of redox for local spread. Plant Cell Environ. 2024. https://doi.org/10.1111/PCE.14805

Alexa A and Rahnenfuhrer J. topGO: Enrichment Analysis for Gene Ontology. 2023.

Araya T, Noguchi K, and Terashima I. Effects of Carbohydrate Accumulation on Photosynthesis Differ between Sink and Source Leaves of Phaseolus vulgaris L. Plant Cell Physiol. 2006:47(5):644–652. https://doi.org/10.1093/PCP/PCJ033

Aung K, Kim P, Li Z, Joe A, Kvitko B, Alfano JR. and He SY. 2020. Pathogenic bacteria target plant plasmodesmata to colonize and invade surrounding tissues. *The Plant Cell*, *32*(3), pp.595-611. https://doi.org/10.1105/tpc.19.00707

Babraham Bioinformatics - FastQC A Quality Control tool for High Throughput Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/. Retrieved July 28, 2023

Bavnhøj L, Driller JH, Zuzic L, Stange AD, Schiøtt B, and Pedersen BP. Structure and sucrose binding mechanism of the plant SUC1 sucrose transporter. Nature Plants 2023 9:6. 2023:9(6):938–950. https://doi.org/10.1038/s41477-023-01421-0

Beaugelin I, Chevalier A, d'Alessandro S, Ksas B, and Havaux M. (2020). Endoplasmic reticulum-mediated unfolded protein response is an integral part of singlet oxygen signalling in plants. The Plant Journal, 102(6), 1266-1280. https://doi.org/10.1111/tpj.14700.

Bellandi A, Papp D, Breakspear A, Joyce J, Johnston MG, de Keijzer J, Raven EC, Ohtsu M, Vincent TR, Miller AJ, et al. Diffusion and bulk flow of amino acids mediate calcium waves in plants. Sci Adv. 2022:8(42):6693. DOI: 10.1126/sciadv.abo6693.

van Butselaar T and Van den Ackerveken G. Salicylic Acid Steers the Growth–Immunity Tradeoff. Trends Plant Sci. 2020:25(6):566–576. https://doi.org/10.1016/J.TPLANTS.2020.02.002

Büttner M and Sauer N. Monosaccharide transporters in plants: structure, function and physiology. Biochimica et Biophysica Acta (BBA) - Biomembranes. 2000:1465(1–2):263–274. https://doi.org/10.1016/S0005-2736(00)00143-7

Caillaud MC, Wirthmueller L, Sklenar J, Findlay K, Piquerez SJM, Jones AME, Robatzek S, Jones JDG, and Faulkner C. The Plasmodesmal Protein PDLP1 Localises to Haustoria-Associated Membranes during Downy Mildew Infection and Regulates Callose Deposition. PLoS Pathog. 2014:10(11):e1004496. https://doi.org/10.1371/JOURNAL.PPAT.1004496

Chen LQ, Hou BH, Lalonde S, Takanaga H, Hartung ML, Qu XQ, Guo WJ, Kim JG, Underwood W, Chaudhuri B, et al. Sugar transporters for intercellular exchange and nutrition of pathogens. Nature 2010 468:7323. 2010:468(7323):527–532. https://doi.org/10.1038/nature09606

Chen LQ, Lin IW, Qu XQ, Sosso D, McFarlane HE, Londoño A, Samuels AL, and Frommer WB. A Cascade of Sequentially Expressed Sucrose Transporters in the Seed Coat and Endosperm Provides Nutrition for the Arabidopsis Embryo. Plant Cell. 2015:27(3):607–619. https://doi.org/10.1105/TPC.114.134585

Cheval C, Samwald S, Johnston MG, Keijzer J de, Breakspear A, Liu X, Bellandi A, Kadota Y, Zipfel C, and Faulkner C. Chitin perception in plasmodesmata characterizes submembrane immune-signaling specificity in plants. Proc Natl Acad Sci U S A. 2020:117(17):9621–9629. https://doi.org/10.1073/PNAS.1907799117

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, and Davies RM. Twelve years of SAMtools and BCFtools. Gigascience. 2021:10(2):1–4. https://doi.org/10.1093/GIGASCIENCE/GIAB008

David LC, Lee SK, Bruderer E, Abt MR, Fischer-Stettler M, Tschopp MA, Solhaug EM, Sanchez K, and Zeeman SC. BETA-AMYLASE9 is a plastidial nonenzymatic regulator of leaf starch degradation. Plant Physiol. 2022:188(1):191–207. https://doi.org/10.1093/PLPHYS/KIAB468

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, and Gingeras TR. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 2013:29(1):15–21. https://doi.org/10.1093/BIOINFORMATICS/BTS635

El-Oirdi M, El-Rahman TA, Rigano L, El-Hadrami A, Rodriguez MC, Daayf F, Vojnov A, and Bouarab K. Botrytis cinerea Manipulates the Antagonistic Effects between Immune Pathways to Promote Disease Development in Tomato. Plant Cell. 2011:23(6):2405. https://doi.org/10.1105/TPC.111.083394

Ewels P, Magnusson M, Lundin S, and Käller M. MultiQC: summarize analysis results for multiple tools and samples in a single report. Bioinformatics. 2016:32(19):3047–3048. https://doi.org/10.1093/BIOINFORMATICS/BTW354

Faulkner C, Petutschnig E, Benitez-Alfonso Y, Beck M, Robatzek S, Lipka V, and Maule AJ. LYM2-dependent chitin perception limits molecular flux via

plasmodesmata. Proc Natl Acad Sci U S A. 2013:110(22):9166–9170.
https://doi.org/10.1073/PNAS.1203458110/-/DCSUPPLEMENTAL

Ferrari S, Plotnikova JM, De Lorenzo G, and Ausubel FM. Arabidopsis local resistance
to Botrytis cinerea involves salicylic acid and camalexin and requires EDS4 and
PAD2, but not SID2, EDS5 or PAD4. The Plant Journal. 2003:35(2):193–205.
https://doi.org/10.1046/J.1365-313X.2003.01794.X

German L, Yeshvekar R, and Benitez-Alfonso Y. Callose metabolism and the
regulation of cell walls and plasmodesmata during plant mutualistic and
pathogenic interactions. Plant Cell Environ. 2023:46(2):391–404.
https://doi.org/10.1111/PCE.14510

Hellemans J, Mortier G, De Paepe A, Speleman F, and Vandesompele J. qBase
relative quantification framework and software for management and automated
analysis of real-time quantitative PCR data. Genome Biol. 2008:8(2):1–14.
https://doi.org/10.1186/GB-2007-8-2-R19/COMMENTS

Howlader P, Bose SK, Jia X, Zhang C, Wang W, and Yin H. Oligogalacturonides
induce resistance in Arabidopsis thaliana by triggering salicylic acid and
jasmonic acid pathways against Pst DC3000. Int J Biol Macromol.
2020:164:4054–4064. https://doi.org/10.1016/J.IJBIOMAC.2020.09.026

Johnson DA, Hill JP, and Thomas MA. The monosaccharide transporter gene family
in land plants is ancient and shows differential subfamily expression and
expansion across lineages. BMC Evol Biol. 2006:6(1):1–20.
https://doi.org/10.1186/1471-2148-6-64/FIGURES/10

Johnston MG, Bellandi A, and Faulkner C. faulknerfalcons/PD_detection: Toolkit for
quantification of callose at plasmodesmata. 2022.
https://doi.org/10.5281/ZENODO.6583765

Johnston MG and Faulkner C. A bootstrap approach is a superior statistical method
for the comparison of non-normal data with differing variances. New
Phytologist. 2021:230(1):23–26. https://doi.org/10.1111/NPH.17159

Kim D, Paggi JM, Park C, Bennett C, and Salzberg SL. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol.
2019:37(8):907–915. https://doi.org/10.1038/S41587-019-0201-4

Kolde R. Package "pheatmap." 2019.

Kraner ME, Link K, Melzer M, Ekici AB, Uebe S, Tarazona P, Feussner I, Hofmann J,
and Sonnewald U. Choline transporter-like1 (CHER1) is crucial for
plasmodesmata maturation in Arabidopsis thaliana. The Plant Journal.
2017:89(2):394–406. https://doi.org/10.1111/TPJ.13393

Kronberg K, Vogel F, Rutten T, Hajirezaei MR, Sonnewald U, and Hofius D. The Silver
Lining of a Viral Agent: Increasing Seed Yield and Harvest Index in Arabidopsis
by Ectopic Expression of the Potato Leaf Roll Virus Movement Protein. Plant
Physiol. 2007:145(3):905–918. https://doi.org/10.1104/PP.107.102806

Krueger F, James F, Ewels P, Afyounian E, and Schuster-Boeckler B.
FelixKrueger/TrimGalore: v0.6.7 - DOI via Zenodo. 2021.
https://doi.org/10.5281/ZENODO.5127899

Lee J-Y, Wang X, Cui W, Sager R, Modla S, Czymmek K, Zybaliov B, Wijk K van,
Zhang C, Lu H, et al. A Plasmodesmata-Localized Protein Mediates Crosstalk
between Cell-to-Cell Communication and Innate Immunity in Arabidopsis. Plant
Cell. 2011:23(9):3353–3373. https://doi.org/10.1105/TPC.111.087742

Levy A, Erlanger M, Rosenthal M, and Epel BL. A plasmodesmata-associated beta-
1,3-glucanase in Arabidopsis. Plant J. 2007:49(4):669–682.
https://doi.org/10.1111/J.1365-313X.2006.02986.X

Li Z, Variz H, Chen Y, Liu SL, Aung K. Plasmodesmata-Dependent Intercellular
Movement of Bacterial Effectors. Front Plant Sci. 2021 Mar 22;12:640277. doi:
10.3389/fpls.2021.640277. PMID: 33959138; PMCID: PMC8095247.

Li Z, Liu SL, Montes-Serey C, Walley JW, Aung K. PLASMODESMATA-LOCATED
PROTEIN 6 regulates plasmodesmal function in Arabidopsis vasculature. Plant
Cell. 2024 Jun 6:koae166. doi: 10.1093/plcell/koae166. Epub ahead of print.
PMID: 38842334.

Liu YH, Song YH, and Ruan YL. Sugar conundrum in plant–pathogen interactions:
roles of invertase and sugar transporters depend on pathosystems. J Exp Bot.
2022:73(7):1910. https://doi.org/10.1093/JXB/ERAB562

Lokhorst J, Venables B, and Turlach B. R package "lasso2." 2021.
https://github.com/cran/lasso2. Retrieved July 28, 2023

Love MI, Huber W, and Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014:15(12):1–21.
https://doi.org/10.1186/S13059-014-0550-8/FIGURES/9

Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet J. 2011:17(1):10–12. https://doi.org/10.14806/EJ.17.1.200

Monroe JD and Storm AR. Review: The Arabidopsis β-amylase (BAM) gene family:
Diversity of form and function. Plant Science. 2018:276:163–170.
https://doi.org/10.1016/J.PLANTSCI.2018.08.016

Ohtsu M, Jennings J, Johnston M, Breakspear A, Liu X, Stark K, Morris RJ, de Keijzer
J, Faulkner C. Assaying Effector Cell-to-Cell Mobility in Plant Tissues Identifies
Hypermobility and Indirect Manipulation of Plasmodesmata. Mol Plant Microbe
Interact. 2024 Feb;37(2):84-92. doi: 10.1094/MPMI-05-23-0052-TA. Epub 2024
Feb 26. PMID: 37942798.

Pantano L. DEGreport: Report of DEG analysis. 2023.
https://doi.org/10.18129/B9.bioc.DEGreport

Paterlini A, Dorussen D, Fichtner F, van Rongen M, Delacruz R, Vojnović A, Helariutta
Y and Leyser O. (2021), Callose accumulation in specific phloem cell types

reduces axillary bud growth in *Arabidopsis thaliana*. New Phytol, 231: 516-523. https://doi.org/10.1111/nph.17398

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, and Salzberg SL. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nature Biotechnology 2015 33:3. 2015:33(3):290–295. https://doi.org/10.1038/nbt.3122

Preiser AL, Banerjee A, Weise SE, Renna L, Brandizzi F, and Sharkey TD. Phosphoglucoisomerase Is an Important Regulatory Enzyme in Partitioning Carbon out of the Calvin-Benson Cycle. Front Plant Sci. 2020:11:580726. https://doi.org/10.3389/FPLS.2020.580726/BIBTEX

R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2022. https://www.R-project.org/

Ross-Elliott TJ, Jensen KH, Haaning KS, Wager BM, Knoblauch J, Howell AH, Mullendore DL, Monteith AG, Paultre D, Yan D, Otero S, Bourdon M, Sager R, Lee JY, Helariutta Y, Knoblauch M, Oparka KJ (2017) Phloem unloading in Arabidopsis roots is convective and regulated by the phloem-pole pericycle eLife 6:e24125 https://doi.org/10.7554/eLife.24125

Ruan YL, Llewellyn DJ, and Furbank RT. The Control of Single-Celled Cotton Fiber Elongation by Developmentally Reversible Gating of Plasmodesmata and Coordinated Expression of Sucrose and K+ Transporters and Expansin. Plant Cell. 2001:13(1):47–60. https://doi.org/10.1105/TPC.13.1.47

Seo PJ, Park JM, Kang SK, Kim SG, and Park CM. An Arabidopsis senescence-associated protein SAG29 regulates cell viability under high salinity. Planta. 2011:233(1):189–200. https://doi.org/10.1007/S00425-010-1293-8

Sevilem I, Miyashima S, and Helariutta Y. (2012). Cell-to-cell communication via plasmodesmata in vascular plants. *Cell Adhesion & Migration*, *7*(1), 27–32. https://doi.org/10.4161/cam.22126

Seyfferth C, Tsuda K, Lu H, and Grant M. Salicylic acid signal transduction: the initiation of biosynthesis, perception and transcriptional reprogramming. 2014. https://doi.org/10.3389/fpls.2014.00697

Shen W, Le S, Li Y, and Hu F. SeqKit: A Cross-Platform and Ultrafast Toolkit for FASTA/Q File Manipulation. PLoS One. 2016:11(10):e0163962. https://doi.org/10.1371/JOURNAL.PONE.0163962

Smith AM and Zeeman SC. Quantification of starch in plant tissues. Nature Protocols 2006 1:3. 2006:1(3):1342–1345. https://doi.org/10.1038/nprot.2006.232

Smith AM and Zeeman SC. Starch: A Flexible, Adaptable Carbon Store Coupled to Plant Growth. https://doi.org/101146/annurev-arplant-050718-100241. 2020:71:217–245. https://doi.org/10.1146/ANNUREV-ARPLANT-050718-100241

Supek F, Bošnjak M, Škunca N, and Šmuc T. REVIGO Summarizes and Visualizes
Long Lists of Gene Ontology Terms. PLoS One. 2011:6(7):e21800.
https://doi.org/10.1371/JOURNAL.PONE.0021800

Tee EE, Johnston MG, and Faulkner C. faulknerfalcons/Tee-etal-2023. 2023a.
https://doi.org/10.5281/ZENODO.7680790

Tee EE, Johnston MG, Papp D, and Faulkner C. A PDLP-NHL3 complex integrates
plasmodesmal immune signaling cascades. Proceedings of the National
Academy of Sciences. 2023b:120(17):e2216397120.
https://doi.org/10.1073/PNAS.2216397120

Tee EE, Samwald S, and Faulkner C. Quantification of Cell-to-Cell Connectivity Using
Particle Bombardment. 2022:263–272. https://doi.org/10.1007/978-1-0716-
2132-5_17

Thomas CL, Bayer EM, Ritzenthaler C, Fernandez-Calvino L, and Maule AJ. Specific
Targeting of a Plasmodesmal Protein Affecting Cell-to-Cell Communication.
PLoS Biol. 2008:6(1):e7. https://doi.org/10.1371/JOURNAL.PBIO.0060007

Tomczynska I, Stumpe M, Doan TG, Mauch F. A Phytophthora effector protein
promotes symplastic cell-to-cell trafficking by physical interaction with
plasmodesmata-localised callose synthases. New Phytol. 2020
Sep;227(5):1467-1478. doi: 10.1111/nph.16653. Epub 2020 Jun 8. PMID:
32396661.

Vatén A, Dettmer J, Wu S, Stierhof YD, Miyashima S, Yadav SR, Roberts CJ,
Campilho A, Bulone V, Lichtenberger R, et al. Callose Biosynthesis Regulates
Symplastic Trafficking during Root Development. Dev Cell. 2011:21(6):1144–
1155. https://doi.org/10.1016/j.devcel.2011.10.006

Velásquez AC, Oney M, Huot B, Xu S, and He SY. Diverse mechanisms of resistance
to Pseudomonas syringae in a thousand natural accessions of Arabidopsis
thaliana. New Phytologist. 2017:214(4):1673–1687.
https://doi.org/10.1111/NPH.14517

Vlot AC, Liu PP, Cameron RK, Park SW, Yang Y, Kumar D, Zhou F, Padukkavidana
T, Gustafsson C, Pichersky E, et al. Identification of likely orthologs of tobacco
salicylic acid-binding protein 2 and their role in systemic acquired resistance in
Arabidopsis thaliana. The Plant Journal. 2008:56(3):445–456.
https://doi.org/10.1111/J.1365-313X.2008.03618.X

Wang C, Dai S, Zhang ZL, Lao W, Wang R, Meng X, and Zhou X. Ethylene and
salicylic acid synergistically accelerate leaf senescence in Arabidopsis. J Integr
Plant Biol. 2021a:63(5):828–833.
https://doi.org/10.1111/JIPB.13075/SUPPINFO

Wang X, Sager R, Cui W, Zhang C, Lu H, and Lee JY. Salicylic Acid Regulates
Plasmodesmata Closure during Innate Immune Responses in Arabidopsis.
Plant Cell. 2013:25(6):2315. https://doi.org/10.1105/TPC.113.110676

Wang Y, Li X, Fan B, Zhu C, and Chen Z. Regulation and Function of Defense-Related
1030   Callose Deposition in Plants. Int J Mol Sci. 2021b:22(5):1–15.
1031   https://doi.org/10.3390/IJMS22052393

1032   Wickham H. ggplot2: Elegant Graphics for Data Analysis (Spring-Verlag New York).

1033   Wickham H, Averick M, Bryan J, Chang W, D' L, Mcgowan A, François R, Grolemund
1034   G, Hayes A, Henry L, et al. Welcome to the Tidyverse. J Open Source Softw.
1035   2019:4(43):1686. https://doi.org/10.21105/JOSS.01686

1036   Wildermuth MC, Dewdney J, Wu G, and Ausubel FM. Isochorismate synthase is
1037   required to synthesize salicylic acid for plant defence. Nature 2001 414:6863.
1038   2001:414(6863):562–565. https://doi.org/10.1038/35107108

1039   Williams LE, Lemoine R, and Sauer N. Sugar transporters in higher plants – a diversity
1040   of roles and complex regulation. Trends Plant Sci. 2000:5(7):283–290.
1041   https://doi.org/10.1016/S1360-1385(00)01681-2

1042   Wilson DC, Kempthorne CJ, Carella P, Liscombe DK, and Cameron RK. Age-related
1043   resistance in arabidopsis thaliana involves the MADS-domain transcription
1044   factor SHORT VEGETATIVE PHASE and direct action of salicylic acid on
1045   pseudomonas syringae. Molecular Plant-Microbe Interactions.
1046   2017:30(11):919–929. https://doi.org/10.1094/MPMI-07-17-0172-
1047   R/ASSET/IMAGES/LARGE/MPMI-07-17-0172-R_F8.JPEG

1048   Wu S, O'Lexy R, Xu M, Sang Y, Chen X, Yu Q, Gallagher KL. Symplastic signaling
1049   instructs cell division, cell expansion, and cell polarity in the ground tissue of
1050   Arabidopsis thaliana roots. Proc Natl Acad Sci U S A. 2016 Oct
1051   11;113(41):11621-11626. doi: 10.1073/pnas.1610358113. Epub 2016 Sep 23.
1052   PMID: 27663740; PMCID: PMC5068303.

1053   Xu B, Cheval C, Laohavisit A, Hocking B, Chiasson D, Olsson TSG, Shirasu K,
1054   Faulkner C, and Gilliham M. A calmodulin-like protein regulates plasmodesmal
1055   closure during bacterial immune responses. New Phytologist. 2017:215:77–84.
1056   https://doi.org/10.1111/nph.14599

1057   Yadav SR, Yan D, Sevilem I, Helariutta Y. Plasmodesmata-mediated intercellular
1058   signaling during plant growth and development. Front Plant Sci. 2014 Feb
1059   17;5:44. doi: 10.3389/fpls.2014.00044. PMID: 24596574; PMCID:
1060   PMC3925825.

1061   Yang L, Li B, Zheng XY, Li J, Yang M, Dong X, He G, An C, and Deng XW. Salicylic
1062   acid biosynthesis is enhanced and contributes to increased biotrophic pathogen
1063   resistance in Arabidopsis hybrids. Nature Communications 2015 6:1.
1064   2015:6(1):1–12. https://doi.org/10.1038/ncomms8309

1065   Zhang Z, Ruan YL, Zhou N, Wang F, Guan X, Fang L, Shang X, Guo W, Zhu S, and
1066   Zhang T. Suppressing a Putative Sterol Carrier Gene Reduces Plasmodesmal
1067   Permeability and Activates Sucrose Transporter Genes during Cotton Fiber
1068   Elongation. Plant Cell. 2017:29(8):2027–2046.
1069   https://doi.org/10.1105/TPC.17.00358

Zhongpeng L, Liu SL, Montes-Serey C, Walley JW, and Aung K. PLASMODESMATA-
LOCATED PROTEIN 6 regulates plasmodesmal function in Arabidopsis
vasculature. Plant Cell. 2024 koae166. https://doi.org/10.1093/plcell/koae166

Zuo J, Niu QW, and Chua NH. Technical advance: An estrogen receptor-based
transactivator XVE mediates highly inducible gene expression in transgenic
plants. Plant J. 2000:24(2):265–273. https://doi.org/10.1046/J.1365-
313X.2000.00868.X

1077

Figures

1079

Figure 1. Characterization of plasmodesmal status in induced LexA::PD-Plug and LexA::icals3m tissue. A) Quantification of aniline blue-stained plasmodesmata-associated callose in 5-week-old plants of LexA::PD-Plug (PD-Plug) or LexA::icals3m (icals3m) 24 h post DMSO or estradiol treatment. Datapoints represent the average of the total PD intensity/plasmodesmata (PD) per image, with n ≥ 14 per genotype/treatment. Bootstrap analysis indicates significant differences between DMSO and estradiol treatment in PD-Plug and icals3m, as indicated by *** $p<0.001$. **B)** Example micrographs of bombardment sites showing movement of GFP from transformed cells indicate cell-to-cell connectivity of uninduced (DMSO) versus restricted cell connectivity in induced (estradiol) leaves of PD-Plug and icals3m. **C)** Quantification of microprojectile bombardment data showing GFP movement into neighboring cells. Leaves of 5-week-old plants were bombarded after 24 h treatment of DMSO or estradiol; n ≥ 62 bombardment sites. Bootstrap analysis with significant differences indicated by a, b and c, with $p<0.01$. Scale bar = 200 µM.

1094

1095

Figure 2. Plasmodesmal closure triggers stress transcriptional responses and salicylic
acid related changes. A) Likelihood ratio test analysis determined number (indicated by n)
of differentially expressed genes (DEGs) significantly differed between plasmodesmal status
(i.e., PD State closed or open), and matches the cluster pattern of upregulated in closed state
in comparison to open state at 12, 24, 48 and 72 h post treatment. Hierarchical clustering
grouped together plasmodesmal state of open (i.e., estradiol treated LexA::PD-Plug and
LexA::icals3m) and closed (DMSO treated LexA::PD-Plug, LexA::icals3m and Col-0, and
estradiol treated Col-0). Expression normalized by row (z-score). **B)** Top 15 defense GO terms
enriched over time when plasmodesmata are closed. **C)** Expression of genes related to the
salicylic acid biosynthesis/catabolism and response over time when plasmodesmata are
closed (i.e., in estradiol treated LexA::PD-Plug and LexA::icals3m). **D)** Quantification of
salicylic acid in 5-week-old plants after 72 h DMSO or estradiol treatment, n ≥ 6 per
genotype/treatment. Significant differences between treatment within a genotype analyzed
using Mann-Whitney test denoted by **, $p < 0.01$, or ***, $p < 0.001$.

1110

Figure 3. Plasmodesmal closure differentially impacts pathogen resistance and immune responses. A) Bacterial growth ($\log_{10}$ CFU/cm^2 leaf area) of *Pst* DC3000 3 dpi on uninduced and induced tissue of 4–5-week-old plants of Col-0, LexA::PD-Plug (PD-Plug) and LexA::icals3m (icals3m). Tissue was treated with DMSO (uninduced) or estradiol (induced) 72 h prior to infection. 4 independent experiments were conducted, with 5 plants per treatment/genotype; N = 20. Independent factors genotype and treatment were significant, with a significant interaction between the two (ANOVA; F = 16.6, df = 2, P < 0.0001; F = 67.0, df = 1, P < 0.0001; F = 8.2, df = 2, P < 0.001). Significant differences denoted by a and b, p < 0.0001. **B)** The average area of *Botrytis cinerea* disease lesions 2 dpi in leaves of 5-week-old plants of induced and uninduced Col-0, LexA::PD-Plug and LexA::icals3m. Tissue was treated with DMSO (uninduced) or estradiol (induced) 72 h prior to infection, with N = 16-17 for all genotypes/treatment combined from three independent repeats. Independent factors genotype and treatment were significant with a significant interaction between the two (ANOVA; F = 24.4, df = 2, P < 0.0001; F = 52.7, df = 1, P < 0.0001; F = 57.0, df = 2, P < 0.0001). Significant differences denoted by a and b, p < 0.0001. **C)** Western blot detection of MAPK-activation by flg22 treatment in 4–5-week-old plants of Col-0, LexA::PD-Plug and LexA::icals3m 72 h post DMSO or estradiol treatment. Minute timings of tissue harvest are denoted, and Coomassie blue staining shown below as loading controls. **D)** Average relative light units (RLU) generated from ROS production in response to flg22 in leaf discs of 5-week-old plants. Leaves were treated with estradiol or DMSO 72 h prior to flg22 treatment and data is combined from two independent experiment with N = 95 per genotype/treatment. Significant differences between treatment determined by bootstrap analysis, with significant differences denoted by **, p < 0.01. **E)** Average RLU generated from ROS production in response to flg22 in seedlings. Plants were treated with estradiol or DMSO 72 h prior to flg22

1137   treatment, with N ≥ 80 per genotype/treatment. Significant differences between
1138   treatment determined by bootstrap analysis, with significant differences denoted by **,
1139 $p < 0.01$.

1140   .

1141

1142

Figure 4. Plasmodesmal closure induces sugar accumulation and impairs leaf growth. A) Leaf area of first two true leaves 6 days post treatment (DMSO or estradiol). Independent factors genotype and treatment were significant, with a significant interaction between the two (ANOVA; F = 82.4, df = 2, P < 0.0001; F = 445.0, df = 1, P < 0.0001; F = 104.2, df = 2, P < 0.0001). Significant differences denoted by a, b and c, p < 0.0001. n = 40. **B)** Starch, sucrose, glucose and fructose content in leaves of 5-week-old plants 72 h post DMSO or estradiol treatment. Independent factors genotype and treatment were significant, with a significant interaction between the two for each metabolite (ANOVA; *starch* F = 4.7, df = 2, P < 0.05, F = 9.2, df = 1, P < 0.05; F = 11.4, df = 2, P < 0.01; *sucrose* F = 26.2, df = 2, P < 0.0001; F = 25.8, df = 1, P < 0.001; F = 26.7, df = 2, P < 0.0001; *glucose* F = 70.0, df = 2, P < 0.0001; F = 74.4, df = 1, P < 0.0001; F = 79.3, df = 2, P < 0.0001; *fructose* F = 28.0, df = 2, P < 0.0001; F = 30.4, df = 1, P < 0.0001; F = 28.8, df = 2, P < 0.0001). Significant differences denoted by a and b, p < 0.01.

1157

Tables

Table 1. Photosynthetic yield as indicated by Y(II) following DMSO or estradiol
treatment of leaves of Col-0, LexA::PD-Plug (PD-Plug) and LexA::icals3m
(icals3m). Pooled average of Y(II) ± SD, with a biological replicate data averaged from
4 leaves per treatment/genotype on a given experimental run where N = 8, combined
from two independent experiments. Independent factors genotype and treatment were
not significant, with no significant interaction between the two (ANOVA; F = 1.3, df =
2, P = 0.29; F = 1.2, df = 1, P = 0.29, F = 0.1, df = 2, P = 0.92).

	Col-0	PD-Plug	icals3m
DMSO	0.79±0.02	0.78±0.02	0.78±0.02
estradiol	0.79±0.01	0.78±0.01	0.79±0.01

1166

1167