Oncogenic Ras-driven Dorsal/NF-κB signaling contributes to tumorigenesis in a *Drosophila* carcinoma model

Caroline Dillard[1,2], Jose Teles Reis[1,2], Ashish Jain, Roland Le Borgne[3], Heinrich Jasper[4],

Tor Erik Rusten [1,2]☒

[1]Center for Cancer Cell Reprogramming, Institute of Clinical Medicine, Faculty of Medicine,

University of Oslo, Oslo, Norway

[2]Department of Molecular Cell Biology, Institute for Cancer Research, Oslo University Hospital,

Oslo, Norway

[3] Univ Rennes, CNRS-UMR 6290, Institut de Génétique et Développement de Rennes, Rennes,

France

[4]Buck Institute for Research on Aging, 8001 Redwood Boulevard, Novato, CA 94945-1400,

USA

*Corresponding authors: t.e.rusten@medisin.uio.no, cmdillar@uio.no

Abstract:

Cancer-driving mutations synergize with inflammatory stress signaling pathways during carcinogenesis. *Drosophila melanogaster* tumour models are increasingly recognized as models to inform conserved molecular mechanisms of tumorigenesis with both local and systemic effects of cancer. Although initial discoveries of the Toll-NFκB signaling pathway in development and immunity was pioneered in *Drosophila*, limited information is available for its role in cancer progression. Using a well-studied cooperative Ras^{V12}-driven epithelial-derived tumour model, we here describe functions of Toll- NF-κB signaling in malignant *Ras^{V12}, scrib$^-$* tumors. The extracellular Toll pathway components ModSP and PGRP-SA and intracellular signaling Kinase, Pelle/IRAK, are rate-limiting for tumor growth. The Toll pathway NFκB protein Dorsal, as well as *cactus/IκB* show elevated expression in tumors with highest expression in invasive cell populations. Oncogenic Ras^{V12}, and not loss of *scribble,* confers increased expression and heterogenous distribution of two Dorsal isoforms, DorsalA and DorsalB in different tumour cell populations. Mechanistic analyses demonstrates that Dorsal drives growth and malignancy by suppressing differentiation, counteracting apoptosis and promoting invasion of *Ras^{V12}, scrib$^-$* tumors genetically dependent on *twist* and *snail*.

Introduction:

Inflammation, defined as the body's response to harmful external inputs, is a hallmark of cancer [1]. This response is mediated by four major inflammatory pathways: the MAPK, PI3K-AKT, JAK-STAT and NF-κB (canonical and non-canonical) signaling pathways which integrate intra- and extracellular alarms and trigger cellular responses [2]. NF-κB activation has been observed in many cancers [3, 4] and correlates with a poor prognosis in breast and non-small cell lung cancer

[5, 6]. NF-κB functions appear cancer-relevant, as a significant body of *in vitro* studies reports its

involvement in cellular processes such as invasion [7-9], proliferation [10, 11], stemness [12],

survival [10-12] and chemoresistance [12-15]. Similarly, a limited number of human cancer

xenograft studies demonstrated a positive effect of NF-κB on cancer proliferation [11], survival

[11] and metastatic behavior [8, 9].

Despite several attempts, pharmaceutical targeting of NF-κB in cancer has proven

ineffective due to toxicity resulting from its essential role in regulating cellular and innate immunity

at the organismal level [16, 17]. Therefore, understanding both the upstream and downstream

mechanisms governing NF-κB activation and functions in cancer, as well as deciphering whether

such mechanisms are specific to cancer cells, is crucial. This understanding may facilitate the

design of NF-κB -targeting cancer therapies that spare healthy cells. *In vivo* studies would be

informative for achieving this goal.

Drosophila melanogaster is increasingly being used to study both cancer cell- autonomous

mechanisms as well as the biology of complex local and systemic paraneoplastic effects

instigated by tumor growth [18]. Among the *Drosophila melanogaster* cancer models, the *Ras^{V12},*

scrib$^{-/-}$ carcinoma model, arising from the concomitant expression of the constitute activated

oncogenic form of the KRAS/NRAS orthologue, Ras85D (here referred to as Ras^{V12}), and the loss

of cell polarity through *scribble* loss-of-function, has been most extensively studied and displays

a remarkable array of hallmarks similar to human cancer [19] [20]. In *Ras^{V12}, scrib$^{-/-}$* tumors,

generated in the larval eye-antennal disc epithelium (EAD), the evolutionarily conserved Ras-

MAPK, PI3K-AKT and JAK-STAT inflammatory pathways all constitute major drivers [19, 20].

Surprisingly, at the organismal level, *Ras^{V12}, scrib$^{-/-}$* tumour-bearing flies also display both

microenvironmental and systemic alterations that mimic human cancer, such as wasting of

adipose and muscle tissues, gut atrophy, microenvironmental extracellular matrix remodeling and

immune cell infiltration [21-24]. Although the discovery and functions of the Toll-NF-κB and IMD-

NF-κB pathways were pioneered in flies and found to control developmental patterning and

infection response of the innate immune system [25-27], less information is available for its

possible roles in carcinogenesis.

Here, we describe the aberrant expression and function of a NF-kB transcription factors in

the *Ras^{V12}; scrib^-* tumour model. We demonstrate that in *Ras^{V12}; scrib^{IR}* tumours, the NF-kB

transcription factor Dorsal (Dl) plays a pro-tumorigenic role by opposing differentiation and

promoting survival and invasion. In this context, Dorsal can amplify JNK signaling activation, a

known driver of invasion in this model. We also found that the known Dorsal targets *snail* and

twist contribute to tumour growth. Interestingly, we report the expression of two different splicing

variants of Dl, DorsalA (DlA) and DorsalB (DlB), in non-overlapping tumour cell populations.

Results

The Toll pathway promotes tumour growth in an autonomous manner

To identify potential tumour-driving mechanisms, we surveyed upregulated genes in

Ras^{V12}; scrib^{-/-} malignant tumours by RNA Sequencing (Fig 1A). We identified 364 genes with

elevated expression levels. These include 14 previously identified genes with defined functions in

driving *Drosophila* tumor growth or paraneoplastic effects, like organ wasting and diuretic

dysfunction (Table 1). Notably, among the upregulated genes - PGRP-SA, GNBP2, Easter, Spz3,

and Spz5 - encode extracellular components of the Toll pathway (Fig 1A). The Toll pathway, one

of the two established NF-kB signaling pathways in *Drosophila*, is differentially activated during

development and in response to fungal or bacterial infection. PGRP-SA (Peptidoglycan

Recognition Protein SA) is a secreted protein that binds to bacteria-derived peptidoglycans upon

bacterial infection, forming a trimeric complex with a member of the GNBP family (Gram-Negative

Binding Protein) and the serine protease ModSP (Modular Serine Protease). This association

initiates a multi-step proteolytic cascade, eventually cleaving the ligand Spz (Spaetzle).

Subsequently Spz binds to the Toll receptor, activating the Toll pathway intracellularly (Fig 1B).

To date, six Spz ligands and nine Toll receptors have been identified and the functions and

associations of the different ligands and receptors is not yet fully understood [27]. Intracellularly,

Toll receptor activation triggers the activation and formation of a complex composed of the

adaptor proteins Myd88 and Tube as well as the serine-threonine protein kinase Pelle. Pelle

subsequently phosphorylates the inhibitor Cactus, the homolog of mammalian IkappaB, which is

targeted to proteasome degradation. As Cactus sequesters the NF-kB Dorsal or Dif into the

cytoplasm, its degradation triggers the translocation of the NF-kBs into the nucleus and the

transcription of their target genes (Fig 1B).

We decided to first investigate whether PGRP-SA is involved in Ras^{V12}; $scrib^{-/-}$

tumorigenesis. To do so, we genetically induced the formation of Ras^{V12}; $scrib^{-/-}$ tumours from

randomly induced single cells in the Eye-Antennal Disc (EAD) of the *Drosophila* larva. This way,

we can knockdown or overexpress any gene of interest within the tumour specifically without

affecting expression in the rest of the body (Fig 1C). To improve RNA interference (IR) expression

levels and processing respectively, knockdown experiments were sometimes performed at 29°C

with the additional expression of Dicer-2, as indicated in the legends.

Knock-down of PGRP-SA ($PGRP-SA^{IR}$) significantly reduced Ras^{V12}; $scrib^{-/-}$ tumour growth

compared to control tumours (Fig 1D). Similarly, knock-down of ModSP ($ModSP^{IR}$) significantly

decreased tumour size (Fig 1E), suggesting that the extracellular components of the Toll pathway

surprisingly play a pro-tumorigenic role in this context. To address whether Toll signaling is

required within tumour cells or in another microenvironmental compartment, we performed knock-

down of the intracellular kinase Pelle/IRAK ($pelle^{IR}$), a hub in Toll signal transduction. $pelle^{IR}$

Ras^{V12}; $scrib^{-/-}$ tumours exhibited a drastic size reduction compared to control tumours (Fig 1F).

These results collectively suggest that Ras^{V12}; $scrib^{-/-}$ tumours produce extracellular Toll-activating

components that contribute to a Toll pathway pro-tumorigenic signal activity in an autonomous

manner.

120 Dorsal is ectopically expressed in oncogenic Ras^{V12}-driven tumours

Toll pathway activation leads to the release and nuclear translocation of the NF-κB

transcription factors Dl and/or Dif, leading to transcription of target genes. While most studies of

Dl and Dif in developmental patterning and infection have focused on their main splicing variants

(DlA and DifA), both NF-kB transcription factors have been reported to also have alternatively

spliced B isoforms (DlB and DifB) [28, 29]. For instance, both DlA (678 amino acids) and DlB (994

amino acids) contain a Relish-homology domain, composed of a DNA binding domain and a

dimerization domain (for homo/heterodimerization and Cactus binding), as well as a

transactivation domain (for transcription activation) [28]. However, the transactivation domain of

DlA is much shorter than Dorsal-B. Noteworthy, only DlA possesses an NLS sequence. Therefore,

DlB cannot translocate to the nucleus by itself but is thought to be able to bind to DlA for

translocating to the nucleus of fat body cells upon infection [28]. Contrary to DlA, DlB is not

expressed and involved in embryonic dorso-ventral patterning. It is rather specifically expressed

in the subsynaptic reticulum of neuromuscular junctions, where it stabilizes Cactus. Similarly, Dif-

B is found in the mushroom body of the CNS, where it also stabilizes Cactus [29]. Although these

observations support the idea that B isoforms play a specific role in nervous system tissues, their

precise functions remain to be elucidated [30].

We began by assessing the expression of Dl in Ras^{V12}; $scrib^{IR}$ tumours through

immunostainings against DlA and DlB (Fig 2A, 2B, 2C, 2D and S1A, S1B Fig). Both Dorsal

isoforms were highly expressed in tumors in the posterior part of the disc that attaches to the brain

through the prospective optic stalk (Fig 2A and 2B), while neither DlA or DlB proteins were

observed in control clones (Fig 2A). DlA and DlB were almost entirely mutually exclusive, in

different cell populations (Fig2A, 2B and S1B Fig). Within Ras^{V12}; $scrib^{IR}$ tumours, cells can be

easily separated according to their levels of DIA mean intensity (Fig 2C). We identified a DIA^{high}

cell population that had four-fold higher DIA protein levels compared to the DIA^{low} remaining cell

population. Importantly, the DIA mean intensity in the DIA^{low} cell population was consistently

higher than in the surrounding *wt* epithelial cells of the EAD (Fig 2C). These observations suggest

that although we observed a peak of DIA expression at the posterior part of the tumour, there is

a general elevation of the basal level of DIA throughout the entire tumour. In contrast, we did not

observe a general elevation of DIB throughout the tumour when compared to *wt* cells of the TME,

but we did observe a local increase in the posterior part of the tumour of a similar magnitude to

DIA (Fig 2B and 2D). Tumour cells located in the posterior area of the tumour are closest to the

CNS, which is the first organ these tumours invade [31-33]. Intriguingly, tumour cells that have

initiated invasion into the optic lobes (OL) of the CNS displayed high levels of DIA or DIB (S1B

Fig), suggestive of a potential role of Toll-NFκB signaling in tumor cell migration.

Finally, a strong enrichment of the repressor Cactus, the ortholog of mammalian IkappaB

(IKKB), was observed in DIA^{high} tumour cells but not in DIB^{high} tumour cells, as assessed with an

endogenous GFP fusion protein reporter, *Cactus::GFP* (S1C Fig). This is particularly interesting

as *cactus* is a positive transcriptional target of the Toll pathway[34]. Cactus upregulation

constitutes a negative feedback loop which could cap Toll pathway activation, a mechanism that

has also been described in mammalian NF-kB signalling pathways [35, 36]. In conclusion,

simultaneous elevation of DIA, DIB and Cactus expression suggests that in the posterior part of

Ras^{V12}; $scrib^{IR}$ tumours, Toll pathway activation is significantly higher than in the rest of the tumour.

Dorsal aberrant accumulation is genetically and spatiotemporally-regulated

We investigated what could be the upstream mechanism leading to aberrant DI

accumulation. Initially, we examined DI isoforms levels in various genetic backgrounds: control

wild type clones, *scrib^{IR}* tumours, *Ras^{V12}* tumours and *Ras^{V12}; scrib^{IR}* tumours. We observed

aberrant expression of DIA as well as DIB exclusively in *Ras^{V12}* and *Ras^{V12}; scrib^{IR}* tumours (Fig

2A), suggesting that *Ras^{V12}* instructs tumour cells competent to produce these isoforms at

abnormal levels. Furthermore, we followed DIA levels over time in *Ras^{V12}; scrib^{IR}* tumours, using

an endogenous Dorsal genetic reporter, that acts as a Dorsal-A specific reporter in this context

(Fig 3A and S2A and S2B Fig). The onset of DIA expression correlated with the appearance of

Elav-positive neurons in the prospective eye region of the tumours around Day5 (Fig 3A).

Interestingly, DIA accumulate in cells surrounding the neurons but was rarely detected in the

neurons themselves (Fig 3A yellow arrows and Fig 3B). Concurrently, DIB expression was

exclusive to the tumour neurons (Fig 3C). Within the tumours, apoptosis was predominantly

observed in the differentiating areas where we observed dying neurons (Fig 3D, yellow arrows).

Therefore, we hypothesized that the local aberrant DI expression might be triggered by cell death-

derived signals. However, efficient blocking of apoptosis in *Ras^{V12}; scrib^{IR}* tumours through Diap1

overexpression (*Diap1^{OE}*) did not affect increased DIA and DIB levels (Fig 3E and 3F).

Together, these observations demonstrate that the aberrant expression NF-kB/DI in

Ras^{V12}; scrib^{IR} tumours is both genetically and locally controlled. General elevated levels of DIA in

the whole tumor population suggests a transcriptional response downstream of oncogenic Ras^{V12}

signaling. On top of this Ras^{V12}-dependent genetic control, a temporal and or local control triggers

elevated levels of DIA, DIB and Cactus specifically in the posterior part of the tumour in the

prospective eye region, as well as in invading tumour cells to the brain.

Dorsal represses Ras^{V12} scrib^{IR} tumour differentiation and death

To investigate the primary functions of DI in *Ras^{V12}; scrib^{IR}* tumours, we conducted

knockdown experiments using two independent RNAis that efficiently target both Dorsal isoforms

(S3A and S3B Fig). *Dorsal^{IR}, Ras^{V12}; scrib^{IR}* tumours exhibited a significant reduction in size

compared to *Cherry^{IR} Ras^{V12}; scrib^{IR}* control tumours (Fig 4A), mimicking the phenotypes

observed upon *PGRP-SA*, *ModSP* and *pelle* knockdowns (Fig1C, 1D and 1E). *Ras^{V12}; scrib^{IR}*

tumour growth results from a combination of cell growth, cell proliferation, inhibition of the

differentiation of epithelial tumour cells into Elav-positive photoneurons and survival [31, 37]. To

delineate the potential functions of Dl in *Ras^{V12}; scrib^{IR}* tumours, we assessed alterations in

proliferation, differentiation and apoptosis. Quantification of mitotic cells suggests that *Dorsal^{IR}*

Ras^{V12}; scrib^{IR} tumours display similar proliferation rates to control tumours, but both their

differentiation and apoptosis rates were significantly higher than in control tumours (Fig4B, 4C

and 4D). Although, a few neurons form in *Ras^{V12}; scrib^{IR}* tumours, they appeared spaced apart

and did not form the full complement of eight neurons per ommatidia. However, upon Dorsal

knockdown, Elav-positive cells increased in number and density, suggesting a partial rescue of

differentiation (Fig4C). The fact that we frequently observed cell death coinciding with areas

where Elav-positive neurons were present (Fig 3D) suggests that cell death might be a

consequence of aberrant tumour cell differentiation. Importantly, *Pelle^{IR} Ras^{V12}; scrib^{-/-}* tumours

also exhibited increased apoptosis and differentiation rates (S3C Fig). This emphasizes the need

for the Toll signalling pathway, and not an alternative way to activate Dorsal/NFkB activity, to

promote tumour growth. Thus, the Toll-Dl signaling pathway promotes tumour growth primarily by

preventing differentiation and promoting survival.

Dorsal is required for invasion and accentuates JNK signaling

We next addressed whether Dl is involved in tumor invasion. In the *Ras^{V12}; scrib^{IR}* tumour

model, cells adopt a migratory behavior by JNK signaling. Specifically, JNK mediates expression

of Matrix MetalloProteinase 1 (MMP1) and Cheerio (Cher-mammalian Filamin), resulting in

basement membrane degradation and cytoskeleton remodeling respectively [22, 32, 37-41]. We

conducted colocalization analyses, assessed by the Pearson's Correlation Coefficient (PCC),

between the Dorsal isoforms and JNK activity in the tumours, assessed either through activated

JNK (phosphor-JNK, pJNK) or a JNK activity reporter, TRE-eGFP. Overall, we found a low

colocalization between DIA and TRE-eGFP (Fig 5A) while we observed a good PCC between DIB

and pJNK within the tumours (Fig 5B). These observations align with the finding that tumor cells

invading the CNS often have high levels of either DIA or DIB.

To test whether Dorsal may function upstream of JNK signaling in Ras^{V12}; $scrib^{IR}$ tumours, we

assessed the levels of the JNK target MMP1 following $Dorsal^{OE}$ and $ModSP^{OE}$ overexpression

which enhances Toll signaling and subsequently Dorsal activity. MMP1 mean intensity was

significantly higher in tumors with elevated *Dorsal* or *ModSP* expression. compared to control

tumours. Therefore, Toll-Dl signaling can augment expression of the JNK transcriptional target,

mmp1 in Ras^{V12}; $scrib^{IR}$ tumours (Fig 5C).

As both DIA, DIB and Cactus are highly expressed in brain lobe proximal and invading cells and

Dl amplifies JNK signalling in the tumours, we investigated the effect of Dorsal knock-down on

tumour cell migration. We scored for three parameters reflecting the mobility of tumour cells at

late stages of tumour development (Day 12): 1/ the VNC invasion score, which assigns a higher

score to tumors with greater migration toward the tip of the ventral nerve cord (VNC) of the CNS,

2/the frequency at which we observe tumour cells within the organs adjacent to the VNC called

leg discs and 3/ the fusion of tumours generated in the two distinct EAD, indicating the ease with

which cells move from one EAD to the other and EADs and come into contact with each other

(Fig 6A). To mitigate the influence of tumour cell number on migration assessment, we compared

control and $Dorsal^{IR}$ tumours of similar size. Upon Dorsal knock-down, at late stages, the three

parameters showed a drastic decrease: the VNC score decreased threefold compared to control,

reaching a low value similar to early-stage control tumours (Fig6B). The frequency of invasion

into leg discs decreased sevenfold and tumors from neighboring EADs often failed to merge (Fig

6A). Similarly, Dorsal overexpression ($Dorsal^{OE}$) increased tumour cell mobility even at early

stages (Day 6), while tumour size is signicantly smaller (Fig 6B), suggesting that 1/ the increase

in invasion is not simply a secondary consequence of higher cell number and 2/ that the balance

between invasion and tumour growth tilted toward invasion when tumour cells have high levels of
Dorsal.

Given the functions of Dorsal in the tumour context, we wondered whether $Dorsal^{OE}$ is sufficient
to induce tumorigenesis. Although, $Dorsal^{OE}$ control clones were not bigger than wt clones, we
systematically observed the formation of dysplasic GFP-labeled $Dorsal^{OE}$ clones within the EADs
that appeared as large-rounded cell masses often located basally (S4A and S4B Fig, yellow
arrow). We hypothetize that they arise from the inhibition of differentiation. Moreover, in this
context, we never observed $Dorsal^{OE}$ cells in the VNC (VNC score in S4A Fig). Thus, Dorsal at
high levels is sufficient to trigger dysplasia but not invasive hyperplastic tumours.

The Epithelial-to-Mesenchymal (EMT) transcription factor Snail has been implicated in
promoting invasion of $Ras^{V12}; lgl^{-/-}$ tumours (a similar tumour model to $Ras^{V12}; scrib^{-/-}$ tumours)
through stimulation of JNK signaling and EMT-like cytoskeleton remodeling [42]. Snail, along
with Twist, another EMT transcription factor, are well-known targets of Dorsal that govern
mesoderm invagination during developmental patterning in cells with the highest Dorsal activity
[43-46]. Interestingly, knockdowns of both Snail and Twist significantly decreased $Ras^{V12}; scrib^{IR}$
tumour size (S4A and S4B Fig), mimicking Dorsal knockdown. At late stages, twistIR $Ras^{V12};$
$scrib^{IR}$ tumour also displayed a lower VNC score compared to control tumours, suggesting
impaired tumour cell migration (S4B Fig). We propose a model where the Dl^{high} tumour cells
activate Twist and Snail expression, provoking migration, in part through JNK signalling
enhancement (Fig 7A).

Taken together, these experiments suggest that Dl promotes tumor growth through
repressing cell death and differentiation as well as promoting EMT like delamination and invasion
dependent on Twist and Snail activity.

Conclusion/Discussion
264

Here, we establish that autonomous activation of the inflammatory Toll pathway plays a pivotal

role in supporting the growth of Ras^{V12}; $scrib^{IR}$ tumours. The NF-kB transcription factors Dorsal,

and key effectors of the Toll pathway, exhibit aberrant expression in this model. In this context,

Dorsal activity facilitates tumour growth by preventing differentiation and promoting survival

without measurable impact on proliferation. Additionally, Dorsal promotes invasion genetically

dependent on the EMT factors Snail and Twist.

An exciting feature in this tumor model is tumor heterogeneity in both NF-kB protein composition

(DIA and DIB) and Toll pathway activity (Dl^{high} cactus^{high} vs Dl^{low}, Cactus^{low}), as judged by classical

target gene expression (*cactus* and *dorsal*) and stimulation of invasion. Dl is unevenly

accumulated within the population of transformed cells in a spatial and regional manner and on

at least two levels: there is a low-level DIA accumulation within the entire region of Ras^{V12}

expression and a heightened level in the posterior region situated in the differentiating prospective

eye region that is physically attached to the brain through the stalk. In this region a notable

accumulation of both splicing variants of Dl, DIA and DIB, is seen within distinct tumour cell

populations.

While NF-kB protein heterogeneity within the same tumour has not been studied yet in

solid mammal tumours, the NF-kB fingerprinting and their subsequent homo-/heterodimerization

within diffuse large B-cell lymphoma hold predictive values for the tumour cell response to

microenvironmental activating cues [47]. With our study, this finding, suggest that NF-kB

composition heterogeneity might also occur in solid mammal cancers and modulate their

response to microenvironmental inflammation.

Moreover, heterogeneity in the level/activity of a single NF-kB mirrors observations in mammalian

cancers. In pancreatic cancer, where oncogenic KRAS is the main driver, tumor heterogeneity is

a central hallmark and frequently associated with variable NF-kB signaling. A central question is

thus, what causes some cells to respond with inflammation whereas others do not? Possibilities

include non-autonomous external input from nearby cells of the microenvironment or circulation

versus intrinsic mechanisms. *In vitro* studies of KRAS[G12V]-transfected immortalized human

pancreatic epithelial derived cells (HPDE) have demonstrated that variable levels of oncogenic

KRAS can produce different outcomes. High KRAS[G12V] expression can direct NF-kB signaling

and EMT phenotypes driven in part by feed forward loop through IL6 signaling, whereas lower

level of KRAS activity retains epithelial character with lower level of NF-kB signaling dependent

on IL1 [48].

In the *Ras[V12], scrib[-/-]* model, a "flip out" cassette turns on Ras[V12] expression under control

of the actin promoter in all cells with the same genetic dose, and by first principle should signal

evenly. Yet, Ras[V12] - driven NF-kB expression is temporally and spatially controlled in a

stereotypic pattern. One possibiity is that this region is prepatterned to respond to increased levels

of Ras[V12] with a higher response. Indeed, in this region, we observe a higher level of pMAPK and

where tumour region neurogenesis is incomplete as we do not observe fully formed ommatidia

emphasizing the inhibitory effect of Dorsal on differentiation. In the prospective eye region, behind

the morphogenetic frurrow, founder R8 neurons recruit further neurons through Receptor Tyrosine

Kinase-MAPK signaling that potentially could add up with Ras[V12]-MAPK signaling to reach a

higher signaling levels [49].

Extrinsic regulation of Toll signaling does have a role in the *Ras[V12], scrib[-/-] t*umor model for growth.

The Toll-activating components implicated in tumour growth, PGRP-SA and ModSP, are part of

a bacterial recognition alert system. The observed overexpression of PGRP-SA in *Ras[V12]*; *scrib[IR]*

tumours may serve to locally increase the concentration of active cleaved Spz ligands, thereby

stimulating Toll activation and promoting tumor growth. The source of activation governed by

PGRP-Sa and ModSP remains unknown. As the posterior part of the disc experience higher levels

of apoptosis, we considered whether apoptotic cell death and debris could active PGRP-

SA/ModSP - driven Toll sigaling in this context through death- induced danger signals and

bacterial component mimicry, but inhibition of death through DIAP1 expression did not alter NFkB

levels.

Notably, previous studies have identified PGRP-SA, Cactus, and Dorsal as positive targets of the

Toll pathway [34]. Furthermore, it has been shown that the CNS serves as a significant source of

Spz ligands that regulates Ras^{V12}; $lgl^{-/-}$ tumour migration tropism through Toll6 [33]. Hence, it is

tempting to think that upon the onset of neurogenesis, tumour cells suddenly gain access to

elevated levels of Spz concentration from the brain, significantly enhancing local Toll activation

and subsequent PGRP-SA and Dorsal expression. This hypothesis may explain the spatial

restriction of aberrant Dorsal expression to the posterior part of the tumor.

We have yet to assess whether DIA and DIB exert different functions within Ras^{V12}; $scrib^{IR}$

tumours, as our knockdowns and overexpression transgenes targeted both isoforms

simultaneously. Consistent with previous reports, DIB expression in the tumor context is restricted

to cell types from the nervous system. Colocalization of DIB with pJNK in neurons that seem prone

to die, allows us to speculate if both isoforms have distinct functions, that DIB may promote

migration through JNK, while DIA may favor stemness and survival. Additionally, we cannot yet

rule out the involvement of other NF-kBs, such as Dif and Relish in Ras^{V12}; $scrib^{IR}$ tumours.

Using splicing variants may be a mechanism to diversify the range of NF-kB actions in

various contexts in Drosophila, as it possesses only three NF-kBs (Dorsal, Dif, and Relish). In

contrast, mammalian NF-kBs are not reported to have splicing variants, although they possess

five different NF-kBs. To our knowledge, the concept of intratumoral NF-kB heterogeneity remains

relatively unexplored in mammalian cancer biology and the fly model offer an experimentally

accessible model to investigate this question *in vivo*. Future research efforts will be crucial in

elucidating the importance and mechanisms underlying intratumoral NF-kB heterogeneity during

malignant transformation.

The Ras^{V12}, scrib tumor model engages five known inflammatory pathways during
transformation. Ras^{V12} directly drives MEK-ERK and PI3K-AKT signaling while engaging Toll-
NFkB, whereas loss of cell polarity through scribbled, initiates TNFR-JNK MAPK stress signaling
and as a result, an Upd/IL6-JAK-STAT autocrine signaling loop altogether favoring malignant
transformation.

In conclusion, the findings herein extends the understanding of how innate immune signaling is
engaged by and cooperates with oncogenic Ras, and opens up for more detailed mechanistic
studies in a genetically tractable preclincial tumor model with clear paralells to human cancer

Material and Methods

Drosophila **lines**

The EyMARCM system is known to generate flip-out clones in the adjacent optic lobes of the
CNS in addition to the eye discs [50]. For assessing tumour cell migration from true carcinoma
cells arising from the eye disc epithelium, we therefore used another genetic system designed in
our lab, called EyaHost, that generates flip-out tumours exclusively in the EAD but not in the
CNS (to be described in detail elsewhere).

Unless otherwise specified, crosses were conducted at 25°C by default; however, for enhanced
knockdown efficiency, most knockdown experiments were carried out at 29°C, as detailed in the
figures and legends.

• EyMARCM Drosophila lines:

Ey,flp; act>STOP>Gal4, UAS-GFP; FRT82B Tub-Gal80ts [32]

Ey,flp, UAS-Dcr2; act>STOP>Gal4, UAS-GFP; FRT82B Tub-Gal80ts

;Sp/CyO, Dfd-YFP; UAS-Ras^{V12} FRT82B scrib[2]/TM6C {Bilder, 2000 #23;Chen, 2012 #24;Perez,

2017 #22;Perez-Garijo, 2009 #25}

; UAS-PGRP-SA[IR]; UAS-Ras^{V12} FRT82B scrib[2]/TM6C (BDSC#60037)

; UAS-modSP[IR]; UAS-Ras^{V12} FRT82B scrib[2]/TM6C (VDRC GD#43972)

; UAS-pelle[IR]; UAS-Ras^{V12} FRT82B scrib[2]/TM6C (BDSC#50715)

; UAS-RFP[IR]; UAS-Ras^{V12} FRT82B scrib[2]/TM6C (BDSC#67852)

;UAS-Ras^{V12} ; FRT82B scrib[1]/TM6C [51]

;UAS-Ras^{V12} ; FRT82B

FRT82B scrib[1]/TM6C

FRT82B

• EyaHost Drosophila lines

; Eya-KD; QUAS-Gal4 (Eya-KD our own lab, QUAS-Gal4 BDSC#83132 {Baena-Lopez, 2018

#32})

QUAS-Ras^{V12} ; Eya-KD; QUAS-Gal4 (QUAS-Ras^{V12} from Peter Galant)

QUAS-Ras^{V12} ; Eya-KD, QUAS-scrib[IR]; QUAS-Gal4 (QUAS-scrib[IR] from our own lab)

QUAS-Ras^{V12} ; Eya-KD, QUAS-scrib[IR];

QUAS-Ras^{V12} ; Eya-KD, QUAS-scrib[IR]; QUAS-Gal4, UAS-Dcr2 (UAS-Dcr2 BDSC#24651)

Act>|STOP>|QF, QUAS-mCD8::GFP;; (Act>|STOP>|QF from our own lab, QUAS-mCD8::GFP

BDSC#30001)

Act>|STOP>|QF, QUAS-mCD8::GFP; UAS-dorsal; (BDSC#9319)

Act>|STOP>|QF, QUAS-mCD8::GFP; UAS-modSP; (B.Lemaître {Dostalova, 2017 #33})

Act>|STOP>|QF, QUAS-mCD8::GFP; UAS-diap1^{OE}; (BDSC#6657)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-cherry^{IR} (BDSC#35785)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-dorsal^{IR-1} (BDSC#32934)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-dorsal^{IR-2} (BDSC#27650)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-snail^{IR} (VDRC GD#6232)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-twist^{IR-1} (BDSC#51164)

Act>|STOP>|QF, QUAS-mCD8::GFP;; UAS-twist^{IR-2} (BDSC#25981)

Act>|STOP>|QF, QUAS-Tag2BFP;; genomic-dorsal::GFP (BDSC#42677)

Act>|STOP>|QF, QUAS-Tag2BFP;; genomic-cactus::GFP (VDRC fTRG 318145)

Act>|STOP>|QF, Ey-flp, QUAS- Tag2BFP; TRE-eGFP; (QUAS-Tag2BFP our own lab, TRE-

eGFP BDSC#59010)

Immunohistochemistry

Inverted larvae heads were fixed 30 min in 4% paraformaldehyde/PBS at room temperature on a

shaker. After washes, fine dissection of single EADs (for early-stage tumours at Day 6) or whole

cephalic complexes (at later stages) was conducted in PBS. The tissues were then incubated with

a solution of primary antibody in 0,5% Triton X-100/PBS overnight at 4°C. Noteworthy, for primary

antibodies that presented penetration defects (marked with * below), we incubated the tissues in

0,5% Triton X-100/PBS overnight at 4°C prior to primary antibody stainings. Similarly, secondary

antibody stainings was done in 0,5% Triton X-100/PBS overnight at 4°C. Samples were then

mounted in Vectashield or equivalent mounting media for imaging.

Primary antibodies were used at the following dilutions: anti-DIA mouse antibody (1:50, DSHB

7A4), anti-DIB rabbit antibody (* 1:1000, S.Wasserman [29]), anti-pH3 rat (1:200 abcam 10543),

anti-Elav rat (1:50, DSHB 7E8A10), anti-cleaved Dcp1 rabbit (1:100, Cell Signaling 9578), anti-

phosphoJNK mouse (* 1:100, Cell Signaling 9255) and anti-MMP1 mouse (* used in a mix 1:1:1,

DSHB 3B8D12, DSHB 3A6B4 and DHSB 5H7B11). Secondary antibodies were all from Jackson

ImmunoResearch.

RNA sequencing and data analysis

Third instar larval eye discs of the indicated genotypes were manually isolated and RNA was

extracted using Qiagen RNAmini kits. TruSeq RNA Library Prep Kit (Illumina) was used to prepare

libraries, and sequencing was performed on an Illumina MiSeq system. Raw data were analyzed

with the Tuxedo suite (RRID:SCR_013194) and reads were mapped to Drosophila genome

release 5.2. Expression was recorded as FPKM: fragments per kilo-base per million reads.

Imaging

Samples were imaged on Zeiss LSM880 and Zeiss LSM980 confocal microscopes as well as on

Nikon HCI SoRa spinning disc microscope. They were further processed on FIJI.

Tumour Volume Analysis

Z-stacks were acquired with a 3µm step for early-stage tumors (Day 6) and a 4µm step for late-

stage tumors (after Day 6). Tumor volume was estimated using a FIJI Macro [52], which

automatically performs threshold-based 3D reconstruction of the GFP+ signal, followed by

volume measurements (3D Object counter plugin). Before 3D reconstruction, the FIJI Macro

applies two consecutive Gaussian blur steps (sigma=2) to reduce noise.

Image processing

• Fluorescence intensity measurements and heatmaps

Fluorescence intensity measurements were performed using a FIJI Macro, which subtracts

background intensity to the signal of interest inside or outside the tumour, using a tumour mask

generated from intensity thresholding. When comparing values across different conditions, they

were normalized to the mean value of the control condition, ensuring the control mean is set at

1. In Fig 2B and Fig 2C, the tumours were split in two compartments "high" and "low", according

to arbitrarily set intensity thresholds (250 for DIA, 10000 for DIB). Normalized mean intensities

were then automatically quantified within the different compartments: tumourHigh, tumourLow

and TME. DIA and DIB intensity heatmaps were generated after background subtraction, using

the Fire LUT in FIJI.

• Colocalization analysis

Colocalization analyses were performed using a FIJI Macro, which automatically computed the

Pearson's Correlation Coefficient (PCC) between two channels of a confocal image using the

"coloc2" plugin. The PCC, ranging from -1 to 1, denotes the degree of colocalization between

signals. A higher absolute value indicates stronger colocalization, with the sign indicating

correlation direction ("-" for anti-correlation, "+" for correlation).

● Proliferation, differentiation and apoptosis ratio

The proliferation, differentiation and apoptosis ratio are defined as the area of the meaningful

marker within the tumour (pH3, Elav and cDcp respectively) divided by the area of the tumour.

These ratios were measured with a FIJI Macro which measures both the tumour area and the

tumour marker area using an automatic-thresholding. For Fig 3A, 3B and 3C, measurements

were conducted on single confocal pictures whereas, for S1 Fig, the apoptosis ratio was

measured on the Z-projection of confocal Z-stacks.

● Migration score

Migration of the VNC and leg discs was manually scored on Z-projection (Max Intensity mode)

of confocal Z-stacks.

Statistical analyses

At least, two biological replicates were performed for each experiment. According to the

normality and size of the datasets, appropriate statistical tests were applied using GraphPad

Prism9 and are indicated in the figure legends. Graphs were also generated on GraphPad

Prism9 and are presented as boxplots (10-90 percentile) or dot plots indicating the mean of the

datasets. Statistical significance was categorized as follows: *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,

****$p < 0.0001$, and n.s when not significant.

Acknowledgement

We thank Drs. Steven A. Wasserman and Amy Kieger, Bruno Lemaitre, the

Bloomington Drosophila Stock Center; the Vienna Drosophila Resource Center and the

Developmental Studies Hybridoma Bank for providing fly

stocks and antibodies. We thank Alicia Alfonso Gomez for help with fly handling, experimental

support and image acquisition. We also thank the Core Facilities for Advanced Light Microscopy

and Advanced Electron Microscopy at Oslo University Hospital and the Microscopy Rennes

Imaging facility of the UMS Biosit, a member of the national infrastructure France-BioImaging,

supported by the French National Research Agency (ANR-10-INBS-04).

C.D, J.T.R., A.J., T.E.R and were financed by The Norwegian Research Council Toppforsk and

Center of Excellence, CanCell grants #262652 and #276070

Author contributions

CD and TER conceived the project and planned experiments. CD, AJ, JTR and H.J. performed
experiments and analysis. H.J., and R.L. B. provided feedback to experimental approaches. T.E.R
Is the leading principal investigator who together with C.D supervised research and edited the
manuscript with input from all coauthors.

1.   Hanahan D. Hallmarks of Cancer: New Dimensions. Cancer Discov. 2022;12(1):31-46.

2.   Zhao H, Wu L, Yan G, Chen Y, Zhou M, Wu Y, et al. Inflammation and tumor progression:
signaling pathways and targeted intervention. Signal Transduct Target Ther. 2021;6(1):263.

3.   Aggarwal BB, Sung B. NF-kappaB in cancer: a matter of life and death. Cancer Discov.
2011;1(6):469-71.

4.   Xia L, Tan S, Zhou Y, Lin J, Wang H, Oyang L, et al. Role of the NFkappaB-signaling pathway in
cancer. Onco Targets Ther. 2018;11:2063-73.

5.   Gu L, Wang Z, Zuo J, Li H, Zha L. Prognostic significance of NF-kappaB expression in non-small
cell lung cancer: A meta-analysis. PLoS One. 2018;13(5):e0198223.

6.   Sarkar DK, Jana D, Patil PS, Chaudhari KS, Chattopadhyay BK, Chikkala BR, et al. Role of NF-
kappaB as a Prognostic Marker in Breast Cancer : A Pilot Study in Indian Patients. Indian J Surg Oncol.
2013;4(3):242-7.

7.   Boukerche H, Su ZZ, Emdad L, Sarkar D, Fisher PB. mda-9/Syntenin regulates the metastatic
phenotype in human melanoma cells by activating nuclear factor-kappaB. Cancer Res. 2007;67(4):1812-
22.

8.   Huang S, Pettaway CA, Uehara H, Bucana CD, Fidler IJ. Blockade of NF-kappaB activity in human
prostate cancer cells is associated with suppression of angiogenesis, invasion, and metastasis.
Oncogene. 2001;20(31):4188-97.

9.   Huber MA, Azoitei N, Baumann B, Grunert S, Sommer A, Pehamberger H, et al. NF-kappaB is
essential for epithelial-mesenchymal transition and metastasis in a model of breast cancer progression. J
Clin Invest. 2004;114(4):569-81.

10.   Biswas DK, Shi Q, Baily S, Strickland I, Ghosh S, Pardee AB, et al. NF-kappa B activation in human
breast cancer specimens and its role in cell proliferation and apoptosis. Proc Natl Acad Sci U S A.
2004;101(27):10137-42.

11.   Cui X, Shen D, Kong C, Zhang Z, Zeng Y, Lin X, et al. NF-kappaB suppresses apoptosis and
promotes bladder cancer cell proliferation by upregulating survivin expression in vitro and in vivo. Sci
Rep. 2017;7:40723.

12.   Talukdar S, Das SK, Pradhan AK, Emdad L, Windle JJ, Sarkar D, et al. MDA-9/Syntenin (SDCBP) Is a
Critical Regulator of Chemoresistance, Survival and Stemness in Prostate Cancer Stem Cells. Cancers
(Basel). 2019;12(1).

13.   Bentires-Alj M, Barbu V, Fillet M, Chariot A, Relic B, Jacobs N, et al. NF-kappaB transcription
factor induces drug resistance through MDR1 expression in cancer cells. Oncogene. 2003;22(1):90-7.

14.   Bednarski BK, Ding X, Coombe K, Baldwin AS, Kim HJ. Active roles for inhibitory kappaB kinases
alpha and beta in nuclear factor-kappaB-mediated chemoresistance to doxorubicin. Mol Cancer Ther.
2008;7(7):1827-35.

15.   Xia YZ, Ni K, Guo C, Zhang C, Geng YD, Wang ZD, et al. Alopecurone B reverses doxorubicin-
resistant human osteosarcoma cell line by inhibiting P-glycoprotein and NF-kappa B signaling.
Phytomedicine. 2015;22(3):344-51.

16.   Baud V, Karin M. Is NF-kappaB a good target for cancer therapy? Hopes and pitfalls. Nat Rev
Drug Discov. 2009;8(1):33-40.

17.   Rasmi RR, Sakthivel KM, Guruvayoorappan C. NF-kappaB inhibitors in treatment and prevention
of lung cancer. Biomed Pharmacother. 2020;130:110569.

18.   Bilder D, Ong K, Hsi TC, Adiga K, Kim J. Tumour-host interactions through the lens of Drosophila.
Nat Rev Cancer. 2021;21(11):687-700.

19.   Dillard C, Reis JGT, Rusten TE. Ras(V12); scrib(-/-) Tumors: A Cooperative Oncogenesis Model
Fueled by Tumor/Host Interactions. Int J Mol Sci. 2021;22(16).

20.   La Marca JE, Richardson HE. Two-Faced: Roles of JNK Signalling During Tumourigenesis in the
Drosophila Model. Front Cell Dev Biol. 2020;8:42.

21.	Pastor-Pareja JC, Wu M, Xu T. An innate immune response of blood cells to tumors and tissue
damage in Drosophila. Dis Model Mech. 2008;1(2-3):144-54; discussion 53.
22.	Cordero JB, Macagno JP, Stefanatos RK, Strathdee KE, Cagan RL, Vidal M. Oncogenic Ras diverts
a host TNF tumor suppressor activity into tumor promoter. Dev Cell. 2010;18(6):999-1011.
23.	Khezri R, Holland P, Schoborg TA, Abramovich I, Takats S, Dillard C, et al. Host autophagy
mediates organ wasting and nutrient mobilization for tumor growth. EMBO J. 2021;40(18):e107336.
24.	Holland P, Quintana EM, Khezri R, Schoborg TA, Rusten TE. Computed tomography with
segmentation and quantification of individual organs in a D. melanogaster tumor model. Sci Rep.
2022;12(1):2056.
25.	Minakhina S, Steward R. Nuclear factor-kappa B pathways in Drosophila. Oncogene.
2006;25(51):6749-57.
26.	Hetru C, Hoffmann JA. NF-kappaB in the immune response of Drosophila. Cold Spring Harb
Perspect Biol. 2009;1(6):a000232.
27.	Valanne S, Wang JH, Ramet M. The Drosophila Toll signaling pathway. J Immunol.
2011;186(2):649-56.
28.	Gross I, Georgel P, Oertel-Buchheit P, Schnarr M, Reichhart JM. Dorsal-B, a splice variant of the
Drosophila factor Dorsal, is a novel Rel/NF-kappaB transcriptional activator. Gene. 1999;228(1-2):233-
42.
29.	Zhou B, Lindsay SA, Wasserman SA. Alternative NF-kappaB Isoforms in the Drosophila
Neuromuscular Junction and Brain. PLoS One. 2015;10(7):e0132793.
30.	Gramates LS, Agapite J, Attrill H, Calvi BR, Crosby MA, Dos Santos G, et al. FlyBase: a guided tour
of highlighted features. Genetics. 2022;220(4).
31.	Brumby AM, Richardson HE. scribble mutants cooperate with oncogenic Ras or Notch to cause
neoplastic overgrowth in Drosophila. EMBO J. 2003;22(21):5769-79.
32.	Pagliarini RA, Xu T. A genetic screen in Drosophila for metastatic behavior. Science.
2003;302(5648):1227-31.
33.	Mishra-Gorur K, Li D, Ma X, Yarman Y, Xue L, Xu T. Spz/Toll-6 signal guides organotropic
metastasis in Drosophila. Dis Model Mech. 2019;12(10).
34.	De Gregorio E, Spellman PT, Tzou P, Rubin GM, Lemaitre B. The Toll and Imd pathways are the
major regulators of the immune response in Drosophila. EMBO J. 2002;21(11):2568-79.
35.	Scott ML, Fujita T, Liou HC, Nolan GP, Baltimore D. The p65 subunit of NF-kappa B regulates I
kappa B by two distinct mechanisms. Genes Dev. 1993;7(7A):1266-76.
36.	Ghosh S, May MJ, Kopp EB. NF-kappa B and Rel proteins: evolutionarily conserved mediators of
immune responses. Annu Rev Immunol. 1998;16:225-60.
37.	Kulshammer E, Mundorf J, Kilinc M, Frommolt P, Wagle P, Uhlirova M. Interplay among
Drosophila transcription factors Ets21c, Fos and Ftz-F1 drives JNK-mediated tumor malignancy. Dis
Model Mech. 2015;8(10):1279-93.
38.	Uhlirova M, Bohmann D. JNK- and Fos-regulated Mmp1 expression cooperates with Ras to
induce invasive tumors in Drosophila. EMBO J. 2006;25(22):5294-304.
39.	Srivastava A, Pastor-Pareja JC, Igaki T, Pagliarini R, Xu T. Basement membrane remodeling is
essential for Drosophila disc eversion and tumor invasion. Proc Natl Acad Sci U S A. 2007;104(8):2721-6.
40.	Kulshammer E, Uhlirova M. The actin cross-linker Filamin/Cheerio mediates tumor malignancy
downstream of JNK signaling. J Cell Sci. 2013;126(Pt 4):927-38.
41.	Doggett K, Turkel N, Willoughby LF, Ellul J, Murray MJ, Richardson HE, et al. BTB-Zinc Finger
Oncogenes Are Required for Ras and Notch-Driven Tumorigenesis in Drosophila. PLoS One.
2015;10(7):e0132987.
42.	Wu C, Ding X, Li Z, Huang Y, Xu Q, Zou R, et al. CtBP modulates Snail-mediated tumor invasion in
Drosophila. Cell Death Discov. 2021;7(1):202.

43.   Jiang J, Kosman D, Ip YT, Levine M. The dorsal morphogen gradient regulates the mesoderm
determinant twist in early Drosophila embryos. Genes Dev. 1991;5(10):1881-91.
44.   Ip YT, Park RE, Kosman D, Yazdanbakhsh K, Levine M. dorsal-twist interactions establish snail
expression in the presumptive mesoderm of the Drosophila embryo. Genes Dev. 1992;6(8):1518-30.
45.   Stathopoulos A, Levine M. Whole-genome analysis of Drosophila gastrulation. Curr Opin Genet
Dev. 2004;14(5):477-84.
46.   Hong JW, Hendrix DA, Papatsenko D, Levine MS. How the Dorsal gradient works: insights from
postgenome technologies. Proc Natl Acad Sci U S A. 2008;105(51):20072-6.
47.   Jayawant E, Pack A, Clark H, Kennedy E, Ghodke A, Jones J, et al. NF-kappaB fingerprinting
reveals heterogeneous NF-kappaB composition in diffuse large B-cell lymphoma. Front Oncol.
2023;13:1181660.
48.   Siddiqui I, Erreni M, Kamal MA, Porta C, Marchesi F, Pesce S, et al. Differential role of
Interleukin-1 and Interleukin-6 in K-Ras-driven pancreatic carcinoma undergoing mesenchymal
transition. Oncoimmunology. 2018;7(2):e1388485.
49.   Treisman JE. Retinal differentiation in Drosophila. Wiley Interdiscip Rev Dev Biol. 2013;2(4):545-
57.
50.   Newsome TP, Asling B, Dickson BJ. Analysis of Drosophila photoreceptor axon guidance in eye-
specific mosaics. Development. 2000;127(4):851-60.
51.   Wu M, Pastor-Pareja JC, Xu T. Interaction between Ras(V12) and scribbled clones induces
tumour growth and invasion. Nature. 2010;463(7280):545-8.
52.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-
source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82.

Figure legends

Fig 1. Extracellular and intracellular Toll signaling components contributes to *Ras^V12*; *scrib^{-/-}* tumour growth.

(A) A transcriptomics analysis comparing the profile of *Ras^V12; scrib^{-/-}* tumour cells to *GMR>GFP* control cells in the eye-antennal disc (EAD) identified 364 genes that are upregulated in the tumours. Among them, *PGRP-SA*, *GNBP2*, *easter*, *spz3* and *spz5* are genes taking part in the extracellular machinery leading to Toll pathway activation. PGRP-SA is highlighted in green because it is also found upregulated in two other tumour models. (B) Simplified version of Toll pathway activation upon bacterial infection. PGRP-SA (Peptidoglycan recognition protein SA) is a carboxypeptidase which recognizes and binds to peptidoglycans upon bacterial infection. Follows the formation of a trimeric complex together with a member of the GNBP family (Gram Negative Binding Protein) and the serine protease (ModSP). This complex initiates a multi-step proteolytic cascade that ultimately leads to the cleavage and activation of the ligand Spz. After the binding of Spz to the Toll receptor, the pathway is activated intracellularly through the activation and formation of a complex composed of the adaptor proteins Myd88 and Tube as well as the serine-threonine protein kinase Pelle. Pelle subsequently phosphorylates the inhibitor cactus, the homolog of mammalian IkappaB, which is targeted to proteasome degradation. As Cactus sequesters the NF-kB Dorsal into the cytoplasm, its degradation triggers the translocation of Dorsal into the nucleus and the transcription of its target genes. (C) Cartoon of the genetic setting used in this study for manipulating *Ras^V12; scrib^{-/-}* tumours. GFP-labelled tumours are generated from randomly-selected single epithelial cells from the EAD. They grow and fuse to form large tumours that start invading the central nervous system (CNS) through the Optic Lobe (OL) first and the Ventral Nerve Cord (VNC) later. Tumour cells also sometimes reach the leg discs (LD). (D) Representative confocal pictures and quantifications of the mean tumour volumes of GFP-labelled control tumours (n=21, m=3,71x10^7 µm^3, SD=±0,47x10^7 µm^3) and *PGRP-SA Ras^V12; scrib^{-/-}* tumours (n=17, m=2,11x10^7 µm^3, SD=±0,19x10^7 µm^3) at day8 after egg laying, statistical significance was determined with an unpaired T-test with Welch's correction. (E) Representative confocal pictures and quantifications of the mean tumour volumes of GFP-labelled control (n=20, m=8,46x10^7 µm^3, SD=±1,77,53x10^7 µm^3) and *ModSP^{IR} Ras^V12; scrib^{-/-} Dcr2* tumours (n=30, m=5,07x10^7 µm^3, SD=±1,00x10^7 µm^3) at Day8 (29°C), statistical significance was determined with an unpaired T-test with Welch's correction. (F) Representative confocal pictures and quantifications of the mean tumour volumes of GFP-labelled control tumours (n=10, m=2,64x10^7 µm^3, SD=±0,76x10^7 µm^3) and *pelle^{IR} Ras^V12; scrib^{-/-}* tumours (n=10, m=0,78x10^7 µm^3, SD=±0,22x10^7 µm^3) at Day6 (29°C), statistical significance was determined with a Mann-Whitney test. Scale bars=50µm.

Fig 2. Two splicing variants of the NF-kB Dorsal are aberrantly expressed in *Ras^V12*-driven tumours.

(A) Immunostainings against Dorsal-A (DlA in green) and Dorsal-B (DlB in magenta) in BFP-labelled *wt* clones (n=20 EADs), *scrib^{IR}* tumours (n=23 EADs), *Ras^V12* tumours (n=20 EADs) and *Ras^V12; scrib^{IR}* tumours (n=24 EADs) at day 6. DlA and DlB accumulation is observed only in *Ras^V12*-driven tumours and appears mainly mutually exclusive. (B) Immunostaining against DlA

(green) and DIB (magenta) in BFP-labelled Ras^{V12}; $scrib^{IR}$ tumours (white) (n=24 EADs) and
close-ups showing that DIA and DIB are not expressed in the same subset of cells. Colocalization
analysis assessed with the Pearson's correlation coefficient reveals a poor colocalization between
DIA and DIB in the tumours (n=24, m=0,11, SD=±0,08). (C) Heat map of DIA intensity within a
representative BFP-labelled Ras^{V12}; $scrib^{IR}$ tumour-bearing EAD at day 6. The tumour is outlined
with the straight white line. An arbitrary intensity threshold set at 250 (gray line on the graph)
allows the clear segregation of a DIAhigh ("tumour High") and a DIAlow ("tumour Low") cell
population within the tumour. Quantifications of the mean DIA intensity of the "tumour High" (n=23,
m=458,7, SD=±28,4), "tumour Low" (n=23, m=135,2, SD=±7,0) and wt Tumour microenvironment
(TME) (n=23, m=82,2, SD=±6,7) demonstrates a general elevation of DIA levels within the tumour
compared to wt cells. Statistical significance was determined with a Tukey's multiple comparison
test. (D) Heat map of DIB intensity within a representative RasV12; scribIR tumour-bearing EAD
at day 6. The tumour is outlined with the straight white line. An arbitrary intensity threshold set at
10000 (gray line on the graph) allows the clear segregation of a DIBhigh ("tumour High") and a
DIBlow ("tumour Low") cell population within the tumour. Quantifications of the mean DIB intensity
of the "tumour High" (n=24, m=20331,9, SD=±3108,7), "tumour Low" (n=24, m=3955,8,
SD=±584,5) and wt Tumour microenvironment (TME) (n=23, m=4660,9, SD=±1187,3).
Normalization was done by subtracting the background intensity. Statistical significance was
determined with a Dunn's multiple comparison test. Scale bars=20µm.

Fig 3. Cell type-dependent Dorsal isoforms accumulation correlates with tumour
neurogenesis and apoptosis.

(A) Representative confocal pictures of BFP-labelled Ras^{V12}; $scrib^{IR}$ tumours, expressing the a
DIA::GFP fusion protein at endogenous levels (green), overtime (Day 4 n=18 EADs, Day 5 n= 14
EADs, Day 6 n=18 EADs, Day 7 n= 17 EADs, Day 8 n=20 EADs and Day 9 n=11 EADs) stained
for the pan-neuronal marker Elav (magenta). The cephalic complex is outlined with the straight
gray line and the tumour is outlined with the straight white line on the close-ups. Yellow arrows
points at neurons that are negative for DIA. Scale bar=50µm. (B) Representative confocal pictures
of GFP-labelled $cherry^{IR}$ Ras^{V12}; $scrib^{IR}$ (n=8 EADs) at day 6 (29°C) stained for Elav (green) and
DIA (magenta). The tumour is outlined with the straight white line. DIA and Elav appears mutually
exclusive. Scale bar=20µm. (C) Representative confocal pictures of GFP-labelled $cherry^{IR}$
Ras^{V12}; $scrib^{IR}$ (n=8 EADs) at day 6 (29°C) stained for Elav (green) and DIB (magenta). The
tumour is outlined with the straight white line. DIB and Elav are expressed in the same tumour
cells. Scale bar=20µm. (D) Representative confocal pictures of GFP-labelled $cherry^{IR}$ Ras^{V12};
$scrib^{IR}$ control tumours (n=62 EADs) at day 6 (29°C) stained for Elav (green) and cDcp1 (magenta).
The tumour is outlined with the straight white line. cDcp1$^+$ cells are mostly found in areas of the
tumour where neurons are present where we often observe neuron apoptosis (yellow arrows).
Scale bar=20µm. (E) Representative confocal pictures of GFP-labelled Ras^{V12}; $scrib^{IR}$ control
tumours (n=7 EADs) and $Diap1^{OE}$ Ras^{V12}; $scrib^{IR}$ (n=9 EADs) stained for cDcp1 (magenta) at day
6. Scale bar=50µm. (F) Representative confocal pictures of GFP-labelled Ras^{V12}; $scrib^{IR}$ control
tumours (n=13 EADs) and $Diap1^{OE}$ Ras^{V12}; $scrib^{IR}$ (n=7 EADs) stained for DIA (green) and DIB
(magenta) at day 6. Scale bar=20µm.

Fig 4. Dorsal represses Ras^{V12} $scrib^{RNAi}$ tumour differentiation and apoptosis.

(A) Representative confocal pictures and quantifications of the mean tumour volumes of GFP-
labelled *Cherry^{IR} Ras^{V12}; scrib^{IR}* control tumours (n=36, m=8,32x10^{7}µm^3, SD=±1,31 x10^{7}µm^3),
dorsal^{IR-1} Ras^{V12}; scrib^{IR} tumours (n=25, m=1,97x10^{7}µm^3, SD=± 1,00x10^{7}µm^3), *Cherry^{IR} Ras^{V12};*
scrib^{IR} Dcr2 control tumours (n=43, m=8,52x10^{7}µm^3, SD=±1,64x10^{7}µm^3) and *dorsal^{IR-2} Ras^{V12};*
scrib^{IR} Dcr2 tumours (n=37, m=4,71x10^{7}µm^3, SD=±1,85x10^{7}µm^3) at Day12 (29°C), statistical
significance was determined with an unpaired T-test with Welch's correction. Scale bar=50µm.
(B) Representative confocal pictures and quantifications of the proliferation ratio of GFP-labelled
Cherry^{IR} Ras^{V12}; scrib^{IR} control tumours (n=36, m=1,08%, SD=±0,41%) and *dorsal^{IR-1} Ras^{V12};*
scrib^{IR} tumours (n=35, m=1,39%, SD=±0,83%) at Day8 (29°C). Proliferation was detected through
immunostainings against the mitotic marker pH3 (phospho-Histone 3, in magenta). Here, the
proliferation ratio is defined as the ratio between the pH3 area within the tumour and the area of
the tumour itself. Scale bar=20µm. (C) Representative confocal pictures and quantifications of the
differentiation ratio of GFP-labelled *Cherry^{IR} Ras^{V12}; scrib^{IR}* control tumours (n=63, m=8,29%,
SD=±2,87%) and *dorsal^{IR-1} Ras^{V12}; scrib^{IR}* tumours (n=57, m=24,40%, SD=±7,09%) at Day8
(29°C). Differentiation was detected through immunostainings against the pan-neuronal marker
Elav (magenta). Here, the differentiation ratio is defined as the ratio between the Elav area within
the tumour and the area of the tumour itself. Scale bar=20µm. (D) Representative confocal
pictures and quantifications of the apoptosis ratio of GFP-labelled *Cherry^{IR} Ras^{V12}; scrib^{IR}* control
tumours (n=63, m=0,71%, SD=±0,29%) and *dorsal^{IR-1} Ras^{V12}; scrib^{IR}* tumours (n=57, m=1,45%,
SD=±0,46%) at Day8 (29°C). Apoptosis was detected through immunostainings against the
cleaved effector caspase, cDcp1 (cleaved-Death Caspase 1, in magenta). Here, the apoptosis
ratio is defined as the ratio between the cDcp1 area within the tumour and the area of the tumour
itself. Statistical significance was determined with an unpaired T-test with Welch's correction.
Scale bar=20µm.

Fig 5. Dorsal amplifies JNK signaling

(A) Representative confocal pictures of BFP-labelled *Ras^{V12}; scrib^{IR}* tumours, expressing the JNK
activity reporter, TRE-eGFP (green) and stained for DIA (magenta). The tumour is outlined with
the straight white line. Colocalization analysis assessed with the Pearson's correlation coefficient
reveals a weak colocalization between DIA and JNK activity in the tumours (n=31, m=0,23,
SD=±0,08). Scale bar=20µm. (B) Representative confocal pictures of GFP-labelled *cherry^{IR}*
Ras^{V12}; scrib^{IR} tumours stained for pJNK (green) and DIB (magenta) at day6 (29°C). The tumour
is outlined with the straight white line. Colocalization analysis assessed with the Pearson's
correlation coefficient reveals a good colocalization between DIB and pJNK in the tumours (n=25,
m=0,41, SD=±0,06). Scale bar=20µm. (C) Representative confocal pictures and quantifications
of the mean MMP1 intensity of GFP-labelled *Ras^{V12}; scrib^{IR}* tumours (n=11, m=1,00, SD=±0,06),
dorsal^{OE} Ras^{V12}; scrib^{IR} tumours (n=11, m=2,85, SD=0,57) and *modSP^{OE} Ras^{V12}; scrib^{IR}* tumours
(n=10, m=1,49, SD=±0,32) at day 6. Intensity normalization was done by subtracting the
background intensity on each picture and finally dividing each data point by the mean intensity
value for the control tumours for each replicate. Statistical significance was determined with
Dunnett's T3 multiple comparisons test. Scale bars=50µm.

Fig 6. Dorsal stimulates tumour migration and the Dorsal targets Snail and Twist contribute
to tumour growth

(A) Z-projections of representative confocal pictures of size-matched GFP-labelled *Cherry^{IR}*
Ras^{V12}; scrib^{IR} tumours (n=3 EADs) and *dorsal^{IR-1} Ras^{V12}; scrib^{IR}* tumours (n=8 EADs) at day 12
(29°C), stained for Hoechst (magenta). Scale bars=50µm. The cephalic complexes (and leg discs

on the small inserts) are outlined with the straight gray line. A cartoon represents the three
invasion parameters assessed in this study: the VNC (Ventral Nerve Cord) invasion score, ranged
between c and 1, is greater the further tumour cells migrate toward the tip of the VNC (green
arrow), indicated as "VNC=" on the Z-projection images; the frequency of leg discs invasion (small
insert with leg discs confocal pictures, Scale bars=20µm) and the fusion of distinct EAD (Eye-
Antennal Disc) tumours (white arrow, dashed line indicates the visible separation of both neighbor
EADs). For the VNC score, statistical significance was determined with Dunnett's T3 multiple
comparisons test. (B) Z-projections of representative confocal pictures of GFP-labelled and
Ras^{V12}; $scrib^{IR}$ (n=15 EADs) and $dorsal^{OE}$ Ras^{V12}; $scrib^{IR}$ tumours (n=24 EADs) at day 6, stained
for Hoechst (magenta). The cephalic complexes (and leg discs on the small inserts) are outlined
with the straight gray line. Quantification of the mean tumour volumes of GFP-labelled Ras^{V12};
$scrib^{IR}$ (n=9, m=9,71x10^6µm^3, SD=±1,53x10^6µm^3) and $dorsal^{OE}$ Ras^{V12}; $scrib^{IR}$ tumours (n=11,
m=6,52x10^6µm^3, SD=±1,12x10^6µm^3) at day 6. OL: Optic Lobe. Statistical significance was
determined with an unpaired T-test with Welch's correction. Scale bars=50µm. (C) Representative
confocal pictures and quantification the mean tumour volume of GFP-labelled $cherry^{IR}$ Ras^{V12};
$scrib^{IR}$ (n=12, m=4,83x10^5µm^3, SD=±1,52x10^5µm^3), and $snail^{IR}$ Ras^{V12}; $scrib^{IR}$ (n=9,
m=1,41x10^5µm^3, SD=±0,65 x10^5µm^3). Statistical significance was determined with a Mann
Whitney's test. (D) Representative confocal pictures and quantification the mean tumour volume
of GFP-labelled $cherry^{IR}$ Ras^{V12}; $scrib^{IR}$ (n=20, m=7,17x10^7µm^3, SD=±1,22x10^7µm^3) and $twist^{IR-1}$
Ras^{V12}; $scrib^{IR}$ (n=7, m=1,81x10^7µm^3, SD=±0,83x10^7µm^3) as well as $Cherry^{IR}$ Ras^{V12}; $scrib^{IR}$ $Dcr2$
control tumours (n=24, m=6,89x10^7µm^3, SD=±1,12x10^7µm^3) and $twist^{IR-2}$ Ras^{V12}; $scrib^{IR}$ $Dcr2$
tumours (n=13, m=3,98x10^7µm^3, SD=±1,43x10^7µm^3). VNC invasion scores are indicated.
Statistical significance was determined with a Mann Whitney's test. Scale bars=50µm.

Fig 7. Working Model

In this working model, we propose that the protein level of the NF-kB Dorsal regulate different
aspect of tumorigenesis. Invasion: In the most posterior part of the tumour, high levels of Dorsal
promote invasion of the tumour cells in the adjacent organs through elevation of JNK signaling
and the EMT transcription factor Snail and Twist that would promote cytoskeleton remodelling.
Tumour growth: In the rest of the tumour where more intermediate levels of Dorsal are observed,
Dorsal would stimulate tumour growth by repressing differentiation and apoptosis. Regulation: At
least two factors seem two trigger aberrant level of Dorsal within the tumours. Genetically, Ras^{V12}
make tumour cells competent to accumulate higher levels of Dorsal throughout the entire tumour.
Locally, a spatiotemporal control is exerted at the posterior part of the tumour where neurons are
formed, to reach very high levels of Dorsal. A possibility is an increase local Toll pathway
activation triggered by high Spz concentration from the brain which will promote Dorsal expression.
There, we observe alternative splicing of DIA versus DIB which segregate in different tumour
subpopulations. We hypothesize that the heterogeneity in Dorsal isoforms and/or levels promote
a balance between tumour growth and invasion favoring tumorigenesis.

S1 Fig. Localization of DIA, DIB and Cactus.

(A) Representative confocal pictures of GFP-labelled Ras^{V12}; $scrib^{-/-}$ tumours (n=10 EADs) at day
6, generated with the conventional EyMARCM genetic system and stained for DIA (magenta) and
Hoechst (blue). Consistent with the data generated with our new EyaHost genetic system in Fig
2, DIA levels are aberrant in the tumours. The tumour is outlined with a straight white line. Scale
bars=50µm. (B) Representative confocal pictures of GFP-labelled Ras^{V12}; $scrib^{IR}$ tumours at day
6, generated with our novel EyaHost genetic system and stained for DIA (green), DIB (magenta)

and Hoechst (blue). The tumour cells that already migrated inside the optic lobe (OL) of the CNS
express high levels of DIA or DIB, suggesting their potential involvement in tumour cell migration.
Scale bars=50μm. (C) Representative confocal pictures of BFP-labelled *genomic-cactus::GFP,*
Ras^{V12}; scrib^{IR} tumours stained for DIB (magenta) at day 6 (upper line) (n=6 EADs) or stained for
DIA (magenta) at day 8 (lower line) (n=6 EADs). The tumour is outlined with a straight white line.
Scale bars=20μm.

S2 Fig. The endogenous DI reporter used in this study is a DIA specific reporter.

(A) Representative confocal pictures of BFP-labelled *genomic-dorsal::GFP Ras^{V12}; scrib^{IR}*
tumours at day 6, stained for DIA (magenta) and DIB (blue). The tumour is outlined with the
straight white line. Scale bars=50μm. (B) Colocalization analyses assessed with the Pearson's
correlation coefficient reveal a good colocalization between DIA and Dorsal::GFP (n=24, m=0,44,
SD=±0,06) and no colocalization between DIB and Dorsal::GFP in the tumours (n=24, m=0,09,
SD=±0,07) suggesting that this reporter is a DIA specific fusion protein. Statistical significance
was determined with an unpaired T-test with Welch's correction.

S3 Fig. Dorsal knockdown efficiencies and pelle knockdown mimicks the function of
Dorsal on differentiation and survival.

(A) Representative confocal pictures and quantification of the mean DIA intensity of GFP-labelled
cherry^{IR} Ras^{V12}; scrib^{IR} (n=6, m=1,00, SD=±0,14) and *dorsal^{IR-1} Ras^{V12}; scrib^{IR}* tumours (n=6,
m=0,54, SD=±0,03) as well as *Cherry^{IR} Ras^{V12}; scrib^{IR} Dcr2* control tumours (n=5, m=1,00,
SD=±0,14) and *dorsal^{IR-2} Ras^{V12}; scrib^{IR} Dcr2* tumours (n=6, m=0,32, SD=±0,05) at day 6 (29°C).
Intensity normalization was done by subtracting the background intensity on each picture and
finally dividing each data point by the mean intensity value for the control tumours for each
replicate. Statistical significance was determined with a Mann-Whitney test. Scale bars=20μm.
(B) Representative confocal pictures and quantification of the mean DIB intensity of GFP-labelled
cherry^{IR} Ras^{V12}; scrib^{IR} (n=7, m=1,00, SD=±0.19) and *dorsal^{IR-1} Ras^{V12}; scrib^{IR}* tumours (n=9,
m=0.40, SD=±0.12) as well as *Cherry^{IR} Ras^{V12}; scrib^{IR} Dcr2* control tumours (n=5, m=1.00,
SD=±0.31) and *dorsal^{IR-2} Ras^{V12}; scrib^{IR} Dcr2* tumours (n=6, m=0.54, SD=±0.15) at day 6 (29°C).
Intensity normalization was done by subtracting the background intensity on each picture and
finally dividing each data point by the mean intensity value for the control tumours for each
replicate. Statistical significance was determined with a Mann-Whitney test. Scale bars=20μm.
(C) Z-projection of representative confocal pictures of GFP-labelled *RFP^{IR} Ras^{V12}; scrib^{IR}* control
tumours and *pelle^{IR} Ras^{V12}; scrib^{IR}* tumours at day 6 (29°C), stained for Elav (green) and cDcp1
(magenta). The tumour is outlined with a straight white line. Quantification of the apoptosis ratio
of *RFP^{IR} Ras^{V12}; scrib^{IR}* control tumours (n=16, m=4,97%, SD=±2,23%) and *pelle^{IR} Ras^{V12}; scrib^{IR}*
tumours (n=15, m=17,84%, SD=±4,16%) at day 6 (29°C). Here, the apoptosis ratio is defined as
the ratio between the cDcp1 area within the tumour and the area of the tumour itself. Statistical
significance was determined with an unpaired T-test with Welch's correction. Quantification of the
differentiation ratio of *RFP^{IR} Ras^{V12}; scrib^{IR}* control tumours (n=5, m=18,04%, SD=±5,99%) and
pelle^{IR} Ras^{V12}; scrib^{IR} tumours (n=8, m=44,14%, SD=±8,83%) at day 6 (29°C). Here, the
differentiation ratio is defined as the ratio between the Elav area within the tumour and the area
of the tumour itself. Statistical significance was determined with a Mann-Whitney test. Scale
bars=50μm.

S4 Fig. Dorsal is sufficient to induce dysplasia and delamination but not overgrowth and
invasion.

(A) Representative confocal pictures and quantification of the mean clone volume of GFP-labelled
wt clones (n=20, m=5,59x10^{5}µm^3, SD=±0,69x10^{5}µm^3) and *dorsal^{OE}* clones (n=30,
m=4,87x10^{5}µm^3, SD=±0,66x10^{5}µm^3) at day 6, stained for DIA (magenta) and Hoechst (blue).
Displasic clones are observed in 100% of the *dorsal^{OE}* clones-bearing EADs. GFP-labelled cells
are never found in the VNC as indicated with the VNC score values. Scale bars=50µm. (B)
Reconstruction of an orthogonal view of *wt* clones and *dorsal^{OE}* clones representing the EAD
tissue (magenta) and the GFP-labelled clones (green). GFP-labelled cells extend from the apical
to the basal side of the pseudostratified epithelium of the EAD of *wt* clones whereas some
Dorsal^{OE} clones spread basally (yellow arrow).

Figure

Figure

Figure

A *Ras^V12; scrib IR>TRE-eGFP*

TRE-eGFP DIA

TRE-eGFP

DIA

DIA/ TRE-eGFP
colocalization

B *Ras^V12; scrib IR>Cherry IR*

pJNK DIB

pJNK

DIB

DIB/ pJNK
colocalization

C *Ras^V12; scrib IR>*

+   *dorsal OE*   *modSP OE*

GFP MMP1 Hoechst

MMP1

Normalized MMP1
Intensity

Figure

MIGRATION

A *Ras^{V12}; scrib IR>*

cherry IR   *dorsal IR-1*

VNC Invasion score
0
0,25
0,5
0,75
1

MaxZ-Projection

GFP Hoechst

OL
VNC

OL
VNC

VNC=0,67   ** VNC=0,22

LD   LD

78%   11%

B *Ras^{V12}; scrib IR>*

+   *dorsal OE*

MaxZ-Projection

EAD
OL
VNC

EAD
OL
VNC

VNC=0,22   *VNC=0,45

Tumour size

Tumour volume (μm^3)

1,5x10^7
1x10^7
5x10^6
0

*

EMT TRANSCRIPTION FACTORS

C *Ras^{V12}; scrib IR, Dcr2>*

cherry IR   *snail IR*

GFP

Tumour size

Tumour volume (μm^3)

1x10^6
5x10^5

**

D *Ras^{V12}; scrib IR>*   *Ras^{V12}; scrib IR, Dcr2>*

cherry IR   *twist IR-1*   *cherry IR*   *twist IR-2*

VNC=0,88   ****VNC=0,25   VNC=0,93   **VNC=0,61

GFP

Tumour size

Tumour volume (μm^3)

1,5x10^8
1x10^8
5x10^7
0

****   **

Figure

A

WORKING MODEL

NF-kB INTRATUMORAL HETEROGENEITY

Tumour Growth

Dorsal
→ Differentiation
→ Cell Death

Invasion

High Dorsal

Snail /Twist → JNK

Migration

Regulation

GENETIC   LOCAL

Ras^V12   Neurons   Toll activating signals?

Alternative splicing
Aberrant levels

Dorsal-A   Dorsal-B
Cactus

Figure

Figure