Splenic follicular B cells promote adverse cardiac remodeling after myocardial infarction via MHC II-dependent antigen presentation.

Obialunanma V. Ebenebe PhD[1*], Jana P. Lovell MD[1*], Carolina Duque BS[2], Yiran Song MSc[1], Sylvie Rousseau BA[1], Kshipra Keole BA[1], Sogol Sedighi MD[1], Aashik Bhalodia BS[1], Kevin Bermea MD[1], Charles D. Cohen PhD[1], Monica R. Mugnier PhD[2], D. Brian Foster PhD[1], Scott Krummey MD PhD[3], Simone Sidoli PhD[4], Luigi Adamo MD PhD[1]

[1]Division of Cardiology, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland.
[2]W. Harry Feinstone Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School of Public Health
[3]Department of Pathology, Johns Hopkins University School of Medicine, Baltimore, Maryland
[4]Department of Biochemistry, Albert Einstein College of Medicine, Bronx, New York

*= these two authors share first authorship

Sources of Funding

Disclosures

L.A. is co-founder of i-Cordis, LLC, a startup company focused on the development of immunomodulatory therapies for heart failure, and he is a consultant for Novo Nordisk and Kiniksa Pharmaceuticals.

Address of Correspondence:

Luigi Adamo, MD, PhD

Division of Cardiology
Johns Hopkins University School of Medicine
720 Rutland Avenue
Ross Research Building 809
Baltimore, MD 21205
Phone: 410-502-2857
Fax: 410-955-1962
@luigiadamomdphd

Abstract

The cardio-splenic axis is a promising therapeutic target in ischemic heart failure (HF), but

splenic–cardiac interactions after myocardial infarction (MI) are poorly understood. Here, we

show that splenic follicular B cells drive ischemic HF via MHC class II–mediated antigen

presentation. Transfer of splenic B cells from mice with ischemic HF induced adverse cardiac

remodeling and modulated myocardial T-cells in naïve recipients. Single-cell RNA sequencing of

mouse and human post-MI B cells revealed upregulation of antigen-presentation related pathways.

Mass spectrometry of the MHC II peptidome demonstrated enrichment of cardiac-derived peptides

in MHC II molecules of splenic B cells from ischemic HF mice. B cell-specific deletion of MHC

II suppressed adverse cardiac remodeling after adoptive transfer of post-MI splenic B cells. These

results broaden our understanding of B-lymphocyte biology and point towards MHC II-mediated

signaling in B-cells as a novel therapeutic target in ischemic HF.

Acute myocardial infarction (MI) triggers a robust activation of inflammatory responses [1, 2].

These inflammatory responses are physiological and necessary to limit tissue injury and induce

tissue repair [3, 4]. However, sustained inflammation contributes to adverse cardiac remodeling

and thus, to the development and progression of ischemic heart failure [5]. Despite advancements

in our understanding of the relationship between chronic inflammation and heart failure (HF),

clinical trials investigating immunomodulatory therapies in HF have yielded largely unsuccessful

results. The failure of these therapies is partly due to the inability to selectively target maladaptive

rather than protective inflammatory responses post-MI [6].

As such, an improved understanding of the sustained maladaptive inflammatory responses

contributing to the pathogenesis of chronic HF is needed. Prior murine and human data indicate

that acute MI induces splenic remodeling and mobilization of immune cells from the spleen to the

injured myocardium [7]. Activation of this "cardio-splenic axis" has been shown to contribute to

adverse cardiac remodeling and the development of chronic HF [8]. Thus, the cardio-splenic axis

has been identified as a promising therapeutic target for developing immunomodulatory treatments

for HF. However, our mechanistic understanding of the processes by which specific splenic

immune cells contribute to adverse cardiac remodeling remains limited.

B cells comprise the majority of immune cells within the spleen of most vertebrates. B cells are

also prevalent in healthy hearts and can recirculate between the spleen and the heart [9-11].

Growing evidence shows that B cells are intricately involved with cardiac function and

dysfunction, and following acute MI, contribute to the development and progression of ischemic

HF [12-16]. However, to date, the precise role of B cells within the cardio-splenic axis in HF has

not been systematically investigated.

Here, we show that B cells, specifically splenic follicular B cells, play an essential role within the

cardio-splenic axis of ischemic heart failure by promoting adverse cardiac remodeling post-MI via

MHC II-dependent antigen presentation.

Results

Splenic B cells contribute to adverse cardiac remodeling in ischemic heart failure.

To investigate whether splenic B cells are involved in the adverse cardiac remodeling of ischemic

HF, we designed adoptive transfer experiments. Unfractionated splenocytes or purified splenic B

cells were transferred from HF mice into naïve mice via intravenous injection. Splenic immune

cells isolated from sham-operated animals were used as controls. The naïve recipient mice were

monitored for eight weeks after adoptive transfer (Fig.1a).

Consistent with previously published data [8], we found that adoptive transfer of splenocytes from

mice with ischemic HF resulted in adverse cardiac remodeling in recipient mice. From serial

echocardiography, recipients of splenocytes from mice with HF had reduced left ventricular

ejection fraction (LVEF) and increased left ventricular end-systolic diameter (LVESD) eight

weeks post adoptive transfer compared to recipients of splenocytes from sham-operated mice

(Fig.1b; mean LVEF 68.1±1.7% vs 77.0±1.3% HF vs. sham recipient, q<0.001 at 8 weeks; mean

LVESD 2.01±0.7mm vs. 1.67±0.02mm HF vs. sham recipient, q<0.001 at 8 weeks).

To investigate the specific role of splenic B cells within the cardio-splenic axis following ischemic

myocardial injury, we performed adoptive transfer of purified splenic B cells from HF or sham-

operated mice (≈95% isolated B cell purity, Supplementary Fig.1). Over the 8-week period, the

adoptive transfer of isolated splenic B cells from mice with HF produced a degree of adverse

remodeling similar to that observed with the adoptive transfer of unfractionated splenocytes

(Fig.1b, bottom panels). Recipients of splenic B cells from mice with HF experienced a progressive

reduction in LVEF and progressive LV dilatation, detected at 4- and 8-weeks post adoptive transfer

(Fig.1b; mean LVEF 67.6±1.6% vs. 77.3±1.3% HF vs. sham recipient, q<0.001 at 8 weeks; mean

LVESD 2.00±0.05mm vs. 1.66±0.06mm HF vs. sham recipient, q<0.001 at 8 weeks). Further

experiments wherein we extended the post adoptive transfer monitoring revealed that this effect

was progressive to 16-weeks post adoptive transfer (Supplementary Fig.2).

Adverse cardiac remodeling in naïve recipients of splenocytes and splenic B cells isolated from

HF mice was further indicated by gravimetric analyses of the heart and spleen, as well as wheat-

germ agglutinin (WGA) staining of myocardial cross-sections. Gravimetric analysis revealed

increased splenic and cardiac mass in recipients of splenic B cells from mice with HF compared

to recipients of B cells from sham-operated mice 8 weeks following adoptive transfer (Fig.1c; for

purified splenic B cells: spleen mean 3.11±0.4 mg/g vs 2.59±0.2 mg/g, p = 0.03; heart mean

4.85±0.05 mg/g vs 4.60±0.2 mg/g, p = 0.03; Supplementary Fig.3 displays the combined

gravimetric data from multiple independent experiments). Furthermore, WGA staining of

myocardial cross-sections showed that increased heart mass was associated with cardiomyocyte

hypertrophy (Fig.1d; p = 0.002).

Adoptively transferred splenic B cells are distributed between the spleen and heart.

To investigate the dynamics of adoptively transferred splenic B cells, we repeated the adoptive

transfer experiments using animals carrying CD45.1 isoform as donors and animals carrying

CD45.2 isoform as recipients (Fig.2a). Donor CD45.1 B cells were present in the peripheral blood

of naïve recipient mice at 1- week and 8- weeks following adoptive transfer (Fig.2b; ~3% and ~2%

of circulating donor-derived B cells at 1- and 8- weeks, respectively). Flow cytometry of the spleen

and heart 8 weeks post-adoptive transfer also revealed the presence of donor CD45.1 B cells,

indicating successful engraftment of CD45.1 B cells in recipient mice (Fig.2b; Supplementary

Fig.4 also includes non-adoptive transfer controls).

**Ischemic myocardial injury results in differential expression of antigen processing and
presentation pathways in splenic B cells.**

To explore the mechanistic relationship between the adoptive transfer of splenic B cells from mice

with ischemic HF and adverse cardiac remodeling in naïve mice, we first analyzed the recipients'

spleen and heart via high-dimensional flow cytometry. At the splenic level, adoptive transfer of

splenic B cells from mice with ischemic HF did not affect the numbers of splenic lymphocytes or

the prevalence of B cell subsets (Fig.2c, Extended data Figure 1); nor were there changes in splenic

CD3 T cells, CD4 T cells, and CD8 T cells numbers relative to recipients from sham-operated

mice (Fig.2c). Closer examination of splenic T cell subsets indicated minor differences. We noted

a downward trend in the percentage of memory CD4 T cells expressing CTLA-4 and FOXP3

($CD44^{hi}$, $CD62L^{lo}$, CTLA-4$^+$, FOXP3$^+$) and a corresponding increase in memory CD4 T cells

negative for CTLA-4 and FOXP3 markers ($CD44^{hi}$, $CD62L^{lo}$, CTLA-4$^-$, FOXP3$^-$, Extended Data

Fig.2). We did not detect differences in subsets of splenic CD8 T cells (Extended Data Fig.3). At

the myocardial level, via flow cytometry, we did not detect differences in cardiac immune cell

populations between recipients of splenic B cells from mice with ischemic HF and recipients of

splenic B cells from sham-operated mice (Extended Data Fig.4).

Since this analysis was unrevealing, we focused our attention on the donors; the population of

splenic B cells that were being adoptively transferred. We first used high-dimensional flow

cytometry. Within the spleen, B cells can be sub-classified based on specific cell markers [17, 18].

There were no changes in the percentage of splenic B cell subtypes isolated from mice with

ischemic HF (4 weeks post-MI surgery) compared to sham-operated mice (Extended Data Fig.5).

We then used single-cell RNA sequencing (scRNAseq) to more deeply probe the molecular

character of the donor splenic B cells from HF mice (Fig.3a). V(D)J analysis did not reveal an

expansion of specific splenic B cell clonotypes in ischemic HF (4 weeks post-MI). Suggesting that

the effects observed at the time of adoptive transfer were not due to splenic B cells' clonal

expansion (Fig.3b; Supplementary Fig.5). UMAP plots were then used to visualize the clustering

of gene expression profiles of splenic B cells (Fig.3c). Comparative gene expression analysis

revealed several genes differentially expressed in splenic B cells from mice with ischemic HF (4

weeks post-MI surgery) (Supplementary Table 1).

To evaluate the biological relevance of these differentially expressed genes (DEGs), we performed

pathway analyses using KEGG pathway annotation [19, 20]. Antigen processing and presentation

was the top biological pathway identified as differentially regulated (q-value = 8.02E-9; Fig.3d).

Some of the specific DEGS included HSPA8, HSP90AA1, HSP90AB1, which encode heat shock

proteins (HSPs) involved in MHC-II mediated antigen presentation [21, 22]; H2-Eb1 and H2-

DMb1 which encode MHC-II proteins and IFI30 (also known as GILT) which is involved in MHC-

II mediated antigen presentation [23]. Further evaluation of B cell subtypes revealed significant

differential regulation of the antigen processing and presentation pathway in the follicular,

germinal center, regulatory, plasma, and memory B cells (Fig.3d) but not in marginal zone B cells

(Supplementary Table 2).

To further corroborate these findings, we expanded our analysis to myocardial B cells and

peripheral blood B cells in animals with ischemic HF (Supplementary Figs.6 and 7). We also

included animals with recent MI (four days post-MI surgery) to gain longitudinal insight

(Supplementary Figs.8-10). These analyses also pointed towards antigen processing and

presentation as one of the top sustained, differentially regulated pathways in myocardial and

circulating B cells in response to myocardial injury. More specifically, antigen processing and

presentation was differentially regulated in multiple cardiac and peripheral blood, B cell subtypes

from mice with ischemic HF (Supplementary Figs.6 and 7), with an especially strong signal in

myocardial and circulating follicular B cells (Supplementary Figs.6C and 7C). Recent MI was

compared to the 8-weeks post sham surgery (which is a suboptimal control); yet this analysis also

highlighted significant differential regulation of antigen processing and presentation in splenic,

cardiac, and peripheral blood B cells with a predilection for follicular B cells (Supplementary

Figs.8-10). Lists of the DEGs associated with antigen processing and presentation in mice with

ischemic HF and mice with recent MI are provided within the spreadsheets included in

Supplementary Table 1.

MHC II molecules of splenic B cells from mice with ischemic HF are loaded with cardiac

peptides.

Collectively, our transcriptomic data suggested that splenic B cells might modulate adverse cardiac

remodeling in ischemic HF via major histocompatibility complex class II (MHC II) mediated

antigen presentation. This was an unexpected finding. B cells have three main functions: antibody

production, cytokine secretion, and antigen presentation. It has been reported that B cells may

contribute to adverse cardiac remodeling following MI via the production of autoantibodies or

chemokine-mediated mobilization of monocytes from the spleen to the heart [12, 24]. However,

to our knowledge, B cell-mediated antigen presentation has never been implicated in the

pathogenesis of cardiac dysfunction. To further investigate the potential role of MHC II-mediated

antigen presentation in ischemic myocardial injury, we isolated MHC II-associated peptides from

splenic B cells of sham-operated and ischemic HF mice (4 weeks post-MI) and analyzed them via

mass spectrometry (Fig.3e). Peptides enriched more than 1.2-fold in MHC II molecules from post-

MI B cells were analyzed *in silico* with a software that predicts peptide binding to the MHC II

isoform expressed on C57BL/6 mice. 54 enriched peptides were identified as predicted binders in

the *in-silico* analysis, corresponding to 45 proteins. 18% of these proteins were cardiac-specific

(Fig.3f). 13 peptides were predicted as strong binders. These included several cardiac-specific

proteins such as cardiac α actin (ACTC1), Titin (TTN) and 2 peptides from myosin heavy chain 6

(MYH6) (Fig.3f bottom panel, Supplementary Table 3).

B cell-specific deletion of MHC II suppresses adverse cardiac remodeling after adoptive
transfer of post-MI splenic B cells.

To then investigate B cell-dependent MHC II-mediated antigen presentation as a possible

mechanistic driver of adverse cardiac remodeling following ischemic myocardial injury, we used

transgenic mice with B cell-specific MHC II deletion ($Cd19^{tm1(cre)Cgn/-}H2-Ab1^{b-tm1Koni/\,b-tm1Koni}$) [25]

and repeated our adoptive transfer experiments (Fig.4a).

Four weeks following MI surgery, we performed adoptive transfer of splenic B cells from MHC

II-deficient HF mice into naïve wild-type (WT) mice. Flow cytometry analysis confirmed the

persistent absence of MHC II on donor B cells at the time of adoptive transfer (Supplementary

Fig.11). In a separate experiment, this time using CD45.1 mice as recipients, because the donor

MHC II-deficient and WT mice naturally express CD45.2, we confirmed the persistent presence

of donor B cells from MHC II-deficient mice 8 weeks following adoptive transfer (Supplementary

Fig.12). Adoptive transfer of splenic B cells from WT ischemic HF mice or sham-operated mice

served as positive and negative controls, respectively. Splenic B cells isolated from MHC II-

deficient mice with ischemic HF did not induce adverse cardiac remodeling upon adoptive transfer

as assessed by echocardiography (Fig.4b; mean LVEF 87.10±1.9% vs. 88.96±2.2% MHC II-

deficient HF vs. WT sham recipient, q = 0.11 at 8 weeks; mean LVESD 1.27±0.1mm vs.

1.19±0.1mm MHC II-deficient HF vs. WT sham recipient, q = 0.17 at 8 weeks). Gravimetric data

revealed borderline increases in cardiac mass and the absence of splenic hypertrophy following

the transfer of splenic B cells from MHC II-deficient HF mice (Fig.4c; spleen mean 2.86±0.3 vs.

2.83±0.2 mg/g MHC II-deficient HF vs. WT sham recipient, q = 0.83; heart mean 4.54±0.2 vs.

4.34±0.1 mg/g MHC II-deficient HF vs. WT sham recipient, q = 0.05). Cardiomyocyte

hypertrophy was also markedly reduced in recipients of splenic B cells from MHC II-deficient HF

mice compared to recipients of splenic B cells from WT HF mice (Fig.4d).

To reinforce the findings from the echocardiographic analysis, we performed proteomics analyses

of left ventricular samples from recipients of splenic B cells from WT sham, WT ischemic HF and

MHC II-deficient ischemic HF. This analysis confirmed adverse cardiac remodeling in recipients

of splenic B cells from WT HF mice relative to recipients from WT sham mice and highlighted

blunted adverse cardiac remodeling in recipients of splenic B cells from MHC II-deficient HF mice

(Fig.5a-b). There was significant upregulation of key proteins involved in adverse cardiac

remodeling (e.g. TPM and TNNT2), cell death and protein turnover (BAG3 and UBB) in recipients

of WT HF splenic B cells relative to MHC II-deficient HF splenic B cells. Interestingly, in

recipients of MHC II-deficient HF splenic B cells, there was significant upregulation of cardiac

repair/stabilization proteins in response to stress like MYOM1, GYG1, CCT2 and NID-1 (Fig.5c).

Follicular splenic B cells mediate MHC II-dependent cardiac adverse remodeling post-MI

We ran several control experiments to support proper interpretation of the results obtained using

MHC II-deficient B cells. First, we repeated the adoptive transfer experiment utilizing

$Cd19^{tm1(cre)Cgn/-I}$ mice as donors and confirmed that there were no off-target effects of the Cre

transgene (i.e. The results from these experiments were comparable to those using WT controls,

Supplementary Fig.13).

We then performed immunofluorescence to assess potential changes in serum antibodies targeting

myocardial proteins. We incubated heart sections from genetically B-cell deficient μMT mice with

serum collected from recipients of isolated splenic B cells from WT sham, WT HF and MHC II-

deficient HF mice 8 weeks post adoptive transfer and stained the histological sections with anti

IgG and IgM antibodies. We observed no differences in IgG-positive and IgM-positive stain

intensities across groups (Supplementary Fig.14).

Finally, we analyzed splenic B cells from MHC II-deficient ischemic HF (4 weeks post- MI) and

sham-operated mice. WT HF mice and sham-operated mice served as positive and negative

controls, respectively. We found that the deletion of MHC II on B cells was associated with an

increase in the percentage of follicular B cells and a decrease in marginal zone B cells post-MI

(Supplementary Fig.15). This raised the possibility that the deletion of MHC II on B cells might

suppress adverse cardiac remodeling post-adoptive transfer simply due to the decrease in marginal

zone B cells. We, therefore, repeated our adoptive transfer experiments, this time transferring

either follicular B cells or marginal zone B cells from HF or sham-operated mice (Fig.6a).

Splenic follicular B cells promoted adverse cardiac remodeling in naïve mice, evidenced by

reduction in LVEF and increase in LVESD relative to sham (Fig.6b); while marginal zone B cells

had no effect (Fig.6b). This identified splenic follicular B cells as the key mediators of adverse

cardiac remodeling along the cardio-splenic axis post-MI and reinforced a direct effect of B cell

MHC II deletion on B cell-dependent adverse cardiac remodeling post-MI.

Single cell RNAseq based analysis indicates that adoptively transferred splenic B cells from
HF mice modulate myocardial T cells via an MHC-II dependent function.

The results presented so far indicate that in ischemic HF, splenic B cells promote adverse cardiac

remodeling through an MHC II-dependent function, likely MHC II-mediated antigen presentation.

We used scRNA-seq to investigate the effect of adoptive transfer of splenic B cells from HF mice

on the cardiac immune cells of recipients. We isolated myocardial immune cells 8 weeks post-

adoptive transfer of splenic B cells from WT-HF, MHC II-deficient HF or WT sham operated mice

and analyzed them via scRNAseq (Fig.7a-b). We focused our analysis on T cell gene expressions

as B cells are known to present antigens to T cells. Comparative gene expression analysis of

recipients of splenic B cells from WT-HF vs WT sham operated mice revealed a list of 70 DEGs

in myocardial CD4 T cells (12 downregulated, 58 upregulated, adjusted p<0.05) and 745 DEGs in

myocardial CD8 T cells (6 downregulated, 739 upregulated, adjusted p<0.05, Supplementary

Table 4). When comparing recipients of MHC II-deficient HF splenic B cells to recipients of WT-

HF splenic B cells, we found minor differences in CD4 T cells, where 33 DEGs were upregulated

and 15 DEGS downregulated (adjusted p<0.05, Fig.7c, left bottom). The difference was more

pronounced in CD8 T cells in which only 2 genes were upregulated, and 84 genes were

downregulated (Fig.7c, right bottom; Supplementary Table 4). To gain insights into the biological

significance of these gene expression changes we used Ingenuity Pathways Analysis (IPA). IPA

showed that adoptive transfer of splenic B cells from HF mice resulted in upregulation of interferon

gamma (IFNγ) related signaling and other downstream signaling pathways in CD8 T cells

(Supplementary Fig.16). Depletion of MHC II from B cells resulted in downregulation of IFNγ

and STAT5B related signaling in CD8 T cells (Supplementary Fig.16). In CD4 T cells, IPA

showed that deletion of MHC II on B cells resulted in suppression of HSF1 mediated signaling, a

signaling pathway that drives the development of regulatory T cells [26].

Taken together, these observations suggest that post-MI splenic follicular B cells promote adverse

cardiac remodeling in ischemic heart failure through MHC II-dependent modulation of myocardial

T cells.

Ischemic heart failure in humans is associated with differential expression of antigen
processing and presentation pathways in B cells.

To evaluate the clinical relevance of our observations, we next sought to assess differentially

expressed biological pathways in B cells isolated from patients who experienced ischemic

myocardial injury. Since our murine data showed that changes in splenic B cells were reflected in

peripheral blood B cells (Supplementary Fig.7) and isolation of splenic B cells from patients was

not feasible, we performed a focused analysis of publicly available peripheral blood scRNAseq

data from patients with ST-elevation myocardial infarction (STEMI) [27] and patients with

established ischemic HF relative to healthy patients [28]. In both datasets, we identified peripheral

blood B cells and performed differential gene expression analysis.

The STEMI dataset included peripheral blood scRNAseq data of 38 patients 24-hours and 8-weeks

post-STEMI in addition to 38 healthy controls [27]. We performed DEG analysis within B cells,

comparing post-STEMI patients at both timepoints with healthy controls (Fig.8a). We identified

459 DEGs at 24-hours post-STEMI and 493 DEGs 8-weeks post-STEMI. KEGG pathway analysis

of these DEGs highlighted antigen processing and presentation as a differentially regulated

pathway both at 24-hours post-STEMI (q = 0.03) and 8-weeks post-STEMI (q = 0.03; Fig.8b),

with genes such as HLA-DRB5 and CIITA being differentially expressed and matching this

pathway. In the second dataset of three patients with established ischemic cardiomyopathy and

one matched healthy control (Fig.8c) [28], we identified 302 DEGs. Antigen processing and

presentation was identified here as a top differentially regulated pathway in patients with ischemic

HF (q = 2.09E-21; Fig.8d). DEGs relevant to these pathways included HLA-DRB1, HLA-DMA,

HLA-DQA1 and -DQA2, HLA-DQB1, which all encode for various MHC II proteins. Complete

lists of the DEGs within the antigen processing and presentation pathway are provided in

Supplementary Table 5.

Discussion
In this study, we report for the first time that splenic follicular B cells promote adverse cardiac

remodeling in ischemic heart failure. We show that splenic B cells present cardiac antigens on

MHC II molecules after MI and that their role in promoting adverse cardiac remodeling along the

cardio-splenic axis is largely dependent on MHC II. These findings change our understanding of

B cell biology in the post-MI period and point to B cell-mediated antigen presentation as a potential

novel and specific therapeutic target to prevent and treat ischemic heart failure.

Previous research has established that immune activation along the cardio-splenic axis is critical

in ischemic heart failure [7, 8, 29-32]. Although B cells are the most prevalent splenic immune

cell across multiple vertebrates, there is limited data on the role of B cells within the cardio-splenic

axis. To address this knowledge gap, we employed the adoptive transfer model developed by

Ismahil et al., which uniquely allows for the isolation of the functional effects of specific immune

cell types within the cardio-splenic axis of ischemic heart failure [8]. The adoptive transfer of

splenic B cells from mice with ischemic heart failure resulted in adverse cardiac remodeling with

evidence of persistent LV dilatation, reduced LV systolic function, and cardiomyocyte

hypertrophy in naïve mice. Our findings thus provide compelling evidence that B cells play a

significant role within the cardio-splenic axis of ischemic heart failure.

Our data corroborates previous observations regarding the dynamics of B cell transit to the heart

[10], in that adoptively transferred splenic B cells are re-distributed between the heart, spleen, and

peripheral blood of recipient mice, suggesting recirculation of transferred splenic B cells between

the heart and spleen. This does not exclude distribution to other organs, which has indeed been

described in prior studies in naïve mice [10]. The observation that MI triggered the differential

regulation of antigen processing and presentation pathways across splenic, myocardial and

circulating B cells further supports the notion that B cells actively recirculate between the spleen

and the heart through the peripheral blood after MI. Notably, the persistent unique gene expression

profile observed in B cells four weeks post-injury suggests that MI induces long-lasting alterations

in B cells even in the absence of observable changes in B cell frequency.

Analyzing the MHC-II associated peptidome of post-MI splenic B cells we found enrichment of

myocardial antigens. MYH-6, a protein that has been previously described as preferential target of

cardiotropic T cells in mice and humans [33, 34] was one of the strong MHC II binders detected,

at 2 distinct regions (two different MYH-6 peptides were identified as enriched). We did not

observe significant clonal expansion of B cells post-MI. We cannot exclude that this might be the

result of insufficient sequencing depth, yet there were no significant changes in the serum

concentration of cardiac autoantibodies. Data from the RITA-MI trial also suggests a lack of

autoantibody involvement, as serum immunoglobulin levels were not significantly changed from

time of MI (baseline) throughout B-cell depletion (Rituximab treatment) [35]. This does therefore

present the possibility that splenic B cells may acquire and present cardiac antigens through B cell

receptor (BCR)-independent mechanisms rather than through typical BCR engagement [36]. BCR-

independent, low-efficiency, antigen uptake has been well documented in B cells [37]. We

speculate that after an MI, release of high concentration of cardiac autoantigens might generate

appropriate conditions for BCR-independent antigen uptake. Our scRNA data highlights the

involvement of pattern recognition receptor (PRR) pathways; thus, there is a possibility that

activation of damage-associated molecular patterns (DAMPs) related mechanisms may be the

preceding step to loading of cardiac antigens on MHC II molecules of splenic B cells. Further

studies will be needed to understand in detail the mechanistic steps that lead to the loading of

myocardial antigens on MHC II molecules of splenic B cells.

The findings from our adoptive transfer experiments, which involved post-MI B cells with either

intact or deficient MHC II expression, underscore the importance of MHC II in mediating adverse

cardiac remodeling along the cardio-splenic axis. Specifically, the absence of MHC II led to a

reduction in adverse cardiac remodeling in recipient mice, as assessed by echocardiography,

gravimetric analysis, and mass spectrometry of myocardial tissue. While MHC II deletion can

influence various B cell functions, including survival, replication, and chemokine production, our

results suggest that MHC II-deficient B cells can survive after adoptive transfer and maintain

comparable frequencies in recipient animals relative to wild-type B cells. Notably, we found that

alterations in the composition of splenic B cells due to MHC II deletion did not account for the

phenotype observed in recipients of MHC II-deficient splenic B cells.

We cannot entirely rule out the impact of MHC II deletion on the secretion of B cell-specific

chemokines. This is a limitation of our study. Yet, collectively, our findings from the MHC-II

associated peptidome analysis and the experiments with MHC-II deficient B cells strongly suggest

that splenic B cells modulate adverse cardiac remodeling after MI through MHC II-dependent

antigen presentation. It bears emphasis that we noted mild effects of post-MI MHC II-deficient

splenic B cells, with trends toward increased cardiac mass, cardiomyocyte cross-sectional area,

(Figure 4 c,d) and mild changes at the proteomic level (Figure 5), indicating some degree of

adverse cardiac remodeling. This may stem from cytokine-dependent, MHC-II independent,

mechanisms by which B cells contribute to cardiac remodeling after MI [12, 38].

B cells typically present antigens via MHC II to CD4 T cells and, under specific conditions, CD8

T cells [39, 40], leading to T cell activation. Prior studies have implicated both CD4 and CD8 T

cells in adverse cardiac remodeling following myocardial injury [41-44]. In the present study, we

found that the adoptive transfer of splenic B cells from ischemic HF modulated gene expression

of both myocardial CD4 T cells and myocardial CD8 T cells in an MHC II dependent manner. In

the CD8 compartment, we observed MHC II dependent activation of IFNγ signaling, a key

mediator of immune mediated myocardial damage [45-47]. The downregulation of the mediator

STAT5B in the absence of MHC II further supports this dependency as altered STAT5B

expression and activation has been implicated in various cardiac pathologies [48, 49]. These

findings therefore suggest that adoptive transfer of post-MI splenic B cells triggers activation of

myocardial T cells, ultimately contributing to adverse cardiac remodeling. Notably, we observed

a stronger signal for MHC II dependent CD8 T cells activation, and minor changes in CD4 T cells.

It remains unclear whether this finding reflects a cascade of activation initiated by CD4 T cells

(with amplification of the activation along the signaling cascade) or if B cells independently

engage both CD4 and CD8 T cells, with a stronger effect on CD8 T cells. Further investigations

will be needed to clarify this interaction.

Our scRNAseq analyses of peripheral blood B cells from patients with ischemic myocardial injury

showed differential regulation of antigen processing and presentation in both the acute and chronic

post MI phases. Several genes involved in this pathway showed shared differential expression

between mice and humans, including multiple MHC complexes, CD74 (which stabilizes MHC II

molecules) [50], and HSPs, which are critical modulators of effector response post antigen

presentation [51, 52].This is noteworthy, as it points to a translational relevance of our findings.

However further work will be needed to assess the direct involvement of these conserved genes

and the extent to which B cell-mediated antigen presentation modulates adverse cardiac

remodeling in patients who experience an MI.

In summary, our findings reveal that MI triggers long-lasting changes in splenic B cells; that

splenic B cells present myocardial antigens via MHC II following myocardial infarction; and that

B cells circulating between the spleen and the heart drive post-MI adverse cardiac remodeling

through MHC II-mediated mechanisms. These observations not only expand our current

understanding of B cell biology but also reshape existing models of immune-cardiac interactions

post-myocardial injury [2, 53-55]. Moreover, they highlight the potential of targeting MHC II-

dependent B cell functions as a novel therapeutic approach to prevent and treat ischemic heart

failure. Further studies will be needed to understand whether our findings are relevant to other

conditions characterized by acute or chronic cardiac damage.

Methods:

All studies were approved by the Institutional Animal Care and Use Committee (ACUC) of the Johns Hopkins University School of Medicine, under protocol numbers MO20M388 and M023M238.

Mouse Models and Surgical Protocol:

Male 10-14- week-old mice were used for all experiments unless otherwise specified. All mice were purchased from Jackson Laboratory (Bar Harbor, ME) or bred in the Johns Hopkins vivarium facility. The following strains were used: C57BL/6J strain N. 000664, B6.SJL-*Ptprc^a Pepc^b*/BoyJ (CD45.1) strain N. 002014, B6.129P2(C)-*Cd19$^{tm1(cre)Cgn}$*/J strain N. 006785, and B6.129X1-*H2-Ab1$^{b-tm1Koni}$*/J strain N. 013181. To generate mice with B cell-specific MHC class II deletion, B6.129P2(C)-*Cd19$^{tm1(cre)Cgn}$*/J and B6.129X1-*H2-Ab1$^{b-tm1Koni}$*/J mice were bred for target *Cd19$^{tm1(cre)Cgn/-}$H2-Ab1$^{b-tm1Koni/b-tm1Koni}$* mice. Mice were randomly grouped into sham donors or HF donors. HF donors underwent left thoracotomy with permanent ligation of the left anterior descending (LAD) coronary artery to induce an MI. For the desired condition of ischemic HF, mice were utilized at four weeks post- MI surgery [56, 57]. Sham donors underwent sham intervention. Experiments referring to wildtype mice include a mix of heterozygous *Cd19$^{tm1(cre)Cgn/-}$H2-Ab$^{-/-}$*, homozygous *Cd19$^{tm1(cre)Cgn/-}$* littermate controls and syngeneic C57BL/6J strain N. 000664.

Adoptive Transfer Studies:

Mice were euthanized four weeks following MI or sham surgery. Animals were gently perfused through heart apex with 3mL Hanks' Balanced salt solution (HBSS) without $CaCl_2$, $MgCl_2$ or $MgSO_4$ (GIBCO). Spleens were harvested and placed into a 35mm tissue culture dish with phosphate-buffered saline (PBS). Splenocytes were isolated as previously published by Ismahil et al [8]. Briefly, the spleen was minced finely using scissors and the end of a syringe plunger. The solution was then pipetted through a 40μm cell strainer into a sterile 50mL tube. The tube was centrifuged at 250g for 5 minutes at 4°C. After decanting the supernatant, the pellet was resuspended with 5mL of ACK lysis buffer (Quality Biological) for 5 minutes at room temperature to remove red blood cells (RBCs). After 5 minutes of lysis, 30mL of PBS was added to each tube and each tube was centrifuged at 250g for 5 minutes at 4°C. The supernatant was decanted, and pelleted splenocytes were then resuspended in sterile 0.9% sodium chloride for a final concentration of ~$1x10^8$ total splenocytes in 150μL per retro-orbital injection into naïve recipient mice.

For adoptive transfer studies with purified B cells, splenic B cells were isolated according to the MojoSort™ Mouse Pan B Cell Isolation Kit protocol (BioLegend), with slight modifications. Briefly, after mincing and filtering each spleen as described above, the pellet of each spleen was resuspended with 4mL of Mojo buffer and then filtered again into fluorescence-activated cell sorting (FACS) tubes. Each tube containing cells from one spleen was then centrifuged at 250g for 5 minutes at 4°C. After decanting the supernatant, the pellet was resuspended with cold 2mL Mojo buffer. Samples were then incubated in 200μL of primary antibody cocktail for 15 minutes on ice, followed by another 15 minutes incubation with 200μL of Nanobeads. Incubation was stopped with 1.5mL Mojo buffer, and magnet sorted for 5 minutes. The cell suspension was decanted into

a fresh FACS tube, magnet incubation was repeated to increase the yield of isolated B cells. The
suspension of each spleen was then centrifuged, and each pellet was resuspended in FACS buffer.
The splenic B cell solutions from mice of the same experimental group were combined into a
single tube and then centrifuged prior to final resuspension in sterile 0.9% sodium chloride for
retro-orbital injection of ~1×10^7 splenic B cells in 150µL into naïve recipient mice. A small sample
was obtained pre- and post-antibody incubation to evaluate purity and yield by flow cytometry
(Supplementary Fig.1). Purified splenic B cells consistently had >95% purity.

For adoptive transfer studies with splenic B cell subtypes, single cell suspensions were prepared
as above and then stained with CD19-BV421, CD21-APC, and CD23-PE antibodies and
propidium iodide. Using MoFlo sorter, marginal zone B (MZB) cells and follicular B cells were
sorted as $CD19^+/CD21^+/CD23^{low}$ and $CD19^+/CD21^{low}/CD23^{high}$, respectively. Following
collection, samples from different mice were pooled, and all sorted and approximately 1.5×10^6
MZB cells and 1×10^7 follicular B cells were injected into naïve recipient mice, maintaining the
proportion of these two cell types in unfractionated splenocytes. These numbers of splenic B cell
subtypes were representative of the proportion relative to the total number of unsorted B cells from
a mouse spleen.

Echocardiography:

Mouse transthoracic echocardiography was performed in conscious mice prior to retro-orbital
injection to assess baseline cardiac structure and function, and again 4-, 8- and 16- weeks post
adoptive transfer. Echocardiography was performed using a VisualSonics Vevo 2100 Imaging
System. M-mode parasternal short images were obtained. LVEF and LVESD were measured using
Vevo Lab software v5.8.2.

Gravimetric and Histological Analyses:

Eight weeks after the adoptive transfer, recipient mice were euthanized, and hearts and spleens
were dissected and weighed. Tibia length and body weight were recorded to normalize the weights
of each mouse heart and spleen. Heart tissue was fixed with 10% neutral buffered formalin, -
embedded in paraffin, and stained with WGA using standard protocols.

Isolation of Myocardial Mononuclear Cells:

To analyze myocardial immune cells via flow cytometry, the heart was digested as previously
described with slight modifications [58]. Each heart was perfused with 3mL of Hank's Buffered
Salt Solution (HBSS) with calcium and magnesium infused into the left ventricle over 1 minute
with aortic clamping. Hearts were then dissected and cut transversely, approximately at the mid-
papillary muscle level, or approximately ~5mm below the atrioventricular groove. The superior
section was used for flow cytometry, and the apical section was used for histological analysis. The
heart tissue for flow cytometry was finely minced and 60mg was suspended in 3mL HBSS. Heart
tissue was digested with 300U DNAse, 625U collagenase II, and 50U hyaluronidase shaken for 30
minutes at 300rpm and 37°C. At the end of the 30 minutes, additional HBSS was added to each
tube to neutralize the enzymatic digestion. Each tube was then centrifuged at 250g for 5 min at
4°C. The supernatant was decanted, and the pellet was resuspended in 5mL of ACK lysis buffer
for 5 minutes at room temperature. The suspension was then diluted with PBS and filtered through

a 40μm strainer. Each cell suspension was centrifuged at 250g for 5 min prior to resuspension in
FACS buffer and filtering through a 35μm mesh strainer into a sterile FACS tube.
523

524   <u>Isolation of Splenocytes for Flow Cytometry:</u>
525   The spleens were removed and prepared as described above for adoptive transfer of splenocytes.
526   Following completion of RBC lysis, cells were resuspended in FACS buffer and filtered through
527   35μm cell strainers into sterile FACS tubes.
528

529   <u>Isolation of Peripheral Blood Cells for Flow Cytometry:</u>
530   Peripheral blood (~100μL) was collected via venipuncture of the facial vein into 1.5mL Eppendorf
531   tubes containing 100μL heparin. RBCs were lysed with 1mL ACK buffer for 5 minutes at room
532   temperature and then centrifuged at 250g for 5 minutes at 4°C. Cells were resuspended in 500μL
533   FACS buffer then filtered through 35μm cell strainers into FACS tubes.
534

<u>Flow Cytometry:</u>
Isolated cell suspensions were stained for 30 minutes on ice with the following fluorescently
conjugated antibodies: CD3-APC/Fire810 (clone 17A2, BioLegend), PD-1-BUV615 (clone
29F.1A12, BD Horizon), CD44-BV737 (clone IM7, Invitrogen), CXCR3-BV421 (clone CXCR3-
173, BioLegend), CD62L-BV510 (clone MEL-14, BioLegend), CD8 (clone 53-6.7, BioLegend),
CD69-BV650 (clone H1.2F3, BioLegend), CD27-BV785 (clone LG.3A10, BioLegend), KLRG-
1-PerCP/Cy5.5 (clone 2F1/KLRG1, BioLegend), CD103-PerCP/eFluor710 (clone 2E7,
Invitrogen), CXCR6-PE (cloneSA051D1, BioLegend), CD127-PE/Dazzle594 (clone A7R34,
BioLegend), ICOS-PE/Cy5 (clone 15F9, BioLegend), Ki-67-AF488 (clone SolA15, Invitrogen),
Granzyme B-Pacific Blue (clone GB11, BioLegend) CD25-BUV395 (clone PC61, BioLegend),
CD4-BV650 (clone GK1.5, BioLegend), CCR6-BV785 (clone29-2L7, BioLegend), CXCR5-
PerCP/eFluor710 (clone SPRCL5, Invitrogen), FOXP3-AF488 (clone MF-14, BioLegend),
CTLA-4-PE (clone UC10-4B9, BioLegend), T-bet-PE/Dazzle594 (clone 4B10, BioLegend),
RORrt-PE/Cy7 (clone Q31-378, BD Pharmingen) CD45-BUV395 (clone 30-F11, BD Horizon),
CD73-BV421 (clone TY/11.8, BioLegend), GL7-Pacific Blue (clone GL7, BioLegend), IgM-
BV605 (clone RMM-1, BioLegend), CD80-BV650 (clone 16-10A1, BioLegend), CD21/CD35-
BV711 (clone 7E9, BioLegend), CD19-BV750 (clone 6D5, BioLegend), I-ab-AF488 (clone KH-
74, BioLegend), CD5-PerCP (clone 53-7.3, BioLegend), CD1d-PerCP/Cy5.5 (clone L363,
BioLegend), CD23-PE (clone B3B4, BioLegend), CD86-PE/Dazzle594 (clone GL-1, BioLegend),
CD38-PE/Fire640 (clone 90, BioLegend), CD11b-PE/Cy5 (clone M1/70, BioLegend), PD-L2-
PE/Cy7 (clone TY25, BioLegend), CD138-PE/Fire810 (clone 281-2, BioLegend), CD93-APC
(clone AA4.1, BioLegend), IgD-APC/Fire 750 (clone 11-26c.2a, BioLegend) CD3-FITC (clone
17A2, Invitrogen), CD3-BV605 (17A2, BioLegend), CD19-BV421 (6D5, BioLegend), CD4-
Spark NIR 685 (GK1.5, BioLegend), CD8-BV785 (53-6.7, BioLegend), CD45-PerCP/Cyanine5.5
(30-F11, BioLegend), CCR2-APC (QA18A56, BioLegend), CD64-PE/Cy7 (X54-5/7.1,
BioLegend), Ly6C-BV650 (HK1.4, BioLegend), MHCII-BV711 (M5/114.15.2, BioLegend),
CD11b-Alexa Fluor 700 (M1/70, BioLegend), Ly6G-APC/Cy7 (1A8, BioLegend), CD45.1-PE
(A20, BioLegend). All antibodies were added at 1:1000 dilution for heart and spleen flow
cytometry and 1:500 for peripheral blood flow cytometry, with the following exceptions. PD-1

antibody was added at 1:1600 dilution. Ki-67, CXCR6, CD21/CD35, CD19, I-ab, and CD38
antibodies were added at 1:800 dilution. CD44, CD27, CD25, CTLA-4, and T-bet, PD-L2
antibodies were added at 1:100 dilution. Dead cells were excluded using the Zombie NIR dye
(BioLegend) at 1:5000 dilution. Flow cytometry was performed on Cytek Aurora Full Spectrum
Flow Cytometric Analyzer. Compensation controls were generated using single-color control
samples and UltraComp eBeads (Invitrogen). Representative gating strategies are shown in
individual figures.
571

Single Cell RNA (scRNA) Sequencing:
C57BL/6J mice 14-18 weeks-old underwent left thoracotomy with either sham surgery (n = 5) or
permanent coronary LAD ligation (n = 11). Blood, heart, and spleen were collected 4 weeks post-
sham or MI surgery ("heart failure"; n = 6) or 4 days post-MI surgery ("recent MI"; n = 5). For
myocardial immune cell scRNA seq, recipient hearts (n=6/group) were collected 8 weeks after
retro-orbital injections of splenic B cells from sham, HF or MHC II KO HF mice; half the heart,
cut along the longitudinal axis were further processed for scRNAseq, remnant stored at -80°C.
Single-cell suspensions for blood, heart, and spleen were prepared as described above. Heart and
spleen samples were stained with 2 drops of Vibrant Dye Cycle Violet Ready Flow Reagent (DCV-
Invitrogen) for 30 minutes, shaking at 300rpm and 37°C. Samples were stained with propidium
iodide (PI), CD45-PE antibodies for all experiments and CD19-APC antibody for specific B cell
subtypes, for 30 minutes on ice. Hashtag antibodies mapped to each experimental group were also
added to cell suspensions at a concentration of 2μL per tube. B cells or CD45$^+$ cells were sorted
using a MoFlo sorter. Equal numbers of the three experimental groups were mixed to reach a total
of 20,000-30,000 cells that were combined in equal proportions within the same single cell library.
Three libraries were prepared for the post-MI surgery experiment: 1) peripheral blood B cells; 2)
splenic B cells; and 3) myocardial B cells. One library was prepared for the myocardial immune
cell experiment.
590

591   Demultiplexing, barcode processing, single cell 5' gene counting, and V(D)J transcript sequence
592   assembly and annotation were performed using Cell Ranger (10x Genomics). Gene specific
593   analysis was performed using *Partek™ Flow™* software (v.11.0) for the post-MI surgery
594   experiment and R (v.4.0.2) for myocardial immune cells experiment.
595

Uniform manifold approximation and projection (UMAP) plots were created on R (v.4.0.2) using
Seurat (v.4.1.1) [59] standard parameters unless otherwise noted. Cells were filtered based on their
distributions to only include those with 200-3200 unique features for blood, 200-3800 for heart
(post-MI experiment), 200-4800 for heart (myocardial immune experiment), and 200-3500 for
spleen, and < 8% mitochondrial read counts. Doublets not identified by hashtag demultiplex were
removed using scDblFinder (v. 1.24.0). Counts were then logNormalized and scaled, and the top
2000 (post-MI experiment) and top 3000 (myocardial immune experiment) most variable genes
were identified using the FindVariableFeatures function. Principal component (PC) analysis was
performed using RunPCA and the first 15 PCs (post-MI) and 30 PCs (myocardial immune) were
used to find the shared nearest neighbors (SNN) using the FindNeighbours function. Clustering
was performed using the FindClusters function with a resolution of 0.2 and visualized using a
UMAP projection. B cell and myocardial immune cell subtypes were classified using ScType [18]
by using a modified list of known murine B cell and myocardial immune markers (Supplementary
Table 6) and mapping mouse gene symbols to their human counterpart using biomart to be

compatible with ScType. Analyses of differential expressed genes between conditions per each
cell type were performed using DEseq2 (1.46.0), with false-discovery rate (FDR) step-up used for
multiple test corrections. Genes with low expression (<10 cells with 1 count) were filtered out
prior to model fitting. For post-MI experiments lists of genes that exhibited differential expression
based on the threshold of raw p-values<0.05 and $\log_2$fold change >1 for upregulation of <-1 for
downregulation were compiled and then analyzed against the Kyoto Encyclopedia of Genes and
Genomes (KEGG) database on the Gene Set Enrichment Analysis (GSEA) website
(https://www.gsea-msigdb.org; [19, 20]). Pathways were filtered using KEGG pathway
annotations for immune-system specificity and were considered as differentially regulated if q-
value from FDR was $\geq$ 0.05. For the myocardial immune experiment DEGs were filtered using
threshold of adjusted p<0.05 and $\log_2$fold change of $\geq$0.02 for upregulation and $\leq$-0.02 for
downregulation. The functional analyses and networks were generated using QIAGEN IPA and
IPA Interpret (QIAGEN Inc., https://digitalinsights.qiagen.com/IPA and
https://digitalinsights.qiagen.com/products-overview/discovery-insights-portfolio/analysis-and-
visualization/qiagen-ipa/qiagen-ipa-interpret/) [60].
625
The Chromium Next GEM Single Cell 5' Kit v2 (10x Genomics) was used to perform single cell
transcriptional analyses. The Chromium Next GEM Single Cell V(D)J Kit was used to assess B
cell receptor (BCR) cDNA and clonality. The Chromium Next GEM Single Cell V(D)J Kit was
used to obtain BCR cDNA. Full-length and productive contigs were analyzed using scRepertoire
(v.2.2.1). Only B cells with fully annotated heavy and light chains were kept for clonotype analysis.
CDR3 nucleotide sequences plus V gene were used to define clonotypes. BCRs with >85%
normalized Levenshtein distance of the nucleotide sequence were considered clones. All codes
used for analyses is publicly available: https://github.com/caro471/Adamo_collaboration_2026.
634

<u>Human scRNA Sequencing Analysis:</u>
Publicly available scRNA sequencing data of peripheral blood from two cohorts were analyzed:
three patients with ischemic cardiomyopathy (ICM) EGA accession: EGAS00001006330 [28] and
38 patients 24 hours and 8 weeks following ST segment elevation MI (STEMI) EGA accession:
EGAD00001010064 [27] relative to healthy controls. These data were processed on R (v.4.0.2)
using Seurat (v.4.1.1) as described above with the following modifications: Cells were filtered
based on their distributions to only include those with 200-8000 unique features and <5%
mitochondrial reads for the ICM dataset, 200 – 4500 unique features and <12% mitochondrial
reads for the STEMI v2 dataset, 200 – 3500 unique features and <8% mitochondrial reads for the
STEMI v3 dataset. 3000 variable features were used for both human datasets. Both datasets had
notable batch effects that were corrected using Harmony [61]. For both datasets, the first 30 PCs
were used for the SNN and clustering. A clustering resolution of 0.5 and 0.4 was used for the ICM
and STEMI datasets respectively. B cells were identified using known human B cell-specific
makers as previously described (Supplementary Table 6) [62]. For the larger dataset, B cells were
pseudobulked per patient and differential gene expression was performed using DEseq2
controlling for sex and 10x chemistry in the regression model. Subsequent KEGG pathway
analyses were performed as described above. For the smaller dataset, the defined threshold used
was p-value <0.05, FDR step up <0.05and $\log_2$fold change threshold of $\leq$-0.58 or $\geq$0.58 with
p<0.05. For the larger dataset, a more stringent $\log_2$fold change threshold of $\leq$-1 or $\geq$1 was used
with p-value<0.05.

Peptidome Analysis:

Wildtype (WT) C57BL/6J and $Cd19^{tm1(cre)Cgn/-}H2-Ab1^{b\text{-tm1Koni}/\,b\text{-tm1Koni}}$ (MHC II-KO) mice (n = 16 total) underwent left thoracotomy with either sham surgery (n = 8) or permanent coronary ligation (n = 8). MHC II KO mice were included to serve as negative controls. Four weeks following surgery, spleens were collected from mice and splenic B cells were isolated, snap frozen in liquid nitrogen, and stored at -80°C. Four spleens from the same experimental condition were combined to get $2x10^6$ million B cells per experimental condition. Frozen splenic B cells were lysed and purified for MHC-II bound peptides as previously described [63]. Briefly cells were lysed in 50mM Tris- 150mM NaCl buffer, supplemented with protease inhibitor cocktail (Roche) and 0.5% NP-40 using a bead homogenizer. Lysates were cleared by centrifugation at 2000g and 4°C for 10 minutes, the supernatant was collected and cleared again by ultra-centrifugation at 100,000g and 4°C for 75 minutes. The supernatant was further cleared by flowing through Protein A/G Sepharose column and then incubated for 2 hours at 4°C in Protein A/G Sepharose column cross-linked with MHC-II mAb (M5/114.15.2, BioLegend) with slow agitation (60-80 rpm). Afterwards flowthrough was decanted, and the peptide-bound column was washed four times: first with a buffer containing, 0.05% NP-40, 50mM Tris, 150mM NaCl, 5mM EDTA, 0.1mM PMSF and 1mM Pepstatin A; second with 50mM Tris and 150mM NaCl; third 50mM Tris and 450mM NaCl and finally 50mM Tris. Immuno-bound MHC molecules were then eluted with 10% acetic acid, lyophilized and shipped to Dr Sidoli's lab and analyzed on an Orbitrap Fusion Lumos (Thermo Scientific) mass spectrometer (details below).

Peptides abundances from WT groups were corrected using peptide abundances from MHC II-KO mice, to eliminate background noise. The average abundance of corrected values was calculated across technical replicates for each peptide. Peptides with mean corrected values above zero were included in further analyses. The fold enrichment for each peptide was determined by dividing values of HF group by sham. The binding affinity (BA) of these peptides were then determined using the NetMHCIIpan-4.3 machine learning tool [64, 65]. Peptides with a fold enrichment of greater than 1.2 and amino acid (AA) sequence length of 9 AAs were submitted into the BA server. Briefly the server is trained on an extensive dataset of naturally occurring peptides covering 650,000 measurements of Binding Affinity and Eluted Ligand Mass spectrometry. The server ranks each peptide according to predicted binding affinity in nanomolar IC50 compared against a set of 100,000 random natural peptides [64, 65].

Cardiac proteomic sample preparation and analyses:

Snap frozen left ventricular tissue collected from recipients of adoptive transfer studies were processed for mass spectrometry analyses using PreOmics® iST-NHS kit according to manufacturer's instructions. Briefly, tissue was homogenized in LYSE-NHS buffer, sonicated and boiled at 95°C for 10 minutes. Homogenates were cleared by centrifugation at 14,000g for 14 minutes, supernatant was collected, and samples were incubated in DIGEST buffer at 37°C for 1-2 hours. Following digest, samples were labelled using TMTpro™ 16plex Label kit (Thermo Fisher), washed, eluted, lyophilized and processed on an Orbitrap Fusion Lumos (Thermo Scientific) mass spectrometer (details below).

Spectral TMT reporter ion intensities were quantified using the median sweep normalization strategy originally described by Herbrich et al [66], following the implementation reported by Foster et al [67], with minor modifications. Briefly, TMT reporter ion intensities were (1) $\log_2$-transformed, (2) median-centered within each channel, followed by (3) median-centering of each individual spectrum across channels, and (4) protein abundance was determined by median summarization, defined as the median of the $\log_2$-transformed, median-centered intensities from all spectra assigned to that protein within a given channel. Differential protein abundance between experimental groups was evaluated using a Bayesian statistical framework implemented in linear models for microarray data (LIMMA) with multigroup comparisons [68]. Pairwise contrasts were also performed. Moderated p-values were adjusted for multiple testing using the positive false discovery rate (q-value) approach described by Storey & Tibshirani [69]. Heat maps were generated using the ComplexHeatmap R package with hierarchical clustering (Euclidean distance, complete linkage). Principal component analysis (PCA) was performed using the PCAtools R package. All codes used for statistical analysis, including PCA and heat map generation, is publicly available at: https://github.com/Frostman300/MHCII_upload.

Mass Spectrometry:

Sample Desalting

Prior to mass spectrometry analysis of both TMT-labeled samples and peptides, samples were desalted using a 96-well plate filter (Orochem) packed with 1 mg of Oasis HLB C-18 resin (Waters). Briefly, the samples were resuspended in 100 µl of 0.1% TFA and loaded onto the HLB resin, which was previously equilibrated using 100 µl of the same buffer. After washing with 100 µl of 0.1% TFA, the samples were eluted with a buffer containing 70 µl of 60% acetonitrile and 0.1% TFA and then dried in a vacuum centrifuge.

LC-MS/MS Acquisition and Analysis

Samples were resuspended in 10 µl of 0.1% TFA and loaded onto a Dionex RSLC Ultimate 300 (Thermo Scientific), coupled online with an Orbitrap Fusion Lumos (Thermo Scientific). Chromatographic separation was performed with a two-column system, consisting of a C-18 trap cartridge (300 µm ID, 5 mm length) and a picofrit analytical column (75 µm ID, 25 cm length) packed in-house with reversed-phase Repro-Sil Pur C18-AQ 3 µm resin. Peptides were separated using a 90 min gradient from 4-30% buffer B (buffer A: 0.1% formic acid, buffer B: 80% acetonitrile + 0.1% formic acid) at a flow rate of 300 nl/min. The mass spectrometer was set to acquire spectra in a data-dependent acquisition (DDA) mode from TMT-labeled samples and data-independent mode (DIA) for peptidome samples. Briefly, the full MS scan was set to 300-1200 m/z in the orbitrap with a resolution of 120,000 (at 200 m/z) and an AGC target of $5\times10e5$. MS/MS was performed in the Orbitrap using the top speed mode (2 secs), an AGC target of $2\times10e5$ and an HCD collision energy of 45. Search tolerance for MS1 was 10ppm and 20ppm for MS2, fixed modifications N-terminus TMT, lysine TMT and cysteine carbamidomethyl and methionine oxidation set for variable modifications.

Proteome raw files were searched using Proteome Discoverer software (v2.5, Thermo Scientific) using SEQUEST search engine and the SwissProt mouse database (updated April 2023). Any peptide cleavage was specified for digestion, so that the software considered all protein fragments. FDR determined by percolator validator. The mass spectrometry proteomics data have been

743 deposited to the ProteomeXchange Consortium via the PRIDE [70] partner repository with the
744 dataset identifier PXD073894
745

<u>Immunofluorescence staining</u>:

Endpoint serum from recipients of adoptive transfer studies were evaluated for cardiac
autoantibodies as previously described [71]. Briefly paraffin embedded cardiac cross-sections
from congenital B cell-deficient μMT mice were rehydrated and prepared using Tris-EDTA
antigen retrieval and incubated with recipient-serum (1:20 dilution) overnight at 4°C. Sections
were then washed and autoantibodies against cardiac antigens were then probed for using Biotin-
goat anti mouse IgG (clone Poly4053, BioLegend) and rat anti-mouse IgM (clone RMM-1,
BioLegend) at 1:1000 dilution. Following washes, the antibodies were detected using streptavidin-
Alexa555 (Invitrogen) and anti-rat-Alexa647 (Life Technologies); cardiomyocyte basement
membrane was stained with WGA-Alexa488 and nuclei stained with DAPI. Fluorescent images
were acquired using the PhenoImager™ HT 3.0, Vectra Polaris (Akoya Biosciences) slide scanner
at 40x objective, with the same exposure time. Images were processed using QuPath and analyzed
with ImageJ, three ROIs were taken for each section and two sections per sample. Aged mouse
serum was used for positive control and negative control were section without serum.
760

<u>Statistical Analysis</u>

Data was initially processed on Microsoft Excel (Microsoft 365). After assessing normality with
the Shapiro-Wilk test, Mann-Whitney or unpaired two-tailed t-tests were used for non-parametric
and parametric comparisons, respectively, of two groups. To compare data across three or more
groups, one-way analysis of variance (ANOVA) or Kruskal- Wallis tests were used for normally
distributed or non-Gaussian distributed data, respectively, followed by pairwise comparisons.
Serial echocardiographic data were analyzed using two-way ANOVA or mixed-effects analysis
followed by pairwise comparisons. WGA data were analyzed using nested t-test for two groups or
nested one-way ANOVA followed by pairwise multiple comparisons for three groups. All pairwise
multiple comparison tests were corrected using the original FDR method of Benjamini and
Hochberg. Outliers were determined using the ROUT method ($Q=1\%$). All statistical tests were
performed on GraphPad Prism version 10.6.1 for Windows, GraphPad Software, Boston,
Massachusetts USA, www.graphpad.com.
774

Acknowledgments

The authors acknowledge the support for the Ross Flow Cytometry core funded by NIH Grant S10OD026859. The authors would also like to thank Dr. Edward Pearce for critical discussions and feedback on the manuscript and Dr. Hao Zhang from the Bloomberg FACS sorting Facility for technical support with FACS sorting. The Sidoli lab gratefully acknowledges for funding the Hevolution Foundation (AFAR), the Einstein-Rockefeller-CUNY-Mount Sinai Center for AIDS Research, the NIH Office of the Director (S10OD030286), and the Aging Biology Foundation.

Author contributions

OVE and JPL designed experiments, performed experiments, analyzed data, and wrote the manuscript. CD assisted with the analysis of murine and human scRNA data. SR, AB, YS, SS and KK assisted with collection and analysis of experimental data. KB, CC, SS, DBF, MM and SK helped with experimental design, review of data and manuscript edits. LA conceived the project, supervised experimental design and analyses, and wrote the manuscript with OVE and JPL.

References

1. Hofmann, U. and S. Frantz, *Immunity strikes: heart failure as a systemic disease.* Eur Heart J, 2014. **35**(6): p. 341-3.
2. Adamo, L., et al., *Reappraising the role of inflammation in heart failure.* Nat Rev Cardiol, 2020. **17**(5): p. 269-285.
3. Nahrendorf, M., et al., *The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions.* J Exp Med, 2007. **204**(12): p. 3037-47.
4. Libby, P., M. Nahrendorf, and F.K. Swirski, *Leukocytes Link Local and Systemic Inflammation in Ischemic Cardiovascular Disease: An Expanded "Cardiovascular Continuum".* J Am Coll Cardiol, 2016. **67**(9): p. 1091-1103.
5. Murphy, S.P., et al., *Inflammation in Heart Failure: JACC State-of-the-Art Review.* J Am Coll Cardiol, 2020. **75**(11): p. 1324-1340.
6. Reina-Couto, M., et al., *Inflammation in Human Heart Failure: Major Mediators and Therapeutic Targets.* Front Physiol, 2021. **12**: p. 746494.
7. Swirski, F.K., et al., *Identification of splenic reservoir monocytes and their deployment to inflammatory sites.* Science, 2009. **325**(5940): p. 612-6.
8. Ismahil, M.A., et al., *Remodeling of the mononuclear phagocyte network underlies chronic inflammation and disease progression in heart failure: critical importance of the cardiosplenic axis.* Circ Res, 2014. **114**(2): p. 266-82.
9. Rocha-Resende, C., et al., *Developmental changes in myocardial B cells mirror changes in B cells associated with different organs.* JCI Insight, 2020. **5**(16).
10. Adamo, L., et al., *Myocardial B cells are a subset of circulating lymphocytes with delayed transit through the heart.* JCI Insight, 2020. **5**(3).
11. Bermea, K.C., et al., *The human myocardium harbors a population of naive B-cells with a distinctive gene expression signature conserved across species.* Front Immunol, 2022. **13**: p. 973211.
12. Zouggari, Y., et al., *B lymphocytes trigger monocyte mobilization and impair heart function after acute myocardial infarction.* Nat Med, 2013. **19**(10): p. 1273-80.
13. Adamo, L., et al., *Modulation of subsets of cardiac B lymphocytes improves cardiac function after acute injury.* JCI Insight, 2018. **3**(11).
14. Wu, L., et al., *IL-10-producing B cells are enriched in murine pericardial adipose tissues and ameliorate the outcome of acute myocardial infarction.* Proc Natl Acad Sci U S A, 2019. **116**(43): p. 21673-21684.
15. Heinrichs, M., et al., *The healing myocardium mobilizes a distinct B-cell subset through a CXCL13-CXCR5-dependent mechanism.* Cardiovasc Res, 2021. **117**(13): p. 2664-2676.
16. Schloss, M.J., et al., *B lymphocyte-derived acetylcholine limits steady-state and emergency hematopoiesis.* Nat Immunol, 2022. **23**(4): p. 605-618.
17. Adamo, L., C. Rocha-Resende, and D.L. Mann, *The Emerging Role of B Lymphocytes in Cardiovascular Disease.* Annu Rev Immunol, 2020. **38**: p. 99-121.
18. Ianevski, A., A.K. Giri, and T. Aittokallio, *Fully-automated and ultra-fast cell-type identification using specific marker combinations from single-cell transcriptomic data.* Nat Commun, 2022. **13**(1): p. 1246.

19.   Mootha, V.K., et al., *PGC-1alpha-responsive genes involved in oxidative*
phosphorylation are coordinately downregulated in human diabetes. Nat Genet, 2003.
34(3): p. 267-73.
20.   Subramanian, A., et al., *Gene set enrichment analysis: a knowledge-based approach for*
interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A, 2005. **102**(43):
p. 15545-50.
21.   Deffit, S.N. and J.S. Blum, *A central role for HSC70 in regulating antigen trafficking and*
MHC class II presentation. Mol Immunol, 2015. **68**(2 Pt A): p. 85-8.
22.   Houlihan, J.L., J.J. Metzler, and J.S. Blum, *HSP90alpha and HSP90beta isoforms*
selectively modulate MHC class II antigen presentation in B cells. J Immunol, 2009.
182(12): p. 7451-8.
23.   Phan, U.T., B. Arunachalam, and P. Cresswell, *Gamma-interferon-inducible lysosomal*
thiol reductase (GILT). Maturation, activity, and mechanism of action. J Biol Chem,
2000. **275**(34): p. 25907-14.
24.   Sun, Y., et al., *Splenic Marginal Zone B Lymphocytes Regulate Cardiac Remodeling*
After Acute Myocardial Infarction in Mice. J Am Coll Cardiol, 2022. **79**(7): p. 632-647.
25.   Parker Harp, C.R., et al., *B cell antigen presentation is sufficient to drive*
neuroinflammation in an animal model of multiple sclerosis. J Immunol, 2015. **194**(11):
p. 5077-84.
26.   Collins, C.B., et al., *Heat shock factor 1 drives regulatory T-cell induction to limit murine*
intestinal inflammation. Mucosal Immunol, 2024. **17**(1): p. 94-110.
27.   van Blokland, I.V., et al., *Single-Cell Dissection of the Immune Response After Acute*
Myocardial Infarction. Circ Genom Precis Med, 2024. **17**(3): p. e004374.
28.   Kuppe, C., et al., *Spatial multi-omic map of human myocardial infarction.* Nature, 2022.
608(7924): p. 766-777.
29.   Emami, H., et al., *Splenic metabolic activity predicts risk of future cardiovascular events:*
demonstration of a cardiosplenic axis in humans. JACC Cardiovasc Imaging, 2015. **8**(2):
p. 121-30.
30.   van der Laan, A.M., et al., *Monocyte subset accumulation in the human heart following*
acute myocardial infarction and the role of the spleen as monocyte reservoir. Eur Heart J,
2014. **35**(6): p. 376-85.
31.   Kercheva, M., et al., *Characteristics of the Cardiosplenic Axis in Patients with Fatal*
Myocardial Infarction. Life (Basel), 2022. **12**(5).
32.   Ismahil, M.A., et al., *Splenic CD169 (+) Tim4 (+) Marginal Metallophilic Macrophages*
Are Essential for Wound Healing After Myocardial Infarction. medRxiv, 2024.
33.   Lv, H., et al., *Impaired thymic tolerance to α-myosin directs autoimmunity to the heart in*
mice and humans. J Clin Invest, 2011. **121**(4): p. 1561-73.
34.   Joachimbauer, A., et al., *Cardiopathogenic T Cells Govern Progression and Functional*
Remodeling in Inflammatory Cardiomyopathy and Chronic Myocarditis. JACC Basic
Transl Sci, 2026. **11**(2): p. 101458.
35.   Zhao, T.X., et al., *Rituximab in patients with acute ST-elevation myocardial infarction:*
an experimental medicine safety study. Cardiovasc Res, 2022. **118**(3): p. 872-882.
36.   Rastogi, I., et al., *Role of B cells as antigen presenting cells.* Front Immunol, 2022. **13**: p.
954936.

37. Rock, K.L., B. Benacerraf, and A.K. Abbas, *Antigen presentation by hapten-specific B*
lymphocytes. I. Role of surface immunoglobulin receptors. J Exp Med, 1984. **160**(4): p.
1102-13.
38. Cohen, C.D., et al., *Myocardial Immune Cells: The Basis of Cardiac Immunology.* J
Immunol, 2023. **210**(9): p. 1198-1207.
39. Do, J.S., et al., *Unexpected role for MHC II-peptide complexes in shaping CD8 T-cell*
expansion and differentiation in vivo. Proc Natl Acad Sci U S A, 2012. **109**(31): p.
12698-703.
40. Migueles, S.A. and M. Connors, *Class II-Restricted CD8s: New Lessons Violate Old*
Paradigms. Immunity, 2016. **45**(4): p. 712-714.
41. Laroumanie, F., et al., *CD4+ T cells promote the transition from hypertrophy to heart*
failure during chronic pressure overload. Circulation, 2014. **129**(21): p. 2111-24.
42. Bansal, S.S., et al., *Activated T Lymphocytes are Essential Drivers of Pathological*
Remodeling in Ischemic Heart Failure. Circ Heart Fail, 2017. **10**(3): p. e003688.
43. Gröschel, C., et al., *T helper cells with specificity for an antigen in cardiomyocytes*
promote pressure overload-induced progression from hypertrophy to heart failure. Sci
Rep, 2017. **7**(1): p. 15998.
44. Santos-Zas, I., et al., *Cytotoxic CD8(+) T cells promote granzyme B-dependent adverse*
post-ischemic cardiac remodeling. Nat Commun, 2021. **12**(1): p. 1483.
45. Finger, S., et al., *A sequential interferon gamma directed chemotactic cellular immune*
response determines survival and cardiac function post-myocardial infarction.
Cardiovasc Res, 2019. **115**(13): p. 1907-1917.
46. Cosper, P.F., P.A. Harvey, and L.A. Leinwand, *Interferon-γ causes cardiac myocyte*
atrophy via selective degradation of myosin heavy chain in a model of chronic
myocarditis. Am J Pathol, 2012. **181**(6): p. 2038-46.
47. Levick, S.P. and P.H. Goldspink, *Could interferon-gamma be a therapeutic target for*
treating heart failure? Heart Fail Rev, 2014. **19**(2): p. 227-36.
48. Vaparanta, K., et al., *STAT5b is a key effector of NRG-1/ERBB4-mediated myocardial*
growth. EMBO Rep, 2023. **24**(5): p. e56689.
49. Jin, G., L. Wang, and J. Ma, *Inhibiting STAT5 significantly attenuated Ang II-induced*
cardiac dysfunction and inflammation. Eur J Pharmacol, 2022. **915**: p. 174689.
50. Schröder, B., *The multifaceted roles of the invariant chain CD74--More than just a*
chaperone. Biochim Biophys Acta, 2016. **1863**(6 Pt A): p. 1269-81.
51. Millar, D.G., et al., *Hsp70 promotes antigen-presenting cell function and converts T-cell*
tolerance to autoimmunity in vivo. Nat Med, 2003. **9**(12): p. 1469-76.
52. Barbera Betancourt, A., et al., *T Cell-Mediated Chronic Inflammatory Diseases Are*
Candidates for Therapeutic Tolerance Induction with Heat Shock Proteins. Front
Immunol, 2017. **8**: p. 1408.
53. Frangogiannis, N.G., *The inflammatory response in myocardial injury, repair, and*
remodelling. Nat Rev Cardiol, 2014. **11**(5): p. 255-65.
54. Westman, P.C., et al., *Inflammation as a Driver of Adverse Left Ventricular Remodeling*
After Acute Myocardial Infarction. J Am Coll Cardiol, 2016. **67**(17): p. 2050-60.
55. Rurik, J.G., H. Aghajanian, and J.A. Epstein, *Immune Cells and Immunotherapy for*
Cardiac Injury and Repair. Circ Res, 2021. **128**(11): p. 1766-1779.
56. Yang, F., et al., *Myocardial infarction and cardiac remodelling in mice.* Exp Physiol,
2002. **87**(5): p. 547-55.

57. Lindsey, M.L., et al., *Guidelines for in vivo mouse models of myocardial infarction.* Am J
Physiol Heart Circ Physiol, 2021. **321**(6): p. H1056-h1073.
58. Bermea, K.C., S.T. Rousseau, and L. Adamo, *Flow Cytometry-Based Quantification and*
Analysis of Myocardial B-cells. J Vis Exp, 2022(186).
59. Hao, Y., et al., *Integrated analysis of multimodal single-cell data.* Cell, 2021. **184**(13): p.
3573-3587.e29.
60. Krämer, A., et al., *Causal analysis approaches in Ingenuity Pathway Analysis.*
Bioinformatics, 2014. **30**(4): p. 523-30.
61. Korsunsky, I., et al., *Fast, sensitive, and accurate integration of single cell data with*
Harmony. bioRxiv, 2018: p. 461954.
62. Bermea, K.C., et al., *Myocardial B cells have specific gene expression and predicted*
interactions in dilated cardiomyopathy and arrhythmogenic right ventricular
cardiomyopathy. Front Immunol, 2024. **15**: p. 1327372.
63. Purcell, A.W., S.H. Ramarathinam, and N. Ternette, *Mass spectrometry-based*
identification of MHC-bound peptides for immunopeptidomics. Nat Protoc, 2019. **14**(6):
p. 1687-1707.
64. Reynisson, B., et al., *Improved Prediction of MHC II Antigen Presentation through*
Integration and Motif Deconvolution of Mass Spectrometry MHC Eluted Ligand Data. J
Proteome Res, 2020. **19**(6): p. 2304-2315.
65. Nilsson, J.B., et al., *Accurate prediction of HLA class II antigen presentation across all*
loci using tailored data acquisition and refined machine learning. Sci Adv, 2023. **9**(47):
p. eadj6367.
66. Herbrich, S.M., et al., *Statistical inference from multiple iTRAQ experiments without*
using common reference standards. J Proteome Res, 2013. **12**(2): p. 594-604.
67. Foster, D.B., et al., *Integrated Omic Analysis of a Guinea Pig Model of Heart Failure*
and Sudden Cardiac Death. J Proteome Res, 2016. **15**(9): p. 3009-28.
68. Smyth, G.K., *Linear models and empirical bayes methods for assessing differential*
expression in microarray experiments. Stat Appl Genet Mol Biol, 2004. **3**: p. Article3.
69. Storey, J.D. and R. Tibshirani, *Statistical significance for genomewide studies.* Proc Natl
Acad Sci U S A, 2003. **100**(16): p. 9440-5.
70. Perez-Riverol, Y., et al., *The PRIDE database at 20 years: 2025 update.* Nucleic Acids
Res, 2025. **53**(D1): p. D543-d553.
71. Ramos, G.C., et al., *Myocardial aging as a T-cell-mediated phenomenon.* Proc Natl Acad
Sci U S A, 2017. **114**(12): p. E2420-e2429.

Figure Legends

Fig. 1.

964   **Fig.1: Splenic B cells promote adverse cardiac remodeling in ischemic heart failure a**)
965   Schema for adoptive transfer studies. Four weeks following permanent LAD coronary ligation or
966   sham surgery, unfractionated splenocytes or isolated splenic B cells were transferred to naïve
967   recipient mice. Recipient mice underwent serial transthoracic echocardiography over 8 weeks
968   followed by tissue harvest. **b**) On echocardiography, adoptive transfer of splenocytes or isolated
969   splenic B cells from heart failure (HF) mice resulted in significant reduction in left ventricular
970   ejection fraction (LVEF) and increase in left ventricular end-systolic diameter (LVESD) in naïve
971   recipient mice. **c**) Quantitative gravimetric data revealed significantly larger hearts and spleens
972   (normalized to body weight) in naive mice 8 weeks post-adoptive transfer of splenocytes or
973   isolated splenic B cells from HF mice relative to recipients of splenocytes or splenic B cells from
974   sham-operated mice. **d**) Cardiomyocyte area was increased in recipients of splenocytes or isolated
975   splenic B cells from HF mice relative to sham recipients; n = 5-6 mice per group. Mean values ±
976   SD are represented. **e**) Representative images of WGA staining of hearts from recipients of isolated
977   splenic B cells from HF or sham-operated mice. LAD- left anterior descending. HF-heart failure.

978

Fig. 2.

Fig.2: Adoptively transferred splenic B cells from mice with ischemic HF are found in the heart and spleen of recipients, and do not alter the immune composition of the spleen of naïve mice a) Schema for adoptive transfer of isolated splenic CD45.1 B cells 4 weeks following LAD ligation or sham surgery to naïve CD45.2 mice. **b**) Flow cytometry of peripheral blood from naïve recipient mice revealed the presence of donor CD45.1 B cells 1 week and 8 weeks following adoptive transfer. Donor CD45.1 B cells were also present in the spleen and heart of CD45.2 mice 8 weeks after adoptive transfer. **c**) Representative gating strategy for flow cytometry of spleens from recipient mice 8 weeks post-adoptive transfer of splenic B cells. Adoptive transfer of splenic B cells from HF mice had no effect on total number of lymphocytes, B cells, CD4$^+$ and CD8$^+$ T cells relative to recipients of sham splenic B cells. LAD- left anterior descending. HF-heart failure. n = 6 mice per group. Mean values are represented.

Fig. 3.

Fig.3: Ischemic myocardial injury is associated with activation of antigen processing and
presentation pathways in splenic B cells and with enrichment of myocardial peptides within
the MHC-II molecules of these B cells a) Schema for single cell sequencing of splenic B cells
from mice 4 weeks following permanent LAD coronary ligation or sham surgery. **b**) V(D)J
analysis of splenic B cells. Ischemic myocardial injury did not induce splenic B cell clonal
expansion. **c**) UMAP plots of splenic B cells from heart failure or sham surgery mice. Splenic B
cells were classified into B cell sub-types. **d**) KEGG pathway analysis of differentially expressed
genes in splenic B cells (fold change >1, p<0.05) revealed significant differential expression of
antigen processing and presentation in multiple B cell sub-types, post-MI. All pathways with FDR
<0.05 are listed. (n = 4-5 mice per group). **e**) Schema for mass spectrometry of MHC-II bound
peptides from splenic B cells four weeks after permanent LAD coronary ligation or sham surgery.
Splenic B cells from MHC II KO (*Cd19tm1(cre)Cgn/-H2-Ab1b*-tm1Koni/ *b*-tm1Koni) mice were
included besides cells from WT (wildtype) mice to allow identification of background signal. **f**)
Categories of proteins both enriched > 1.2 folds in the MHC-II molecules of splenic B cells isolated
from mice with ischemic HF and *in silico* confirmed binding to the C57Bl/6 specific MHC-II
molecule H2-I-Ab. 18% of these proteins are cardiac-specific. The peptides/corresponding
proteins identified as strong binders are reported in the heat map. The cardiac proteins titin (TTN),
myosin heavy chain 6 (MYH6) and α-actin (ACTC1) were identified among the top predicted

strong binding peptides. MYH-6 is repeated twice because two different peptides from MYH-6
were detected. LAD- left anterior descending. HF-heart failure.

1013

Fig. 4.

1014
Fig.4: Transfer from ischemic HF mice of splenic B cells deficient in MHC class II does not
produce adverse cardiac remodeling in naïve recipient mice at the echocardiographic,
gravimetric or histological level a) Schema for adoptive transfer studies of isolated splenic B
cells from *Cd19tm1(cre)Cgn/-H2-Ab1b*-tm1Koni/ *b*-tm1Koni mice and wildtype mice following
permanent LAD coronary ligation or sham surgery. **b**) Naïve recipients of splenic B cells isolated
from HF mice with B cell-specific MHC class II deletion did not experience adverse cardiac
remodeling on serial echocardiography over an 8-week period post-adoptive transfer. **c**)
Gravimetric data of recipient mice. Recipient mice of B cells deficient in MHCII from HF mice
did not experience significant splenic or cardiac remodeling by gravimetric analysis. **d**)
Representative images of WGA staining of hearts from recipient mice. Transfer of isolated
MHCII-deficient splenic B cells did not induce cardiomyocyte hypertrophy. LAD- left anterior
descending. HF-heart failure. n= 8 mice per group. Mean values ± SD are represented.

1027

1028

Fig.5: Mass spectrometry of left ventricle shows that transfer of splenic B cells from mice
with ischemic HF into naïve recipients results in adverse cardiac remodeling, which is
blunted when splenic B cells are deficient in MHC class II a) Schema for adoptive transfer
studies of isolated splenic B cells from *Cd19tm1(cre)Cgn/-H2-Ab1b*-tm1Koni/*b*-tm1Koni mice
and wildtype (WT) mice following permanent coronary artery ligation or sham surgery. LV-Left
ventricle. **b**) PCA plot of protein clusters within each animal across the three groups and heatmap
of proteins significantly changed between WT heart failure (WT-HF) and sham. Transfer of
isolated MHC II-deficient splenic B cells from heart failure mice (MHC II KO-HF) shifts cardiac
proteome of naïve recipients towards control. **c**) Volcano plot of proteomics comparison between
WT and MHC II-deficient heart failure recipient mice. Protein markers of adverse cardiac
remodeling are down regulated in naïve recipients of heart failure MHC II-deficient splenic B cells
relative to wildtype recipients. n=3-4 mice per group. LIMMA statistical test.

1041

1042

Fig.6: Splenic Follicular B cells promote adverse cardiac remodeling in ischemic HF a)
Schema for adoptive transfer studies. Four weeks following permanent LAD coronary ligation,
follicular splenic B cells or marginal zone splenic B (MZB) cells were transferred to naïve recipient
mice. The adoptive transfer of unsorted splenic B cells from mice 4 weeks following coronary
artery ligation (HF control) or sham surgery served as positive and negative controls, respectively.
Recipient mice underwent serial transthoracic echocardiography over 8 weeks. **b)** On
echocardiography, adoptive transfer of splenic follicular B cells from heart failure (HF) mice
resulted in a reduction in left ventricular ejection fraction (LVEF) (p<0.001) and an increase in left
ventricular end-systolic diameter (LVESD) in naïve recipient mice over eight weeks (p=0.05)
compared to recipients of splenic B cells from sham-operated mice. The adoptive transfer of
marginal zone B cells from HF mice did not result in LVEF reduction or LV dilatation. n = 8 mice
per group. Mean values ± SD are represented.

1055

1056

1057

Fig.7: Adoptive transfer of splenic B cells from mice with ischemic HF produces MHC-II dependent changes in myocardial T cells 8 weeks post transfer a) Schema for scRNA-seq based analysis of chronic changes in myocardial immune cells mediated by splenic B cells in the context of ischemic HF. 4 weeks post-MI splenic B cells from WT or *Cd19tm1(cre)Cgn/-H2-Ab1b*-tm1Koni/ *b*-tm1Koni (MHC II KO) mice were isolated and transferred into naïve recipients. 8 weeks later myocardial immune cells were isolated from recipient animals and analyzed via scRNA-seq. **b**) UMAP plots of myocardial immune cells 8 weeks post adoptive transfer of splenic B cells from WT sham, WT HF or MHC II deficient HF mice. **c**) Ingenuity Pathway Analysis (IPA) generated volcano plots of significantly differentially expressed genes; adjusted p<0.05, (n = 6 mice per group). HF= heart failure.

1068

1069

1070

Fig.8: Antigen processing and presentation is differentially expressed in circulating B cells in humans post-STEMI or with ischemic cardiomyopathy **a**) Single cell sequencing of peripheral blood B cells from 38 patients 8-weeks or 24-hours post-STEMI and 38 healthy controls. UMAP plots are shown with B cells identified. **b**) KEGG pathway analysis of differentially expressed genes ($\log_2$ foldchange$\leq$-1$\geq$1, p<0.05) revealed significant differential regulation of antigen processing and presentation in peripheral B cells of patients at both 24 hours and 8 weeks post STEMI. **c**) Single cell sequencing of peripheral blood B cells from 3 patients with ischemic cardiomyopathy (ICM) and 1 healthy patient. UMAP plots are shown with B cells identified. **d**) KEGG pathway analysis of differentially expressed genes ($\log_2$ foldchange$\leq$-0.58$\geq$0.58, FDR<0.05, p<0.05) revealed significant differential expression of antigen processing and presentation in peripheral blood B cells from ICM patients. All immune system pathways with FDR <0.05 are listed. Th = T helper. NOD = nucleotide oligomerization domain. IL = interleukin.

1083