Dataset factors influencing age-related changes in brain structure and function in neurodevelopmental conditions

Marlee M. Vandewouw[1,2*], Yifan (Julia) Ye[1,3], Jennifer Crosbie[4,5], Russell J. Schachar[4,5], Alana Iaboni[1], Stelios Georgiades[6], Robert Nicolson[7], Elizabeth Kelley[8,9,10], Muhammad Ayub[10], Jessica Jones[9,10], Paul D. Arnold[11], Margot J. Taylor[12,13,14,15], Jason P. Lerch[12,16,17], Evdokia Anagnostou[1,12,18], Azadeh Kushki[1,2]

Autism Research Centre, Bloorview Research Institute, Holland Bloorview Kids Rehabilitation Hospital, Toronto, ON, Canada
Institute of Biomedical Engineering, University of Toronto, Toronto, Canada
Division of Engineering Science, University of Toronto, Toronto, Canada
Department of Psychiatry, University of Toronto, Toronto, Canada
Department of Psychiatry, The Hospital for Sick Children, Toronto, Ontario, Canada
Department of Psychiatry and Behavioural Neurosciences, McMaster University, Hamilton, Canada
Department of Psychiatry, Western University, London, Canada
Department of Psychology, Queen's University, Kingston, Canada
Centre for Neuroscience Studies, Queen's University, Kingston, Canada
Department of Psychiatry, Queen's University, Kingston, Canada
The Mathison Centre for Mental Health Research & Education, Cumming School of Medicine, University of Calgary, Calgary, Canada
Program in Neurosciences & Mental Health, The Hospital for Sick Children, Toronto, Canada
Department of Diagnostic Imaging, The Hospital for Sick Children, Toronto, Canada
Department of Psychology, University of Toronto, Toronto, Canada
Department of Medical Imaging, University of Toronto, Toronto, Canada
Wellcome Centre for Integrative Neuroimaging, FMRIB, Nuffield Department of Clinical Neurosciences, University of Oxford, Oxford, United Kingdom,
Department of Medical Biophysics, University of Toronto, Toronto, Canada
Institute of Medical Science, University of Toronto, Toronto, Canada

* Corresponding author:
Azadeh Kushki
Autism Research Centre, Holland Bloorview Kids Rehabilitation Hospital
150 Kilgour Road, Toronto, ON M4G 1R8
Phone: 416-425-6220, extension 3563
Email: akushki@hollandbloorview.ca

Data availability statement:

Funding statement: Funding was provided by the Ontario Brain Institute, Canadian Institutes of Health Research, and New Frontiers in Research Fund.

Conflict of interest disclosure: R. Nicolson has received grants from Brain Canada, Hoffman La Roche, Otsuka Pharmaceuticals, and Maplight Therapeutics. E. Anagnostou has received grants from Roche and Anavex, served as a consultant to Roche, Quadrant Therapeutics, Ono, and Impel Pharmaceuticals, has received in-kind support from AMO Pharma and CRA-Simons Foundation, holds a patent for the device,

"Anxiety Meter", has received royalties from APPI and Springer, and has received an editorial honorarium from Wiley. A. Kushki received grants from the National Science and Engineering Research Council and holds a patent for Anxiety Meter with royalties paid from Awake Labs. R. Schachar has served as a consultant to Ehave, Lilly Corporation, and Highland Therapeutics. The remaining authors have reported no biomedical financial interests or potential conflicts of interest.

Ethics approval statement: The current study was approved by the institution's research ethics board, and both consortia studies were approved by the appropriate boards.

Abstract

With brain structure and function undergoing complex changes throughout childhood

and adolescence, age is a critical consideration in neuroimaging studies, particularly for

those of individuals with neurodevelopmental conditions. However, despite the

increasing use of large, consortium-based datasets to examine brain structure and

function in neurotypical and neurodivergent populations, it is unclear whether age-

related changes are consistent between datasets, and whether inconsistencies related

to differences in sample characteristics, such as demographics and phenotypic

features, exist. To address this, we built models of age-related changes of brain

structure (cortical thickness and surface area; N=1,579) and function (resting-state

functional connectivity strength; N=1,792) in two neurodiverse datasets: the Province of

Ontario Neurodevelopmental network (POND) and the Healthy Brain Network (HBN).

We examined whether deviations from these models differed between the datasets, and

explored whether these deviations were associated with demographic and clinical

variables. We found significant differences between the two datasets for measures of

cortical surface area and functional connectivity strength throughout the brain. For

cortical surface area, the patterns of differences were associated with intelligence, while

for functional connectivity strength, positive associations were observed with head

motion. Our findings highlight that patterns of age-related changes in the brain may be

influenced by demographic and phenotypic characteristics, and thus future studies

should consider these when examining or controlling forage effects in analyses.

Introduction

The brain's structure and function undergo complex and protracted changes

throughout childhood and adolescence due to neurobiological processes such as

synaptogenesis, myelination, and synaptic pruning (Grayson & Fair, 2017; Mills &

Tamnes, 2020; Norbom et al., 2021). Neurodevelopmental differences, such as those

associated with autism spectrum disorder (autism), attention-deficit/hyperactivity

disorder (ADHD), and obsessive-compulsive disorder (OCD), can further impact the

presentation of age-related changes, particularly given the heterogeneity in etiology,

biology, and behaviour (Bruin et al., 2020; Lombardo et al., 2019; Wolfers et al., 2019).

As such, age is a critical consideration in neuroimaging studies of neurotypical and

neurodivergent populations.

While many studies directly examine age-related effects, age is often treated as

a nuisance covariate (Hyatt et al., 2020). With increasing awareness of the importance

of replicability in neuroimaging studies (Poldrack et al., 2017), it is becoming more

common to leverage multiple consortium-based datasets to examine brain structure and

function in neurotypical and neurodivergent populations (Abrol et al., 2023; Grotzinger

et al., 2023; Marek et al., 2022; Nicolaisen-Sobesky et al., 2022; Romer et al., 2019;

Vandewouw et al., 2023). However, these datasets differ in composition in terms factors

which may influence age-related effects, such as participants' demographics (e.g., age,

sex and gender, race and ethnicity, socioeconomic status), diagnosis, co-occurring

conditions, and other phenotypic variables. Thus, treating age in the same manner in

datasets with differing characteristics may not be appropriate.

Studies have examined the consistency of age-related changes in brain structure

and function across neurotypical cohorts (Cao et al., 2017; Herting et al., 2018; Madan

& Kensinger, 2017; Mills et al., 2016; Morandini et al., 2021; Tamnes et al., 2017; Wang

et al., 2021). For example, annual percentage changes in cortical thickness and surface

area across childhood and adulthood have been shown to differ between four samples

of typically developing individuals, with the precise patterns varying across lobes of the

brain (Tamnes et al., 2017). White matter volume has also been shown to peak at

different age ranges in different cohorts, with those from the United States showing

decreased volume across age compared to European samples (Mills et al., 2016).

These findings have important implications for those which not only are directly

examining age-related effects, but also those which treat age as a covariate, as

controlling for age in the same manner does not consider these differences. Yet, no

studies have examined whether these changes are consistent between cohorts of

neurodivergent children, and if not, what factors may be contributing to the

inconsistencies. This is particularly critical to investigate given the large variability in

demographic and phenotypic characteristics of samples across different cohorts, even

when the same diagnosis labels are studied. To this end, we examined age-related

changes of brain structure (cortical thickness and surface area) and function (functional

connectivity strength) in two datasets of neurodiverse children and adolescents, namely

the Province of Ontario Neurodevelopmental network (POND) and the Healthy Brain

Network (HBN). We first built dataset-specific developmental models and evaluated

whether deviations from these models differed between the POND and HBN individuals,

separately for males and females. Next, we explored whether the patterns of between-

dataset differences were associated with demographic and phenotypic variables.

Methods

Participants

70   Neuroimaging and phenotype data were obtained from two independently

collected datasets: the Province of Ontario Neurodevelopmental network (POND;

exported April 2023), and the Healthy Brain Network (HBN; Release 9). For both

datasets, participants who were between 5-19 years of age and who were either

neurotypical or had diagnoses of autism, attention-deficit/hyperactivity disorder (ADHD),

or obsessive-compulsive disorder (OCD) were selected. Structural magnetic resonance

imaging (sMRI) data were available for 868 POND (255 ADHD, 377 autism, 75 OCD,

161 TD) and 1,505 HBN (1,100 ADHD, 162 autism, 36 OCD, 207 TD) participants and

resting-state functional magnetic imaging (rs-fMRI) data were available for 864 POND

(211 ADHD, 385 autism, 92 OCD, 185 TD) and 1,459 HBN (1,068 ADHD, 160 autism,

29 OCD, 202 TD) participants. Informed consent and/or assent was obtained from the

participants and/or caregivers. The current study was approved by the institution's

research ethics board, and both consortia studies were approved by the appropriate

boards.

Data acquisition

For both datasets, T1-weighted magnetization-prepared rapid gradient echo

(MPRAGE) sequences (sMRI data) and five minutes of resting-state data with an echo

planar imaging (EPI) sequence (rs-fMRI data). The POND the rs-fMRI data were

obtained as participants watched either a movie of their choosing or the naturalistic

movie paradigm akin to a screensaver, *Inscapes* (Vanderwal et al., 2015), while HBN

participants viewed a fixation cross. While HBN collected two five-minute resting-state

acquisitions, only one was selected to be consistent with POND. Acquisition parameters

for the sMRI and rs-fMRI protocols for each POND and HBN site are presented in

Supplemental Table 1.

Biological sex was collected as part of both studies and used in the current

analysis; insufficient data on gender was available at the time of export. Phenotypic

measures of core and co-occurring differences for the diagnostic categories (see

Supplemental Table 2), socioeconomic status, and race and ethnicity data were also

available for both datasets and were used to characterize each sample. Socioeconomic

status was determined using the caregiver highest level of education and annual

household income. Race and ethnicity data were collected for both datasets according

to each country's census guidelines. Due to low sample size, categories for both

datasets were collapsed into minoritized race/ethnicity and white for statistical tests.

Image processing

The sMRI data were processed using the CIVET pipeline (version 2.1.0; (Ad-

Dab'bagh et al., 2006)) on CBRAIN (Sherif et al., 2014) with the automatic quality

control pipeline, details of which are described elsewhere (Hammill et al., 2021; Kushki

et al., 2019, 2021). Of the remaining participants that passed quality control, propensity

score matching (Rosenbaum & Rubin, 1985) was used to identify a smaller subset of

participants to ensure the datasets showed no significant difference in median age or

sex proportion (see (Vandewouw et al., 2023), for further details). Measures of cortical

thickness and surface area were extracted for 76 brain regions defined using the

Automated Anatomical Labelling (AAL) atlas (Tzourio-Mazoyer et al., 2002). Across

both datasets, ComBat harmonization (Fortin et al., 2018; Johnson et al., 2007) was

used to adjust all measures for site effects, followed by the regression of total gray

matter volume, using subsampling (10,000 iterations, each selecting 63.2% of the

sample) for both steps to increase robustness (Da-ano et al., 2020).

The rs-fMRI data preprocessing was performed using fMRIPrep (Esteban et al.,

2019, 2020), and is described in detail elsewhere (Vandewouw et al., 2023).

Participants were required to have at least 2/3 of their data within the recommended

motion thresholds to pass quality control. Of the remaining participants, propensity

score matching was used to identify a smaller subset of individuals for which there was

not a statistically significant between-dataset differences in median age, sex proportion,

and median head motion (Vandewouw et al., 2023). The Schaefer cortical (Schaefer et

al., 2018) and Melbourne subcortical (Tian et al., 2020) atlases were used to define 232

brain regions, and pairwise Pearson correlations were computed between region-

averaged time series to construct rs-fMRI connectomes. ComBat harmonization with

subsampling (Da-ano et al., 2020; Fortin et al., 2018; Johnson et al., 2007) was used to

adjust the connectivity values for site effects across both datasets. The resulting

connectomes were thresholded with orthogonal minimum spanning trees to remove

spurious connections (Dimitriadis et al., 2017), and the functional connectivity strength

of each brain region was extracted.

Modeling age-related changes

Computational models relating age to the regional sMRI (area and thickness) and

rs-fMRI (connectivity strength) measures were built using Bayesian linear regression

(BLR; (Huertas et al., 2017)). Unlike traditional regression, BLR estimates the posterior

probability of the regression coefficient, capturing the expected variation or uncertainty

around the predicted age-related change; this enables the computation of a statistical *z*-

score, or deviation score, that is similar to a residual but also considers the uncertainty

around the prediction (Huertas et al., 2017; Rutherford et al., 2022). BLR is comparable

but less computationally expensive than similar techniques such as Gaussian process

regression (Fraza et al., 2021; Huertas et al., 2017). We used the BLR implementation

from the Predictive Clinical Neuroscience toolkit (version 0.25;

https://pcntoolkit.readthedocs.io/en/latest) in Python (version 3.9), applying a B-spline

basis expansion of age with three evenly spaced knots to model non-linearities, using

warping to model normal and non-normal distributions in the brain measures, and using

the Powell method for optimization of the Bayesian hyper-parameters.

Dataset- and sex-specific models were built. To promote generalizability, 80% of

participants of a given subgroup were used to build a model, with 20% withheld for

testing. This procedure was repeated 10,000 times, each time with a different randomly

selected subset of participants. Deviation scores (z-scores) were calculated from a built

model under two test scenarios: the model built on 80% of a given subgroup was tested

on 1) the remaining 20% of participants from the same subgroup, and 2) all other

participants. **Figure 1** depicts an example of this process where the model is built using

80% of the female participants in POND and tested on the remaining 20% of female

participants from POND, 100% of male participants from POND, and 100% of the HBN

participants (male and female). The models were also used to predict the expected

brain measure at 7-, 11-, and 15-years of age (approximately the first, second, and third

age quartiles). Median predictions and deviation scores were computed across all

iterations. This procedure was performed with both the full samples and smaller,

matched subsamples.

Figure 1: An example for the POND females of the subsampling approach for comparing the dataset- and sex-specific BLR models.

Statistics

First, differences between the two datasets in their demographic and clinical
characteristics were examined, separately for males and females. Chi-squared tests,
Kruskal-Wallis tests (given non-normality according to Shapiro-Wilks tests), and ordinal
regressions were used for categorical, continuous, and ordinal variables, respectively,
reporting Cramer's *v*, eta-squared, and odds ratio effect sizes; in all cases, significance
was held at p<0.05.

For each of the brain measures, we compared the deviations between the
primary (e.g., POND) and secondary (e.g., HBN) datasets; this was performed with both
POND and HBN defined as the primary dataset, and separately for males and females.
One-way ANOVAs or Kruskal-Wallis tests were used where appropriate, resulting *p*-
values were adjusted using the false discovery rate (FDR) method, holding significance
at *q*<0.05 and reporting eta-squared effect sizes. Results for the full and matched
samples are presented in the main text and **Supplemental information**, respectively.

Given differences in demographic (socioeconomic status and race/ethnicity) and

clinical (diagnosis and phenotypic measures) variables between POND and HBN, we

explored what factors may be contributing to the between-dataset differences. To do so,

for each brain measure, we extracted the deviations for each participant from their

dataset- and sex-specific model and used generalized linear mixed-effects models to

examine their associations with these variables; dataset and sex were modeled as

random factors. Resulting p-values were FDR-corrected across the brain regions, and

significance was held at $q < 0.05$.

Results

Participants

For the sMRI dataset, 754 HBN (214 female, 540 male) and 825 POND (270

female, 555 male) children and adolescents were included in the analysis; participant

demographics are presented in **Table 1**, with phenotypic measures, socioeconomic

status, and race/ethnicity for each dataset are presented in **Table 2**. Given the

difference in age between the two datasets (females: $H(1)=104.35$, $p<.001$, $\eta^2=.04$;

males: $H(1)=203.75$, $p<.001$, $\eta^2=.03$), a smaller, matched subset of participants was

identified for which no significant difference remained, with each dataset consisting of

609 (178 females, 431 males) individuals (**Supplemental Table 3** and **4**).

Table 1: Participant demographics for the full sMRI sample.

			HBN	POND	Test-statistic [1]	p-value	Effect size
Female	***N***		214	270	–	–	–
	Dx (*N*)	**ADHD**	185	56	104.35	<.001	.46
		Autism	16	70			
		OCD	57	60			
		TD	12	28			
	Age² (years)		10.0 [8.1, 13.4]	12.0 [9.6, 15.0]	21.53	<.001	.04
Male	***N***		540	555	–	–	–
	Dx (*N*)	**ADHD**	399	167	203.75	<.001	.43
		Autism	74	245			
		OCD	69	86			
		TD	13	42			
	Age² (years)		10.2 [7.9, 13.1]	11.8 [9.6, 14.6]	38.30	<.001	.03

Note. HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; Dx: diagnosis; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder; TD: typically developing; [1]Kruskal-Wallis H-statistic for age with eta-squared effect size, and chi-squared statistic for diagnosis with Cramer's v effect size. [2]Age is reported as median [IQR].

Table 2: Socioeconomic status, race/ethnicity, and phenotypic measures for the full sMRI sample.

		HBN	POND	Test-statistic[1]	*p*-value	Effect size
Females	**Race/ethnicity[2]**					
	White	130	114	31.53	<.001	.28
	Other	119	27			
	Caregiver income[3]					
	<$50k/year	44	23	0.56	.578	1.12
	$50k-$150k	90	67			
	>$150k	86	44			
	Caregiver education[4]					
	< High school	9	1	3.75	<.001	2.07
	High school	34	19			
	College	87	88			
	Graduate	137	35			
	SCQ	6.0 [3.0, 9.0]	5.0 [2.0, 15.0]	0.06	.800	.02
	RBS-R	8.5 [2.0, 28.0]	10.0 [2.0, 28.0]	0.86	.353	.01
	SWAN-I	2.0 [0.0, 5.5]	2.0 [0.0, 6.0]	0.05	.823	<.01
	SWAN-HI	0.0 [0.0, 2.0]	1.0 [0.0, 3.0]	4.16	.041	.01
	TOCS	–	-5.0 [-39.5, 5.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 5.0]	–	–	–	–
	FSIQ	102.0 [88.0, 114.5]	106.0 [94.0, 115.2]	3.31	.069	0.01
	CBCL-I	58.0 [48.0, 66.0]	61.5 [52.0, 69.0]	8.43	.004	0.02
	CBCL-E	56.0 [47.0, 65.0]	54.0 [44.0, 64.0]	1.71	.191	<.01
Males	**Race/ethnicity[2]**					
	White	255	224	49.25	<.001	.24
	Other	264	78			
	Caregiver income[3]					
	<$50k/year	81	61	3.41	.001	1.62
	$50k-$150k	170	150			
	>$150k	186	79			
	Caregiver education[4]					
	< High school	10	5	4.04	<.001	1.70
	High school	97	45			
	College	187	184			
	Graduate	251	75			
	SCQ	8.0 [5.0, 11.0]	10.0 [4.0, 20.0]	25.42	<.001	.07
	RBS-R	17.0 [4.0, 45.0]	16.0 [5.0, 31.0]	4.25	.039	<.01
	SWAN-I	3.0 [0.0, 6.0]	4.0 [1.0, 7.0]	7.48	.006	.01
	SWAN-HI	1.0 [0.0, 4.0]	2.0 [0.0, 6.0]	14.65	<.001	.01
	TOCS	–	-15.0 [-45.0, 2.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 5.0]	–	–	–	–
	FSIQ	101.0 [86.0, 111.0]	100.0 [88.2, 111.0]	0.00	.945	<.01
	CBCL-I	58.0 [50.0, 67.0]	61.0 [52.0, 68.0]	10.25	.001	.01
	CBCL-E	57.0 [48.0, 66.0]	56.0 [46.0, 65.0]	2.21	.137	<.01

sMRI: structural magnetic resonance imaging; HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; TOCS: Toronto Obsessive-Compulsive Scale; CBCL-OCS: Child Behaviour Checklist obsessive-compulsive subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale; [1]Kruskal-Wallis *H*-statistic for the non-normally distributed continuous variables with eta-squared effect sizes, chi-squared statistic for race/ethnicity with Cramer's *v* effect size, and ordinal regression with odds ratios for the ordinal variables; [2]Collected according to the respective country's census guidelines; [3]In the respective country's currency; [4]Collect includes associate and bachelor's degrees

For the fMRI dataset, 1072 HBN (349 females, 723 males) and 720 POND (207

females, 513 males) participants were included in the analysis (**Table 3**); phenotypic

measures, socioeconomic status, and race/ethnicity for each dataset are shown in

Tables 4. Significant between-dataset differences in age (females: $H(1)=21.34$, $p<.001$,

$\eta^2=.04$; males: $H(1)=40.82$, $p<.001$, $\eta^2=.03$) and motion (females: $H(1)=36.50$, $p<.001$,

$\eta^2=.03$; males: $H(1)=29.50$, $p<.001$, $\eta^2=.01$) were observed. Thus, a smaller, matched

subset of children and adolescents with no such differences was identified, with each

dataset consisting of 627 (189 female, 438 male) individuals (**Supplemental Table 5**

and **6**).

Table 3: Participant demographics for the full fMRI sample.

			HBN	POND	Test-Statistic[1]	p-value	Effect size
Females	**N**		349	207	–	–	–
	Dx (N)	ADHD	232	39	142.43	<.001	.51
		Autism	24	71			
		OCD	13	31			
		TD	80	66			
	Age[2] (years)		10.2 [8.5, 13.1]	12.0 [9.5, 14.8]	21.34	<.001	.04
	Motion[2] (mm)		0.2 [0.1, 0.3]	0.1 [0.1, 0.2]	36.50	<.001	.03
Males	**N**		723	5135	–	–	–
	Dx (N)	ADHD	528	135	290.41	<.001	.48
		Autism	99	235			
		OCD	12	57			
		TD	84	86			
	Age[2] (years)		10.6 [8.6, 13.0]	12.1 [9.8, 14.9]	40.82	<.001	.03
	Motion[2] (mm)		0.2 [0.1, 0.3]	0.2 [0.1, 0.3]	29.50	<.001	.01

HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder; TD: typically developing; [1]Kruskal-Wallis H-statistic for age with eta-squared effect size, and chi-squared statistic for diagnosis with Cramer's v effect size; [2]Age and motion is reported as median [IQR].

Table 4: Socioeconomic status, race/ethnicity, and phenotypic measures for the full fMRI sample.

		HBN	POND	Test-statistic[1]	*p*-value	Effect size
Females	**Race/ethnicity[2]**					
	White	175	112	30.81	<.001	.26
	Other	153	27			
	Caregiver income[3]					
	<$50k/year	55	24			
	$50k-$150k	117	67	0.69	.489	1.14
	>$150k	136	57			
	Caregiver education[4]					
	< High school	7	1			
	High school	42	18	3.79	<.001	2.00
	College	113	86			
	Graduate	183	42			
	SCQ	5.5 [3.0, 9.0]	5.0 [2.0, 15.0]	0.08	.782	.03
	RBS-R	8.0 [2.0, 27.5]	9.5 [2.0, 28.0]	0.59	.441	.01
	SWAN-I	2.0 [0.0, 6.0]	2.0 [0.0, 6.0]	0.15	.694	<.01
	SWAN-HI	0.0 [0.0, 2.0]	1.0 [0.0, 3.0]	4.53	.033	.01
	TOCS	–	-6.0 [-40.0, 6.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 4.0]	–	–	–	–
	FSIQ	102.0 [90.5, 113.0]	106.0 [94.0, 115.0]	3.03	.082	<.01
	CBCL-I	56.0 [48.0, 65.2]	62.0 [52.0, 70.0]	15.17	<.001	.03
	CBCL-E	55.0 [47.0, 64.0]	54.0 [44.0, 63.0]	2.41	.121	.01
Males	**Race/ethnicity[2]**					
	White	332	223	59.40	<.001	.25
	Other	352	74			
	Caregiver income[3]					
	<$50k/year	123	53			
	$50k-$150k	240	143	1.27	.205	1.18
	>$150k	276	110			
	Caregiver education[4]					
	< High school	15	5			
	High school	125	44	3.52	<.001	1.55
	College	243	175			
	Graduate	329	84			
	SCQ	7.0 [4.0, 11.0]	9.0 [3.2, 19.0]	23.86	<.001	.06
	RBS-R	15.0 [3.0, 45.0]	15.0 [5.0, 28.0]	3.78	.052	.01
	SWAN-I	3.0 [0.0, 6.0]	4.0 [1.0, 7.0]	7.23	.007	.01
	SWAN-HI	1.0 [0.0, 3.0]	2.0 [0.0, 5.0]	30.42	<.001	.03
	TOCS	–	-14.5 [-45.0, 2.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 4.0]	–	–	–	–
	FSIQ	100.0 [87.0, 110.0]	102.0 [89.8, 112.0]	1.35	.245	<.01
	CBCL-I	58.0 [50.0, 66.0]	61.0 [52.0, 68.0]	11.50	.001	.01
	CBCL-E	55.0 [48.0, 64.0]	55.0 [46.0, 65.0]	0.05	.820	<.01

fMRI: functional magnetic resonance imaging; HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; TOCS: Toronto Obsessive-Compulsive Scale; CBCL-OCS: Child Behaviour Checklist obsessive-compulsive subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale; [1]Kruskal-Wallis *H*-statistic for the non-normally distributed continuous variables with eta-squared effect sizes, chi-squared statistic for race/ethnicity with Cramer's *v* effect size, and ordinal regression with odds ratios for the ordinal variables; [2]Collected according to the respective country's census guidelines; [3]In the respective country's currency; [4]Collect includes associate and bachelor's degrees

Dataset differences in deviations

The deviation scores for the age models were compared between datasets,

separately for the males and females, with significant differences presented in **Figure 2**

for the full samples, and **Supplemental Figure 1** for the matched samples. To explore

what factors may be contributing to the observed differences, associations between

deviations and the demographic variables (age, sex, race/ethnicity, caregiver highest

level of education, and annual household income) and phenotypic variables (social

communication difficulties, repetitive behaviours, inattention, hyperactivity/impulsivity,

obsessive-compulsive traits, intelligence, internalizing problems, and externalizing

problems) were investigated.

For cortical surface area (**Figure 2A** and **Supplemental Figure 1A**), deviation

scores differed between the datasets throughout the brain, with consistent findings

between the full and matched samples. When examining the median deviation scores

for the significant regions, deviations for the POND participants were negative when the

models were built using the HBN participants, indicating that the HBN models

overestimated surface area for the POND individuals; on the other hand, POND models

underestimated surface area in the HBN participants (positive deviation scores). In both

the full and matched samples, significant positive associations between deviations and

intelligence were observed (**Supplemental Figure 2A**); all other associations were

insignificant (**Supplemental Table 7**).

Figure 2: Effect sizes (eta-squared) for significant between-dataset differences in deviations from age-related models for the females (left) and males (right) from the full sample when the models were built using each dataset (HBN: Healthy Brain Network, POND: Province of Ontario Neurodevelopmental network).

For cortical thickness, few regions showed significant between-dataset

differences in the deviation scores in the full sample (**Figure 2B**), with no regions

showing significant effects in the matched sample (**Supplemental Figure 1B**).

For functional connectivity strength (**Figure 2C**; **Supplemental Figure 1C**),

significant differences in the deviation scores between POND and HBN were observed

throughout the brain, with more widespread effects observed in the males (HBN: 232

regions, POND: 232 regions) compared to females (HBN: 137 regions, POND: 151

regions). When models were built using the HBN individuals, median deviation scores

indicated that they underestimated functional connectivity strength in POND;

conversely, POND models overestimated connectivity strength in the HBN participants.

Deviation scores were significantly positively correlated with head motion

(**Supplemental Figure 2B**; **Supplemental Table 8**).

Discussion

In this study, using data from two consortia, namely POND and HBN, we built

dataset-specific models that describe age-related changes in measures of brain

structure and function. We determined that deviations from the models were

significantly different between the two datasets for measures of cortical surface area

and functional connectivity strength throughout the brain, but not cortical thickness;

these differences were observed within both males and females. For cortical surface

area, the patterns of differences were associated intelligence, suggesting that it may be

contributing to the between-dataset findings. On the other hand, for functional

connectivity strength, positive associations were observed with head motion.

Overall, our findings highlight that patterns of age-related changes in the brain

may not be consistent between cohorts of neurodiverse individuals, and these

inconsistencies may be related to differences in the demographic and phenotypic

composition of the sample. While previous work has identified factors such as

methodology and medication (Bednarz & Kana, 2018) as contributors to the lack of

convergence in the literature on the nature and directionality of brain differences in

neurodevelopmental conditions, we hypothesize that differences in age-related changes

between cohorts, potentially driven by differences in measures like intelligence and

head motion, are also contributing to the lack of consistent findings. Even though age

may not be the focus of a particular study, adjusting for age in the same manner in

multiple cohorts despite the presence of cohort-specific age-related effects can produce

downstream effects in subsequent analyses.

For cortical surface area, differences in deviations from age models differed

between POND and HBN throughout the brain, irrespective of which group was used to

construct the model. Deviations from age models of cortical surface area were also

shown to be positively associated with cortical surface area. The association between

cortical surface area and intelligence is well established (Lett et al., 2020; Schnack et

al., 2015; Tadayon et al., 2020; Yang et al., 2023), with indication that intelligence is

associated with both the magnitude and timing of cortical surface area development

(Schnack et al., 2015).

While between-dataset differences in deviations from age models were observed

for cortical surface area, this finding did not extend to cortical thickness. The

development of cortical thickness and surface area are driven by different genetic and

biological processes and show distinct maturational trajectories (Geschwind & Rakic,

2013; Panizzon et al., 2009; Tamnes et al., 2017; Vijayakumar et al., 2016; Wierenga et

al., 2014). Surface area has also been shown to relate to general psychopathology,

while this association is not observed with cortical thickness (Jalbrzikowski et al., 2019;

Mewton et al., 2022; Romer et al., 2023; Schmaal et al., 2016), and the genetic

association between brain volume and intelligence is driven by surface area and not

thickness (Vuoksimaa et al., 2015). Thus, given the differences between POND and

HBN in the core and co-occurring domains of neurodevelopmental conditions, we

hypothesize that the unique relationship between psychopathology and intelligence with

surface area is why differences in deviations are restricted to this structural measure

rather than cortical thickness.

We also observed differences between POND and HBN in deviations from

models describing changes in functional connectivity strength with age. Deviations from

the models of age-related effects were also shown to be positively associated with head

motion. Head motion is known to increase functional connectivity (Satterthwaite et al.,

2013), and head motion is different between POND and HBN, possibly due to the use of

a fixation-cross resting-state compared to movie paradigms (Vanderwal et al., 2015).

Thus, we suggest that head motion may be driving deviations in age-related changes of

functional connectivity; this conclusion is supported by decreased effect sizes in the

motion-matched subsample, although this is only observed in the males.

The results of this study should be interpreted considering several limitations.

This study is cross-sectional, and future studies should leverage longitudinal data to

validate our developmental findings. We also only constructed age models that were

specific to a binarized measure of biological sex; future work should also examine

gender-specific patterns. Finally, given the sample size required to establish

reproducible brain-behaviour associations (Marek et al., 2022), the contribution of the

identified factors to deviations from age models should be validated in larger samples.

The replicability crisis has posed a significant challenge to the neuroscience field

and to the broader scientific community. In this context, we have questioned the extent

to which age-related changes of brain structure and function are consistent between two

independent cohorts of individuals with neurodevelopmental conditions. While deriving

the neurobiological changes that occur during development is often a specific research

goal, the presence or absence of replicable developmental effects is also highly relevant

for all investigations which include age at any step in the methodology. Given our

findings that age-related differences are not consistent across phenotypic and

acquisition variables, these need to be considered as covariates when examining, or

correcting, for age.

Acknowledgements: We would like to thank all participants and their families for their
involvement in this study. We would also like to thank all individuals who work for POND
for their invaluable assistance in data collection.

References

Abrol, A., Fu, Z., Du, Y., Wilson, T. W., Wang, Y. P., Stephen, J. M., & Calhoun, V. D. (2023). Developmental and aging resting functional magnetic resonance imaging brain state adaptations in adolescents and adults: A large N (>47K) study. *Human Brain Mapping*, *1*, 0. https://doi.org/10.1002/HBM.26200

Ad-Dab'bagh, Y., Einarson, D., Lyttelton, O., Muehlboeck, J.-S., Mok, K., Ivanov, O., Vincent, R. D., Lepage, C., Lerch, J., Fombonne, E., & Evans, A. C. (2006). The CIVET Image-Processing Environment: A Fully Automated Comprehensive Pipeline for Anatomical Neuroimaging Research. *Proceedings of the 12th Annual Meeting of the Organization for Human Brain Mapping*.

Bednarz, H. M., & Kana, R. K. (2018). Advances, challenges, and promises in pediatric neuroimaging of neurodevelopmental disorders. *Neuroscience & Biobehavioral Reviews*, *90*, 50–69. https://doi.org/10.1016/J.NEUBIOREV.2018.03.025

Bruin, W. B., Taylor, L., Thomas, R. M., Shock, J. P., Zhutovsky, P., Abe, Y., Alonso, P., Ameis, S. H., Anticevic, A., Arnold, P. D., Assogna, F., Benedetti, F., Beucke, J. C., Boedhoe, P. S. W., Bollettini, I., Bose, A., Brem, S., Brennan, B. P., Buitelaar, J. K., … van Wingen, G. A. (2020). Structural neuroimaging biomarkers for obsessive-compulsive disorder in the ENIGMA-OCD consortium: medication matters. *Translational Psychiatry 2020 10:1*, *10*(1), 1–12. https://doi.org/10.1038/s41398-020-01013-y

Cao, B., Mwangi, B., Passos, I. C., Wu, M. J., Keser, Z., Zunta-Soares, G. B., Xu, Di., Hasan, K. M., & Soares, J. C. (2017). Lifespan Gyrification Trajectories of Human Brain in Healthy Individuals and Patients with Major Psychiatric Disorders. *Scientific Reports 2017 7:1*, *7*(1), 1–8. https://doi.org/10.1038/s41598-017-00582-1

Da-ano, R., Masson, I., Lucia, F., Doré, M., Robin, P., Alfieri, J., Rousseau, C., Mervoyer, A., Reinhold, C., Castelli, J., De Crevoisier, R., Rameé, J. F., Pradier, O., Schick, U., Visvikis, D., & Hatt, M. (2020). Performance comparison of modified ComBat for harmonization of radiomic features for multicenter studies. *Scientific Reports*, *10*(1), 1–12. https://doi.org/10.1038/s41598-020-66110-w

Dimitriadis, S. I., Antonakakis, M., Simos, P., Fletcher, J. M., & Papanicolaou, A. C. (2017). Data-Driven Topological Filtering Based on Orthogonal Minimal Spanning Trees: Application to Multigroup Magnetoencephalography Resting-State Connectivity. *Brain Connectivity*, *7*(10), 661–670. https://doi.org/10.1089/brain.2017.0512

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: a robust

preprocessing pipeline for functional MRI. *Nature Methods, 16*(1), 111–116. https://doi.org/10.1038/s41592-018-0235-4

Esteban, O., Markiewicz, C. J., Goncalves, M., & DuPre, Elizabeth Kent, James D. Salo, Taylor Ciric, Rastko Pinsard, Basile Blair, Ross W. Poldrack, Russell A. Gorgolewski, K. J. (2020). *fMRIPrep: a robust preprocessing pipeline for functional MRI* (20.2.0). Zenodo. https://doi.org/10.5281/zenodo.4252786

Fortin, J. P., Cullen, N., Sheline, Y. I., Taylor, W. D., Aselcioglu, I., Cook, P. A., Adams, P., Cooper, C., Fava, M., McGrath, P. J., McInnis, M., Phillips, M. L., Trivedi, M. H., Weissman, M. M., & Shinohara, R. T. (2018). Harmonization of cortical thickness measurements across scanners and sites. *NeuroImage, 167*, 104–120. https://doi.org/10.1016/J.NEUROIMAGE.2017.11.024

Fraza, C. J., Dinga, R., Beckmann, C. F., & Marquand, A. F. (2021). Warped Bayesian linear regression for normative modelling of big data. *NeuroImage, 245*, 118715. https://doi.org/10.1016/J.NEUROIMAGE.2021.118715

Geschwind, D. H., & Rakic, P. (2013). *Perspective Cortical Evolution: Judge the Brain by Its Cover*. https://doi.org/10.1016/j.neuron.2013.10.045

Grayson, D. S., & Fair, D. A. (2017). Development of large-scale functional networks from birth to adulthood: A guide to the neuroimaging literature. *NeuroImage, 160*, 15–31. https://doi.org/10.1016/j.neuroimage.2017.01.079

Grotzinger, A. D., Mallard, T. T., Liu, Z., Seidlitz, J., Ge, T., & Smoller, J. W. (2023). Multivariate genomic architecture of cortical thickness and surface area at multiple levels of analysis. *Nature Communications 2023 14:1, 14*(1), 1–13. https://doi.org/10.1038/s41467-023-36605-x

Hammill, C., Lerch, J. P., Taylor, M. J., Ameis, S. H., Chakravarty, M. M., Szatmari, P., Anagnostou, E., & Lai, M. C. (2021). Quantitative and Qualitative Sex Modulations in the Brain Anatomy of Autism. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging, 6*(9), 898–909. https://doi.org/10.1016/J.BPSC.2021.03.001

Herting, M. M., Johnson, C., Mills, K. L., Vijayakumar, N., Dennison, M., Liu, C., Goddings, A. L., Dahl, R. E., Sowell, E. R., Whittle, S., Allen, N. B., & Tamnes, C. K. (2018). Development of subcortical volumes across adolescence in males and females: A multisample study of longitudinal changes. *NeuroImage, 172*, 194. https://doi.org/10.1016/J.NEUROIMAGE.2018.01.020

Huertas, I., Oldehinkel, M., van Oort, E. S. B., Garcia-Solis, D., Mir, P., Beckmann, C. F., & Marquand, A. F. (2017). A Bayesian spatial model for neuroimaging data based on biologically informed basis functions. *NeuroImage, 161*, 134–148. https://doi.org/10.1016/J.NEUROIMAGE.2017.08.009

Hyatt, C. S., Owens, M. M., Crowe, M. L., Carter, N. T., Lynam, D. R., & Miller, J. D. (2020). The quandary of covarying: A brief review and empirical examination of

covariate use in structural neuroimaging studies on psychological variables. *NeuroImage*, *205*, 116225. https://doi.org/10.1016/J.NEUROIMAGE.2019.116225

Jalbrzikowski, M., Freedman, D., Hegarty, C. E., Mennigen, E., Karlsgodt, K. H., Olde Loohuis, L. M., Ophoff, R. A., Gur, R. E., & Bearden, C. E. (2019). Structural Brain Alterations in Youth With Psychosis and Bipolar Spectrum Symptoms. *Journal of the American Academy of Child & Adolescent Psychiatry*, *58*(11), 1079–1091. https://doi.org/10.1016/J.JAAC.2018.11.012

Johnson, W. E., Li, C., & Rabinovic, A. (2007). Adjusting batch effects in microarray expression data using empirical Bayes methods. *Biostatistics*, *8*(1), 118–127. https://doi.org/10.1093/biostatistics/kxj037

Kushki, A., Anagnostou, E., Hammill, C., Duez, P., Brian, J., Iaboni, A., Schachar, R., Crosbie, J., Arnold, P., & Lerch, J. P. (2019). Examining overlap and homogeneity in ASD, ADHD, and OCD: a data-driven, diagnosis-agnostic approach. *Translational Psychiatry*, *9*(1). https://doi.org/10.1038/s41398-019-0631-2

Kushki, A., Cardy, R. E., Panahandeh, S., Malihi, M., Hammill, C., Brian, J., Iaboni, A., Taylor, M. J., Schachar, R., Crosbie, J., Arnold, P., Kelley, E., Ayub, M., Nicolson, R., Georgiades, S., Lerch, J. P., & Anagnostou, E. (2021). Cross-Diagnosis Structural Correlates of Autistic-Like Social Communication Differences. *Cerebral Cortex*, *31*(11), 5067–5076. https://doi.org/10.1093/cercor/bhab142

Lett, T. A., Vogel, B. O., Ripke, S., Wackerhagen, C., Erk, S., Awasthi, S., Trubetskoy, V., Brandl, E. J., Mohnke, S., Veer, I. M., Nöthen, M. M., Rietschel, M., Degenhardt, F., Romanczuk-Seiferth, N., Witt, S. H., Banaschewski, T., Bokde, A. L. W., Büchel, C., Quinlan, E. B., … Walter, H. (2020). Cortical Surfaces Mediate the Relationship Between Polygenic Scores for Intelligence and General Intelligence. *Cerebral Cortex*, *30*(4), 2708–2719. https://doi.org/10.1093/CERCOR/BHZ270

Lombardo, M. V., Lai, M. C., & Baron-Cohen, S. (2019). Big data approaches to decomposing heterogeneity across the autism spectrum. *Molecular Psychiatry 2019 24:10*, *24*(10), 1435–1450. https://doi.org/10.1038/s41380-018-0321-0

Madan, C. R., & Kensinger, E. A. (2017). Age-related differences in the structural complexity of subcortical and ventricular structures. *Neurobiology of Aging*, *50*, 87–95. https://doi.org/10.1016/J.NEUROBIOLAGING.2016.10.023

Marek, S., Tervo-Clemmens, B., Calabro, F. J., Montez, D. F., Kay, B. P., Hatoum, A. S., Donohue, M. R., Foran, W., Miller, R. L., Hendrickson, T. J., Malone, S. M., Kandala, S., Feczko, E., Miranda-Dominguez, O., Graham, A. M., Earl, E. A., Perrone, A. J., Cordova, M., Doyle, O., … Dosenbach, N. U. F. (2022). Reproducible brain-wide association studies require thousands of individuals. *Nature*, *603*(7902), 654. https://doi.org/10.1038/S41586-022-04492-9

Mewton, L., Lees, B., Squeglia, L. M., Forbes, M. K., Sunderland, M., Krueger, R., Koch, F. C., Baillie, A., Slade, T., Hoy, N., & Teesson, M. (2022). The relationship between brain structure and general psychopathology in preadolescents. *Journal of Child Psychology and Psychiatry*, *63*(7), 734–744. https://doi.org/10.1111/JCPP.13513

Mills, K. L., Goddings, A. L., Herting, M. M., Meuwese, R., Blakemore, S. J., Crone, E. A., Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., & Tamnes, C. K. (2016). Structural brain development between childhood and adulthood: Convergence across four longitudinal samples. *NeuroImage*, *141*, 273–281. https://doi.org/10.1016/J.NEUROIMAGE.2016.07.044

Mills, K. L., & Tamnes, C. K. (2020). Longitudinal structural and functional brain development in childhood and adolescence. *The Oxford Handbook of Developmental Cognitive Neuroscience*. https://doi.org/10.1093/OXFORDHB/9780198827474.013.4

Morandini, H. A. E., Rao, P., Hood, S. D., Zepf, F. D., Silk, T. J., & Griffiths, K. R. (2021). Age-related resting-state functional connectivity of the Vigilant Attention network in children and adolescents. *Brain and Cognition*, *154*, 105791. https://doi.org/10.1016/J.BANDC.2021.105791

Nicolaisen-Sobesky, E., Mihalik, A., Kharabian-Masouleh, S., Ferreira, F. S., Hoffstaedter, F., Schwender, H., Maleki Balajoo, S., Valk, S. L., Eickhoff, S. B., Yeo, B. T. T., Mourao-Miranda, J., & Genon, S. (2022). A cross-cohort replicable and heritable latent dimension linking behaviour to multi-featured brain structure. *Communications Biology 2022 5:1*, *5*(1), 1–11. https://doi.org/10.1038/s42003-022-04244-5

Norbom, L. B., Ferschmann, L., Parker, N., Agartz, I., Andreassen, O. A., Paus, T., Westlye, L. T., & Tamnes, C. K. (2021). New insights into the dynamic development of the cerebral cortex in childhood and adolescence: Integrating macro- and microstructural MRI findings. *Progress in Neurobiology*, *204*, 102109. https://doi.org/10.1016/J.PNEUROBIO.2021.102109

Panizzon, M. S., Fennema-Notestine, C., Eyler, L. T., Jernigan, T. L., Prom-Wormley, E., Neale, M., Jacobson, K., Lyons, M. J., Grant, M. D., Franz, C. E., Xian, H., Tsuang, M., Fischl, B., Seidman, L., Dale, A., & Kremen, W. S. (2009). Distinct Genetic Influences on Cortical Surface Area and Cortical Thickness. *Cerebral Cortex*, *19*(11), 2728–2735. https://doi.org/10.1093/CERCOR/BHP026

Poldrack, R. A., Baker, C. I., Durnez, J., Gorgolewski, K. J., Matthews, P. M., Munafò, M. R., Nichols, T. E., Poline, J. B., Vul, E., & Yarkoni, T. (2017). Scanning the horizon: towards transparent and reproducible neuroimaging research. *Nature Reviews Neuroscience 2017 18:2*, *18*(2), 115–126. https://doi.org/10.1038/nrn.2016.167

Romer, A. L., Knodt, A. R., Sison, M. L., Ireland, D., Houts, R., Ramrakha, S., Poulton, R., Keenan, R., Melzer, T. R., Moffitt, T. E., Caspi, A., & Hariri, A. R. (2019). Replicability of structural brain alterations associated with general psychopathology: evidence from a population-representative birth cohort. *Molecular Psychiatry 2019 26:8*, *26*(8), 3839–3846. https://doi.org/10.1038/s41380-019-0621-z

Romer, A. L., Ren, B., & Pizzagalli, D. A. (2023). Brain Structure Relations With Psychopathology Trajectories in the Adolescent Brain Cognitive Development Study. *Journal of the American Academy of Child & Adolescent Psychiatry*. https://doi.org/10.1016/J.JAAC.2023.02.002

Rosenbaum, P. R., & Rubin, D. B. (1985). Constructing a control group using multivariate matched sampling methods that incorporate the propensity score. *American Statistician*, *39*(1), 33–38. https://doi.org/10.1080/00031305.1985.10479383

Rutherford, S., Mostafa Kia, S., Wolfers, T., Fraza, C., Zabihi, M., Dinga, R., Berthet, P., Worker, A., Verdi, S., Ruhe, H. G., Beckmann, C. F., & Marquand, A. F. (2022). The normative modeling framework for computational psychiatry. *Nature Methods*, *17*, 1711–1734. https://doi.org/10.1038/s41596-022-00696-5

Satterthwaite, T. D., Elliott, M. A., Gerraty, R. T., Ruparel, K., Loughead, J., Calkins, M. E., Eickhoff, S. B., Hakonarson, H., Gur, R. E. R. C., Gur, R. E. R. C., & Wolf, D. H. (2013). An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *NeuroImage*. https://doi.org/10.1016/j.neuroimage.2012.08.052

Schaefer, A., Kong, R., Gordon, E. M., Laumann, T. O., Zuo, X. N., Holmes, A. J., Eickhoff, S. B., & Yeo, B. T. T. (2018). Local-global parcellation of the human cerebral cortex from intrinsic functional connectivity mri. *Cerebral Cortex*, *28*(9), 3095–3114. https://doi.org/10.1093/cercor/bhx17

Schmaal, L., Hibar, D. P., Sämann, P. G., Hall, G. B., Baune, B. T., Jahanshad, N., Cheung, J. W., Van Erp, T. G. M., Bos, D., Ikram, M. A., Vernooij, M. W., Niessen, W. J., Tiemeier, H., Hofman, A., Wittfeld, K., Grabe, H. J., Janowitz, D., Bülow, R., Selonke, M., … Veltman, D. J. (2016). Cortical abnormalities in adults and adolescents with major depression based on brain scans from 20 cohorts worldwide in the ENIGMA Major Depressive Disorder Working Group. *Molecular Psychiatry 2016 22:6*, *22*(6), 900–909. https://doi.org/10.1038/mp.2016.60

Schnack, H. G., Van Haren, N. E. M., Brouwer, R. M., Evans, A., Durston, S., Boomsma, D. I., Kahn, R. S., & Hulshoff Pol, H. E. (2015). Changes in Thickness and Surface Area of the Human Cortex and Their Relationship with Intelligence. *Cerebral Cortex*, *25*(6), 1608–1617. https://doi.org/10.1093/CERCOR/BHT357

Sherif, T., Rioux, P., Rousseau, M.-E., Kassis, N., Beck, N., Adalat, R., Das, S., Glatard, T., & Evans, A. C. (2014). CBRAIN: a web-based, distributed computing platform for collaborative neuroimaging research. *Frontiers in Neuroinformatics*, *8*, 54. https://doi.org/10.3389/fninf.2014.00054

Tadayon, E., Pascual-Leone, A., & Santarnecchi, E. (2020). Differential Contribution of Cortical Thickness, Surface Area, and Gyrification to Fluid and Crystallized Intelligence. *Cerebral Cortex*, *30*(1), 215–225. https://doi.org/10.1093/CERCOR/BHZ082

Tamnes, C. K., Herting, M. M., Goddings, A. L., Meuwese, R., Blakemore, S. J., Dahl, R. E., Güroğlu, B., Raznahan, A., Sowell, E. R., Crone, E. A., & Mills, K. L. (2017). Development of the Cerebral Cortex across Adolescence: A Multisample Study of Inter-Related Longitudinal Changes in Cortical Volume, Surface Area, and Thickness. *Journal of Neuroscience*, *37*(12), 3402–3412. https://doi.org/10.1523/JNEUROSCI.3302-16.2017

Tian, Y., Margulies, D. S., Breakspear, M., & Zalesky, A. (2020). Topographic organization of the human subcortex unveiled with functional connectivity gradients. *Nature Neuroscience*, *23*(11), 1421–1432. https://doi.org/10.1038/s41593-020-00711-6

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, *15*, 273–289. https://doi.org/10.1006/nimg.2001.0978

Vanderwal, T., Kelly, C., Eilbott, J., Mayes, L. C., & Castellanos, F. X. (2015). Inscapes: A movie paradigm to improve compliance in functional magnetic resonance imaging. *NeuroImage*, *122*, 222–232. https://doi.org/10.1016/j.neuroimage.2015.07.069

Vandewouw, M. M., Brian, J., Crosbie, J., Schachar, R. J., Iaboni, A., Georgiades, S., Nicolson, R., Kelley, E., Ayub, M., Jones, J., Taylor, M. J., Lerch, J. P., Anagnostou, E., & Kushki, A. (2023). Identifying Replicable Subgroups in Neurodevelopmental Conditions Using Resting-State Functional Magnetic Resonance Imaging Data. *JAMA Network Open*, *6*(3), e232066–e232066. https://doi.org/10.1001/JAMANETWORKOPEN.2023.2066

Vijayakumar, N., Allen, N. B., Youssef, G., Dennison, M., Yücel, M., Simmons, J. G., & Whittle, S. (2016). Brain development during adolescence: A mixed-longitudinal investigation of cortical thickness, surface area, and volume. *Human Brain Mapping*, *37*(6), 2027–2038. https://doi.org/10.1002/HBM.23154

Vuoksimaa, E., Panizzon, M. S., Chen, C.-H., Fiecas, M., Eyler, L. T., Fennema-Notestine, C., Hagler, D. J., Fischl, B., Franz, C. E., Jak, A., Lyons, M. J., Neale, M. C., Rinker, D. A., Thompson, W. K., Tsuang, M. T., Dale, A. M., & Kremen, W. S.

(2015). The Genetic Association Between Neocortical Volume and General Cognitive Ability Is Driven by Global Surface Area Rather Than Thickness. *Cerebral Cortex*, *25*(8), 2127–2137. https://doi.org/10.1093/cercor/bhu018

Wang, Q., Hu, K., Wang, M., Zhao, Y., Liu, Y., Fan, L., & Liu, B. (2021). Predicting brain age during typical and atypical development based on structural and functional neuroimaging. *Human Brain Mapping*, *42*(18), 5943–5955. https://doi.org/10.1002/HBM.25660

Wierenga, L. M., Langen, M., Oranje, B., & Durston, S. (2014). Unique developmental trajectories of cortical thickness and surface area. *NeuroImage*, *87*, 120–126. https://doi.org/10.1016/J.NEUROIMAGE.2013.11.010

Wolfers, T., Beckmann, C. F., Hoogman, M., Buitelaar, J. K., Franke, B., & Marquand, A. F. (2019). Individual differences v. the average patient: Mapping the heterogeneity in ADHD using normative models. *Psychological Medicine*, *50*(2), 314–323. https://doi.org/10.1017/S0033291719000084

Yang, S., Ma, X., Xia, X., Qiao, Z., Huang, M., Wang, N., Hu, X., Zhang, X., Deng, W., Kang, L., Li, X., Hao, G., Xi, J., Meng, H., Li, T., Hou, X., & Fu, Y. (2023). A Bivariate Twin Study of Cortical Surface Area and Verbal and Nonverbal Intellectual Skills in Adolescence. *Neuroscience*. https://doi.org/https://doi.org/10.1016/j.neuroscience.2023.04.009

Supplemental information

Supplemental Table 1: Acquisition parameters for the sMRI and rs-fMRI protocols for each site in the POND and HBN datasets.

Dataset	Scanner	Scan type	Sequence	TR (ms)	TE (ms)	FA (°)	FOV (mm)	Voxel size (mm)	Scan time (min)
sMRI protocol									
	SK: 3T TimTrio	–	MPRAGE	2300	2.96	9	192×240×256	1	5.0
	QU: 3T TimTrio	–	MPRAGE	2300	3.14	9	192×240×256	0.8	6.2
	SK: 3T Prisma^fit	–	MPRAGE	1870	3.14	9	192×240×256	0.8	5.0
	QU: 3T Prisma^fit	–	MPRAGE	1870	3.10	9	192×240×256	0.8	5.0
POND	HB: 3T Prisma^fit	–	MPRAGE	1870	3.10	9	192×240×256	0.8	5.0
	CBIC: 3T Prisma^fit	–	MPRAGE	2500	3.15	8	179×256×256	0.8	7.00
	CUNY: 3T Prisma^fit	–	MPRAGE	2500	3.15	8	179×256×256	0.8	7.00
	RU: 3T TimTrio	–	MPRAGE	2500	3.15	8	179×256×256	0.8	7.00
HBN	SI: 1.5T Avanto	–	MPRAGE	2730	1.64	7	176×256×256	1	6.53
fMRI protocol									
	SK: 3T TimTrio	Movies	EPI	2340	30	70	224×224×140	3.5	5.1
	QU: 3T TimTrio	Inscapes	EPI	2340	30	70	224×224×140	3.5	5.1
	SK: 3T Prisma^fit	Inscapes	EPI	1500	30	70	222×222×150	3.0	5.1
	QU: 3T Prisma^fit	Inscapes	EPI	1500	30	70	222×222×150	3.0	5.1
POND	HB: 3T Prisma^fit	Inscapes	EPI	1500	30	70	222×222×150	3.0	5.1
	CBIC: 3T Prisma^fit	Fixation	EPI	800	30	31	202×202×144	2.4	2×5.1
	CUNY: 3T Prisma^fit	Fixation	EPI	800	30	31	202×202×144	2.4	2×5.1
	RU: 3T TimTrio	Fixation	EPI	800	30	31	202×202×144	2.4	2×5.1
	SI: 1.5T Avanto	Fixation	EPI	1450	40	55	195×195×135	2.5	10.3
	SK: 3T TimTrio	Movies	EPI	2340	30	70	224×224×140	3.5	5.0
HBN	QU: 3T TimTrio	Inscapes	EPI	2340	30	70	224×224×140	3.5	5.0

sMRI: structural magnetic resonance imaging; rs-fMRI: resting-state functional magnetic resonance imaging; POND: Province of Ontario Neurodevelopmental network; HBN: Healthy Brain Network; SK: Hospital for Sick Children; QU: Queen's University; HB: Holland Bloorview Kids Rehabilitation Hospital; CBIC: CitiGroup Cornell Brain Imaging Center; CUNY: the City University of New York Advanced Science Research Center; RU: Rutgers University; SI: Staten Island; TR: repetition time; TE: echo time; FA: flip angle; FOV: field of view

Supplemental Table 2: Summary of the phenotypic measures of core and co-occurring differences for the diagnostic categories.

Domain	Assessment	Score	Range	Interpretation[1]	Acronym
Social communication difficulties	Social Communication Questionnaire (Rutter et al., 2003)	Total score	0-39	More severe difficulties with social communication	SCQ
Repetitive behaviours	Repetitive Behaviours Scale – Revised (Bodfish et al., 2000)	Total score	0-129	More severe repetitive behaviours	RBS-R
Inattention	Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour (Swanson et al., 2012)	Inattention subscale	0-9	More severe difficulties with inattention	SWAN-I
Hyperactivity	Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour (Swanson et al., 2012)	Hyperactivity subscale	0-9	More severe difficulties with hyperactivity/impulsivity	SWAN-HI
Obsessive-compulsive traits	Toronto Obsessive-Compulsive Scale (Nelson et al., 2001)	Total score	-63-63	More severe obsessive-compulsive traits	TOCS
	Child Behaviour Checklist (Nelson et al., 2001)	Obsessive-compulsive subscale	0-16	More severe obsessive-compulsive traits	CBCL-OCS
Intelligence	Age-appropriate Wechsler or Stanford-Binet scales	Full-scale intelligence quotient	40-160	Higher intelligence	FSIQ
Internalizing problems	Child Behaviour Checklist (Achenbach & Ruffle, 2000)	Internalizing problems subscale	40-75	Worse internalizing problems	CBCL-I
Externalizing problems	Child Behaviour Checklist (Achenbach & Ruffle, 2000)	Externalizing problems subscale	40-75	Worse externalizing problems	CBCL-E

[1]Interpretation: relative to higher scores

Supplemental Table 3: Participant demographics for the matched sMRI sample.

			HBN	POND	Test-Statistic[1]	p-value	Effect size
	N		178	178	–	–	–
Females		ADHD	124	51			
		Autism	7	56	76.33	<.001	.46
		OCD	38	47			
	Dx (*N*)	TD	9	27			
	Age[2] (years)		10.1 [8.5, 13.3]	11.1 [9.0, 13.3]	2.61	.107	<.01
	N		431	431	–	–	–
Males		ADHD	304	145			
	Dx (*N*)	Autism	55	190	141.73	<.001	.41
		OCD	61	63			
		TD	11	33			
	Age[2] (years)		10.6 [8.6, 13.0]	10.9 [9.0, 13.0]	2.07	.150	<.01

HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder; TD: typically developing; [1]Kruskal-Wallis *H*-statistic for age with eta-squared effect size, and chi-squared statistic for diagnosis with Cramer's *v* effect size; [2]Age is reported as median [IQR].

Supplemental Table 4: Socioeconomic status, race/ethnicity, and phenotypic measures for the matched sMRI sample.

		HBN	POND	Test-Statistic[1]	p-value	Effect size
Females	Race/ethnicity[2]					
	White	86	92	21.18	<.001	.28
	Other	78	24			
	Caregiver income[3]					
	<$50k/year	31	18			
	$50k-$150k	65	57	0.03	.974	0.99
	>$150k	51	34			
	Caregiver education[4]					
	< High school	6	1			
	High school	19	15	3.88	<.001	2.44
	College	58	72			
	Graduate	94	27			
	SCQ	6.0 [4.0, 8.0]	5.0 [2.0, 14.2]	0.18	.675	.02
	RBS-R	11.0 [2.0, 28.0]	11.5 [2.0, 29.0]	0.22	.639	.01
	SWAN-I	2.0 [0.0, 5.0]	2.0 [0.0, 7.0]	0.04	.843	<.01
	SWAN-HI	0.0 [0.0, 2.0]	1.0 [0.0, 4.0]	7.16	.007	.02
	TOCS	–	-6.0 [-40.0, 6.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 4.8]	–	–	–	–
	FSIQ	102.0 [87.0, 113.5]	106.0 [94.0, 115.0]	1.90	.168	.01
	CBCL-I	56.0 [48.0, 65.0]	60.0 [52.0, 69.0]	6.91	.009	.02
	CBCL-E	57.0 [47.0, 64.8]	55.0 [44.0, 64.0]	0.80	.372	<.01
Males	Race/ethnicity[2]					
	White	205	168	26.58	<.001	.20
	Other	202	68			
	Caregiver income[3]					
	<$50k/year	67	47			
	$50k-$150k	133	123	3.09	.002	1.64
	>$150k	146	58			
	Caregiver education[4]					
	< High school	6	4			
	High school	79	30	3.39	.001	1.66
	College	143	149			
	Graduate	196	57			
	SCQ	8.0 [5.0, 11.0]	10.0 [4.0, 20.0]	19.42	<.001	.06
	RBS-R	14.5 [4.0, 44.0]	16.0 [5.0, 31.0]	0.77	.381	<.01
	SWAN-I	3.0 [0.0, 6.0]	4.0 [1.0, 7.0]	11.37	<.001	.01
	SWAN-HI	1.0 [0.0, 4.0]	2.0 [0.0, 6.0]	22.60	<.001	.03
	TOCS	–	-15.0 [-45.0, 2.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 5.0]	–	–	–	–
	FSIQ	100.5 [87.0, 111.0]	101.0 [88.0, 111.0]	0.07	.794	<.01
	CBCL-I	58.0 [50.0, 67.0]	62.0 [52.0, 68.0]	10.87	.001	.01
	CBCL-E	56.0 [48.0, 65.0]	56.0 [48.0, 66.0]	0.06	.814	<.01

sMRI: structural magnetic resonance imaging; HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; TOCS: Toronto Obsessive-Compulsive Scale; CBCL-OCS: Child Behaviour Checklist obsessive-compulsive subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale; [1]Kruskal-Wallis *H*-statistic for the non-normally distributed continuous variables with eta-squared effect sizes, chi-squared statistic for race/ethnicity with Cramer's *v* effect size, and ordinal regression with odds ratios for the ordinal variables; [2]Collected according to the respective country's census guidelines; [3]In the respective country's currency; [4]Collect includes associate and bachelor's degrees

Supplemental Table 5: Participant demographics for the matched fMRI sample.

			HBN	POND	Test-Statistic[1]	p-value	Effect size
	N		189	189	–	–	–
Females	Dx (N)	ADHD	130	38	107.87	<.001	.53
		Autism	7	65			
		OCD	12	31			
		TD	40	55			
	Age[2] (years)		10.7 [8.7, 13.4]	11.7 [9.4, 14.0]	2.77	.096	.01
	Motion[2] (mm)		0.1 [0.1, 0.2]	0.1 [0.1, 0.2]	2.09	.148	<.01
	N		438	438	–	–	–
Males	Dx (N)	ADHD	321	121	202.91	<.001	.48
		Autism	53	192			
		OCD	7	48			
		TD	57	77			
	Age[2] (years)		10.9 [9.0, 13.1]	11.4 [9.4, 13.6]	2.75	.097	<.01
	Motion[2] (mm)		0.2 [0.1, 0.2]	0.2 [0.1, 0.3]	0.53	.466	<.01

HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; ADHD: attention-deficit/hyperactivity disorder; OCD: obsessive-compulsive disorder; TD: typically developing; [1]Kruskal-Wallis H-statistic for age with eta-squared effect size, and chi-squared statistic for diagnosis with Cramer's v effect size; [2]Age and motion is reported as median [IQR].

Supplemental Table 6: Socioeconomic status, race/ethnicity, and phenotypic measures for the matched fMRI sample.

		HBN	POND	Test-statistic[1]	p-value	Effect size
Females	Race/ethnicity[2]					
	White	102	100	14.34	<.001	.22
	Other	74	27			
	Caregiver income[3]					
	<$50k/year	32	20			
	$50k-$150k	68	62	0.00	.999	1.00
	>$150k	71	52			
	Caregiver education[4]					
	< High school	2	1			
	High school	16	16	4.04	<.001	2.45
	College	69	77			
	Graduate	102	38			
	SCQ	5.0 [3.0, 8.5]	5.0 [2.0, 14.2]	0.01	.917	.03
	RBS-R	10.0 [2.0, 27.2]	11.0 [2.0, 29.0]	0.11	.741	.01
	SWAN-I	2.0 [0.0, 5.8]	2.0 [0.0, 6.0]	0.00	.987	<.01
	SWAN-HI	0.0 [0.0, 2.0]	1.0 [0.0, 3.0]	6.81	.009	.03
	TOCS	–	-5.0 [-39.8, 7.8]	–	–	–
	CBCL-OCS	2.0 [1.0, 5.0]	–	–	–	–
	FSIQ	104.0 [91.0, 116.0]	106.0 [94.0, 115.0]	0.37	.544	<.01
	CBCL-I	56.0 [48.0, 65.0]	62.0 [52.0, 70.0]	10.73	.001	.03
	CBCL-E	54.0 [47.0, 64.2]	54.0 [44.0, 64.0]	0.94	.333	<.01
Males	Race/ethnicity[2]					
	White	219	185	30.07	<.001	.21
	Other	202	67			
	Caregiver income[3]					
	<$50k/year	66	46			
	$50k-$150k	158	124	1.37	.171	1.23
	>$150k	169	97			
	Caregiver education[4]					
	< High school	5	4			
	High school	77	33	3.24	.001	1.60
	College	144	155			
	Graduate	205	73			
	SCQ	7.0 [4.0, 10.0]	9.0 [3.0, 18.5]	14.08	<.001	.06
	RBS-R	12.0 [2.0, 40.8]	15.0 [5.0, 27.0]	0.51	.474	<.01
	SWAN-I	3.0 [0.0, 6.0]	4.0 [1.0, 7.0]	5.85	.016	.01
	SWAN-HI	0.0 [0.0, 3.0]	2.0 [0.0, 6.0]	38.64	<.001	.05
	TOCS	–	-15.0 [-44.0, 2.0]	–	–	–
	CBCL-OCS	2.0 [1.0, 4.0]	–	–	–	–
	FSIQ	99.0 [86.0, 110.0]	101.0 [89.5, 111.5]	1.32	.250	<.01
	CBCL-I	58.0 [49.0, 67.0]	61.0 [52.0, 68.0]	8.84	.003	.01
	CBCL-E	54.0 [46.0, 63.0]	56.0 [46.0, 66.2]	1.97	.161	<.01

fMRI: functional magnetic resonance imaging; HBN: Healthy Brain Network; POND: Province of Ontario Neurodevelopmental network; SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; TOCS: Toronto Obsessive-Compulsive Scale; CBCL-OCS: Child Behaviour Checklist obsessive-compulsive subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale; [1]Kruskal-Wallis *H*-statistic for the non-normally distributed continuous variables with eta-squared effect sizes, chi-squared statistic for race/ethnicity with Cramer's *v* effect size, and ordinal regression with odds ratios for the ordinal variables; [2]Collected according to the respective country's census guidelines; [3]In the respective country's currency; [4]Collect includes associate and bachelor's degrees

Supplemental Table 7: Correlation between the effect sizes for the between-dataset difference in deviation scores and associations with demographic and clinical variables for cortical surface area in the full sample.

	Correlation value	*p*-value
Dx	.16	.007
Race/ethnicity	.02	.674
Caregiver income	.00	.975
Caregiver education	.07	.223
SCQ	.08	.155
RBS-R	-.05	.422
SWAN-I	.14	.014
SWAN-HI	<.01	.990
FSIQ	.21	<.001
CBCL-I	.03	.576
CBCL-E	.09	.101

SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale

Supplemental Table 8: Correlation between the effect sizes for the between-dataset difference in deviation scores and associations with demographic and clinical variables for functional connectivity strength in the full sample.

	Correlation value	*p*-value
Dx	-.05	.127
Motion	.20	<.001
Race/ethnicity	-.18	<.001
Caregiver income	-.13	<.001
Caregiver education	-.03	.417
SCQ	-.01	.803
RBS-R	-.12	<.001
SWAN-I	.02	.576
SWAN-HI	.03	.352
FSIQ	-.06	.072
CBCL-I	-.06	.050
CBCL-E	-.06	.054

SCQ: social communication questionnaire; RBS-R: repetitive Behaviours Scale – Revised; SWAN-I: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour inattention subscale; SWAN-HI: Strengths and Weaknesses of ADHD-symptoms and Normal Behaviour hyperactivity/impulsivity subscale; FSIQ: full-scale intelligence quotient; CBCL-I: Child Behaviour Checklist internalizing problems subscale; CBCL-E: Child Behaviour Checklist externalizing problems subscale

Supplemental Figure 1: Effect sizes (eta-squared) for significant between-dataset differences in deviations from age-related models for the females (left) and males (right) from the matched sample when the models were built using each dataset (HBN: Healthy Brain Network, POND: Province of Ontario Neurodevelopmental network).

Factors associated with deviations

Supplemental Figure 2: Significant association between (A) the deviations from age models of cortical surface area and intelligence (FSIQ) and (B) the deviations from connectivity strength models and head motion. FSIQ: full-scale intelligence quotient

References

Achenbach, T. M., & Ruffle, T. M. (2000). The Child Behavior Checklist and Related Forms for Assessing Behavioral/Emotional Problems and Competencies. *Pediatrics In Review*, *21*(8), 265–271. https://doi.org/10.1542/PIR.21-8-265

Bodfish, J. W., Symons, F. J., Parker, D. E., & Lewis, M. H. (2000). Varieties of repetitive behavior in autism: Comparisons to mental retardation. *Journal of Autism and Developmental Disorders*, *30*(3), 237–243. https://doi.org/10.1023/A:1005596502855

Nelson, E. C., Hanna, G. L., Hudziak, J. J., Botteron, K. N., Heath, A. C., & Todd, R. D. (2001). Obsessive-compulsive scale of the child behavior checklist: specificity, sensitivity, and predictive power. *Pediatrics*, *108*(1). https://doi.org/10.1542/PEDS.108.1.E14

Rutter, M., Bailey, A., & Lord, C. (2003). *The social communication questionnaire*. Western Psychological Services.

Swanson, J. M., Schuck, S., Porter, M. M., Carlson, C., Hartman, C. A., Sergeant, J. A., Clevenger, W., Wasdell, M., McCleary, R., Lakes, K., & Wigal, T. (2012). Categorical and Dimensional Definitions and Evaluations of Symptoms of ADHD: History of the SNAP and the SWAN Rating Scales. *The International Journal of Educational and Psychological Assessment*, *10*(1), 51–70. http://www.ncbi.nlm.nih.gov/pubmed/26504617%0Ahttp://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4618695