Mitochondrial Phosphopantetheinylation is Required for Oxidative Function

Pieter R. Norden[1, *], Riley J. Wedan[1, *], Jacob Z. Longenecker[1], Samuel E. J. Preston[1],

Naomi Graber[1], Olivia A. Pentecost[1], Morgan Canfield[1], Elizabeth McLaughlin[1], and Sara M.

Nowinski[1]

[1]Department of Metabolism and Nutritional Programming, Van Andel Institute, Grand

Rapids, MI

*These authors contributed equally

Correspondence:

Sara.nowinski@vai.org (S.N.)

Abstract

4'-phosphopantetheinyl (4'PP) groups are essential co-factors added to target proteins by

phosphopantetheinyl transferase (PPTase) enzymes. Although mitochondrial 4'PP-modified

proteins have been described for decades, a mitochondrially-localized PPTase has never

been found in mammals. We discovered that the cytoplasmic PPTase aminoadipate

semialdehyde dehydrogenase phosphopantetheinyl transferase (AASDHPPT) is required for

mitochondrial respiration and oxidative metabolism. Loss of AASDHPPT results in failed 4'-

PP modification of the mitochondrial acyl carrier protein and blunted activity of the

mitochondrial fatty acid synthesis (mtFAS) pathway. We found that in addition to its

cytoplasmic localization, AASDHPPT localizes to the mitochondrial matrix via an N-terminal

mitochondrial targeting sequence contained within the first 13 amino acids of the protein.

Our data show that this novel mitochondrial localization of AASDHPPT is required to support

mtFAS activity and oxidative function. We further identify two variants of uncertain

significance in *AASDHPPT* that are likely pathogenic in humans due to loss of mtFAS

activity.

Keywords:

Mitochondria, fatty acid synthesis, electron transport chain, reductive carboxylation,

metabolism, respiration, compartmentalization, phosphopantetheine

Abbreviations:

4'PP – 4'-phosphopantetheine/yl, AASDHPPT – aminoadipate semialdehyde

dehydrogenase phosphopantetheinyl transferase, ALDH1L2 – Aldehyde Dehydrogenase 1

Family Member L2, CoA – coenzyme A, DHRS2 – mitochondrial Dehydrogenase/Reductase

2, ETC – electron transport chain, FAS – fatty acid synthesis, FASN – fatty acid synthase,

LYRM – leucine tyrosine arginine motif, mtFAS – mitochondrial fatty acid synthesis, NAD+ -

nicotinamide adenine dinucleotide (oxidized form), OGDH – oxoglutarate dehydrogenase,

PAGE – polyacrylamide gel electrophoresis, PDH – pyruvate dehydrogenase, PPT –

mammalian AASDHPPT, PPTase – phosphopantetheinyl transferase, PTM –

posttranslational modification, SDHB – succinate dehydrogenase subunit B, TCA –

tricarboxylic acid, VUS – variant of uncertain significance

Main

4'-phosphopantentheinylation is an essential post translational modification (PTM)

whereby a 4'-phosphopantetheine (4'PP) group, derived from coenzyme A (CoA), is covalently

attached to target proteins by a phosphopantetheinyltransferase (PPTase) enzyme[6]. A total of

six proteins in the mammalian proteome have been found to have a 4'-PP-modified serine:

three cytoplasmic proteins (Fatty Acid Synthase, FASN; Aminoadipate semialdehyde

dehydrogenase, AASDH; and Aldehyde Dehydrogenase 1 Family Member L1, AL1L1), and

three mitochondrial proteins (NADH:Ubiquinone Oxidoreductase Subunit AB1, NDUFAB1;

Aldehyde Dehydrogenase 1 Family Member L2, AL1L2; and mitochondrial

Dehydrogenase/Reductase 2, DHRS2)[6]. In mammalian genomes, however, there is only one

annotated PPTase: <u>a</u>mino<u>a</u>dipate-<u>s</u>emialdehyde <u>d</u>e<u>h</u>ydrogenase- <u>p</u>hospho<u>p</u>antetheinyl

<u>t</u>ransferase (AASDHPPT), which was thought to exist solely in the cytoplasm[8]. It was therefore

unclear how 4'PP is attached to mitochondrial proteins.

Since no mammalian mitochondrial PPTase had ever been described, we questioned

whether an unidentified PPTase existed in mammals, or conversely if AASDHPPT (hereafter

abbreviated PPT) was important for mitochondrial function regardless of its reported

cytoplasmic localization. We therefore set out to understand if PPT was required for

mitochondrial function and, if so, how it modifies mitochondrial targets despite existing in

separate cellular compartments.

To test whether PPT is required for mitochondrial oxidative function, we used a

CRISPR/Cas-9-based strategy to mutate *Aasdhppt* in C2C12 mouse skeletal muscle

myoblasts. Complete loss of PPT was incompatible with growth, however we generated two

clonal cell lines with hypomorphic mutations in *Aasdhppt* (Extended Data Fig. 1A) that resulted

in decreased PPT expression (PPT-1 and PPT-2, Fig. 1A). We next asked whether PPT
deficiency resulted in altered respiratory-dependent growth by culturing cells in media with
either glucose or galactose, the latter requiring functional mitochondrial respiration. Clonal
mutant PPT-2 showed significantly reduced growth in glucose-containing media (Fig. 1B,
Extended Data Fig. 1B), while clonal mutant PPT-1 showed a modest, yet non-statistically
significant reduction in growth compared to control cells. In contrast, both PPT mutants
showed markedly reduced proliferation in galactose-containing media (Fig. 1C, Extended
Data Fig. 1B). We next performed a Seahorse mitochondrial stress test and observed that
PPT-deficient mutants exhibited a reduction in both basal respiration rate and spare
respiratory capacity compared to controls (Fig. 1D). To assess whether PPT-deficient
respiratory function was associated with perturbed OXPHOS complex expression and
assembly, we performed blue-native PAGE on isolated mitochondria from PPT-deficient and
control cells. PPT-deficient mutants displayed reduced levels of fully assembled ETC complex
I (CI), complex II (CII) and complex IV (CIV) (Fig. 1E). Complex V (CV) was largely unaffected.
Additionally, complex III (CIII) was not affected by PPT-deficiency but reductions in CIII-
containing supercomplexes (SC) were observed (Fig. 1E). Together, these data demonstrate
that PPT deficiency results in dysfunctional mitochondrial respiratory and ETC deficient
phenotypes.
Many ETC impairments affect mitochondrial metabolism by dysregulating the NAD^+/NADH
ratio, causing a disruption in TCA cycle flux and nutrient utilization. To assess the activity of
the TCA cycle we cultured cells in fully labeled ^{13}C-glutamine for up to 6 hours and evaluated
carbon entry into TCA cycle intermediates to assess reductive (blue) versus oxidative (red)
use of glutamine (Fig. 1F). At steady state, PPT-deficient cells demonstrated an increase in
the pool size of succinate and a decrease in downstream metabolites such as fumarate,
malate, and aspartate (Fig. 1G, Extended Data Fig. 1C-E). Pool sizes of citrate and α-
ketoglutarate were variable (Extended Data Fig. 1F and G); however, the ratio of malate to
citrate was consistently reduced in PPT mutants compared to controls (Fig. 1H). The reduction
of TCA metabolites downstream of succinate relative to citrate suggests a block in oxidative
(NAD^+ consuming) TCA cycle activity.
To assess whether this was indeed the case, we examined citrate labeling patterns.
Citrate M+4 results from the oxidation of glutamine to oxaloacetate and consumes NAD^+,
whereas Citrate M+5 results from the reductive carboxylation of glutamine to citrate and
regenerates NAD^+. PPT-deficient cells showed a reduction in M+4 citrate and an increase in
M+5 citrate compared to controls, indicating increased reductive carboxylation of glutamine

(Extended Data Fig. 1H). This is further supported by the increase in the ratio of citrate

M+5:citrate M+4 at isotopic steady state (Fig. 1I).

Citrate produced in the mitochondria can also be transported to the cytosol to be broken

down by ATP-citrate lyase (ACLY) into malate and acetyl-CoA, with citrate M+4 producing

malate M+4 and citrate M+5 producing malate M+3 (Fig. 1F). PPT-deficient cells displayed a

reduction in malate M+4, which likely results from the loss of M+4 citrate (Extended Data Fig.

1I). However, the PPT-deficient cells did not display a concordant increase in M+3 malate,

indicating that differences in malate labeling patterns result from mitochondrial malate, and

conversion of citrate to malate in the cytosol is similar or reduced in PPT mutants (Extended

Data Fig. 1I). In sum, the glutamine labeling patterns in PPT-deficient cells support a model

in which loss of oxidative metabolism in the mitochondria results in increased reductive

carboxylation of glutamine that is typical of cells with loss of ETC activity[14]. The finding that

malate M+3 is not increased in PPT-deficient cells also supports the notion that the reductive

carboxylation seen in PPT-deficient cells is driven by a need for NAD^+ regeneration and not

by a demand for cytoplasmic acetyl-CoA.

As *Aasdhppt* was clearly important for mitochondrial oxidative function, we next asked

which mitochondrial targets were likely to be most prominently involved in this phenotype. As

mentioned above, there are three known 4'PP-modified proteins in mammalian mitochondria,

NDUFAB1, AL1L2, and DHRS2[6]. To explore which of these putative targets displayed the

strongest genetic relationships to PPT, we analyzed gene co-dependency relationships from

the DepMap Public 23Q4+ Score Chronos dataset[15] to identify gene functionality within the

same biological pathway[16]. Among the three known mitochondrial 4'PP-modified proteins,

AASDHPPT gene effect was positively correlated only with *NDUFAB1* (Pearson r = 0.24)

(Extended Data Fig. 2A-C).

Unique among 4'PP-modified proteins, NDUFAB1 is a structural subunit of complex I of

the electron transport chain (ETC), as well as an integral part of the mitochondrial fatty acid

synthesis (mtFAS) pathway. mtFAS builds *de novo* fatty acyl chains on NDUFAB1, which are

required both as precursors for the biosynthesis of lipoic acid (LA), an important TCA cycle

enzyme co-factor, as well as to stabilize ETC assembly factors and iron-sulfur (FeS) cluster

biogenesis complexes via protein-protein interactions with NDUFAB1[12,13]. Because the newly

synthesized fatty acids are added to the terminal sulfhydryl group of the 4'PP PTM, the

addition of a 4'PP to NDUFAB1 is absolutely necessary for mtFAS to occur and to support

these downstream pathways[9]. Of note, the mtFAS pathway gene *OXSM* ranked third overall

for co-dependency with *AASDHPPT* (Pearson r = 0.42)[17] out of all genes in the Depmap

database (Extended Data Fig. 2D). Additional mtFAS pathway genes also ranked highly in

terms of *AASDHPPT* co-dependency, including *MCAT* and *MECR* (Pearson r = 0.36 and 0.32

respectively) (Extended Data Fig. 2E and F). Given that the 4'-PP modification is necessary

for acyl synthesis on NDUFAB1, these data suggested that impaired mitochondrial respiration

and TCA cycle metabolism in PPT-deficient cells is primarily attributable to altered mtFAS

function.

To further explore the genetic relationships between *AASDHPPT* and mtFAS, we

performed clustering and two-dimensional mapping of pairwise gene essentiality score

correlations with *AASDHPPT*. We included genes associated with the GO term "pantothenate

metabolism" from the DepMap project database as well as all genes encoding known

phosphopantetheinylated proteins, mtFAS pathway genes, and genes involved in downstream

lipoic acid synthesis and ETC assembly - such as the <u>l</u>eucine, <u>t</u>yrosine, <u>a</u>rginine <u>m</u>otif (LYRM)

assembly factors which are regulated by mtFAS[13]. The analyzed genes grouped into four

distinct clusters based on gene essentiality (Extended Data Fig. 2G). Two-dimensional

mapping of correlation distance showed that *AASDHPPT* most strongly correlated with mtFAS

pathway genes, as well as several LYRM family members, further suggesting that PPT may

act as a critical regulator of mtFAS functional activity (Fig. 2A).

To test this hypothesis, we examined whether PPT-deficiency altered known mtFAS

pathway endpoints, including protein lipoylation and succinate dehydrogenase subunit B

(SDHB) stability, which is a reliable surrogate of the LYRM-dependent ETC assembly

mechanism we previously described[13]. PPT-deficient cells exhibited reduced lipoylation of

pyruvate dehydrogenase (PDH) and α-ketoglutarate dehydrogenase (OGDH) subunits (DLAT

and DLST, respectively) along with reduced stability of SDHB (Fig. 2B), resembling the

molecular phenotype of mtFAS-deficient cells.

Although one prior study demonstrated that PPT is able to add a 4'PP group to NDUFAB1

in vitro[8], its requirement for NDUFAB1 phosphopantetheinylation had never been tested in

any *in vivo* setting. We monitored 4'PP-modification of NDUFAB1 by immunoblot, as

unmodified apo-NDUFAB1 has a lower molecular weight and thus migrates faster than 4'PP-

modified holo-NDUFAB1[8]. As hypothesized, we found that apo-NDUFAB1 accumulated in

PPT-deficient cells while holo-NDUFAB1 was depleted, indicated by a downward shift on the

blot (Fig. 2B). PDH and OGDH lipoylation, SDHB stability, and detection of holo-NDUFAB1

were all rescued by transient re-expression of full length AASDHPPT with a C-terminal GFP-

Flag tag (PPT-GFP) compared to a control vector expressing mitochondrially localized DSRed

(mtDSRed) (Fig. 2C). Additionally, basal respiration and spare respiratory capacity were also

both restored by re-expression of PPT-GFP in both PPT-deficient cell lines (Fig. 2D). From

these data, we conclude that PPT is required for 4'-PP modification of NDUFAB1 and therefore

for downstream mtFAS activity, protein lipoylation, ETC assembly, and mitochondrial

respiration[11].

The question then became: if PPT is required for modification of mitochondrial proteins,

does it act in the cytoplasm, prior to mitochondrial import, or is there an undescribed

mitochondrial pool of PPT? To address this question, we performed immunofluorescence

staining for endogenous PPT and mitochondrial ATP synthase subunit ATP5a and assessed

co-localization (Fig. 3A, Extended Data Fig. 3A). As expected, PPT expression was

significantly reduced in PPT-deficient cells compared to controls, and exhibited a punctate,

perinuclear expression pattern that co-localized with ATP5a staining in control cells (Pearson

r = 0.60 and 0.62) (Fig. 3B, Extended Data Fig. 3B). Additionally, PPT expression was

surprisingly observed in the nucleus of some, but not all, cells; a finding that we did not pursue

further in the present study, but which raises intriguing questions for future work.

We further verified the observed mitochondrial localization using subcellular fractionation

and Western blotting in control and PPT-deficient cells. We detected endogenous PPT

expression in both mitochondria and post-mitochondrial supernatant (PMS)(Fig. 3C). PPT

expression was reduced in both fractions, and near-completely absent from mitochondria in

PPT-deficient cells (Fig. 3C). We also characterized subcellular localization of PPT-GFP, and

found strong signal overlap with ATP5a, in agreement with our endogenous immunostaining

observations (Fig. 3D).

We then asked whether this pool of mitochondrially-localized PPT was mitochondrially

associated or inside the mitochondrial matrix, where NDUFAB1 and the rest of the mtFAS

machinery reside. To tackle this question, we employed a bimolecular fluorescence

complementation (BiFC) system[21] to visualize the sub-mitochondrial localization of PPT. We

used a split-Venus tag system in which the N-terminus of the Venus fluorophore (VN, residues

1-172) was attached to either the N- or C-terminus of PPT, while the C-terminus of Venus (VC,

residues 155-238) was attached to the C-terminus of a control protein with known localization

(MFN2 or CS) or NDUFAB1. This system allows us to observe fluorescent signal only when

both halves of the split-Venus tag are able to associate within a compartment[21]. We quantified

Venus fluorescence in each of our BiFC conditions by flow cytometry, and found that in cells

expressing C-terminally tagged PPT (PPT-VN), significant Venus signal was only observed

when co-expressed with either CS-VC or NDUFAB1-VC (Fig. 3E-G, see Extended data Fig.

3C for gating strategy), indicating that PPT-VN does in fact localize to the mitochondrial matrix.

Most mitochondrial proteins are encoded by nuclear genes and are targeted to and

imported into mitochondria via an N-terminal mitochondrial targeting sequence (MTS),

consisting of approximately 20 to 40 amino acid residues[22]. Surprisingly, in contrast, cells

expressing N-terminally tagged PPT (VN-PPT), positive Venus fluorescence was only

detected when co-expressed with MFN2-VC (Fig. 3E-G), indicating that N-terminally tagged

VN-PPT is mitochondrially associated on the OMM. Because our BiFC data showed that

tagging PPT on the N-terminus prevented its localization in the mitochondrial matrix (Fig. 3E-

G), we hypothesized that mitochondrial localization of PPT may be driven by a N-terminal

MTS. To test this, we generated N-terminal truncations of PPT to the next two methionine

residues at positions 14 and 37 (PPT$_{\Delta1-13}$-GFP and PPT$_{\Delta1-36}$-GFP, respectively) and examined

their localization using fluorescence microscopy.

While PPT-GFP co-localized strongly with ATP5a, PPT$_{\Delta1-13}$-GFP exhibited diffuse

localization throughout the nucleus and cytoplasm (Fig. 4A-B). PPT$_{\Delta1-36}$-GFP exhibited weak

fluorescent expression mostly within the nucleus, suggesting that this larger N-terminal

truncation destabilizes PPT (Fig. 4A-B). Sub-cellular fractionation and Western blotting

supported our microscopy results, showing that while PPT-GFP and PPT$_{\Delta1-13}$-GFP are both

present in post-mitochondrial supernatant (PMS) and mitochondrial fractions, PPT$_{\Delta1-13}$-GFP

accumulates in the PMS (Fig. 4C), suggesting difficulty in localizing to mitochondria. We also

performed the converse experiment, in which we added only the first 36 amino acids of PPT

to GFP, to test the sufficiency of the putative MTS to drive GFP to the mitochondria

(PPT_MTS-GFP). PPT_MTS-GFP showed strong overlap with ATP5a staining by

fluorescence microscopy (Extended data Fig. 4A), as well as strong expression in the

mitochondrial fraction via Western blot (Extended data Fig. 4B), indicating that the first 36

amino acids of PPT comprise a bona fide MTS.

To test whether this newfound mitochondrial localization is required for PPT to modify

mitochondrial target proteins, we transduced PPT-deficient cells with mtDSRed, PPT-GFP,

PPT$_{\Delta1-13}$-GFP, or PPT$_{\Delta1-36}$-GFP, and assessed mtFAS pathway activity. Expression of PPT-

GFP rescued protein lipoylation and SDHB stability, whereas PPT$_{\Delta1-13}$-GFP and PPT$_{\Delta1-36}$-GFP

both failed to rescue either of these mtFAS-dependent endpoints (Fig. 4D). These data

demonstrate that mitochondrial localization of PPT is required for 4'PP-modification of

NDUFAB1. To further investigate whether mitochondrial localization of PPT is required for

OXPHOS activity, we performed growth assays as before in glucose- and galactose-

containing media, with cells PPT-deficient cells expressing mtDSRed, PPT-GFP, or PPT$_{\Delta1-13}$-

GFP. We found that growth in galactose, which requires functional mitochondrial respiration,

was partially rescued by expression of PPT-GFP, and that PPT$_{\Delta 1\text{-}13}$-GFP rescued less well

than the full length PPT-GFP construct (Extended Data Fig. 4C-F). Together, these data

demonstrate that mitochondrial localization of PPT is necessary for phosphopantetheinylation

of NDUFAB1, mtFAS pathway activity, and respiration-dependent growth.

We recently described a pathogenic variant in the mtFAS gene *MCAT* c.812T>C; p.T271I

(rs760294168; NM_173467.4)[23]. As our data demonstrate that perturbed *Aasdhppt*

expression results in similar phenotypes as mtFAS loss, we wondered whether there may be

patients with pathological variants in *AASDHPPT*. We searched the ClinVar database[24] and

identified several missense variants of uncertain significance (VUS) in *AASDHPPT* that were

associated with suspected inborn genetic disease. We selected six of these variants for

analysis: c.343C>T; p.P12S (NM_015423.3), c.140G>A; pR47H (NM_015423.3), c.214G>T;

pA72T (NM_015423.3), c.235T>G; pW79G (NM_015423.3), c.247C>T; pR83C

(NM_015423.3), and c.286A>G; pK96E (NM_015423.3).

We created each of these missense mutations in our full length PPT-GFP construct and

assessed their ability to rescue perturbed mtFAS activity in our PPT-deficient cells. We found

that PPT-GFP R47H and PPT-GFP W79G exhibited weaker rescue effects compared to the

other variants (Fig. 4E, Extended Data Fig. 4G). In addition, cells expressing the PPT-GFP

W79G variant showed low levels of protein expression compared to the other variants, despite

expression from the same promoter (EF-1α promoter), suggesting that this variant may have

reduced protein stability. These data identify R47H and W79G as potential clinically observed

pathologic variants of *AASDHPPT*, with molecular phenotypes attributable to altered mtFAS

activity and perturbed cellular respiration.

Our data position PPT as a critical mediator of mitochondrial oxidative function, due to its

requirement for proper function of the mtFAS pathway. The discovery of distinct pools of PPT

raises questions about what factors control PPT activity in each compartment. For instance,

PPT uses CoA as the substrate for 4'PP-modification of target proteins. How perturbations in

CoA availability affect PPT enzymatic activity may have important implications for PPT targets

such as mtFAS in diseases with CoA dysregulation[15]. Moreover, our data demonstrate that

PPT is targeted to mitochondria via a N-terminal MTS, but how this localization is controlled

in cells to produce two distinct pools is completely unstudied. Whether PPT localization is

dynamic will also be an interesting and important area for future study. Of note, we identified

two probable pathogenic patient variants in *AASDHPPT*. Our results predict that patients

harboring these mutant alleles likely exhibit impaired mtFAS, leading to diminished

mitochondrial function. These data add to the growing list of patients with pathogenic variants

in mtFAS and lipoic acid synthesis genes and support the notion that impairments in CoA

metabolism may cause disease via mtFAS dysfunction[16,17].

Methods

Aasdhppt mutant cell lines, growth assays, and viral transduction

C2C12 immortalized mouse skeletal myoblasts (ATCC CRL-1772, verification provided by

ATCC) were grown in DMEM supplemented with 4.5 g/L glucose, glutamine, and sodium

pyruvate (Corning, 10-013-CV) and 10% FBS (Sigma, F0926) in humidified tissue culture

incubators at 37°C, 5% CO_2. To generate *Aasdhppt* mutant clones, four sgRNA sequences

(**Table 1)** were designed targeting exon 1 or exon 2 of the mouse *Aasdhppt* gene and

subcloned into the pLentiCRISPR v2 vector using previously described methods[18]. Parental

C2C12s were transfected with pLentiCRISPRv2_sgAasdhppt or empty vector. 48 hours after

transfection, cells were sorted on GFP signal intensity and plated into 96-well plates to obtain

single cell colonies for expansion and establishment of stable, clonal cell lines. Clones were

screened for AASDHPPT protein expression levels and two control clones and two *Aasdhppt*

mutant clones were selected for further experiments. For growth assays 10,000 cells/well

were plated in 12-well plates or 1,000 cells/well in 48-well plates and grown in DMEM with

either no sodium pyruvate, 4 mM glutamine, and 25 mM glucose or no sodium pyruvate, 4

mM glutamine, and 10 mM galactose for 3 days in technical triplicates. Tissue culture plates

were placed in an IncuCyte® chamber (Sartorius) and percent confluency was measured from

sixteen, 10X magnification fields per well for 12-well plates, or nine, 10X magnification fields

for 48-well plates every 6 hours. For co-localization and rescue experiments, control and

Aasdhppt mutant clonal lines were transiently infected with retrovirus harboring a control

transfer plasmid expressing mtDSRed (pQXCIP mtDSRed) or pMMLV retrovirus vectors

custom designed, synthesized, and purchased from VectorBuilder (Chicago, IL, USA),

harboring full length, or N-terminal truncations of residues 1-13 or 1-36, of AASDHPPT with

C-terminal GFP and Flag tag- fusion. For generation of retrovirus constructs harboring single

point mutations in the full length AASDHPPT-GFP-Flag construct, primers in **Table 2** were

used with a New England Biolabs Q5 Site-Directed Mutagenesis Kit. Plasmid reactions were

transformed into competent *E. coli* and selected colonies were sequence validated for

successful incorporation of point mutations.

Genomic Sequence Analysis of AASDHPPT Mutants

Approximately five million cells of each clonal mutant cell line, PPT-1 and PPT-2, were pelleted and washed with PBS then frozen at -80ºC. Genomic DNA was extracted using a column-based kit assay (Qiagen 69504). Aasdhppt was amplified using primers described in **Table 3**, and sanger sequenced using the forward primer. Sanger sequencing peaks were deconvoluted using the Indigo web-based software (https://www.gear-genomics.com)[19].

Crude mitochondrial isolation

Cells were harvested and processed as previously described[13]. Briefly, cells were harvested by trypsinization, washed once with cold, sterile PBS (Gibco, 10010-023), then stored at -80°C. At the time of mitochondrial isolation, cell pellets were thawed and resuspended in 1 mL CP-1 buffer (100 mM KCl, 50 mM Tris-HCL, 2 mM EGTA, pH 7.4) supplemented with mammalian protease inhibitor cocktail (mPIC, Millipore Sigma P8340) and mechanically lysed by 10 passes through a 1 mL insulin syringe with 26-guage needle and centrifuged at 700 x g for 10 minutes to pellet unlysed cells and debris. Supernatant was transferred to a new microcentrifuge tube and centrifuged at 10,000 x g for 10 minutes to pellet the crude mitochondrial fraction. Post-mitochondrial supernatant (PMS) was used for Western blot analysis or discarded. Mitochondrial pellets were resuspended in a small volume of either RIPA buffer (10 mM Tris-HCl, pH 8.0, 1 mM EDTA, 0.5 mM EGTA, 1% Triton X-100, 0.1% Sodium Deoxycholate, 0.1% SDS, 140 mM NaCl) supplemented with mPIC, or CP-1 buffer, equal to approximately twice the pellet volume. Resuspended pellets were then used for applications described below.

SDS-PAGE and immunoblotting

Cultured cells were scraped from tissue culture plates directly into RIPA buffer supplemented with mammalian protease inhibitor cocktail (mPIC, Millipore Sigma) and Pierce Universal Nuclease, incubated at 4°C with constant agitation on a microcentrifuge tube orbital shaker, then centrifuged at 17,000 x g for 20 minutes at 4°C to remove insoluble material. Supernatant was then transferred to a new microcentrifuge tube to be saved as whole cell lysate (WCL). For protein analysis, WCL, PMS, and/or crude mitochondrial fractions were first normalized for total protein content via Pierce BCA Protein assay (Thermo Fisher Scientifc 23225). Samples were resolved by SDS-PAGE and transferred to nitrocellulose membranes. Immunoblotting was performed using the indicated primary antibodies (**Table 4**) with

visualization and analysis of protein signal using Rabbit and Mouse IgG Antibody DyLight 680,

or 800, Conjugated secondary antibodies and Bio-Rad ChemiDoc Imaging System.

Seahorse Extracellular Flux Rate Assay

Extracellular flux rate assays were performed on a Seahorse XFe96 Analyzer (Agilent).

Cells from the indicated clones were plated in approximately 22 wells of a 96-well seahorse

plate at a density of 10,000 cells/well in DMEM +10% FBS and allowed to adhere overnight.

The morning of the assay, cells were washed twice and media was replaced with non-buffered

seahorse media (Agilent 103680) supplemented with 25 mM glucose, 2 mM glutamine, and 1

mM sodium pyruvate. A standard mitochondrial stress test was then performed using 1 μM

oligomycin, 3 μM FCCP, and 0.5 μM Rotenone + 0.5 μM Antimycin A. Baseline oxygen

consumption was measured three times, followed by acute injection of oligomycin, followed

by FCCP and finally Rotenone + Antimycin A. Measurements were standard, with acute

injection of drug followed by 3 minutes of mixing then measurements of oxygen levels over 3

minutes, with three measurements per phase. Data were normalized with crystal violet

absorption for approximation of cell number. Results were analyzed in WAVE software.

Blue Native-PAGE

Crude mitochondrial fractions were prepared and normalized for total protein content as

previously described. 100 μg of mitochondria were pelleted at 10,000 x g for 10 minutes at

4°C. Supernatant was removed and pellets were resuspended in 60 μL of 1x pink lysis buffer

(Invitrogen BN2003). Digitonin (GoldBio D-180-2.5) was added to samples at a final

concentration of 1% mass/volume and samples were incubated on ice for 15 min. Insoluble

material was pelleted by centrifugation at 17,000 x g for 20 minutes and supernatant was

transferred to new microcentrifuge tubes. 6 μL of NativePAGE sample buffer (Invitrogen

BN2004) was added to samples and 10 μL of sample (~15 μg of crude mitochondrial fraction)

was loaded per lane on pre-cast 3-12%, Bis-Tris NativePAGE gels (Invitrogen BN1001BOX)

with NativePAGE anode buffer (Invitrogen 2001) and dark blue cathode buffer (Invitrogen

BN2002) at 150V for 30-40 minutes then switched to light blue cathode buffer and run at 20

V overnight. Gels were then transferred to PVDF membranes at 100 V for 2 hours and blotted

with the indicated primary antibodies (**Table 4**. Secondary anti-mouse HRP antibody and

SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific 34096)

was used to visualize bands with a Bio-Rad ChemiDoc Imaging System.

Glutamine Tracing

The indicated clonal cell lines were plated in 6-well plates in triplicate at a density of 200K

for PPT-1 and PPT-2, and 175K for Ctrl-1 and Ctrl-2 to account for differences in growth rate.

Cells were allowed to adhere for two hours, at which point the media was changed to

unlabeled media (Corning DMEM 17-207-CV with 1mM sodium pyruvate, 5mM glucose, and

4mM glutamine). After twenty hours in unlabeled media, wells were changed to labeled media

at the indicated time points prior to harvest (Corning DMEM 17-207-CV with 1mM sodium

pyruvate, 5mM glucose, and 4mM U^{13}C-glutamine). Three wells per cell line were left

unlabeled for natural isotope abundance correction, and another three per cell line were used

to determine cell number at the time of freezing. 24 hours after cells were in unlabeled media,

all wells were washed with normal saline once and frozen at -80°C. Plates were then extracted

for metabolites using a volume of 40:20:20 methanol:acetonitrile:water scaled to the number

of cells determined by the cell count wells (average from three wells). Lysates were sonicated

and incubated on ice for 60 minutes. At this point any insoluble material was spun out of the

sample at 17,000 x g for 10 minutes at 4°C. 1000 µL of supernatant was dried using a

SpeedVac concentrator. Dried metabolomics extracts were next resuspended in 100% water

containing 25 µg/mL D5-Glutamate (DLM-556, Cambridge) such that resuspension sample

equivalents were 5.82E+3 cells/µL. 2 µL of resuspended samples were injected on column, a

process blank was analyzed before and after experimental samples, and a pooled sample

was injected twice before experimental samples for column conditioning, and every 7-10

sample injections thereafter. Data dependent MS2 (ddMS2) data was collected from pooled

experimental replicates to identify group-specific compounds.

Data were collected on a Vanquish liquid chromatography system coupled to an Orbitrap

Exploris 240 (Thermo Fisher Scientific) using a heated electrospray ionization (H-ESI) source

in ESI negative mode. All samples were run through a 24-minute reversed-phase

chromatography ZORBAX extend-C18 column (1.8 µm, 2.1mm × 150mm, 759700-902,

Agilent, California, USA) combined with a guard column (1.8 µm, 2.1 mm × 5 mm, 821725-

907, Agilent). Mobile phase A was LC/MS H2O (W6-4, Fisher) with 3% LC/MS grade MeOH

(A456-4, Fisher), mobile phase B was LC/MS grade methanol (A456-4, Fisher), and both

mobile phases contained 10 mM tributylamine (90780-100ML, Sigma), 15 mM acetic acid,

and 0.01% medronic acid (v/v, 5191-4506, Agilent). For the re-equilibration gradient, mobile

phase A was kept the same, and mobile phase B was 99% LC/MS grade acetonitrile (A955-

4, Fisher). Column temperature was kept at 35ºC, flow rate 0.25 mL/min, and the solvent

gradient was as follows: 0-2.5 min held at 0% B, 2.5-7.5 min from 0% B to 20% B, 7.5-13 min

from 20% B to 45% B, 13-20 min from 45% B to 99% B, and 20-24 min held at 99% B. The

analytical solvent gradient was followed by a 16 min re-equilibration gradient to prep the

column before the next sample injection that went as follows: 0-0.05 min held at 99% B at

0.25 mL/min, 0.05-1 min from 99% B to 50% B and 0.25 mL/min to 0.1 mL/min, 1-11 min held

at 50% B and 0.1 mL/min, 11-11.05 min from 50% B to 0% B at 0.1 mL/min, 11.05-14 min held

at 0% B at 0.1 mL/min, 14-14.05 min held at 0% B and increased flow rate from 0.1 mL/min

to 0.25 mL/min, and 14.05-16 min held at 0% B and 0.25 mL/min. The mass spectrometer

acquisition settings were as follows: source voltage -2,500V, sheath gas 60, aux gas 19,

sweep gas 1, ion transfer tube temperature 320ºC, and vaporizer temperature 250ºC. Full

scan data were collected with a scan range of 70-800 m/z at a mass resolution of 240,000.

Fragmentation data was collected using a data-dependent MS2 (ddMS2) acquisition method

with MS1 mass resolution at 120,000, MS2 mass resolution at 15,000, and HCD collision

energy fixed at 30%.

Natural isotope abundance correction was done using the FluxFix Isotopologue Analysis

Tool*14* (version 0.1) web-based application[20]. Data analysis was conducted in Skyline (version

23.1.0.268), consisting of peak picking and integration, using an in-house curated compound

database of accurate mass MS1 and retention time derived from analytical standards and/or

MS2 spectral matches for the specified chromatography method.

AASDHPPT co-essentiality analysis and network modeling

Gene essentiality scores and gene co-dependency Pearson correlation values were

obtained from the DepMap Portal Project Achilles 23Q4 release[9,21–23] in which genome-wide

CRISPR loss-of-function screens were performed in 1100 human cancer cell lines. Co-

Dependency plots between AASDHPPT and other genes were generated from gene

essentiality scores for each cancer line obtained directly from the DepMap Portal. For two-

dimensional network analysis, gene essentiality correlations for a custom list of genes were

used to create a data matrix (see Supplementary Excel File**)**. From this matrix, methodologies

and code originally published by Arnold et al.[24] and available at: https://github.com/finley-

lab/coessentiality-network were first used to generate a correlation matrix heatmap of

codependent gene modules. Co-dependent gene modules were then visualized as a network

diagram and genes with low correlation scores (Pearson r < 0.2) were filtered out as previously

described[24]. Graph edges were weighted according to the strength of pairwise gene

correlations.

Cell immunostaining and microscopy analysis

Cell immunostaining and microscopy analysis

Cells were seeded into chambered cover glass slides and cultured overnight. The

following day, cells were fixed in ice-cold 4% paraformaldehyde (PFA) in PBS solution for 10

minutes at room temperature, followed by 3 washes in 1X PBS solution. Cells were then

permeabilized and blocked in PBS solution containing 0.5% BSA, 5% goat serum, and 0.3%

Triton X-100 for 1 hour at room temperature. Cells permeabilized/blocked with blocking buffer

were then incubated with denoted primary antibodies overnight at 4°C followed by washes

with 1X PBS prior to incubation with Alexa Fluor conjugated secondary antibodies (**Table 4**)

for 2 hours at room temperature. Washes were repeated and cells were then stained with

Hoechst Dye for 10 minutes at room temperature. After subsequent washes, samples were

then imaged on a Zeiss LSM 880 with Airyscan and non-linear optics microscope, using a 40X

water objective and 1.2 NA to collect z-stacks.

For quantification of endogenous AASDHPPT expression in control and *Aasdhppt* mutant

clonal lines, regions of interest (ROIs) were manually drawn with Fiji software for individual

cells from channels with fluorescent ATP5a expression. Area was then measured for each

ROI with corresponding mean fluorescent intensity of signal from the AASDHPPT fluorescent

channel. Mean fluorescent intensity per cell area was then calculated. For statistical analysis,

GraphPad Prism v10 softwared was used to remove outliers from the data set by the ROUT

method, Q = 1%.

For colocalization analysis of endogenous AASDHPPT or AASDHPPT-GFP fusion

constructs with mitochondrial proteins in maximum intensity projections of each imaged field,

nuclear signal detected by Hoechst Dye staining was first used to create ROIs in Fiji. Pixels

with signal detected in nuclear ROIs in red and green channels were then removed from

images of the same field prior to merging of red and green channels so that colocalization

analysis would be exclusive of nuclei and limited to subcellular organelles and cytoplasmic

regions. Merged channels were then processed by the BIOP JACoP Fiji plugin

(https://c4science.ch/w/bioimaging_and_optics_platform_biop/image-

processing/imagej_tools/jacop_b/) using the Otsu method of thresholding for each channel to

generate Pearsons Correlation values.

Bimolecular fluorescence complementation (BiFC) assay

To generate BiFC cell lines, five split-Venus constructs were designed and synthesized by

VectorBuilder in a pMMLV retroviral backbone. Constructs were designed to contain either the

C' terminal half of the Venus tag with blasticidin resistance (VC), or the N' terminal half of the

Venus tag with puromycin resistance (VN). VC constructs were tagged to the C' terminal end of *citrate synthase* (CS-VC), *mitofusin 2* (MFN2-VC) or *Ndufab1* (NDUFAB1-VC). VN constructs were tagged to either the C' or N' terminal end of *Aasdhppt* (PPT-VN and VN-PPT, respectively). Following retroviral packaging in HEK293T cells, 40,000 wildtype C2C12 cells were transduced to express either PPT-VN or VN-PPT in combination with each VC construct. Negative control lines were generated by transducing cells with individual VC or VN constructs alongside an empty vector control (PQXCIP for VC constructs, and PQXCIB for VN constructs). This ensured that all cells were transduced with an approximately equal total viral load. Stable populations were derived via combined puromycin and blasticidin selection (2 µg/mL each), yielding 11 cell lines in total; 6 with combined VC/VN constructs plus 5 single construct controls.

For measurement of fluorescence, approximately 100,000 cells were trypsinized and resuspended in 200 µL PBS. Cell suspensions were then assessment on an Accuri bench-top cytometer, recording a minimum of 20,000 events per sample. The percentage of Venus-positive cells was analyzed using FlowJo (v. 10.10.0). Fluorescence intensity histograms were generated using FlowJo Layout Editor. A representative flow cytometry gating strategy is shown in Extended data Fig. 3C. Visual confirmation of fluorescence achieved by imaging all BiFC cell lines on a Nikon Eclipse Ti2 microscope. Briefly, 20,000 cells per cell line were seeded in chamber cover slides (Cellvis, C4-1.5H-N). After 24 hours, cells were fixed in 4% PFA for 30 mins at room temperature before imaging the following day (20X objective, 0.75 NA).

Statistics

Statistical analysis was performed using GraphPad Prism 10. Unless otherwise noted, data were analyzed by one-way ANOVA followed by two-sided Dunnett's multiple comparison test (when compared to only control) or Šídák's or Tukey's multiple comparison test (when comparing all groups). A p-value of <0.05 was considered statistically significant. Data that included a third vairable (ex. time) was analyzed by two-way ANOVA followed by Tukey's multiple comparison test.

Table 1

sgRNA cloned into pLentiCRISPRv2	sequence
PPT 1	GATGCGGTCTATCCAGCCGG
PPT 2	GCAGACGGATATGATCCCAA

PPT 3	AGCGGACCGAGCCGGTCGTA
PPT 4	GGAACACCATACGACCGGCT

Table 2

Primers for site-directed mutagenesis	sequence
P12S Fwd	GTGCGTGGTGAGCTCCATGGAGG
P12S Rev	AGCCTCTTAGCGGGG
R47H Fwd	GGAGAAGGAGCATATCGGCAAGTTC
R47H Rev	TCCGGCTGGATAGAC
A72T Fwd	GAAATTAGTTACCGAGAAATTGAATATCCC
A72T Rev	CTTATCATCAAACGACCAG
W79G Fwd	GAATATCCCTGGCGATCATATCCGTC
W79G Rev	AATTTCTCTGCAACTAATTTC
R83C Fwd	GGATCATATCTGCCTGCAGAGAAC
R83C Rev	CAAGGGATATTCAATTTCTC
K96E Fwd	GGTTCTTGCGGAAGACTCGTTGA
K96E Rev	GGCTTTCCCTTTGAAGTTCTC

Table 3

Primers for Genomic Amplification of *Aasdhppt*	
Forward Primer	TACCTGGCTTGAGGATCAGC
Reverse Primer	GGTTCTGCAGCAAGAACTGCAT

Table 4

Antibodies	Source and reference number
AASDHPPT	Proteintech 11244-1-AP
β-actin	Proteintech 66009-1-Ig
GRIM19	Abcam ab110240
SDHA	Abcam ab14715
UQCRFS1	Abcam ab14746

MTCO1	Abcam ab110270
ATP5a	Abcam ab14748
Lipoic acid	Millipore 437695
SDHB	Abcam ab14714
NDUFAB1	Abcam ab96230 and ab181021
GFP	Abcam ab290
Goat anti-Mouse IgG (H+L) AlexaFluor 568	Thermo Fisher A-11004
Goat anti-Rabbit IgG (H+L) AlexaFluor 568	Thermo Fisher A-11011
Goat anti-Mouse IgG (H+L), HRP	Thermo Fisher A28177
Goat anti-Mouse IgG (H+L) Dylight 800	Rockland Immunochemicals 610-145-002-0.5
Goat anti-Rabbit IgG (H+L) Dylight 800	Rockland Immunochemicals 611-145-002-0.5
Goat anti-Mouse IgG (H+L) AlexaFluor 680	Thermo Fisher A-21057
Goat anti-Rabbit IgG (H+L) AlexaFluor 680	Thermo Fisher A10043

Acknowledgments

This work was supported by funding from the NIH (R35GM151245 to SMN) and the Van Andel Institute – Metabolism & Nutrition (MeNu) Program. Stipend support for RJW and OAP was provided by the Van Andel Institute Graduate School (VAIGS). RW was also supported by NIH F30GM154476. We thank the Meijer Foundation for financial support of MC and EM. We also thank the Van Andel Institute's Flow Cytometry Core (RRID:SCR_022685) for their assistance with single cell sorting for generation of clonal cell lines, the Metabolomics and Mass Spectrometry Core (RRID:SCR_024903) for their assistance with labeled glutamine tracing experiments and analysis, the Optical Imaging Core (RRID:SCR_021968) for their assistance with airyscan microscopy, and the Bioinformatics and Biostatistics Core (RRID:SCR_024762) for their assistance with gene essentiality score correlation network analysis.

The authors declare no competing interests.

Data Availability

All data needed to evaluate the conclusions in the manuscript are present in the

manuscript and/or Supplementary Information. Additional data related to this paper may be

requested from the authors.

Figure Legends

Figure 1. AASDHPPT is required for mitochondrial oxidative metabolism.

(A) Whole cell lysates were immunoblotted for the denoted targets. Data are representative

of three biological replicates. (B-C) Cells were grown in either 25 mM glucose (B) or 10 mM

galactose (C) in an IncuCyte® system with images captured every 6 hours for 3 days. Data

represent the mean confluency; error bars denote SEM * = p <0.05. ns = no significance. Data

are representative of 3 biological replicates (D) Seahorse mitochondrial stress test for Oxygen

Consumption Rate (OCR). Data are representative of three independent experiments. Error

bars represent SD. (E) Blue-native PAGE separation of isolated mitochondrial protein

complexes from denoted cell lines, followed by immunoblot with the indicated antibodies. Data

are representative of three biological replicates. (F) Schematic of isotopomeric labeling

patterns upon substitution of $U^{13}C$-glutamine for unlabeled glutamine. Open circles indicate

^{12}C carbons, where circles with grey, red, or blue, indicate ^{13}C carbons. (G-H) Technical

triplicate samples (n=3) of the indicated cell lines cultured in $U^{13}C$-glutamine for the indicated

time points, harvested, and analyzed via LC-MS. Steady state metabolite pool sizes and ratios

are shown for the 4-hour labeling time point. (I) Ratio of Citrate M+5 to Citrate M+4 at the four-

hour time point. Error bars represent SEM (B, C), SD (D, G, H, I). * = p<0.05 ** = p<0.01, ***

= p<0.001, **** = p<0.0001 as determined by ANOVA followed by Tukey's multiple

comparisons test.

Figure 2. AASDHPPT is required for mitochondrial fatty acid synthesis pathway

function.

(A) Two-dimensional network diagram representing gene essentiality score correlations

between selected genes associated with *AASDHPPT* and known substrate targets, mtFAS

pathway members, LYRM family members, and the GO term "pantothenate metabolism" using

a minimum threshold correlation score > 0.2. Different colors represent genes associated with

clusters as shown in Extended Data Fig. 2G. Correlation strength is represented by the length

and thickness of the connecting edge. (B) Whole cell lysates immunoblotted for the denoted

targets. Data are representative of three biological replicates. # denotes non-specific bands.

Red asterisk denotes positive PPT expression band in clonal controls that is reduced in PPT-

deficient mutants. (C) Whole cell lysates from indicated cell lines expressing either mtDSRed

or PPT-GFP fusion protein were immunoblotted for protein expression of the denoted targets.

Data are representative of three biological replicates. (D) Seahorse standard mitochondrial

stress test to determine Oxygen Consumption Rate (OCR). Data are representative of three

independent experiments. Error bars represent SD.

Figure 3. AASDHPPT localizes to the mitochondrial matrix.

(A) Representative 40X HPF images of *Aasdhppt* mutant and control cell lines

immunostained with Hoechst dye to visualize nuclei and antibodies against AASDHPPT (red)

and ATP5a (green). Scale bars = 100μm. (B) Quantification of AASDHPPT and ATP5a signal

colocalization outside of the nucleus from n = 8 40X HPF per group assessed by Pearson's

correlation coefficient.(C) Subcellular fractionation of *Aasdhppt* mutant cell lines and control.

Post-mitochondrial supernatant (PMS) and mitochondrial lysate (Mito) was isolated from the

denoted cell lines and immunoblotted for AASDHPPT, ATP5a, and GAPDH. Data are

representative of three biological replicates. (D) Representative 40X HPF images of WT

skeletal muscle myoblasts expressing AASDHPPT-GFP (PPT-GFP) stained with Hoechst dye

and antibody against ATP5a (red). Scale bars represent 100 μm. Dashed outlines denote

magnified regions shown in panels with solid outlines. For all statistical analysis data represent

the mean; error bars denote SD. **** = p<0.0001 as determined by one-way ANOVA with

Šidák's multiple comparisons test. (E) Representative histograms of Venus (FITC)

fluorescence in the indicated BiFC cell lines. (F) Percentage of Venus-positive cells in the

indicated cell lines as measured by flow cytometry. Data represent the mean; error bars

denote standard deviation (n = 3 per group). Statistical significance was determined by one-

way ANOVA with Tukey's multiple comparison test. (G) Representative 20X images of

endogenous Venus fluorescence in VC + VN BiFC cell lines. Scale bars represent 50 μm.

Figure 4. Mitochondrial AASDHPPT is required for mtFAS function.

(A) Representative 40X HPF images of WT skeletal muscle myoblasts expressing

mtDSRed, PPT-GFP, 13 residue N-terminal truncation of PPT (PPT$_{\Delta1\text{-}13}$-GFP), or 36 residue

N-terminal truncation of PPT (PPT$_{\Delta1\text{-}36}$-GFP) constructs, stained with Hoechst dye and

antibody against ATP5a. Scale bars represent 100 μm. Dashed outlines denote magnified

regions shown in panels with solid outlines. Arrows denote weak expression of GFP signal

detected. (B) Quantification of PPT-GFP and ATP5a signal colocalization outside of the

nucleus from n = 8 40X HPF assessed by Pearson's correlation coefficient from experiment
shown in (A). Data represent the mean; error bars denote SD. **** = p<0.0001, determined
by one-way ANOVA with Dunnett's multiple comparisons test. (C) Subcellular fractionation of
WT cells expressing PPT-GFP, PPT$_{\Delta1\text{-}13}$-GFP, or PPT$_{\Delta1\text{-}36}$-GFP. Post-mitochondrial
supernatant (PMS) and mitochondrial lysate (Mito) was isolated from the denoted cell lines
and immunoblotted for the indicated targets. Blots are representative of three biological
replicates. (D) Whole cell lysates from cells expressing mtDSRed, PPT-GFP, PPT$_{\Delta1\text{-}13}$-GFP,
or PPT$_{\Delta1\text{-}36}$-GFP were immunoblotted for the denoted targets. Data are representative of n >
3 biological replicates. (E) Whole cell lysates from clonal control and *Aasdhppt*-deficient cells
transduced with retrovirus expressing mtDSRed, PPT-GFP P12S, PPT-GFP R47H, PPT-GFP
A72T, PPT-GFP W79G, PPT-GFP R83C, or PPT-GFP K96E were separated by SDS-PAGE
and immunoblotted for the denoted targets. Immunoblots are representative of three biological
replicates.

Extended Data Fig. 1. AASDHPPT is required for mitochondrial oxidative metabolism.
(A) Sequence analysis of sgRNA-targeted region in clonal Aasdhppt mutant cell lines, and
corresponding mutations generated. (B) Representative IncuCyte® phase contrast images
from cells denoted in Fig. 1B and C at 72 hours grown in glucose or galactose supplemented
DMEM. Scale bars represent 200 μm. (C-G) Pool size of the indicated metabolites from steady
state metabolomics analysis at the four-hour labeling time point. Data from n=3 technical
replicates of the indicated cell lines. Error bars represent SD. Statistical analysis was done
using a one-way ANOVA followed by Tukey's multiple comparisons test. (H-I) Fractional
enrichment of given isotopologues is represented as percentage of the total pool at each
indicated time point. Error bars represent SD. Significance was determined by two-way
ANOVA followed by Tukey's multiple comparisons test. ** = p <0.01. *** = p <0.001. **** = p
<0.0001. NS = not significant.

Extended Data Fig. 2. AASDHPPT is required for mitochondrial fatty acid synthesis
pathway function.
(A-F) Codependency plots of gene essentiality scores comparing *AASDHPPT* with
NDUFAB1 (A), *ALDH1L2* (B), *DHRS2* (C)*, OXSM* (D), *MCAT* (E), and *MECR* (F) from DepMap
CRISPR screens[9,21–23,25]. **(**G) Heatmap depicting hierarchical clustering of pairwise gene
essentiality score correlations of PPT, known PPT substrate targets, mtFAS pathway

members, LYRM family members, and genes associated with the GO term "pantothenate

metabolism."

Extended Data Fig. 3. AASDHPPT localizes to the mitochondrial matrix

(A) Representative 40X HPF images of second clones of *Aasdhppt* mutants and controls

stained with Hoechst dye to visualize nuclei and antibodies against AASDHPPT (red) and

ATP5a (green). Scale bars = 100μm. (B) Quantification of mean fluorescent intensity of

AASDHPPT expression per cell area (μm^2) in *Aasdhppt* mutants (PPT-1 n = 170 and PPT-2

n =159) and control clonal cell lines (Ctrl-1 n = 186 and Ctrl-2 n = 118) as represented in (A

and Fig. 3A).

Extended Data Fig. 4. Mitochondrial localization of AASDHPPT is required for mtFAS

function

(A) Representative 40X images of cells expressing PPT_MTS-GFP stained with Hoechst

dye and antibody against ATP5a. Scale bars represent 20 μm. (B) Subcellular fractionation of

cells expressing PPT_MTS-GFP. Post-mitochondrial supernatant (PMS) and mitochondrial

lysate (Mito) was isolated and immunoblotted for the indicated targets. Blots are

representative of three biological replicates. (C,E) Cells were seeded in either 25 mM glucose

or 10 mM galactose and grown in an IncuCyte® system with images captured every 6 hours

for 3 days. Percent confluency measured by IncuCyte® software. Statistical analysis for

measurements at 72 hrs was performed with one-way ANOVA and Tukey's multiple

comparisons test. * = p <0.05. *** = p <0.001. **** = p <0.0001. Data presented are mean ±

SEM from one experiment representative of three biological replicates. (D,F) Representative

IncuCyte® phase images from cells denoted in C and E at 72 hours grown in glucose or

galactose supplemented DMEM. Scale bars represent 200 μm. For all statistical analysis data

represent the mean; error bars denote SD. * = p <0.05. ** = p <.01. **** = p<0.0001, ns = not

significant as determined by one-way ANOVA with Dunnett's multiple comparisons test. (G)

Whole cell lysates from second clones of control and *Aasdhppt*-deficient cells transduced with

retrovirus expressing mtDSRed, PPT-GFP P12S, PPT-GFP R47H, PPT-GFP A72T, PPT-GFP

W79G, PPT-GFP R83C, or PPT-GFP K96E were separated by SDS-PAGE and

immunoblotted for the denoted targets. Immunoblots are representative of three biological

replicates.

666
667

References

1.   Chen, N., Liu, Y., Li, Y. & Wang, C. Chemical Proteomic Profiling of Protein 4'-

Phosphopantetheinylation in Mammalian Cells. *Angew. Chem. Int. Ed Engl.* **59**, 16069–

16075 (2020).

2.   Joshi, A. K., Zhang, L., Rangan, V. S. & Smith, S. Cloning, Expression, and

Characterization of a Human 4′-Phosphopantetheinyl Transferase with Broad Substrate

Specificity*. *J. Biol. Chem.* **278**, 33142–33149 (2003).

3.   Mullen, A. R. *et al.* Reductive carboxylation supports growth in tumour cells with defective

mitochondria. *Nature* **481**, 385–388 (2011).

4.   Tsherniak, A. *et al.* Defining a cancer dependency map. *Cell* **170**, 564-576.e16 (2017).

5.   Wainberg, M. *et al.* A genome-wide atlas of co-essential modules assigns function to

uncharacterized genes. *Nat. Genet.* **53**, 638–649 (2021).

6.   Solmonson, A. & DeBerardinis, R. J. Lipoic acid metabolism and mitochondrial redox

regulation. *J. Biol. Chem.* **293**, 7522–7530 (2018).

7.   Nowinski, S. M. *et al.* Mitochondrial fatty acid synthesis coordinates oxidative metabolism

in mammalian mitochondria. *Elife* **9**, e58041 (2020).

8.   Van Vranken, J. G. *et al.* ACP Acylation Is an Acetyl-CoA-Dependent Modification

Required for Electron Transport Chain Assembly. *Mol. Cell* **71**, 567-580.e4 (2018).

9.   DepMap, B. DepMap 23Q4 Public. Figshare+

https://doi.org/10.25452/FIGSHARE.PLUS.24667905.V2 (2023).

10. Wedan, R. J., Longenecker, J. Z. & Nowinski, S. M. Mitochondrial fatty acid synthesis is

an emergent central regulator of mammalian oxidative metabolism. *Cell Metab.* **36**, 36–

47 (2024).

11. Kerppola, T. K. Design and implementation of bimolecular fluorescence complementation

(BiFC) assays for the visualization of protein interactions in living cells. *Nat. Protoc.* **1**,

1278–1286 (2006).

12. Omura, T. Mitochondria-targeting sequence, a multi-role sorting sequence recognized at

all steps of protein import into mitochondria. *J. Biochem.* **123**, 1010–1016 (1998).

13. Webb, B. D. *et al.* Recessive pathogenic variants in MCAT cause combined oxidative

phosphorylation deficiency. *Elife* **12**, (2023).

14. Landrum, M. J. *et al.* ClinVar: improving access to variant interpretations and supporting

evidence. *Nucleic Acids Res.* **46**, D1062–D1067 (2018).

15. Naquet, P., Kerr, E. W., Vickers, S. D. & Leonardi, R. Regulation of coenzyme A levels by

degradation: the "Ins and Outs." *Prog. Lipid Res.* **78**, 101028 (2020).

16. Kastaniotis, A. J., Autio, K. J. & R Nair, R. Mitochondrial fatty acids and neurodegenerative

disorders. *Neuroscientist* **27**, 143–158 (2021).

17. Lambrechts, R. A. *et al.* CoA-dependent activation of mitochondrial acyl carrier protein

links four neurodegenerative diseases. *EMBO Mol. Med.* **11**, e10488 (2019).

18. Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide libraries for

CRISPR screening. *Nat. Methods* **11**, 783–784 (2014).

19. Rausch, T., Fritz, M. H.-Y., Untergasser, A. & Benes, V. Tracy: basecalling, alignment,

assembly and deconvolution of sanger chromatogram trace files. *BMC Genomics* **21**, 230

(2020).

20. Trefely, S., Ashwell, P. & Snyder, N. W. FluxFix: automatic isotopologue normalization for

metabolic tracer analysis. *BMC Bioinformatics* **17**, 485 (2016).

21. Meyers, R. M. *et al.* Computational correction of copy number effect improves specificity

of CRISPR–Cas9 essentiality screens in cancer cells. *Nat. Genet.* **49**, 1779–1784 (2017).

22. Dempster, J. M. *et al.* Extracting biological insights from the project Achilles genome-

scale CRISPR screens in cancer cell lines. *bioRxiv* (2019) doi:10.1101/720243.

23. Dempster, J. M. *et al.* Chronos: a CRISPR cell population dynamics model. *bioRxiv*

(2021) doi:10.1101/2021.02.25.432728.

24. Arnold, P. K. *et al.* A non-canonical tricarboxylic acid cycle underlies cellular identity.

Nature **603**, 477–481 (2022).

25. Pacini, C. *et al.* Integrated cross-study datasets of genetic dependencies in cancer. *Nat.*

Commun. **12**, 1661 (2021).

FIGURE 1: AASDHPPT is required for mitochondrial oxidative metabolism

FIGURE 2: AASDHPPT is required for mtFAS pathway function

A **2D network:** Gene essentiality score correlations

LYRM7
NDUFB9
LIPT2
NDUFA6
LIPT1
MECR
LIAS
MCAT
OXSM
AASDHPPT
NDUFAB1
LYRM4
ACSF3
PANK2
ETFRF1
LYRM2
ACACA
FASN

Cluster 1
Cluster 2
Cluster 3
Cluster 4

B

\#
*
\#
AASDHPPT
DLAT
Lipoic Acid
DLST
SDHB
holo
apo
NDUFAB1
ATP5a
β-Actin

CTL-1 CTL-2 PPT-1 PPT-2

C

mtDSRed   PPT GFP

GFP
DLAT
Lipoic Acid
DLST
SDHB
holo
apo
NDUFAB1
ATP5a
β-Actin

CTL-1 CTL-2 PPT-1 PPT-2 PPT-1 PPT-2

D Oxygen consumption

Oligomycin   FCCP   Rotenone/Antimycin A

CTL-1
PPT-1
PPT-2
PPT-1 + PPT
PPT-2 + PPT

OCR (pmol/min/norm. unit)
Time (mins)

FIGURE 3: AASDHPPT localizes to the mitochondrial matrix

FIGURE 4: Mitochondrial AASDHPPT is required for mtFAS function

EXTENDED DATA FIGURE 1

A PPT-1/2 mutations

Cell line	NC_000075.7	Protein
PPT-1	c.342_344delAGC c.398G>C c.478C>G c.479T>G c.739C>T c.740T>G c.744C>T c.745G>T c.752G>A c.754T>A c.758G>A c.764A>T c.767Tdel c.771G>A c.773G>A c.779G>A	p.Q41del p.A60P p.A62G p.R64F p.M66I p.I67K p.L70F p.A72Q p.E73K p.K74N p.L75*
	c.342_346delAGCCG	p.0? (Frameshift)
PPT-2	c.366G>T c.394G>C c.757G>T c.771G>T	p.G49V p.K58N p.R68M p.E73*
	c.342_346delAGCCG	p.0? (Frameshift)

B Brightfield: Proliferation (72 hrs)

Glucose: CTL-1, PPT-1, PPT-2
Galactose: CTL-1, PPT-1, PPT-2

C Fumarate
D Malate
E Aspartate
F Citrate
G α-Ketoglutarate

H Citrate
I Malate

A

B

C

Representative flow cytometry gating strategy: