Genetic variation in trophic avoidance shows fruit flies are generally attracted to bacterial pathogens

Katy M. Monteith[1], Phoebe Thornhill[1], Pedro F. Vale[1]*

[1]*Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh, Edinburgh EH9 3FL, United Kingdom.*

*Correspondence

pedro.vale@ed.ac.uk

Keywords: Pathogen avoidance, GWAS, genetic variation, bacterial infection, Immune deficiency (IMD) pathway, feeding

Abstract

Pathogen avoidance behaviours are often assumed to be an adaptive host defence from

infection. However, there is limited experimental data on the prerequisite for this assumption:

heritable, intrapopulation phenotypic variation for avoidance. We investigated trophic

pathogen avoidance in 122 inbred *Drosophila melanogaster* lines, and in a derived outbred

population. Using the FlyPAD system, we tracked the feeding choice that flies made between

substrates that were either clean or contained a bacterial pathogen. We uncovered significant,

but weakly heritable variation in the preference index among fly lines. However, instead of

avoidance, most lines demonstrated a preference for several bacterial pathogens, showing

avoidance only for extremely high bacterial concentrations. Bacterial preference was not

associated with susceptibility to infection and was retained in flies with disrupted immune

signalling. Phenotype-genotype association analysis indicated several novel genes (*CG2321*,

CG2006, *ptp99A*) associated with increased preference for the bacterial substrate, while the

amino-acid transporter *sobremesa* was associated with greater aversion. Together with

previous work showing fitness benefits of consuming high-protein diets, our results suggest

that bacterial attraction may instead reflect a dietary preference for protein over carbohydrate.

More work quantifying intrapopulation variation in avoidance behaviours is needed to fully

assess its importance in host-pathogen evolutionary ecology.

Background

Behavioural immunity is an animal's first line of defence against infection and is characterized by behaviours that help avoid contact with infectious environments or infected conspecifics [1–4]. Invertebrates have evolved sophisticated mechanisms to detect and avoid pathogens involving the peripheral nervous system, specifically the gustatory and olfactory systems [5–11]. For example, geosmin, a microbial odorant, activates a specific subclass of olfactory neurons and inhibits oviposition, chemotaxis, and feeding, inducing pathogen avoidance [11]. More direct detection of pathogens also occurs via the gustatory sensory system of Drosophila, which detects bacterial cell wall components like lipopolysaccharide (LPS) and peptidoglycan (PGN). LPS is detected by bitter gustatory receptor neurons in the fly oesophagus expressing *TrpA1*, triggering feeding and oviposition avoidance [12,13], whereas PGN triggers grooming behaviour on stimulation of wing margins and legs [14] and learned avoidance of bacteria during feeding [15].

In contrast to the great detail learned about the mechanistic basis of avoidance behaviours in a few model systems [9], we currently know little about the extent of intrapopulation phenotypic or genetic variation in pathogen avoidance behaviours, the extent to which such variation is heritable, and therefore if it is likely to respond to selection [2]. Investigating the genetic and environmental factors that contribute to within-population variation in avoidance behaviours is important to understand its ecological and evolutionary consequences [2,16,17]. For instance, foraging and feeding are essential components of host ecology and are crucial for organismal reproduction and fitness, but they also provide a major route of pathogen transmission [18–22]. Characterising variability in host behaviours that contribute to individual heterogeneity in pathogen acquisition and spread is therefore a major focus of disease ecology and epidemiology [17,18,23–25].

Intrapopulation variation in avoidance behaviours is also likely to have evolutionary

consequences for both hosts and pathogens [2,26]. The ability to avoid contact with pathogens

allows healthy individuals to evade the pathology resulting from infection and further prevents

the activation of the immune response, which may be metabolically costly or even cause

immunopathology, with detrimental effects on host fitness [27–29]. Given that individuals vary

widely in susceptibility and in the extent to which they experience losses in fitness during

infection [30–33], the potential costs and benefits of avoiding infection are therefore likely to

vary between individuals [2,17]. Such variable costs, in turn, may contribute to the

maintenance of standing genetic variation in avoidance via fluctuating selection [34–36], and

affect the predicted evolutionary trajectories of pathogens [37,38]. However, without empirical

measurements of the phenotypic and genetic variation in pathogen avoidance behaviours, it

is difficult to assess their true impact on host-pathogen ecology and evolution[2,26].

In the present work, we aimed to quantify the intrapopulation variation in behavioural pathogen

avoidance in *Drosophila melanogaster*, a widely used model organism for host-pathogen

interactions and behavioural genetics[39–41]. We designed a two-choice feeding assay using

the FlyPAD system for tracking, recording, and analysing the feeding behaviour of fruit flies in

real-time [42]. We assayed the preference for (or aversion of) several bacterial pathogens in

a panel of 122 inbred DGRP lines and in a DGRP-derived outbred population of flies. Our

assay offered flies a choice of feeding from a clean substrate or an identical substrate to which

an exponentially growing inoculum of bacteria had been added. This assay therefore

mimicked a frequent natural scenario where insects must make foraging decisions between

food sources that may or may not contain sources of pathogenic infection [19,21,22,43].

Results

Variation in fly aversion to a bacterial feeding substrate

Fruit flies are naturally attracted to bacterial volatiles [44–47], but previous work has also

established that adult *D. melanogaster* actively avoid sources of pathogenic bacterial infection

[13,15,48]. Our starting point was therefore to investigate the extent to which pathogen

avoidance varied within a population, and how much of this variation could be explained by all

sources of genetic variation, that is, the broad sense heritability H^2 [49,50].

We exposed 122 DGRP lines to a choice between a clean substrate of 5% sucrose and an

identical substrate containing 5% of an overnight culture (OD_{600}= 1) of the bacterial pathogen

Pseudomonas entomophila. We observed substantial among line genetic variation (Line effect

p <0.001), with mean preference index scores ranging from to 0.22 (SE ± 0.16 – RAL-380)

indicating pathogen avoidance, to -0.66 ± 0.06 (RAL-879) indicating strong attraction to food

contaminated with *P. entomophila*. Contrary to our initial expectation, we found that the

majority of DGRP lines (108 / 122) showed a preference for the pathogen-contaminated

substrate, with a mean preference index across all 122 lines of -0.22 (median -0.31) (**Figure

1A ,Table 1**). We detected a non-zero, but weak, genetic component underlying this variation

in preference, with broad sense heritability estimate of 0.07-0.08 (Table 1).

Using the same feeding data, we were also able to calculate the total number of sips taken on

both substrates by each fly line, as a measure of feeding activity. Feeding activity is relevant

in the context of pathogen exposure because more active feeders may be more likely to

acquire infection orally if they do not avoid contaminated substrates[51]. This analysis also

revealed substantial genetic variation in feeding (Line effect p <0.001) ranging from a mean

of 35.5±14 sips (RAL-790) to 1047±140 sips (RAL 059) measured over a 30-minute period

(**Figure 1B**). These results are consistent with previous measures of genetic variation in food

intake in the DGRP panel [52]. We identified a significant (but weak) negative relationship

between the number of sips and the preference index (R= -0.2, p=0.029), indicating that fly

lines with more active feeding also showed a stronger preference for the pathogen-

contaminated substrate – although many low feeders also showed a preference for the

pathogen-contaminated substrate (**Figure 1C**).

Variance component	Binomial model using the number of sips on each substrate as the response	Linear model with preference index as the response
Mean	-0.22	-
Median	-0.31	-
Genetic Variance (V_G)	0.25	0.25
Phenotypic Variance (V_P)	3.53	3.29
Environmental Variance (V_E)	3.28	3.04
Broad-sense Heritability H^2	0.071	0.077

Table 1. Descriptive statistics and variance components of fly preference index. We

calculated the broad-sense heritability by partitioning the variance into genetic variance and

residual variance caused by other factors. We calculated broad-sense heritability as $H^2 =$

$V_G/(V_G + V_E)$ in two separate but similar models. In one analysis we fit a model with a binomial

error structure, where the response variable included the total number of sips taken by flies

on each substrate; in a second model, we modelled the preference index in a linear model.

Both approaches yielded very similar estimates of variance components.

Figure 1. Genetic variation. A. We measured the choice index in 122 lines from the

Drosophila Genetic Reference Panel (DGRP). We measured pathogen avoidance as a

preference index, calculated over 30 minutes, as $(sips_{clean} - sips_{pathogen}) / sips_{total}$. Positive

values of the preference index indicate pathogen avoidance, where a value of 1 indicates

complete preference for the clean substrate; negative values suggest a pathogen preference,

where -1 indicates complete preference for the pathogen-contaminated substrate; 0 indicates

an equal number of sips on each substrate. Each line was replicated between 16-24 times. B)

Line means for the total number of sips taken during a 30-min FlyPAD feeding assay. C. The

relationship between the preference index in A and the total number of sips shown in B. Each

data point is the line mean for each trait. R is the Pearson correlation coefficient, and the line

shows the linear relationship, shaded by the 95% CI. FlyPAD image from https://flypad.rocks/.

Drosophila are attracted to several bacterial pathogen species

To investigate the generality of the preference for pathogenic substrates observed in our previous experiments, we tested whether flies also preferred substrates contaminated with other pathogens. To this end, we repeated the avoidance assays with three Gram-negative bacterial pathogens (*P. entomophila*; *P. aeruginosa* -PA14; *Providencia rettgeri)* and with the Gram-positive *Enterococcus faecalis*. Instead of repeating these assays on the set of 122 lines, we instead employed an advanced outcrossed population derived from the DGRP panel called the Ashworth Outcrossed Population (AOx Population). Briefly, this outbred population was established from 100 pairwise crosses of 113 DGRP lines in order to obtain an outbred population, with similar levels of genetic diversity present within the DGRP [49,53], which at the time of the experiment had been maintained for at least 90 generations of outcrossing. This approach was useful, as it allowed us to test pathogen avoidance on fewer flies and showed a mean response that was comparable to that of the 122 DGRP lines (Figures 1A and 2). Further, while the results using the individual DGRP lines included only female flies, here we included males and females which allowed us to test if pathogen preference was a female-specific response or common in both sexes.

Overall, we observed similar outcomes to the initial DGRP choice assay, and both male and female flies showed a general preference for the pathogen-contaminated substrate, particularly when exposed to *P. entomophila* (preference index (PI) = -0.33± 0.06; test if mean is different from zero: t = -5.5022, df = 119, p = 2.186e-07), *P. rettgeri* (PI= -0.34±0.05; t = -7.1034, df = 168, p = 3.294e-11), and *P. aeruginosa* (PI = -0.16±0.05; t = -3.0766, df = 158, p = 0.0025)(**Figure 2**). We found no evidence for pathogen preference (or avoidance) when flies were exposed to *E. faecalis* (PI= -0.05±0.060; t = -0.925, df = 158, p = 0.3564). We did however, observe clear pathogen avoidance when we increased the concentration of the PA14 from OD1 to OD100 (PI= 0.61± 0.06; t = 10.756, df = 111, p- < 0.001). This demonstrates that there are conditions under which flies will find bacterial cultures unpalatable, but such high concentrations of bacteria are unlikely to be common in nature. Under more realistic

conditions, where concentrations are that of an overnight bacterial culture, flies did not avoid

bacterial pathogens and in some cases showed a strong preference for pathogen-

contaminated substrates. We considered the possibility that the starvation period experienced

by flies prior to the FlyPAD assay may have influenced their preference [42], but we found

similar preference patterns when we compared two experiments using either a short (4-6-

hour) starvation period or a longer (18-24-hour) starvation period (Fig S1).

Figure 2. The preference index measured in response to different bacterial pathogens.

Two-choice feeding assays were carried out as described for Figure 1 but using flies from an

DGRP-derived advanced outbred population. We calculated the preference index (PI) in

males and females given a choice of a clean substrate (5% sucrose) or OD_{600}=1 *P.*

entomophila (n= 55 females; 65 males), *P. aeruginosa* (n= 78 females; 81 males)*, Providencia*

rettgeri (n= 83 females; 86 males)*, Enterococcus faecalis* (n= 73 females; 86 males)*, or P.*

aeruginosa (OD600=100; n= 56 females; 56 males). Each data point is the PI of an individual

fly. Points are coloured according to the PI (blue indicates pathogen avoidance and red

indicates pathogen preference). Black dots and error bars are the mean±SE of each group.

FlyPAD image from https://flypad.rocks/.

Variation in pathogen preference index is not driven by differences in susceptibility to bacterial infection

We hypothesized that the strong preference for pathogenic substrates could be driven by differences in susceptibility between fly lines, whereby lines with a strong preference for pathogenic substrates might also be less susceptible to bacterial infection. We tested this hypothesis in two ways. First, we chose the fly lines with either the highest or lowest preference index (five lines per extreme) and we exposed these flies to an oral *P. entomophila* infection ($OD_{600}= 50$) as an assay of infection susceptibility[54]. Most flies died within two days of infection and almost all flies had died within 96 hours. We could not distinguish between flies showing weak or strong preference for the pathogen substrate based on their mortality profiles (hazard ratio: 1.099± 0.07 SE; z=1.34, p=0.18), suggesting that lines with a very high attraction to pathogen substrates when feeding did not have greater capacity to resist or clear a bacterial infection (**Figure 3A**).

We went further to test the relationship between the measured preference index and the susceptibility of flies to an orally acquired, enteric *P. entomophila* infection. Here, we were able to use previously published data on susceptibility to *P. entomophila* oral infection in the DGRP, which contained 73 DGRP lines in common to our experiment [30]. In concordance with the oral infection of the lines showing extreme preference (**Figure 3A**), we found no significant association between the line means for the pathogen preference we measured, and the line means for survival of flies three days following oral infection (**Figure 3B**). Taken together, these analyses of infection data suggest that variation between fly lines in the preference for pathogenic substrates is unlikely to be driven by differences in susceptibility.

Figure 3. Testing associations between the preference index and infection

susceptibility. A) Five lines with the highest (red) and five lines with the lowest preference

index (blue) were challenged with an oral *P. entomophila* infection (as described in [54,55]) at

a concentration of $OD_{600} = 50$; n = 7 replicate groups, with 10 flies per group (70 flies) per

treatment. Crosses indicate censored flies that remained alive at 96h post infection. B) The

relationship between the preference index and the percentage of flies that died after 3 days of

oral infection with *P. entomophila*. Each data point is the line mean of preference measured

for each DGRP line in Figure 1 and the survival of the same line as measured in [30]. R is the

Pearson correlation coefficient, and the line shows the linear relationship, shaded by the 95%

CI.

A functional Immune Deficiency (IMD) pathway is not required for pathogen attraction

Both infection assays described above were designed to guarantee that flies acquire an

infection, and therefore reflect susceptibility once the pathogen has established infection.

However, we might expect the main immune-related driver (if any) of pathogen avoidance

would be the susceptibility to acquire infection via the oral infection route, which may be under

potentially distinct immune control. To further investigate a potential role of immunity in

shaping the observed preference for pathogenic substrates, we took a functional genetic

approach and performed the same two-choice assay using transgenic Drosophila lines with

disrupted IMD-pathway function, which is the primary immune pathway involved in the

response to gram-negative bacterial infection in Drosophila [56]. These experiments were also

motivated by previous work showing that the IMD pathway, particularly the peptidoglycan

receptors PGRP-LC and PGRP-LB, play a direct role in the detection of pathogen cues and

subsequent neuroimmune signalling involved in pathogen avoidance behaviours [7–9].

We found that the control line w^{1118} displayed a preference for the pathogenic substrate (P.

$aeruginosa$ in this case), as observed in the DGRP panel, and the DGRP-derived AOx

population. Disrupting the function of $Relish,$ the main NF-kB transcription factor in the IMD

pathway (**Figure 4A**), or the peptidoglycan receptors $PGRP$-LB or $PGRP$-LC did not

significantly alter this preference (Figure 4), suggesting that IMD-mediated immunity is not

involved in the observed preference for pathogenic substrates.

Figure 4. Pathogen preference measured in IMD loss-of-function lines. We used three

transgenic lines with loss-of-function of specific IMD-pathway signalling components. Each

line was isogenised onto w^{1118}, which acts as a control with complete IMD signalling.

Preference assays were carried out as described for experiments above (Fig 1 and Fig 2),

negative values indicate a preference for the bacterial substrate. ns – non-significant

difference tested using pairwise contrasts between w^{1118} and each loss-of-function line.

Genotype-phenotype associations reveal novel genetic variants associated with

pathogen preference

The DGRP is often employed as a powerful resource for identifying genetic variants underlying

complex traits through genome-wide association studies (GWAS)[49,57,58]. Despite the low

broad-sense heritability (Table 1) and that the 122 lines used in this study are likely unpowered

to detect small effect variants [49], we decided to run a GWAS analysis in an attempt to identify

any putative large-effect candidate variants associated with the phenotypic variation in the

feeding preference.

GWAS analysis revealed a clear peak on chromosome 3R showing a strong association (p <

10^{-8}) between variation in the preference index and several SNPs (Figure 5). To refine our

analysis to the most promising candidate variants, we focused on variants with a p-value < 10^{-6}

, which resulted a list of 48 SNPs (**Table S1**). Notably, the top ten associations ranked by p-

value all mapped to a gene-dense region spanning approximately 2000 bp on chromosome

3R. SNPs in this region mainly caused synonymous changes in three adjacent genes:

CG2321 and *CG2006* are coding proteins with nuclear expression but which are otherwise

poorly described and have currently unknown functions; one of the SNPs in CG2006 also

maps to *spase12*, which is essential for cell differentiation and development in Drosophila [59].

These genes are directly downstream of *protein tyrosine phosphatase 99A (ptp99A)*, also

included in the top 48 SNPs (Table S1). *ptp99A* has previously described functions in axon

growth and guidance during neuronal development [60]. *ptp99A* has also been described as

having a role in the regulation of food intake in a GWAS of Drosophila feeding behaviour[52].

Despite resulting in synonymous changes, all variants in this region of chromosome 3R were

associated with a decrease in preference index, that is, attraction to the pathogen substrate

(Figure 5).

Variants associated with increased avoidance of bacteria (a positive preference index) were

much less common, though this is to be expected given the predominance of DGRP lines

showing bacterial preference. However, within the top 48 variants, two independent SNPs

were associated with increased avoidance, both located on the X chromosome and

specifically on gene *CG9413*, also known as *sobremesa* (Figure 5). *sobremesa* encodes an

SLC7 transmembrane amino acid transporter, which plays a crucial role in regulating amino

acid homeostasis and feeding into signalling pathways that control developmental timing,

growth, and feeding [61,62].s*obremesa* is expressed in larval glial cells and has been shown

to modulate brain development in larval stage *D. melanogaster* [63,64]. While the function of

sobremesa has only recently been described to be involved in the regulation of larval growth,

it is also expressed in adult head and gut tissues (Flybase FBgn0030574; FB2024_02; [65]),

so it is plausible that it functions as an amino-acid sensor in adult flies and could therefore

modulate behavioural changes in response to their amino-acid environment.

Figure 5. GWAS. A) The QQ plot shows quantile distribution of observed variant-phenotype

association p-values compared to the distribution of expected p-values by chance; deviations

above the diagonal line (and λ >1.00) indicate associations with a smaller p-value (more

significant) than would be expected by chance. B) Manhattan plot showing single-SNP GWAS,

testing the statistical association between the preference index and the SNPs in the genomes

of the 122 lines tested. Each point is a SNP found along the genome, and fly chromosomes

are labelled on the x-axis; the y-axis shows the -log10 p-values, where higher points indicate

stronger associations. C) The preference index of DGRP lines carrying either the reference or

variant alleles for the most significant SNPs on three genes clustered on chromosome 3R –

CG2321, *CG2006*, *ptp99A*. We also plot *CG9314/ sobremesa* as the most significant variant

associated with an increased preference index (greater pathogen avoidance). See Table S1

for a list of 48 SNPs with p-values $<10^{-6}$.

Potential fitness benefits of pathogen preference

Given the lack of association with immunity, and that the GWAS analysis indicated two

different genes (*ptp99A* and *sobremesa*) with putative roles in feeding [52] or nutrient

sensing[63], one possibility is that the preference for pathogen-contaminated substrates we

observe may in fact be driven by a preference for amino-acids. Due the make-up of bacterial

cell-wall components such as peptidoglycans, amino-acids were more abundant in the

substrate containing the bacterial inoculum relative to the sucrose-only substrate. For

example, the peptidoglycans of *Pseudomonas aeruginosa* are typically composed of an

amino-acid chain including L-alanine, D-glutamic acid, Diaminopimelic acid (DAP), and D-

alanine [66]. A plausible hypothesis is that by choosing a bacteria-contaminated substrate,

flies are in fact seeking out protein over sugar and that this diet-driven choice is presumably

adaptive.

A strength of DGRP as a community resource is that many phenotypes have been carefully

measured in the same lines in multiple labs [49,57,58]. We therefore analysed data from a

previous study which had investigated the macronutrient tolerance in the DGRP by measuring

the percentage of larvae that pupated and eventually emerged as adult flies under six different

diets [67]. This work included a <u>H</u>igh <u>S</u>ugar <u>D</u>iet (HSD), and a <u>H</u>igh <u>P</u>rotein <u>D</u>iet (HPD), that

are comparable to our 5% sucrose agar (clean) substrate and our pathogen substrate,

respectively. This data shows that both aspects of fly developmental viability are generally

higher in the HPD when compared to the HSD, regardless of the preference index we

measured (Figure 6). This is broadly in agreement with our observation that almost all DGRP

lines showed a preference for pathogen contaminated food which was relatively richer in

protein (Figure 1A).

Figure 6. The effects of diet on Drosophila developmental traits. A) The percentage of

larvae pupating and B) the percentage of eclosing adults, each on high protein vs a high sugar

diet. Each data point is the line mean of a single DGPR line for each trait/diet. Data from [67].

The colour of each point indicates the preference index for each DGRP line as shown in Figure

1, where red indicates pathogen preference and blue indicates pathogen avoidance. The

diagonal line indicated a correlation of 1 if traits are equal on both diets. Points above the

diagonal indicate a higher trait value on the high protein diet.

Discussion

Avoidance behaviours are often assumed to be adaptive, but it is notable how we know so

little about the basic requisites for this assumption[2]. To be adaptive, avoidance must be

amenable to respond to selection, which requires heritable phenotypic variation that this

variation is associated with differences in fitness [50,68]. Yet, while elegant functional

neurogenetic experiments have dissected the mechanisms underlying avoidance behaviours,

especially in invertebrates [9], we lack measurements of phenotypic and genotypic variation

in avoidance behaviours is most species [2,17,69]. Taking our results together, we find

substantial intrapopulation variation in the likelihood to avoid infection, measured across 122

genetic backgrounds, although we find that this variation is only weakly heritable. Contrary to

our initial expectation however, we found that most genotypes did not avoid the bacterial

substrate, but instead preferred to feed on it.

We also hypothesised that behavioural avoidance would be associated with immune-mediated

susceptibility. First, we might we expect an avoidance-resistance trade-off where lines that

are strongly attracted to pathogenic substrates may show higher resistance to infection as

found previously in salmon [70] and sheep [71]. However, we did not find any evidence for

trade-offs. Further, previous work has implicated specific components of the IMD signalling

pathway, such as PGRPs, in the detections and signalling of bacteria-derived cues [7,8].

However, we did not detect any substantial effect of disrupting immune signalling on the

preference for pathogen substrates. Instead, we argue that this preference is most likely driven

by a dietary preference for protein. This is supported by the outcome of the genotype-

phenotype association analysis, which implicated the role of *ptp99A,* previously linked to

variation in food intake and *sobremesa*, an amino acid transporter, and also by the general

fitness advantage of high protein diets compared a high sugar diet.

A potential limitation of our experimental design is the precise choice flies were presented

with, which explicitly tested a preference between clean, sugar-only substrate, and a substrate

replacing the sugar with an overnight culture of bacteria. This design does not allow

distinguishing between a fly preference for consuming bacteria, or a bacteria-induced

attraction. For example, some bacterial volatiles like ammonia and amines may attract flies,

while short-chain fatty acids derived from bacteria also induce attraction [44–47]. However,

while previous work has shown that flies detect and avoid odours associated with pathogenic

infection [11], other work has found, as we have here, that flies initially prefer the odour of

pathogenic bacteria - even when given a choice of two strains that are identical except for the

production of a virulence factor [15]. While this bacterial preference was reduced if flies were

fed with bacteria before the choice assay, this was the case whether or not the fed bacteria

were pathogenic, suggesting that the main driver of the change in preference was not a

learned avoidance of pathogens, but more likely a feeding/satiety-related response[15].

These results are concordant with ours, indicating that flies are generally attracted to bacterial

substrates and that these choices are likely driven by a fly's dietary amino-acid requirements.

Independently of the precise mechanism driving the preference for bacterial pathogens, this

pattern of variation is likely to have important implications for the ecology and evolution of

infection. In the simplest sense, broad attraction to pathogen-contaminated substrates will

increase the likelihood of contact between flies and infectious sources. Further, given that we

find there is no association with susceptibility, this should translate into a higher prevalence of

infection and its associated fitness costs. In support of this prediction, a model of behavioural

resistance to parasites – which under some assumptions is equivalent to pathogen avoidance

– shows that broadly, lower average levels of avoidance are expected to lead to higher

infection prevalence [38]. Other work investigating the role of individual trait heterogeneity on

the population-level epidemic outcomes found that populations with high levels of behavioural

variance (related to avoidance and relevant for contact rates between infectious and non-

infectious individuals) were more likely to experience more severe epidemics[24].   The

evolutionary implications of higher infection prevalence, in turn, will depend on the relative

costs of infection and of both behavioural and other physiological and immune defences

[36,38].

In sum, avoidance behaviours no doubt play an important role in host defence by avoiding the

detrimental effects of infection, but also by allowing individuals to avoid the physiological costs

of immune deployment [28,72]. It is important to recognise, though, that individuals are likely

to vary in their propensity to avoid sources of infection, particularly when these come at the

cost of not feeding or reproducing. In some cases, what is perceived as avoidance may

instead be the by-product of selection on these other fitness-enhancing activities [19,69].

Understanding the ecological and evolutionary drivers of avoidance, and how avoidance is

likely to drive the ecology and evolution of hosts and pathogens therefore remains an

important and understudied topic of research, and requires more widespread measurement

of the extent to which individuals vary in avoidance and how these behavioural choices may

impact fitness [2,16,26,73,74].

Methods

Fly lines and outbred population

We measured the choice index in individual, mated female flies from 122 lines from the

Drosophila Genetic Reference Panel (DGRP), a panel of fully-sequenced fly lines originating

from the same population, and therefore offers a useful snapshot of natural genetic variation

in pathogen avoidance behaviour [49,58]. These specific lines have been maintained in our

lab since 2013 and differ from the original DGRP in that they have been previously cleared of

from the endosymbiont *Wolbachia* by rearing with the antibiotic Tetracycline for two

generations [75]. Other lines used were w^{1118} (Vienna *Drosophila* Resource Center); Rel^{E20}

(*relish - IMD* pathway regulator [76]); PGRP-LC (Peptidoglycan receptor protein-LC;

BDSC_12500)[77] ; PGRP-LB (BDSC_55715)[78].

In some assays, we employed an outbred fly population. The Ashworth Advanced Outcrossed

(AOx) population is a large, lab adapted, outbred *Drosophila melanogaster* population derived

from the DGRP panel [58]. The AOx was originally established in October 2014 by setting up

100 crosses of 113 DGRP lines [79], and has been maintained since then as an outbred

population on a 30-day generation cycle with census size of between 3000-4000 adults in

each generation. Every generation, eggs are collected from a population cage (a plastic

container: 30 cm × length × 20 cm width × 20 cm height) and dispensed into 6oz bottles (57

mm diameter x 103 mm height containing cornmeal-agar-yeast medium[80] at a density of

100 (±20) eggs per bottle. 25 bottles were set up at each generation and incubated at 25 °C,

12:12 hour LD cycle. The development time of these flies is 9-10 days. Four weeks post egg

collection, all adults present in bottles are transferred to a population cage and provided with

two fresh apple agar plates (in a 90 mm Petri dish) supplemented with ad libitum yeast paste

in the centre of each plate. 6-8 hours later all eggs are collected by careful washing and

distributed to new bottles to begin the next generation.

441   Bacterial strains and culture conditions

Depending on the experiment we used the following bacterial strains: *Pseudomonas*

entomophila [55,81], *P. aeruginosa* [82,83]*, Providencia rettgeri [84,85], Enterococcus*

faecalis [86]. Unless otherwise stated, all bacterial cultures were prepared in the same way.

A single colony stored at -70ºC in 100µl of 25% glycerol was inoculated in 10mL Luria-Bertani

(LB) Broth and left to grow overnight in an orbital shaker at 37°C, 140 rpm. 10mL of the culture

was reinoculated at 1:10 and placed at 37°C, 140 rpm until $OD_{600nm}=1$. In experiments using

higher bacterial concentrations, overnight cultures were grown until exponential growth phase

was reached (OD_{600nm} 0.6-0.8). This culture was then divided equally into 50mL falcon tubes

and centrifuged at 2500 rpm for 15 minutes to form pellets and the supernatant removed. The

pellets were resuspended by shaking and recombined in a single falcon tube, which was

centrifuged again. The supernatant was removed, and the pellet resuspended in the

appropriate volume of LB broth to reach $OD_{600nm}=50$ (in the experiment with immune deletion

lines) or $OD_{600nm}=100$ (for one of the *P. aeruginosa* feeding assays).

FlyPAD choice assay

We used the FlyPAD system for tracking, recording, and analysing the feeding behaviour of fruit flies in real-time [42]. Two-to-six-day-old flies were wet-starved overnight for 18-22 hours prior to each assay. Following the starvation period, flies were placed individually within a FlyPAD arena previously prepared with two choice substrates, one on each electrode. The "clean" choice substrate contained (LB broth, 2% agar & 5% sucrose); the infectious substrate was identical, but the sucrose was replaced with 5% of an overnight bacterial culture, adjusted to OD_{600}=1. Each FlyPAD arena received 1ul of each substrate, carefully pipetted onto each electrode, and choice assays lasted 30 min [42]. Each DGRP line was replicated between 16-24 times, in blocks of 32 flies. We carried our four blocks each day, two between 9h-12h and a third and fourth block between 14h-17h, randomising the genotypes between blocks to minimise any potential time-of-day effects. The order in which the substrate (clean or bacterial) was pipetted on to the FlyPAD was also alternated between blocks to account for any potential pipetting bias. We measured pathogen avoidance as a preference index, calculated over 30 minutes, as $(sips_{clean} - sips_{pathogen}) / sips_{total}$, that is, the difference in sips taken by an individual fly on either pathogen-contaminated substrate relative or a clean alternative, relative to the total number of sips. Positive values of the preference index are suggestive of pathogen avoidance, where a value of 1 indicates complete preference for the clean substrate; negative values suggest a pathogen preference, where -1 indicates complete preference for the pathogen-contaminated substrate; 0 indicates an equal number of sips on each substrate.

Phenotype-Genotype Association Analysis

We carried out Phenotype-Genotype Association Analysis using well established analysis pipelines[49,58]. Specifically, we used the GWAS analysis as implemented by DGRPool (https://dgrpool.epfl.ch/ [57], which uses Plink2 v2.00a3LM (1 July 2021) and runs all variants in the DGRP database, using the dm3 genome assembly (4,438,427 variants: 3,963,420 SNPs, 293,363 deletions, 169,053 insertions and 12,591 MNPs). While the DGRP lines used in this study have been previously cleared of the endosymbiont *Wolbachia* (see above), the

DGRPool implementation of GWAS includes the *Wolbachia* status of the lines and five major

insertions as covariates in the model[87]. We evaluated single marker associations using

preference index line means with common variants and then used the resulting quantile-

quantile QQ plots and Manhattan plots to identify strong associations between genomic

variants and variation in each phenotype. We focused on candidate variants with $p < 10^{-6}$ that

mapped to coding genes or that that occurred within 5Kb of a coding gene and a minor allele

frequency (MAF) >5% [57].

Data analysis

All data and analysis code is available at https://doi.org/10.5281/zenodo.11149149 [88]. All data

was analysed in R (v. 4.2.2) and RStudio (2023.12.1+402), using *tidyverse* for data

manipulation and *ggplot2* to plot all figures[89]. We used the package *lme4* [90] to fit

generalized linear mixed models (GLMM- using the function "*glmer*") and linear mixed models

using the function "*lmer*"; the package *lmerTest* was used to extract *p*-values from models[91].

We used the package *performance* [92], using the "*check_model*" function to evaluate the

model fits, to test for severe deviations from gaussian-distributed residuals, and to test for

overdispersion. To analyse the preference index among 122 DGRP lines, we fitted a constant

(intercept-only) linear mixed model to predict '*preference.index*'. The model included *line*, *day*,

and *replicate* (individual fly) as random effects:

503                    $preference.index \sim 1 + (1 | line) + (1 | day) + (1 | replicate)$

504   Variance components were extracted using the *VarCorr* package, and broad-sense

505   heritability was calculated as the genetic variance *Vg* (the variance explained by DGRP line)

506   divided by the total phenotypic variance *Vp* (the residual model variance), where *Vp = Vg +*

507   *Ve* (the environmental variance)[49,50]. We used the package *coxme* to fit cox proportional

508   hazards models with random effects [93]:

509                    $coxme(Surv(hpi, censor) \sim pathogen.preference + (1 | infection.day)$

Differences between lines with a low or high preference for pathogen substrates were

inferred from the significance of *exp(coef)* from the coxme summary, which provide the

hazard ratio. Correlations between traits were carried out on the line means for each trait by

testing the significance of the Pearson correlation coefficient.

Acknowledgments

We thank Angela Reid and Alison Fulton for help with media preparation. We acknowledge

funding and support from the Branco Weiss fellowship and a Chancellor's Fellowship to PFV.

PT was funded by a University of Edinburgh scholarship, through the Wellcome Trust PhD

Programme in Hosts, Pathogens and Global Health awarded to Keith Matthews (grant

218492/Z/19/Z). For the purpose of open access, the author has applied a Creative Commons

Attribution (CC BY) license to any Author Accepted Manuscript version arising from this

submission.

Author Contributions: KMM and PFV conceived and designed the experiments. KMM and

PT performed the experiments. PFV analysed the data. PFV wrote the paper with additional

inputs from KMM and PT.

Data accessibility statement: All data and analysis code is available at

https://doi.org/10.5281/zenodo.11149149

Competing Interest Statement: The authors declare no competing interests

References

1.   Curtis VA. 2014 Infection-avoidance behaviour in humans and other animals. *Trends*
Immunol. **35**, 457–464. (doi:10.1016/j.it.2014.08.006)

2.   Gibson AK, Amoroso CR. 2022 Evolution and Ecology of Parasite Avoidance. *Annual*
Review of Ecology, Evolution and Systematics **53**, 47–67. (doi:10.1146/annurev-ecolsys-
102220-020636)

3.   Parker BJ, Barribeau SM, Laughton AM, de Roode JC, Gerardo NM. 2011 Non-
immunological defense in an evolutionary framework. *Trends in Ecology and Evolution*
26, 242–248. (doi:10.1016/j.tree.2011.02.005)

4.   Vale PF, Siva-Jothy J, Morrill A, Forbes M. 2018 The influence of parasites on insect
behavior. In *Insect Behavior: from mechanisms to ecological and evolutionary*
consequences, pp. 274–291. OUP Oxford.

5.   Kacsoh BZ, Lynch ZR, Mortimer NT, Schlenke TA. 2013 Fruit Flies Medicate Offspring
After Seeing Parasites. *Science* **339**, 947–950. (doi:10.1126/science.1229625)

6.   Kavaliers M, Choleris E, Ågmo A, Pfaff DW. 2004 Olfactory-mediated parasite
recognition and avoidance: linking genes to behavior. *Hormones and Behavior* **46**, 272–
283. (doi:10.1016/j.yhbeh.2004.03.005)

7.   Kurz CL, Charroux B, Chaduli D, Viallat-Lieutaud A, Royet J. 2017 Peptidoglycan sensing
by octopaminergic neurons modulates Drosophila oviposition. *eLife* **6**, e21937.
(doi:10.7554/eLife.21937)

8.   Masuzzo A, Manière G, Grosjean Y, Kurz L, Royet J. 2022 Bacteria-Derived Peptidoglycan
Triggers a Noncanonical Nuclear Factor-κB-Dependent Response in Drosophila
Gustatory Neurons. *J. Neurosci.* **42**, 7809–7823. (doi:10.1523/JNEUROSCI.2437-21.2022)

9.   Masuzzo A, Montanari M, Kurz L, Royet J. 2020 How Bacteria Impact Host Nervous
System and Behaviors: Lessons from Flies and Worms. *Trends Neurosci* **43**, 998–1010.
(doi:10.1016/j.tins.2020.09.007)

10. Meisel JD, Kim DH. 2014 Behavioral avoidance of pathogenic bacteria by Caenorhabditis
elegans. *Trends in Immunology* **35**, 465–470. (doi:10.1016/j.it.2014.08.008)

11. Stensmyr MC *et al.* 2012 A Conserved Dedicated Olfactory Circuit for Detecting Harmful
Microbes in Drosophila. *Cell* **151**, 1345–1357. (doi:10.1016/j.cell.2012.09.046)

12. Kim SH, Lee Y, Akitake B, Woodward OM, Guggino WB, Montell C. 2010 Drosophila
TRPA1 channel mediates chemical avoidance in gustatory receptor neurons. *Proc Natl*
Acad Sci U S A **107**, 8440–8445. (doi:10.1073/pnas.1001425107)

13. Soldano A *et al.* 2016 Gustatory-mediated avoidance of bacterial lipopolysaccharides via
TRPA1 activation in Drosophila. *Elife* **5**, e13133. (doi:10.7554/eLife.13133)

14. Yanagawa A, Neyen C, Lemaitre B, Marion-Poll F. 2017 The gram-negative sensing
receptor PGRP-LC contributes to grooming induction in Drosophila. *PLoS One* **12**,
e0185370. (doi:10.1371/journal.pone.0185370)

15. Kobler JM, Rodriguez Jimenez FJ, Petcu I, Grunwald Kadow IC. 2020 Immune Receptor
Signaling and the Mushroom Body Mediate Post-ingestion Pathogen Avoidance. *Curr*
Biol **30**, 4693-4709.e3. (doi:10.1016/j.cub.2020.09.022)

16. Buck JC, Weinstein SB, Young HS. 2018 Ecological and Evolutionary Consequences of
Parasite Avoidance. *Trends Ecol Evol* **33**, 619–632. (doi:10.1016/j.tree.2018.05.001)

17. Poirotte C, Charpentier MJE. 2022 Inter-individual variation in parasite avoidance
behaviors and its epidemiological, ecological, and evolutionary consequences. In *Animal*
Behavior and Parasitism (eds V Ezenwa, SM Altizer, R Hall), p. 0. Oxford University Press.
(doi:10.1093/oso/9780192895561.003.0015)

18. Sarabian C, Wilkinson A, Sigaud M, Kano F, Tobajas J, Darmaillacq A-S, Kalema-Zikusoka
G, Plotnik JM, MacIntosh AJJ. 2023 Disgust in animals and the application of disease
avoidance to wildlife management and conservation. *Journal of Animal Ecology* **in press**.
(doi:10.1111/1365-2656.13903)

19. Siva-Jothy JA, Monteith KM, Vale PF. 2018 Navigating infection risk during oviposition
and cannibalistic foraging in a holometabolous insect. *Behav. Ecol.* **29**, 1426–1435.
(doi:10.1093/beheco/ary106)

20. Vale PF, Jardine MD. 2017 Infection avoidance behavior: Viral exposure reduces the
motivation to forage in female Drosophila melanogaster. *Fly (Austin)* **11**, 3–9.
(doi:10.1080/19336934.2016.1207029)

21. Fouks B, Lattorff HMG. 2011 Recognition and avoidance of contaminated flowers by
foraging bumblebees (Bombus terrestris). *PLoS ONE* **6**.
(doi:10.1371/journal.pone.0026328)

22. Tillman Jr FE, Adelman JS. 2023 Searching while sick: How does disease affect foraging
decisions and contact rates? *Functional Ecology* **37**, 838–844. (doi:10.1111/1365-
2435.14207)

23. Barron D, Gervasi S, Pruitt J, Martin L. 2015 Behavioral competence: how host behaviors
can interact to influence parasite transmission risk. *Current Opinion in Behavioral*
Sciences **6**, 35–40. (doi:10.1016/j.cobeha.2015.08.002)

24. White LA, Siva-Jothy JA, Craft ME, Vale PF. 2020 Genotype and sex-based host variation
in behaviour and susceptibility drives population disease dynamics. *Proceedings of the*
Royal Society B: Biological Sciences **287**, 20201653. (doi:10.1098/rspb.2020.1653)

25. White LA, Forester JD, Craft ME. 2018 Covariation between the physiological and
behavioral components of pathogen transmission: host heterogeneity determines
epidemic outcomes. *Oikos* **127**, 538–552.

26. Hawley DM, Gibson AK, Townsend AK, Craft ME, Stephenson JF. 2021 Bidirectional
interactions between host social behaviour and parasites arise through ecological and
evolutionary processes. *Parasitology* **148**, 274–288. (doi:10.1017/S0031182020002048)

27. Lazzaro BP, Tate AT. 2022 Balancing sensitivity, risk, and immunopathology in immune
regulation. *Curr Opin Insect Sci* **50**, 100874. (doi:10.1016/j.cois.2022.100874)

28. Nystrand M, Dowling DK. 2020 Effects of immune challenge on expression of life-history
and immune trait expression in sexually reproducing metazoans—a meta-analysis. *BMC*
Biology **18**, 135. (doi:10.1186/s12915-020-00856-7)

29. Schwenke RA, Lazzaro BP, Wolfner MF. 2016 Reproduction-Immunity Trade-Offs in
Insects. *Annu. Rev. Entomol.* **61**, 239–256. (doi:10.1146/annurev-ento-010715-023924)

30. Bou Sleiman MS, Osman D, Massouras A, Hoffmann AA, Lemaitre B, Deplancke B. 2015
Genetic, molecular and physiological basis of variation in Drosophila gut
immunocompetence. *Nature Communications* **6**, 7829. (doi:10.1038/ncomms8829)

31. Vale PF, Jardine MD. 2015 Sex-specific behavioural symptoms of viral gut infection and
Wolbachia in Drosophila melanogaster. *J Insect Physiol* **82**, 28–32.
(doi:10.1016/j.jinsphys.2015.08.005)

32. Kutzer M, Gupta V, Neophytou K, Doublet V, Monteith KM, Vale PF. 2023 Intraspecific
genetic variation in host vigor, viral load, and disease tolerance during Drosophila C
Virus infection. *Open Biology* , 2022.06.09.495537. (doi:10.1101/2022.06.09.495537)

33. Wang JB, Lu H-L, Leger RJS. 2017 The genetic basis for variation in resistance to infection
in the Drosophila melanogaster genetic reference panel. *PLOS Pathogens* **13**, e1006260.
(doi:10.1371/journal.ppat.1006260)

34. Johnson OL, Tobler R, Schmidt JM, Huber CD. 2023 Fluctuating selection and the
determinants of genetic variation. *Trends in Genetics* **0**. (doi:10.1016/j.tig.2023.02.004)

35. Antonovics J, Thrall PH. 1997 The cost of resistance and the maintenance of genetic
polymorphism in host—pathogen systems. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences **257**, 105–110. (doi:10.1098/rspb.1994.0101)

36. Boots M, Best A, Miller MR, White A. 2009 The role of ecological feedbacks in the
evolution of host defence: what does theory tell us? *Philos Trans R Soc Lond B Biol Sci*
364, 27–36. (doi:10.1098/rstb.2008.0160)

37. McLeod DV, Day T. 2015 Pathogen evolution under host avoidance plasticity. *Proc. Biol.*
Sci. **282**. (doi:10.1098/rspb.2015.1656)

38. Amoroso CR, Antonovics J. 2020 Evolution of behavioural resistance in host–pathogen
systems. *Biology Letters* **16**, 20200508. (doi:10.1098/rsbl.2020.0508)

39. Troha K, Buchon N. 2019 Methods for the study of innate immunity in *Drosophila*
melanogaster. *Wiley Interdisciplinary Reviews: Developmental Biology* , e344.
(doi:10.1002/wdev.344)

40. Siva-Jothy JA, Vale PF. 2021 Dissecting genetic and sex-specific sources of host
heterogeneity in pathogen shedding and spread. *PLOS Pathogens* **17**, e1009196.
(doi:10.1371/journal.ppat.1009196)

41. Dubnau J. 2014 *Behavioral Genetics of the Fly (Drosophila Melanogaster)*. Cambridge
University Press.

42. Itskov PM, Moreira J-M, Vinnik E, Lopes G, Safarik S, Dickinson MH, Ribeiro C. 2014
Automated monitoring and quantitative analysis of feeding behaviour in *Drosophila*.
Nature Communications **5**, ncomms5560. (doi:10.1038/ncomms5560)

43. Aartsma Y, Cusumano A, Fernández de Bobadilla M, Rusman Q, Vosteen I, Poelman EH.
2019 Understanding insect foraging in complex habitats by comparing trophic levels:
insights from specialist host-parasitoid-hyperparasitoid systems. *Current Opinion in*
Insect Science **32**, 54–60. (doi:10.1016/j.cois.2018.11.001)

44. Keesey IW, Koerte S, Khallaf MA, Retzke T, Guillou A, Grosse-Wilde E, Buchon N, Knaden
M, Hansson BS. 2017 Pathogenic bacteria enhance dispersal through alteration of
Drosophila social communication. *Nature Communications* **8**, 265. (doi:10.1038/s41467-
017-00334-9)

45. Min S, Ai M, Shin SA, Suh GSB. 2013 Dedicated olfactory neurons mediating attraction
behavior to ammonia and amines in Drosophila. *Proceedings of the National Academy of*
Sciences **110**, E1321–E1329. (doi:10.1073/pnas.1215680110)

46. Venu I, Durisko Z, Xu J, Dukas R. 2014 Social attraction mediated by fruit flies'
microbiome. *Journal of Experimental Biology* **217**, 1346–1352. (doi:10.1242/jeb.099648)

47. von Hoermann C *et al.* 2022 Linking bacteria, volatiles and insects on carrion: the role of
temporal and spatial factors regulating inter-kingdom communication via volatiles. *R Soc*
Open Sci **9**, 220555. (doi:10.1098/rsos.220555)

48. Babin A, Kolly S, Schneider F, Dolivo V, Zini M, Kawecki TJ. 2014 Fruit flies learn to avoid
odours associated with virulent infection. *Biol Lett* **10**, 20140048.
(doi:10.1098/rsbl.2014.0048)

49. Mackay TFC, Huang W. 2018 Charting the genotype–phenotype map: lessons from the
Drosophila melanogaster Genetic Reference Panel. *WIREs Developmental Biology* **7**,
e289. (doi:10.1002/wdev.289)

50. Walsh B, Lynch M. 2018 *Evolution and Selection of Quantitative Traits*. Oxford: Oxford
University Press. (doi:10.1093/oso/9780198830870.001.0001)

51. Pfenning-Butterworth AC, Vetter RE, Hite JL. 2023 Natural variation in host feeding
behaviors impacts host disease and pathogen transmission potential. *Ecol Evol* **13**,
e9865. (doi:10.1002/ece3.9865)

52. Garlapow ME, Huang W, Yarboro MT, Peterson KR, Mackay TFC. 2015 Quantitative
680   Genetics of Food Intake in Drosophila melanogaster. *PLoS One* **10**, e0138129.
681   (doi:10.1371/journal.pone.0138129)

682   53. Savola E, Montgomery C, Waldron FM, Monteith KM, Vale P, Walling C. 2021 Testing
683   evolutionary explanations for the lifespan benefit of dietary restriction in fruit flies
684   (Drosophila melanogaster). *Evolution* **75**, 450–463. (doi:10.1111/evo.14146)

685   54. Siva-Jothy JA, Prakash A, Vasanthakrishnan RB, Monteith KM, Vale PF. 2018 Oral
686   Bacterial Infection and Shedding in Drosophila melanogaster. *J Vis Exp*
687   (doi:10.3791/57676)

688   55. Prakash A, Monteith KM, Vale PF. 2022 Mechanisms of damage prevention, signalling
689   and repair impact disease tolerance. *Proceedings of the Royal Society B: Biological
690   Sciences* **289**, 20220837. (doi:10.1098/rspb.2022.0837)

691   56. Myllymäki H, Valanne S, Rämet M. 2014 The Drosophila Imd Signaling Pathway. *J
692   Immunol* **192**, 3455–3462. (doi:10.4049/jimmunol.1303309)

693   57. Gardeux V, Bevers RPJ, David FPA, Rosschaert E, Rochepeau R, Deplancke B. 2023
694   DGRPool: A web tool leveraging harmonized Drosophila Genetic Reference Panel
695   phenotyping data for the study of complex traits. *eLife* **12**. (doi:10.7554/eLife.88981.1)

696   58. Mackay TFC *et al.* 2012 The Drosophila melanogaster Genetic Reference Panel. *Nature*
697   **482**, 173.

698   59. Gilbert EH, Kwak S-J, Chen R, Mardon G. 2013 Drosophila Signal Peptidase Complex
699   Member Spase12 Is Required for Development and Cell Differentiation. *PLOS ONE* **8**,
700   e60908. (doi:10.1371/journal.pone.0060908)

701   60. Hatzihristidis T, Desai N, Hutchins AP, Meng T-C, Tremblay ML, Miranda-Saavedra D.
702   2015 A *Drosophila*-centric view of protein tyrosine phosphatases. *FEBS Letters* **589**, 951–
703   966. (doi:10.1016/j.febslet.2015.03.005)

704   61. Fotiadis D, Kanai Y, Palacín M. 2013 The SLC3 and SLC7 families of amino acid
705   transporters. *Mol Aspects Med* **34**, 139–158. (doi:10.1016/j.mam.2012.10.007)

706   62. Miguel-Aliaga I. 2012 Nerveless and gutsy: intestinal nutrient sensing from invertebrates
707   to humans. *Semin Cell Dev Biol* **23**, 614–620. (doi:10.1016/j.semcdb.2012.01.002)

708   63. Galagovsky D, Depetris-Chauvin A, Manière G, Geillon F, Berthelot-Grosjean M, Noirot E,
709   Alves G, Grosjean Y. 2018 Sobremesa L-type Amino Acid Transporter Expressed in Glia Is
710   Essential for Proper Timing of Development and Brain Growth. *Cell Reports* **24**, 3156-
711   3166.e4. (doi:10.1016/j.celrep.2018.08.067)

712   64. Manière G, Alves G, Berthelot-Grosjean M, Grosjean Y. 2020 Growth regulation by
713   amino acid transporters in Drosophila larvae. *Cell. Mol. Life Sci.* **77**, 4289–4297.
714   (doi:10.1007/s00018-020-03535-6)

65. Jenkins VK, Larkin A, Thurmond J. 2022 Using FlyBase: A Database of Drosophila Genes
and Genetics. In *Drosophila: Methods and Protocols* (ed C Dahmann), pp. 1–34. New
York, NY: Springer US. (doi:10.1007/978-1-0716-2541-5_1)

66. Anderson EM, Sychantha D, Brewer D, Clarke AJ, Geddes-McAlister J, Khursigara CM.
2020 Peptidoglycomics reveals compositional changes in peptidoglycan between
biofilm- and planktonic-derived Pseudomonas aeruginosa. *J Biol Chem* **295**, 504–516.
(doi:10.1074/jbc.RA119.010505)

67. Havula E *et al.* 2022 Genetic variation of macronutrient tolerance in Drosophila
melanogaster. *Nat Commun* **13**, 1637. (doi:10.1038/s41467-022-29183-x)

68. Kingsolver JG, Pfennig DW. 2007 Patterns and Power of Phenotypic Selection in Nature.
BioScience **57**, 561–572. (doi:10.1641/B570706)

69. Amoroso CR, Shepard LL, Gibson AK. 2024 Genetic variation in parasite avoidance, yet
no evidence for constitutive fitness costs. *Evolution* , qpae030.
(doi:10.1093/evolut/qpae030)

70. Klemme I, Hyvärinen P, Karvonen A. 2020 Negative associations between parasite
avoidance, resistance and tolerance predict host health in salmonid fish populations.
Proc Biol Sci **287**, 20200388. (doi:10.1098/rspb.2020.0388)

71. Hutchings MR, Knowler KJ, McAnulty R, McEwan JC. 2007 Genetically resistant sheep
avoid parasites to a greater extent than do susceptible sheep. *Proc Biol Sci* **274**, 1839–
1844. (doi:10.1098/rspb.2007.0398)

72. Kutzer MAM, Cornish B, Jamieson M, Zawistowska O, Monteith KM, Vale PF. 2024
Mitochondrial background can explain variable costs of immune deployment. *Journal of
Evolutionary Biology* , voae027. (doi:10.1093/jeb/voae027)

73. Lopes PC, French SS, Woodhams DC, Binning SA. 2022 Infection avoidance behaviors
across vertebrate taxa: Patterns, processes, and future directions. In *Animal Behavior
and Parasitism* (eds V Ezenwa, SM Altizer, R Hall), p. 0. Oxford University Press.
(doi:10.1093/oso/9780192895561.003.0014)

74. Stockmaier S, Stroeymeyt N, Shattuck EC, Hawley DM, Meyers LA, Bolnick DI. 2021
Infectious diseases and social distancing in nature. *Science* **371**, eabc8881.
(doi:10.1126/science.abc8881)

75. Magwire MM, Fabian DK, Schweyen H, Cao C, Longdon B, Bayer F, Jiggins FM. 2012
Genome-Wide Association Studies Reveal a Simple Genetic Basis of Resistance to
Naturally Coevolving Viruses in Drosophila melanogaster. *PLoS Genet* **8**, e1003057.
(doi:10.1371/journal.pgen.1003057)

76. Hedengren M, BengtÅsling, Dushay MS, Ando I, Ekengren S, Wihlborg M, Hultmark D.
1999 Relish, a Central Factor in the Control of Humoral but Not Cellular Immunity in
Drosophila. *Molecular Cell* **4**, 827–837. (doi:10.1016/S1097-2765(00)80392-5)

77. Rämet M, Manfruelli P, Pearson A, Mathey-Prevot B, Ezekowitz RAB. 2002 Functional
genomic analysis of phagocytosis and identification of a Drosophila receptor for E. coli.
Nature **416**, 644–648. (doi:10.1038/nature735)

78. Paredes JC, Welchman DP, Poidevin M, Lemaitre B. 2011 Negative Regulation by
Amidase PGRPs Shapes the Drosophila Antibacterial Response and Protects the Fly from
Innocuous Infection. *Immunity* **35**, 770–779. (doi:10.1016/j.immuni.2011.09.018)

79. Waldron F, Monteith KM, Vale P. 2019 Creation and maintenance of Ashworth
outcrossed DGRP population V.1. (doi:10.17504/protocols.io.zbuf2nw)

80. Lewis E. 2014 A new standard food medium. 1960 Drosophila Information Service. in.
Cold Spring Harb Protoc **2014**, pdb.rec081414. (doi:10.1101/pdb.rec081414)

81. Dieppois G, Opota O, Lalucat J, Lemaitre B. 2015 Pseudomonas entomophila: A Versatile
Bacterium with Entomopathogenic Properties. In *Pseudomonas: Volume 7: New Aspects*
of Pseudomonas Biology (eds J-L Ramos, JB Goldberg, A Filloux), pp. 25–49. Dordrecht:
Springer Netherlands. (doi:10.1007/978-94-017-9555-5_2)

82. Apidianakis Y, Rahme LG. 2009 Drosophila melanogaster as a model host for studying
Pseudomonas aeruginosa infection. *Nature Protocols* **4**, 1285–1294.
(doi:10.1038/nprot.2009.124)

83. Gupta V, Vasanthakrishnan RB, Siva-Jothy J, Monteith KM, Brown SP, Vale PF. 2017 The
route of infection determines Wolbachia antibacterial protection in Drosophila.
Proceedings of the Royal Society B: Biological Sciences **284**.
(doi:10.1098/rspb.2017.0809)

84. Galac MR, Lazzaro BP. 2011 Comparative pathology of bacteria in the genus Providencia
to a natural host, Drosophila melanogaster. *Microbes and Infection* **13**, 673–683.
(doi:10.1016/j.micinf.2011.02.005)

85. Prakash A, Fenner F, Shit B, Salminen TS, Monteith KM, Khan I, Vale PF. 2024 IMD-
mediated innate immune priming increases Drosophila survival and reduces pathogen
transmission. , 2023.02.22.529244. (doi:10.1101/2023.02.22.529244)

86. Lazzaro BP, Sackton TB, Clark AG. 2006 Genetic variation in Drosophila melanogaster
resistance to infection: A comparison across bacteria. *Genetics* **174**, 1539–1554.
(doi:10.1534/genetics.105.054593)

87. Huang W *et al.* 2014 Natural variation in genome architecture among 205 Drosophila
melanogaster Genetic Reference Panel lines. *Genome Res.* **24**, 1193–1208.
(doi:10.1101/gr.171546.113)

88. Vale P. 2024 Data for 'Genetic variation in trophic avoidance shows fruit flies are
generally attracted to bacterial pathogens'. (doi:10.5281/zenodo.11149149)

89. Wickham H, Chang W, RStudio. 2016 ggplot2: Create Elegant Data Visualisations Using
the Grammar of Graphics.

90. Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* **67**, 1–48. (doi:10.18637/jss.v067.i01)

91. Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software* **82**. (doi:10.18637/jss.v082.i13)

92. Lüdecke D, Ben-Shachar M, Patil I, Waggoner P, Makowski D. 2021 performance: An R
Package for Assessment, Comparison and Testing of Statistical Models. *JOSS* **6**, 3139.
(doi:10.21105/joss.03139)

93. Therneau TM. 2015 coxme: Mixed Effects Cox Models.

797