Closely related facultative and constitutive CAM phenotypes show little transcriptomic

overlap in the subgenus *Tillandsia*

Clara Groot Crego[1,2], Sarah Saadain[1,2,3], Marylaure De La Harpe[1], Jacqueline Hess[1,4], Michael H.J.

Barfuss[1], Walter Till[1], Christian Lexer[1,†], Ovidiu Paun[1]

[1]Department of Botany and Biodiversity Research, University of Vienna, 1030 Vienna, Austria

[2]Vienna Graduate School of Population Genetics, Vienna, Austria

[3]Institut für Populationsgenetik, Vetmeduni Vienna, 1210 Vienna, Austria

[4]Cambrium GmbH, Max-Urich-Str. 3, 13055 Berlin, Germany

[†]deceased

Correspondence to: clara.groot.crego@univie.ac.at, ORCID: 0000-0002-4547-3608

Word count for Introduction: **1,057**

Word count for Methods: **2,022**

Word count for Results: **2,571**

Word count for Discussion: **1,263**

Word count for Conclusion**: 234**

Word count for main body: 7,147

Number of Figures: **7**

Number of Tables: **0**

Number of SI: **6 figures, 8 tables**

1. **Summary**

- The evolutionary trajectory of Crassulacean acid metabolism (CAM) remains elusive, as it has evolved repeatedly, yet represents a complex trait requiring changes on several organismal levels. It is debated whether distinct forms of CAM embody a single quantitative trait, or entirely separate, categorical phenotypes. The subgenus *Tillandsia* provides an excellent opportunity to compare the genetic basis of different forms of CAM, as species are closely related, yet display a broad range of CAM phenotypes.

- By combining physiological and transcriptomic analyses of well-watered and drought-stressed accessions of a *Tillandsia* species pair, we achieved a comprehensive characterization of their photosynthetic phenotypes and the transcriptomic underpinnings and overlap of distinct CAM phenotypes in *Tillandsia*.

- We found that a species previously identified as C3 displays an active CAM cycle under drought stress, yet the effect of drought was markedly increased compared to the constitutive CAM species . The CAM-specific transcriptomic response to day-night changes and to drought showed little overlap between the two species, with most DE genes showing distinct regulation patterns or being unique to one species.

- Our study reveals that the two species do not exhibit a shared transcriptomic basis of CAM expressed at distinct levels, but that different transcriptomic routes can lead to similar phenotypes, supporting the view that CAM is a highly polygenic trait.

2. Introduction

The repeated evolution of complex traits has been a long-standing matter of study for evolutionary biologists. Although one can anticipate that traits governed by a single mutation arise regularly during the course of evolution, it remains puzzling how some complex traits, necessitating alterations across multiple organismal layers, have evolved frequently and repeatedly across the tree of life.

One explanation for this is that similar phenotypes can be obtained through different underlying sets of mutations, also referred to as "genetic routes", assuming complex traits are highly polygenic and the underlying genes show a significant level of redundancy (Goldstein and Holsinger, 1992; Nowak *et al.*, 1997; Barghi, Hermisson and Schlötterer, 2020; Láruson, Yeaman and Lotterhos, 2020). Indeed, instances where similar phenotypes have evolved without a common genetic basis have been observed across a broad spectrum of life including beach mice (Steiner *et al.*, 2009), conifers (Yeaman *et al.*, 2016), lab-reared *Drosophila* (Barghi *et al.*, 2019) and cacao plants. (Hämälä *et al.*, 2020)

An example of a complex but highly evolvable trait is Crassulacean Acid Metabolism (CAM), a carbon-concentrating mechanism that has evolved independently in at least 38 plant families, amounting to a minimum of 66 independent origins (Gilman *et al.*, 2023). CAM plants restructure part of the photosynthetic metabolism across the diel cycle by assimilating CO_2 at night, rather than during the day as C3 plants do. A key difference between CAM and C3 plants is that CO_2 is temporarily converted to malate by phosphoenolpyruvate carboxylase (PEPC). The malate is stored in the vacuole overnight, and reconverted to CO_2 during the day for the Calvin Benson cycle. The CO_2 accumulated overnight allows the plant to close its stomata during the day, which has two effects: it increases the cellular concentration of CO_2 compared to O_2 in the cell, making Rubisco more efficient (Osmond, 1978), and it significantly decreases daytime evapotranspiration, increasing the plants' water use efficiency (Borland *et al.*, 2014). This is especially beneficial under high temperature or drought-stress conditions, which increase Rubisco's oxygenation activity (Heyduk, Ray and Leebens-Mack, 2021).

CAM has long been considered a complex trait, not only because of the concerted changes of temporal expression required to set the alternate biochemical pathway in motion, but also because certain anatomical features appear associated with CAM and may therefore be required for a plant to express CAM to its fullest (Nelson, Sage and Sage, 2005). Examples of such

anatomical features are larger vacuoles (Heyduk *et al.*, 2016; Males, 2018), tighter cell packaging

(Nelson and Sage, 2008) and succulence (Gibson, 1982; Griffiths *et al.*, 2008), but see (Martin,

Hsu and Lin, 2009).

Initially, CAM was regarded as a discrete trait with a bi-modal distribution of phenotypes

(a plant is either C3 or CAM), in part because the main method of measuring CAM activity using

carbon isotope measurements lacks sensitivity to detect intermediate forms (Pierce, Winter and

Griffiths, 2002; Winter and Holtum, 2002). In the meantime, other methods of measuring CAM

activity have revealed the existence of many intermediate forms, which show that at least low-

level or episodic expression of CAM can occur without major anatomical changes (Silvera,

Santiago and Winter, 2005; Heyduk, Ray and Leebens-Mack, 2021). These findings have caused

a shift towards thinking of CAM as a continuum. However, there exists a discrepancy between the

idea of CAM being a continuous trait ranging from not being expressed at all to being fully

expressed, and the many 'types' of CAM that have been described in the literature (e.g. "facultative

CAM", "weak CAM", "CAM-cycling", "strong CAM", "constitutive CAM", etc.) which are not

consistent with an ordered set of continuous phenotypes (Edwards, 2023).

It is, in other words, still unclear whether CAM is in fact a complex trait that can be

achieved by multiple routes of adaptation, or a true continuous trait with a shared genetic basis

that is expressed to different degrees among plant species. Comparative genetic analyses between

different CAM phenotypes help understand to what degree CAM evolution has a constrained

evolutionary path or a large genetic redundancy with multiple genetic routes to achieve similar

phenotypic outcomes. For example, Heyduk *et al.* (2018) found both shared and variable patterns

of gene expression in CAM-related genes among members of the Agavoideae with weak and

strong CAM phenotypes. More studies on the molecular basis of CAM on closely related taxa

expressing distinct CAM phenotypes are needed to obtain a comprehensive view of CAM

evolution.

The subgenus *Tillandsia* (genus *Tillandsia*, Bromeliaceae) is an emerging model to study

repeated evolution of ecological traits within a plant adaptive radiation. As part of one of the fastest

diversifying lineages in the plant kingdom (Givnish *et al.*, 2014), the subgenus is considered an

adaptive radiation that generated more than 250 species over the last 7 million years (Barfuss *et*

al., 2016; Mendoza *et al.*, 2017; Yardeni *et al.*, 2023). Numerous key innovation traits characterise

this radiation, including CAM, which occurs in a broad range of phenotypes ranging from C3-like
to constitutive CAM (Crayn, Winter and Smith, 2004; De La Harpe *et al.*, 2020).

In this study, we test whether facultative and constitutive CAM species have a shared
genetic basis that is expressed at different levels - suggesting that CAM is a continuous trait - or,
alternatively, achieve CAM by distinct genetic routes. We subjected plants to experimental
drought stress followed by physiological and transcriptomic analyses contrasting the
photosynthetic phenotypes of two *Tillandsia* species. The species in question, *T. leiboldiana* and
T. vanhyningii (originally *T. ionantha var. vanhyningii*, recently reassigned to species level by
Beutelspacher & Garcia-Martinez (2021)), display some of the most disparate carbon isotope
measurements in the group (De La Harpe et al. 2020). *T. leiboldiana'*s photosynthetic phenotype
was studied in depth in standard conditions by a recent study using metabolomic, comparative
genomic and transcriptomic analyses (Groot Crego *et al.*, 2024), which revealed CAM-like
expression profiles of relevant genes, yet a C3-like phenotype. This led to the suggestion that it
may be a facultative CAM plant that performs a C3 cycle in normal conditions but might have the
ability to activate a weak CAM cycle, for example under drought conditions. Our study describes
the photosynthetic phenotypes of these *Tillandsia* species in unprecedented detail and elucidate
the extent of drought tolerance of different CAM phenotypes. Our analyses reveal a lack of overlap
between species in the transcriptomic response of CAM-related genes to both day-night changes
and to drought, despite both displaying CAM under drought conditions.
135

3. Material and Methods

3.1. Experimental set-up and sampling

We designed an experiment that captures both CAM-related and drought-stress-related changes in physiology and gene expression. For this, two species of the subgenus *Tillandsia* were selected: *T. vanhyningii* and *T. leiboldiana*. While reported carbon isotope ratios classify *T. vanhyningii* as a strong and constitutive CAM plant ($\delta^{13}C$ = -13.9, (De La Harpe *et al.*, 2020)), and *T. leiboldiana* as a C3 plant ($\delta^{13}C$ = -31.3), the close relatedness of the two lineages within a clade known for multiple C3/CAM switches makes it likely that previously reported C3 species may in fact be intermediate CAM/C3 performing weak CAM, CAM idling or facultative CAM. Features of *T. leiboldiana*'s phenotype uncovered by (Groot Crego *et al.*, 2024) further indicate that it may be able to perform some form of weak CAM. We therefore designed an experimental set-up that tests the respective phenotypes of both species in normal and drought-stress conditions.

The experiment comprised six clonal accessions per species (Table S1), which were placed under identical greenhouse conditions in October 2018 at the greenhouse facilities of the Botanical Garden of the University of Vienna. Prior to the experiment, the accessions were allowed to acclimate to greenhouse conditions for 6 weeks, before being placed in a regime of 12 hours of light / 12 hours of dark for 14 days, by complementing daylight with artificial growing lights. During this time, three accessions of each species were placed in a drought regime for 14 days by not watering. The remaining three accessions were kept in normal watering conditions used to maintain the plants in the botanical garden (1x every two days, control). After 14 days, all accessions were sampled every 6 hours (Four timepoints: Day + 4 hours (10 am), Day + 10 hours (4 pm), Night + 4 hours (10 pm), Night +10 hours (4 am)), generating three replicates for the combination of treatments (species x time point x watering regime (Fig. 1). To minimise wounding, leaf material was collected at each timepoint by delicately pulling out an entire leaf from the base of the plant. The tip and base of the leaf were then removed, and the middle part was immediately placed in liquid nitrogen and stored at -80 °C.

*Figure 1. **A)** Accession of* Tillandsia leiboldiana *at the Botanical Garden of the University of Vienna (Photograph by Michael H.J. Barfuss, 2023). **B)** Accession of Tillandsia vanhyningii at the Botanical Garden of the University of Vienna (Photograph by Manuel Lasserus, 2024). **C)** Experimental setup, consisting of six clonal accessions per species, of which three specimens are used per watering treatment, sampled at four time points, resulting in three samples per species, time point and treatment. Titratable acidity was quantified for each condition, whereas RNA-seq data was analysed for the time points indicated with* [RNA].

3.2.   Titratable acidity measurements

Frozen leaf material for all four timepoints, two watering regimes and two species (48 samples in total) was weighed and then ground in a pre-cooled mortar to a powder, then added to a 5 ml solution of 80 % EtOH. The samples were then incubated for one hour in an 80 °C water bath with occasional vortexing. Tubes were centrifuged at 5,000 rpm for 5 minutes, after which the supernatant was transferred and the pellet discarded. The extractions were then let to cool to room temperature. Using a pH526 WTM Sentix mic metre, each sample was measured by titrating 0.1 N NaOH to the solution until reaching a pH of 8.2 exactly and recording the volume of added NaOH. Titratable acidity (TA) in μ-equivalents per gram of fresh weight (g FW) was calculated as μl NaOH x 0.1 mol/L/g FW.

To test whether titratable acidity significantly fluctuated across time within each species and watering regime, we constructed a Linear Mixed-Effects Model (LMM) for each species as follows:

$$titratable\ acid \sim watering\ regime \times time\ point + (\ 1\ |\ replicate\ )$$

Normality of titratable acidity measurements within each species was tested beforehand. In the LMM model, titratable acid represents the dependent variable, while watering regime and time point are fixed effects, and their interaction is included into the model. To control for random variation in titratable acidity measurements across replicates of the same time point and watering regime, the model includes a random intercept for each replicate. The analysis was performed using the *lmer* function from R package *lme4* (Bates *et al.*, 2015). The goodness of fit of the model was evaluated by plotting a QQ-Plot and the fitted values to residuals for each species. The significance of fixed effects was evaluated by obtaining a p-value using the R package *lmerTest* (Kuznetsova, Brockhoff and Christensen, 2017). A Tukey HSD test in the R package *emmeans* was performed to assess the effects of individual watering regimes and time points on titratable acidity.

To directly test for a difference between species in titratable acidity, we calculated the maximum difference in titratable acidity (ΔTA) between day and night by subtracting the TA of the two timepoints with the largest difference, which resulted to be TA_{N+10} - TA_{D+10} for all groups (species and watering regime, see Fig. 2A). Significant differences in ΔTA for watering regime and species were tested with ANOVA and Tukey HSD. We performed Shapiro-Wilk and Levene's

200 tests to ascertain that the assumptions of normality and homogeneity of variances were met. The

201 model used for ANOVA was ΔTA ~ species * watering regime.

3.3. RNA extraction and sequencing

RNA-seq data was obtained for samples at timepoints 10 am (D+4) and 10 pm (N+4) for

two watering regimes and two species (24 samples in total). Total RNA was extracted using the

QIAGEN RNeasy® Mini Kit. We evaluated the purity and concentration of the extractions using

Nanodrop, RIN and fragmentation profiles were obtained with a BioAnalyzer™ (Table S1). RNA

library preparation and poly-A capture were performed at the Vienna Biocenter Core Facilities

(VBCF) using a NEBNext stranded mRNA kit. The libraries were sequenced on one lane of

Illumina NovaSeq SP 150 PE flowcells.

3.4. RNA-seq data processing

Raw sequence quality was assessed using FastQC v. 0.11.8

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and MultiQC (Ewels *et al.*, 2016).

Adapters and low-quality bases were trimmed off using AdapterRemoval v. 2.3.1 (Schubert,

Lindgreen and Orlando, 2016) with settings --trimns --trimqualities --minquality 20 --

trimwindows 12 --minlength 36. The trimmed data was aligned to the *T. fasciculata* reference

genome (GenBank accession: GCA_029168755.2, Groot Crego et al. (2024) with STAR v. 2.7.3a

(Dobin *et al.*, 2013) in one pass using a GFF file to specify exonic regions.

3.5. Differential gene expression analysis

Read counts were quantified per exon for paired-end and reversely stranded reads (options

-p, -s 2) in the subread package FeatureCounts v. 2.0.2 (Liao, Smyth and Shi, 2014), after which

the counts of all exons of a gene were summed up to obtain per-gene counts. Differential gene

expression (DGE) analyses were performed for each species separately using the Bioconductor

package edgeR v. 3.40.0 (Robinson, McCarthy and Smyth, 2009).

We built a design matrix incorporating the two investigated treatments (timepoint and

watering regime). Genes where three or fewer samples of the same species had a Counts Per

Million (cpm) value smaller or equal to 1 were filtered out. Library size normalisation was done

using edgeR's default trimmed mean of M-values (TMM) normalisation method (Robinson and

Oshlack, 2010). The Cox-Reid profile-adjusted likelihood (CR) method was used to estimate

dispersion. Genes were tested for differential expression between timepoints and between watering
regimes separately for each species by fitting a negative binomial General Linear Model (GLM)
with the following contrasts: Day versus Night at control = Day_control - Night_control, Day
versus Night at drought = Day_drought - Night_drought, Drought vs Control at day = Drought_day
- Control_day, Drought vs Control at night = Drought_night - Control_night.

To test the fit of these GLM's, we conducted a quasi-likelihood F-test. The level of
significance was adjusted with the Benjamini-Hochberg correction method to account for multiple
testing. Genes were considered differentially expressed when the adjusted p-value was below 0.05
and the expression had an absolute log-fold change of at least 1.5. The total number of
differentially expressed genes for each treatment (timepoint and watering regime) was then
visualised and compared between species using ggplot2.

240   3.6.   Venn-diagrams

To understand the extent of shared response to drought between both species, differentially
expressed genes were visualised with Venn diagrams using the R package VennDiagram (Chen
and Boutros, 2011). Diagrams were made for the overlap of differentially expressed (DE) genes
between watering regimes (control and drought) at different timepoints, and for the overlap in DE
genes between timepoints (day and night) at different watering regimes.

To test whether overlaps in DE genes between species, conditions and timepoints were
larger or smaller than expected by random sampling, we performed a hypergeometric test in the R
package SuperExactTest (Wang, Zhao and Zhang, 2015). Since separate DE analyses were
performed for each species, the starting set of filtered genes differed between species. However,
the size of the starting set of genes was highly comparable between the two species, with 17,899
filtered genes in *T. vanhyningii* and 17,676 in *T. leiboldiana*, of which 92.5 % overlap. We
therefore performed the hypergeometric test by taking the average filtered gene set size between
the two species, and obtaining p-values for every overlap present in the Venn-diagrams.

254   3.7.   Functional interpretation of DE genes

Gene Ontology (GO) enrichment analyses were performed for both sets of DE genes (time
point and watering regime) within both species using the Bioconductor package topGO v. 2.50.0.
(Alexa, Rahnenführer and Lengauer, 2006). Significance of enrichment was evaluated with the

Fisher exact test and the algorithm "weight01". The top 100 significant GO terms were then

compiled sorted by decreasing significance using a custom-made script available at the repository

https://github.com/cgrootcrego/Tillandsia_CAM-Drought.

Aside from performing GO enrichment analyses on the full subset of DE genes, we also

performed enrichment analyses on each subset of genes representing a section in the Venn-

diagrams whenever it contained more than 25 genes, to functionally describe the overlap between

species and conditions.

Complementary to a GO enrichment analysis, we identified genes previously described as

related to CAM, starch metabolism and glycolysis in the same manner as described in Groot Crego

et al. 2024. Briefly, the extensive list of *A. comosus* gene ID's representing genes underlying

CAM, starch metabolism or glycolysis compiled by (2021) based on a diverse set of studies was

used to obtain the ID of *T. fasciculata* orthologous genes. This resulted in a total of 815 genes. The

genes from this subset that were differentially expressed in any of the comparisons were then

grouped by their overlap between species as indicated in the Venn diagrams.

The abundance of RNA reads for a select group of 74 CAM-related genes was plotted in a

heatmap using the R package ComplexHeatmap (Gu, Eils and Schlesner, 2016) for all 24 samples.

This subset includes 12 *T. fasciculata* orthologs of all pineapple genes putatively described as part

of the CAM carbon-fixation module (hereafter referred to as "Core CAM genes") (Ming *et al.*,

2015), regardless of their differential expression status. The remaining 62 genes include

aquaporins (5), circadian clock regulators (15), drought resistance genes (5), malate transporters

(2), photosystem-related genes (9), vacuolar proton pumps (13), genes related to stomatal

movement (7), vacuolar sugar invertases (3) and sugar transporters (3). Reads were normalised

with the TPM (Transcripts Per Million) method using total exonic length of each gene, and

converted to a log10 scale, then mean-centred with a z-score transformation. A separate heatmap

was made exclusively for homologs of the *PEPC* and *PPCK* genes.

3.8.　Gene trees of PPC1, PPC2 and PPCK

We identified homologs of the two subfamilies of PEPC, PPC1, PPC2 and of PPCK by

running BLASTP on the available gene models of *A. comosus* (V3), *Sorghum bicolor* (v3.1.1),

Oryza sativa (v7.0), *Zea mays* (RefGen v.4), *Asparagus officinalis* (V1.1) and *Arabidopsis*

thaliana (TAIR10) on Phytozome (https://phytozome-next.jgi.doe.gov/), using the known

Tillandsia homologs of pineapple PEPC and PPCK genes (Groot Crego *et al.*, 2024). Hits were
filtered based on identity and target length for each respective gene. For PPC1, we selected all
targets longer than 500 aa and with more than 80% identity, resulting in 56 protein sequences
across the seven species. For PPC2, the identity threshold was lowered to 70 %, resulting in eight
sequences. For PPCK, we selected all sequences with an identity greater than 60 % regardless of
target length, resulting in 17 sequences.

Protein sequences were aligned using mafft v.7.520 (Katoh *et al.*, 2002) with options *--op*
1.5 --ep 0.1 --maxiterate 1000 --localpair. Gene trees were constructed using IQTREE v.2.2.5
(Minh *et al.*, 2020) with options *-B 1000 -T AUTO*. The resulting consensus tree was used for
visualisation, collapsing branches leading to Poaceae genes.

4.   Results

4.1.   Titratable acidity measurements

In control conditions, titratable acidity displays a CAM-like diel curve in *T. vanhyningii*,
with an overnight accumulation of acids and depletion throughout the day (Figure 2A, Table S2).
Our LMM model shows TA fluctuates significantly between timepoints both under control and
drought conditions, yet there is no significant difference in TA values between watering regimes
at the same time point (Fig. S1, Table S3). In *T. leiboldiana*, TA does not appear to accumulate
overnight in control conditions, as differences between time points in the LMM model are not
significant. Under drought conditions, TA values between D+10 and N+10 are significantly
different, indicating a slight fluctuation in leaf acidity. *T. leiboldiana* exhibits a markedly different
temporal curve in TA accumulation between control and drought conditions, as the difference in
TA values between watering regimes at each time point is significantly different in all but one time
point (Table S3)t.

The maximum difference in TA (ΔTA) reflects the degree of acid accumulation over time
and is significantly different between species (ANOVA p-value = 0.00406, Table S3) but not
between watering regimes (p-value = 0.473). Overall, our analysis of TA indicates that diel
metabolism in the two species is substantially different, and so is their response to drought. While
T. vanhyningii's TA curve shows strong oscillation under control conditions, which is
characteristic for constitutive CAM, it appears nearly identical under drought except for an earlier
accumulation at N+4. *T. leiboldiana*'s TA curve shows no significant fluctuation under control

conditions, which is characteristic of a C3 plant, yet some diel fluctuation becomes apparent under
drought, indicating a weak, facultative CAM cycle.

321
Figure 2: Titratable Acidity (TA) measurements of T. vanhyningii *and* T. leiboldiana. ***A)*** *Titratable acidity*
measured at four time points across a diel cycle under two watering regimes. The significance level of
within-species, between-watering regime comparisons at a given time point from an LMM model are shown
*with following thresholds: ns - p-value > 0.05; * - p-value < 0.05; ** - p-value < 0.01; *** - p-value < 1E-*
03. ***B)*** *Maximum difference in titratable acidity across a diel cycle for both species in two watering regimes.*
The bar plots represent the average ΔTA measured across replicates, error bars represent the standard
deviation. The significance level from the ANOVA test is shown with the same thresholds as in A).

4.2. Differential gene expression

Mapping to the *T. fasciculata* reference genome was highly uniform between species, with
the median proportion of reads mapping uniquely being the same for *T. leiboldiana* and *T.*
vanhyningii (Fig. S2). After assigning RNA-seq reads to genes, we obtain on average 49,266,700
± 5,734,213 uniquely assigned reads per sample (Table S4). On average, 69 % ± 1 % of the full *T.*
fasciculata annotated gene set had non-zero counts per individual. The variance in read counts per
individual is largely explained by species-specific differences, as the first component of the PCA

separates samples clearly by species and explains 44.9 % of variance (Fig. S3), exceeding

differences in gene expression between day and night, or between watering regimes. Therefore,

DGE analyses were performed separately for each species.

339   DGE analyses reveal widely different counts of DE genes along the time and watering

regime axes. In both species, many genes are DE between day and night, regardless of watering

Figure 3: Counts of up- and down-regulated DE genes (adjusted p-value < 0.05, |FDR| ≥ 1.5) in four

comparisons of different conditions: diel differences in two watering regimes (first four bars), and

differences due to watering regimes at two timepoints (last four bars). The analyses were conducted in each

species separately. Up-regulation reflects increased expression at day compared to night for timewise

comparisons, and in drought compared to control conditions for comparisons of watering regimes.

regime (Fig. 3). However, *T. vanhyningii* consistently exhibits a closer resemblance in day-

versus-night activity compared to *T. leiboldiana*. The transcriptomic activity in *T. vanhyningii*

under control conditions is comparable between day and night, with only a 2.5 % increase in

upregulated genes at night than at day. In comparison, *T. leiboldiana* has 2.4x more genes

upregulated at day than at night in control conditions. This difference could reflect the more equal

task division between day and night in CAM plants compared to plants with a C3-like metabolism,

where most of the photosynthetic apparatus operates during the day. Under drought conditions,

transcriptomic activity shifts towards the night in both species. While this pattern is noticeable in

T. vanhyningii, with 58 % of DE genes upregulated at night under drought compared to 51 % in

control, it is especially pronounced in *T. leiboldiana*, where 47 % of genes is upregulated at night

under drought compared to 30 % in control.

In comparison with time-wise analyses, much fewer genes are DE between watering

regimes, except in *T. leiboldiana* at night. Here, *T. vanhyningii* appears to show a particularly

similar expression between the watering regimes independent on the time point, as opposed to *T.*

leiboldiana. DE gene counts are low for *T. vanhyningii* both at day and at night, suggesting that

most temporal expression patterns detected in the time-wise DGE analysis are constitutive.

Compared to *T. vanhyningii*, *T. leiboldiana* manifests a much more noticeable response to

drought, especially at night. During the day, transcriptomic activity seems decreased under

drought, as only 20 % of DE genes are upregulated. The total number of DE genes between

watering conditions at night is 2.8x larger than at day, which is a contrary trend to *T. vanhyningii*.

4.3.   Overlap in transcriptomic response

The hypergeometric test revealed that all detected overlaps between species, time points

and watering regimes are larger than expected by random chance, indicating that the same

biological processes were captured between species and watering regimes by our RNA-seq

experiment (Table S5). The timewise transcriptomic response has a large overlap between species

and watering regimes, with 53 % of all DE genes being unique to one species and watering regime

(Fig. 4A). While this is to be expected given all plants experience day-night changes, only 5.9 %

of all timewise DE genes are shared between all groups. The largest overlap is found within species

between conditions, as only 19 % of all time-wise DE genes are shared between species (Table

S6). The overlap in timewise DE genes between watering regimes is greater in *T. vanhyningii* (44

%) than in *T. leiboldiana* (35 %). Conversely, the two species share more timewise DE genes under drought conditions (17 %) than in control conditions (14 %).

In comparison, the response to watering regime is largely non-shared between groups, with 77 % of all DE genes being unique to one species and timepoint and only two DE genes being shared among all groups (Fig. 4B). *T. vanhyningii* shows little overlap between time points in the response to drought (8 %), while in *T. leiboldiana* 22 % of genes are DE between watering regimes at both timepoints (Table S6). Due to the small number of DE genes in *T. vanhyningii* between watering regimes, the overlap between species is negligibly small (2.4 %) in proportion to the number of DE genes called in *T. leiboldiana*.

In summary, both the overall transcriptomic response to day-night changes and to watering regimes shows little overlap between species, yet diel expression between species seems to overlap slightly more under drought than under control, indicating an increase in the resemblance of the two species' metabolism under drought stress.

4.4. CAM-related transcriptomic overlap and differences between species, timepoints and conditions

4.4.1. Overlap between species

A CAM-related change in expression is likely to be associated both with differential expression between time points as between watering regimes in *T. leiboldiana*, which seems to activate a CAM cycle under drought conditions. When considering all DE genes summed up across species, time points and watering regimes, 582 were DE in at least one of the two timewise DE analyses across species and in at least one of the two watering-regime DE analyses. This number represents 25 % of all genes with time-structured activity in at least one species, 43 % of all genes that are DE between watering conditions in at least one species and 17 % of all DE genes combined.

Of these 582 genes, the largest group is DE in *T. leiboldiana* for both time and watering regime, but not DE in *T. vanhyningii* (Fig. 5). The second largest group consists of 151 genes (26 %) which display a time-structured activity in *T. vanhyningii* while being DE between watering regimes in *T. leiboldiana*, and includes nine previously described CAM-related genes, among which the sugar transporter SWEET14. 137 genes (24 %) show a time-structured expression in both species but are only DE between watering conditions in *T. leiboldiana*. Among these genes

A

B

were 16 previously described CAM-related genes, yet only four were upregulated at drought in *T. leiboldiana*, including core CAM gene encoding for PEPC kinase (*PPCK2*), a drought resistance gene YLS9 (Müller *et al.*, 2017), and two genes encoding enzymes associated with glycolysis and gluconeogenesis. This shows that the overlap of CAM-related response to drought between species is limited mostly to core CAM genes.

We also searched for CAM-related genes among the larger set of DE genes called across time and watering regime separately. The set of timewise DE genes contained 101 and 86 CAM-related genes in *T. vanhyningii* and *T. leiboldiana* respectively, and reveal a myriad of underlying functions such as circadian clock, glycolysis, stomatal movement, drought resistance, and water and sugar transport (Table S7). Overlap between species and watering regimes of these genes follows approximately the same trend as the full set of DE genes, with 49 % of genes being unique to a given species and watering regime (Table S6). The twelve CAM-related genes that are differentially expressed between time points in all groups are related to the circadian clock, photosynthesis, and photosystem activity, which are expected to constitutively show diel expression in plants. In *T. vanhyningii,* 45 % of CAM timewise DE genes are shared between watering conditions. On the other hand, *T. leiboldiana* only shares 36 % of genes between watering conditions, showing a clearer shift in photosynthetic metabolism when drought-stressed. The proportion of timewise CAM DE genes shared between species is similar in both watering regimes (21 % at control and 22 % at drought), which is slightly higher than across all DE genes, but remains a minority of the total set of CAM-related DE genes.

Figure 5: Overview of the overlap between DE analyses across time points and across watering regimes.
The bar plots represent the number of genes overlapping between DE analyses, sorted in descending order.
Below, the linked dots represent the analyses which display each overlap. The vertically oriented bar plot
on the left-side shows the total number of DE genes called in each analysis. DE analyses were performed
between time points and watering regimes in each species (VAN = T. vanhyningii*, LEI =* T. leiboldiana*)*

For DE genes between watering regimes, seven and 78 CAM-related genes were present in *T. vanhyningii* and *T. leiboldiana* respectively (Table S6, Table S8). In *T. vanhyningii*, only one gene was shared between timepoints, whereas in *T. leiboldiana* 28 % overlapped between timepoints. Only one gene overlapped between the species, namely a gene encoding for a probable linoleate 9S-lipoxygenase enzyme, which was found differentially expressed between C3 and CAM *Tillandsia* species in a previous study (De La Harpe *et al.*, 2020). Lipoxygenases are involved in the biosynthesis of jasmonates, which play an important role in response to various biotic and abiotic stressors (Bachmann *et al.*, 2002).

4.4.2. CAM-related expression in *Tillandsia vanhyningii*

In control conditions, *T. vanhyningii* shows timewise differential expression of core CAM gene PPCK and of genes underlying CAM-related functions such as stomatal movement, vacuolar proton pumps, malate transport, sugar transport and glycolysis (Table S7). Several genes encoding for Late Embryogenesis Abundant (LEA) proteins , which are linked to drought tolerance, also show diel expression in control conditions, for example.

The set of CAM-related functions showing diel expression in drought conditions is similar to that of control conditions and has an overlap of 46 genes, including PPCK, circadian clock regulator REVEILLE1, a beta-amylase (starch breakdown), and all LEA proteins that are DE in control. AVP1, a secondary proton pump that aids in the vacuolar transport of malate, is only DE in control conditions, whereas a malate dehydrogenase is exclusively showing diel expression in drought conditions. While sugar transporter SWEET16 shows diel expression exclusively under control conditions, SWEET14 is exclusively DE in drought conditions. An additional set of LEA proteins also become timewise DE in drought conditions. These findings suggest that, while the CAM pathway of *T. vanhyningii* remains largely similar regardless of watering conditions, different genes underlying the same functions are being co-opted under specific circumstances.

Very few CAM-related genes were DE in *T. vanhyningii* between watering regimes. They include an aquaporin PIP2-4 (water transport) and several enzymes associated with glycolysis and gluconeogenesis. None of these genes are part of the core CAM pathway, further showing the lack of response to drought in the constitutive CAM species regarding its photosynthetic metabolism.

4.4.3. CAM-related expression in *Tillandsia leiboldiana*

In *T. leiboldiana* at control conditions, identified CAM-related DE genes underlie functions that are typically also expected to fluctuate across time in C3 plants (glycolysis and gluconeogenesis, sugar transport, circadian clock regulation). Beta-carbonic anhydrase 2 is timewise DE in control conditions, but is upregulated during the day (Fig. S4), which is contrary to the expected nighttime activity in CAM. Under drought conditions, 31 DE genes overlap with control conditions (Table S7), yet many additional CAM-related functions become timewise DE under drought conditions, such as malate transport, stomatal movement, sugar metabolism, water transport and starch metabolism. Genes encoding Alpha-carbonic anhydrase 1 (*α-CA*) and phosphoenolpyruvate carboxylase kinase (*PPCK2*) are upregulated at night, along with circadian clock regulator XAP5 CIRCADIAN TIMEKEEPER (XCT), which has been detected as

differentially expressed between CAM and C3-like *Tillandsia* in several studies (De La Harpe *et*
al., 2020; Groot Crego *et al.*, 2024). These findings further emphasise that a CAM cycle is likely
activated in *T. leiboldiana* under drought conditions.

The comparison of gene expression between watering regimes at a given time point
confirms this finding even further. Most CAM-related genes that are DE between watering regimes
can be found in *T. leiboldiana* at night, and include the core CAM genes phosphoenolpyruvate
carboxylase (*PEPC*) and *PPCK* (Table S8).

4.4.4.   Diel expression of PPC and PPCK homologs between species and
watering regimes

The transcript abundances of PEPC and PPCK homologs reveal a variety of expression
patterns. Gene *Tfasc_v1.16311-RA*, which is the ortholog of *PPC1-M1* (Fig. S5A) and was
previously identified as the CAM-related fluctuating copy of PEPC in *T. fasciculata* (Groot Crego
et al., 2024) does not show clear diel expression changes in *T. vanhyningii* (Fig. 6). In *T.*
leiboldiana a slight increase is noticeable at night but it is not significant in either condition.
This could be explained by the time of sampling not being synchronous with the peaking and
troughing times of PEPC, and aligns with other core CAM enzymes not being timewise DE in this
experiment (Fig. S3). However, the enzyme is noticeably more highly expressed in *T. vanhyningii*
compared to *T. leiboldiana*. The remaining known *PEPC* genes are generally less expressed than
PPC1-M1 in both species, but appear to become activated under drought conditions, namely in *T.*
vanhyningii for *PPC1-M2* (not significant), and in *T. leiboldiana* for *PPC2*.

For PEPC kinase (PPCK), diel expression stands out in one gene (*PPCK2,* Fig. S6) in *T.*
vanhyningii under both watering regimes. While not strongly expressed at night in control
conditions in *T. leiboldiana*, the gene copy becomes highly expressed under drought conditions,
which is perhaps the most direct indication of an activated CAM cycle in *T. leiboldiana* under
drought. *PPCK1* shows mild diel fluctuation in *T. vanhyningii* but is overall lowly expressed in *T.*
leiboldiana.

Figure 6: Transcript abundance in "Transcripts per Million" (TPM) for all homologous genes encoding for PEPC and PEPC kinase (PPCK) in T. vanhyningii *(left panel) and* T. leiboldiana *(right panel) at two timepoints and watering regimes. TPM values are transformed to a logarithmic scale. The colour of gene names reflects their differential expression status: blue = DE in Drought vs Control comparisons, red is DE in both comparisons and black = not DE. If a gene is DE, the species initial in which it was detected is mentioned in superscript behind its name.*

4.5. Description of the general response to drought

As described above, the quantitative response to drought is far greater in *T. leiboldiana* than in *T. vanhyningii* (Fig. 3), suggesting suggests that *T. leiboldiana*'s metabolism is being severely affected by drought, while *T. vanhyningii*'s seems largely the same between conditions. This demonstrates inherently the drought-tolerant nature of a constitutive CAM plant and contrasts it to the marked stress-response of a closely related weak CAM plant.

GO term enrichment of gene sets that are DE in *T. leiboldiana* between watering regimes further point at a drought response underlying the large-scale expression changes observed in this

species (Fig. 7A), with genes involved in the response to water deprivation and to the stress-hormone abscisic acid showing the strongest overall up-regulation at drought of all DE genes. Processes that are strongly under-regulated during drought reveal a general shutdown of metabolism, such as photosynthesis, transport, and biosynthetic processes.

Figure 7: Summary of significantly enriched GO terms representing a Biological Process (BP) in DE genes between watering regimes in T. leiboldiana *(A) and* T. vanhyningii *(B). GO term enrichment at day and night were combined for each bar plot. The bars show the significance of each enriched GO term as the p-value adjusted for multiple-testing, on a logarithmic scale. The colour gradient shows the degree of overall up- and down-regulation of the genes underlying each GO term, as a z-score of the fold-change on a*

 logarithmic scale. The GO terms are shown in descending order from most up-regulated in drought
conditions to most up-regulated in control conditions.

This contrasts strongly with enriched GO terms found among the few DE genes in *T.*
vanhyningii, where a response to drought is not immediately apparent - photosynthesis does not
seem to be regulated differently between watering regimes and no orchestrated response to water
deprivation is apparent (Fig. 7B). However, a few enriched GO terms do suggest a limited response
to drought, such as metabolism of inositol, jasmonate and molybdate.

5.   **Discussion**

Crassulacean Acid Metabolism is associated with diversification in many plant families,
and described as a key innovation trait in Bromeliaceae (Benzing, 2000; Ogburn and Edwards,
2009; Silvera *et al.*, 2009; Quezada and Gianoli, 2011), yet the genetic architecture of the trait
itself, its degree of redundancy and genetic overlap between different forms, and its evolutionary
trajectory are not well understood (Edwards, 2023). The subgenus *Tillandsia* exhibits a broad
range of CAM phenotypes reflected in carbon isotope values (Crayn *et al.*, 2015; De La Harpe *et*
al., 2020) and is an excellent system for studying CAM evolution, but the exact range of
phenotypes within the subgenus has not been characterised beyond carbon isotope ratios and
transcriptomic studies under control conditions. In this study, we compared physiological and
transcriptomic responses to drought between two species with some of the most extreme reported
carbon isotope ratios within the subgenus to further characterise the phenotypic range in the group
and elucidate the degree of shared transcriptomic response underlying two different CAM
phenotypes and their response to drought.

551   ***Tillandsia leiboldiana* is a weak CAM plant**

Both our time-wise titratable acidity measurements and differential gene expression
analysis indicate the presence of a latent CAM cycle in *T. leiboldiana* which becomes activated
under severe drought stress. The most direct evidence for this can be found in the nighttime
activation of *PPCK2* in *T. leiboldiana* in drought conditions, while it does not significantly
fluctuate over time in control conditions. Reported carbon isotope ratios for *T. leiboldiana* are well
within the C3 range ($\delta^{13}C$ = -28.0 / -31.3; Caryn et al. (2015), De La Harpe et al. (2020)), indicating
yet again the importance of characterising photosynthetic phenotypes at the hand of multiple

measurements performed in various environmental and developmental conditions. Instead of being a strong C3 plant as the reported carbon isotope ratios suggest, *T. leiboldiana* is much more likely to be a weak, facultative CAM plant performing C3 photosynthesis in optimal conditions but with pre-existing CAM-like expression patterns in certain genes (Groot Crego *et al.*, 2024) and a weak overnight accumulation of malate under drought conditions.

Due to a lack of gas exchange measurements, we can't yet establish whether *T. leiboldiana* performs CAM cycling, i.e. malate production and storage starting from CO_2 accumulated during the day, or instead, assimilates CO_2 overnight. The presence of several genes associated with stomatal function among differentially expressed genes between watering regimes in *T. leiboldiana* suggests a change in stomatal behaviour at drought conditions. However, this could be a general response to drought rather than a CAM-specific response. For example, TRAB1, a member of the abscisic acid signalling pathway, which is generally activated under drought stress (Hobo, Kowyama and Hattori, 1999; Zhu, 2002), is upregulated in *T. leiboldiana* under drought conditions at night (Fig. S4). On the other hand, the guard cell anion channel SLAC1, which controls stomatal closure (Vahisalu *et al.*, 2008), is down-regulated at night in *T. leiboldiana* under drought (Fig. S4), a potential indication that stomata open at night. Nonetheless, gas exchange measurements are needed to fully understand stomatal activity and timing of CO_2 absorption in *T. leiboldiana*.

Response to drought is minimal in *Tillandsia vanhyningii*

Our physiological and transcriptomic analyses indicate very limited functional changes in *T. vanhyningii* under drought conditions, especially compared to *T. leiboldiana*. Titratable acidity is accumulated in a very similar manner among watering regimes, with the exception that accumulation appears to occur faster overnight under drought conditions. This agrees with previous studies on sister species *T. ionantha*, which showed that its physiology was little affected even after six months of drought, and attributed its ability to maintain leaf water levels to day-time closing of stomata, low stomatal densities and its enhanced ability to absorb water from the atmosphere thanks to a dense layer of absorptive trichomes, (Nowak and Martin, 1997; Ohrui *et al.*, 2007).

A very small number of genes were found to be DE between watering regimes in *T. vanhyningii*, while a major response was detected in *T. leiboldiana*. Upregulated enriched GO

terms at drought in *T. vanhyningii* included metabolism of inositol, jasmonate, molybdate and

proline, all compounds known to play a role in drought response (Loewus and Murthy, 2000; Kim

et al., 2004; Wasternack, 2014). This indicates that while *T. vanhyningii* shows minor signs of

response to drought, it is able to maintain most of its metabolism stable under drought stress, which

is not the case for *T. leiboldiana*.

These findings further support the view that "atmospheric" or "grey" *Tillandsia* such as *T.*

vanhyningii are drought-adapted, which has been shown in physiological studies (Martin, 1994;

Stiles and Martin, 1996; Nowak *et al.*, 1997; Castillo, Cervera and Navarro-Alberto, 2016), but to

our knowledge not on a broad transcriptomic or genomic level. Atmospheric *Tillandsia* represent

the most extreme forms of adaptation to an epiphytic lifestyle among Bromeliaceae, having lost

almost entirely the ability to absorb water with roots (Benzing, 2000; Crayn, Winter and Smith,

2004). They occur in arid habitats and are all constitutive CAM plants, leading to the suggestion

that CAM is a key innovation trait in this group (Crayn, Winter and Smith, 2004). Our findings

show indeed that *T. vanhyningii* is only mildly affected by drought conditions and points at

constitutive drought tolerance on the transcriptomic and physiological level in atmospheric

Tillandsia.

Both response to drought and photosynthetic metabolism show little overlap between

species on the transcriptomic level

T. vanhyningii and *T. leiboldiana* are closely related members of a recent radiation with an

estimated maximum divergence time of 3.22 Mya (Barfuss *et al.*, 2016; Mendoza *et al.*, 2017;

Yardeni *et al.*, 2023). Despite the relatively short time in which these species have diverged, our

analyses demonstrate clear differentiation on the physiological and transcriptomic level, resulting

in distinct photosynthetic phenotypes. On the transcriptomic level, both the set of time-structured

genes and the set of genes that are DE between watering regimes show little overlap between

species, which has also been reported in other CAM lineages such as Agavoideae (Heyduk,

McAssey and Leebens-Mack, 2022).

The majority of time-structured genes previously described as CAM-related are uniquely

time-structured within one species. This suggests that, despite *T. leiboldiana* appearing to activate

CAM under drought conditions, different sets of genes underlie this phenotype compared to *T.*

vanhyningii. A few notable exceptions of CAM-related genes that are shared between *T.*

vanhyningii and drought-stressed *T. leiboldiana* are *PPCK2* (Fig. 6), tonoplast dicarboxylate transporter (malate transport) and protein YLS9 (drought resistance). The minor overlap in the facultatively activated CAM cycle of *T. leiboldiana* and the constitutive cycle of *T. vanhyningii* suggests a strong distinction in the transcriptomic basis leading to the two forms of CAM, rather than a shared pathway which is expressed at lower levels in the weak CAM plant.

This is even more strongly supported by the distinct expression patterns of different gene family members of PEPC. Under drought conditions, the homologous copy of *PPC2* appears to be up-regulated in *T. leiboldiana*, while being lowly expressed at all conditions in *T. vanhyningii*. Because none of the PEPC copies are time-structured in this experiment in any species or condition, likely due to the time of sampling not being synchronous with the peak activity times of this enzyme, it is difficult to confirm that *PPC2* is assimilated in CAM in *T. leiboldiana* under drought conditions. While in the majority of CAM plant lineages, including *T. fasciculata* (Groot Crego *et al.*, 2024) and *A. comosus* (Ming *et al.*, 2015), the PEPC gene copy recruited into the CAM pathway is *PPC1*, recruitment of *PPC2* has been suggested in *Yucca aloifolia* (Heyduk, McAssey and Leebens-Mack, 2022) and in *Isoetes taiwanensis* (Wickell *et al.*, 2021).

Conclusion

Taken together, this study characterises in detail the photosynthetic phenotypes of two species of the subgenus *Tillandsia*, and illustrates that the species with one of the most C3-like carbon isotope value recorded in the group so far is not a C3 plant, but a weak CAM intermediate. While our findings suggest that the subgenus *Tillandsia* likely doesn't contain candidate C3 species, and therefore doesn't cover the full range of existing phenotypes in the C3-CAM continuum known from other plant groups, our analyses also demonstrate an important distinction on the transcriptomic level between weak and constitutive CAM. While on the phenotypic level, both plants exhibit a similar phenotype at different rates, we find little overlap in the underlying transcriptomic response. This highlights the polygenic nature of CAM and provides a hypothesis for the high rate at which this trait has evolved independently across the plant kingdom; namely the presence of multiple genetic routes to achieve similar phenotypes. Our findings also put in question the idea of CAM being a trait with a single transcriptomic basis expressed to different degrees in the phenotypic continuum between weak and strong CAM phenotypes. There is still a broad diversity of intermediate CAM phenotypes and genotypes to be characterised in further

detail, and the subgenus *Tillandsia* with its wide range of intermediate CAM species forms an excellent candidate group for future exciting research into the CAM continuum and its evolution.

6. Acknowledgements

This work was financially supported by the Austrian Science Fund (FWF) with grant https://doi.org/10.55776/W1225 to a faculty team including C.L. and O.P., and grant https://doi.org/10.55776/P35275 to O.P. Additionally, this work was funded by the professorship start-up grant of Christian Lexer at the University of Vienna BE772002. We thank Viktor Vagovic at the Core Facility Botanical Garden of the University of Vienna for caretaking and maintenance of the plant accessions used in this study and assistance with the drought experiment. We also thank Aglaia Szukala, Marta Pelizzola, Nina Flowers and Simon Reynaert for advice and discussion on transcriptome analysis.

7. Competing interests

The authors declare no conflict of interests.

8. Author Contributions

This study was conceived by CL, JH, MDH, OP, and CGC. The greenhouse experiment and sampling was conducted by MDH and MHJB. Laboratory work including titratable acidity measurements and DNA extraction and library prep was conducted by MDH, OP and MHJB. Analyses were performed by CGC, MDH and SS. The manuscript was primarily written by CGC with contributions from SS, and amended following the dedicated reading and feedback of all co-authors.

9. Data availability

Titratable acidity measurements, detailed information on statistical testing and accession information is provided in the supplementary tables of this manuscript. The raw RNA-seq reads will be deposited on the NCBI SRA database upon publication of this manuscript. Scripts, read counts and extended tables of GO term enrichment analyses will be made available on the following repository: https://github.com/cgrootcrego/Tillandsia_CAM-Drought.

10. References

681 682 683 Alexa, A., Rahnenführer, J. and Lengauer, T. (2006) "Improved scoring of functional groups from gene expression data by decorrelating GO graph structure," *Bioinformatics* , 22(13), pp. 1600–1607.

685 Bachmann, A. *et al.* (2002) "Jasmonate-induced lipid peroxidation in barley leaves initiated by distinct 13-LOX forms of chloroplasts," *Biological chemistry*, 383(10), pp. 1645–1657.

687 688 Barfuss, M. H. J. *et al.* (2016) "Taxonomic revision of bromeliaceae subfam. Tillandsioideae based on a multi-locus DNA sequence phylogeny and morphology," *Phytotaxa*, 279(1), pp. 1–97.

690 Barghi, N. *et al.* (2019) "Genetic redundancy fuels polygenic adaptation in Drosophila," *PLoS biology*, 17(2), p. e3000128.

692 Barghi, N., Hermisson, J. and Schlötterer, C. (2020) "Polygenic adaptation: a unifying framework to understand positive selection," *Nature reviews. Genetics*, 21(12), pp. 769–781.

694 Bates, D. *et al.* (2015) "Fitting Linear Mixed-Effects Models Using lme4," *Journal of statistical software*, 67, pp. 1–48.

696 Benzing, D. H. (2000) *Bromeliaceae: Profile of an Adaptive Radiation*. Cambridge University Press.

698 699 Beutelspacher, C. R. and García-Martínez, R. (2021) "Descripción de nuevos híbridos y una nueva combinación en Tillandsia (Bromeliaceae) para Chiapas, México," *Lacandonia*, 15(2), pp. 19–28.

701 Borland, A. M. *et al.* (2014) "Engineering crassulacean acid metabolism to improve water-use efficiency," *Trends in plant science*, 19(5), pp. 327–338.

703 704 Castillo, R. J., Cervera, J. C. and Navarro-Alberto, J. (2016) "Drought and extreme temperature tolerance for Tillandsia dasyliriifolia, an epiphytic bromeliad from the northern coastal dune scrubland in Yucatan, Mexico," *Botanical Sciences*, 94(1), pp. 121–126.

706 707 Chen, H. and Boutros, P. C. (2011) "VennDiagram: a package for the generation of highly-customizable Venn and Euler diagrams in R," *BMC bioinformatics*. bmcbioinformatics.biomedcentral …, 12, p. 35.

709 710 711 Crayn, D. M. *et al.* (2015) "Photosynthetic pathways in Bromeliaceae: Phylogenetic and ecological significance of CAM and C3 based on carbon isotope ratios for 1893 species," *Botanical journal of the Linnean Society. Linnean Society of London*. Blackwell Publishing Ltd, 178(2), pp. 169–221.

713 714 Crayn, D. M., Winter, K. and Smith, J. A. C. (2004) "Multiple origins of crassulacean acid metabolism and the epiphytic habit in the Neotropical family Bromeliaceae," *Proceedings of the National Academy of Sciences*. National Academy of Sciences, 101(10), pp. 3703–3708.

De La Harpe, M. *et al.* (2020) "Genomic footprints of repeated evolution of CAM photosynthesis in a Neotropical species radiation," *Plant, cell & environment*, 43(12), pp. 2987–3001.

Dobin, A. *et al.* (2013) "STAR: Ultrafast universal RNA-seq aligner," *Bioinformatics* . Oxford Academic, 29(1), pp. 15–21.

Edwards, E. J. (2023) "Reconciling continuous and discrete models of C4 and CAM evolution," *Annals of botany*. doi: 10.1093/aob/mcad125.

Ewels, P. *et al.* (2016) "MultiQC: summarize analysis results for multiple tools and samples in a single report," *Bioinformatics* , 32(19), pp. 3047–3048.

Gibson, A. C. (1982) "The anatomy of succulence," *Crassulacean acid metabolism*. American Society of Plant Physiologists Rockville, MD, USA, pp. 1–17.

Gilman, I. S. *et al.* (2023) "The CAM lineages of planet Earth," *Annals of botany*, 132(4), pp. 627–654.

Givnish, T. J. *et al.* (2014) "Adaptive radiation, correlated and contingent evolution, and net species diversification in Bromeliaceae," *Molecular phylogenetics and evolution*. Academic Press, 71(1), pp. 55–78.

Goldstein, D. B. and Holsinger, K. E. (1992) "Maintenance of polygenic variation in spatially structured populations: roles for local mating and genetic redundancy," *Evolution; international journal of organic evolution*, 46(2), pp. 412–429.

Griffiths, H. *et al.* (2008) "Leaf succulence determines the interplay between carboxylase systems and light use during Crassulacean acid metabolism in Kalanchoë species*," *Journal of experimental botany*. Oxford Academic, 59(7), pp. 1851–1861.

Groot Crego, C. *et al.* (2024) "CAM evolution is associated with gene family expansion in an explosive bromeliad radiation," *The Plant cell*. doi: 10.1093/plcell/koae130.

Gu, Z., Eils, R. and Schlesner, M. (2016) "Complex heatmaps reveal patterns and correlations in multidimensional genomic data," *Bioinformatics* , 32(18), pp. 2847–2849.

Hämälä, T. *et al.* (2020) "Gene Expression Modularity Reveals Footprints of Polygenic Adaptation in Theobroma cacao," *Molecular biology and evolution*, 37(1), pp. 110–123.

Heyduk, K. *et al.* (2016) "Gas exchange and leaf anatomy of a C3-CAM hybrid, Yucca gloriosa (Asparagaceae)," *Journal of experimental botany*, 67(5), pp. 1369–1379.

Heyduk, K. *et al.* (2018) "Shifts in gene expression profiles are associated with weak and strong Crassulacean acid metabolism," *American journal of botany*, 105(3), pp. 587–601.

Heyduk, K., McAssey, E. V. and Leebens-Mack, J. (2022) "Differential timing of gene expression and recruitment in independent origins of CAM in the Agavoideae (Asparagaceae),"

The New phytologist, 235(5), pp. 2111–2126.

Heyduk, K., Ray, J. N. and Leebens-Mack, J. (2021) "Leaf anatomy is not correlated to CAM
function in a C3+CAM hybrid species, Yucca gloriosa," *Annals of botany*, 127(4), pp. 437–449.

Hobo, T., Kowyama, Y. and Hattori, T. (1999) "A bZIP factor, TRAB1, interacts with VP1 and
mediates abscisic acid-induced transcription," *Proceedings of the National Academy of Sciences*
of the United States of America, 96(26), pp. 15348–15353.

Katoh, K. *et al.* (2002) "MAFFT: a novel method for rapid multiple sequence alignment based
on fast Fourier transform," *Nucleic acids research*, 30(14), pp. 3059–3066.

Kim, T.-H. *et al.* (2004) "De novo protein synthesis in relation to ammonia and proline
accumulation in water stressed white clover," *Functional plant biology: FPB*, 31(8), pp. 847–
855.

Kuznetsova, A., Brockhoff, P. B. and Christensen, R. H. B. (2017) "lmerTest Package: Tests in
Linear Mixed Effects Models," *Journal of statistical software*, 82, pp. 1–26.

Láruson, Á. J., Yeaman, S. and Lotterhos, K. E. (2020) "The Importance of Genetic Redundancy
in Evolution," *Trends in ecology & evolution*, 35(9), pp. 809–822.

Liao, Y., Smyth, G. K. and Shi, W. (2014) "FeatureCounts: An efficient general purpose
program for assigning sequence reads to genomic features," *Bioinformatics* . Oxford University
Press, 30(7), pp. 923–930.

Loewus, F. A. and Murthy, P. P. N. (2000) "myo-Inositol metabolism in plants," *Plant science:*
an international journal of experimental plant biology, 150(1), pp. 1–19.

Males, J. (2018) "Concerted anatomical change associated with crassulacean acid metabolism in
the Bromeliaceae," *Functional plant biology: FPB*, 45(7), pp. 681–695.

Martin, C. E. (1994) "Physiological ecology of the Bromeliaceae," *The Botanical review;*
interpreting botanical progress, 60(1), pp. 1–82.

Martin, C. E., Hsu, R. C.-C. and Lin, T.-C. (2009) "The relationship between CAM and leaf
succulence in two epiphytic vines, Hoya carnosa and Dischidia formosana (Asclepiadaceae), in a
subtropical rainforest in northeastern Taiwan," *Photosynthetica*, 47(3), pp. 445–450.

Mendoza, C. G. *et al.* (2017) "Geographic structure in two highly diverse lineages of Tillandsia
(Bromeliaceae)," *Botany*, 95(7), pp. 641–651.

Ming, R. *et al.* (2015) "The pineapple genome and the evolution of CAM photosynthesis,"
Nature genetics. Nature Publishing Group, 47(12), pp. 1435–1442.

Minh, B. Q. *et al.* (2020) "IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era," *Molecular biology and evolution*, 37(5), pp. 1530–1534.

Müller, M. *et al.* (2017) "De novo transcriptome assembly and analysis of differential gene
expression in response to drought in European beech," *PloS one*, 12(9), p. e0184167.

Nelson, E. A. and Sage, R. F. (2008) "Functional constraints of CAM leaf anatomy: tight cell
packing is associated with increased CAM function across a gradient of CAM expression,"
Journal of experimental botany, 59(7), pp. 1841–1850.

Nelson, E. A., Sage, T. L. and Sage, R. F. (2005) "Functional leaf anatomy of plants with
crassulacean acid metabolism," *Functional plant biology: FPB*, 32(5), pp. 409–419.

Nowak, E. J. and Martin, C. E. (1997) "Physiological and Anatomical Responses to Water
Deficits in the Cam Epiphyte Tillandsia ionantha (Bromeliaceae)," *International journal of plant*
sciences. The University of Chicago Press, 158(6), pp. 818–826.

Nowak, M. A. *et al.* (1997) "Evolution of genetic redundancy," *Nature*, 388(6638), pp. 167–171.

Ogburn, R. M. and Edwards, E. J. (2009) "Anatomical variation in Cactaceae and relatives: Trait
lability and evolutionary innovation," *American journal of botany*, 96(2), pp. 391–408.

Ohrui, T. *et al.* (2007) "Foliar trichome- and aquaporin-aided water uptake in a drought-resistant
epiphyte Tillandsia ionantha Planchon," *Planta*, 227(1), pp. 47–56.

Osmond, C. B. (1978) "Crassulacean Acid Metabolism: A Curiosity in Context," *Annual review*
of plant physiology. Annual Reviews, 29(1), pp. 379–414.

Pierce, S., Winter, K. and Griffiths, H. (2002) "Carbon isotope ratio and the extent of daily CAM
use by Bromeliaceae," *The New phytologist*. Wiley, 156(1), pp. 75–83.

Quezada, I. M. and Gianoli, E. (2011) "Crassulacean acid metabolism photosynthesis in
Bromeliaceae: an evolutionary key innovation," *Biological journal of the Linnean Society.*
Linnean Society of London. Oxford Academic, 104(2), pp. 480–486.

Robinson, M. D., McCarthy, D. J. and Smyth, G. K. (2009) "edgeR: A Bioconductor package for
differential expression analysis of digital gene expression data," *Bioinformatics* . Oxford
University Press, 26(1), pp. 139–140.

Robinson, M. D. and Oshlack, A. (2010) "A scaling normalization method for differential
expression analysis of RNA-seq data," *Genome biology*, 11(3), p. R25.

Schubert, M., Lindgreen, S. and Orlando, L. (2016) "AdapterRemoval v2: rapid adapter
trimming, identification, and read merging," *BMC research notes*. BioMed Central, 9(1), p. 88.

Silvera, K. *et al.* (2009) "Crassulacean acid metabolism and epiphytism linked to adaptive
radiations in the Orchidaceae," *Plant physiology*, 149(4), pp. 1838–1847.

Silvera, K., Santiago, L. S. and Winter, K. (2005) "Distribution of crassulacean acid metabolism
in orchids of Panama: evidence of selection for weak and strong modes," *Functional plant*
biology: FPB, 32(5), pp. 397–407.

Steiner, C. C. *et al.* (2009) "The genetic basis of phenotypic convergence in beach mice: similar
pigment patterns but different genes," *Molecular biology and evolution*, 26(1), pp. 35–45.

Stiles, K. C. and Martin, C. E. (1996) "Effects of drought stress on CO2 exchange and water
relations in the CAM epiphyte Tillandsia utriculata (Bromeliaceae)," *Journal of plant*
physiology, 149(6), pp. 721–728.

Vahisalu, T. *et al.* (2008) "SLAC1 is required for plant guard cell S-type anion channel function
in stomatal signalling," *Nature*, 452(7186), pp. 487–491.

Wang, M., Zhao, Y. and Zhang, B. (2015) "Efficient Test and Visualization of Multi-Set
Intersections," *Scientific reports*, 5, p. 16923.

Wasternack, C. (2014) "Action of jasmonates in plant stress responses and development--applied
aspects," *Biotechnology advances*, 32(1), pp. 31–39.

Wickell, D. *et al.* (2021) "Underwater CAM photosynthesis elucidated by Isoetes genome,"
Nature Communications 2021 12:1. Nature Publishing Group, 12(1), pp. 1–13.

Winter, K. and Holtum, J. A. M. (2002) "How closely do the delta(13)C values of Crassulacean
Acid metabolism plants reflect the proportion of CO(2) fixed during day and night?," *Plant*
physiology, 129(4), pp. 1843–1851.

Yardeni, G. *et al.* (2021) "Taxon-specific or universal? Using target capture to study the
evolutionary history of rapid radiations," *Molecular ecology resources*. doi: 10.1111/1755-
0998.13523.

Yardeni, G. *et al.* (2023) "The explosive radiation of the Neotropical Tillandsia subgenus
Tillandsia (Bromeliaceae) has been facilitated by pervasive hybridization," *bioRxiv*. doi:
10.1101/2023.11.16.567341.

Yeaman, S. *et al.* (2016) "Convergent local adaptation to climate in distantly related conifers,"
Science, 353(6306), pp. 1431–1433.

Zhu, J.-K. (2002) "Salt and drought stress signal transduction in plants," *Annual review of plant*
biology, 53, pp. 247–273.

11.   Supporting information

Figure S1: Diagnostic plots for Linear Mixed-Effects Models constructed in each species to assess

the effect of time and watering regime on titratable acidity measurements.

Figure S2: Mapping rates of across 24 samples of different species and watering regimes.

Figure S3: Principal component analysis of TMM-normalised read counts in log(CPM).

Figure S4: Transcript abundance in "Transcripts per Million" (TPM) for a subset of CAM-related

genes in T. vanhyningii (left panel) and T. leiboldiana (right panel) at two timepoints and watering

regimes.

Figure S5: Gene trees of PPC1 (A) and PPC2 (B).

Figure S6: Gene trees of PPCK with the isozymes PPCK1 and PPCK2.

Table S1: Sampling overview and summary of RNA extractions for 24 samples.

Table S2: Titratable acidity measurements for Tillandsia vanhyningii and Tillandsia leiboldiana at

four timepoints spread over 6-hour intervals.

Table S3: Statistical models and results for titratable acidity measurements.

Table S4: Summary of per-gene read counting with FeatureCounts.

Table S5: Statistical results of the hypergeometric test of overlaps in DE genes between species

and conditions

Table S6: Overlap of DE genes betwen species, watering regimes and time points.

Table S7: Description of CAM genes in sections of the Venn diagram A (Day vs Night).

Table S8: Description of CAM genes in sections of the Venn diagram A (Drought vs Control).